Delineating antibody escape from Omicron sublineages

Henning Gruell[1,*], Kanika Vanshylla[1,*], Michael Korenkov[1], Pinkus Tober-Lau[2], Matthias

Zehner[1], Friederike Münn[2], Hanna Janicki[1], Max Augustin[3], Philipp Schommers[1,3], Leif Erik

Sander[2], Florian Kurth[2,4], Christoph Kreer[1], Florian Klein[1,5,6]

1   Laboratory of Experimental Immunology, Institute of Virology, Faculty of Medicine and

University Hospital Cologne, University of Cologne, Cologne, Germany

2   Department of Infectious Diseases and Respiratory Medicine, Charité-Universitätsmedizin Berlin,

Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany

3   Department I of Internal Medicine, Faculty of Medicine and University Hospital Cologne, University of

Cologne, Cologne, Germany

4   Department of Tropical Medicine, Bernhard Nocht Institute for Tropical Medicine and Department of

Medicine I, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

5   German Center for Infection Research (DZIF), Partner site Bonn-Cologne, Cologne, Germany

6   Center for Molecular Medicine Cologne (CMMC), University of Cologne, Cologne, Germany

*   These authors contributed equally

Correspondence should be addressed to florian.klein@uk-koeln.de.

Abstract

SARS-CoV-2 neutralizing antibodies play a critical role in prevention and treatment of COVID-19 but are challenged by viral evolution and antibody evasion, exemplified by the highly resistant Omicron BA.1 sublineage.[1-12] Importantly, the recently identified Omicron sublineages BA.2.12.1 and BA.4/5 with differing spike mutations are rapidly emerging in various countries. By determining polyclonal serum activity of 50 convalescent or vaccinated individuals against BA.1, BA.1.1, BA.2, BA.2.12.1, and BA.4/5, we reveal a further reduction of BA.4/5 susceptibility to vaccinee sera. Most notably, delineation of the sensitivity to an extended panel of 163 antibodies demonstrates pronounced antigenic differences of individual sublineages with distinct escape patterns and increased antibody resistance of BA.4/5 compared to the most prevalent BA.2 sublineage. These results suggest that the antigenic distance from BA.1 and the increased resistance compared to BA.2 may favor immune escape-mediated expansion of BA.4/5 after the first Omicron wave. Finally, while most monoclonal antibodies in clinical stages are inactive against all Omicron sublineages, we identify promising novel antibodies with high pan-Omicron neutralizing potency. Our study provides a detailed understanding of the antibody escape from the most recently emerging Omicron sublineages that can inform on effective strategies to prevent and treat COVID-19.

Main

Despite increasing levels of immunity against SARS-CoV-2 induced by vaccination and infection, the Omicron variant resulted in a global surge of infections that reflects its high transmissibility and immune evasion mediated by the highly mutated spike protein.[1,4-15] Although prolonged vaccine dosing intervals and booster immunizations based on the ancestral Wu01 strain elicit Omicron neutralizing serum activity, titers against Omicron

remain considerably lower compared to those against other variants.[4,6,11,16-18] Moreover,

the spike mutations of Omicron have rendered several therapeutic monoclonal antibodies

ineffective.[7,11,19] Most experimental evidence on the resistance of Omicron to antibody-

mediated neutralization is limited to analyses of the initial BA.1 strain. However, novel

sublineages are rapidly emerging and associated with increasing numbers of

infections.[20,21] Determining their antibody escape properties is therefore of critical

importance to effectively guide preventive and therapeutic measures. To this end, we

analyzed in detail antibody-mediated neutralization of prevalent and emerging Omicron

sublineages (BA.1, BA.1.1, BA.2, BA.2.12.1, and BA.4/5) both on a polyclonal and

monoclonal level.

Results

Rapid spread of Omicron sublineages

The spike protein of BA.1 differs in 39 amino acid residues from the ancestral Wu01 strain

of SARS-CoV-2 (**Fig. 1a**). While other Omicron sublineages share several mutations with

BA.1, they diverge at various amino acid positions including in critical antibody epitopes

(**Fig. 1a,b**). For example, the BA.1.1 spike protein contains an R346K substitution in the

receptor-binding domain (RBD) that was also observed in the SARS-CoV-2 Mu variant and

associated with escape from neutralizing antibodies (**Fig. 1a**).[22] While BA.1 and BA.1.1

dominated the initial Omicron surge, they were rapidly outcompeted by BA.2 (**Fig. 1c**).

BA.2 shares 21 of its 31 spike protein changes with BA.1 but differs considerably in both

the N-terminal domain (NTD) and the RBD, regions targeted by the most potent SARS-

CoV-2 neutralizing antibodies (**Fig. 1a,b**).

Following a decline in cases from the BA.1-/BA.2-related first Omicron wave, new

sublineages are rapidly emerging (**Fig. 1c**). For example, BA.2.12.1 outcompetes other

BA.2 sublineages in the United States resulting in an increase of infections. Moreover,

within weeks of their identification, the BA.4 and BA.5 sublineages with an identical spike

protein have become dominant in South Africa and Portugal and driven rising case

numbers (**Fig. 1c**).[21] BA.2.12.1 and BA.4/5 share the majority of their spike mutations

with BA.2 (94% and 90%, respectively), but contain additional changes at sites associated

with antibody escape. These include substitutions at residue 452 (L452Q in BA.2.12.1 and

L452R in BA.4/5, respectively) that were also recorded for the Lambda and Delta variants

(**Fig. 1a**).[23,24] In addition, the BA.4/5 spike protein harbors the F486V RBD mutation that

can reduce monoclonal antibody sensitivity but had not yet been observed in other

variants of interest or concern.[25] Thus, newly emerged Omicron sublineages differ from

BA.1 in key residues of the spike protein. Given their apparent growth advantages

compared to BA.1 and the prevalent BA.2, the sublineages BA.2.12.1 and BA.4/5 are likely

to be amongst the dominant variants of the SARS-CoV-2 pandemic in the near future.

Reduced serum neutralizing activity against Omicron sublineages

To determine Omicron sublineage escape profiles, we measured neutralizing activity

against the ancestral Wu01 strain as well as BA.1, BA.1.1, BA.2, BA.2.12.1, and BA.4/5

using lentivirus-based pseudovirus assays.[26,27] In order to assess escape from polyclonal

serum activity, we collected samples from two longitudinal cohorts of i.) SARS-CoV-2-

convalescent individuals (*n*=20) and ii.) vaccinated health care workers (*n*=30)

(**Extended Data Table 1**).[27,28] In both cohorts, individuals received an mRNA vaccine

booster immunization (BNT162b2) after a median of 14 months and 9 months following

infection or two doses of BNT162b2, respectively.

Convalescent individuals had a median age of 51 years (interquartile range [IQR]

35-60) and were diagnosed with mild or asymptomatic SARS-CoV-2 infection. All

individuals were infected during the initial phase of the pandemic between February and

April 2020, prior to the emergence of designated viral variants of concern. Early post-

infection samples (V1) were collected at a median of 48 days (IQR 34-58) after disease

onset or diagnosis and neutralizing activity was assessed by determining the 50%

inhibitory serum dilutions ($ID_{50}s$) (**Fig. 2a**). Neutralization of the Wu01 strain was

detected in all samples (100%) obtained early after infection, with individual ID_{50} values

ranging from 16 to 2,607 (geometric mean ID_{50} [$GeoMeanID_{50}$] of 264) (**Fig. 2b**). In

contrast, serum activity against Omicron sublineages was strongly reduced and only

detectable in 0-15% for BA.1 and BA1.1, and 45-50% for BA.2, BA.2.12.1, and BA.4/5 (**Fig.**

2b). However, analysis of sera obtained at a median of 33 days (IQR 27-54) after a single

BNT162b2 booster immunization revealed neutralizing activity against all tested

Omicron sublineages (**Fig. 2b, Extended Data Fig. 1a**) with $GeoMeanID_{50}s$ ranging from

1,456 against BA.4/5 to 2,103 against BA.2 (**Fig. 2b, Extended Data Fig. 1b,c**).

In addition, we determined Omicron sublineage neutralizing activity induced

solely by vaccination (**Fig. 2c**). At a median of 28 days (IQR 27-32) after completion of the

initial two-dose course of BNT162b2 (V1), Wu01 neutralizing serum activity was detected

in all 30 individuals with a $GeoMeanID_{50}$ of 561 (**Fig. 2d**). Although Omicron sublineage

neutralization was detectable in 43-73% of vaccinated individuals at V1, $GeoMeanID_{50}s$

were very low (ranging from 8-17) (**Fig. 2d**). Follow-up samples obtained at a median of

29 days (IQR 26-35) after booster immunization showed 8-fold higher activity against

Wu01 ($GeoMeanID_{50}$ of 4,773) and strongly increased activity against all Omicron

sublineages (27- to 70-fold $GeoMeanID_{50}$ increases) (**Fig. 2d, Extended Data Fig. 1d-f**).

Similar to the convalescent cohort, the lowest $GeoMeanID_{50}$ after booster immunization

was observed against the BA.4/5 sublineages (GeoMeanID$_{50}$ of 312, **Fig. 2d**). Of note,

amongst vaccinated individuals, the neutralizing activity against BA.4/5 was statistically

significantly lower compared to that against BA.1, BA.1.1, and BA.2 (GeoMeanID$_{50}$s of 648,

557, and 582, respectively) (**Extended Data Fig. 1e**).

We conclude that booster immunizations are critical to elicit neutralizing serum

activity against all prevalent Omicron sublineages in vaccinated as well as convalescent

individuals. Notably, BA.4/5 consistently demonstrated the lowest sensitivity of all

sublineages against polyclonal serum. These results suggest increased antigenic escape of

the recently emerged BA.4/5 sublineages mediated by the small number of spike protein

differences relative to BA.2.

Dissecting escape of Omicron sublineages

To decode Omicron escape from neutralizing antibodies on a molecular level, we

produced and tested 158 monoclonal antibodies isolated from SARS-CoV-2-convalescent

individuals against all sublineages. These included 67 randomly selected antibodies from

the Coronavirus Antibody Database (CoV-AbDab),[29] 79 antibodies isolated in our

previous work,[30,31] as well as 12 antibodies that were clinically tested. In total, this

antibody panel originated from at least 43 different individuals out of 19 independent

studies (**Fig. 3a**, **Extended Data Table 2**). All antibodies were identified in unvaccinated

individuals that were infected in the early phase of the pandemic prior to the emergence

of Omicron. The selection encompassed a broad spectrum of SARS-CoV-2 neutralizing

antibodies with 92 V_H/V_L combinations and a diverse range of CDR3 lengths and V gene

mutations (**Fig. 3a**, **Extended Data Table 2**). Most antibodies targeted the RBD (97%)

and the panel included previously described public clonotypes such as the V_H3-53/3-66

subgroup (**Fig. 3a**, **Extended Data Table 2**).

While all 158 antibodies neutralized the Wu01 strain, only 18%, 17%, 22%, 23%

and 18% remained active against BA.1, BA.1.1, BA.2, BA.2.12.1, and BA.4/5, respectively

(**Fig. 3b, Extended Data Fig. 2a, Extended Data Table 2**). Moreover, amongst antibodies

with retained neutralizing activity, the overall potency against Omicron sublineages was

6- to 14-fold lower compared to Wu01 with GeoMeanIC$_{50}$s of 0.42 (BA.1), 0.52 (BA.1.1),

0.17 (BA.2), 0.18 (BA.2.12.1), and 0.28 μg/ml (BA.4/5) (**Fig. 3c**).

The neutralization profiles revealed both commonalities and diversity in antibody

sensitivity of the different Omicron sublineages (**Fig. 3d-f, Extended Data Fig. 2b**).

Between the most closely related sublineages (BA.1 and BA.1.1; BA.2 and BA.2.12.1), only

small differences in antibody activity were observed (r$_s$ of 0.81 and 0.86, respectively)

(**Fig. 3e,f Extended Data Fig. 2c**). In contrast, comparisons between the primary

sublineages BA.1, BA.2 and BA.4/5 demonstrated higher degrees of variation in antibody

susceptibility (**Fig. 3e,f**). For example, BA.2 was more resistant (>1 log$_{10}$ IC$_{50}$ difference)

than BA.1 to only a single antibody (2%), but more sensitive to ten antibodies (23%) (**Fig.**

3e). Moreover, higher fractions of antibodies showed >10-fold higher resistance against

BA.4/5 relative to other variants (**Fig. 3e**). While the comparison of BA.4/5 to BA.1 and

BA.1.1 revealed heterogeneity in sensitivity to the antibody panel with no correlation, the

sensitivity of BA.4/5 and BA.2 or BA.2.12.1 more strongly correlated (r$_s$ of 0.58 and 0.68,

respectively). Importantly, however, whereas BA.4/5 were more sensitive to only 0-5%

of antibodies compared to BA.2 and BA.2.12.1, BA.4/5 showed higher resistance to 21%

of the antibodies (**Fig. 3e**).

Based on the analysis of the sublineage neutralization profiles, three prevalent

classes of Omicron neutralizing antibodies became apparent: i.) antibodies with broadly

comparable activity against Omicron sublineages (21/43, 49%), ii.) antibodies with

strongly reduced activity against BA.1 and BA.1.1 compared to BA.2, BA.2.12.1, and

BA.4/5 (7/43, 16%), and iii.) antibodies with strongly reduced activity against BA.4/5

compared to BA.1, BA.1.1, as well as BA.2 and/or BA.2.12.1 (6/43, 14%).

We conclude that in contrast to the limited differences in sensitivity to polyclonal

serum, Omicron sublineages considerably differ in their sensitivity to monoclonal

antibodies. Importantly, BA.4/5 demonstrated a strong bias to higher resistance

compared to the prevalent BA.2 sublineages.

Minor sequence variations critically affect antibody activity

Antibodies with Omicron neutralizing activity carried a modestly higher number of amino

acid mutations in the heavy and light chain variable gene compared to antibodies only

neutralizing Wu01 (median of 6 vs. 4 and 4 vs. 3 mutations; p=0.0143 and p=0.0170,

respectively), suggesting that higher sequence diversification might be overall favorable

for Omicron neutralization (**Fig. 3f, Extended Data Fig. 2e**).

Antibody responses against SARS-CoV-2 are highly convergent across individuals

based on the identification of public clonotypes with conserved sequence characteristics

and neutralization mechanisms.[32,33,34] Amongst the analyzed antibody panel, 18

sequences from 11 individuals could be assigned to the prominent V_H3-53/3-66|V_K1-9

clonotype (**Fig. 3g, Extended Data Fig. 2f**).[30,35,36] Although these antibodies are highly

conserved on a sequence level, they substantially differed in their Omicron neutralizing

capacity (**Fig. 3g, Extended Data Fig. 2f**). For example, antibodies R207-1C4 and R568-

2G5 harbor eight amino acid mutations in their V_H gene segment (without CDRH3) of

which five are at the same position and three are identical. While both antibodies had

similar Wu01 neutralizing activity, R207-1C4 did not neutralize any Omicron sublineage,

whereas R568-2G5 showed neutralizing activity against all variants. Notably, antibody

C140, another member of this clonotype which has the identical CDRH3 motif as R568-

2G5, did not neutralize any Omicron variant.

We conclude that minimal variation in antibody sequences can tip the scale

between Omicron neutralization and resistance. Thus, sequence- or antibody class-based

predictions of Omicron neutralization are difficult and experimental assessment of

individual monoclonal antibody activity remains essential.

Impact on clinical and novel monoclonal antibodies

SARS-CoV-2 neutralizing monoclonal antibodies can reduce morbidity and mortality in

infected individuals and are critical for passive immunization to protect individuals that

do not mount an adequate immune response upon vaccination.[2,3,37-41] To determine how

spike protein mutations of Omicron sublineages affect antibodies in clinical use, we

analyzed 9 monoclonal antibodies that received authorization for clinical use (**Fig. 4a,b**)

and 9 that are advanced in clinical development (**Fig. 4b, Extended Data Fig. 3**). All

antibodies target the RBD of the SARS-CoV-2 spike protein and were tested against Wu01

and the Omicron sublineages (**Fig. 4b**).

Most antibodies showed highly potent neutralizing activity against Wu01 with

IC_{50}s below 0.005 µg/ml (**Fig. 4b**). Less potent and incomplete Wu01 neutralizing activity

was observed for sotrovimab, consistent with previous reports showing reduced activity

against pseudoviruses lacking the dominant D614G spike mutation (**Fig. 4a**).[42,43] Only five

(28%) out of the 18 tested antibodies neutralized BA.1 with IC_{50}s <10 µg/ml, and of these,

three had strongly reduced potency (**Fig. 4b**). While the antibody neutralization profile of

BA.1.1 was generally similar to that of BA.1, there were some differences. For example,

DZIF-10c neutralized BA.1 with an IC_{50} of 0.046 µg/ml but completely lost activity against

BA.1.1 (**Fig. 4b, Extended Data Fig. 3**).

Although the number of antibodies in clinical use with neutralizing activity against

BA.2 (5 out 18, 28%), BA.2.12.1 (7 out 18, 39%), and BA.4/5 (6 out 18, 33%) was similarly

small, the neutralization profiles did differ (**Fig. 4b**). For example, cilgavimab neutralized

BA.2 and BA.2.12.1 with high potency (IC_{50}s of 0.008 and 0.014 µg/ml, respectively) but

was >480-fold less potent against BA.1 and BA.1.1 (**Fig. 4a,b**). While, cilgavimab

maintained high potency against BA.4/5 (IC_{50} of 0.085 µg/ml), the activity of antibody

DZIF-10c against BA.4/5 was strongly reduced when compared to BA.2 and BA.2.12.1

(8.64 vs. 0.03 µg/ml) (**Fig. 4b**). In contrast, whereas antibody imdevimab showed no

appreciable activity against BA.1 and BA.1.1, neutralization of BA.2, BA.2.12.1, and BA.4/5

was detectable at low levels (**Fig. 4a,b**).

Out of all clinical antibodies tested, only bebtelovimab maintained high

neutralization potency against all Omicron sublineages (IC_{50} <0.005 µg/ml; **Fig. 4a,b**). To

identify additional promising candidates, we examined antibodies previously isolated

from individuals with 'elite' IgG neutralizing activity.[30] Remarkably, antibodies R207-

2F11, R568-1G9, and R200-1F9 demonstrated highly potent neutralizing activity against

previously circulating variants (Wu01, B.1, Alpha, Beta, Delta, Epsilon, and Kappa), as well

as against BA.1, BA.1.1, BA.2, BA.2.12.1, and BA.4/5 with IC_{50}s <0.05 µg/ml (**Fig. 4d,**

Extended Data Table 2). These antibodies were identified from three different

convalescent individuals within weeks of infection during the early phase of the

pandemic.[30] Thus, the results highlight the capability of the immune system to generate

potent antibodies against the highly divergent Omicron variants in response to contact

with the ancestral spike protein.

We conclude that the prevalent and newly emerged Omicron sublineages,

including BA.2.12.1 and BA.4/5, escape from most monoclonal antibodies in clinical use.

Omicron sensitivity to these antibodies can, however, strongly differ on the sublineage

level.

Discussion

Antibody-mediated immunity is a critical component of prophylactic and therapeutic

measures against SARS-CoV-2 infection.[2,3,44,45] Throughout the COVID-19 pandemic, it has

however been confronted by the periodic emergence of antibody escape variants

developing in the context of increasing population immunity.[46] Omicron sublineages with

diverse spike proteins gaining prevalence suggest further increase in immune escape

and/or transmissibility.[20,21,47] Thus, establishing the impact of the recent BA.2.12.1 and

BA.4/5 sublineages on polyclonal and monoclonal immunity is critical to guide antibody-

mediated strategies for prevention and treatment.

Our results demonstrate that Wu01-based mRNA vaccine boosters are effective in

eliciting serum neutralizing activity against the diverse Omicron sublineages. However,

this activity is strongly limited compared to Wu01 neutralization and lower BA.4/5

neutralizing titers indicate even more pronounced immune evasion relative to previous

Omicron sublineages. Interestingly, the relative reduction in activity against BA.4/5 after

booster immunization was higher in infection-naïve than in convalescent individuals. This

might indicate differences in the breadth and potency of the neutralizing antibody

response elicited by infection only compared to those induced by "hybrid immunity".

While the differences in polyclonal serum sensitivity between the Omicron

sublineages were overall modest, our analysis of an extended antibody panel revealed

distinct patterns in sublineage sensitivity on a monoclonal level. Comparing Omicron

sublineages revealed the largest antigenic distance between BA.4/5 and BA.1/BA.1.1 as

indicated by the lack of correlation in antibody sensitivity and notable fractions of

antibodies showing higher sensitivity to one or the other sublineage. Whereas BA.4/5 and

BA.2/BA.2.12.1 showed more similar antibody susceptibility profiles, almost all

differences manifested as higher antibody resistance of BA.4/5.

Notably, minor differences in antibody sequence strongly affected Omicron

activity of antibodies with high potency against Wu01, highlighting the relative difficulty

of inducing antibodies with optimized Omicron neutralization after single contact with

Wu01-related spike. Accordingly and consistent with previous analyses on BA.1 and BA.2,

[48,49] almost all antibodies in clinical investigation showed strongly reduced or abolished

activity against all Omicron sublineages. However, we identify novel antibodies isolated

after ancestral infection with remarkable activity against all tested variants that may

provide options for immune-mediated treatment and prevention in the Omicron era of

the pandemic.

Our results suggest that outcompetition of BA.1(.1) by BA.2 may have been driven

mostly by higher transmissibility rather than by immune evasion. In contrast, higher

antigenic distance and antibody resistance of BA.4/5 may support its expansion after the

initial Omicron wave. The rapid emergence of Omicron sublineages with reduced

antibody susceptibility highlights the challenges posed by constant viral evolution and the

critical need for genomic surveillance and sensitivity assessments to inform on antibody-

based prophylactic and therapeutic measures.

Acknowledgments

We are grateful to all study participants for their dedication to our research. We thank F.

Dewald, L. Gieselmann, B. Kurt, C. Lehmann, P. Mayer, N. Riet, S. Salomon, M. Schlotz, R.

Schröder, R. Stumpf, and H. Wüstenberg, as well as the members of the EICOV/COVIM

Study Group (Y. Ahlgrimm, L. Bardtke, K. Behn, N. Bethke, H. Bias, D. Briesemeister, C.

Conrad, V. M. Corman, C. Dang-Heine, S. Dieckmann, C. Eroglu, D. Frey, J.-A. Gabelich, J.

Gerdes, U. Gläser, A. Hetey, L. Hasler, E. T. Helbig, D. Hillus, W. Hirst, A. Horn, C. Hülso, S.

Jentzsch, C. von Kalle, L. Kegel, A. Krannich, W. Koch, P. Kopankiewicz, P. Kroneberg, H. Le,

L. J. Lippert, C. Lüttke, P. de Macedo Gomes, B. Maeß, C. Matthaiou, J. Michel, A. Nitsche, A.-

M. Ollech, C. Peiser, A. Pioch, C. Pley, K. Pohl, C. Rubisch, L. Ruby, A. Sanchez Rezza, I.

Schellenberger, V. Schenkel, J. Schlesinger, S. Schmidt, G. Schwanitz, T. Schwarz, J. Seybold,

A. Solarek, A. Stege, S. Steinbrecher, P. Stubbemann, C. Thibeault, and D. Treue) for sample

acquisition and processing; L. Buchholz and M. Wunsch for help with antibody production

and preparation; and J.J. Malin for providing clinical antibody stocks. We thank all

laboratories obtaining and submitting SARS-CoV-2 sequencing information to GISAID for

facilitating rapid analyses of global variant distributions. This work was supported by

grants from COVIM: NaFoUniMedCovid19 (FKZ: 01KX2021) (to L.E.S. and F. Klein), the

Federal Institute for Drugs and Medical Devices (V-2021.3 / 1503_68403 / 2021–2022)

(to L.E.S. and F. Kurth), the German Center for Infection Research (DZIF) (to F. Klein), and

the Deutsche Forschungsgemeinschaft (DFG) CRC1310 (to C.K. and F. Klein).

311   Author contributions

Conceptualization, H.G., K.V., C.K., and F. Klein; methodology, H.G., K.V., L.E.S., F. Kurth, C.K.,

and F. Klein; investigation, H.G. and K.V.; resources, P.T-L., M.K., M.Z., F.M., H.J., M.A., and

P.S.; formal analysis, H.G., K.V., and C.K.; writing-original draft, H.G., K.V., C.K., and F. Klein;

writing-review and editing, M.K., P.T-L., and F. Kurth; visualization, H.G., K.V., C.K., and F.

Klein; supervision, L.E.S., F. Kurth, and F. Klein; funding acquisition, L.E.S., F. Kurth, and F.

Klein.

Figure legends

Fig. 1. Omicron sublineage differences.

a, Spike amino acid changes in the BA.1, BA.1.1, BA.2, BA.2.12.1, and BA.4/5 Omicron sublineages relative to Wu01. **b,** Locations of changed amino acids in Omicron sublineages highlighted on the SARS-CoV-2 spike (PDB: 6XR8), with colors indicating mutations shared between different sublineages. Representations at the bottom show RBD and NTD outlined in black. Changes exclusive to BA.1.1 are indicated in green and those to BA.4/5 in brown, residue 452 (mutated in BA.2.12.1 and BA.4/5) is indicated in purple. **c,** Top panels indicate weekly reported SARS-CoV-2 infections as aggregated by the COVID-19 Data Repository by the Center for Systems Science and Engineering at Johns Hopkins University and http://ourworldindata.org. Relative proportions of infections by variant per week were extrapolated from the proportion of sequences submitted to the GISAID SARS-CoV-2 database per variant and week (accessed on May 24, 2022) shown in the bottom panels. NTD, N-terminal domain; RBD, receptor-binding domain; PDB, Protein Data Bank.

Fig. 2. Omicron sublineage neutralizing serum activity in vaccinated and convalescent individuals.

a, Study scheme in COVID-19-convalescent individuals. Samples were collected after infection occurring between February and April, 2020 (V1), and after a BNT162b2 booster immunization (V2). **b,** Fifty-percent inhibitory serum dilutions (ID_{50}s) against Wu01, BA.1, BA.1.1, BA.2.12.1, and BA.4/5 determined by pseudovirus neutralization assays in convalescent individuals. Bars indicate geometric mean ID_{50}s with 95% confidence intervals (CIs) at V1 (left) and V2 (right). Numbers indicate geometric mean ID_{50}s and percentage of individuals with detectable neutralizing activity (ID_{50} >10) in parentheses. **c,** Study scheme in vaccinated individuals. Samples were collected after the second dose of BNT162b2 (V1) and after the third dose of BNT162b2 (V2). **d,** Serum ID_{50}s against Wu01, BA.1, BA.1.1, BA.2.12.1, and BA.4/5 determined by pseudovirus neutralization assays in vaccinated individuals. Bars indicate geometric mean ID_{50}s with 95% CIs at V1 (left) and V2 (right). Numbers indicate geometric mean ID_{50}s and percentage of individuals with detectable neutralizing activity (ID_{50} >10) in parentheses. In **b** and **d**, ID_{50}s below the lower limit of quantification (LLOQ, ID_{50} of 10; indicated by black dotted lines) were imputed to ½ LLOQ (ID_{50}=5), and ID_{50}s above the upper limit of quantification (21,870) were imputed to 21,871.

Fig. 3. Determining Omicron sublineage immune escape using monoclonal antibodies.

a, SARS-CoV-2 neutralizing monoclonal antibodies (*n*=158) derived from 19 studies and ≥43 convalescent individuals were analyzed. Bar charts indicate number of antibodies per heavy chain variable gene segment (V_H), amino acid (aa) length of the heavy chain complementarity-determining region 3 (CDRH3), and number of V_H aa mutations relative

to the V_H germline gene. Pie chart indicates antibody epitopes with slice sizes proportional

to antibody fractions. RBD, receptor-binding domain. **b,** Fractions of antibodies from the

panel (*n*=158) displaying activity (IC_{50} <10 μg/ml) against Wu01 and Omicron

sublineages. **c,** IC_{50}s of neutralizing antibodies against individual variants/sublineages

(Wu01, *n*=158; BA.1, *n*=29; BA.1.1, *n*=27; BA.2, *n*=34; BA.2.12.1, *n=36*; BA.4/5, *n=29*). Solid

lines depict geometric mean IC_{50}s and 95% confidence intervals, and dashed lower and

upper lines indicate lower (LLOQ, 0.005 μg/ml) and upper limit of quantification (ULOQ,

10 μg/ml), respectively. IC_{50}s <LLOQ were imputed to ½ LLOQ (IC_{50}=0.0025). **d,** Heat map

depicting IC_{50}s of the subset of antibodies (*n*=43) with neutralizing activity (IC_{50} <10

μg/ml) against at least one of the Omicron sublineages. Antibodies are sorted based on

their potency against BA.1. **e,** Log_{10} IC_{50} ratios of antibodies neutralizing any Omicron

sublineage for indicated sublineage comparisons. Within each panel, antibodies are

sorted by increasing IC_{50} ratios. Ratios <-1 log_{10} are highlighted in green, ratios between

-1 and 1 log_{10} in grey, and ratios >1 log_{10} in red. Numbers indicate relative fractions. IC_{50}

<LLOQ were imputed to ½ LLOQ (IC_{50}=0.0025) and IC_{50} values >ULOQ were imputed to

2x ULOQ (IC_{50}=20). **f,** Spearman's rank correlation coefficients (rho) for comparisons of

antibodies with neutralizing capacity against any Omicron sublineage (*n*=43) as shown in

Fig. S2c. Numbers in bold indicate values with *p*<0.05. **g,** Heavy chain V amino acid

mutations of antibodies with neutralizing activity against Wu01 only and antibodies

neutralizing both Wu01 and at least one omicron sublineage. Lines indicate medians and

interquartile ranges. Groups were compared using the two-tailed Mann-Whitney U test.

h, Phylogenetic tree and heavy chain sequence alignment of antibodies of the V_H3-53/3-

66|V_K1-9 public clonotype. Letters indicate aa mutations relative to the V_H germline gene.

Number of aa mutations from corresponding germline allele and neutralizing activity

against Omicron sublineages are indicated on the right. Germline V_H represents the

consensus of all identified germline alleles of depicted antibodies (V_H3-53*01, V_H3-53*04, and V_H3-66*01). Number of aa mutations from corresponding germline allele and neutralizing activity against Wu01 and Omicron sublineages are indicated on the right.

Fig. 4. Omicron sublineage neutralizing activity of monoclonal antibodies in clinical use.

a, Dose response curves showing percentage neutralization of monoclonal antibodies against Wu01 and Omicron sublineages in pseudovirus neutralization assay. Circles show averages and error bars indicate standard deviation. Dotted lines indicate 50% neutralization (IC_{50}). **b,** IC_{50}s of monoclonal antibodies with current or previous authorization for clinical use or in clinical development. Symbols indicate whether clinical products or parental antibodies produced as human IgG1 were used. **c,** Dose response curves showing percentage neutralization of novel monoclonal antibodies against Wu01 and Omicron sublineages in pseudovirus neutralization assay. Circles show averages and error bars indicate standard deviation. Dotted lines indicate 50% neutralization (IC_{50}). **d,** IC_{50}s of novel monoclonal antibodies shown in **c** against Wu01 and Omicron sublineages (upper rows) and previously circulating variants (lower rows, as previously determined [30]).

Extended Data Figure legends

Extended Data Fig. 1. Serum neutralization of Omicron sublineages.

a, Serum ID_{50}s against Wu01 and Omicron sublinages in the cohort of convalescent individuals after infection (V1) and BNT162b2 booster immunization (V2) as in **Fig. 2**. Lines connect ID_{50}s of individual participants at V1 and V2. Dashed lines indicate lower

limit of quantification (LLOQ, $ID_{50}=10$). **b,** Numbers in black indicate Spearman's rank

correlation coefficients (rho) for comparisons at V2 shown in **c**. Numbers in red indicate

p values determined by Friedman test followed by Dunn's multiple comparison test for

comparisons at V2, with statistically significant differences indicated in bold. **c,**

Correlation plots of $\log_{10}$ serum ID_{50}s against indicated viruses in convalescent

individuals at V2. **d,** Serum ID_{50}s against Wu01 and Omicron sublineages in the cohort of

BNT162b2-vaccinated individuals after the second (V1) and the third vaccine dose (V2)

as in **Fig. 2**. Lines connect ID_{50}s of individual participants at V1 and V2. Dashed lines

indicate lower limit of quantification (LLOQ, $ID_{50}=10$). **e,** Numbers in black indicate

Spearman's rank correlation coefficients (rho) for comparisons at V2 shown in **f**. Numbers

in red indicate *p* values determined by Friedman test followed by Dunn's multiple

comparison test for comparisons at V2, with statistically significant differences indicated

in bold. **f,** Correlation plots of $\log_{10}$ serum ID_{50}s against indicated viruses in vaccinated

individuals at V2. In **a** and **d**, Serum ID_{50}s <LLOQ were imputed to ½ LLOQ ($ID_{50}=5$).

Extended Data Fig. 2. Neutralization profile of Omicron neutralizing antibodies.

a, Spider plot of IC_{50}s for all antibodies against Wu01 and Omicron sublineages.

Antibodies are sorted arbitrarily but equally for each virus. Circles indicate IC_{50}s (from

outer to inner: 0.005, 0.05, 0.5, and 5 µg/ml). **b,** Bar charts of antibodies with neutralizing

activity (IC_{50} <10 µg/ml) against any Omicron sublineage (*n*=43). In each chart, antibodies

are sorted by BA.1-neutralizing activity. Dotted lines indicate lower (LLOQ, 0.005 µg/ml)

and upper limits of quantification (ULOQ; 10 µg/ml). **c,** IC_{50} correlation plots for

antibodies with neutralizing capacity against any Omicron sublineage (*n*=43). Colors

indicate epitopes as in **Fig. 3a**. Grey dashed lines represent identity lines and black dashed

lines indicate limits of quantification. In **a-c**, IC_{50}s <LLOQ were imputed to ½ LLOQ

(IC_{50}=0.0025) and IC_{50}s >ULOQ were imputed to 2x ULOQ (IC_{50}=20). **d,** Light chain V

amino acid mutations of antibodies with neutralizing activity against Wu01 only and

antibodies neutralizing both Wu01 and at least one omicron sublineage. Lines indicate

medians and interquartile ranges and groups were compared using the two-tailed Mann-

Whitney U test. **e,** Light chain sequence alignment of antibodies of the V_H3-53/3-66|V_K1-

9 public clonotype. Letters indicate amino acid (aa) mutations relative to the light chain

germline gene. Germline V_K represents the consensus of all identified germline alleles of

the depicted antibodies. Number of aa mutations from corresponding germline allele and

neutralizing activity against Wu01 and Omicron sublineages are indicated on the right.

Extended Data Fig. 3. Omicron sublineage neutralizing activity of monoclonal

antibodies in clinical testing.

Dose response curves showing % neutralization of monoclonal antibodies against Wu01

and Omicron sublineages in a pseudovirus neutralization assay. Circles show averages

and error bars indicate standard deviation. Dotted lines indicate 50% neutralization

(IC_{50}).

Extended Data Tables

Extended Data Table 1: Study cohorts

Extended Data Table 2: Monoclonal antibody panel analyses

Methods

Study cohort and sample collection

Serum samples from COVID-19-convalescent individuals were collected at the University

Hospital Cologne under study protocols approved by the ethics committee (EC) of the

Medical Faculty of the University of Cologne (16-054 and 20-1187). Between April and

May, 2020, individuals with a history of SARS-CoV-2 infection confirmed by polymerase

chain reaction (documented through a written test certificate or as reported to study

investigators by the participants) were enrolled within eight weeks of symptom onset

and/or diagnosis. As all participants were enrolled early during the pandemic (i.e., prior

to the emergence of variants of concern, as designated by the World Health Organization),

most individuals are likely to have been infected with an early viral strain similar to Wu01.

Participants were followed longitudinally to analyze long-term immunity to SARS-CoV-2.

Follow-up samples after booster immunization were collected between May and August,

2021. No additional SARS-CoV-2 infections between the sampling time points were

reported by any of the participants.

Serum samples from vaccinated individuals were collected under protocols

approved by the EC of Charité - Universitätsmedizin Berlin (EICOV, EA4/245/20) as well

as the EC of the Federal State of Berlin and the Paul Ehrlich Institute (COVIM, EudraCT-

No. 2021-001512-28). Study participation irrespective of medical conditions was offered

to health-care workers vaccinated at the Charité – Universitätsmedizin (Berlin, Germany).

All serum samples were tested for antibodies targeting the SARS-CoV-2 nucleocapsid

using the SeraSpot Anti-SARS-CoV-2 IgG microarray-based immunoassay (Seramun

Diagnostica). Samples from individuals with a history of SARS-CoV-2 infection, a positive

SARS-CoV-2 nucleic acid amplification test (performed at sampling), or detectable anti-

nucleocapsid antibodies were not included in this analysis. Serum samples after two

vaccinations were collected in February and March 2021 (for 29/30 participants; from

one participant, the early serum sample was obtained in July 2021); samples after booster

immunization were collected in December 2021 and January 2022.

All study participants provided written informed consent. Vaccinations in both

cohorts were performed as part of routine care outside of the observational studies.

Selection of participants and samples for analysis was based on receipt of identical

vaccines and comparable sampling time points relative to vaccinations. Serum samples

were collected after centrifugation and stored at -80°C until analysis.

SARS-CoV-2 pseudovirus constructs

All SARS-CoV-2 spike proteins were expressed using codon-optimized expression

plasmids. Wu01 (EPI_ISL_406716), BA.2.12.1, and BA.4/5 pseudoviruses were produced

using an expression plasmid that incorporated a C-terminal deletion of 21 cytoplasmic

amino acids that result in increased pseudovirus titers. Expression plasmids for Omicron

sublineage spike proteins were produced by assembling and cloning codon-optimized

overlapping gene fragments (Thermo Fisher) into the pCDNA3.1/V5-HisTOPO vector

(Thermo Fisher) using the NEBuilder HiFi DNA Assembly Kit (New England Biolabs), and

included the full spike protein amino acid sequences with the following amino acid

changes relative to Wu01:

BA.1: A67V, Δ69-70, T95I, G142D, Δ143-145, N211I, Δ212, ins215EPE, G339D, S371L,

S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R,

N501Y, Y505H, T547K, D614G, H655Y, N679K, P681H, N764K, D796Y, N856K, Q954H,

N969K, and L981F.

BA.1.1: As for BA.1 with an additional R346K mutation.

BA.2: T19I, Δ24-26, A27S, A67V, G142D, V213G, G339D, S371F, S373P, S375F, T376A,

D405N, R408S, K417N, N440K, S477N, T478K, E484A, Q493R, Q498R, N501Y, Y505H,

D614G, H655Y, N679K, P681H, N764K, D796Y, Q954H, N969K.

BA.2.12.1: As for BA.2 with additional L452Q and S704L mutations.

BA.4/5: As for BA.2 with additional Δ69-70, L452R, and F486V mutations, but lacking the

Q493R mutation.

All plasmid sequences were verified by sequencing.

Monoclonal antibodies

Monoclonal antibodies previously isolated in our lab had been obtained by single cell-

sorting of SARS-CoV-2 spike-specific B cells followed by reverse transcription, PCR

amplification and cloning of antibody variable regions.[30,31] For monoclonal antibodies

derived from the CoV-AbDab [29], variable region amino acid sequences were reverse

translated with the reverse translate tool from the Sequence Manipulation Suite[50] using

the *Homo sapiens* codon table obtained from the Codon Usage Database,[51] and sequences

were ordered as gene fragments from Integrated DNA Technologies (IDT) with 5' and 3'

overhangs. The variable regions were inserted into heavy and light chain expression

plasmids[52] by sequence- and ligation-independent cloning (SLIC). For antibodies ADG-2,

COV2-2130 (cilgavimab), COV2-2196 (tixagevimab), COV2-2381, MAD0004J08, and P2C-

1F11 (BRII-196), gene fragments based on the nucleotide sequences published in

GenBank were ordered at IDT and cloned as above. For antibodies C135, CT-P59

(regdanvimab), and LY-CoV1404 (bebtelovimab), gene fragments based on antibody

structures deposited in the Protein Data Bank (accession nos. 7K8Z, 7CM4, and 7MMO)

were ordered at IDT after codon optimization using the IDT Codon Optimization Tool and

cloned as above. For antibodies 47D11, BD-368-2, C144, and P2B-2F6, amino acid

sequences were derived from CoV-AbDaB, corresponding nucleotide sequences

generated and codon-optimized using the IDT Codon Optimization Tool, and gene

fragments cloned as above.

Monoclonal antibody production was performed using 293-6E cells (National

Research Council of Canada) by co-transfection of heavy and light chain expression

plasmids using 25 kDa branched polyethylenimine (Sigma-Aldrich). Culture supernatants

were harvested after an incubation period of 6-7 days at 37°C and 6% CO_2 under constant

shaking in FreeStyle Expression Medium supplemented with penicillin (20 U/ml) and

streptomycin (20 μg/ml) (all Thermo Fisher). Clarified cell supernatants were incubated

with Protein G Sepharose 4 FastFlow (Cytiva) overnight at 4°C . After centrifugation,

antibodies bound to Protein G beads were eluted in chromatography columns (Bio-Rad)

using 0.1 M glycine (pH=3.0) and buffered in 1 M Tris (pH=8.0). Buffer exchange to PBS

was performed using centrifugal filter units (Millipore). For antibodies bamlanivimab,

casirivimab, DZIF-10c, etesevimab, imdevimab, and sotrovimab, aliquots from clinical

stocks were used.

Pseudovirus neutralization assays

Neutralization assays were performed using lentivirus-based pseudoviruses and ACE2-

expressing 293T cells.[26,27] Pseudovirus particle production was performed in HEK293T

cells by co-transfection of individual expression plasmids encoding for the SARS-CoV-2

spike protein, HIV-1 Tat, HIV-1 Gag/Pol, HIV-1 Rev, and luciferase-IRES-ZsGreen using

FuGENE 6 Transfection Reagent (Promega). Culture supernatants were exchanged with

fresh medium (high glucose DMEM supplemented with 2 mM L-glutamine, 100 IU/ml

penicillin, 100 μg/ml streptomycin, 1 mM sodium pyruvate (all Thermo Fisher), and 10%

FBS (Sigma-Aldrich)) 24 h post transfection. Pseudovirus-containing supernatants were

harvested between 48-72 h after transfection, centrifuged, clarified using a 0.45 μm filter,

and stored at -80°C. Pseudoviruses were titrated by infection of 293T-ACE2 cells and

luciferase activity was determined after a 48-hour incubation at 37°C and 5% CO_2 by

addition of luciferin/lysis buffer (10 mM $MgCl_2$, 0.3 mM ATP, 0.5 mM coenzyme A, 17 mM

IGEPAL CA-630 (all Sigma-Aldrich), and 1 mM D-Luciferin (GoldBio) in Tris-HCL) using a

microplate reader (Berthold).

Serum samples were heat-inactivated at 56°C for 45 min before use. Three-fold

serial dilutions of serum (starting at 1:10) and monoclonal antibodies (starting at 10

μg/ml) were prepared in culture medium and co-incubated with pseudovirus

supernatants for one hour at 37°C and 5% CO_2 prior to addition of 293T-ACE2 cells.

Following a 48-hour incubation at 37°C and 5% CO_2, luciferase activity was determined

as described above. Average background relative light units (RLUs) of non-infected cells

were subtracted, and serum ID_{50}s and antibody IC_{50}s were determined as the serum

dilutions and antibody concentrations resulting in a 50% RLU reduction compared to the

average of virus-infected untreated controls cells using a non-linear fit model plotting an

agonist vs. normalized dose response curve with variable slope using the least squares

fitting method in Prism 7.0 (GraphPad). All serum and monoclonal antibody samples were

tested in duplicates. Imputation rules for values outside the limits of quantification are

described below (see Statistical methods).

SARS-CoV-2 neutralizing antibody panel and sequence analysis

The panel of 158 SARS-CoV-2 neutralizing monoclonal antibodies isolated from SARS-

CoV-2 convalescent individuals included in the analysis in **Fig. 3** is based on 79 antibodies

obtained in our previous work,[30,31] 67 randomly selected (retrieved on January 1, 2021)

human SARS-CoV-2 neutralizing antibodies deposited at CoV-AbDab,[29] and 12 antibodies

in clinical use or development. We did not include five of the antibodies in clinical development shown in **Fig. 4** into this analysis, as they were obtained from individuals infected with SARS-CoV (ADG-2, sotrovimab), from immunized mice harboring human immunglobulin gene repertoires (47D11, casirivimab), or using phage display technology that does not ensure native pairing of antibody heavy and light chains (regdanvimab).

Antibody amino acid sequences were annotated with IgBLASTp[53] based on the IMGT database.[54] For sequence statistics, top V gene calls were counted without individual alleles, CDR3 lengths are reported according to the IMGT numbering system and numbers of V mutations refer to the top V gene call from IgBLASTp. Phylogenetic analysis of antibodies belonging to the V_H3-53/3-66|V_K1-9 public clonotype was performed by alignment of amino acid sequences with the MAFFT algorithm[55] via the EMBL-EBI search and sequence analysis tools API[56] and the Tree Builder tool from Geneious Prime 2020.0.4 (Biomatters) using the Jukes-Cantor distance model for tree building with the neighbour-joining method without resampling. Data aggregation and visualization was performed with the Python libraries pandas (v1.1.5), NumPy (v1.19.2), SciPy (v1.5.2), Matplotlib (v3.3.4) with Python (v3.6.8), as well as Microsoft Excel 2011 for Mac (v14.7.3), and Adobe Illustrator.

Visualization of SARS-CoV-2 spike amino acid changes

Amino acid changes relative to the Wu01 spike protein were visualized on a cryo-electron microscopy 3D-reconstruction of the SARS-CoV-2 spike protein (PDB ID: 6XR8, ref. [57]) using ChimeraX (v. 1.3).[58,59]

SARS-CoV-2 variant distribution

Numbers of weekly cases as curated by the COVID-19 Data Repository by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University[60] was retrieved from http://ourworldindata.org (accessed on May 24, 2022). GISAID-curated clade and lineage statistics of sequences submitted to the GISIAD database[61-63] were retrieved from GISAID (accessed on May 24, 2022) and frequency of individual variants was plotted as fraction of all submitted sequences per week and variant. For visualization, all BA.1 sublineages except for BA.1.1 (and sublineages of BA.1.1) were classified as BA.1, and all BA.2 sublineages except for BA.2.12.1 were classified as BA.2

Statistical Methods

For graphical representation and statistical evaluation of serum samples in **Fig. 2** and **Extended Data Fig. 1**, samples that did not achieve 50% inhibition at the lowest tested dilution of 10 (lower limit of quantification, LLOQ) were imputed to ½ of the LLOQ (ID_{50}=5) and serum samples with ID_{50} >21,870 (upper limit of quantification) were imputed to ID_{50}=21,871. For graphical representation and statistical analysis of monoclonal neutralizing antibodies in **Fig. 3** and **Extended Data Fig. 2**, IC_{50} values of antibodies with an IC_{50} <0.005 µg/ml (LLOQ) were imputed to ½ LLOQ (IC_{50}=0.0025), and IC_{50} values >10 µg/ml (ULOQ) were imputed to 2x ULOQ (IC_{50}=20 µg/ml).

Testing for statistical significance of differences in serum neutralization titers against different variants/sublinages was performed with the Friedman test with Dunn's multiple comparison using Prism 7.0 (GraphPad). Spearman's rank correlation coefficients (Rho) were determined using Prism 7.0 (GraphPad). Amino acid mutations relative to germline between antibodies neutralizing Wu01 only and those neutralizing

any Omicron sublineage were compared with two-sided Mann-Whitney U tests using

Prism 7.0 (GraphPad).

Data and code availability

Requests for data or materials should be directed to the corresponding author and may

be subject to restrictions based on data and privacy protection regulations and/or may

require a Material Transfer Agreement (MTA). The paper does not report original code.

651
References

1   Andrews, N. *et al.* Covid-19 Vaccine Effectiveness against the Omicron (B.1.1.529)
Variant. *N Engl J Med* **386**, 1532-1546, doi:10.1056/NEJMoa2119451 (2022).
2   Corti, D., Purcell, L. A., Snell, G. & Veesler, D. Tackling COVID-19 with neutralizing
monoclonal antibodies. *Cell* **184**, 3086-3108, doi:10.1016/j.cell.2021.05.005 (2021).
3   Gruell, H. *et al.* Antibody-Mediated Neutralization of SARS-CoV-2. *Immunity*,
doi:10.1016/j.immuni.2022.05.005 (2022).
4   Garcia-Beltran, W. F. *et al.* mRNA-based COVID-19 vaccine boosters induce
neutralizing immunity against SARS-CoV-2 Omicron variant. *Cell* **185**, 457-466 e454,
doi:10.1016/j.cell.2021.12.033 (2022).
5   Carreno, J. M. *et al.* Activity of convalescent and vaccine serum against SARS-CoV-2
Omicron. *Nature* **602**, 682-688, doi:10.1038/s41586-022-04399-5 (2022).
6   Schmidt, F. *et al.* Plasma Neutralization of the SARS-CoV-2 Omicron Variant. *N Engl J*
Med **386**, 599-601, doi:10.1056/NEJMc2119641 (2022).
7   Liu, L. *et al.* Striking antibody evasion manifested by the Omicron variant of SARS-
CoV-2. *Nature* **602**, 676-681, doi:10.1038/s41586-021-04388-0 (2022).
8   Planas, D. *et al.* Considerable escape of SARS-CoV-2 Omicron to antibody
neutralization. *Nature* **602**, 671-675, doi:10.1038/s41586-021-04389-z (2022).
9   Cele, S. *et al.* Omicron extensively but incompletely escapes Pfizer BNT162b2
neutralization. *Nature* **602**, 654-656, doi:10.1038/s41586-021-04387-1 (2022).
10   Hoffmann, M. *et al.* The Omicron variant is highly resistant against antibody-
mediated neutralization: Implications for control of the COVID-19 pandemic. *Cell* **185**,
447-456 e411, doi:10.1016/j.cell.2021.12.032 (2022).
11   Gruell, H. *et al.* mRNA booster immunization elicits potent neutralizing serum activity
against the SARS-CoV-2 Omicron variant. *Nat Med* **28**, 477-480, doi:10.1038/s41591-
021-01676-0 (2022).
12   Tseng, H. F. *et al.* Effectiveness of mRNA-1273 against SARS-CoV-2 Omicron and Delta
variants. *Nat Med*, doi:10.1038/s41591-022-01753-y (2022).
13   Altarawneh, H. N. *et al.* Protection against the Omicron Variant from Previous SARS-
CoV-2 Infection. *N Engl J Med* **386**, 1288-1290, doi:10.1056/NEJMc2200133 (2022).
14   Viana, R. *et al.* Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in
southern Africa. *Nature* **603**, 679-686, doi:10.1038/s41586-022-04411-y (2022).

15 Madhi, S. A. *et al.* Population Immunity and Covid-19 Severity with Omicron Variant
in South Africa. *N Engl J Med* **386**, 1314-1326, doi:10.1056/NEJMoa2119658 (2022).
16 Zhao, X. *et al.* Effects of a Prolonged Booster Interval on Neutralization of Omicron
Variant. *N Engl J Med* **386**, 894-896, doi:10.1056/NEJMc2119426 (2022).
17 Wratil, P. R. *et al.* Three exposures to the spike protein of SARS-CoV-2 by either
infection or vaccination elicit superior neutralizing immunity to all variants of
concern. *Nat Med* **28**, 496-503, doi:10.1038/s41591-022-01715-4 (2022).
18 Perez-Then, E. *et al.* Neutralizing antibodies against the SARS-CoV-2 Delta and
Omicron variants following heterologous CoronaVac plus BNT162b2 booster
vaccination. *Nat Med* **28**, 481-485, doi:10.1038/s41591-022-01705-6 (2022).
19 VanBlargan, L. A. *et al.* An infectious SARS-CoV-2 B.1.1.529 Omicron virus escapes
neutralization by therapeutic monoclonal antibodies. *Nat Med* **28**, 490-495,
doi:10.1038/s41591-021-01678-y (2022).
20 Yamasoba, D. *et al.* Virological characteristics of the SARS-CoV-2 Omicron BA.2 spike.
Cell, doi:10.1016/j.cell.2022.04.035 (2022).
21 Tegally, H. *et al.* Continued Emergence and Evolution of Omicron in South Africa:
New BA.4 and BA.5 lineages. *medRxiv*, doi:10.1101/2022.05.01.22274406 (2022).
22 Greaney, A. J. *et al.* Mapping mutations to the SARS-CoV-2 RBD that escape binding
by different classes of antibodies. *Nat Commun* **12**, 4196, doi:10.1038/s41467-021-
24435-8 (2021).
23 Kimura, I. *et al.* The SARS-CoV-2 Lambda variant exhibits enhanced infectivity and
immune resistance. *Cell Rep* **38**, 110218, doi:10.1016/j.celrep.2021.110218 (2022).
24 Planas, D. *et al.* Reduced sensitivity of SARS-CoV-2 variant Delta to antibody
neutralization. *Nature* **596**, 276-280, doi:10.1038/s41586-021-03777-9 (2021).
25 Baum, A. *et al.* Antibody cocktail to SARS-CoV-2 spike protein prevents rapid
mutational escape seen with individual antibodies. *Science* **369**, 1014-1018,
doi:10.1126/science.abd0831 (2020).
26 Crawford, K. H. D. *et al.* Protocol and Reagents for Pseudotyping Lentiviral Particles
with SARS-CoV-2 Spike Protein for Neutralization Assays. *Viruses* **12**,
doi:10.3390/v12050513 (2020).
27 Vanshylla, K. *et al.* Kinetics and correlates of the neutralizing antibody response to
SARS-CoV-2 infection in humans. *Cell Host Microbe* **29**, 917-929 e914,
doi:10.1016/j.chom.2021.04.015 (2021).
28 Hillus, D. *et al.* Safety, reactogenicity, and immunogenicity of homologous and
heterologous prime-boost immunisation with ChAdOx1 nCoV-19 and BNT162b2: a
prospective cohort study. *The Lancet Respiratory Medicine* **9**, 1255-1265,
doi:10.1016/s2213-2600(21)00357-x (2021).
29 Raybould, M. I. J., Kovaltsuk, A., Marks, C. & Deane, C. M. CoV-AbDab: the
coronavirus antibody database. *Bioinformatics* **37**, 734-735,
doi:10.1093/bioinformatics/btaa739 (2021).
30 Vanshylla, K. *et al.* Discovery of ultrapotent broadly neutralizing antibodies from
SARS-CoV-2 elite neutralizers. *Cell Host Microbe* **30**, 69-82 e10,
doi:10.1016/j.chom.2021.12.010 (2022).
31 Kreer, C. *et al.* Longitudinal Isolation of Potent Near-Germline SARS-CoV-2-
Neutralizing Antibodies from COVID-19 Patients. *Cell* **182**, 843-854 e812,
doi:10.1016/j.cell.2020.06.044 (2020).
32 Robbiani, D. F. *et al.* Convergent antibody responses to SARS-CoV-2 in convalescent
individuals. *Nature* **584**, 437-442, doi:10.1038/s41586-020-2456-9 (2020).

33   Nielsen, S. C. A. *et al.* Human B Cell Clonal Expansion and Convergent Antibody
Responses to SARS-CoV-2. *Cell Host Microbe* **28**, 516-525 e515,
doi:10.1016/j.chom.2020.09.002 (2020).
34   Yuan, M. *et al.* Structural basis of a shared antibody response to SARS-CoV-2. *Science*
369, 1119-1123, doi:10.1126/science.abd2321 (2020).
35   Cao, Y. *et al.* Potent Neutralizing Antibodies against SARS-CoV-2 Identified by High-
Throughput Single-Cell Sequencing of Convalescent Patients' B Cells. *Cell* **182**, 73-84
e16, doi:10.1016/j.cell.2020.05.025 (2020).
36   Zhang, Q. *et al.* Potent and protective IGHV3-53/3-66 public antibodies and their
shared escape mutant on the spike of SARS-CoV-2. *Nat Commun* **12**, 4210,
doi:10.1038/s41467-021-24514-w (2021).
37   Weinreich, D. M. *et al.* REGEN-COV Antibody Combination and Outcomes in
Outpatients with Covid-19. *N Engl J Med* **385**, e81, doi:10.1056/NEJMoa2108163
(2021).
38   Gupta, A. *et al.* Early Treatment for Covid-19 with SARS-CoV-2 Neutralizing Antibody
Sotrovimab. *N Engl J Med* **385**, 1941-1950, doi:10.1056/NEJMoa2107934 (2021).
39   Cohen, M. S. *et al.* Effect of Bamlanivimab vs Placebo on Incidence of COVID-19
Among Residents and Staff of Skilled Nursing and Assisted Living Facilities: A
Randomized Clinical Trial. *JAMA* **326**, 46-55, doi:10.1001/jama.2021.8828 (2021).
40   Group, R. C. Casirivimab and imdevimab in patients admitted to hospital with COVID-
19 (RECOVERY): a randomised, controlled, open-label, platform trial. *The Lancet* **399**,
665-676, doi:10.1016/s0140-6736(22)00163-5 (2022).
41   Levin, M. J. *et al.* Intramuscular AZD7442 (Tixagevimab-Cilgavimab) for Prevention of
Covid-19. *N Engl J Med*, doi:10.1056/NEJMoa2116620 (2022).
42   Liu, C. *et al.* Reduced neutralization of SARS-CoV-2 B.1.617 by vaccine and
convalescent serum. *Cell* **184**, 4220-4236 e4213, doi:10.1016/j.cell.2021.06.020
(2021).
43   Weissman, D. *et al.* D614G Spike Mutation Increases SARS CoV-2 Susceptibility to
Neutralization. *Cell Host Microbe* **29**, 23-31 e24, doi:10.1016/j.chom.2020.11.012
(2021).
44   Khoury, D. S. *et al.* Neutralizing antibody levels are highly predictive of immune
protection from symptomatic SARS-CoV-2 infection. *Nat Med* **27**, 1205-1211,
doi:10.1038/s41591-021-01377-8 (2021).
45   Feng, S. *et al.* Correlates of protection against symptomatic and asymptomatic SARS-
CoV-2 infection. *Nat Med* **27**, 2032-2040, doi:10.1038/s41591-021-01540-1 (2021).
46   Harvey, W. T. *et al.* SARS-CoV-2 variants, spike mutations and immune escape. *Nat*
Rev Microbiol **19**, 409-424, doi:10.1038/s41579-021-00573-0 (2021).
47   Lyngse, F. P. *et al.* Transmission of SARS-CoV-2 Omicron VOC subvariants BA.1 and
BA.2: Evidence from Danish Households. *medRxiv*, doi:10.1101/2022.01.28.22270044
(2022).
48   Bruel, T. *et al.* Serum neutralization of SARS-CoV-2 Omicron sublineages BA.1 and
BA.2 in patients receiving monoclonal antibodies. *Nat Med*, doi:10.1038/s41591-022-
01792-5 (2022).
49   Iketani, S. *et al.* Antibody evasion properties of SARS-CoV-2 Omicron sublineages.
Nature, doi:10.1038/s41586-022-04594-4 (2022).
50   Stothard, P. The sequence manipulation suite: JavaScript programs for analyzing and
formatting protein and DNA sequences. *Biotechniques* **28**, 1102, 1104,
doi:10.2144/00286ir01 (2000).

51 Nakamura, Y., Gojobori, T. & Ikemura, T. Codon usage tabulated from international
DNA sequence databases: status for the year 2000. *Nucleic Acids Res* **28**, 292,
doi:10.1093/nar/28.1.292 (2000).

52 Tiller, T. *et al.* Efficient generation of monoclonal antibodies from single human B
cells by single cell RT-PCR and expression vector cloning. *J Immunol Methods* **329**,
112-124, doi:10.1016/j.jim.2007.09.017 (2008).

53 Ye, J., Ma, N., Madden, T. L. & Ostell, J. M. IgBLAST: an immunoglobulin variable
domain sequence analysis tool. *Nucleic Acids Res* **41**, W34-40,
doi:10.1093/nar/gkt382 (2013).

54 Lefranc, M. P. IMGT, the International ImMunoGeneTics Information System. *Cold*
Spring Harb Protoc **2011**, 595-603, doi:10.1101/pdb.top115 (2011).

55 Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid
multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30**,
3059-3066, doi:10.1093/nar/gkf436 (2002).

56 Madeira, F. *et al.* The EMBL-EBI search and sequence analysis tools APIs in 2019.
Nucleic Acids Res **47**, W636-W641, doi:10.1093/nar/gkz268 (2019).

57 Cai, Y. *et al.* Distinct conformational states of SARS-CoV-2 spike protein. *Science* **369**,
1586-1592, doi:10.1126/science.abd4251 (2020).

58 Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers,
educators, and developers. *Protein Sci* **30**, 70-82, doi:10.1002/pro.3943 (2021).

59 Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization and
analysis. *Protein Sci* **27**, 14-25, doi:10.1002/pro.3235 (2018).

60 Dong, E., Du, H. & Gardner, L. An interactive web-based dashboard to track COVID-19
in real time. *The Lancet Infectious Diseases* **20**, 533-534, doi:10.1016/s1473-
3099(20)30120-1 (2020).

61 Khare, S. *et al.* GISAID's Role in Pandemic Response. *China CDC Wkly* **3**, 1049-1051,
doi:10.46234/ccdcw2021.255 (2021).

62 Elbe, S. & Buckland-Merrett, G. Data, disease and diplomacy: GISAID's innovative
contribution to global health. *Glob Chall* **1**, 33-46, doi:10.1002/gch2.1018 (2017).

63 Shu, Y. & McCauley, J. GISAID: Global initiative on sharing all influenza data - from
vision to reality. *Euro Surveill* **22**, doi:10.2807/1560-7917.ES.2017.22.13.30494
(2017).

Figure 1

Fig. 1. Omicron sublineage differences.
a, Spike amino acid changes in the BA.1, BA.1.1, BA.2, BA.2.12.1, and BA.4/5 Omicron sublineages relative to Wu01. **b,** Locations of changed amino acids in Omicron sublineages highlighted on the SARS-CoV-2 spike (PDB: 6XR8), with colors indicating mutations shared between different sublineages. Representations at the bottom show RBD and NTD outlined in black. Changes exclusive to BA.1.1 are indicated in green and those to BA.4/5 in brown, residue 452 (mutated in BA.2.12.1 and BA.4/5) is indicated in purple. **c,** Top panels indicate weekly reported SARS-CoV-2 infections as aggregated by the COVID-19 Data Repository by the Center for Systems Science and Engineering at Johns Hopkins University and http://ourworldindata.org. Relative proportions of infections by variant per week were extrapolated from the proportion of sequences submitted to the GISAID SARS-CoV-2 database per variant and week (accessed on May 24, 2022) shown in the bottom panels. NTD, N-terminal domain; RBD, receptor-binding domain; PDB, Protein Data Bank.

Figure 2

Fig. 2. Omicron sublineage neutralizing serum activity in vaccinated and convalescent individuals.
a, Study scheme in COVID-19-convalescent individuals. Samples were collected after infection occurring between February and April, 2020 (V1), and after a BNT162b2 booster immunization (V2). **b,** Fifty-percent inhibitory serum dilutions (ID$_{50}$s) against Wu01, BA.1, BA.1.1, BA.2.12.1, and BA.4/5 determined by pseudovirus neutralization assays in convalescent individuals. Bars indicate geometric mean ID$_{50}$s with 95% confidence intervals (CIs) at V1 (left) and V2 (right). Numbers indicate geometric mean ID$_{50}$s and percentage of individuals with detectable neutralizing activity (ID$_{50}$ >10) in parentheses. **c,** Study scheme in vaccinated individuals. Samples were collected after the second dose of BNT162b2 (V1) and after the third dose of BNT162b2 (V2). **d,** Serum ID$_{50}$s against Wu01, BA.1, BA.1.1, BA.2.12.1, and BA.4/5 determined by pseudovirus neutralization assays in vaccinated individuals. Bars indicate geometric mean ID$_{50}$s with 95% CIs at V1 (left) and V2 (right). Numbers indicate geometric mean ID$_{50}$s and percentage of individuals with detectable neutralizing activity (ID$_{50}$ >10) in parentheses. In **b** and **d**, ID$_{50}$s below the lower limit of quantification (LLOQ, ID$_{50}$ of 10; indicated by black dotted lines) were imputed to ½ LLOQ (ID$_{50}$=5), and ID$_{50}$s above the upper limit of quantification (21,870) were imputed to 21,871.

Figure 3

Fig. 3. Determining Omicron sublineage immune escape using monoclonal antibodies.
a, SARS-CoV-2 neutralizing monoclonal antibodies (n=158) derived from 19 studies and ≥43 convalescent individuals were analyzed. Bar charts indicate number of antibodies per heavy chain variable gene segment (V_H), amino acid (aa) length of the heavy chain complementarity-determining region 3 (CDRH3), and number of VH aa mutations relative to the V_H germline gene. Pie chart indicates antibody epitopes with slice sizes proportional to antibody fractions. RBD, receptor-binding domain. **b,** Fractions of antibodies from the panel (n=158) displaying activity (IC50 <10 µg/ml) against Wu01 and Omicron sublineages. **c,** IC50s of neutralizing antibodies against individual variants/sublineages (Wu01, n=158; BA.1, n=29; BA.1.1, n=27; BA.2, n=34; BA.2.12.1, n=36; BA.4/5, n=29). Solid lines depict geometric mean IC50s and 95% confidence intervals, and dashed lower and upper lines indicate lower (LLOQ, 0.005 µg/ml) and upper limit of quantification (ULOQ, 10 µg/ml), respectively. IC50s <LLOQ were imputed to ½ LLOQ (IC50=0.0025). **d,** Heat map depicting IC50s of the subset of antibodies (n=43) with neutralizing activity (IC50 <10 µg/ml) against at least one of the Omicron sublineages. Antibodies are sorted based on their potency against BA.1. **e,** Log10 IC50 ratios of antibodies neutralizing any Omicron sublineage for indicated sublineage comparisons. Within each panel, antibodies are sorted by increasing IC50 ratios. Ratios <-1 log10 are highlighted in green, ratios between -1 and 1 log10 in grey, and ratios >1 log10 in red. Numbers indicate relative fractions. IC50 <LLOQ were imputed to ½ LLOQ (IC50=0.0025) and IC50 values >ULOQ were imputed to 2x ULOQ (IC50=20). **f,** Spearman's rank correlation coefficients (rho) for comparisons of antibodies with neutralizing capacity against any Omicron sublineage (n=43) as shown in Fig. S2c. Numbers in bold indicate values with p<0.05. **g,** Heavy chain V amino acid mutations of antibodies with neutralizing activity against Wu01 only and antibodies neutralizing both Wu01 and at least one omicron sublineage. Lines indicate medians and interquartile ranges. Groups were compared using the two-tailed Mann-Whitney U test. **h,** Phylogenetic tree and heavy chain sequence alignment of antibodies of the V_H3-53/3-66|V_K1-9 public clonotype. Letters indicate aa mutations relative to the V_H germline gene. Number of aa mutations from corresponding germline allele and neutralizing activity against Omicron sublineages are indicated on the right. Germline V_H represents the consensus of all identified germline alleles of depicted antibodies (V_H3-53*01, V_H3-53*04, and V_H3-66*01). Number of aa mutations from corresponding germline allele and neutralizing activity against Wu01 and Omicron sublineages indicated on the right.

Figure 4

a

Antibody		IC50 (µg/ml)					
Name		Wu01	BA.1	BA.1.1	BA.2	BA.2.12.1	BA.4/5
Bamlanivimab*		<0.005	>10	>10	>10	>10	>10
Etesevimab*		0.018	>10	>10	>10	>10	>10
Bebtelovimab†		<0.005	<0.005	<0.005	<0.005	<0.005	<0.005
Imdevimab*		<0.005	>10	>10	4.173	4.624	3.392
Casirivimab*		<0.005	>10	>10	>10	>10	>10
Sotrovimab*		0.976	0.105	0.127	1.488	3.949	3.671
Tixagevimab†		<0.005	>10	>10	>10	3.438	>10
Cilgavimab†		<0.005	6.848	>10	0.008	0.014	0.085
Regdanvimab†		<0.005	>10	>10	>10	>10	>10
DZIF-10c*		<0.005	0.046	>10	0.033	0.034	8.638
BRII-196†		0.008	>10	>10	>10	5.398	0.820
C135†		<0.005	>10	>10	>10	>10	>10
C144†		<0.005	>10	>10	>10	>10	>10
COV2-2381†		0.008	>10	>10	>10	>10	>10
ADG-2†		<0.005	0.212	0.194	>10	>10	>10
47D11†		0.386	>10	>10	>10	>10	>10
BD-368-2†		<0.005	>10	>10	>10	>10	>10
MAD0004J08†		<0.005	>10	>10	>10	>10	>10

Monoclonal antibody used for neutralization assay:

* Clinical product used.

† Parental antibody (Fc-unmodified IgG1) used.

d

Antibody	IC50 (µg/ml)					
Name	Wu01	BA.1	BA.1.1	BA.2	BA.2.12.1	BA.4/5
R207-2F11	<0.005	<0.005	0.006	0.006	<0.005	0.011
R568-1G9	0.007	0.006	0.006	0.005	<0.005	0.045
R200-1F9	<0.005	0.023	0.025	0.026	0.017	0.017
	B.1	Alpha	Beta	Delta	Epsilon	Kappa
R207-2F11	0.003	0.004	0.013	0.004	0.002	0.011
R568-1G9	0.002	0.003	0.001	0.002	0.002	0.006
R200-1F9	0.001	0.002	0.002	0.004	0.003	0.003

Fig. 4. Omicron sublineage neutralizing activity of monoclonal antibodies in clinical use.
a, Dose response curves showing percentage neutralization of monoclonal antibodies against Wu01 and Omicron sublineages in pseudovirus neutralization assay. Circles show averages and error bars indicate standard deviation. Dotted lines indicate 50% neutralization (IC50). **b,** IC50s of monoclonal antibodies with current or previous authorization for clinical use or in clinical development. Symbols indicate whether clinical products or parental antibodies produced as human IgG1 were used. **c,** Dose response curves showing percentage neutralization of novel monoclonal antibodies against Wu01 and Omicron sublineages in pseudovirus neutralization assay. Circles show averages and error bars indicate standard deviation. Dotted lines indicate 50% neutralization (IC50). **d,** IC50s of novel monoclonal antibodies shown in **c** against Wu01 and Omicron sublineages (upper rows) and previously circulating variants (lower rows, as previously determined[30]).