1

Conduction velocity along a key white matter tract is associated with autobiographical memory recall ability

Ian A. Clark[1], Siawoosh Mohammadi[2], Martina F. Callaghan[1], Eleanor A. Maguire[1]*

[1]Wellcome Centre for Human Neuroimaging, Department of Imaging Neuroscience, UCL Queen Square Institute of Neurology, University College London, London, UK

[2]Institute of Systems Neuroscience, University Medical Centre Hamburg-Eppendorf, Hamburg, Germany

*For correspondence: e.maguire@ucl.ac.uk

Competing interests: The authors declare no competing interests.

Abstract

Conduction velocity is the speed at which electrical signals travel along axons and is a crucial determinant of neural communication. Inferences about conduction velocity can now be made in vivo in humans using a measure called the magnetic resonance (MR) g-ratio. This is the ratio of the inner axon diameter relative to that of the axon plus the myelin sheath that encases it. Here, in the first application to cognition, we found that variations in MR g-ratio, and by inference conduction velocity, of the parahippocampal cingulum bundle were associated with autobiographical memory recall ability in 217 healthy adults. This tract connects the hippocampus with a range of other brain areas. We further observed that the association seemed to be with inner axon diameter rather than myelin content. The extent to which neurites were coherently organised within the parahippocampal cingulum bundle was also linked with autobiographical memory recall ability. Moreover, these findings were specific to autobiographical memory recall, and were not apparent for laboratory-based memory tests. Our results offer a new perspective on individual differences in autobiographical memory recall ability, highlighting the possible influence of specific features of white matter microstructure when recalling detailed memories of our real-life past experiences.

Introduction

Communication between neurons in the brain is critical for cognition, and depends upon action potentials being conveyed along axons within white matter tracts. The speed at which these electrical signals travel along axons is known as the conduction velocity. It has been suggested that faster axonal conduction velocity promotes better cognition. For example, an increase in axonal conduction velocity is hypothesised to underpin the greater cognitive processing ability of vertebrates, in particular primates and humans (Brancucci, 2012; Miller, 1994), compared to invertebrates (Arancibia-Cárcamo et al., 2017; Bullock et al., 1984; Nave, 2010). In rats (Aston-Jones et al., 1985) and cats (Xi et al., 1999), faster axonal conduction velocity has been observed in younger compared to older animals. In a similar vein, axonal degradation that can lead to reduced conduction velocity has been identified in older compared to younger monkeys (Peters et al., 2000; Peters and Sethares, 2002).

Echoing these findings from non-humans, estimates of conduction velocity in humans from the latency of visual evoked potentials recorded over primary visual cortex (Reed and Jensen, 1992) and between the thalamus and parietal cortex (Reed and Jensen, 1993) have been positively correlated with nonverbal intelligence quotients. In addition, faster axonal conduction velocities are thought to better explain increases in intelligence compared to absolute and relative brain volumes (Dicke and Roth, 2016).

The conduction velocity of an axon is dependent upon the axon diameter, the presence and thickness of a myelin sheath, the distance between the nodes of Ranvier (periodic gaps in the myelin that facilitate action potential propagation), inter-nodal spacing, and electrical properties of the axonal and myelin membranes (Arancibia-Cárcamo et al., 2017; Drakesmith et al., 2019; Gasser and Grundfest, 1939; Hursh, 1939; Huxley and Stämpeli, 1949; Rushton, 1951). A number of these features are not yet measurable in humans in vivo. However, seminal

electrophysiological work has derived a relationship between axon morphology and conduction

velocity using only axon diameter and myelin sheath thickness (Rushton, 1951). These two

metrics are particularly key because a larger axon diameter results in less resistance to the

action potential ion flow, resulting in faster conduction velocity. The presence of a myelin

sheath around an axon is beneficial in two ways. First, the myelin sheath acts like an electrical

insulating layer, reducing ion loss and preserving the action potential. Second, the presence of

unmyelinated gaps in the myelin sheath (the nodes of Ranvier) enables a process called

saltatory propagation to take place. As the majority of the axon is wrapped in myelin, the nodes

of Ranvier are the only locations where action potentials can occur. This increases the strength

of electrical signals because all the ions gather at these nodes instead of being dispersed along

the length of the axon. Stronger action potentials are therefore sent along the myelinated portion

of the axon at higher speeds, with this signal being boosted on arrival at the next node of

Ranvier by another action potential, which helps to maintain a fast conduction velocity.

The conduction velocity of an axon is, therefore, not determined only by the axon

diameter, but also by the relationship between the axon diameter and the thickness of the

surrounding myelin sheath, a measure known as the g-ratio (Chomiak and Hu, 2009; Rushton,

1951; Schmidt and Knösche, 2019). Specifically, Rushton (1951) derived an equation:

conduction velocity $\propto d\sqrt{-\ln(g)}$ where d is the inner axon diameter, g is the g-ratio = d/D,

and D is the outer fibre (axon plus myelin sheath) diameter. In other words, the g-ratio is

computed as the ratio of the inner axon diameter relative to that of the axon plus the myelin

sheath that encases it (Figure 1).

Figure 1. Schematic of a myelinated axon showing how the g-ratio is calculated.

Until recently, g-ratio measurements were restricted to invasive studies in non-human

animals. However, by combining diffusion magnetic resonance imaging (MRI) with

quantitative structural MRI scans optimised to assess myelination (e.g. magnetisation transfer

saturation; Weiskopf et al., 2013), it is now possible to estimate the g-ratio in vivo in humans

across the whole brain (Drakesmith et al., 2019; Mohammadi et al., 2015; Mohammadi and

Callaghan, 2021; Stikov et al., 2015). This is achieved by measuring an aggregate g-ratio,

which is an area-weighted ensemble average across a voxel of an underlying distribution of

microscopic g-ratios of axons (Stikov et al., 2015; West et al., 2016). Whole brain MR g-ratio

maps enable the investigation of the MR g-ratio of white matter fibre pathways at the group

level. These MR g-ratio estimates have been optimised (Ellerbrock and Mohammadi, 2018;

Jung et al., 2018; West et al., 2018) and used to investigate white matter development

(Cercignani et al., 2017), changes in the g-ratio during aging (Berman et al., 2018), and as a

potential neuroimaging marker in patients with multiple sclerosis (Yu et al., 2019). Of note,

the MR g-ratio has been found to associate well with estimates of axonal conduction velocity

(Berman et al., 2019; Drakesmith et al., 2019). Consequently, the MR g-ratio provides a non-

invasive MRI method that can associate in vivo structural neuroimaging of humans with axonal

conduction velocity.

Given that the MR g-ratio encompasses information about the inner axon diameter and

myelin thickness, the identification of statistically significant relationships with the MR g-ratio

can also guide as to which of these two features might be influencing variations in conduction

velocity (Caeyenberghs et al., 2016; Kaller et al., 2017; Lakhani et al., 2016; Waxman, 1980;

Xin and Chan, 2020). This is particularly useful in relation to the inner axon diameter. While

measures of myelination can be obtained using conventional MRI scanners (e.g. via

magnetisation transfer saturation maps), estimates of inner axon diameter cannot, and instead

require ultra-strong gradient systems – 300 mT/m compared to the 10s of mT/m of

conventional MRI scanners (Jones et al., 2018; Veraart et al., 2020). However, only four ultra-

strong gradient Connectom MRI scanners exist in the world. Consequently, measuring the MR

g-ratio with conventional MRI scanners can be useful in guiding inferences about inner axon

diameter.

Adjudicating between the possible influence of inner axon diameter or myelination

relies on knowledge, or an assumption, about whether the associated change in conduction

velocity is faster or slower. This is because significant associations with the MR g-ratio only

indicate the existence of a relationship with conduction velocity but not the direction. As noted

previously, faster conduction velocity is often held to promote better cognition (e.g. Brancucci,

2012; Dicke and Roth, 2016; Miller, 1994; Reed and Jensen, 1992). Therefore, in Figure 2A,

inner axon diameter, myelin thickness and (MR) g-ratio are plotted together to illustrate how

different changes in the MR g-ratio are related to the underlying microstructural properties,

given a faster conduction velocity. Myelin thickness is represented by the gradient in

background colour and contours, with thinnest myelin at the bottom right, and thickest on the

top left. The direction of the arrows describes the change in g-ratio for the microstructural

variations presented. There are three main scenarios. (1) A decrease in MR g-ratio values. This

would suggest that faster conduction velocity is due to greater thickness of the myelin sheath,

with the inner axon diameter remaining constant (Figure 2A blue arrow, and Figure 2B). (2) A

decrease in MR g-ratio values, but to a lesser extent than that observed in the first scenario.

This would suggest that faster conduction velocity is due primarily to greater myelin sheath

thickness, but one that is also accompanied by a larger inner axon diameter (Figure 2A red

arrow, and Figure 2C). (3) An increase in MR g-ratio values. This would suggest that faster

conduction velocity is predominantly due to a larger inner axon diameter, with only small

differences, if any, in myelin thickness being present (Figure 2A orange arrow, and Figure 2D).

145   We note, for completeness, that a fourth scenario also exists where constant MR g-ratio

values could be associated with faster conduction velocity. This would occur when both the

inner axon diameter and myelin thickness change proportionally to each other (Figure 2A black

arrow, and Figure 2E). However, this scenario could also mean that there is no variation in

conduction velocity. One way to increase interpretability in this situation, is by examining

scans optimised to assess myelination (such as magnetisation transfer saturation values).

Observing no relationship with the MR g-ratio, but a relationship with magnetisation transfer

saturation values, would suggest a proportional change in the underlying myelin and,

consequently, variation in conduction velocity. By contrast, no relationship with either the MR

g-ratio or magnetisation transfer saturation values would suggest that there was no change in

the underlying microstructure and therefore no variation in conduction velocity.

Figure 2. Illustration of how the MR g-ratio relates to specific microstructural properties given a faster conduction velocity. (**A**) Graphical representation of the relationships between myelin thickness, inner axon diameter and MR g-ratio, assuming a faster conduction velocity (see Appendix 1 for details of the simulation). Myelin thickness is represented by the gradient in background colour and contours on the graph, with the thinnest myelin at the bottom right and thickest at the top left. The direction of the arrows describes the change in g-ratio for each microstructural variation presented in B-E. The positioning and colours of the arrows correspond to the text box outline colours in B-E. (**B-E**) Illustrations of how changes in the MR g-ratio relate to the underlying axonal microstructure, given a faster conduction velocity.

The MR g-ratio has, therefore, the potential to provide a number of novel insights into

human cognition. One area where the MR g-ratio may be particularly helpful is in probing

individual differences. Our particular interest is in the ability to recall past experiences from

real life, known as autobiographical memories. The detailed recall of autobiographical

memories is a critical cognitive function that serves to sustain our sense of self, enable

independent living, and prolong survival (Tulving, 2002). While some healthy individuals can

recollect decades-old autobiographical memories with great richness and clarity, others

struggle to recall what they did last weekend (LePort et al., 2012; Palombo et al., 2015). In the

context of the healthy population, we currently lack a clear biological explanation for the basis

of these individual differences (Palombo et al., 2018).

There is no doubt that the hippocampus is central to the processing of autobiographical

memories, and hippocampal damage is linked with autobiographical memory impairments

(McCormick et al., 2018; Scoville and Milner, 1957; Winocur and Moscovitch, 2011).

However, no consistent relationship between autobiographical memory recall ability and

hippocampal grey matter volume or microstructure has been identified in healthy individuals

(Clark et al., 2020; Clark et al., 2021a; LePort et al., 2012; Maguire et al., 2003; Van Petten,

2004). The hippocampus does not act alone, and functional neuroimaging studies have revealed

that a distributed set of brain areas supports autobiographical memory recall along with the

hippocampus, including the parahippocampal, retrosplenial, parietal and medial prefrontal

cortices (Andrews-Hanna et al., 2014; Maguire, 2001; Spreng et al., 2009; Svoboda et al.,

2006). Changes in MR g-ratio and, by inference, conduction velocity, might affect

communication between these brain regions and so influence individual differences in

autobiographical memory recall within the healthy population.

Three white matter pathways in particular enable communication with the hippocampal

region – the fornix, the uncinate fasciculus and the parahippocampal cingulum bundle. The

fornix (Figure 3A) is a major pathway in and out of the hippocampus and connects it to the

orbital and medial prefrontal cortices, the basal forebrain, the anterior thalamus, the

hypothalamus and the mammillary bodies (Aggleton et al., 2015; Croxson et al., 2005). The

uncinate fasciculus (Figure 3B) originates in the uncus, entorhinal and parahippocampal

cortices and passes over the lateral nucleus of the amygdala, arcs around the Sylvian fissure,

terminating in various locations throughout the prefrontal cortex (Croxson et al., 2005; Von

Der Heide et al., 2013). The parahippocampal cingulum bundle (Figure 3C) links the

hippocampus with the entorhinal, parahippocampal, retrosplenial and parietal cortices, as well

as providing another route between the hippocampus and anterior thalamus (Bubb et al., 2018;

Jones et al., 2013b). It also links to the prefrontal cortex via its connections with other parts of

the cingulum bundle.

Fornix

Uncinate Fasciculus

Parahippocampal Cingulum Bundle

Figure 3. The three white matter tracts of interest, given their relationship with the hippocampal region. The fornix was defined using the ICBM-DTI-81 white-matter labels atlas (Mori et al., 2008). The uncinate fasciculus and parahippocampal cingulum bundle were defined using the Johns Hopkins probabilistic white matter tractography atlas (Hua et al., 2008), with the minimum probability threshold set to 25%.

In the current study we calculated the MR g-ratio within these three pathways to ascertain whether this was related to autobiographical memory recall ability. A significant relationship with the MR g-ratio would suggest that variation in the conduction velocity of a pathway is associated with autobiographical memory recall ability. The relationship of MR g-ratio to the underlying microstructure as outlined in Figure 2, would further guide us as to whether any significant effects were more likely to be associated with the extent of myelination

or the size of the inner axon diameter of the fibres in these three white matter tracts. These

analyses were augmented by examining whether magnetisation transfer saturation values of the

pathways (assessing myelination) were associated with autobiographical memory recall ability.

In addition, we also used the neurite orientation dispersion and density imaging

(NODDI; Zhang et al., 2012) biophysical model to derive two complementary biological

measures that could provide further insights into the arrangement of neurites in a voxel. A

neurite is any projection from a neuron's cell body, such as an axon or a dendrite. The neurite

orientation dispersion index is an estimate of the organisation of the neurites in a voxel, where

a small orientation dispersion index value indicates a low dispersion of neurites, in other words,

that the neurites are coherently organised. The second property, neurite density, is a measure

of the density of the neurites in a voxel.

For completeness, the commonly reported physical parameters from standard diffusion

tensor imaging (DTI; e.g. fractional anisotropy and mean diffusivity; Basser, 1995) that are

often derived from diffusion data were also computed (Oeschger et al., 2021). However, these

metrics lack biological specificity (Jensen and Helpern, 2010; Jones et al., 2013a) and,

consequently, could not speak to our research questions. For example, fractional anisotropy is

a very general measure. Variations in fractional anisotropy values can occur for numerous

reasons, including, but not limited to, changes in the extent of myelination, axon coherence,

axon density, and the level of astrocytes. Consequently, standard DTI parameter results are

only briefly summarised in the main text, with full details provided in Appendix 1 and

Supplementary files 1 and 2.

To ensure an appropriate sample size and a wide range of autobiographical memory

recall ability, we examined a large group of healthy young adults from the general population

($n = 217$; 109 female, 108 male; mean age of 29.0 years, $SD = 5.60$; age was restricted to

between 20-41 years to limit the possible effects of aging).

All participants underwent the widely-used Autobiographical Interview (Levine et al.,

2002). This task, which is the gold standard in the field, provides a detailed metric

characterising a person's ability to recall real-life past experiences, as well as providing a useful

control measure. We also examined performance on another eight standard laboratory-based

memory tests. Their inclusion allowed us to ascertain whether any relationships with

conduction velocity were specific to recollecting detailed autobiographical memories from real

life, or were also applicable to the recall of more constrained laboratory-based stimuli.

Diffusion and magnetisation transfer saturation MRI scans were obtained for each

person to enable calculation of the MR g-ratio and the other measures. Our analyses were

performed using weighted means from each of the three white matter tracts of interest rather

than voxel-wise across the whole brain, reducing the potential for false positives (Marek et al.,

2022).

Focusing on the fornix, uncinate fasciculus and parahippocampal cingulum bundle, we

predicted that variations in the MR g-ratio from some or all of these tracts would be associated

with autobiographical memory recall ability. Such a finding would, for the first time, suggest

a link between variations in white matter tract conduction velocity and individual differences

in autobiographical memory recall.

Under the assumption that a significant relationship between the MR g-ratio and

autobiographical memory recall represents an association with faster conduction velocity (e.g.

Brancucci, 2012; Dicke and Roth, 2016; Miller, 1994; Reed and Jensen, 1992), and aided by

the analysis of the magnetisation transfer saturation values, we further sought to evaluate the

scenarios presented in Figure 2. Specifically, we asked whether autobiographical memory

recall ability was more likely to be associated with inner axon diameters or myelin thickness.

A negative relationship between the MR g-ratio and autobiographical memory recall ability

(Figure 2A blue and red arrows, and Figures 2B and 2C), along with a positive relationship

between magnetisation transfer saturation and autobiographical memory recall ability, would

suggest that myelin thickness was more relevant. By contrast, a positive relationship between

the MR g-ratio and autobiographical memory recall ability (Figure 2A orange arrow, and

Figure 2D) would highlight the potential relevance of inner axon diameters. Observing no

associations between the MR g-ratio or magnetisation transfer saturation values and

autobiographical memory recall, would suggest no relationships with the underlying

microstructure or conduction velocity, and would speak against the possibility of a proportional

change in microstructure (Figure 2A black arrow, and Figure 2E).

Finally, given that a previous functional MRI meta-analysis identified different neural

substrates associated with autobiographical memory recall and the recall of laboratory-based

stimuli (McDermott et al., 2009; see also Maguire, 2001, 2012; Miller et al., 2022; Mobbs et

al., 2021; Nastase et al., 2020; Spiers and Maguire, 2007), we expected that associations with

the MR g-ratio might be specific to autobiographical memory recall. We reasoned that detailed,

multimodal, autobiographical memories may rely on inter-regional connectivity to a greater

degree than simpler, more constrained laboratory-based memory tests.

Results

Autobiographical memory recall scores

We employed the widely-used Autobiographical Interview (Levine et al., 2002) to score

autobiographical memory recall (see Materials and Methods for full details). The main measure

of autobiographical memory recall ability was the mean number of "internal" details from the

freely recalled autobiographical memories. Internal details are those that describe the specific

past event in question, and are considered to reflect episodic information. Across the

participants, the mean number of internal details provided per memory was 23.95 ($SD = 7.25$;

range = 4.60–44.60).

As a control measure, the mean number of "external" details was also calculated from

the autobiographical memory descriptions. External details pertain to semantic information

about the past event, and other non-event information. Across the participants, the mean

number of external details provided per memory was 5.35 ($SD = 3.20$; range = 0.8–17.40).

Laboratory-based memory test performance

While our main interest was in autobiographical memory recall, eight commonly used

laboratory-based memory tasks were also administered. Their inclusion allowed us to establish

whether any associations identified with the main microstructure measure, MR g-ratio, were

specific or not to the recollection of real-life autobiographical memories.

The ability to recall a short narrative was examined using the immediate and delayed

recall tests of the Logical Memory subtest of the Wechsler Memory Scale IV (Wechsler, 2009).

Across participants, the mean immediate recall scaled score was 12.95 ($SD = 2.09$, range = 6-

18) and the mean delayed recall scaled score was 12.58 ($SD = 2.62$, range = 6-19). Verbal list

recall ability was assessed using the immediate and delayed recall of the Rey Auditory Verbal

Learning Test (see Strauss et al., 2006). The mean immediate recall (aggregate) score was

58.82 ($SD = 7.42$, range = 33-73) and the mean delayed recall score was 12.92 ($SD = 2.17$,

range = 6-15). Visuospatial recall ability was examined using the delayed recall of the Rey–

Osterrieth Complex Figure (Rey, 1941), with a mean delayed recall score of 22.28 ($SD = 5.71$,

range = 8.5-35). Recognition memory ability was tested using the Warrington Recognition

Memory Tests for Words and Faces (Warrington, 1984). The mean recognition memory scaled

score for words was 12.75 ($SD = 2.06$, range = 3-15) and for faces was 11.00 ($SD = 3.33$, range

= 3-18). Finally, participants also completed the "Dead or Alive" task which probes general

knowledge about whether famous people have died or are still alive, providing a measure of

semantic memory (Kapur et al., 1989). The mean accuracy performance on this test was

81.32% ($SD = 8.44$, range = 57.14%-97.26%).

No relationships between memory measures and fornix or uncinate fasciculus
microstructure

We first investigated the fornix and uncinate fasciculus. None of the biophysical measures from

either tract were significantly associated with autobiographical memory recall ability. This was

the case when using a corrected $p < 0.017$ (see Materials and Methods) or an uncorrected $p <$

0.05 threshold. Full details of these results are provided in Appendix 1 (Appendix 1–figures 1

and 2; Appendix 1–tables 1-4), with the source data available in Supplementary file 1.

Standard DTI parameters (e.g. fractional anisotropy and mean diffusivity) were also

extracted from the fornix and uncinate fasciculus. None of the standard DTI parameters, from

either tract, were significantly associated with autobiographical memory recall ability, even

when using an uncorrected $p < 0.05$ threshold (see Appendix 1–tables 1-4 for full details, source

data are available in Supplementary file 1).

There were no significant associations between fornix or uncinate fasciculus MR g-

ratios or magnetisation transfer saturation values and any of the laboratory-based memory tests

(see Appendix 1– tables 5 and 6 for full details, source data are available in Supplementary file

1).

Of note, and for completeness, we also performed exploratory analyses in six additional

white matter tracts: the anterior thalamic radiation, the dorsal cingulum bundle, the forceps

minor, the inferior longitudinal fasciculus, the inferior occipitofrontal fasciculus and the

superior longitudinal fasciculus. However, as with the fornix and the uncinate fasciculus, none

of the metrics from any of these tracts were associated with autobiographical memory recall

ability, even when using an uncorrected $p < 0.05$ threshold (see Appendix 1– figures 3-8 and

Appendix 1– tables 7-18 for full details).

The parahippocampal cingulum bundle

We found that variations in autobiographical memory recall ability were uniquely related to

the microstructure of the parahippocampal cingulum bundle. This tract connects the

hippocampus with the entorhinal, parahippocampal, retrosplenial and parietal cortices, and the

anterior thalamus (Figure 4). Moreover, via other subdivisions of the cingulum bundle, it is

indirectly connected with prefrontal regions including the medial prefrontal cortex. The

parahippocampal cingulum bundle is, therefore, well positioned for information transfer

between the key regions involved in autobiographical memory recall (Andrews-Hanna et al.,

2014; Maguire, 2001; Spreng et al., 2009; Svoboda et al., 2006).

354
355   **Figure 4.** Simplified schematic of the location and main connections of the parahippocampal
356   cingulum bundle. The blue lines indicate direct connections, and the dashed blue line an
357   indirect connection.
358

As with the other tracts (see Materials and Methods), the parahippocampal cingulum

bundle region of interest (ROI) was defined bilaterally using the Johns Hopkins probabilistic

white matter tractography atlas (Hua et al., 2008). To reduce partial volume effects, we used a

conservative minimum probability of 25%, and the tract ROI was refined for each participant

to ensure the mask was limited to each person's white matter. The mean number of voxels in

the parahippocampal cingulum bundle ROI was 129.11 (*SD* = 25.68), and the variance in

number of voxels across individuals was accounted for in our analyses. As before, a corrected

statistical threshold of $p < 0.017$ was applied (see Materials and Methods). Table 1 shows the

summary statistics for the microstructure measures, with the source data for the

parahippocampal cingulum bundle available in Supplementary file 2.

Table 1. Means and standard deviations for the microstructure measures from the
parahippocampal cingulum bundle.
372

Microstructure Measure	Mean	Standard Deviation
MR g-ratio	0.647	0.043
Magnetisation transfer saturation	0.959	0.007
Neurite dispersion (ODI)	0.189	0.038
Neurite density	0.480	0.051

Note. ODI = Orientation Dispersion Index

Variation in MR g-ratio of the parahippocampal cingulum bundle was associated with

autobiographical memory recall ability

We first investigated whether the MR g-ratio of the parahippocampal cingulum bundle was

associated with autobiographical memory recall ability, performing partial correlation analyses

with age, gender, scanner and the number of voxels in the ROI included as covariates. A

significant positive association was observed between the parahippocampal cingulum bundle

MR g-ratio and the number of internal details (Figure 5A; $r(211) = 0.18$, $p = 0.008$, 95% CI =

0.05, 0.29). This relationship was specific to internal details, with no association evident for

the external details control measure (Figure 5B; $r(211) = -0.09$, $p = 0.17$, 95% CI = -0.21,

0.019). Direct comparison of the correlations confirmed there was a significantly larger

correlation between the MR g-ratio and internal details than for external details (Figure 5C;

mean r difference = 0.28 (95% CI = 0.11, 0.45), $z = 3.21$, $p = 0.0013$). This suggests that

variations in the conduction velocity of the parahippocampal cingulum bundle are associated

with individual differences in autobiographical memory recall ability. The positive nature of

the correlation highlights the potential relevance of inner axon diameter rather than myelin

thickness (Figure 2A orange arrow, and Figure 2D).

392
Figure 5. MR g-ratio and the parahippocampal cingulum bundle. The relationship between
parahippocampal cingulum bundle MR g-ratio and autobiographical memory recall ability
(internal details), and the control measure (external details) are shown. **(A)** There was a
significant positive correlation between the MR g-ratio and internal details (dashed lines
indicate the confidence intervals). **(B)** There was no significant relationship between the MR
g-ratio and external details. **(C)** Bar chart showing the partial correlation coefficients (with
standard errors) between the MR g-ratio and internal and external details. There was a
significant difference between the correlations when they were directly compared; ***$p <$
0.001. Data points for this figure are provided in Figure 5-source data 1.
402

To further aid the interpretation of the relationship with MR g-ratio in terms of the

underlying axonal microstructure, we also investigated whether the magnetisation transfer

saturation values (assessing myelination) of the parahippocampal cingulum bundle were

associated with autobiographical memory recall ability. We performed partial correlation

analyses with the same covariates as before. Magnetisation transfer saturation values were not

significantly related to either internal ($r(211) = 0.00$, $p = 1.0$, 95% CI = -0.12, 0.12), or external

($r(211) = 0.14$, $p = 0.048$, 95% CI = -0.06, 0.30) details.

Overall, therefore, under the assumption that the significant relationship between the

MR g-ratio and autobiographical memory recall ability is associated with faster conduction

velocity, Rushton's model of conduction velocity would indicate that better autobiographical

memory recall is more likely to be related to parahippocampal cingulum bundle axons having

larger inner axon diameters rather than thicker myelin sheaths (see Figure 2A orange arrow,

and Figure 2D).

Variation in neurite dispersion within the parahippocampal cingulum bundle was

associated with autobiographical memory recall

In addition to the MR g-ratio, we also examined the relationship between autobiographical

memory recall ability and two complementary biophysical measures, the neurite orientation

dispersion index and neurite density maps estimated using the NODDI biophysical model

(Zhang et al., 2012). Partial correlations revealed a significant negative correlation between the

neurite orientation dispersion index (a small orientation dispersion index value indicates low

dispersion) and internal details (Figure 6A; $r(211) = -0.19$, $p = 0.005$, 95% CI = -0.32, -0.06).

This was again specific to internal details with no significant relationship between the neurite

orientation dispersion index and external details (Figure 6B; $r(211) = 0.07$, $p = 0.28$, 95% CI

= -0.05, 0.20). Direct comparison of the correlations revealed a significantly larger correlation

between the neurite orientation dispersion index and internal details than for external details

(Figure 6C; mean r difference = 0.27 (95% CI = 0.44, 0.10), z = 3.13, p = 0.0017). Neurite

density was not significantly related to either internal (r(211) = 0.04, p = 0.60, 95% CI = -0.09,

0.16), or external (r(211) = 0.01, p = 0.93, 95% CI = -0.12, 0.13) details. Therefore, in addition

to a higher MR g-ratio, when neurites in the parahippocampal cingulum bundle were less

dispersed and thus more coherently organised, this was associated with better autobiographical

memory recall.

436
Figure 6. Neurite dispersion and the parahippocampal cingulum bundle. The relationship
between parahippocampal cingulum bundle neurite dispersion (orientation dispersion index)
and autobiographical memory recall ability (internal details), and the control measure (external
details) are shown. (A) There was a significant negative correlation between neurite dispersion
and internal details (dashed lines indicate the confidence intervals). (B) There was no
significant relationship between neurite dispersion and external details. (C) Bar chart showing
the partial correlation coefficients (with standard errors) between neurite dispersion and
internal and external details. There was a significant difference between the correlations when
they were directly compared; **$p < 0.01$. Data points for this figure are provided in the Figure
6-source data 1.
447

Standard DTI parameters of the parahippocampal cingulum bundle and autobiographical memory recall ability

Partial correlations using the same covariates as the previous analyses were also performed using the standard DTI parameters. As these metrics lack biological specificity (Jensen and Helpern, 2010; Jones et al., 2013a), and could not speak to our research questions, the results are summarised here, with full details in Appendix 1–tables 19 and 20 and Appendix 1–figures 9 and 10, and the source data are available in Supplementary file 2.

Two significant relationships were identified between the standard DTI parameters and autobiographical memory recall ability. First, there was a positive correlation between fractional anisotropy and the number of internal details (Appendix 1–figure 9A; $r(211) = 0.20$, $p = 0.003$, 95% CI = 0.07, 0.32). Second, a positive correlation was evident between diffusivities parallel and the number of internal details (Appendix 1–figure 10A; $r(211) = 0.19$, $p = 0.005$, 95% CI = 0.06, 0.32). As with the microstructure metrics, these relationships were specific to internal details with no relationships identified with external details for either fractional anisotropy ($r(211) = -0.06$, $p = 0.39$, 95% CI = -0.19, 0.07) or diffusivities parallel ($r(211) = -0.048$, $p = 0.49$, 95% CI = -0.19, 0.10). No significant relationships were found between mean diffusivity, mean kurtosis or diffusivities perpendicular and either internal or external details.

No relationship between microstructure measures of the parahippocampal cingulum bundle and scores on laboratory-based memory tests

Finally, we tested for associations between the key microstructure measures and performance on the eight laboratory-based memory tasks. No relationships with the MR g-ratio or magnetisation transfer saturation values were evident for any task, even when using an uncorrected $p < 0.05$ threshold (Appendix 1-table 21, source data are available in

Supplementary file 2). This suggests that the parahippocampal cingulum bundle MR g-ratio,

and by inference conduction velocity, was specifically associated with recall of

autobiographical memories from real life rather than performance on the more constrained

laboratory-based memory tests.

Discussion

The conduction velocity of action potentials along axons is crucial for neural communication.

Until recently it was not possible to examine metrics associated with axonal conduction

velocity, such as the g-ratio, in vivo in the human brain, with this being largely the preserve of

studies involving non-human animals. However, by combining diffusion MRI with quantitative

structural MRI scans optimised to assess myelination (Callaghan et al., 2014), it is now possible

to estimate the MR g-ratio in vivo in humans (Drakesmith et al., 2019; Mohammadi et al.,

2015; Mohammadi and Callaghan, 2021; Stikov et al., 2015). Here, in the first application to

human cognition, we found that variations in the MR g-ratio specifically in the

parahippocampal cingulum bundle were associated with individual differences in

autobiographical memory recall ability in a large sample of healthy adults. Under various

modelling assumptions (detailed below) this was associated with larger inner axon diameters

rather than myelin content. Moreover, we identified a possible link between autobiographical

memory recall ability and more coherently organised neurites. Our findings were also specific

to autobiographical memory retrieval and were not evident for laboratory-based memory tests.

These results offer a new perspective on individual differences in autobiographical memory

recall ability. This is especially welcome given the current lack of a clear biological explanation

for such variations in the healthy population (Clark et al., 2020; Clark et al., 2021a; LePort et

al., 2012; Maguire et al., 2003; Palombo et al., 2018; Van Petten, 2004).

A key property of the MR g-ratio is that it provides a non-invasive methodology with

which to relate in vivo structural neuroimaging of humans with axonal conduction velocity

(Berman et al., 2019; Drakesmith et al., 2019). Moreover, the identification of a statistically

significant relationship with the MR g-ratio can help to guide as to which of inner axon

diameter or myelin thickness is more likely to be associated with variations in conduction

velocity (Caeyenberghs et al., 2016; Kaller et al., 2017; Lakhani et al., 2016; Waxman, 1980;

Xin and Chan, 2020).

Our inferences rested upon the assumption that a significant relationship with the MR

g-ratio reflected faster conduction velocity, given previous work suggesting that faster

conduction velocity might be associated with better cognition (e.g. Brancucci, 2012; Dicke and

Roth, 2016; Miller, 1994; Reed and Jensen, 1992). As shown in Figure 2, there are three main

scenarios describing how changes in the MR g-ratio are related to the underlying

microstructural properties, given a faster conduction velocity. The positive relationship we

observed between the parahippocampal cingulum bundle MR g-ratio and autobiographical

memory recall ability suggests that this effect was associated predominantly with larger inner

axon diameter (Figure 2A orange arrow, and Figure 2D). By contrast, had a negative correlation

between the MR g-ratio and autobiographical memory recall ability been identified, we could

instead have inferred that myelination was the relevant microstructural feature. The lack of

relationship between parahippocampal cingulum bundle magnetisation transfer saturation

values (optimised to assess myelination) and autobiographical memory recall ability provided

further corroboration that myelin may not have been influential in this context. Greater

myelination is often held to be a prominent influence on behavioural and cognitive performance

(Caeyenberghs et al., 2016; Fields and Bukalo, 2020; Kaller et al., 2017; Lakhani et al., 2016;

Waxman, 1980; Xin and Chan, 2020). By contrast, our results highlight the potentially

important role that the inner axon diameter could be playing in autobiographical memory recall.

We also found that lower neurite dispersion, suggesting more coherent neurite

organisation, was related to better autobiographical memory recall ability. Our measure of

neurite dispersion was obtained using the NODDI biophysical model (Zhang et al., 2012),

which aims to isolate the organisation of the neurites in a voxel from the density of the neurites

in a voxel. While neurite dispersion was significantly related to autobiographical memory

recall, no relationship was observed with neurite density.

A larger inner axon diameter reduces resistance to action potential signals, enabling

greater conduction velocities (Gasser and Grundfest, 1939; Hursh, 1939), and coherently

organised fibres may decrease the distance signals need to travel, further reducing

communication times (Salami et al., 2003; Steriade, 1995). This combination of features might

optimise a fibre bundle for faster communication, and this may benefit autobiographical

memory recall.

These results were specific to one white matter tract, the parahippocampal cingulum

bundle (Bubb et al., 2018). MR g-ratio measures from no other tract, including the dorsal part

of the cingulum bundle, showed any association with autobiographical memory recall ability.

The parahippocampal cingulum bundle directly connects the hippocampus, parahippocampal,

retrosplenial and parietal cortices, the anterior thalamus and, through other subdivisions of the

cingulum bundle, the medial prefrontal cortex. These regions are typically engaged during

fMRI studies of autobiographical memory recall (Andrews-Hanna et al., 2014; Maguire, 2001;

Spreng et al., 2009; Svoboda et al., 2006), and damage to them is often associated with

autobiographical memory impairments (Berryhill et al., 2007; McCormick et al., 2018;

Scoville and Milner, 1957; Vann et al., 2009). The retrosplenial and parahippocampal cortices

are thought to provide visuospatial elements of autobiographical memories (Dalton and

Maguire, 2017; Epstein and Higgins, 2007; Mullally and Maguire, 2011; Vann et al., 2009),

while the medial prefrontal cortex may initiate autobiographical retrieval and support schema-

guided recall (Gilboa and Marlatte, 2017; McCormick et al., 2018; McCormick et al., 2020).

The parahippocampal cingulum bundle is, therefore, uniquely positioned as a transmission

highway enabling this information to reach the hippocampus, where memories can be

reconstructed (Bartlett, 1932; Hassabis and Maguire, 2007; Schacter et al., 2012). Larger inner

axon diameters and coherently organised neurites could facilitate rapid information flow along

the parahippocampal cingulum bundle leading to more memory elements being available

simultaneously, which in turn may result in increased detail and better integration of a memory

representation.

There have only been a small number of previous studies investigating white matter

tracts and individual differences in autobiographical memory recall ability in healthy people.

These focused on standard DTI parameters, such as fractional anisotropy and mean diffusivity,

and not the MR g-ratio. Two such studies were suggestive of a relationship between

parahippocampal cingulum bundle fractional anisotropy and autobiographical memory recall

(Irish et al., 2014; Memel et al., 2020). Our analyses of the standard DTI parameters, therefore,

replicated these previous results. Direct comparison between these studies and our findings is

difficult, however, given the relatively small sample sizes, older age of participants in some of

the extant studies, and the testing of mixed groups of healthy older people and patients with

dementia. In addition, fractional anisotropy and other physical parameters lack biological

specificity (Jensen and Helpern, 2010; Jones et al., 2013a) and, consequently, they cannot

speak to questions concerning axonal conduction velocity, which is our main interest here. In

a similar vein, we are not aware of any reports of bilateral lesions that selectively compromise

or sever the parahippocampal cingulum bundle in humans. Given our findings, we would

predict that such lesions would adversely affect the ability to recall autobiographical memories.

Despite its prime location and connectivity at the heart of the brain's autobiographical

memory system, the dearth of studies in humans and the rarity of selective bilateral lesions to

the parahippocampal cingulum bundle have perhaps obscured its importance when compared to other, more celebrated, memory-related white matter tracts. We also examined two such tracts, the fornix and the uncinate fasciculus, but in both cases no significant relationships between any of our neuroimaging metrics and autobiographical memory recall ability were evident in our large cohort of young healthy adults. As outlined in Figure 2, there was one possible scenario where a null relationship with the MR g-ratio could reflect an association with conduction velocity – if both the inner axon diameter and myelin thickness increased proportionally to each other (Figure 2A black arrow, and Figure 2E). However, we found no associations between autobiographical memory recall and both the MR g-ratio *and* magnetisation transfer saturation values of the fornix and uncinate fasciculus, which speaks against this explanation. In terms of standard DTI parameters, unlike a previous study (Hodgetts et al., 2017), we did not find a relationship between fornix fractional anisotropy and autobiographical memory recall ability, which may be due to the larger sample that we examined.

Both the fornix and parahippocampal cingulum bundle are vulnerable to partial volume effects (Concha et al., 2005). However, to mitigate this issue we took steps to ensure the data were extracted only from white matter voxels (see Materials and Methods). Diffusion data are also susceptible to distortions which can particularly affect the uncinate fasciculus. We addressed this challenge by using a new technique that improves distortion correction in this region (Clark et al., 2021b). The absence of fornix findings in our study could echo those relating to hippocampal volume (Clark et al., 2020; Clark et al., 2021a), whereby the structure is widely acknowledged to be involved in autobiographical memory recall, and damage impedes retrieval (Aggleton et al., 2000; D'Esposito et al., 1995; Gaffan and Gaffan, 1991; Tsivilis et al., 2008), but microstructural variations have limited impact within the healthy adult population. Regarding the uncinate fasciculus, unilateral lesions do not seem to significantly

impair performance on laboratory-based memory tests (Papagno et al., 2011) or the recall of

premorbid autobiographical memories (Levine et al., 2009). Bilateral uncinate fasciculus

lesions are very rare in humans, but might result in greater memory impairment. Alternatively,

hippocampal-prefrontal connections may be better served by other pathways, for example, via

the fornix or parts of the cingulum bundle.

Our findings were not only specific to the parahippocampal cingulum bundle, but also

to the internal details of autobiographical memories, which reflect the episodicity of past

experiences. By contrast, our control measure of external details, also from the same task but

which concerns non-episodic information in the autobiographical memories, did not correlate

with any white matter microstructure metrics. Moreover, no relationships were identified with

any of the eight laboratory-based memory tasks examined, highlighting the specific nature of

the relationship between parahippocampal cingulum bundle MR g-ratio and, by inference,

conduction velocity, and autobiographical memory recall ability. It may be that vivid, detailed,

multimodal, autobiographical memories rely on inter-regional connectivity, particularly that

supported by the parahippocampal cingulum bundle, to a greater degree than simpler, more

constrained laboratory-based memory tests. This result aligns with previous work involving an

fMRI meta-analysis which showed that recall of autobiographical memories and laboratory-

based memory stimuli were associated with substantially different neural substrates

(McDermott et al., 2009). More generally, our findings add to the increasingly-recognised

importance of studying real-world cognition in order to fully characterise brain-behaviour

relationships (Maguire, 2001, 2012; Miller et al., 2022; Mobbs et al., 2021; Nastase et al., 2020;

Spiers and Maguire, 2007).

Age, gender, scanner and the number of voxels within an ROI were included as

covariates in all analyses, limiting the potential confounding effects of these variables. Our

analyses were also performed using weighted means from each of the three white matter tracts

of interest rather than voxel-wise across the whole brain, reducing the potential for false

positives (Marek et al., 2022). We also tested a sample of over 200 participants, which has been

suggested as sufficient for correlational neuroimaging research such as that performed here

(Cecchetti and Handjaras, pre print; DeYoung et al., pre print).

Nevertheless, as with all neuroimaging techniques, methodological limitations need to

be considered when measuring the MR g-ratio (see Campbell et al., 2018; Mohammadi and

Callaghan, 2021 for in depth methodological reviews). First, the MR g-ratio is an area-

weighted average of all microscopic axons in an MRI-voxel that is slightly weighted towards

larger axons. Second, MR proxies are required to estimate myelin and axonal volumes for the

calculation of the MR g-ratio. Multiple methodologies are available to estimate myelin and

axonal volumes, with no consensus yet reached as to the best combination (Mohammadi and

Callaghan, 2021). Third, a calibration step is required to more closely align the MR proxies

with the estimated volume fractions. Here, we used the standard single-point calibration

method to estimate the slope of the myelin-based proxy, assuming that the offset can be

neglected. Future work may be able to improve these calibrations. But, of note, since a non-

negligible offset in the myelin-based proxy can increase the error in the MR g-ratio, any

calibration improvements would likely serve to increase the observed correlation. Fourth,

although the model of the MR g-ratio allows for fibre dispersion (Stikov et al., 2015), this is

not accounted for in mapping to conduction velocity. Going forward, more advanced models

may help to further elaborate on these relationships. Finally, our myelin measure can be

influenced by any factor, cognitive or demographic, that leads to a difference in absolute

myelination. In principle, unexplained variance could obscure a true underlying relationship

with autobiographical memory. To mitigate this, our analyses controlled for age, gender and

any potential scanner-related differences. The MR g-ratio, however, is less affected by this

limitation because it is a relative measure and does not depend on absolute myelin content but

on the balance between absolute myelin and axonal volumes.

While we examined the parahippocampal cingulum bundle as a unitary pathway, it

comprises both long and short association fibres with differing connectivity (Bubb et al., 2018).

Some fibres will form long range connections between, for example, the hippocampus and

retrosplenial cortex, whereas others will make shorter range connections between neighbouring

regions. The use of Connectom MRI scanners (Jones et al., 2018) is starting to make

examination of short range "u-fibres" possible in vivo in humans (Movahedian Attar et al.,

2020; Shastin et al., 2022). While there are currently only four of these scanners in the world,

future studies using Connectom MRI could seek to identify specific connections *within* the

parahippocampal cingulum bundle that relate to individual differences in autobiographical

memory recall.

In conclusion, white matter microstructure measures related to conduction velocity are

now possible to derive in vivo in humans. This has the potential to provide novel insights into

how the brain processes and integrates information (Berman et al., 2019; Drakesmith et al.,

2019), deepening our understanding of the information flow that underpins critical cognitive

functions such as autobiographical memory recall.

Materials and Methods

Participants

Two hundred and seventeen healthy people took part in the study, including 109 females and

108 males. The age range was restricted to 20-41 years old to limit the possible effects of aging

(mean age = 29.0 years, *SD* = 5.60). Participants had English as their first language and reported

no history of psychological, psychiatric or neurological conditions. Our aim was to assess

people from the general population who would not be classed as having extreme expertise on

classic hippocampal tasks, as this could affect hippocampal structure (Maguire et al., 2000;

Woollett and Maguire, 2011). Consequently, people with vocations such as taxi driving (or

those training to be taxi drivers), ship navigators, aeroplane pilots, or those with regular hobbies

including orienteering, or taking part in memory sports and competitions, were excluded. Of

the approximately 2000 people who contacted us, 23 were explicitly excluded on this basis.

Participants were reimbursed £10 per hour for taking part which was paid at study completion.

All participants gave written informed consent and the study was approved by the University

College London Research Ethics Committee (project ID: 6743/001).

A sample size of 217 was determined during study design to be robust to employing

different statistical approaches when answering multiple questions of interest. Specifically, the

sample allowed for sufficient power to identify medium effect sizes when conducting

correlation analyses at alpha levels of 0.01 and when comparing correlations at alpha levels of

0.05 (Cohen, 1992). Samples of over 200 participants have also been suggested as sufficient

for correlational neuroimaging research similar to that performed here (Cecchetti and

Handjaras, pre print; DeYoung et al., pre print).

Procedure

Participants completed the study over multiple visits. Diffusion imaging and magnetisation

transfer saturation scans were acquired on two separate days, and the Autobiographical

Interview was conducted during a third visit. All participants completed all parts of the study.

The Autobiographical Interview

This widely-used test (Levine et al., 2002) was employed to measure autobiographical memory

recall ability. Participants are asked to provide autobiographical memories from a specific time

and place over four time periods – early childhood (up to age 11), teenage years (aged from

11-17), adulthood (from age 18 years to 12 months prior to the interview; two memories are

requested) and the last year (a memory from the last 12 months); therefore, five memories in

total are harvested. Recordings of the memory descriptions are transcribed for later scoring.

The main outcome measure of the Autobiographical Interview is the mean number of

internal details included in the description of an event from across the five autobiographical

memories. Internal details are those describing the event in question (i.e. episodic details) and

include event, place, time and perceptual information, as well as thoughts and emotions relating

to the event itself. We used the secondary outcome measure of the Autobiographical Interview,

the mean number of external details included in the five autobiographical memories, as a

control measure. External details include semantic information concerning the event, or other

non-event information, and are not considered to reflect autobiographical memory recall

ability.

Double scoring was performed on 20% of the data. Inter-class correlation coefficients,

with a two-way random effects model looking for absolute agreement were calculated for both

internal and external details. This was performed both for individual memories and as an

average of all five memories across each participant. For internal details the coefficients were

0.94 and 0.97 respectively, and for external details they were 0.84 and 0.87 respectively. For

reference, a score of 0.8 or above is considered excellent agreement beyond chance.

Laboratory-based memory tests

Eight laboratory-based memory tasks were also administered to participants. These are

standard memory tests that are often used in neuropsychological settings. Tasks were

performed and scored in line with their standardised and published protocols.

The ability to recall a short narrative was examined using the immediate and delayed

recall tests of the Logical Memory subtest of the Wechsler Memory Scale IV (Wechsler, 2009).

Verbal list recall was assessed using the immediate and delayed recall tests of the Rey Auditory

Verbal Learning Test (see Strauss et al., 2006). Visuospatial recall was examined using the

delayed recall of the Rey–Osterrieth Complex Figure (Rey, 1941). Recognition memory was

investigated using the Warrington Recognition Memory Tests for Words and Faces

(Warrington, 1984). Finally, participants also underwent the "Dead or Alive" task which probes

general knowledge about whether famous individuals have died or are still alive, providing a

measure of semantic memory (Kapur et al., 1989).

Diffusion MRI data acquisition

Three MRI scanners were used to collect the neuroimaging data. All scanners were Siemens

Magnetom TIM Trio systems with 32 channel head coils and were located at the same

neuroimaging centre, running the same software. The sequences were loaded identically onto

the individual scanners. Participant set-up and positioning followed the same protocol for each

scanner.

Diffusion-weighted images were collected using the multiband accelerated EPI pulse

sequence developed by the Centre for Magnetic Resonance Research at the University of

Minnesota (R012a-c, R013a on VB17, https://www.cmrr.umn.edu/multiband/; Feinberg et al.,

2010; Xu et al., 2013). Acquisition parameters were: resolution = 1.7 mm isotropic; FOV =

220 mm × 220 mm × 138 mm; 60 directions with 6 interleaved b0 images, echo time (TE) =

112 ms, repetition time (TR) = 4.84 s, with a multiband acceleration factor of 3. The sequence

was performed 4 times – twice with b-values of 1000 and twice with b-values of 2500. The

first acquisition of each set of b-values was performed with phase-encoding in the anterior to

posterior direction (blip-up), the second in the posterior to anterior direction (blip-down). The

total acquisition time was 22 minutes.

Magnetisation transfer saturation data acquisition

The specific scanner used to collect a participant's diffusion-weighted images was also used to

obtain their magnetisation transfer saturation map.

Whole brain structural maps of magnetisation transfer saturation, at an isotropic

resolution of 800 μm, were derived from a multi-parameter mapping quantitative imaging

protocol (Callaghan et al., 2015; Callaghan et al., 2019; Weiskopf et al., 2013). This protocol

consisted of the acquisition of three multi-echo gradient-echo acquisitions with either proton

density, T1 or magnetisation transfer weighting. Each acquisition had a TR of 25 ms. Proton

density weighting was achieved with an excitation flip angle of 6^0, which was increased to 21^0

to achieve T1 weighting. Magnetisation transfer weighting was achieved through the

application of a Gaussian RF pulse 2 kHz off resonance with 4 ms duration and a nominal flip

angle of 220^0. This acquisition had an excitation flip angle of 6^0. The field of view was 256

mm head-foot, 224 mm anterior-posterior, and 179 mm right-left. The multiple gradient echoes

per contrast were acquired with alternating readout gradient polarity at eight equidistant echo

times ranging from 2.34 to 18.44 ms in steps of 2.30 ms using a readout bandwidth of 488

Hz/pixel. Only six echoes were acquired for the magnetisation transfer weighted volume to

facilitate the off-resonance pre-saturation pulse and subsequent spoiling gradient within the

TR. To accelerate the data acquisition, partially parallel imaging using the GRAPPA algorithm

was employed in each phase-encoded direction (anterior-posterior and right-left) with forty

integrated reference lines and a speed up factor of two. Calibration data were also acquired at

the outset of each session to correct for inhomogeneities in the RF transmit field (Lutti et al.,

2010). The total acquisition time was 27 minutes.

Diffusion MRI pre-processing

The diffusion MRI data were processed using the ACID toolbox (www.diffusiontools.com) within SPM12 (www.fil.ion.ucl.ac.uk/spm). The weighted average consecutive HySCO pipeline described in Clark et al. (2021b) was followed, with the addition of multi-shell Position-Orientation Adaptive Smoothing (msPOAS; Becker et al., 2012) and Rician bias correction (André et al., 2014). In brief, the blip-up and blip-down data were first separately corrected for motion and eddy current artefacts. Next, msPOAS was performed, followed by correction for susceptibility-related distortion artefacts using the HySCO2 module (Macdonald and Ruthotto, 2018; Ruthotto et al., 2012). Tensor fitting (Mohammadi et al., 2013) was then implemented separately on each of the distortion corrected blip-up and blip-down datasets to estimate FA maps. HySCO2 was then repeated using the distortion corrected and brain-masked FA maps as input instead of b0 images; the second HySCO2 field map being consecutively applied to the "pre-corrected" diffusion MRI data. Finally, Rician bias noise correction was employed on the distortion corrected data (André et al., 2014), before the data were combined using a weighted average to minimise information loss due to susceptibility distortion blurring induced by local spatial compression.

Magnetisation transfer saturation pre-processing

The magnetisation transfer saturation data were processed for each participant using the hMRI toolbox (Tabelow et al., 2019) within SPM12. The default toolbox configuration settings were used, with the exception that correction for imperfect spoiling was additionally enabled (Corbin and Callaghan, 2021). The output magnetisation transfer saturation map quantified the degree of saturation of the steady state signal induced by the application of the off-resonance pre-

pulse, having accounted for spatially varying T_1 times and RF field inhomogeneity (Weiskopf

et al., 2013).

Each participant's magnetisation transfer saturation map was segmented into white

matter probability maps using the unified segmentation approach (Ashburner and Friston,

2005), but with no bias field correction (since the magnetisation transfer saturation map does

not suffer from any bias field modulation) and using the tissue probability maps developed by

Lorio et al. (2016).

Diffusion model fitting

The MR g-ratio was calculated according to Ellerbrock and Mohammadi (2018):

$$g_{MR} = \sqrt{1 - \frac{MVF_{MR}}{MVF_{MR} + AVF_{MR}}}$$

with MVF_{MR} being the myelin-volume fraction estimated from the magnetisation transfer

saturation map and AVF_{MR} being the axonal-volume fraction. The AVF_{MR} was estimated as

$AVF_{MR} = (1 - MVF_{MR})$ AWF according to Stikov et al. (2015), where AWF was obtained by

combining the intra-cellular fraction (ν_{icvf}) and isotropic fraction (ν_{iso}) maps from NODDI

(Zhang et al., 2012) as $AWF = (1 - \nu_{iso})\,\nu_{icvf}$. The magnetisation transfer saturation map

was obtained from the hMRI toolbox as described above. For calibration of the magnetisation

transfer saturation map to a myelin-volume fraction map ($MVF_{MR} = \alpha\,MT_{sat}$), we used the g-

ratio based calibration method as reported in Ellerbrock and Mohammadi (2018) and

Mohammadi and Callaghan (2021), with $\alpha = 0.1683$.

The NODDI biophysical model (Zhang et al., 2012) was also used to obtain maps of

the neurite orientation dispersion index and neurite density using the NODDI toolbox

(http://mig.cs.ucl.ac.uk/index.php?n=Tutorial.NODDImatlab).

Finally, for completeness, Axial-Symmetric DKI (Oeschger et al., 2021) was performed

on the pre-processed diffusion data using the ACID toolbox to generate maps of the more

commonly reported physical parameters of fractional anisotropy, mean diffusivity, mean

kurtosis, diffusivities parallel and diffusivities perpendicular.

Microstructure data extraction

Microstructure data extraction was performed in Montreal Neurological Institute (MNI) space.

The diffusion and magnetisation transfer saturation maps were transformed from native to MNI

space using the hMRI toolbox (Tabelow et al., 2019). This involved performing inter-subject

registration using DARTEL (Ashburner, 2007) on the segmented magnetisation transfer

saturation grey and white matter probability maps, with the resulting DARTEL template and

deformations then used to normalize the diffusion and magnetisation transfer saturation maps

to MNI space at 1.5 x 1.5 x 1.5mm.

Bilateral tract ROIs were defined using the Johns Hopkins probabilistic white matter

tractography atlas (Hua et al., 2008), with the exception of the fornix which was defined using

the ICBM-DTI-81 white-matter labels atlas (Mori et al., 2008) as the fornix is not available in

the probabilistic atlas. Our primary foci were the fornix, uncinate fasciculus and

parahippocampal cingulum bundle. However, we also performed exploratory analyses on six

other tracts - the anterior thalamic radiation, the dorsal cingulum bundle, forceps minor, inferior

longitudinal fasciculus, inferior occipitofrontal fasciculus and superior longitudinal fasciculus.

To reduce partial volume effects, for all tracts (with the exception of the fornix as the available

fornix tract was not probabilistic) the minimum probability threshold was set to 25%. In

addition, all of the tract ROIs were refined for each participant using their segmented

magnetisation transfer saturation white matter probability map, with a minimum probability of

90% to limit the mask to white matter. This also served to remove any residual mis-alignment

from the maps being transformed into MNI space, as no smoothing was performed to preserve

the quantitative values. As this resulted in differing tract ROI sizes for each participant, the

number of voxels in each tract for each participant was calculated. Mean values of the extracted

microstructure metrics from the tract ROIs were determined using a weighted average, where

voxels with higher white matter probabilities contributed more to the mean.

847   We used the well-established ROI approach (e.g. Ellerbrock and Mohammadi, 2018;

Memel et al., 2020) rather than, for example, a tract-based pipeline, in order to reduce the

influence of seed and target region selection. This is because small changes to these selections

can result in different tracts being identified. We also wanted to avoid the inclusion of excess

grey matter in the tracts themselves, because it is not possible to estimate the MR g-ratio in

grey matter tissue. In addition, an ROI approach reduced the potential for false positives in

comparison to performing tract-based voxel-wise analyses.

Statistical analyses

Analyses were performed in SPSS v27 unless otherwise stated. Data were summarised using

means and standard deviations. There were no missing data, and no data needed to be removed

from any analysis.

859   As we had different tract ROI sizes for each participant, we first assessed whether there

were any relationships between the number of voxels in the tract ROIs and autobiographical

memory recall ability. We performed partial correlations for each tract between the number of

voxels in the tract ROI and the number of internal details on the Autobiographical Interview,

with age, gender and scanner as covariates. No significant relationships were identified (all $r <$

0.12, all $p > 0.1$). However, to ensure no residual effects were present, the number of voxels in

a tract ROI was included as a covariate in the analyses.

In our main analyses, we first investigated the relationships between each

microstructure measure and the number of internal details from the Autobiographical Interview

using partial correlations, with bootstrapping performed 10,000 times to calculate confidence

intervals. Four covariates were included in each partial correlation: age, gender, scanner and

the number of voxels in a tract ROI. For these primary analyses, similar partial correlations

were performed for the external details control measure. If an internal details correlation was

significant, the internal and external details correlations were then directly compared in order

to test for statistical difference using the technique described by Meng et al. (1992). This

approach extends the Fisher z transformation, allowing for more accurate testing and

comparison of two related correlations. The correlation comparison was performed using the

R cocor package v1.1.3 (Diedenhofen and Musch, 2015).

We also investigated the relationship between the MR g-ratio and the magnetisation

transfer saturation values and eight laboratory-based memory tests. As with the main analyses,

partial correlations were performed between the MR g-ratio and magnetisation transfer

saturation values and the outcome measures of the memory tests with age, gender, scanner and

the number of voxels in a tract ROI included as covariates, and bootstrapping performed 10,000

times to calculate confidence intervals.

As the microstructure measures were investigated across several tracts, we corrected

for the repeated testing of the same measures across our three main tracts of interest (the fornix,

the uncinate fasciculus and the parahippocampal cingulum bundle) using the Bonferroni

method; dividing alpha = 0.05 by 3. Consequently, associations with a two-sided p-value <

0.017 were considered significant. As the comparison of correlations was performed only when

a significant correlation was identified, a two-sided p-value < 0.05 was deemed significant.

Acknowledgments

Thanks to Anna Monk, Victoria Hotchin, Gloria Pizzamiglio and Alice Liefgreen for assistance with data collection and scoring. Thanks also to Mohammad Ashtarayeh and Jan Malte Oeschger for their support with implementing the diffusion model fitting and g-ratio calculations.

Additional information

Funding

Funder: Wellcome

Grant reference numbers: 101759/Z/13/Z; 210567/Z/18/Z; 203147/Z/16/Z

Author: Eleanor A. Maguire

Funder: ERA-NET NEURON

Grant reference number: hMRI-ofSCI

Author: Martina F. Callaghan, Siawoosh Mohammadi

Funder: German Federal Ministry of Education and Research

Grant reference number: 01EW1711A and B

Author: Siawoosh Mohammadi

Funder: German Research Foundation

Grant reference numbers: MO 2397/5-1; MO 2397/4-1

Author: Siawoosh Mohammadi

Funder: Forschungszentrums Medizintechnik Hamburg

Grant reference numbers: 01fmthh2017

Author: Siawoosh Mohammadi

Funder: MRC and Spinal Research Charity

Grant reference number: MR/R000050/1

Author: Martina F. Callaghan

The funders had no role in study design, data collection and interpretation, or the decision to

submit the work for publication.

Author contributions

Ian A. Clark: conceptualization, methodology, investigation, data scoring, formal analysis,

writing – original draft, writing - review & editing; Siawoosh Mohammadi: formal analysis,

writing - review & editing; Martina F. Callaghan: Formal analysis, writing - review & editing;

Eleanor A. Maguire: conceptualization, methodology, funding acquisition, supervision, formal

analysis, writing – original draft, writing - review & editing

Author ORCIDs

Ian A. Clark https://orcid.org/0000-0002-5678-2190

Siawoosh Mohammadi https://orcid.org/0000-0003-1311-9636

Martina F. Callaghan https://orcid.org/0000-0003-0374-1659

Eleanor A. Maguire https://orcid.org/0000-0002-9470-6324

Ethics

Human subjects: All participants gave written informed consent and the study was approved

by the University College London Research Ethics Committee (project ID: 6743/001).

Additional files

• Appendix 1. Details of the simulation performed for Figure 2A. Investigation of the

MR g-ratio, magnetisation transfer saturation values, neurite orientation dispersion

index, neurite density and standard DTI parameters in the fornix and uncinate fasciculus

– no significant relationships were identified. Investigation of associations between the

MR g-ratio and magnetisation transfer saturation values of the fornix and uncinate

fasciculus and laboratory-based memory tests – no significant relationships were

identified. Exploratory analyses of the MR g-ratio, magnetisation transfer saturation

values, neurite orientation dispersion index, neurite density and standard DTI

parameters in other white matter tracts – no significant relationships were identified.

Investigation of the standard DTI parameters extracted from the parahippocampal

cingulum bundle. Investigation of the parahippocampal cingulum bundle MR g-ratio

and magnetisation transfer saturation values and the laboratory-based memory tests –

no significant relationships were identified.

• Supplementary file 1. Microstructure and standard DTI parameters data for the fornix

and uncinate fasciculus.

• Supplementary file 2. Microstructure and standard DTI parameters data for the

parahippocampal cingulum bundle and source data for Appendix 1-figures 9 and 10.

• Figure 5-source data 1

• Figure 6-source data 1

• Materials Design Analysis Reporting (MDAR) Form

Data availability

The data for every participant are provided in Figure 5-source data 1, Figure 6-source data 1,

Supplementary file 1 and Supplementary file 2.

References

Aggleton JP, McMackin D, Carpenter K, Hornak J, Kapur N, Halpin S, Wiles CM, Kamel H,
Brennan P, Carton S, Gaffan D (2000) **Differential cognitive effects of colloid cysts**
in the third ventricle that spare or compromise the fornix *Brain* **123**:800-815.
http://dx.doi.org/10.1093/brain/123.4.800
Aggleton JP, Wright NF, Rosene DL, Saunders RC (2015) **Complementary patterns of direct**
amygdala and hippocampal projections to the macaque prefrontal cortex *Cerebral*
Cortex **25**:4351-4373. http://dx.doi.org/10.1093/cercor/bhv019
André ED, Grinberg F, Farrher E, Maximov II, Shah NJ, Meyer C, Jaspar M, Muto V, Phillips
C, Balteau E (2014) **Influence of noise correction on intra- and inter-subject**
variability of quantitative metrics in diffusion kurtosis imaging *PloS One* **9**:e94531.
http://dx.doi.org/10.1371/journal.pone.0094531
Andrews-Hanna JR, Saxe R, Yarkoni T (2014) **Contributions of episodic retrieval and**
mentalizing to autobiographical thought: Evidence from functional
neuroimaging, resting-state connectivity, and fMRI meta-analyses *NeuroImage*
91:324-335. http://dx.doi.org/10.1016/j.neuroimage.2014.01.032
Arancibia-Cárcamo IL, Ford MC, Cossell L, Ishida K, Tohyama K, Attwell D (2017) **Node of**
Ranvier length as a potential regulator of myelinated axon conduction speed *eLife*
6:e23329. http://dx.doi.org/10.7554/eLife.23329
Ashburner J, Friston KJ (2005) **Unified segmentation** *NeuroImage* **26**:839-851.
http://dx.doi.org/10.1016/j.neuroimage.2005.02.018
Ashburner J (2007) **A fast diffeomorphic image registration algorithm** *NeuroImage* **38**:95-
113. http://dx.doi.org/10.1016/j.neuroimage.2007.07.007

Aston-Jones G, Rogers J, Shaver RD, Dinan TG, Moss DE (1985) **Age-impaired impulse flow from nucleus basalis to cortex** *Nature* **318**:462-464. http://dx.doi.org/10.1038/318462a0

Bartlett FC (1932) **Remembering: A study in experimental and social psychology**. Cambridge University Press, Cambridge, England.

Basser PJ (1995) **Inferring microstructural features and the physiological state of tissues from diffusion-weighted images** *NMR in Biomedicine* **8**:333-344. http://dx.doi.org/10.1002/nbm.1940080707

Becker SMA, Tabelow K, Voss HU, Anwander A, Heidemann RM, Polzehl J (2012) **Position-orientation adaptive smoothing of diffusion weighted magnetic resonance data (POAS)** *Medical Image Analysis* **16**:1142-1155. http://dx.doi.org/10.1016/j.media.2012.05.007

Berman S, West KL, Does MD, Yeatman JD, Mezer AA (2018) **Evaluating g-ratio weighted changes in the corpus callosum as a function of age and sex** *NeuroImage* **182**:304-313. http://dx.doi.org/10.1016/j.neuroimage.2017.06.076

Berman S, Filo S, Mezer AA (2019) **Modeling conduction delays in the corpus callosum using MRI-measured g-ratio** *NeuroImage* **195**:128-139. http://dx.doi.org/10.1016/j.neuroimage.2019.03.025

Berryhill ME, Phuong L, Picasso L, Cabeza R, Olson IR (2007) **Parietal lobe and episodic memory: Bilateral damage causes impaired free recall of autobiographical memory** *Journal of Neuroscience* **27**:14415. http://dx.doi.org/10.1523/JNEUROSCI.4163-07.2007

Brancucci A (2012) **Neural correlates of cognitive ability** *Journal of Neuroscience Research* **90**:1299-1309. https://doi.org/10.1002/jnr.23045

Bubb EJ, Metzler-Baddeley C, Aggleton JP (2018) **The cingulum bundle: Anatomy, function, and dysfunction** *Neuroscience & Biobehavioral Reviews* **92**:104-127. http://dx.doi.org/10.1016/j.neubiorev.2018.05.008

Bullock TH, Moore JK, Fields RD (1984) **Evolution of myelin sheaths: Both lamprey and hagfish lack myelin** *Neuroscience Letters* **48**:145-148. http://dx.doi.org/10.1016/0304-3940(84)90010-7

Caeyenberghs K, Metzler-Baddeley C, Foley S, Jones DK (2016) **Dynamics of the human structural connectome underlying working memory training** *Journal of Neuroscience* **36**:4056. http://dx.doi.org/10.1523/JNEUROSCI.1973-15.2016

Callaghan MF, Freund P, Draganski B, Anderson E, Cappelletti M, Chowdhury R, Diedrichsen J, FitzGerald THB, Smittenaar P, Helms G, Lutti A, Weiskopf N (2014) **Widespread age-related differences in the human brain microstructure revealed by quantitative magnetic resonance imaging** *Neurobiology of Aging* **35**:1862-1872. http://dx.doi.org/10.1016/j.neurobiolaging.2014.02.008

Callaghan MF, Josephs O, Herbst M, Zaitsev M, Todd N, Weiskopf N (2015) **An evaluation of prospective motion correction (PMC) for high resolution quantitative MRI** *Frontiers in Neuroscience* **9**. http://dx.doi.org/10.3389/fnins.2015.00097

Callaghan MF, Lutti A, Ashburner J, Balteau E, Corbin N, Draganski B, Helms G, Kherif F, Leutritz T, Mohammadi S, Phillips C, Reimer E, Ruthotto L, Seif M, Tabelow K, Ziegler G, Weiskopf N (2019) **Example dataset for the hMRI toolbox** *Data in Brief* **25**:104132. http://dx.doi.org/10.1016/j.dib.2019.104132

Campbell JSW, Leppert IR, Narayanan S, Boudreau M, Duval T, Cohen-Adad J, Pike GB, Stikov N (2018) **Promise and pitfalls of g-ratio estimation with MRI** *NeuroImage* **182**:80-96. http://dx.doi.org/10.1016/j.neuroimage.2017.08.038

Cecchetti L, Handjaras G (pre print) **Reproducible brain-wide association studies do not**
necessarily require thousands of individuals *PsyArXiv*.
http://dx.doi.org/10.31234/osf.io/c8xwe
Cercignani M, Giulietti G, Dowell NG, Gabel M, Broad R, Leigh PN, Harrison NA, Bozzali
M (2017) **Characterizing axonal myelination within the healthy population: a**
tract-by-tract mapping of effects of age and gender on the fiber g-ratio
Neurobiology of Aging **49**:109-118.
http://dx.doi.org/10.1016/j.neurobiolaging.2016.09.016
Chomiak T, Hu B (2009) **What is the optimal value of the g-ratio for myelinated fibers in**
the rat CNS? A theoretical approach *PloS One* **4**:e7754.
http://dx.doi.org/10.1371/journal.pone.0007754
Clark IA, Monk AM, Hotchin V, Pizzamiglio G, Liefgreen A, Callaghan MF, Maguire EA
(2020) **Does hippocampal volume explain performance differences on**
hippocampal-dependent tasks? *NeuroImage* **221**:117211.
http://dx.doi.org/10.1016/j.neuroimage.2020.117211
Clark IA, Callaghan MF, Weiskopf N, Maguire EA (2021a) **The relationship between**
hippocampal-dependent task performance and hippocampal grey matter
myelination and iron content *Brain and Neuroscience Advances* **5**:1-8.
http://dx.doi.org/10.1177/23982128211011923
Clark IA, Callaghan MF, Weiskopf N, Maguire EA, Mohammadi S (2021b) **Reducing**
susceptibility distortion related image blurring in diffusion MRI EPI data
Frontiers in Neuroscience **15**:706473. https://dx.doi.org/10.3389/fnins.2021.706473
Cohen J (1992) **A power primer** *Psychological Bulletin* **112**:155-159.
http://dx.doi.org/10.1037/0033-2909.112.1.155
Concha L, Gross DW, Beaulieu C (2005) **Diffusion tensor tractography of the limbic system**
American Journal of Neuroradiology **26**:2267-2274.
Corbin N, Callaghan MF (2021) **Imperfect spoiling in variable flip angle T1 mapping at**
7T: quantifying and minimising impact *Magnetic Resonance in Medicine* **86**:693-
708. http://dx.doi.org/10.1002/mrm.28720
Croxson PL, Johansen-Berg H, Behrens TEJ, Robson MD, Pinsk MA, Gross CG, Richter W,
Richter MC, Kastner S, Rushworth MFS (2005) **Quantitative investigation of**
connections of the prefrontal cortex in the human and macaque using probabilistic
diffusion tractography *Journal of Neuroscience* **25**:8854-8866.
http://dx.doi.org/10.1523/JNEUROSCI.1311-05.2005
D'Esposito M, Verfaellie M, Alexander MP, Katz DI (1995) **Amnesia following traumatic**
bilateral fornix transection *Neurology* **45**:1546.
http://dx.doi.org/10.1212/WNL.45.8.1546
Dalton MA, Maguire EA (2017) **The pre/parasubiculum: a hippocampal hub for scene-**
based cognition? *Current Opinion in Behavioral Sciences* **17**:34-40.
http://dx.doi.org/10.1016/j.cobeha.2017.06.001
DeYoung CG, Sassenberg TA, Abend R, Allen T, Beaty R, Bellgrove M, Blain SD, Bzdok D,
Chavez RS, Engel SA, Ma F, Fornito A, Genç E, Goghari V, Grazioplene R, Hanson
JL, Haxby J, Hilger K, Homan P, Joyner K, Kaczkurkin A, Latzman RD, Martin EA,
Passamonti L, Pickering A, Safron A, Servaas M, Smillie L, Spreng RN, Tiego J,
Viding E, Wacker J (pre print) **Reproducible between-person brain-behavior**
associations do not always require thousands of individuals *PsyArXiv*.
http://dx.doi.org/10.31234/osf.io/sfnmk

Dicke U, Roth G (2016) **Neuronal factors determining high intelligence** *Philosophical*
Transactions of the Royal Society of London. Series B: Biological Sciences
371:20150180. http://dx.doi.org/10.1098/rstb.2015.0180
Diedenhofen B, Musch J (2015) **cocor: A comprehensive solution for the statistical**
comparison of correlations *PloS One* **10**:e0121945.
http://dx.doi.org/10.1371/journal.pone.0121945
Drakesmith M, Harms R, Rudrapatna SU, Parker GD, Evans CJ, Jones DK (2019) **Estimating**
axon conduction velocity in vivo from microstructural MRI *NeuroImage*
203:116186. http://dx.doi.org/10.1016/j.neuroimage.2019.116186
Ellerbrock I, Mohammadi S (2018) **Four in vivo g-ratio-weighted imaging methods:**
Comparability and repeatability at the group level *Frontiers in Neuroscience* **39**:24-
41. http://dx.doi.org/10.1002/hbm.23858
Epstein RA, Higgins JS (2007) **Differential parahippocampal and retrosplenial**
involvement in three types of visual scene recognition *Cerebral Cortex* **17**:1680-
1693. http://dx.doi.org/10.1093/cercor/bhl079
Feinberg DA, Moeller S, Smith SM, Auerbach E, Ramanna S, Glasser MF, Miller KL, Ugurbil
K, Yacoub E (2010) **Multiplexed echo planar imaging for sub-second whole brain**
fMRI and fast diffusion imaging *PloS One* **5**:e15710.
http://dx.doi.org/10.1371/journal.pone.0015710
Fields RD, Bukalo O (2020) **Myelin makes memories** *Nature Neuroscience* **23**:469-470.
http://dx.doi.org/10.1038/s41593-020-0606-x
Gaffan D, Gaffan EA (1991) **Amnesia in man following transection of the fornix: A review**
Brain **114**:2611-2618. http://dx.doi.org/10.1093/brain/114.6.2611
Gasser HS, Grundfest H (1939) **Axon diameters in relation to the spike dimensions and the**
conduction velocity in mammalian a fibers *American Journal of Physiology* **127**:393-
414. http://dx.doi.org/10.1152/ajplegacy.1939.127.2.393
Gilboa A, Marlatte H (2017) **Neurobiology of schemas and schema-mediated memory**
Trends in Cognitive Sciences **21**:618-631. http://dx.doi.org/10.1016/j.tics.2017.04.013
Hassabis D, Maguire EA (2007) **Deconstructing episodic memory with construction** *Trends*
in Cognitive Sciences **11**:299-306. http://dx.doi.org/10.1016/j.tics.2007.05.001
Hodgetts CJ, Postans M, Warne N, Varnava A, Lawrence AD, Graham KS (2017) **Distinct**
contributions of the fornix and inferior longitudinal fasciculus to episodic and
semantic autobiographical memory *Cortex* **94**:1-14.
http://dx.doi.org/10.1016/j.cortex.2017.05.010
Hua K, Zhang J, Wakana S, Jiang H, Li X, Reich DS, Calabresi PA, Pekar JJ, van Zijl PCM,
Mori S (2008) **Tract probability maps in stereotaxic spaces: Analyses of white**
matter anatomy and tract-specific quantification *NeuroImage* **39**:336-347.
http://dx.doi.org/10.1016/j.neuroimage.2007.07.053
Hursh JB (1939) **Conduction velocity and diameter of nerve fibers** *American Journal of*
Physiology **127**:131-139. http://dx.doi.org/10.1152/ajplegacy.1939.127.1.131
Huxley AF, Stämpeli R (1949) **Evidence for saltatory conduction in peripheral myelinated**
nerve fibres *Journal of Physiology* **108**:315-339.
https://doi.org/10.1113/jphysiol.1949.sp004335
Irish M, Hornberger M, El Wahsh S, Lam BYK, Lah S, Miller L, Hsieh S, Hodges JR, Piguet
O (2014) **Grey and white matter correlates of recent and remote autobiographical**
memory retrieval – Insights from the dementias *PloS One* **9**:e113081.
http://dx.doi.org/10.1371/journal.pone.0113081

Jensen JH, Helpern JA (2010) **MRI quantification of non-Gaussian water diffusion by kurtosis analysis** *NMR in Biomedicine* **23**:698-710. http://dx.doi.org/10.1002/nbm.1518

Jones DK, Knösche TR, Turner R (2013a) **White matter integrity, fiber count, and other fallacies: The do's and don'ts of diffusion MRI** *NeuroImage* **73**:239-254. http://dx.doi.org/10.1016/j.neuroimage.2012.06.081

Jones DK, Christiansen KF, Chapman RJ, Aggleton JP (2013b) **Distinct subdivisions of the cingulum bundle revealed by diffusion MRI fibre tracking: Implications for neuropsychological investigations** *Neuropsychologia* **51**:67-78. http://dx.doi.org/10.1016/j.neuropsychologia.2012.11.018

Jones DK, Alexander DC, Bowtell R, Cercignani M, Dell'Acqua F, McHugh DJ, Miller KL, Palombo M, Parker GJM, Rudrapatna US, Tax CMW (2018) **Microstructural imaging of the human brain with a 'super-scanner': 10 key advantages of ultra-strong gradients for diffusion MRI** *NeuroImage* **182**:8-38. http://dx.doi.org/10.1016/j.neuroimage.2018.05.047

Jung W, Lee J, Shin H-G, Nam Y, Zhang H, Oh S-H, Lee J (2018) **Whole brain g-ratio mapping using myelin water imaging (MWI) and neurite orientation dispersion and density imaging (NODDI)** *NeuroImage* **182**:379-388. http://dx.doi.org/10.1016/j.neuroimage.2017.09.053

Kaller MS, Lazari A, Blanco-Duque C, Sampaio-Baptista C, Johansen-Berg H (2017) **Myelin plasticity and behaviour—connecting the dots** *Current Opinion in Neurobiology* **47**:86-92. http://dx.doi.org/10.1016/j.conb.2017.09.014

Kapur N, Young A, Bateman D, Kennedy P (1989) **Focal retrograde amnesia: A long term clinical and neuropsychological follow-up** *Cortex* **25**:387-402. http://dx.doi.org/10.1016/S0010-9452(89)80053-X

Lakhani B, Borich MR, Jackson JN, Wadden KP, Peters S, Villamayor A, MacKay AL, Vavasour IM, Rauscher A, Boyd LA (2016) **Motor skill acquisition promotes human brain myelin plasticity** *Neural Plasticity*:7526135. http://dx.doi.org/10.1155/2016/7526135

LePort AKR, Mattfeld AT, Dickinson-Anson H, Fallon JH, Stark CEL, Kruggel F, Cahill L, McGaugh JL (2012) **Behavioral and neuroanatomical investigation of Highly Superior Autobiographical Memory (HSAM)** *Neurobiology of Learning and Memory* **98**:78-92. http://dx.doi.org/10.1016/j.nlm.2012.05.002

Levine B, Svoboda E, Hay JF, Winocur G, Moscovitch M (2002) **Aging and autobiographical memory: Dissociating episodic from semantic retrieval** *Psychology and Aging* **17**:677-689. http://dx.doi.org/10.1037/0882-7974.17.4.677

Levine B, Svoboda E, Turner GR, Mandic M, Mackey A (2009) **Behavioral and functional neuroanatomical correlates of anterograde autobiographical memory in isolated retrograde amnesic patient M.L** *Neuropsychologia* **47**:2188-2196. http://doi.org/10.1016/j.neuropsychologia.2008.12.026

Lorio S, Fresard S, Adaszewski S, Kherif F, Chowdhury R, Frackowiak RS, Ashburner J, Helms G, Weiskopf N, Lutti A, Draganski B (2016) **New tissue priors for improved automated classification of subcortical brain structures on MRI** *NeuroImage* **130**:157-166. http://dx.doi.org/10.1016/j.neuroimage.2016.01.062

Lutti A, Hutton C, Finsterbusch J, Helms G, Weiskopf N (2010) **Optimization and validation of methods for mapping of the radiofrequency transmit field at 3T** *Magnetic Resonance in Medicine* **64**:229-238. http://dx.doi.org/doi:10.1002/mrm.22421

Macdonald J, Ruthotto L (2018) **Improved susceptibility artifact correction of echo-planar MRI using the alternating direction method of multipliers** *Journal of Mathematical Imaging and Vision* **60**:268-282. http://dx.doi.org/10.1007/s10851-017-0757-x

Maguire EA, Gadian DG, Johnsrude IS, Good CD, Ashburner J, Frackowiak RSJ, Frith CD (2000) **Navigation-related structural change in the hippocampi of taxi drivers** *Proceedings of the National Academy of Sciences USA* **97**:4398-4403. http://dx.doi.org/10.1073/pnas.070039597

Maguire EA (2001) **Neuroimaging studies of autobiographical event memory** *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* **356**:1441-1451. http://dx.doi.org/10.1098/rstb.2001.0944

Maguire EA, Valentine ER, Wilding JM, Kapur N (2003) **Routes to remembering: the brains behind superior memory** *Nature Neuroscience* **6**:90-95. http://dx.doi.org/10.1038/nn988

Maguire EA (2012) **Studying the freely-behaving brain with fMRI** *NeuroImage* **62**:1170-1176. http://dx.doi.org/10.1016/j.neuroimage.2012.01.009

Marek S, Tervo-Clemmens B, Calabro FJ, Montez DF, Kay BP, Hatoum AS, Donohue MR, Foran W, Miller RL, Hendrickson TJ, Malone SM, Kandala S, Feczko E, Miranda-Dominguez O, Graham AM, Earl EA, Perrone AJ, Cordova M, Doyle O, Moore LA, Conan GM, Uriarte J, Snider K, Lynch BJ, Wilgenbusch JC, Pengo T, Tam A, Chen J, Newbold DJ, Zheng A, Seider NA, Van AN, Metoki A, Chauvin RJ, Laumann TO, Greene DJ, Petersen SE, Garavan H, Thompson WK, Nichols TE, Yeo BTT, Barch DM, Luna B, Fair DA, Dosenbach NUF (2022) **Reproducible brain-wide association studies require thousands of individuals** *Nature* **603**:654-660. http://dx.doi.org/10.1038/s41586-022-04492-9

McCormick C, Ciaramelli E, De Luca F, Maguire EA (2018) **Comparing and contrasting the cognitive effects of hippocampal and ventromedial prefrontal cortex damage: A review of human lesion studies** *Neuroscience* **274**:295-318. http://dx.doi.org/10.1016/j.neuroscience.2017.07.066

McCormick C, Barry DN, Jafarian A, Barnes GR, Maguire EA (2020) **vmPFC drives hippocampal processing during autobiographical memory recall regardless of remoteness** *Cerebral Cortex* **30**:5972-5987. http://dx.doi.org/10.1093/cercor/bhaa172

McDermott KB, Szpunar KK, Christ SE (2009) **Laboratory-based and autobiographical retrieval tasks differ substantially in their neural substrates** *Neuropsychologia* **47**:2290-2298. http://dx.doi.org/10.1016/j.neuropsychologia.2008.12.025

Memel M, Wank AA, Ryan L, Grilli MD (2020) **The relationship between episodic detail generation and anterotemporal, posteromedial, and hippocampal white matter tracts** *Cortex* **123**:124-140. http://dx.doi.org/10.1016/j.cortex.2019.10.010

Meng X-L, Rosenthal R, Rubin DB (1992) **Comparing correlated correlation coefficients** *Psychological Bulletin* **111**:172-175. http://dx.doi.org/10.1037/0033-2909.111.1.172

Miller CT, Gire D, Hoke K, Huk AC, Kelley D, Leopold DA, Smear MC, Theunissen F, Yartsev M, Niell CM (2022) **Natural behavior is the language of the brain** *Current Biology* **32**:R482-R493. http://dx.doi.org/10.1016/j.cub.2022.03.031

Miller EM (1994) **Intelligence and brain myelination: A hypothesis** *Personality and Individual Differences* **17**:803-832. http://dx.doi.org/10.1016/0191-8869(94)90049-3

Mobbs D, Wise T, Suthana N, Guzmán N, Kriegeskorte N, Leibo JZ (2021) **Promises and challenges of human computational ethology** *Neuron* **109**:2224-2238. http://dx.doi.org/10.1016/j.neuron.2021.05.021

Mohammadi S, Freund P, Feiweier T, Curt A, Weiskopf N (2013) **The impact of post-processing on spinal cord diffusion tensor imaging** *NeuroImage* **70**:377-385. http://dx.doi.org/10.1016/j.neuroimage.2012.12.058

Mohammadi S, Carey D, Dick F, Diedrichsen J, Sereno MI, Reisert M, Callaghan MF, Weiskopf N (2015) **Whole-brain in-vivo measurements of the axonal g-ratio in a group of 37 healthy volunteers** *Frontiers in Neuroscience* **9**. http://dx.doi.org/10.3389/fnins.2015.00441

Mohammadi S, Callaghan MF (2021) **Towards in vivo g-ratio mapping using MRI: unifying myelin and diffusion imaging** *Journal of Neuroscience Methods*:108990. https://dx.doi.org/10.1016/j.jneumeth.2020.108990

Mori S, Oishi K, Jiang H, Jiang L, Li X, Akhter K, Hua K, Faria AV, Mahmood A, Woods R, Toga AW, Pike GB, Neto PR, Evans A, Zhang J, Huang H, Miller MI, van Zijl P, Mazziotta J (2008) **Stereotaxic white matter atlas based on diffusion tensor imaging in an ICBM template** *NeuroImage* **40**:570-582. http://dx.doi.org/10.1016/j.neuroimage.2007.12.035

Movahedian Attar F, Kirilina E, Haenelt D, Pine KJ, Trampel R, Edwards LJ, Weiskopf N (2020) **Mapping short association fibers in the early cortical visual processing stream using in vivo diffusion tractography** *Cerebral Cortex* **30**:4496-4514. http://dx.doi.org/10.1093/cercor/bhaa049

Mullally SL, Maguire EA (2011) **A new role for the parahippocampal cortex in representing space** *The Journal of Neuroscience* **31**:7441-7449. http://dx.doi.org/10.1523/JNEUROSCI.0267-11.2011

Nastase SA, Goldstein A, Hasson U (2020) **Keep it real: rethinking the primacy of experimental control in cognitive neuroscience** *NeuroImage* **222**:117254. http://dx.doi.org/10.1016/j.neuroimage.2020.117254

Nave K-A (2010) **Myelination and support of axonal integrity by glia** *Nature* **468**:244-252. http://dx.doi.org/10.1038/nature09614

Oeschger J, Tabelow K, Mohammadi S (2021) **Reducing rician noise bias in axial-symmetric diffusion kurtosis imaging and biophysical tissue models**, Proceedings of the International Society for Magnetic Resonance in Medicine Scientific Meeting and Exhibition, p. 3654.

Palombo DJ, Alain C, Söderlund H, Khuu W, Levine B (2015) **Severely deficient autobiographical memory (SDAM) in healthy adults: A new mnemonic syndrome** *Neuropsychologia* **72**:105-118. http://dx.doi.org/10.1016/j.neuropsychologia.2015.04.012

Palombo DJ, Sheldon S, Levine B (2018) **Individual differences in autobiographical memory** *Trends in Cognitive Sciences* **22**:583-597. http://dx.doi.org/10.1016/j.tics.2018.04.007

Papagno C, Miracapillo C, Casarotti A, Romero Lauro LJ, Castellano A, Falini A, Casaceli G, Fava E, Bello L (2011) **What is the role of the uncinate fasciculus? Surgical removal and proper name retrieval** *Brain* **134**:405-414. http://dx.doi.org/10.1093/brain/awq283

Peters A, Moss MB, Sethares C (2000) **Effects of aging on myelinated nerve fibers in monkey primary visual cortex** *Journal of Comparative Neurology* **419**:364-376. https://doi.org/10.1002/(SICI)1096-9861(20000410)419:3<364::AID-CNE8>3.0.CO;2-R

Peters A, Sethares C (2002) **Aging and the myelinated fibers in prefrontal cortex and corpus callosum of the monkey** *Journal of Comparative Neurology* **442**:277-291. http://dx.doi.org/10.1002/cne.10099

Reed ET, Jensen AR (1993) **A somatosensory latency between the thalamus and cortex also correlates with level of intelligence** *Intelligence* **17**:443-450. http://dx.doi.org/10.1016/0160-2896(93)90011-S

Reed TE, Jensen AR (1992) **Conduction velocity in a brain nerve pathway of normal adults correlates with intelligence level** *Intelligence* **16**:259-272. http://dx.doi.org/10.1016/0160-2896(92)90009-G

Rey A (1941) **L'examen psychologique dans les cas d'encéphalopathie traumatique. (Les problems.). [The psychological examination in cases of traumatic encepholopathy. Problems.]** *Archives de Psychologie* **28**:215-285.

Rushton WAH (1951) **A theory of the effects of fibre size in medullated nerve** *Journal of Physiology* **115**:101-122. https://doi.org/10.1113/jphysiol.1951.sp004655

Ruthotto L, Kugel H, Olesch J, Fischer B, Modersitzki J, Burger M, Wolters CJPiM, Biology (2012) **Diffeomorphic susceptibility artifact correction of diffusion-weighted magnetic resonance images** *Physics in Medicine and Biology* **57**:5715. http://dx.doi.org/10.1088/0031-9155/57/18/5715

Salami M, Itami C, Tsumoto T, Kimura F (2003) **Change of conduction velocity by regional myelination yields constant latency irrespective of distance between thalamus and cortex** *Proceedings of the National Academy of Sciences* **100**:6174. http://dx.doi.org/10.1073/pnas.0937380100

Schacter DL, Addis DR, Hassabis D, Martin VC, Spreng RN, Szpunar KK (2012) **The future of memory: Remembering, imagining, and the brain** *Neuron* **76**:677-694. http://dx.doi.org/10.1016/j.neuron.2012.11.001

Schmidt H, Knösche TR (2019) **Action potential propagation and synchronisation in myelinated axons** *PLoS Computational Biology* **15**:e1007004. http://dx.doi.org/10.1371/journal.pcbi.1007004

Scoville WB, Milner B (1957) **Loss of recent memory after bilateral hippocampal lesions** *Journal of Neurology, Neurosurgery and Psychiatry* **20**:11-21. http://dx.doi.org/10.1136/jnnp.20.1.11

Shastin D, Genc S, Parker GD, Koller K, Tax CMW, Evans J, Hamandi K, Gray WP, Jones DK, Chamberland M (2022) **Surface-based tracking for short association fibre tractography** *NeuroImage*. http://dx.doi.org/10.1016/j.neuroimage.2022.119423

Spiers HJ, Maguire EA (2007) **Decoding human brain activity during real-world experiences** *Trends in Cognitive Sciences* **11**:356-365. http://dx.doi.org/10.1016/j.tics.2007.06.002

Spreng RN, Mar RA, Kim ASN (2009) **The common neural basis of autobiographical memory, prospection, navigation, theory of mind, and the default mode: A quantitative meta-analysis** *Journal of Cognitive Neuroscience* **21**:489-510. http://dx.doi.org/10.1162/jocn.2008.21029

Steriade M (1995) **Two channels in the cerebellothalamocortical system** *Journal of Comparative Neurology* **354**:57-70. https://doi.org/10.1002/cne.903540106

Stikov N, Campbell JSW, Stroh T, Lavelée M, Frey S, Novek J, Nuara S, Ho M-K, Bedell BJ, Dougherty RF, Leppert IR, Boudreau M, Narayanan S, Duval T, Cohen-Adad J, Picard P-A, Gasecka A, Côté D, Pike GB (2015) **In vivo histology of the myelin g-ratio with magnetic resonance imaging** *NeuroImage* **118**:397-405. http://dx.doi.org/10.1016/j.neuroimage.2015.05.023

Strauss E, Sherman EM, Spreen O (2006) **A compendium of neuropsychological tests: administration, commentary and norms**, 3rd ed. Oxford University Press, New York.

Svoboda E, McKinnon MC, Levine B (2006) **The functional neuroanatomy of autobiographical memory: A meta-analysis** *Neuropsychologia* **44**:2189-2208. http://doi.org/10.1016/j.neuropsychologia.2006.05.023

Tabelow K, Balteau E, Ashburner J, Callaghan MF, Draganski B, Helms G, Kherif F, Leutritz T, Lutti A, Phillips C, Reimer E, Ruthotto L, Seif M, Weiskopf N, Ziegler G, Mohammadi S (2019) **hMRI – A toolbox for quantitative MRI in neuroscience and clinical research** *NeuroImage* **194**:191-210. http://dx.doi.org/10.1016/j.neuroimage.2019.01.029

Tsivilis D, Vann SD, Denby C, Roberts N, Mayes AR, Montaldi D, Aggleton JP (2008) **A disproportionate role for the fornix and mammillary bodies in recall versus recognition memory** *Nature Neuroscience* **11**:834-842. http://dx.doi.org/10.1038/nn.2149

Tulving E (2002) **Episodic memory: From mind to brain** *Annual Review of Psychology* **53**:1-25. http://dx.doi.org/10.1146/annurev.psych.53.100901.135114

Van Petten C (2004) **Relationship between hippocampal volume and memory ability in healthy individuals across the lifespan: review and meta-analysis** *Neuropsychologia* **42**:1394-1413. http://dx.doi.org/10.1016/j.neuropsychologia.2004.04.006

Vann SD, Aggleton JP, Maguire EA (2009) **What does the retrosplenial cortex do?** *Nature Reviews: Neuroscience* **10**:792-802. http://dx.doi.org/10.1038/nrn2733

Veraart J, Nunes D, Rudrapatna U, Fieremans E, Jones DK, Novikov DS, Shemesh N (2020) **Noninvasive quantification of axon radii using diffusion MRI** *eLife* **9**:e49855. http://dx.doi.org/10.7554/eLife.49855

Von Der Heide RJ, Skipper LM, Klobusicky E, Olson IR (2013) **Dissecting the uncinate fasciculus: disorders, controversies and a hypothesis** *Brain* **136**:1692-1707. http://dx.doi.org/10.1093/brain/awt094

Warrington EK (1984) **Recognition Memory Test: Manual.** NFER-Nelson, Berkshire, UK.

Waxman SG (1980) **Determinants of conduction velocity in myelinated nerve fibers** *Muscle & Nerve* **3**:141-150. http://dx.doi.org/10.1002/mus.880030207

Wechsler D (2009) **WMS-IV.: Wechsler memory scale-Administration and scoring manual**. Pearson Assessment, London, UK.

Weiskopf N, Suckling J, Williams G, Correia M, Inkster B, Tait R, Ooi C, Bullmore E, Lutti A (2013) **Quantitative multi-parameter mapping of R1, PD*, MT, and R2* at 3T: a multi-center validation** *Frontiers in Neuroscience* **7**. http://dx.doi.org/10.3389/fnins.2013.00095

West KL, Kelm ND, Carson RP, Does MD (2016) **A revised model for estimating g-ratio from MRI** *NeuroImage* **125**:1155-1158. http://dx.doi.org/10.1016/j.neuroimage.2015.08.017

West KL, Kelm ND, Carson RP, Alexander DC, Gochberg DF, Does MD (2018) **Experimental studies of g-ratio MRI in ex vivo mouse brain** *NeuroImage* **167**:366-371. http://dx.doi.org/10.1016/j.neuroimage.2017.11.064

Winocur G, Moscovitch M (2011) **Memory transformation and systems consolidation** *Journal of the International Neuropsychological Society* **17**:766-780. http://dx.doi.org/10.1017/s1355617711000683

Woollett K, Maguire EA (2011) **Acquiring "the Knowledge" of London's layout drives structural brain changes** *Current Biology* **21**:2109-2114. http://dx.doi.org/10.1016/j.cub.2011.11.018

Xi MC, Liu RH, Engelhardt JK, Morales FR, Chase MH (1999) **Changes in the axonal**
conduction velocity of pyramidal tract neurons in the aged cat *Neuroscience*
92:219-225. http://dx.doi.org/10.1016/S0306-4522(98)00754-4
Xin W, Chan JR (2020) **Myelin plasticity: sculpting circuits in learning and memory** *Nature*
Reviews Neuroscience **21**:682-694. http://dx.doi.org/10.1038/s41583-020-00379-8
Xu J, Moeller S, Auerbach EJ, Strupp J, Smith SM, Feinberg DA, Yacoub E, Uğurbil K (2013)
Evaluation of slice accelerations using multiband echo planar imaging at 3T
NeuroImage **83**:991-1001. http://dx.doi.org/10.1016/j.neuroimage.2013.07.055
Yu F, Fan Q, Tian Q, Ngamsombat C, Machado N, Bireley JD, Russo AW, Nummenmaa A,
Witzel T, Wald LL, Klawiter EC, Huang SY (2019) **Imaging g-ratio in multiple**
sclerosis using high-gradient diffusion MRI and macromolecular tissue volume
American Journal of Neuroradiology **40**:1871. http://dx.doi.org/10.3174/ajnr.A6283
Zhang H, Schneider T, Wheeler-Kingshott CA, Alexander DC (2012) **NODDI: Practical in**
vivo neurite orientation dispersion and density imaging of the human brain
NeuroImage **61**:1000-1016. http://dx.doi.org/10.1016/j.neuroimage.2012.03.072

1386

Appendix 1

- **Details of the simulation performed for Figure 2A.**

- **Investigation of the MR g-ratio, magnetisation transfer saturation values, neurite orientation dispersion index, neurite density and standard DTI parameters in the fornix and uncinate fasciculus.**

- **Investigation of associations between the MR g-ratio and magnetisation transfer saturation values of the fornix and uncinate fasciculus and the laboratory-based memory tests.**

- **Exploratory analyses of the MR g-ratio, magnetisation transfer saturation values, neurite orientation dispersion index, neurite density and standard DTI parameters in other white matter tracts.**

- **Investigation of the standard DTI parameters extracted from the parahippocampal cingulum bundle.**

- **Investigation of the parahippocampal cingulum bundle MR g-ratio and magnetisation transfer saturation values and the laboratory-based memory tests.**

Details of the simulation performed for Figure 2A

In Figure 2A, inner axon diameter, myelin thickness and g-ratio are plotted together. The range of g-ratio values used in the simulation spanned two standard deviations about the mean MR g-ratio observed for the parahippocampal cingulum bundle in the current study (mean = 0.647, standard deviation = 0.043). The axon diameter was computed for this MR g-ratio range by re-arranging the equation presented in Berman et al. (2019) such that axon diameter = $\exp(\frac{\text{g-ratio}-0.506}{0.22})$. Fibre dimeter was then calculated as fibre diameter = $\frac{\text{axon diamter}}{\text{g-ratio}}$, enabling myelin thickness to be computed as myelin thickness = $\frac{\text{fibre diameter}-\text{axon diameter}}{2}$.

We note that discrepancies between reported microscopic parameters (i.e. g-ratio, modelled axon diameter and myelin thickness) derived from in vivo and ex vivo histology may arise for two reasons. (1) The in vivo MR g-ratio is computed from volume-fractions unlike the microscopic g-ratio measured with histology. (2) The heuristic equation by Berman et al. (2019) that is relating the in vivo MR g-ratio to axon diameter is rather capturing the tail of the axon radii distribution.

1427 **Investigation of the MR g-ratio, magnetisation transfer saturation values,**
1428 **neurite orientation dispersion index, neurite density and standard DTI**
1429 **parameters in the fornix and uncinate fasciculus**
1430

1431 As can be observed in the tables below, there were no significant correlations between

1432 microstructural measures or standard DTI parameters and autobiographical memory recall

1433 ability for either of the tracts when using the corrected ($p < 0.017$) threshold.

1434

1435 **Fornix**

1436 The mean number of voxels in the region of interest (ROI) was 249.55 (SD = 23.09).

1437
1438 **Appendix 1–figure 1.** The location of the fornix.
1439

1440 **Appendix 1–table 1.** Means and standard deviations for the microstructure measures and
1441 standard DTI parameters extracted from the fornix.
1442

Measure	Mean	Standard Deviation
MR g-ratio	0.720	0.017
Magnetisation transfer saturation	0.973	0.003
Neurite dispersion (ODI)	0.143	0.021
Neurite density	0.601	0.045
Fractional anisotropy	0.605	0.035
Mean diffusivity (10^{-3} mm^2/s)	0.901	0.036
Mean kurtosis	0.909	0.100
Diffusivities parallel (10^{-3} mm^2/s)	1.645	0.072
Diffusivities perpendicular (10^{-3} mm^2/s)	0.529	0.042

1443 Note. ODI = Orientation Dispersion Index

1444 **Appendix 1–table 2.** Partial correlations between the microstructure measures or standard DTI
1445 parameters extracted from the fornix and autobiographical memory recall ability (internal
1446 details).
1447

Measure	r(211)	p	95% Confidence Interval Lower	Upper
MR g-ratio	-0.04	0.53	-0.16	0.08
Magnetisation transfer saturation	-0.08	0.23	-0.20	0.03
Neurite dispersion (ODI)	-0.03	0.62	-0.16	0.10
Neurite density	-0.03	0.69	-0.15	0.10
Fractional anisotropy	0.02	0.80	-0.12	0.15
Mean diffusivity	0.07	0.35	-0.08	0.21
Mean kurtosis	-0.03	0.66	-0.16	0.10
Diffusivities parallel	0.05	0.46	-0.09	0.19
Diffusivities perpendicular	0.02	0.74	-0.12	0.15

1448 Note. ODI = Orientation Dispersion Index

1449

1450

1451

Uncinate fasciculus

The mean number of voxels in the ROI was 191.98 (SD = 29.77).

Appendix 1–figure 2. The location of the uncinate fasciculus.

Appendix 1–table 3. Means and standard deviations for the microstructure measures and standard DTI parameters extracted from the uncinate fasciculus.

Measure	Mean	Standard Deviation
MR g-ratio	0.723	0.016
Magnetisation transfer saturation	0.972	0.003
Neurite dispersion (ODI)	0.189	0.022
Neurite density	0.562	0.045
Fractional anisotropy	0.512	0.036
Mean diffusivity (10^{-3} mm^2/s)	0.877	0.304
Mean kurtosis	0.913	0.100
Diffusivities parallel (10^{-3} mm^2/s)	1.456	0.055
Diffusivities perpendicular (10^{-3} mm^2/s)	0.588	0.039

Note. ODI = Orientation Dispersion Index

1466 **Appendix 1–table 4.** Partial correlations between the microstructure measures or standard DTI
1467 parameters extracted from the uncinate fasciculus and autobiographical memory recall ability
1468 (internal details).
1469

Measure	r(211)	p	95% Confidence Interval	
			Lower	Upper
MR g-ratio	0.10	0.15	-0.03	0.22
Magnetisation transfer saturation	-0.02	0.81	-0.14	0.11
Neurite dispersion (ODI)	-0.01	0.94	-0.14	0.13
Neurite density	0.01	0.89	-0.14	0.16
Fractional anisotropy	-0.01	0.94	-0.14	0.13
Mean diffusivity	-0.01	0.87	-0.14	0.13
Mean kurtosis	-0.01	0.91	-0.13	0.18
Diffusivities parallel	-0.01	0.87	-0.14	0.13
Diffusivities perpendicular	0.00	0.96	-0.15	0.14

1470 Note. ODI = Orientation Dispersion Index

1471

Investigation of associations between the MR g-ratio and magnetisation transfer saturation values of the fornix and uncinate fasciculus and the laboratory-based memory tests

As can be observed in the tables below, there were no significant correlations between either

the MR g-ratio or magnetisation transfer saturation values of the fornix or uncinate fasciculus

and any of the laboratory-based memory tasks for either of the tracts, even when using a p <

0.05 uncorrected threshold.

Appendix 1–table 5. Partial correlations between the fornix MR g-ratio and magnetisation transfer saturation values and the laboratory-based memory tests.

Measure	r(211)	p	95% Confidence Interval	
			Lower	Upper
MR g-ratio				
Logical Memory immediate recall	-0.01	0.85	-0.14	0.11
Logical Memory delayed recall	-0.07	0.33	-0.19	0.06
RAVLT immediate recall	-0.06	0.42	-0.20	0.09
RAVLT delayed recall	0.00	0.98	-0.14	0.15
Rey-Osterrieth Complex Figure delayed recall	0.10	0.16	-0.04	0.23
Warrington RMT for Words	0.04	0.59	-0.08	0.16
Warrington RMT for Faces	-0.05	0.52	-0.17	0.09
Dead or Alive Test	-0.05	0.50	-0.16	0.09
Magnetisation transfer saturation				
Logical Memory immediate recall	0.04	0.53	-0.11	0.19
Logical Memory delayed recall	0.11	0.11	-0.04	0.26
RAVLT immediate recall	0.01	0.92	-0.11	0.13
RAVLT delayed recall	0.03	0.71	-0.09	0.15
Rey-Osterrieth Complex Figure delayed recall	0.07	0.33	-0.06	0.19
Warrington RMT for Words	0.06	0.41	-0.05	0.17
Warrington RMT for Faces	0.04	0.61	-0.09	0.16
Dead or Alive Test	0.08	0.25	-0.06	0.21

Note. RAVLT = Rey Auditory Verbal Learning Test; RMT = Recognition Memory Test

1486 **Appendix 1–table 6.** Partial correlations between the uncinate fasciculus MR g-ratio and
1487 magnetisation transfer saturation values and the laboratory-based memory tests.
1488

Measure	r(211)	p	95% Confidence Interval	
			Lower	Upper
MR g-ratio				
Logical Memory immediate recall	0.07	0.34	-0.06	0.19
Logical Memory delayed recall	0.04	0.34	-0.09	0.17
RAVLT immediate recall	0.00	0.98	-0.14	0.13
RAVLT delayed recall	0.06	0.40	-0.10	0.21
Rey-Osterrieth Complex Figure delayed recall	0.01	0.94	-0.13	0.15
Warrington RMT for Words	-0.01	0.86	-0.16	0.15
Warrington RMT for Faces	-0.07	0.34	-0.20	0.06
Dead or Alive Test	-0.04	0.60	-0.15	0.08
Magnetisation transfer saturation				
Logical Memory immediate recall	-0.04	0.53	-0.17	0.09
Logical Memory delayed recall	-0.05	0.49	-0.18	0.09
RAVLT immediate recall	-0.13	0.06	-0.26	0.01
RAVLT delayed recall	-0.16	0.02	-0.27	-0.03
Rey-Osterrieth Complex Figure delayed recall	-0.07	0.33	-0.20	0.07
Warrington RMT for Words	-0.14	0.04	-0.26	-0.01
Warrington RMT for Faces	-0.05	0.44	-0.18	0.09
Dead or Alive Test	-0.01	0.93	-0.14	0.13

1489 Note. RAVLT = Rey Auditory Verbal Learning Test; RMT = Recognition Memory Test

1490
1491

Exploratory analyses of the MR g-ratio, magnetisation transfer saturation values, neurite orientation dispersion index, neurite density and standard DTI parameters in other white matter tracts

As can be observed in the tables below, there were no significant correlations between microstructural measures or standard DTI parameters and autobiographical memory recall ability for any of the tracts when using the corrected ($p < 0.017$) threshold.

Anterior thalamic radiation

The mean number of voxels in the ROI was 2090.21 ($SD = 67.43$).

Appendix 1–figure 3. The location of the anterior thalamic radiation.

Appendix 1–table 7. Means and standard deviations for the microstructure measures and standard DTI parameters extracted from the anterior thalamic radiation.

Measure	Mean	Standard Deviation
MR g-ratio	0.724	0.014
Magnetisation transfer saturation	0.986	0.001
Neurite dispersion (ODI)	0.252	0.017
Neurite density	0.598	0.041
Fractional anisotropy	0.430	0.028
Mean diffusivity (10^{-3} mm^2/s)	0.861	0.029
Mean kurtosis	0.986	0.063
Diffusivities parallel (10^{-3} mm^2/s)	1.328	0.041
Diffusivities perpendicular (10^{-3} mm^2/s)	0.627	0.034

Note. ODI = Orientation Dispersion Index

1509 **Appendix 1–table 8.** Partial correlations between the microstructure measures or standard DTI
1510 parameters extracted from the anterior thalamic radiation and autobiographical memory recall
1511 ability (internal details).
1512

Measure	r(211)	p	95% Confidence Interval	
			Lower	Upper
MR g-ratio	0.05	0.47	-0.06	0.17
MT sat	-0.08	0.26	-0.20	0.04
Neurite dispersion (ODI)	-0.05	0.46	-0.18	0.09
Neurite density	0.10	0.16	-0.04	0.23
Fractional anisotropy	0.08	0.26	-0.06	0.21
Mean diffusivity	0.03	0.67	-0.11	0.17
Mean kurtosis	0.07	0.28	-0.07	0.22
Diffusivities parallel	0.09	0.20	-0.04	0.21
Diffusivities perpendicular	-0.02	0.77	-0.16	0.12

1513 Note. MT sat = Magnetisation Transfer saturation; ODI = Orientation Dispersion Index

1514

1515 **Dorsal cingulum bundle**

1516 The mean number of voxels in the ROI was 611.48 (*SD* = 25.29).

1517

1518
1519 **Appendix 1–figure 4.** The location of the dorsal cingulum bundle.
1520

1521

1522 **Appendix 1–table 9.** Means and standard deviations for the microstructure measures and
1523 standard DTI parameters extracted from the dorsal cingulum bundle.
1524

Measure	Mean	Standard Deviation
MR g-ratio	0.710	0.016
Magnetisation transfer saturation	0.984	0.003
Neurite dispersion (ODI)	0.147	0.019
Neurite density	0.560	0.038
Fractional anisotropy	0.570	0.039
Mean diffusivity (10^{-3} mm^2/s)	0.862	0.026
Mean kurtosis	0.862	0.093
Diffusivities parallel (10^{-3} mm^2/s)	1.531	0.061
Diffusivities perpendicular (10^{-3} mm^2/s)	0.527	0.040

1525 Note. ODI = Orientation Dispersion Index

1526

1527

1528

1529 **Appendix 1–table 10.** Partial correlations between the microstructure measures or standard
1530 DTI parameters extracted from the dorsal cingulum bundle and autobiographical memory recall
1531 ability (internal details).
1532

Measure	r(211)	p	95% Confidence Interval	
			Lower	Upper
MR g-ratio	0.09	0.22	-0.04	0.21
MT sat	0.08	0.23	-0.04	0.21
Neurite dispersion (ODI)	0.09	0.17	-0.05	0.23
Neurite density	0.09	0.17	-0.05	0.23
Fractional anisotropy	0.01	0.85	-0.13	0.16
Mean diffusivity	0.06	0.43	-0.08	0.19
Mean kurtosis	0.13	0.05	0.00	0.27
Diffusivities parallel	0.05	0.48	-0.08	0.18
Diffusivities perpendicular	0.01	0.91	-0.14	0.15

1533 Note. MT sat = Magnetisation Transfer saturation; ODI = Orientation Dispersion Index

1534

1535

1536 **Forceps minor**

1537 The mean number of voxels in the ROI was 4613.67 (*SD* = 78.02).

1538
1539 **Appendix 1–figure 5.** The location of the forceps minor.
1540

1541

1542 **Appendix 1–table 11.** Means and standard deviations for the microstructure measures and
1543 standard DTI parameters extracted from the forceps minor.
1544

Measure	Mean	Standard Deviation
MR g-ratio	0.698	0.026
Magnetisation transfer saturation	0.993	0.002
Neurite dispersion (ODI)	0.201	0.016
Neurite density	0.601	0.042
Fractional anisotropy	0.500	0.027
Mean diffusivity (10^{-3} mm^2/s)	0.881	0.031
Mean kurtosis	0.947	0.128
Diffusivities parallel (10^{-3} mm^2/s)	1.467	0.052
Diffusivities perpendicular (10^{-3} mm^2/s)	0.588	0.034

1545 Note. ODI = Orientation Dispersion Index

1546

1547

1548

1549

1550 **Appendix 1–table 12.** Partial correlations between the microstructure measures or standard
1551 DTI parameters extracted from the forceps minor and autobiographical memory recall ability
1552 (internal details).
1553

Measure	r(211)	p	95% Confidence Interval	
			Lower	Upper
MR g-ratio	0.04	0.53	-0.07	0.18
MT sat	0.07	0.34	-0.06	0.19
Neurite dispersion (ODI)	-0.06	0.38	-0.21	0.07
Neurite density	0.09	0.19	-0.04	0.22
Fractional anisotropy	0.10	0.17	-0.03	0.23
Mean diffusivity	-0.04	0.56	-0.17	0.09
Mean kurtosis	0.00	0.97	-0.11	0.21
Diffusivities parallel	0.03	0.67	-0.09	0.18
Diffusivities perpendicular	-0.07	0.32	-0.19	0.05

1554 Note. MT sat = Magnetisation Transfer saturation; ODI = Orientation Dispersion Index

1555

1556

1557 **Inferior longitudinal fasciculus**

1558 The mean number of voxels in the ROI was 2844.03 (*SD* = 52.97).

1559
1560 **Appendix 1–figure 6.** The location of the inferior longitudinal fasciculus.
1561

1562

1563 **Appendix 1–table 13.** Means and standard deviations for the microstructure measures and
1564 standard DTI parameters extracted from the inferior longitudinal fasciculus.
1565

Measure	Mean	Standard Deviation
MR g-ratio	0.724	0.013
Magnetisation transfer saturation	0.994	0.002
Neurite dispersion (ODI)	0.184	0.017
Neurite density	0.558	0.041
Fractional anisotropy	0.486	0.028
Mean diffusivity (10^{-3} mm^2/s)	0.902	0.029
Mean kurtosis	0.918	0.061
Diffusivities parallel (10^{-3} mm^2/s)	1.473	0.045
Diffusivities perpendicular (10^{-3} mm^2/s)	0.616	0.035

1566 Note. ODI = Orientation Dispersion Index

1567

1568

1569

1570

1571 **Appendix 1–table 14.** Partial correlations between the microstructure measures or standard
1572 DTI parameters extracted from the inferior longitudinal fasciculus and autobiographical
1573 memory recall ability (internal details).
1574

| Measure | r(211) | p | 95% Confidence Interval | |
			Lower	Upper
MR g-ratio	0.07	0.35	-0.06	0.19
MT sat	0.07	0.35	-0.05	0.18
Neurite dispersion (ODI)	-0.03	0.64	-0.17	0.10
Neurite density	0.07	0.35	-0.07	0.19
Fractional anisotropy	0.10	0.14	-0.04	0.23
Mean diffusivity	-0.01	0.87	-0.15	0.13
Mean kurtosis	0.10	0.16	-0.03	0.23
Diffusivities parallel	0.07	0.29	-0.07	0.21
Diffusivities perpendicular	-0.07	0.34	-0.19	0.07

1575 Note. MT sat = Magnetisation Transfer saturation; ODI = Orientation Dispersion Index

1576

Inferior occipitofrontal fasciculus

The mean number of voxels in the ROI was 3344.31 (*SD* = 46.0).

Appendix 1–figure 7. The location of the inferior occipitofrontal fasciculus.

Appendix 1–table 15. Means and standard deviations for the microstructure measures and standard DTI parameters extracted from the inferior occipitofrontal fasciculus.

Measure	Mean	Standard Deviation
MR g-ratio	0.729	0.011
Magnetisation transfer saturation	0.995	0.001
Neurite dispersion (ODI)	0.179	0.012
Neurite density	0.563	0.036
Fractional anisotropy	0.508	0.024
Mean diffusivity (10^{-3} mm^2/s)	0.889	0.026
Mean kurtosis	0.912	0.057
Diffusivities parallel (10^{-3} mm^2/s)	1.488	0.039
Diffusivities perpendicular (10^{-3} mm^2/s)	0.590	0.031

Note. ODI = Orientation Dispersion Index

1591 **Appendix 1–table 16.** Partial correlations between the microstructure measures or standard
1592 DTI parameters extracted from the inferior occipitofrontal fasciculus and autobiographical
1593 memory recall ability (internal details).
1594

| Measure | r(211) | p | 95% Confidence Interval | |
			Lower	Upper
MR g-ratio	-0.01	0.93	-0.12	0.12
MT sat	0.08	0.22	-0.03	0.19
Neurite dispersion (ODI)	-0.01	0.89	-0.14	0.13
Neurite density	0.06	0.42	-0.07	0.18
Fractional anisotropy	0.06	0.38	-0.08	0.20
Mean diffusivity	0.00	0.99	-0.13	0.13
Mean kurtosis	0.07	0.31	-0.06	0.19
Diffusivities parallel	0.05	0.50	-0.09	0.18
Diffusivities perpendicular	-0.03	0.62	-0.17	0.10

1595 Note. MT sat = Magnetisation Transfer saturation; ODI = Orientation Dispersion Index

1596

1597

Superior longitudinal fasciculus

The mean number of voxels in the ROI was 4243.22 (*SD* = 39.80).

Appendix 1–figure 8. The location of the superior longitudinal fasciculus.

Appendix 1–table 17. Means and standard deviations for the microstructure measures and standard DTI parameters extracted from the superior longitudinal fasciculus.

Measure	Mean	Standard Deviation
MR g-ratio	0.751	0.011
Magnetisation transfer saturation	0.996	0.001
Neurite dispersion (ODI)	0.218	0.012
Neurite density	0.634	0.034
Fractional anisotropy	0.471	0.026
Mean diffusivity (10^{-3} mm^2/s)	0.825	0.025
Mean kurtosis	1.029	0.041
Diffusivities parallel (10^{-3} mm^2/s)	1.324	0.035
Diffusivities perpendicular (10^{-3} mm^2/s)	0.575	0.041

Note. ODI = Orientation Dispersion Index

1611 **Appendix 1–table 18.** Partial correlations between microstructure measures or standard DTI
1612 parameters extracted from the superior longitudinal fasciculus and autobiographical memory
1613 recall ability (internal details).

1614

Measure	r(211)	p	95% Confidence Interval	
			Lower	Upper
MR g-ratio	0.08	0.28	-0.05	0.20
MT sat	-0.04	0.54	-0.16	0.08
Neurite dispersion (ODI)	-0.09	0.17	-0.23	0.05
Neurite density	0.12	0.09	-0.01	0.24
Fractional anisotropy	0.13	0.05	0.01	0.26
Mean diffusivity	-0.02	0.80	-0.14	0.11
Mean kurtosis	0.11	0.12	-0.03	0.24
Diffusivities parallel	0.12	0.09	-0.03	0.25
Diffusivities perpendicular	-0.08	0.23	-0.20	0.03

1615 Note. MT sat = Magnetisation Transfer saturation; ODI = Orientation Dispersion Index

1616

 Investigation of the standard DTI parameters extracted from the
 parahippocampal cingulum bundle

1619

1620   As reported in the main text, significant correlations between a number of microstructural

1621   measures from the parahippocampal cingulum bundle and autobiographical memory recall

1622   ability were evident. This was also the case for several of the standard DTI parameters – see

1623   Appendix 1–tables 19 and 20 and Appendix 1–figures 9 and 10 below.

1624

1625   **Appendix 1–table 19.** Means and standard deviations for the standard DTI parameters
1626   extracted from the parahippocampal cingulum bundle.
1627

Measure	Mean	Standard Deviation
Fractional anisotropy	0.466	0.053
Mean diffusivity (10^{-3} mm^2/s)	0.931	0.041
Mean kurtosis	0.779	0.122
Diffusivities parallel (10^{-3} mm^2/s)	1.479	0.071
Diffusivities perpendicular (10^{-3} mm^2/s)	0.656	0.057

1628
1629   **Appendix 1–table 20.** Partial correlations between the standard DTI parameters extracted from
1630   the parahippocampal cingulum bundle and autobiographical memory recall ability (internal
1631   details).
1632

Measure	r(211)	p	95% Confidence Interval	
			Lower	Upper
Fractional anisotropy	0.20	0.003[*]	0.07	0.32
Mean diffusivity	-0.02	0.72	-0.15	0.11
Mean kurtosis	0.08	0.23	-0.05	0.21
Diffusivities parallel	0.19	0.005[*]	0.06	0.32
Diffusivities perpendicular	0.15	0.03	-0.27	-0.02

1633   [*]$p < 0.017$ (two-sided Bonferroni corrected threshold)
1634

Specifically, partial correlation analyses, with age, gender, scanner and the number of

voxels in the ROI included as covariates revealed a significant positive correlation between

internal details and fractional anisotropy (FA) (Appendix 1–figure 9A; $r(211) = 0.20$, $p = 0.003$,

95% CI = 0.07, 0.32). This relationship was specific to internal details, and was not evident for

the external details control measure (Appendix 1–figure 9B; $r(211)$ = -0.06, p = 0.39, 95% CI

= -0.19, 0.07). Direct comparison of the two correlations confirmed a significant difference

between them, showing that parahippocampal cingulum bundle FA was related to internal

details to a greater extent than external details (Appendix 1–figure 9C; mean r difference =

0.26 (95% CI = 0.10, 0.44), z = 3.08, p = 0.002).

1644

1645
Appendix 1–figure 9. Fractional anisotropy (FA) and the parahippocampal cingulum bundle.
The relationships between parahippocampal cingulum bundle FA and autobiographical
memory recall ability (internal details), and the control measure (external details) are shown.
(A) There was a significant positive correlation between FA and internal details (dashed lines
indicate the confidence intervals). **(B)** There was no significant relationship between FA and
external details. **(C)** Bar chart showing the partial correlation coefficients (with standard errors)
between FA and internal and external details. There was a significant difference between the
correlations when they were directly compared; **p < 0.01. Data points for this figure are
provided in Supplementary file 2.
1655

In addition, a significant positive correlation between diffusivities parallel and the

number of internal details was also apparent (Appendix 1–figure 10A; $r(211) = 0.19$, $p = 0.005$,

95% CI = 0.06, 0.32). As with FA, no significant relationship was observed between

diffusivities parallel and external details (Appendix 1–figure 10B; $r(211) = -0.048$, $p = 0.49$,

95% CI = -0.19, 0.10). Direct comparison of the correlations confirmed that diffusivities

parallel was related to internal details to a greater extent than external details (Appendix 1–

figure 10C; mean r difference = 0.24 (95% CI = 0.07, 0.41), $z = 2.81$, $p = 0.0049$).

1663
Appendix 1–figure 10. Dpara and the parahippocampal cingulum bundle. The relationships
between parahippocampal cingulum bundle diffusivities parallel (Dpara) and autobiographical
memory recall ability (internal details), and the control measure (external details) are shown.
(A) There was a significant positive correlation between Dpara and internal details (dashed
lines indicate the confidence intervals). **(B)** There was no significant relationship between
Dpara and external details. **(C)** Bar chart showing the partial correlation coefficients (with
standard errors) between Dpara and internal and external details. There was a significant
difference between the correlations when they were directly compared; **$p < 0.01$. Data points
for this figure are provided in Supplementary file 2.

In contrast, no relationships were observed when examining the partial correlations

between either internal or external details and mean diffusivity (internal: $r(211) = -0.02$, $p =$

0.72, 95% CI = -0.15, 0.11; external: $r(211) = 0.01$, $p = 0.86$, 95% CI = -0.11, 0.14), the mean

kurtosis (internal: $r(211) = 0.08$, $p = 0.23$, 95% CI = -0.05, 0.21; external: $r(211) = 0.03$, $p =$

0.62, 95% CI = -0.01, 0.16) or diffusivities perpendicular (internal: $r(211) = -0.15$, $p = 0.03$,

95% CI = -0.27, -0.02; external: $r(211) = 0.44$, $p = 0.52$, 95% CI = -0.08, 0.17), when using

the corrected ($p < 0.017$) threshold. This suggests that none of these parameters were strongly

associated with individual differences in autobiographical memory recall.

1681

1682 **Investigation of the parahippocampal cingulum bundle MR g-ratio and**

1683 **magnetisation transfer saturation values and the laboratory-based memory**

1684 **tests**

1685
1686 Investigation of associations between the MR g-ratio and magnetisation transfer saturation

1687 values and the eight laboratory-based memory tests was also undertaken. No relationships were

1688 evident for any test, even when using an uncorrected p < 0.05 threshold, see Appendix 1-table

1689 21.

1690

1691 **Appendix 1–table 21.** Partial correlations between the parahippocampal cingulum bundle MR
1692 g-ratio and magnetisation transfer saturation values and the laboratory-based memory tests.
1693

Measure	r(211)	p	95% Confidence Interval	
			Lower	Upper
MR g-ratio				
Logical Memory immediate recall	0.01	0.93	-0.13	0.13
Logical Memory delayed recall	-0.05	0.46	-0.18	0.08
RAVLT immediate recall	-0.06	0.41	-0.18	0.07
RAVLT delayed recall	-0.09	0.17	-0.21	0.04
Rey-Osterrieth Complex Figure delayed recall	0.07	0.34	-0.06	0.19
Warrington RMT for Words	0.02	0.80	-0.11	0.16
Warrington RMT for Faces	-0.04	0.53	-0.17	0.10
Dead or Alive Test	0.02	0.79	-0.11	0.15
Magnetisation transfer saturation				
Logical Memory immediate recall	0.07	0.35	-0.06	0.19
Logical Memory delayed recall	0.01	0.91	-0.12	0.14
RAVLT immediate recall	-0.01	0.90	-0.14	0.14
RAVLT delayed recall	-0.02	0.80	-0.14	0.11
Rey-Osterrieth Complex Figure delayed recall	-0.09	0.21	-0.23	0.05
Warrington RMT for Words	-0.09	0.21	-0.22	0.04
Warrington RMT for Faces	-0.05	0.49	-0.17	0.09
Dead or Alive Test	0.11	0.11	-0.01	0.23

1694 Note. RAVLT = Rey Auditory Verbal Learning Test; RMT = Recognition Memory Test