Secreted dengue virus NS1 from infection is predominantly dimeric and in complex with high-density lipoprotein

Bing Liang Alvin Chew[1,2]†, An Qi Ngoh [3]†, Wint Wint Phoo[4]†, Kitti Wing Ki Chan[3]†, Zheng Ser[4], Nikhil K Tulsian[5,6], Shiao See Lim [3], Mei Jie Grace Weng[1,2], Satoru Watanabe[3], Milly M. Choy[3], Jenny G. Low[3,8], Eng Eong Ooi[3,9,10], Christiane Ruedl[11], Radoslaw M. Sobota[4], Subhash G. Vasudevan[3,7,12]*, Dahai Luo[1,2]*

Affiliations:

[1]Lee Kong Chian School of Medicine, Nanyang Technological University; EMB 03-07, 59 Nanyang Drive, Singapore 636921

[2]NTU Institute of Structural Biology, Nanyang Technological University; EMB 06-01, 59 Nanyang Drive, Singapore 636921, Singapore

[3]Program in Emerging Infectious Diseases, Duke-NUS Medical School; 8 College Road, Singapore 169857, Singapore

[4]Functional Proteomics Laboratory, Institute of Molecular and Cell Biology, Agency for Science, Technology and Research (A*STAR); 61 Biopolis Drive, Proteos, Singapore 138673, Singapore

[5]Department of Biological Sciences, National University of Singapore, Singapore,

[6]Singapore Centre for Life Sciences, Department of Biochemistry, National University of Singapore, Singapore, Singapore

[7]Department of Microbiology and Immunology, National University of Singapore; 5 Science Drive 2, Singapore 117545, Singapore

[8]Department of Infectious Diseases, Singapore General Hospital, Singapore, Singapore.

[9]Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore.

[10]Saw Swee Hock School of Public Health, National University of Singapore, Singapore, Singapore.

[11]School of Biological Sciences, Nanyang Technological University, Singapore, Singapore

[12]Institute for Glycomics (G26), Griffith University Gold Coast Campus, QLD 4222, Australia.

† These four authors contributed equally to the work.

*Corresponding author. Email: subhash.vasudevan@duke-nus.edu.sg (S.V.); luodahai@ntu.edu.sg (D.L.)

Abstract

Severe dengue infections are characterized by endothelial dysfunction shown to be associated with the secreted nonstructural protein 1 (sNS1), making it an attractive vaccine antigen and biotherapeutic target. To uncover the biologically relevant structure of sNS1, we obtained infection-derived sNS1 (isNS1) from DENV-infected Vero cells through immunoffinity purification instead of recombinant sNS1 (rsNS1) overexpressed in insect or mammalian cell lines. We found that isNS1 appeared as an approximately 250 kDa complex of NS1 and ApoA1 and further determined the cryoEM structures of isNS1 and its complex with a monoclonal antibody/Fab. Indeed, we found that the major species of isNS1 is a complex of the NS1 dimer partially embedded in a High-Density Lipoprotein (HDL) particle. Cross-linking mass spectrometry (XL-MS) studies confirmed that the isNS1 interacts with the major HDL component ApoA1 through interactions that map to the NS1 wing and hydrophobic domains. Furthermore, our studies demonstrated that the sNS1 in sera from DENV-infected mice and a human patient form a similar complex as isNS1. Our results report the molecular architecture of a biological form of sNS1 which may have implications for the molecular pathogenesis of dengue.

Summary

CryoEM structures of secreted dengue virus NS1 protein reveal dimers in complex with high-density lipoprotein.

Introduction

Dengue virus (DENV) is a member of the flavivirus genus and is the cause of significant healthcare problems and economic burden worldwide, partially due to the lack of effective therapeutics and the limited efficacy of licensed vaccines, Dengvaxia® and QDENGA®[1,2]. While the majority of DENV infections are mild or asymptomatic, severe dengue infections, marked by vascular leakage, can be life-threatening and even fatal. The severity of dengue infections has been demonstrated to be correlated to high sera levels of sNS1 in clinical studies[3,4]. The viral nonstructural protein 1 (NS1) is a highly conserved and multifunctional protein that exists as intracellular membrane-associated NS1, a presumed GPI-anchored outer plasma membrane-associated NS1, and secreted NS1 (sNS1) during viral infection[5]. NS1 could

interact with a myriad of proteins associated with its immune evasion and virotoxin roles[6]. The interactions of NS1 with viral E protein[7] and the NS4A-2K-NS4B precursor protein[8] have been shown to be important in virus production and replication respectively. Intracellular NS1 found in the ER lumen is an essential part of the membranous viral RNA replication compartment[7]. Upon release from the cells, sNS1 enters the blood circulation where *in vitro* and *in vivo* mouse evidence have demonstrated its ability to induce vascular permeability, a hallmark of severe dengue, either independently[9] or through inducing pro-inflammatory responses[10].

The high-resolution crystal structures of NS1 dimer have been reported for DENV[11], WNV[11], and ZIKV[12,13] which provide a conserved molecular view of flaviviral NS1. Mature NS1 is 352 amino acids long with an apparent molecular mass between 40 – 50 kDa depending on its glycosylation state. NS1 has a three-domain architecture, a hydrophobic β-roll (residues 1-29), an α/β wing (38-151), and a β-ladder (181-352). The connector segments between the wing and β-ladder domains, residues 30-37 and 152-180, form a 3-stranded β-sheet. The dimer has a distinct cross shape with the wings extending from the central β-ladder which has an extended β-sheet that faces the hydrophobic β-roll and a "spaghetti loop" on the opposite hydrophilic outer face that lacks structured elements[11-13]. sNS1 was reported to be a barrel-shaped hexamer with lipid cargo held together by hydrophobic interactions based on biophysical and low-resolution EM analysis[14-16]. Recent cryoEM structures of recombinant sNS1 (rsNS1) from DENV-2 showed mainly tetrameric rsNS1 with only 3% of the population found to be in the hexameric form[17]. While earlier studies by Gutsche et al (2011) found that the lipid profile of DENV-1 infection-derived sNS1 to be similar to high-density lipoprotein (HDL)[15], the same research group now showed that DENV-2 rsNS1 could dock onto HDL with direct visualization using negative stain EM analysis as reported by Benfrid et al. (2022)[18]. Additionally, sNS1 appears to utilize the scavenger receptor B1, a known receptor for HDL, as the cognate receptor in cultured cells[19]. Since the interactions of sNS1 with the target cells implicated in pathogenesis require the hydrophobic β-roll side to be exposed[11], the mechanism by which the oligomeric sNS1 with its lipid load in the central channel dissociates and associates have been a topic of intense research interest[18,20,21]. However, the native structure of the extracellular NS1 during viral infection remains unclear.

To closely mimic of sNS1 circulating in dengue patients' sera, we obtained infection-derived

sNS1 (isNS1) from the culture supernatant of Vero cells infected with either the DENV2 WT

(isNS1wt) or T164S mutant (isNS1ts). The T164S mutation in the greasy finger loop between

the wing and β-ladder interdomain of NS1 was identified from a DENV2 epidemic in Cuba in

1997, where a correlation with enhanced clinical disease severity was observed[22]. We

demonstrated that this single mutation in NS1 could directly cause lethality in mice and

increased sNS1 secretion[23], which served as an epidemiologically relevant tool to study the

biochemical and structural characteristics of sNS1 in this study.

We determined that the isNS1 is a complex of the NS1 dimer embedded on a single HDL

particle composed of ApoA1. By applying integrative structural tools, we report the cryoEM

structural model of the purified isNS1:HDL complex at ~8 Å resolution with protein-protein

interactions between NS1 and ApoA1 defined by crosslinking mass spectrometry (XL-MS).

Furthermore, the Ab56.2 binding site model in the cryoEM model of isNS1:ApoA1 complex was

elucidated using hydrogen deuterium exchange mass spectrometry (HDX-MS) as well as NS1

peptide competition ELISA. Interrogation of DENV-infected mouse sera and a human patient

suggests that the sNS1:HDL complex exists *in vivo* as a similar complex to isNS1.

Results

Native isNS1 is a complex of NS1 with ApoA1

We purified sNS1 using a monoclonal antibody 56.2 from the supernatant of the infected Vero

cell cultures (Fig. 1a and Supplementary Fig. 1a-b). Initial negative stain screening of the

isNS1wt (Supplementary Fig. 1c) showed some 2D classes of NS1 dimers presumably docked

onto a spherical density, mirroring the *in vitro* reconstitution of crsNS1 with HDL complex

negative stain microscopy results by Benfrid et al. (2022)[18]. Both immunoaffinity purified

isNS1wt (Fig. 1b and Supplementary Fig. 2a) and isNS1ts (Supplementary Fig. 2a and 3b)

retained a molecular size of ~250 kDa as detected with Coomassie Blue (Fig. 1b and

Supplementary Fig. 2a) and on a Western blot using Ab56.2 following separation on a Native-

PAGE (Fig. 1b), as previously reported[23]. The isNS1wt migrated slower than the rsNS1 protein

(a gift from Shu et al. 2022[17]), indicating possible conformational or oligomeric differences

between the two NS1 species (Fig. 1b). On a reducing SDS-PAGE, we identified two major

bands of approximately 50 kDa and 25 kDa in isNS1wt, the former corresponding to the single

monomeric NS1 band seen in rsNS1 (Fig. 1c). Preliminary mass ID identified the 25 kDa protein
as bovine apolipoprotein A1 (ApoA1), a major component of HDL in bovine serum-containing
culture media, which has a reported molecular weight of 28 kDa[24]. Hence, the two bands were
further validated on a Western blot, which confirmed the 50 kDa band as NS1 and the 25 kDa
band as ApoA1 respectively (Fig. 1c). The same observations were seen in isNS1ts
(Supplementary Fig. 3).

Fig. 1. Composition of the secreted NS1 from DENV-infected Vero cells. DENV 2 WT cell
culture supernatant was filtered, supplemented with protease inhibitor cocktail and 0.05%
sodium azide, concentrated using a 100 kDa MWCO Vivaflow cassette and purified using 56.2
anti-NS1 antibody immunoaffinity chromatography. The eluted isNS1wt was dialysed against
PBS, concentrated, and stored at -80°C until further use. **(a)** Schematic of isNS1 purification to
illustrate the samples used for gel analyses. % NS1 is measured by the total amount of NS1
(quantified using the anti-NS1 ELISA kit (Bio-rad) as a percentage of total protein (quantified
using the Bradford assay) found in each sample. Details of the % enrichment in NS1 along the
purification process is as shown in Supplementary Fig. S1 B. **(b)** Coomassie blue detection of
proteins from Crude, Wash and Elute immunoaffinity fractions for isNS1wt, with the
recombinant sNS1 (rsNS1) obtained from Shu et al (2022)[17] as a positive control, after

separation on a 10% Native-PAGE gel (left). The Crude and Elute fractions contain 1 µg of total
protein. The Wash fraction contains approximately 100 ng of total protein in maximum well
volume of the gel. The same set of samples were also subjected to a western blot detection of
NS1 using 56.2 anti-NS1 antibody after separation on a 10% Native-PAGE (right). The Crude
and Elute fractions contain 500 ng of total protein. The Wash fraction contains approximately
100 ng of total protein in maximum well volume of the gel. **(c)** Coomassie blue detection of
proteins from Crude, Wash and Elute immunoaffinity fractions for isNS1wt and rsNS1[17], after
separation on a 4-20% reducing SDS-PAGE gel. The Crude and Elute fractions contain 1 µg of
total protein. The Wash fraction contains approximately 100 ng of total protein in maximum well
volume of the gel. Similarly, the same set of samples were also subjected to a western blot
detection of NS1 and ApoA1 using 56.2 anti-NS1 antibody or ApoA1 antibody (Biorbyt,
orb10643) respectively, after separation on a 4-20% reducing SDS-PAGE (right). **(d)** In-gel
protein identification of the purified isNS1wt by liquid chromatography mass spectrometry (LC-
MS). Proportion of NS1, ApoA1 and other unidentified proteins quantified in total ion intensity,
obtained from the following samples: Elute in solution (boxed in blue), 250 kDa gel band (boxed
in purple), 50 kDa gel band (boxed in red) and 25 kDa gel band (green). The boxed gel bands are
from representative gels showing the different protein species found while the actual gel bands
used for protein identification by LC-MS are as shown in Supplementary Fig. 2a-b.
164

Next, we elucidated the composition of the prominent 250 kDa band observed in the Native-

PAGE by label-free quantification (LFQ) analysis of proteins using liquid chromatography–mass

spectrometry (LC-MS). We obtained LC-MS data for the ~250 kDa protein excised from the

Native-PAGE separation of the immunoaffinity purified isNS1wt (represented in Fig. 1b, purple

box as Gel Band 1); an aliquot of the elute fraction from immunoaffinity purification (depicted in

Fig. 1a blue box and referred to as "Elute") and the 50 kDa and 25 kDa bands excised after

separation of the elute fraction on a reducing SDS-PAGE (Fig. 1c, represented by red and green

boxed labelled as Gel Band 2 and 3 respectively) as positive controls. The raw data for the ion

intensities of the samples tested are as detailed in Supplementary File 1. We found that the

combined LFQ intensity of NS1 and ApoA1 accounted for more than 77.6% and 88.6% of the

total ion intensity count (TIC) detected from the elute fraction (Fig. 1a, blue box) and the 250

kDa band (Fig. 1b, the purple box labelled '1') respectively, indicating that NS1 and ApoA1

were the major proteins in the elute fraction and the ~250 kDa band from immunoaffinity

purified isNS1wt (Fig. 1d). Accordingly, as depicted in Fig. 1d, NS1 accounted for 96.85% of

the TIC for the 50 kDa band (Fig. 1c, red box '2'), while the intensity of ApoA1 accounted for

96.23% of the TIC for the 25 kDa band (Fig. 1c, green box '3'). Similar LFQ analysis of LC-MS

data for the purified isNS1ts mutant showed that NS1 and ApoA1 were the major components

(Supplementary Fig. 2) of the respective Elute and ~250 kDa gel band on Native-PAGE

(Supplementary Fig. 2a; 3a, purple box). The corresponding 50 kDa and 25 kDa gel bands on the

reducing SDS-PAGE (Supplementary Fig. 2b, 3b) identified as NS1 and ApoA1 by
immunoblotting with the respective antibodies (Supplementary Fig. 3b), were further confirmed
by LC-MS as NS1 and ApoA1 (Supplementary Fig. 2c, 3c, Supplementary file 1). Taken
together, the results from the Native PAGE and LFQ by LC-MS strongly indicated that the
isNS1 purified from the supernatant of infected cells *in vitro* is an approximately 250 kDa
complex of NS1 and ApoA1.

Fig. 2. CryoEM analysis of secreted NS1 in complex with antibody Ab56.2. (a) Size
exclusion chromatography was run on a Superdex 200 increase 3.2/300 GL column connected to
the ÄKTA purifier with a flow rate of 0.075 mL/min in PBS (pH 7.4) for purified isNS1wt
(gray), Ab56.2 (faded red), Fab56.2 (faded blue), isNS1wt:Ab56.2 (red) and isNS1wt:Fab56.2
(blue). A slight leftward shift in elution volume was observed for isNS1wt upon complexing
with Ab56.2 and Fab56.2. 2D class averages of **(b)** isNS1wt:Ab56.2 and **(c)** isNS1wt:Fab56.2
showing representative sub-class of the Fab56.2:isNS1:HDL particles with black scale bar, 50 Å.
The corresponding number of particles and percentages are listed below the respective boxes.
Red dashed line boxes highlight two rare views consisting of 1033 particles (3.5%) only seen in
isNS1wt:Fab56.2 sample. **(d)** Model of isNS1wt:Fab56.2 predicted structures rigid body fitted in
the CryoEM map of Fab56.2:isNS1:HDL (grey, contoured at 0.14) with correlation value of 0.75
to the fitted regions (map simulated from atoms at 5 Å). isNS1wt is coloured by its three
domains, namely the β-roll (orange), wing (blue), and β-ladder (cyan). Fab56.2 is coloured by its
heavy chain (dark green) and light chain (light green).
205

CryoEM structure of isNS1

To gain insights into the molecular organization of this native isNS1, we determined the cryoEM structure of isNS1 (Supplementary Table 1 and Supplementary Fig. 4-7). The 2D classes of isNS1ts are observed to be heterogenous spheres with some internal features, some of which resembling the cross-shaped NS1 protruding from the sphere (Supplementary Fig. 4b). However, the 3D reconstruction of isNS1ts alone was observed to have no discernible features beyond a spherical-like density with varying dimensions averaging around 106 Å by 77 Å (Supplementary Fig. 4b-c). To obtain higher resolution structural information on the complex, we collected data for the ternary complexes of the isNS1wt:Ab56.2 isNS1wt:Fab56.2 (Fig. 2a and Supplementary Fig. 5-6). The use of antibodies as fiduciary markers is a well-proven approach for solving the structures of small proteins[25,26]. Overall, both samples resulted in similarly distinguishable 2D class averages (Fig. 2b-c) which show asymmetrical binding of one unit of Fab56.2 to the dimeric cross-shaped sNS1 associated with a lower-density sphere, presumably HDL.

In the isNS1wt:Fab56.2 ternary complex dataset, we further observed two apparent but rare sub-classes of free sNS1wt dimer bound to two units of Fab56.2 representing 3.5% of the total population of picked particles (Fig. 2c, red dashed boxes). While the cryoEM map reconstructions remain limited at low resolutions (Supplementary Fig. 5-6), they could be fitted with the NS1 dimer and Fab56.2 models (Fig. 2d) predicted separately using AlphaFold2[27,28], with a correlation value of 0.75 to the fitted regions using a simulated 5 Å map from the NS1 dimer and Fab models. The AlphaFold2 predicted model of the NS1 dimer was used despite the availability of the crystallographic dimer model (PBS ID: 6WER)[29] given their high similarity (RMSD: 0.67) and that the latter had a few unmodelled residues (aa 118-128). The map-model fitting showed that the epitope sites were at the tip of the C-terminal β-ladder region. The approximate dimensions of the spherical HDL could also be measured at 82 Å by 65 Å (Supplementary Fig. 5-6) which aligned with existing reports of spherical HDL being measured at 70-120 Å in diameter with predominantly homodimeric ApoA1 arranged as an anti-parallel double-belt stabilized by intermolecular salt bridges[30,31]. We predicted a model of an antiparallel dimer of ApoA1 using AlphaFold2[27,28] without the first 58 residues of ApoA1 as the N-terminal domain is highly flexible[30]. The model has a cross-sectional distance of approximately 78 Å and fits into the spherical density in the cryoEM map. The NS1 dimer appears semi-embedded on the HDL particle through its hydrophobic surface, with a conformation that exposed only one end of

the β-ladder domain to which Ab56.2 and Fab56.2 bind. Overall, the cryoEM model indicated

that the purified ~250 kDa isNS1 is a complex of a NS1 dimer and HDL, consistent with the gel

and LFQ analysis of LC-MS data presented above.

Similarly, we determined the structures of the ternary complex of isNS1ts mutant with Fab56.2

(Fig. 3a). Interestingly, 2D class averages of isNS1ts:Fab56.2 dataset showed a significant sub-

population of a free NS1ts dimer bound to two units of Fab56.2 (53.7%), in addition to the HDL

spheres (23.6%) and isNS1ts dimer:HDL:Fab56.2 (22.7%) classes (Fig. 3b and Supplementary

Fig. 7). The predicted structure of the isNS1ts dimer bound to Fab56.2 could be fitted with an

overall correlation value of 0.5 using map simulated from atoms at 5 Å (Fig. 3c). Achieving

atomic resolution to map the Fab binding interface with the NS1 β-ladder domain remains a

challenge due to the preferred orientation (Supplementary Fig. 7). A comparison of the cryoEM

density maps of free isNS1ts dimer:Fab56.2 (Fig. 3D, grey) and isNS1ts dimer:HDL:Fab56.2

ternary complex (Fig. 3d, yellow) fitted with a correlation value of 0.71 based on the NS1 map

region, revealed a pose difference of the Fab56.2 between the free isNS1ts dimer and the HDL-

bound isNS1ts dimer (Fig. 3d, inset). Further comparison of the density maps of the isNS1

dimer:HDL population in the Fab56.2 ternary complexes of isNS1wt (Fig. 3e, purple) and

isNS1ts (Fig. 3e, yellow) revealed conformational similarities of the isNS1 dimer:HDL complex.

257
258
Fig. 3. Secreted NS1 forms free dimers in complex with antibody Ab56.2. (a) Size exclusion
chromatography was run on a Superdex 200 increase 3.2/300 GL column connected to the
AKTA purifier with a flow rate of 0.075mL/min in PBS (pH 7.4) for purified isNS1ts (gray),
Fab56.2 (faded blue) and isNS1ts:Fab56.2 (red). A similar leftward shift in elution volume was
also observed for isNS1ts upon complexing with Fab56.2. **(b)** 2D class averages of
isNS1ts:Fab56.2 dataset with 326,548 particles picked and separated into 3 distinct populations,
HDL spheres, Fab56.2:isNS1:HDL, and free isNS1:Fab56.2. Black scale bar, 50 Å, as indicated.
The corresponding number of particles and percentages are listed below the respective boxes. **(c)**
Model of isNS1ts dimer and Fab56.2 predicted structures rigid body fitted in the
isNS1ts:Fab56.2 density map (grey, contoured at 0.14) with correlation value of 0.53 (overall,
map simulated from atoms at 5 Å). isNS1ts is coloured by its three domains, namely the β-roll
(orange), wing (blue), and β-ladder (cyan). Fab56.2 is coloured by its heavy chain (dark green)

and light chain (light green). **(d)** Density map fitting between isNS1ts:Fab56.2 (grey, contoured at 0.14) to Fab56.2:NS1ts:HDL (yellow, contoured at 0.1) with correlation value of 0.53 (overall) and 0.7137 (on D2NS1 map region only). Inset shows the difference in the pose of Fab from the free NS1 form to the HDL-bound form. (e) Density map fitting between Fab56.2:isNS1wt:HDL (purple, contoured at 0.05) to Fab56.2:NS1ts:HDL (yellow) with correlation value of 0.72 (overall).

XL-MS maps the inter- and intra-molecular architecture of the isNS1

Next to refine our understanding of the NS1 dimer:HDL complex of the isNS1 indicated by the LFQ of LC-MS (Fig. 1d and Supplementary Fig. 3d) and cryoEM results (Fig. 2d and 3d), we probed the interactions between NS1 and ApoA1 - the major protein moiety of HDL - using crosslinking mass spectrometry (XL-MS)[32] on the immunoaffinity purification eluate fraction (Fig. 4a, Supplementary Fig. 8a-b). We identified 28 NS1:NS1 and 29 ApoA1:ApoA1 intramolecular crosslinks, as well as 25 NS1:ApoA1 intermolecular crosslinks for isNS1wt (Fig. 4b; Supplementary File 2a-b). As depicted in Fig. 4b-c, the NS1 residues that are involved in both intra- and inter-molecular crosslinks were located mainly on the β-roll and wing domains, with ~34% of the NS1:ApoA1 crosslinks being located on the β-roll domain. The NS1:NS1 intramolecular crosslink pairs were further validated on the predicted NS1 dimer model by implementing a 30 Å cut-off distance for the Cα atoms of crosslinked residues, a criterion that was met by 85% of the NS1:NS1 intramolecular crosslinks (24 out of 28; Fig. 4d). The NS1:ApoA1 intermolecular crosslinks for the immunoaffinity purified isNS1wt were found to be located between the β-roll and wing domains of NS1 and the helices 3, 4, 8 and 10 of ApoA1 (Fig. 4b, d). Applying the same cut-off value of <30 Å distance between the Cα atoms of the NS1:ApoA1intermolecular crosslinked residues and validating it on the cryoEM model of isNS1wt in this study (Fig. 4d-e) satisfied 76% of the crosslinks (19 out of 25 intermolecular crosslinks between NS1 dimer and ApoA1). In the case of ApoA1:ApoA1 intramolecular crosslinks, the <30 Å distance between their Cα atoms cut-off satisfied 62% of the crosslinks (18 out of 29 ApoA1:ApoA1 intramolecular crosslinks). The higher number of violated crosslinks in ApoA1:ApoA1 compared to those in NS1:NS1 could be due to the flexible and dynamic nature of ApoA1 dimers which may prevent it from assuming a fixed relative position among the ApoA1 dimer population in the HDL. Overall, the XL-MS data and structural validation agreed with the cryo-EM model of isNS1 being a complex of the NS1 dimer embedded on a HDL particle composed of an ApoA1 dimer (Fig. 4d-e).

Additionally, the XL-MS mapping of isNS1ts mutant (Supplementary Fig. 8c-d) followed by rigid-body fitting into the corresponding cryoEM map (Supplementary Fig. 8e) revealed a similar crosslink pattern to isNS1wt, except that there were fewer crosslink sites (Supplementary Fig. 8f, Supplementary File. 2a). This indicated that isNS1ts dimer may have a weaker affinity to HDL in comparison to the isNS1wt dimer. This finding is consistent with the cryoEM data for isNS1ts where we found that free isNS1ts dimers bound to Fab56.2 formed a significant sub-population (34.5%) of the total picked particles (Fig. 3b, Supplementary Fig. 7b).

Given the reported structures of hexameric and tetrameric rsNS1 from mammalian expression systems, insect cells[11,15,16,29] and Expi293 HEK cells[17], we further examined whether rsNS1 oligomerizes in the presence of exogenously added human HDL, using XL-MS. The crosslinked rsNS1 alone appeared as higher oligomer bands of dimers, tetramers, and hexamers on the SDS-PAGE analysis (Fig. 4f, lane 3) which differed from the crosslinked isNS1 (Fig. 4f, lane 3). Interestingly, the addition of human HDL and crosslinker resulted in a smearing band ranging from 30-150 kDa with the loss of the well-defined monomeric rsNS1 band (Fig. 4f, lane 4). In the crosslinked rsNS1 in the absence of HDL, we identified 17 NS1:NS1 intramolecular crosslinks at the β-roll and wing domains (Fig. 4f-g, Supplementary File 3a-b). In the presence of human HDL, we identified 30 ApoA1:ApoA1, 4 NS1:NS1, and 6 NS1:ApoA1 crosslinks (Supplementary File 3a-b). Most crosslinks identified are ApoA1:ApoA1 intramolecular crosslinks in the rsNS1:ApoA1 samples (Fig. 4g). Only 1 out of 6 rsNS1:ApoA1 intermolecular crosslinks was within the 30 Å cutoff distance when mapped to the NS1:ApoA1 cryoEM model (Fig. 4e). This was in stark contrast with the isNS1 samples where the number of crosslinks between ApoA1:ApoA1, sNS1:sNS1, and sNS1:ApoA1 was similar (Fig. 4b). Overall, this suggests that rsNS1 may form oligomers of dimers but not a complex with HDL under the tested conditions.

330
Fig. 4. Interaction sites of NS1:ApoA1 complex identification by crosslinking mass
spectrometry. (a) SDS-PAGE analysis of isNS1wt with or without the addition of DSSO
crosslinker. **(b)** The identified crosslinks are visualised on the NS1 and bovine ApoA1
constructs. The intramolecular (NS1:NS1 and ApoA1:ApoA1) crosslinks are in magenta. The
intermolecular NS1:ApoA1 crosslinks are in green. **(c)** The isNS1 residues that are involved in
NS1:NS1 interactions are visualised on the NS1 dimer model. The NS1 cartoon model is
coloured by its three domains, namely the β-roll (orange), wing (blue), and β-ladder (cyan) with
the intramolecular (magenta) and intermolecular (green) crosslinking sites depicted as spheres.

(d) The overall model interpretation of NS1:ApoA1 complex within the crosslinker theoretical
distance cut-off at =< 30 Å as depicted. ApoA1 dimer cartoon model with its conserved helices
as labelled coloured in intervals of grey and light purple. **(e)** The NS1:ApoA1 dimer model with
validated crosslinks were fitted into the cryoEM envelope. **(f)** SDS-PAGE analysis of
crosslinked rsNS1 alone or with human HDL **(Lanes 3-5)**. Non-crosslinked rsNS1 and Human
HDL are the control **(Lanes 1-2)**. The crosslinked rsNS1 can be seen in higher oligomers **(Lane
3)**. **(g)** Identified crosslinks are mapped on the NS1 and ApoA1 constructs, coloured as per panel
B.
347

Ab56.2 binds at the C-terminal β-ladder region of NS1

The cryoEM map for isNS1wt:Fab56.2 reliably placed the epitope site of Ab/Fab56.2 at the C-

terminal b-ladder region, although the precise identification of the epitope site was hindered by

the limited resolution of the cryoEM maps (Fig. 2-3, Supplementary Fig. 4-7). Attempts to

improve the resolution were challenging so we resorted to using a truncated form of recombinant

secreted NS1 consisting of only the C-terminal region (rsNS1c; residues 172-352) to determine

Ab56.2 epitope. The rsNS1c fragment of residues 172-352 was previously used for structural

studies of dengue[33,34] and other flaviviruses, West Nile virus[33], Zika virus[34,35], and Japanese

encephalitis virus[36]. The epitope region of Ab56.2 was first confirmed by overlapping 15-mer

NS1 peptide competition ELISA for Ab56.2 binding (Supplementary Fig. 9a, Supplementary

Table 2). The greatest competition was observed at the NS1 peptide spanning residues 316-330

out of 301-340 which showed a decrease in OD450 absorbance (Supplementary Fig. 9b). We

further applied the hydrogen-deuterium exchange mass spectrometry (HDX-MS) to monitor the

conformational dynamics between Fab56.2 and NS1c to obtain higher peptide-level resolution

for the binding interface. Comparative HDX results revealed global scale protection against

deuterium exchange of NS1c in the presence of Fab56.2 (Fig. 5a, Supplementary Fig. 10a),

indicative of reduced conformational dynamics. Large-scale differences in deuterium exchange

were primarily observed across peptides spanning the dimer interface (residues 174-186) and at

the tip of the C-terminal β-ladder (residues 286-347) of NS1c (Fig. 5b). Multiple overlapping

peptides, spanning residues 286-347 (Fig. 5b), showed protection against deuterium exchange

even at the longer labeling times (100 min) in NS1c:Fab56.2 complex (Supplementary Fig. 10a,

Supplementary File 4). These results suggest that Fab56.2 bound stably with NS1c. Coupled with

the observation that the most significant change in the differential plot was at residues 300-310

(Fig. 5a), these results collectively indicate that NS1c binds stably to Fab56.2 at the primary

epitope site of residues 300-310.

Comparative HDX results of NS1c-bound and free Fab56.2 revealed protection against
deuterium exchange of all three complementarity-determining regions (CDR) of the heavy chain
(Fig. 5c, Supplementary Fig. 10b) and CDRL2 of the light chain (Fig. 5d, Supplementary Fig.
10c). Upon binding to NS1c, peptides spanning CDRH3 (residues 89-116) showed the largest
magnitude decrease in deuterium exchange at all labeling time points, while CDRH2 showed
minor significant differences. In addition, only the CDRL2 loop (residues 49-56) of the light
chain of Fab56.2 showed significantly decreased deuterium exchange. Mapping of the deuterium
exchange profiles of heavy and light chains onto a model of Fab56.2 (Fig. 5e) revealed that
CDRH1-3 and CDRL2 loops were spatially co-localized to form the paratope site to bind NS1c,
with the heavy chain being the primary anchor. Collectively, these results indicated that residues
300-310 of NS1 act as the primary epitope site to bind Fab56.2 and Fab56.2 binds stably with
NS1c.

387
Fig. 5. Interaction interface of rsNS1c-Fab562 complex characterized by HDX-MS. (a)
Woods differential plot showing the differences in deuterium exchange (y-axis) between
Fab562-bound and free NS1c protein across various residues (x-axis) at 10 min labeling
timepoint. Negative differences indicate protection against deuterium exchange across NS1c
peptides in the presence of Fab562, as compared to free NS1c. A p-value <0.05 is considered as
significance threshold, which identified 6 non-significant peptides (grey lines) and 48 protected
peptides (blue lines). **(b)** Cartoon representation of NS1c dimer model showing differences in
deuterium exchange at 10 min labeling as indicated in key. Dashed line distinguishes the two

monomers in the dimer. Peptide spanning residues 300-310 showing the highest protection are indicated. **(c, d)** Woods differential plot comparing the differences in deuterium exchange of Fab562 in the presence and absence of NS1c for various peptides of **(c)** heavy chain, and **(d)** light chain. A p-value <0.05 is considered as significance threshold, which identified deprotected (red lines), non-significant, and protected peptides as indicated. **(e)** Cartoon representation of Fab562 model showing deuterium exchange differences at 10 min mapped for Fab562-NS1c complex, as per key. Plots were generated using Deuteros 2.0, while cartoon structures were generated using PyMoL.

Circulating NS1 is in the form of NS1:HDL complex

Lastly, we sought to provide *in vivo* evidence of NS1 and ApoA1 interaction using sera samples from DENV-infected mice and a human patient. Since we previously showed that DENV2 T164S mutant virus led to more severe disease in the AG129 mouse model[23], we used the same infection model to obtain pooled mice serum for immunoaffinity purification of sNS1ts as described in Supplementary Fig. 1. The mouse sera immunoaffinity purified sNS1ts was detected at ~250 kDa by both anti-ApoA1 and anti-NS1 antibodies in a Western blot analysis following separation on a Native PAGE (Fig. 6a), supporting our mass spectrometry finding that ApoA1 is part of the ~250 kDa isNS1 protein complex purified from infected Vero cells (Fig. 1d). We next examined if the isNS1:HDL complex can be detected in dengue patients using serum samples obtained from the CELADEN clinical trial[37]. A primary infection sample with high levels of sNS1 (>15 µg/ml) detected by commercial Platelia™ NS1 ELISA was selected for this purpose to avoid the confounding factor of the presence of anti-NS1 Abs in the secondary infected patients and to pull-down sufficient sNS1 from the patient serum. The patient serum was first subjected to a pre-clearing step using Protein AG resin to reduce the abundance of human IgGs that may interfere with the downstream analysis. We then subjected the pre-cleared patient serum to immunoprecipitation using commercial polyclonal anti-ApoA1 antibodies to pull down and detect ApoA1 and possibly sNS1. Increasing the anti-ApoA1 antibody from 10 to 50 µg coupled onto Protein AG resin for pull-down resulted in a corresponding increase in the sNS1 amount detected by ELISA (Fig. 6b), suggesting a specific association between sNS1 and ApoA1. Collectively, despite the limited amount of sNS1 in sera samples from DENV-infected mice or humans, our data points to the association of ApoA1 with sNS1 *in vivo*.

427
428
Fig. 6. sNS1 associates with ApoA1 in DENV infected mouse and human serum. (a) AG129
mice (n = 10) were infected with DENV2 NS1 T164S mutant virus and the pooled infected sera
collected on day 4 post-infection was subjected to NS1 immunoaffinity purification using anti-
NS1 56.2 coupled resin as in Supplementary Fig. 1. 2 mg of the purified eluate was then
subjected to Western blot analysis after separation and transfer from Native PAGE for detection
of ApoA1 and NS1. ApoA1 was detected using the mouse monoclonal anti-ApoA1 clone 513
(Invitrogen, MIA1404) **(left panel)** and the oligomeric NS1 was detected using anti-NS1 56.2
IgG clone **(right panel)**. **(b)** Protein AG resin (Pierce) pre-cleared DENV1 infected patient
serum (n = 1) from the CELADEN trial[37] was immunoprecipitated with 10 or 50 mg of rabbit
polyclonal anti-ApoA1 antibody (Biorbyt, orb10643) to detect association between ApoA1 and
NS1 by ELISA. The amount of ApoA1 and NS1 in the immunoprecipitated sample was
determined by human ApoA1 (Abcam, ab189576) and Platelia[TM] NS1 Ag (Bio-Rad) ELISAs.
441

Discussion

In this study, by integrating biochemical and biophysical approaches including immunoaffinity

purification, cryoEM, XL-MS and HDX-MS, we have successfully purified and characterised a

native population of isNS1. We determined the cryo-EM structure of a natively secreted NS1

from DENV2-infected cells as a ternary complex of NS1 dimer embedded on a single HDL

particle. Our in-solution cross-linking mass spectrometry (XL-MS) analysis captured interactions

within a cut-off distance of 30 Å imposed by the linker-spacer between the β-roll and wing

domains of NS1 and helices 3, 5, 8, and 10 of (bovine) ApoA1 – the main protein moiety of

HDL. More importantly, we were able to detect NS1:HDL complexes *in vivo* in both the sera of

DENV2-infected mice and in the serum of a DENV1-infected patient, highlighting the biological

relevance of the NS1:HDL complexes in dengue infection.

Benfrid et al. (2022) recently reconstituted complexes of rsNS1 dimers bound to the surface of a

spheroid human HDL particle *in vitro* at 1:1 to 3:1 ratio when up to 400 μg/mL of rsNS1 were

used[18]. Similarly, in our studies with infection-derived sNS1 (isNS1), we observed up to two

NS1 dimers docked onto an HDL particle (Fig. 2-3, Supplementary Fig. 1c, 4-7). It is noteworthy

that the concentration of isNS1:HDL in our study (~30 μg/mL) is closer to the maximum

clinically reported concentration of sNS1 in the sera of DENV patients (50 μg/mL)[3,38,39].

Therefore, the ratio of sNS1 to HDL varies depending on the concentration of the proteins used

for complex formation. Furthermore, the observation that NS1 interacts with HDL regardless of

the source of mammalian ApoA1 is unsurprising, given the high sequence similarity of bovine

and mouse HDL to human HDL of approximately 79% and 65% respectively (Supplementary

Fig. 11) and that the structure of ApoA1 is highly conserved [40]

Acknowledging the limitation of our study that we used only one antibody (56.2) for our

immunoaffinity purification, our key take-home message in terms of structure is that our data

does not support the widely-held view that sNS1 is composed of a trimer of NS1 dimers with a

~28 Å wide central channel filled with lipid as established in several seminal publications[14-16]. In

fact, our data confirms the observations reported in Benfrid et al.[18] and in the rare instances

where trimer of dimers have been experimentally shown by cryoEM of rNS1 structures[17], only a

single lipid molecule can be accommodated in the narrow central channel and unlikely to be

reflective of sNS1 circulating in serum. We argue that the amount of lipid associated with

infection derived sNS1[15,23] is consistent with NS1 dimer(s) associated with HDL. The interaction

with HDL requires an exposed hydrophobic face of NS1 which is incompatible to the hexamer

model of sNS1. Indeed, it is agreeable that the oligomeric state of NS1 is dynamic which may

vary depending on the context and environment. This point requires further independent studies.

Our previous functional studies comparing isNS1wt with isNS1ts (T164S) showed that

incubating immunoaffinity-purified sNS1 with human PBMCs from 3 independent human

donors triggered the production of proinflammatory cytokines IL6 and TNFa in a concentration

dependent manner and that it was more pronounced with the mutant than the WT[23]. Furthermore,

we have provisional transendothelial electrical resistance (TEER) assay data with isNS1 proteins

used in the structural studies with human umbilical vascular endothelial cells (hUVEC) which

show that acid-eluted isNS1 proteins in this study induces endothelial hyperpermeability as

shown in previous studies[41]. The presence of a significant population of free isNS1ts dimers in

complex with Fab56.2 and fewer crosslinks identified between isNS1ts and ApoA1 point

towards the possibility that isNS1ts is more dynamic in its association with host proteins than

isNS1wt. The disease severity and increased complement protein expression in AG129 mice

liver infected with NS1T164S carrying virus compared with WT that was observed in our

previous study [23] can be ascribed to weakly bound mutant NS1 with fast on/off rate with HDL

being transported to the liver where specific receptors bind to free sNS1 and interact with

effector proteins such as complement to drive inflammation and associated pathology. This

notion also requires further validation with other mutations that can impact sNS1 – HDL

interaction.

Overall, our work provided a refined understanding of the native structure of sNS1 in DENV

pathogenesis. The functional role of HDL-bound NS1 protein in disease is now a topic of

contention but one that will be resolved by more data obtained with infection-derived sNS1 by

scientists in the field. Contrary to the known protective roles of HDL such as against endotoxic

shock caused by lipopolysaccharides[42-44], Benfrid et al. (2022) showed that sNS1:HDL complex

triggers proinflammatory signals when tested with primary human macrophages[18]. The presence

of sNS1:HDL complexes as a predominant population of sNS1 in our *in vitro* infection model

corroborates with a previous report of an interaction between insect-derived recombinant NS1

and the classical HDL/ApoA1-binding receptor known as SR-BI[19], which is primarily expressed

on hepatocytes[45]. This interaction in hepatocytes have been suggested to trigger endocytosis of

sNS1 and subsequent downstream effects including vascular permeability, which is a

characteristic feature of severe dengue[19,46,47]. Internalisation of sNS1 is critical for increased

virus production in hepatocytes[46] and NS1-mediated permeability in endothelial cells[47].

Furthermore, NS1:HDL complexes have been shown to induce the production of pro-

inflammatory cytokines in macrophages and their presence has been detected in the sera of

hospitalised DENV patients[18]. Yet, this hypothesis is confounded by other reports. While rsNS1

has been shown to trigger a TLR4-mediated release of proinflammatory cytokines to induce

vascular leakage[48,49], Coelho et al. (2021) further showed that lipid-free recombinant ApoA1

neutralizes the proinflammatory effects caused by insect-derived recombinant intracellular

NS1[49]. Furthermore, exogenous administration of purified HEK-293 derived NS1 failed to

worsen *in vivo* vascular leakage in sublethally infected mice[50]. This underscores the need for

physiologically relevant studies to reconcile these discrepancies and gain a comprehensive

understanding of the role of NS1-HDL complex in DENV pathogenesis.

In conclusion, despite the study limitations that our isNS1wt and isNS1ts were immunoaffinity

purified with one specific monoclonal antibody (Ab56.2), our structural model, obtained by

CryoEM and verified by XL-MS, of infection-derived sNS1 shows a NS1 dimer embedded on a

single HDL particle composed of ApoA1 dimer. Future investigation of this work using infected

mouse sera from other DENV strains and different flaviviruses would inform the structural and

functional relevance of the HDL-bound NS1 protein complex amongst flaviviral NS1 as well as

provide a basis for the development of NS1-directed therapeutics.

References

1   Bhatt, S. *et al.* The global distribution and burden of dengue. *Nature* **496**, 504-507
(2013). https://doi.org:10.1038/nature12060
2   Diamond, M. S. & Pierson, T. C. Molecular Insight into Dengue Virus Pathogenesis and
Its Implications for Disease Control. *Cell* **162**, 488-492 (2015).
https://doi.org:10.1016/j.cell.2015.07.005
3   Libraty, D. H. *et al.* High Circulating Levels of the Dengue Virus Nonstructural Protein
NS1 Early in Dengue Illness Correlate with the Development of Dengue Hemorrhagic
Fever. *J Infect Dis* **186**, 1165-1168 (2002). https://doi.org:10.1086/343813
4   Paranavitane, S. A. *et al.* Dengue NS1 antigen as a marker of severe clinical disease.
BMC Infect Dis **14**, 570 (2014). https://doi.org:10.1186/s12879-014-0570-8
5   Glasner, D. R., Puerta-Guardo, H., Beatty, P. R. & Harris, E. The Good, the Bad, and the
Shocking: The Multiple Roles of Dengue Virus Nonstructural Protein 1 in Protection and
Pathogenesis. *Annu Rev Virol* **5**, 227-253 (2018). https://doi.org:10.1146/annurev-
virology-101416-041848
6   Watterson, D., Modhiran, N. & Young, P. R. The many faces of the flavivirus NS1
protein offer a multitude of options for inhibitor design. *Antiviral Res* **130**, 7-18 (2016).
https://doi.org:10.1016/j.antiviral.2016.02.014
7   Scaturro, P., Cortese, M., Chatel-Chaix, L., Fischl, W. & Bartenschlager, R. Dengue
Virus Non-structural Protein 1 Modulates Infectious Particle Production via Interaction
with the Structural Proteins. *PLOS Pathogens* **11**, e1005277 (2015).
https://doi.org:10.1371/journal.ppat.1005277
8   Plaszczyca, A. *et al.* A novel interaction between dengue virus nonstructural protein 1
and the NS4A-2K-4B precursor is required for viral RNA replication but not for
formation of the membranous replication organelle. *PLoS Pathog* **15**, e1007736 (2019).
https://doi.org:10.1371/journal.ppat.1007736
9   Puerta-Guardo, H., Glasner, D. R. & Harris, E. Dengue Virus NS1 Disrupts the
Endothelial Glycocalyx, Leading to Hyperpermeability. *PLoS Pathog* **12**, e1005738
(2016). https://doi.org:10.1371/journal.ppat.1005738
10   Beatty, P. R. *et al.* Dengue virus NS1 triggers endothelial permeability and vascular leak
that is prevented by NS1 vaccination. *Sci Transl Med* **7**, 304ra141 (2015).
https://doi.org:10.1126/scitranslmed.aaa3787
11   Akey, D. L. *et al.* Flavivirus NS1 Structures Reveal Surfaces for Associations with
Membranes and the Immune System. *Science* **343**, 881-885 (2014).
https://doi.org:doi:10.1126/science.1247749
12   Xu, X. *et al.* Contribution of intertwined loop to membrane association revealed by Zika
virus full-length NS1 structure. *EMBO J* **35**, 2170-2178 (2016).
https://doi.org:10.15252/embj.201695290
13   Brown, W. C. *et al.* Extended surface for membrane association in Zika virus NS1
structure. *Nature Structural & Molecular Biology* **23**, 865-867 (2016).
https://doi.org:10.1038/nsmb.3268
14   Flamand, M. *et al.* Dengue Virus Type 1 Nonstructural Glycoprotein NS1 Is Secreted
from Mammalian Cells as a Soluble Hexamer in a Glycosylation-Dependent Fashion.
Journal of Virology **73**, 6104-6110 (1999). https://doi.org:doi:10.1128/JVI.73.7.6104-
6110.1999

15   Gutsche, I. *et al.* Secreted dengue virus nonstructural protein NS1 is an atypical barrel-
shaped high-density lipoprotein. *Proc Natl Acad Sci U S A* **108**, 8003-8008 (2011).
https://doi.org:10.1073/pnas.1017338108

16   Muller, D. A. *et al.* Structure of the dengue virus glycoprotein non-structural protein 1 by
electron microscopy and single-particle analysis. *J Gen Virol* **93**, 771-779 (2012).
https://doi.org:10.1099/vir.0.039321-0

17   Shu, B. *et al.* CryoEM structures of the multimeric secreted NS1, a major factor for
dengue   hemorrhagic   fever.   *Nature   Communications*   **13**,   6756   (2022).
https://doi.org:10.1038/s41467-022-34415-1

18   Benfrid, S. *et al.* Dengue virus NS1 protein conveys pro-inflammatory signals by docking
onto   high-density   lipoproteins.   *EMBO   Rep*   **23**,   e53600   (2022).
https://doi.org:10.15252/embr.202153600

19   Alcala, A. C. *et al.* Dengue Virus NS1 Uses Scavenger Receptor B1 as a Cell Receptor in
Cultured Cells. *J Virol* **96**, e0166421 (2022). https://doi.org:10.1128/JVI.01664-21

20   Avirutnan, P. *et al.* Secreted NS1 of dengue virus attaches to the surface of cells via
interactions with heparan sulfate and chondroitin sulfate E. *PLoS Pathog* **3**, e183 (2007).
https://doi.org:10.1371/journal.ppat.0030183

21   Alcala, A. C. *et al.* The dengue virus non-structural protein 1 (NS1) is secreted from
infected mosquito cells via a non-classical caveolin-1-dependent pathway. *J Gen Virol*
98, 2088-2099 (2017). https://doi.org:10.1099/jgv.0.000881

22   Rodriguez-Roche, R. *et al.* Virus evolution during a severe dengue epidemic in Cuba,
1997. *Virology* **334**, 154-159 (2005). https://doi.org:10.1016/j.virol.2005.01.037

23   Chan, K. W. K. *et al.* A T164S mutation in the dengue virus NS1 protein is associated
with   greater   disease   severity   in   mice.   *Sci   Transl   Med*   **11**   (2019).
https://doi.org:10.1126/scitranslmed.aat7726

24   Kitti, C. W. K. *THE ROLE OF GENETIC VARIANTS OF NON-STRUCTURAL*
PROTEIN 1 IN DENGUE PATHOGENESIS, National University of Singapore, (2019).

25   Wu, S. *et al.* Fabs enable single particle cryoEM studies of small proteins. *Structure*
(London, England : 1993) **20**, 582-592 (2012). https://doi.org:10.1016/j.str.2012.02.017

26   Bloch, J. S. *et al.* Structure and mechanism of the ER-based glucosyltransferase ALG6.
Nature **579**, 443-447 (2020). https://doi.org:10.1038/s41586-020-2044-z

27   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021). https://doi.org:10.1038/s41586-021-03819-2

28   Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nature Methods*
19, 679-682 (2022). https://doi.org:10.1038/s41592-022-01488-1

29   Biering, S. B. *et al.* Structural basis for antibody inhibition of flavivirus NS1-triggered
endothelial   dysfunction.   *Science*   **371**,   194-200   (2021).
https://doi.org:10.1126/science.abc0476

30   Silva, R. A. *et al.* Structure of apolipoprotein A-I in spherical high density lipoproteins of
different   sizes.   *Proc   Natl   Acad   Sci   U   S   A*   **105**,   12176-12181   (2008).
https://doi.org:10.1073/pnas.0803626105

31   He, Y. *et al.* Apolipoprotein A1 Forms 5/5 and 5/4 Antiparallel Dimers in Human High-
density Lipoprotein. *Molecular & Cellular Proteomics* **18**, 854-864 (2019).
https://doi.org:10.1074/mcp.RA118.000878

32   Kao, A. *et al.* Development of a Novel Cross-linking Strategy for Fast and Accurate
Identification of Cross-linked Peptides of Protein Complexes. *Molecular & Cellular*
Proteomics **10** (2011). https://doi.org:10.1074/mcp.M110.002212

621	33	Edeling, M. A., Diamond, M. S. & Fremont, D. H. Structural basis of Flavivirus NS1
622		assembly and antibody recognition. *Proc Natl Acad Sci U S A* **111**, 4285-4290 (2014).
623		https://doi.org:10.1073/pnas.1322036111
624	34	Modhiran, N. *et al.* A broadly protective antibody that targets the flavivirus NS1 protein.
625		*Science* **371**, 190-194 (2021). https://doi.org:10.1126/science.abb9425
626	35	Wang, D. *et al.* A Mutation Identified in Neonatal Microcephaly Destabilizes Zika Virus
627		NS1 Assembly in Vitro. *Sci Rep* **7**, 42580 (2017). https://doi.org:10.1038/srep42580
628	36	Poonsiri, T. *et al.* Structural Study of the C-Terminal Domain of Nonstructural Protein 1
629		from Japanese Encephalitis Virus. *J Virol* **92** (2018). https://doi.org:10.1128/jvi.01868-17
630	37	Low, J. G. *et al.* Efficacy and safety of celgosivir in patients with dengue fever
631		(CELADEN): a phase 1b, randomised, double-blind, placebo-controlled, proof-of-
632		concept trial. *The Lancet Infectious Diseases* **14**, 706-715 (2014).
633		https://doi.org:10.1016/s1473-3099(14)70730-3
634	38	Young, P. R., Hilditch, P. A., Bletchly, C. & Halloran, W. An Antigen Capture Enzyme-
635		Linked Immunosorbent Assay Reveals High Levels of the Dengue Virus Protein NS1 in
636		the Sera of Infected Patients. *Journal of Clinical Microbiology* **38**, 1053-1057 (2000).
637		https://doi.org:doi:10.1128/jcm.38.3.1053-1057.2000
638	39	Alcon, S. *et al.* Enzyme-Linked Immunosorbent Assay Specific to Dengue Virus Type 1
639		Nonstructural Protein NS1 Reveals Circulation of the Antigen in the Blood during the
640		Acute Phase of Disease in Patients Experiencing Primary or Secondary Infections.
641		*Journal of Clinical Microbiology* **40**, 376-381 (2002).
642		https://doi.org:doi:10.1128/jcm.40.02.376-381.2002
643	40	Malajczuk, C. J., Gandhi, N. S. & Mancera, R. L. Structure and intermolecular
644		interactions in spheroidal high-density lipoprotein subpopulations. *J Struct Biol X* **5**,
645		100042 (2021). https://doi.org:10.1016/j.yjsbx.2020.100042
646	41	Puerta-Guardo, H. *et al.* Flavivirus NS1 Triggers Tissue-Specific Vascular Endothelial
647		Dysfunction Reflecting Disease Tropism. *Cell Rep* **26**, 1598-1613 e1598 (2019).
648		https://doi.org:10.1016/j.celrep.2019.01.036
649	42	Trinder, M., Walley, K. R., Boyd, J. H. & Brunham, L. R. Causal Inference for
650		Genetically Determined Levels of High-Density Lipoprotein Cholesterol and Risk of
651		Infectious Disease. *Arteriosclerosis, Thrombosis, and Vascular Biology* **40**, 267-278
652		(2020). https://doi.org:doi:10.1161/ATVBAHA.119.313381
653	43	Tall, A. R. HDL in Morbidity and Mortality: A 40+ Year Perspective. *Clinical Chemistry*
654		**67**, 19-23 (2020). https://doi.org:10.1093/clinchem/hvaa148
655	44	Harslof, M., Pedersen, K. M., Afzal, S., Davey Smith, G. & Nordestgaard, B. G. Lower
656		levels of small HDL particles associated with increased infectious disease morbidity and
657		mortality: a population-based cohort study of 30 195 individuals. *Cardiovasc Res* **119**,
658		957-968 (2023). https://doi.org:10.1093/cvr/cvac194
659	45	Acton, S. *et al.* Identification of scavenger receptor SR-BI as a high density lipoprotein
660		receptor. *Science* **271**, 518-520 (1996). https://doi.org:10.1126/science.271.5248.518
661	46	Alcon-LePoder, S. *et al.* The Secreted Form of Dengue Virus Nonstructural Protein NS1
662		Is Endocytosed by Hepatocytes and Accumulates in Late Endosomes: Implications for
663		Viral Infectivity. *Journal of Virology* **79**, 11403-11411 (2005).
664		https://doi.org:doi:10.1128/jvi.79.17.11403-11411.2005
665	47	Wang, C. *et al.* Endocytosis of flavivirus NS1 is required for NS1-mediated endothelial
666		hyperpermeability and is abolished by a single N-glycosylation site mutation. *PLoS*
667		*Pathog* **15**, e1007938 (2019). https://doi.org:10.1371/journal.ppat.1007938

48   Modhiran, N. *et al.* Dengue virus NS1 protein activates cells via Toll-like receptor 4 and
disrupts endothelial cell monolayer integrity. *Sci Transl Med* **7**, 304ra142 (2015).
https://doi.org:10.1126/scitranslmed.aaa3863
49   Coelho, D. R. *et al.* ApoA1 Neutralizes Proinflammatory Effects of Dengue Virus NS1
Protein and Modulates Viral Immune Evasion. *J Virol* **95**, e0197420 (2021).
https://doi.org:10.1128/JVI.01974-20
50   Lee, P. X. *et al.* Relative contribution of nonstructural protein 1 in dengue pathogenesis. *J*
Exp Med **217** (2020). https://doi.org:10.1084/jem.20191548
51   Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass
spectrometry-based proteomics evidences. *Nucleic Acids Research* **50**, D543-D552
(2021). https://doi.org:10.1093/nar/gkab1038
679
680
681

Acknowledgments

We thank the scientific facility support from NTU Institute of Structural Biology and Protein Product Platform. The authors acknowledge the Cryo-Electron Microscopy Facility at Center for Bioimaging Science, Department of Biological Science, National University of Singapore the scientific and technical assistance. Recombinant DENV2 NS1 protein is a generous gift from Dr Shu Bo and Professor Lok Shee Mei. We thank members of the DL and SV labs for their support.

Funding: This research is supported by the Singapore Ministry of Education under its Singapore Ministry of Education Academic Research Fund Tier 2 (T2EP30220-0020) to DL and National Medical Research Council of Singapore (MOH-OFIRG18may-0006; MOH-OFIRG20nov-0002) to SV, A*STAR core funding and Singapore National Research Foundation under its NRF-SIS "SingMass" scheme to R.M.S, Career Development Award 2021 (A*STAR BMRC) to W.W.P., Career Development Fund 2021 (A*STAR BMRC) to S.Z., and LKCMedicine Dean's Postdoctoral Fellowship to C.B.L.A.

Author contributions: Conceptualization: SV and DL; Investigation: BLAC, AQN, WWP, KWKC, ZS, SSL, MJGW, MMC, JGL, EEO, SV and DL; Writing: BLAC, AQN, SV, and DL with input from all authors.

Competing interests: Authors declare that they have no competing interests.

Data availability:
The data that support this study are available from the corresponding authors upon reasonable request. The cryoEM density maps have been deposited in the Electron Microscopy Data Bank (EMDB) under accession codes EMD-36483 (isNS1ts:Fab56.2) and EMD-36480 (Fab56.2:isNS1ts:HDL). The protein model files for isNS1ts:Fab56.2 and Fab56.2:isNS1ts:HDL model are available upon request. Crosslinking MS raw files and the search results can be downloaded from https://repository.jpostdb.org/preview/14869768463bf85b347ac2 with the access code: 3827. The HDX-MS data is deposited to the ProteomeXchange consortium via PRIDE partner repository[51] with the dataset identifier PXD042235.