Discovery of a non-canonical prototype long-chain monoacylglycerol lipase through a

structure-based endogenous reaction intermediate complex

Nikos Pinotsis[1,$,#], Anna Krüger[2,#], Nicolas Tomas[3], Spyros D Chatziefthymiou[1], Claudia Litz[1], Simon

Arnold Mortensen[1], Mamadou Daffé[3], Hedia Marrakchi[3], Garabed Antranikian[2] and Matthias

Wilmanns[1,4]*

[1] European Molecular Biology Laboratory, Hamburg Unit, Notkestrasse 85, 22607 Hamburg, Germany

[2] Hamburg University of Technology, Kasernenstrasse 12, 21073 Hamburg, Germany

[3] Institut de Pharmacologie et de Biologie Structurale, IPBS, Université de Toulouse, CNRS, UPS,

Toulouse, France

[4] University Hamburg Clinical Center Hamburg-Eppendorf, Martinistrasse 52, 20251 Hamburg, Germany

[$] Current address: Department of Chemistry, National and Kapodistrian University of Athens, Zografou,

Greece

[#] Equal contribution

Corresponding author: Matthias Wilmanns, Email: matthias.wilmanns@embl-hamburg.de

ABSTRACT

The identification and characterization of enzyme function is largely lacking behind the rapidly increasing

availability of large numbers of sequences and associated high-resolution structures. This is often

hampered by lack of knowledge on *in vivo* relevant substrates. Here, we present a case study of a high-

resolution structure of an unusual orphan lipase in complex with an endogenous C18 monoacylglycerol

ester reaction intermediate from the expression host, which is insoluble under aqueous conditions and

thus not accessible for studies in solution. The data allowed its functional characterization as a prototypic

long-chain monoacylglycerol lipase, which uses a minimal lid domain to position the substrate through a

hydrophobic tunnel directly to the enzyme's active site. Knowledge about the molecular details of the

substrate binding site allowed us to modulate the enzymatic activity by adjusting protein/substrate

interactions, demonstrating the potential of our findings for future biotechnology applications.

INTRODUCTION

The annotation of relevant enzyme function under physiological conditions remains a major challenge,

despite the availability of millions of protein sequences and experimentally determined or predicted

molecular structures associated with them [1,2]. Enzymes often have different and broader substrate profiles

within their natural cellular environment than experimentally accessible under *in vitro* conditions or by

using purified protein samples. Hence, albeit powerful tools have been developed to predict enzymatic

function, options to prove them *in vivo* remain limited [3-8]. High-resolution structural biology allows the

direct visualization of enzyme-ligand complexes at atomic detail and thus presents a highly attractive tool

for enzyme function discovery, but it requires the availability of pure compounds for structural analysis.

Although protein structures in the presence of large hydrophobic substrates or ligands have been

published, often with a requirement of applying target-tailored solubilization protocols [9-12], they still

remain exceptional and hence available structure repositories remain mostly populated with protein

structures of soluble, small ligands. Alternatively, enzyme structures could also be determined in the

presence of endogenous substrates or catalysis intermediates taken up from their cellular environment for

direct evidence of enzyme-substrate relationships, but successful instances have remained rare to date [13].

This requires conditions where substrate turnover is inhibited by genetic modification of enzyme targets

or experimental conditions that lead to lack of catalytic activity. In this study, we demonstrate the power

of an endogenous substrate uptake for structural and subsequent functional characterization of an orphan

enzyme with putative lipase/esterase activity, which otherwise would not have been accessible due to lack

of solubility.

Lipases are enzymes that reversibly catalyze the hydrolysis and synthesis of a large variety of glycerol

esters [14-16]. Knowledge about these enzymes has allowed the development of numerous lipase-catalyzed

processes in medical biotechnology, detergent industry, organic synthesis, biodiesel production,

agrochemical industry, flavor and aroma industry, and food production [17-19]. In contrast to many other

carboxyl ester hydrolases, lipases are active mainly at water-lipid interfaces, a process known as

interfacial activation [20,21]. In essence, long-chain glycerol esters are delivered in non-monomeric

emulsions rather than water-soluble substrates. Interfacial activation of lipases is thought to require an

additional α-helical domain referred to as "lid", alternatively also named "cap", which contains several

hydrophilic residues in the vicinity of the active site [22]. In contrast to the catalytic domain (CD), there is

only little sequence conservation of the lid. Available structural data on lipases show highly diverse lid

topology and structure, suggesting variable mechanisms to promote active site opening for access of acyl

substrates and rendering precise prediction of enzymatic substrate specificity challenging [12,23,24]. Lipases

from extremophilic microorganisms are of superior interest for applications in industrial processes since

demanding conditions such as high temperature or high salt concentration do not impair their activity [25-27].

Most of the previous molecular activity studies have been limited to isolated or recombinant lipases.

Given that quantitative lipase activity measurements are generally hampered by the limited solubility of

long-chain glycerol esters under those experimental conditions it is plausible to assume that the substrate

profile differs under cellular conditions.

Based on turnover measurements with a broad spectrum of substrates, an enzyme from the thermophilic

anaerobic bacterium *Thermoanaerobacter thermohydrosulfuricus (Tth)* was tentatively assigned as a

lipase [28]. This strain was originally isolated from Solar Lake located on the Sinai peninsula, which has an

extreme marine environment with a temperature range from 16 to 60°C and high level of salinity. These

environmental conditions have given rise to complex biochemical phenomena that are linked to cycles of

evaporation and infiltration of water external sources. Isolates from Solar Lake have shown remarkable

biochemical processes related to degradation of starch, amylose, and pullulan [29,30]. As this enzyme

revealed a remarkable level of robustness against a variety of solvents and temperatures, it represents an

attractive target for potential biotechnology applications. However, underlying molecular mechanisms

explaining the enzyme's activity properties and substrate promiscuity remained unknown.

In this work, we determined the structure of the enzyme revealed an unexpected long-chain C18

monoacylglycerol (MAG) bound to the active site, which was taken up from its expression host. Its distal

part is embedded by a tunnel formed by an unconventional lid domain with minimal α-helix-β-hairpin-α-

helix (HBH) topology. In cell extracts that overexpress the enzyme, we found high levels of C16/18 fatty

acid (FA) products, in agreement with our structural data, demonstrating that the *Tth* enzyme functions as

a long-chain MAG lipase. *Tth* MAG lipase prototypes a new lipase/esterase family with HBH lid domains

comprising sequences inserted into the HBH β-hairpin that are highly divergent in length and structure.

Our findings demonstrate a powerful example of using high-resolution structural biology where complex

formation with an endogenous reaction intermediate has become achievable to elucidate its function.

RESULTS AND DISCUSSION

Tth MAG lipase reveals an α/β hydrolase fold with a minimal HBH lid domain

To provide insight into the *Tth* MAG lipase active-site topography and shed light on its unique substrate

promiscuity, we characterized the enzyme structurally and functionally. Purified *Tth* MAG lipase elutes

as a dimer **(Fig. 1a).** Its fold melting temperature measured by nano differential scanning fluorimetry

(nanoDSF) is at 77°C, which is within expectation in light of the thermophilic origin of *Tth* **(Fig. 1b,**

Supplementary Table 1). Next, we determined its high-resolution crystal structure, both in the presence

and absence of the covalent active site serine protease inhibitor phenylmethylsulfonyl fluoride (PMSF)

(Fig. 1c & e & 2, **Supplementary Table 2**). The overall fold of dimeric *Tth* MAG lipase can also be

reliably modeled using the AlphaFold2 and MMseqs2 package **(Supplementary Fig. 1)** [31,32]. However, as

there are significant differences in the areas of the bound ligands and the dimeric interface, in the

remaining contribution we refer to our experimental structures only.

Both *Tth* MAG lipase structures – the active and phenylmethylsulfonyl (PMS)-inhibited – reveal a two-

fold symmetric dimer, in agreement with its dimeric assembly in solution **(Fig. 1a & e)**. The *Tth* MAG

lipase dimer is formed by an almost flat interface area which represents about 10 % of the total solvent

accessible area **(Supplementary Fig. 2).** The interface is established through an intricate network of

charged interactions involving residues from both domains, with the majority of the interactions

originating from residues of the CD. It forms a large cleft that connects the two active sites of the two

protomers **(Fig. 1e).** Converting the charge of one of the most central interface residues (E72R) renders a

monomeric version of the protein, confirming that the interface observed in the structure is relevant **(Fig.**

1a).

A minimal lid domain comprising the residues 141-183 is inserted between strand β6 and helix αE on top

of the CD active-site area **(Fig. 1c-e, Supplementary Fig. 2)**. Different from most other lipases, in which

the lid is generally formed by a bundle of α-helices [22,24], in *Tth* MAG lipase this domain is compact. It is

formed by two small α-helices, followed by a β-hairpin and an additional α-helix, referred to as α-

helix/β-hairpin/α-helix (HBH) **(Fig. 1c, Supplementary Fig. 2)**. In both *Tth* MAG lipase structures, the

orientation of the HBH lid to the CD is basically identical **(Fig. 2d-e)**, dismissing any large-scale

opening/closing mechanisms previously shown for other lid domain-containing lipases [22,24]. Despite the

presence of the lid domain, the MAG lipase active site remains largely accessible **(Fig. 1e)**.

 Long chain MAG reaction intermediate is held by a hydrophobic lid tunnel

In the active site of both *Tth* MAG lipase structures, with and without PMS, we observed literally

identical density of a glycerol molecule, presenting one of the two final products of esterase/lipase

catalysis **(Fig. 2a-b, 3, Supplementary Fig. 3a).** The glycerol molecule tightly interacts via its two C2

and C3 hydroxyl groups with the carboxylate group of a non-conserved active-site residue Glu43, specific

to *Tth* MAG lipase **(Fig. 2a-b, Supplementary Fig. 5a)**. As di- and triacylglycerol (DAG, TAG) esters

would require linkages through these two hydroxyl groups, only MAG ester binding via the C1 hydroxyl

group of the glycerol backbone is sterically possible. The dihedral conformation of this residue is in an

energetically less favored area of the Ramachandran plot in both *Tth* MAG lipase structures

(**Supplementary Fig. 6**), suggesting that its role in glycerol binding may supersede any other lower

energy conformation.

In both protomers of the active *Tth* MAG lipase structure, we found additional extended ligand electron

density pointing away from the active site crossing through a 15 Å long tunnel formed by the HBH lid

domain (**Supplementary Fig. 3a**). As we did not add any substrate to the crystallization buffer, bound

ligands can only originate from the *Escherichia (E.) coli* expression host FA pool. To rule out that the

presence of glycerol in the *Tth* MAG lipase structure is due to the addition of glycerol in the cryo-

protectant buffer, we determined an additional *Tth* MAG lipase structure without using glycerol in any

sample preparation step and cryo-protection prior to X-ray data collection (**Supplementary Table 2**). In

this structure, glycerol is found in a literally identical active site position in five out of six protein chains

of the respective asymmetric unit, confirming its origin to be independent from a specific crystallization

and cryo-protection protocol **(Supplementary Fig. 7**). To independently identify the ligand content of the

extended electron density, we analyzed the crystals of the active enzyme by reversed phase liquid

chromatography followed by mass spectrometry (LC-MS). A single peak found in the elution diagram

corresponded to the glycerol monostearate (2,3-dihydroxypropyloctadecanoate) (**Supplementary Fig.**

3b), suggesting that under the given sample preparation and crystallization conditions *Tth* MAG lipase

products cannot be released.

Based on these data, we modeled a monostearate (C18) acyl ligand into the electron density of the active

Tth MAG lipase structure **(Fig. 2a & Supplementary Fig. 3a)**. Since we observed significant connecting

density between the terminal C1 carbon of the C18 monoacyl ligand and both the γ-hydroxyl oxygen of

Ser113 and the O1 oxygen of the glycerol moiety, unrestrained refinement produced distances between

these atoms in the range of 2.2-2.6 Å. The resulting model, which was confirmed by an omit electron

density map, supports the presence of a loosely coordinated tetrahedral MAG reaction intermediate [14,33,34],

closely matching the geometry of covalently bound tetrahedral reaction mimics in a previously

established MAG lipase [12] (**Fig. 3 & Supplementary Fig. 4**). We also noticed that a conserved solvent

molecule required for ester hydrolysis next to the active site triad residue H233 detected in those

structures [12] is missing in our *Tth* MAG reaction ligand complex. As *Tth* MAG lipase is exposed to an

acidic water/organic solvent mixture in LC-MS experiments after its re-solubilization from crystals

favoring esterification [35], our most plausible explanation is a reverse process from the C18 reaction

intermediate observed in the crystal structure to glycerol monostearate substrate detected in LC-MS

experiments. All subsequent experiments were designed by interpreting our structural and mass

spectrometry-based findings as C18 MAG.

The distal part of the C18 MAG reaction intermediate involving carbon atoms C11 to C18 is held by a

hydrophobic tunnel across the *Tth* MAG lipase HBH lid domain, which is open at both ends (**Fig. 2a & c**

& e). Whereas the proximal opening towards the *Tth* MAG lipase active site can be viewed as an

extension of the active site area (**Fig. 2a**), the distal lid tunnel exit is exposed and defined by a ring-like

structure of side chains from hydrophobic residues (**Fig. 2d-e**). In the presence of the C18 MAG reaction

intermediate, there are significant alterations in the packing arrangement of several residues within the lid

core when comparing to the PMSF-inhibited structure (**Fig. 2c-e**). The largest spatial deviations are found

in the N-terminal part of the lid domain, which includes the first two lid helices α-LA and α-LB (**Fig. 2d,**

Supplementary Fig. 8). In the *Tth* MAG lipase ternary complex structure these two helices are separated

by a short loop, whereas in the PMSF-inhibited structure both merge into one long helix (residues 142-

161) with a sharp kink at residue Gln153 (**Fig. 2d**). As a result, in the PMSF-inhibited structure the side

chains of Ser150 and Tyr154 next to the kink block the formation of the lid tunnel (**Fig. 2c-e**). In the

presence of the C18 MAG reaction intermediate, the aromatic ring plane of Tyr154 becomes part of the

lid tunnel surface. In conclusion, our structural data are strongly suggestive of the hydrophobic lid tunnel

to be crucial for positioning long-chain MAG esters to allow efficient turnover by lipase hydrolysis.

Mechanism for variable length MAG ester turnover

To investigate our structure-based findings suggesting the *Tth* enzyme to function as a MAG lipase, we

first measured MAD, DAG and TAG ester hydrolysis using an established coupled *in vitro* assay [12]. For

C8 MAG, we found a turnover of 16.4 +/- 1.3 U mg^{-1} by the wild-type (wt) *Tth* enzyme. In contrast, only

residual turnover was observed for the corresponding C8 DAG and TAG esters (**Fig. 4b, left panel,**

Supplementary Table 3)**,** demonstrating that the *Tth* enzyme is indeed highly specific for MAGs.

Furthermore, we only found residual activity for long chain C18:1 MAG, presumably due to limited

solubility.

We then screened a number of *Tth* MAG lipase mutants for quantitative analysis using C8 MAG as

substrate, (**Fig. 4b, right panel; Supplementary Table 3**). As expected, no activity was found for the

S113A active-site mutant, which was subsequently used as negative control. Testing further *Tth* MAG

lipase mutants without the lid domain (Δ140-183) and abolishing dimerization (E72R) also resulted in no

measurable activity, demonstrating the enzyme to be highly sensitive to structural alterations beyond the

catalytic center (**Fig. 2a-b**). We also probed two further residues, we considered to play roles in MAG

turnover. The first one was Glu43, which directly interacts with the O1 and O2 positions of the glycerol

moiety **(Fig. 2b)** and thus most likely contributes to the high level of MAG specificity as opposed to

DAG and TAG substrates. While we found about 1/3 of wt activity for the E43A variant, mutation of the

same residue to lysine (E43K) led to complete abolishment of MAG activity. This is supported by our

structural data **(Fig. 2b)**, suggesting that steric interference with the intricate interaction network between

the glycerol moiety of the MAG ester and residues from the *Tth* enzyme may impair its proper positioning

and thus substrate turnover. In contrast, removing the tyrosine group of Tyr154 (Y154A) or adding

another bulky side chain (Y154R), designed to test possible effect by residues from the *Tth* lid tunnel

(Fig. 2c-e), yielded only relative minor activity effects. These findings suggest that there is no significant

contribution of the lid to the turnover of C8 MAG, which is supported by our structural observations that

only the distal part of the long-chain C18 MAG is bound by the *Tth* lid tunnel **(Fig. 2a).** All *Tth* MAG

lipase variants were validated by circular dichroism to be properly folded as well as analyzed for thermal

stability and dimeric assembly **(Supplementary Tables 1 and 4, Supplementary Fig. 9 and 10)**.

For quantitative characterization of *Tth* MAG lipase *in vitro* activity, we subsequently employed a

spectrophotometric monoacyl *p*-nitrophenyl acyl ester (*p*NP) assay [28,36]. We used this assay for

investigating substrates with variable acyl length ranging from C2 to C18, which is equivalent to the C18

MAG reaction intermediate we found in the *Tth* MAG lipase structure (**Fig. 4a**), as a convenient approach

for direct detection of turnover and thus allowing reliable quantification. We observed a skewed activity

profile for C2-C18 esters, with the highest activities found for C4-C8 esters (**Fig. 4c, Supplementary**

Table 3). For C16 and C18 esters, there was only residual activity at a level where quantitative

significance estimates were not possible. When directly comparing C8 *p*NP ester hydrolysis with C8

MAG turnover (**Fig. 4b, left panel**), the activity was about 15%, indicating that the presence of an

unnatural *p*NP leaving group replacing the glycerol leaving group in MAG substrates could have a

diminishing effect on *Tth* enzyme substrate turnover.

At this point, we suspected that the established *Tth* activity profiles are largely convolutes of diverting

length-dependent substrate binding affinities and catalytic turnover. To address this question, we

employed steady-state Michaelis-Menten kinetics for quantitatively determining catalytic efficiency

k_{cat}/K_M, again using the *p*NP ester turnover assay due to its superior statistical reliability (**Fig. 4d-e;**

Supplementary Table 6). We found stronger binding affinities with increasing *p*NP ester length in wt

Tth MAG lipase, ranging from residual C2 ester binding (> 2 mM) to substantially higher binding

affinities of around 0.1 mM for the longest measurable *p*NP esters (C12, C14) **(Fig. 4d, left panel).** As

expected, the catalytic rate constants (k_{cat}) for C2-C14 *p*NP esters show a similar profile as for overall

activity turnover measurements **(Fig. 4d, central panel; Supplementary Table 6)**. As a result, the

catalytic efficiencies for C4-C12 *p*NP esters are all in the same range of 12-21 s^{-1} mM^{-1}, with insignificant

differences of less than a factor of 2 **(Fig. 4c, right panel; Supplementary Table 6)**. The turnover

efficiency was considerably less for the shortest substrate used in these experiments (C2) and medium- to

long-chain *p*NP esters (> C12), indicative that the catalytic efficiency of medium- to long-chain MAG

esters is hampered due to insufficient solubility under the given experimental conditions (**Supplementary**

Table 5).

Finally, we investigated the release of FAs by different *Tth* MAG lipase variants expressed in cells (*E.*

coli through the analysis of their respective fatty acid methyl esters (FAMEs). When analyzing the FA

composition using gas chromatography–mass spectrometry (GC-MS), wt and S113A mutant *Tth* MAG

lipase as well as void vector displayed the same composition with mostly hexadecanoic (palmitic) acid

(C16:0), cis-Δ11-octadecenoic (cis-vaccenic) acid (C18:1) and smaller contributions of ethylenic

(C16:1)/cyclopropanic C17 (9-10 methylene hexadecanoic) and C19 (11-12 methylene octadecanoic)

acids **(Supplementary Fig. 11).** Further, we used high performance thin layer chromatography **(Fig. 4f,**

left panel; Supplementary Fig. 11) to assess the FAMEs released in MAG lipase variants. When

comparing the data from wt *Tth* MAG lipase with the S113A negative control variant, wt *Tth* MAG lipase

displayed about two-fold higher FA amounts, demonstrating the enzyme's ability to turn over the

respective acyl esters under cellular conditions **(Fig. 4f, right panel; Supplementary Table 7).**

Interestingly, the *Tth* MAG lipase E43A variant showed a significant two-fold increase of FA turnover

with respect to wt *Tth* MAG lipase, while the same *Tth* variant in the *in vitro* MAG turnover assay

showed reduced activity when compared with the wt enzyme. In agreement with our *in vitro* data as well,

expression of the (Δ140-183) *Tth* MAG lipase variant without the lid domain led to only residual level of

FA turnover, similar to the S113A control mutant. Interestingly, the *Tth* MAG lipase Y154R variant also

showed a tendency for increased FA turnover.

Tth MAG lipase is a minimal prototype for HBH lid containing lipases

Moving beyond the specific functional properties of *Tth* MAG lipase, we wondered about the enzyme's

relations to other previously characterized lipases and esterases. A search against the UniProtKB50

protein sequence data base revealed the top scoring 1000 sequence families to be almost exclusively from

bacterial and archaeal phyla (**Supplementary Fig. 12a, Supplementary Data 1**). Amongst these

sequences were only two with known 3D structures: cinnamoyl esterase from *Lactobacillus johnsonii*

(UNIPROT code D3YEX6 [https://www.uniprot.org/uniprotkb/D3YEX6/entry], PDB code 3S2Z

[https://doi.org/10.2210/pdb3S2Z/pdb], plus additional codes of the same sequence), and feruloyl esterase

from *Butyrivibrio proteoclasticus* (UNIPROT code D2YW37 [https://www.uniprot.org/uniprotkb/

D2YW37/entry], PDB code 2WTM [https://doi.org/10.2210/pdb2WTM/pdb], plus additional codes of the

same sequence). Cinnamoyl and feruloyl esters are found as CoA ester intermediates in phenylpropanoids

metabolic pathways [37]. Even though these enzymes share a high degree of similarity with *Tth* MAG lipase

minor active site sequence differences can explain the inability of *Tth* MAG lipase to hydrolyze

phenylpropanoid ester substrates.

The relations of cinnamoyl esterase from *Lactobacillus johnsonii* (UNIPROT code D3YEX6

[https://www.uniprot.org/uniprotkb/D3YEX6/entry], PDB code 3S2Z

[https://doi.org/10.2210/pdb3S2Z/pdb], plus additional codes of the same sequence) and feruloyl esterase

from *Butyrivibrio proteoclasticus* (UNIPROT code D2YW37 [https://www.uniprot.org/uniprotkb/

D2YW37/entry], PDB code 2WTM [https://doi.org/10.2210/pdb2WTM/pdb], plus additional codes of the

same sequence) to the sequence and structure of *Tth* MAG lipase are at a comparable level, as illustrated

by rmsd values of 1.2–1.3 Å and structure-based sequence identities of 30-32 % (**Fig. 5, Supplementary**

Fig. 5, Supplementary Table 8). The lid domain in these three structures comprises a virtually identical

two-stranded antiparallel β-hairpin (residues 163-165 and 170-172 in the *Tth* MAG lipase), which is

connected by a short hairpin motif of five or less residues (**Fig. 1c, Fig. 5a**). This hairpin shields a highly

conserved histidine-glycine-phenylalanine (HGF) motif within the catalytic domain loop connecting β-

strand 3 and α-helix B (residues 35-37 in the *Tth* MAG lipase structure) and the small α-helix B'

(residues 77-79 in the *Tth* MAG lipase structure) next to the active site (**Fig. 5b, Supplementary Fig. 5**).

Helix B' comprises a phenylalanine-serine-glutamate/aspartate (FSE/D) motif that is conserved in the

sequences of the related esterases from *L. johnsonii, B. proteoclasticus*, and *T. thermohydrosulfuricus*

(*Tth*) but not in other related esterase/lipases **(Supplementary Fig. 5).** Phe37, Phe77 and Phe/Tyr175

from the C-terminal lid domain helix α-LC form a highly conserved aromatic residue cluster in these

structures and play a crucial role in shaping the substrate-binding sites of the respective enzymes (**Fig. 2a,**

Fig. 6, Supplementary Fig. 5).

Based on these relations it is surprising that *Tth* MAG lipase did not show measurable activity against

phenylpropanoid FA ester substrates of the esterases from *L. johnsonii* and *B. proteoclasticus* (**Fig. 5a**)

[40,41]. Inspection of their structures revealed a plausible explanation of the distinct substrate specificities:

First, in these esterases, there is an additional second small β-hairpin interaction following the first lid

helix α-LA, which is not found in any other HBH esterase/lipase lid domains including the structure of

Tth MAG lipase (**Fig. 5a, Fig. 6**)**.** This hairpin hovers over the central part of the active site of these two

esterases to shield bound substrates and thereby further restricts the size of the substrate binding pocket.

In the *Tth* MAG lipase this hairpin structure is replaced by a second lid helix α-LB, which is more distal

to the enzyme's active site **(Fig. 6a).** Secondly, there are two residues in the *Tth* MAG lipase lid structure,

Leu145 from the N-terminal lid helix α-LA and Phe175 from the C-terminal lid helix α-LC that are

replaced by residues with functional polar groups, aspartate (Asp138, only in the sequence of the *L.*

johnsonii esterase sequence) and tyrosine (Tyr169 in the *L. johnsonii* and *B. proteoclasticus* esterase

sequences) respectively **(Fig. 6).** Structures of the esterase from *L. johnsonii* in the presence of several

ferulate derivatives (3PFB [https://doi.org/10.2210/pdb3PFB /pdb], 3PFC

[https://doi.org/10.2210/pdb3PFC/pdb], 3S2Z [https://doi.org/10.2210/pdb3S2Z/pdb]) demonstrate the

distinct roles of these residues in substrate binding, and removal of the polar side chain of Asp138

strongly impairs substrate turnover [41]. In conclusion, the absence of these functional residue groups in the

Tth MAG lipase well explains the lack of any measurable activity towards hydroxycinnamic esters.

Next, we searched for more distantly related esterase/lipase sequences by exploiting available structural

data. The structure-based identities of all 16 selected target sequences (except those from *L. johnsonii* and

B. proteoclasticus with closer relations) are in the range of 15-23 % and the root mean square deviation

values of the superimposed structures are between 1.9 and 2.2 Å, suggesting their relations to be distant

with uncertain phylogenetic relations **(Supplementary Fig. 12b, Supplementary Table 8)**. Invariant

residue positions are confined to the active site including the catalytic triad and four glycine positions

(Gly71, Gly111, Gly115, Gly116), which also contribute to the active site oxyanion hole, another

hallmark of α/β hydrolases **(Supplementary Fig. 5)** [24,38].

Remarkably, we found an accumulation of esterase/lipase sequences with non-canonical HBH lid

domains **(Fig. 5, Supplementary Table 8, Supplementary Fig. 5).** This motif is not present in most

previously characterized MAG lipases that comprise lid domains with one or more helices without a β-

sheet inserted [22], we define here as canonical. These include human MAG lipase, which has been used as

model in various studies [15,39] **(Fig. 5a)**. Comparison of these HBH lid structures showed that the terminal

helices and adjacent loops function as flexible hinges, even in structures of the same sequence with

different ligands **(Fig. 5b)** [22,24]. On the other hand, the hydrogen bond pattern found in the conserved lid

β-hairpin provides an additional element of local fold rigidity, which is not found in other more generic

lid domains that are formed from α-helices only. While the small antiparallel β-hairpin motif is conserved

in HBH lid-containing esterases/lipases, there is substantial divergence with respect to the length of

inserted sequences **(Fig. 5)**. Different from the short hairpins found in the *Tth* MAG lipase and the most

closely related esterases from *L. johnsonii* and *B. proteoclasticus*, in other more remotely related

sequences there are various levels of extensions, allowing to cover extensive surface areas of the CD

active site. The most extreme case is found in the hydrolase-1 domain-containing protein from *B.*

bassiana, which contains an insert of 43 residues with a 12-residue helix and an extensive loop structure,

allowing to shield the complete active site of this enzyme (PDB code 7D79

[https://doi.org/10.2210/pdb7D79/pdb]) **(Fig. 5a, Supplementary Table 9**).

We finally investigated to what extent esterase/lipases with a HBH lid domain may resemble properties

we observed in the structure of *Tth* MAG lipase. Remarkably, none of the structures with a short HBH lid

domain **(Fig. 5a),** except for *Tth* MAG lipase, reveals any tunnel-like structure across the lid domain. This

is in accordance with data indicating most efficient substrate turnover for esters of small organic

compounds with additional polar groups, as well demonstrated for hydroxycinnamic acid family esterases

[40,41]. In contrast, HBH lid tunnels are observed in at least two structures with long HBH lid domains the

Paleococcus ferrophilus MAG lipase (6QE2 [https://doi.org/10.2210/pdb6QE2/pdb]) and the *Bacillus* sp.

H257 lipase (4KE8 [https://doi.org/10.2210/pdb4KE8/pdb]/4KE9

[https://doi.org/10.2210/pdb4KE9/pdb]), which both have been annotated as long-chain MAG lipases

(Supplementary Table 8). In one of these structures of the MAG lipase from *Bacillus* sp. (4KE9

[https://doi.org/10.2210/pdb4KE9/pdb]), the distal end of a 16-carbon MAG analogue binds through a

short tunnel of its HBH lid domain [12], with some resemblance of the HBH lid path found in the structure

of *Tth* MAG lipase. As key residues of the *Bacillus* sp. enzyme involved in regulation of MAG lipase

turnover (Ile145, Glu156) are positioned on segments of the long HBH lid loop insert, which is missing in

Tth MAG lipase **(Fig. 5b),** the precise mechanism of MAG lipase turnover regulation appears to be

unrelated. A more detailed structural comparison is not possible as there is only very little conservation of

residues contributing to long-chain MAG binding.

Taking the data together, HBH lid domains seem to play a crucial role in esterase/lipase function.

Supported by our *Tth* MAG lipase structural and functional data, they promote binding of especially long

chain MAG substrates. MAG lipase binding and turnover is promoted by a toolbox of specific lid domain

features: largely varying levels of extensive substrate-shielding areas as demonstrated by vastly diverting

HBH β-hairpin sequence inserts, and the formation of lid domain tunnels to hold especially the distal part

of long-chain MAG esters. Ultimately, our structural data on *Tth* MAG lipase demonstrate that the

presence of a minimal HBH lid domain is sufficient to allow turnover of long-chain MAG esters. Thus,

the structure of *Tth* MAG lipase presents a minimal prototype for a growing number of esterases and

lipases with HBH lid domains of different length. Our data illustrate that a profound understanding of

MAG lipase catalysis goes well beyond classical enzymology with well-defined active site / substrate

relations. Knowledge on enzyme-specific contributions by lid structure and dynamics next and topping

the active site are essential for our understanding of substrate specificity and possible future applications

on lipase design with further improved turnover properties. Our structure data also reveal the importance

on the way how the glycerol backbone of lipase acyl substrates is bound by specific active site residues,

ultimately defining its activity profile as MAG lipase, which is not permissive to act as a DAG or TAG

lipase. HBH lid lipases with the ability for tunnel formation, as defined in this contribution, may offer

distinct opportunities for additional evolvement to recognize promiscuous lipidic substrates of variable

length, which may even be extracted directly from lipid membranes [21,39].

In conclusion, this study presents a rare case where a high-resolution enzyme structure in the presence of

an endogenous long-chain lipid reaction intermediate led to its functional annotation as a MAG lipase.

We attribute the opportunity of capturing such complex to the extremophilic properties of its natural host

Tth, which are sufficiently different from the growth conditions of the target enzyme's heterologous

expression host (*E. coli*). Due to the high level of lid diversity and dynamics, homology modeling tools

successfully advocated for various other discoveries of enzymatic function [1,2] would have failed, as the

closest related enzymes with known structures to *Tth* MAG lipase have specific phenylpropanoid FA

esterase activities. Ultimately, this study highlights the strength of high-resolution structural biology to

identify and characterize unexplored enzymatic functions under physiological conditions that allowed

effective inhibition of substrate turnover. It may inspire future investigations of enzyme function by

systematically searching for suitable conditions, under which endogenous substrates could be directly

captured.

METHODS

Protein expression and purification

For X-ray crystallography studies, wt *Tth* MAG lipase was expressed and purified according to a

previously established protocol [28]. In this contribution, the protein was expressed using a pETBlue-1

(Novagen) vector in *E. coli* (DE3) strain. The purification was performed by liquid chromatography using

a Sepharose column followed by Size Exclusion Chromatography (SEC) step. For SEC profile analysis, a

Superdex 75 increase 3.2/300 column (Cytiva) was used. The protein was eluted in a 50 mM Tris-HCl pH

8.0 buffer containing 170 mM NaCl and the pure fractions were pooled together and concentrated up to

20 mg ml^{-1}. For production of the seleno-L-methionine (SeMet)-incorporated *Tth* MAG lipase, the

corresponding expression vector was used to transform the methionine auxotroph *E. coli* strain

B834(DE3). A preculture grown in LB broth medium was used for the expression of the protein in 2 L

M9 minimal medium including trace elements. After a 3-hour starvation step a L-aminoacid mixture was

added (40 mg ml^{-1} each except methionine) and SeMet (60 mg ml^{-1}). The culture was induced with 1 mM

isopropyl-β-D-thiogalactopyranoside (IPTG) when entering the logarithmic growth phase and was further

grown overnight. The SeMet protein was purified using the same protocol as the native protein.

To generate different protein variants for *in vitro* assays we sub-cloned the gene encoding *Tth* MAG

lipase into a pETM-14 vector (EMBL). For all single residue *Tth* variants used in this study, we

performed PCR-based mutagenesis on wt *Tth* MAG lipase. For the *Tth* MAG lipase (Δ140-183) construct

two primers were designed, the first one in forward orientation for the C-terminal segment of the catalytic

domain expressing residues 183-259 and the second one in reverse orientation for the N-terminal segment

of the catalytic domain expressing residues 1-140 (**Supplementary Table 9**). Both primers generated a

unique KpnI restriction site. PCR amplification followed by KpnI restriction and ligation yielded the *Tth*

MAG lipase (Δ140-183) construct.

The genes encoding wt *Tth*, E43A, E43K, E72R, Y154A, and Y154R were cloned into the pETM-14

expression vector, and for expression *E. coli* BL21-CodonPlus(DE3)-RIL cells (Agilent Technologies)

were used. Cells were grown at 37 °C in lysogeny broth (LB-Lennox) supplemented with kanamycin and

chloramphenicol. Cells were induced by addition of 0.5 mM IPTG at an OD_{600} of 0.6-0.8 and harvested

after continuous shaking overnight at 18 °C. For purification, the cells were resuspended in lysis buffer

(25 mM Tris-HCl (pH 8.0), 300 mM NaCl, 20 mM imidazole, and DNaseI), sonicated, and centrifuged at

40.000 g for 30 minutes at 4 °C. The supernatant was filtered and loaded onto Ni-NTA beads, which had

been equilibrated in wash buffer (25 mM Tris-HCl (pH 8.0), 300 mM NaCl, 20 mM imidazole). After

loading, beads were washed with wash buffer and eluted in elution buffer (25 mM Tris-HCl (pH 8.0), 300

mM NaCl, 500 mM imidazole). Each *Tth* variant was concentrated using an Amicon® Ultra concentrator

with a 10 kDa cutoff (membrane - regenerated cellulose) and further purified by size exclusion

chromatography, using a Superdex 75 10/300 column (GE Healthcare), equilibrated in 25 mM Tris-HCl

(pH 8.0), 50 mM NaCl.

Tth MAG lipase reductive methylation

Tth MAG lipase, purified protein was concentrated to 5 mg ml^{-1} and for each 1 ml of enzyme solution, 20

μl 1 M dimethylaminoborane (DMAB) was added followed by 40 μl 1 M formaldehyde, as previously

described [42]. DMAB and formaldehyde were added once more after 2 hours of incubation. After another 2

hours, another 10 μl 1 M DMAB was added and incubated for 18 hours. During the entire incubation time

the sample was kept in the dark and at 4 °C. To terminate the reaction, DMAB and formaldehyde were

removed via a three-step centrifugation buffer exchange to a 50 mM Tris-HCl pH 8.0 and 100 mM NaCl

buffer using a filter with a 10 kDa molecular-weight cutoff. Methylated and non-methylated *Tth* MAG

lipase were compared by SDS-PAGE (**Supplementary Fig. 13**). The level of methylation of *Tth* MAG

lipase was quantified by quadrupole time of flight (Q-TOF) mass spectrometry **(Supplementary Table**

11).

For these experiments, protein samples were acidified using 1% trifluoroacetic acid prior to injection onto

the Acquity UPLC System (Waters Corporation). Approximately 2 µg of each sample were loaded onto a

protein separation column (Acquity UPLC Protein BEH C4 column, 2.1 mm x 150 mm, 1.7 µm). The

outlet of the analytical column was coupled directly to a Q-TOF Premier mass spectrometer

(Waters/Micromass) using the standard Electrospray Ionisation source in positive ion mode. Solvent A

was water, 0.1% formic acid, and solvent B was acetonitrile, 0.1% formic acid. For Q-TOF experiments,

a spray voltage of 3.5 kV was applied with a cone voltage of 35 V and extraction cone at 5 V. The

desolvation temperature was set at 350 °C, with source temperature 120 °C. Desolvation gas was nitrogen

at a flow rate of 500 L/min. Collision energy was set at 5 eV with argon in the collision cell at a pressure

of $4.5 \cdot 10^{-5}$ mbar. Data were acquired in continuum mode over the mass range 500–3500 m/z with a scan

time of 0.5 s and an interscan delay of 0.1 s. Data were externally calibrated against an intact protein

reference standard (Thermo Fisher), acquired immediately prior to sample data acquisition. Spectra from

the chromatogram protein peak were summed and intact mass was calculated using the MaxEnt1

maximum entropy algorithm (Waters/Micromass) to give the zero charge deconvoluted molecular weight.

Catalytic activity against C8 *p*-nitrophenyl MAG ester using the assay as described in section "*In vitro*

assay to determine lipolytic enzyme activity" was close to non-methylated *Tth* MAG lipase (2,3 U mg^{-1},

cf. **Fig. 4b, left panel, and Supplementary Table 3)**.

Tth MAG lipase crystallization

Crystallization experiments using commercial screens for the sitting drop vapor diffusion technique were

performed at the EMBL-Hamburg crystallization facility at 20 ˚C [43]. Protein solution at 20 mg ml^{-1} was

mixed with the precipitant solution to a final volume of 0.4 nl (0.2 + 0.2 nl). Single rod-shaped crystals of

about 100 µm in length grew from precipitant solution containing 5% PEG-6000 in 0.1M Tris-HCl (pH

8.0) buffer.

We also used a methylated version of the protein at 20 mg ml^{-1} for *Tth* MAG lipase crystallization. Well

diffracting crystals with a maximum size of 50 x 50 x 100 µm appeared after 7-10 days using precipitant

solution of 2.4 M ammonium sulfate and 0.1 M Tris (pH 7.5) buffer. For the PMS-bound structure, the

protein was incubated for 2 hours with 10 mM PMSF followed by the same crystallization protocol.

Identical crystallization conditions were used for methylated SeMet-substituted protein.

X-ray data collection, structure solution, and validation

Prior to X-data collection, the methylated *Tth* MAG lipase crystals were immersed in mother liquor

containing 30 % (v/v) glycerol and flash cooled in liquid nitrogen. All data sets were collected at 100 K.

An anomalous dispersion experiment of a peak (λ=0.97873 Å) and an inflection (λ=0.97903 Å) data-set

using a SeMet-containing methylated *Tth* MAG lipase crystal was carried out on the tunable beamline

BM14 (EMBL/ESRF, Grenoble, France). X-ray data of PMS-inhibited *Tth* MAG lipase were also

collected on the same beamline (λ=0.95372 Å). The SeMet crystal belongs to the space group $P4_32_12$ while

the PMS-bound to the space group $P4_322$, both estimated to contain a protein dimer per asymmetric unit.

All data sets were integrated, scaled and merged using the HKL suite [44]. Selenium atoms were located

using the SHELXD program and initial density modification was performed by SHELXE [45]. The correct

heavy atom substructure was then used together with the anomalous data from HKL as input for the

PHENIX AutoSol wizard, which recalculated the experimental phases and carried out density

modification [46]. The density modified phases where then merged with the high resolution data-set from

the same SeMet crystal using the program CAD and were then used as an input for model building by

ARP/wARP [47,48]. The produced model was further improved by manual corrections and modifications in

COOT [49]. ARP/wARP was used again for adding ordered solvent molecules. Additional ligands were

added manually after inspecting the electron density maps in COOT. Molrep was used to determine the

remaining structures (PMS-bound and native)[50]. Refinement using molecular dynamics and TLS was

performed by PHENIX [51]. The data collection and refinement statistics are reported in **Supplementary**

Table 2.

Native *Tth* MAG lipase crystals were immersed in crystallization solution containing in addition 15 %

MPD as cryoprotectant and then immediately mounted at the beamline's goniometer cryo-cooled at 100

K. X-ray data to 2.42 Å resolution were collected at a wavelength λ=0.9763 Å at the EMBL/DESY

beamline BW7A at the DORIS III synchrotron storage ring integrated and scaled as above. We

determined initial phases by molecular replacement using the Se-Met substituted and lysine methylated

Tth MAG lipase coordinates (7Q4J). Structure refinement was carried out as described above with the

addition of non-crystallographic symmetry restraints.

The quality of all models was assessed by Ramachandran plots with 97.5 %, 97.2 %, 97.6% of the

residues in the most favored positions for the active Se-Met, PMS-inhibited and native structures while no

residues were present in disallowed positions (**Supplementary Fig. 6**).

A structure-based sequence alignment was computed by PDBeFold [52]. Figures of the 3D structural model

were prepared using PYMOL v.2.5.

AlphaFold2 (ColabFold version1.5.2) was used for modeling the *Tth* MAG lipase structure from the

sequence, by forcing a dimeric arrangement. The resulting models superimposed onto the *Tth* MAG lipase

crystal structures with a root mean square deviation of about 0.45 Å.

Analysis of crystal content

About 30 crystals of Se-Met *Tth* MAG lipase were carefully washed with water and dissolved in 200 µl 1

M NaCl. A chloroform/methanol mixture of 250/500 µl was then added to the solution. After 10 minutes

of intensive vortexing, 250 µl chloroform were added and the solution mixed for 1 minute. Then, 250 µl

water were added and the solution mixed again for 1 minute. The mixture was centrifuged for 10 minutes

at maximum speed and the upper water phase was carefully discarded. The organic phase was evaporated

under vacuum centrifugation at room temperature and resuspended in 20 µl hexafluoroisopropanol. LC-

MS measurements injecting 20 µl of the sample in a C18 column and applying a 50-95% acetonitrile

gradient in water, revealed a single peak eluted from the column, which in the MS spectrum gave a peak

at 359 in positive ionization mode (358+1).

Monoacylglycerol (MAG) ester assay

Enzyme activity towards MAG esters was measured as follows [53]. The protein concentration of the

purified enzyme variants was adjusted with sample buffer comprising 50 mM Tris (pH 8.0) and 100 mM

NaCl to a suitable concentration range of 0.05 to 0.005 mg ml^{-1} to perform measurements in the linear

range of the assay. 1-Octanoyl-*rac*-glycerol and 1-oleoyl-*rac*-glycerol were used as short- and long-chain

MAG substrates, respectively. 1,2-Dioctanoyl-*sn*-glycerol and tricaprylin were used as DAG and TAG

substrates. All substrates were purchased from Sigma-Aldrich if not stated otherwise. Prior to use, the

substrate stocks were diluted with assay buffer comprising 50 mM Tris/HCl (pH 8.0) to a final

concentration of 1 mM and supplemented with defatted bovine serum albumin at a molar ratio of 1:1. All

values were determined in triplicates and corrected by blank subtraction. The activity assay was

performed at 40 °C for 10 minutes followed by addition of 100 μl chloroform, centrifugation (10,000xg

for 10 minutes) and determination of the released glycerol in 10 μl of the upper phase with a commercial

Glycerol Assay Kit according to the manufacturer's instructions (Sigma-Aldrich).

Mono p-nitrophenyl acyl ester (pNP) assay

Cleavage of *p*-nitrophenyl acyl (*p*NP) esters was determined as in previous studies [28]. The protein

concentration of the purified enzyme variants was adjusted with sample buffer (25 mM Tris, pH 8.0 and

50 mM NaCl) to a suitable concentration ranging from 0.02 to 0.05 mg ml^{-1} to perform measurements in

the linear range of the assay. The substrates were stored as 10 mM stock solutions in 100 % ethanol

protected from light at -20 °C. Prior to use, the substrate stocks were diluted with assay buffer (50 mM

Tris-HCl, pH 8.0) additionally containing 1 mg ml^{-1} gum arabicum (Sigma Aldrich). All values were

determined in quadruplicates and corrected for autohydrolysis using the experimentally determined pH-

dependent extinction coefficient of *p*-nitrophenol in microtiter plates ($\varepsilon = 9.3024$ mL μmol^{-1}). The activity

assay was performed at 40 °C in a semi-automated high-throughput setup in micro-scale using clear, flat-

bottom 96-well plates (Corning). Applying a Freedom EVO® screening robot equipped with a Safire2

plate reader (Tecan Group Ltd.), pNP substrates were used at a final concentration of 0.9 mM, and

continuous measurements were performed at $\lambda = 410$ nm for 10 minutes in a total assay volume of 200 μl

per well. Blanks with sample buffer, but without enzyme, were simultaneously measured for each

substrate.

Tth MAG lipase Michaelis-Menten kinetics were performed by quadruple measurements of enzyme

activity with pNP esters at different substrate concentrations (0.005 mM, 0.0075 mM, 0.01 mM, 0.025

mM, 0.05 mM, 0.075 mM, 0.1 mM, 0.25 mM, 0.5 mM, 0.75 mM and 1 mM). The respective activity

values were fitted to the Michaelis-Menten equation applying a non-linear regression tool in Prism

version 9.1.2. Obtained K_M and v_{max} values of each Michaelis-Menten fit were then used to calculate

averages and standard deviations of these basic kinetic parameters. Finally, k_{cat} and K_M yielded the

catalytic efficiency **(Supplementary Table 6)**.

Cellular Tth MAG lipase activity by cell extract analysis

Wt *Tth* and Y154A were cloned into the pETBlue-1 vector, transformed into *E. coli* Tuner™ (DE3) pLacI

cells and grown at 37 °C in lysogeny broth (LB-Lennox) supplemented with ampicillin and

chloramphenicol. Cells were induced by addition of 1 mM IPTG at an OD_{600} of 0.6-0.8 and harvested

after continuous shaking at 37 °C for 1 hour. Cell pellets were washed with PBS prior to lipid extraction.

For lipid isolation and analysis, total lipid extracts (TLE) were prepared from wet *E. coli* cell pellets by

three successive extractions using distinct chloroform/methanol mixtures (1:2, 1:1, and 2:1 v/v). The

fractions were pooled, washed with water, and dried under air stream. For chromatography analyses, FAs

were methylated using TMS-diazomethane (Sigma), purified on a florisil batch column and analyzed by

thin layer chromatography (TLC) (Merck silica Gel 60) with dichloromethane as developing solvent.

For lipid semi-quantitative analysis, equivalent masses of total lipid fraction from each strain were spotted

onto a high-performance TLC (HPTLC) silica gel 60 plate (Merck) with a Camag ATS4 apparatus. The

plates were developed in dichloromethane using a Camag ADC2 device and stained by immersion using a

Camag CID3 apparatus with 10 % primuline in acetone/water (84:16). Lipids were revealed and

quantified by determining their relative percentage using the Chemidoc Imaging System and ImageLab

software (Biorad)**.**

To determine FA composition by GC-MS, the FAMEs were purified by preparative thin-layer

chromatography (TLC) using a silica gel coated plate G60 (Merck) with dichloromethaneide as

developing solvent. Visualization was performed by spraying the plate with rhodamine B, and FAMEs

were scraped off the plate and extracted twice with diethyl ether. The purified FAMEs were then analyzed

by GC-MS (Thermo Scientific gas chromatograph, fitted with a 15 m x 0.25 mm TG1MS fused silica

capillary column and connected to an ISQ single quadrupole mass spectrometer), and identified according

to their retention time and their fragmentation pattern [54].

Nano differential scanning fluorimetry

Temperature-induced aggregation curves were measured using Prometheus NT.48 (NanoTemper

Technologies GmbH) equipped with a scattering module. Proteins were prepared in 25 mM Tris-HCl (pH

8.0), 50 mM NaCl and measured in three technical replicates at a concentration of 1.5 mg ml^{-1}. Excitation

power was pre-adjusted to get fluorescence readings above 2000 RFU for F330 and F350, and samples

were heated from 15 °C to 95 °C with a slope of 1 °C per minute. The processed data were exported from

the manufacturer's software (PR.ThermControl) and analyzed with MoltenProt to obtain aggregation

temperatures [55].

Circular dichroism

Far-UV CD spectrums for the *Tth* samples were recorded on a Chirascan Circular Dichroism

Spectrometer upgraded with an Active Nitrogen Management System (Applied Photophysics).

Measurements were performed at a controlled temperature of 20 °C in a 1 mm QS cuvette, between 280-

190 nm using a bandwidth of 1 nm. The *Tth* samples were measured at a concentration of 0.25 mg ml^{-1} in

20 mM Tris-HCl (pH 8.0) in three technical replicates. Signal from buffer was subtracted for all measured

samples. The estimation of the secondary structure contents was calculated using CONTIN [56].

Statistics and Reproducibility

559 All enzymatic experiments were carried out at least in triplicate as follows: MAG ester assay: n=3 or 4;

560 *p*NP assay: n=3 or 4; cellular *Tth* MAG lipase activity by cell extract analysis: n=3, 4, 6 or 7; nano

561 differential scanning fluorimetry: n=3; circular dichroism: n=3. Statistical significance of the cell extract

562 analysis data **(Fig. 4f)** was evaluated using two-tailed Student's t-test and p values of less than 0.05 were

563 deemed significant. Statistical analysis was performed using GraphPad Prism 5.04.

DATA AVAILABILITY

The coordinates of the X-ray structures determined in this contribution have been deposited in the Protein

Data Bank under accession codes 7Q4J [https://doi.org/10.2210/pdb7Q4J/pdb], 7Q4H

[https://doi.org/10.2210/pdb7Q4H/pdb], 8B9S [https://doi.org/10.2210/pdb8B9S/pdb]. All other data

generated in this study are provided in the Supplementary Information and a Source Data file.

The coordinates of the following X-ray structures used for analyzing the results in this work are also

available from the Protein Data Bank with accession codes: 3S2Z [https://doi.org/10.2210/pdb3S2Z/pdb],

2WTM [https://doi.org/10.2210/pdb2WTM/pdb], 7D79 [https://doi.org/10.2210/pdb7D79/pdb], 3HJU

[https://doi.org/10.2210/pdb3HJU/pdb], 3LLC [https://doi.org/10.2210/pdb3LLC/pdb], 4KE8

[https://doi.org/10.2210/pdb4KE8/pdb], 4KE9 [https://doi.org/10.2210/pdb4KE9/pdb], 3PFB

[https://doi.org/10.2210/pdb3PFB/pdb], 3PFC [https://doi.org/10.2210/pdb3PFC/pdb], 5CML

[https://doi.org/10.2210/pdb5CML/pdb], and 6QE2 [https://doi.org/10.2210/pdb6QE2/pdb]).

REFERENCES

1.   Jacobson, M.P., Kalyanaraman, C., Zhao, S. & Tian, B. Leveraging structure for enzyme function
prediction: methods, opportunities, and challenges. *Trends Biochem Sci* **39**, 363-71 (2014).

2.   Mills, C.L., Beuning, P.J. & Ondrechen, M.J. Biochemical functional predictions for protein
structures of unknown or uncertain function. *Comput Struct Biotechnol J* **13**, 182-91 (2015).

3.   Bastard, K. et al. Revealing the hidden functional diversity of an enzyme family. *Nat Chem Biol*
10, 42-9 (2014).

4.   Calhoun, S. et al. Prediction of enzymatic pathways by integrative pathway mapping. *Elife* **7**(2018).

5.   Zhao, S. et al. Discovery of new enzymes and metabolic pathways by using structure and genome
context. *Nature* **502**, 698-702 (2013).

6.   Carter, M.S. et al. Functional assignment of multiple catabolic pathways for D-apiose. *Nat Chem*
Biol **14**, 696-705 (2018).

7.   Sévin, D.C., Fuhrer, T., Zamboni, N. & Sauer, U. Nontargeted in vitro metabolomics for high-
throughput identification of novel enzymes in Escherichia coli. *Nat Methods* **14**, 187-194 (2017).

8.   Zhao, S. et al. Prediction and characterization of enzymatic activities guided by sequence similarity
and genome neighborhood networks. *Elife* **3**(2014).

9.   Lee, C.W. et al. Structural basis for the ligand-binding specificity of fatty acid-binding proteins
(pFABP4 and pFABP5) in gentoo penguin. *Biochem Biophys Res Commun* **465**, 12-8 (2015).

10.   Lücke, C., Qiao, Y., van Moerkerk, H.T., Veerkamp, J.H. & Hamilton, J.A. Fatty-acid-binding
protein from the flight muscle of Locusta migratoria: evolutionary variations in fatty acid binding.
Biochemistry **45**, 6296-305 (2006).

11.   Vasiliauskaité-Brooks, I. et al. Structural insights into adiponectin receptors suggest ceramidase
activity. *Nature* **544**, 120-123 (2017).

12.   Rengachari, S. et al. Conformational plasticity and ligand binding of bacterial monoacylglycerol
lipase. *J Biol Chem* **288**, 31093-104 (2013).

13.   Kim, J. et al. Structure-guided discovery of the metabolite carboxy-SAM that modulates tRNA
function. *Nature* **498**, 123-6 (2013).

14.   Jaeger, K.E., Dijkstra, B.W. & Reetz, M.T. Bacterial biocatalysts: molecular biology, three-
dimensional structures, and biotechnological applications of lipases. *Annu Rev Microbiol* **53**, 315-
51 (1999).

15.   Labar, G., Wouters, J. & Lambert, D.M. A review on the monoacylglycerol lipase: at the interface
between fat and endocannabinoid signalling. *Curr Med Chem* **17**, 2588-607 (2010).

16.   Salhi, A., Carriere, F., Grundy, M.M.L. & Aloulou, A. Enzymes Involved in Lipid Digestion. in
Bioaccessibility and Digestibility of Lipids from Food (eds. Grundy, M.M.L. & Wilde, P.J.) 3-28
(Springer International Publishing, Cham, 2021).

17.   Jaeger, K.E. & Eggert, T. Lipases for biotechnology. *Curr Opin Biotechnol* **13**, 390-7 (2002).

18.   Borrelli, G.M. & Trono, D. Recombinant Lipases and Phospholipases and Their Use as Biocatalysts
613   for Industrial Applications. *Int J Mol Sci* **16**, 20774-840 (2015).

614   19.   Anobom, C.D. et al. From structure to catalysis: recent developments in the biotechnological
615   applications of lipases. *Biomed Res Int* **2014**, 684506 (2014).

616   20.   Reis, P., Watzke, H., Leser, M., Holmberg, K. & Miller, R. Interfacial mechanism of lipolysis as
617   self-regulated process. *Biophys Chem* **147**, 93-103 (2010).

618   21.   Aloulou, A. et al. Exploring the specific features of interfacial enzymology based on lipase studies.
619   *Biochim Biophys Acta* **1761**, 995-1013 (2006).

620   22.   Khan, F.I. et al. The Lid Domain in Lipases: Structural and Functional Determinant of Enzymatic
621   Properties. *Front Bioeng Biotechnol* **5**, 16 (2017).

622   23.   Aschauer, P. et al. Crystal structure of the Saccharomyces cerevisiae monoglyceride lipase Yju3p.
623   *Biochim Biophys Acta* **1861**, 462-70 (2016).

624   24.   Bauer, T.L., Buchholz, P.C.F. & Pleiss, J. The modular structure of α/β-hydrolases. *Febs j* **287**,
625   1035-1053 (2020).

626   25.   López-López, O., Cerdán, M.E. & González Siso, M.I. New extremophilic lipases and esterases
627   from metagenomics. *Curr Protein Pept Sci* **15**, 445-55 (2014).

628   26.   Salameh, M.A. & Wiegel, J. Purification and characterization of two highly thermophilic alkaline
629   lipases from Thermosyntropha lipolytica. *Appl Environ Microbiol* **73**, 7725-31 (2007).

630   27.   Littlechild, J.A. Improving the 'tool box' for robust industrial enzymes. *J Ind Microbiol Biotechnol*
631   **44**, 711-720 (2017).

632   28.   Royter, M. et al. Thermostable lipases from the extreme thermophilic anaerobic bacteria
633   Thermoanaerobacter thermohydrosulfuricus SOL1 and Caldanaerobacter subterraneus subsp.
634   tengcongensis. *Extremophiles* **13**, 769-83 (2009).

635   29.   Klingeberg, M., Hippe, H. & Antranikian, G. Production of novel pullulanases at high
636   concentrations by two newly isolated thermophilic clostridia. *FEMS Microbiol Lett* **57**, 145-52
637   (1990).

638   30.   Jørgensen, B.B. & Cohen, Y. Solar Lake (Sinai). 5. The sulfur cycle of the bcnthic cyanobacterial
639   mats 1. *Limnology and oceanography* **22**, 657-666 (1977).

640   31.   Mirdita, M., Steinegger, M. & Söding, J. MMseqs2 desktop and local web server app for fast,
641   interactive sequence searches. *Bioinformatics* **35**, 2856-2858 (2019).

642   32.   Jumper, J. et al. Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589
643   (2021).

644   33.   Holmquist, M. Alpha/Beta-hydrolase fold enzymes: structures, functions and mechanisms. *Curr
645   Protein Pept Sci* **1**, 209-35 (2000).

646   34.   Reis, P., Holmberg, K., Watzke, H., Leser, M.E. & Miller, R. Lipases at interfaces: a review. *Adv
647   Colloid Interface Sci* **147-148**, 237-50 (2009).

648   35.   Fischer, F., Mutschler, J. & Zufferey, D. Enzyme catalysis with small ionic liquid quantities. *J Ind
649   Microbiol Biotechnol* **38**, 477-87 (2011).

650   36.   Winkler, U.K. & Stuckmann, M. Glycogen, hyaluronate, and some other polysaccharides greatly
651   enhance the formation of exolipase by Serratia marcescens. *J Bacteriol* **138**, 663-70 (1979).

652   37.   Barros, J. et al. Role of bifunctional ammonia-lyase in grass cell wall biosynthesis. *Nat Plants* **2**,
653   16050 (2016).

38. Nardini, M. & Dijkstra, B.W. Alpha/beta hydrolase fold enzymes: the family keeps growing. *Curr*
Opin Struct Biol **9**, 732-7 (1999).

39. Scalvini, L., Piomelli, D. & Mor, M. Monoglyceride lipase: Structure and inhibitors. *Chem Phys*
Lipids **197**, 13-24 (2016).

40. Goldstone, D.C. et al. Structural and functional characterization of a promiscuous feruloyl esterase
(Est1E) from the rumen bacterium Butyrivibrio proteoclasticus. *Proteins* **78**, 1457-69 (2010).

41. Lai, K.K. et al. An inserted α/β subdomain shapes the catalytic pocket of Lactobacillus johnsonii
cinnamoyl esterase. *PLoS One* **6**, e23269 (2011).

42. Rypniewski, W.R., Holden, H.M. & Rayment, I. Structural consequences of reductive methylation
of lysine residues in hen egg white lysozyme: an X-ray analysis at 1.8-A resolution. *Biochemistry*
32, 9851-8 (1993).

43. Mueller-Dieckmann, J. The open-access high-throughput crystallization facility at EMBL
Hamburg. *Acta Cryst*. **D62**, 1446-1452 (2006).

44. Otwinowski, Z. & Minor, W. Processing of X-ray diffraction data collected in oscillation mode.
Macromolecular Crystallography, Pt A **276**, 307-326 (1997).

45. Sheldrick, G.M. A short history of SHELX. *Acta Crystallogr A* **64**, 112-22 (2008).

46. Terwilliger, T.C. et al. Iterative model building, structure refinement and density modification with
the PHENIX AutoBuild wizard. *Acta Crystallographica Section D* **64**, 61-69 (2008).

47. Langer, G., Cohen, S.X., Lamzin, V.S. & Perrakis, A. Automated macromolecular model building
for X-ray crystallography using ARP/wARP version 7. *Nat Protoc* **3**, 1171-9 (2008).

48. Winn, M.D. et al. Overview of the CCP4 suite and current developments. *Acta Crystallogr D Biol*
Crystallogr **67**, 235-42 (2011).

49. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Crystallogr D*
Biol Crystallogr **60**, 2126-32 (2004).

50. Vagin, A. & Teplyakov, A. Molecular replacement with MOLREP. *Acta Crystallogr D Biol*
Crystallogr **66**, 22-5 (2010).

51. Afonine, P.V. et al. Towards automated crystallographic structure refinement with phenix.refine.
Acta Crystallogr D Biol Crystallogr **68**, 352-67 (2012).

52. Krissinel, E. & Henrick, K. Secondary-structure matching (SSM), a new tool for fast protein
structure alignment in three dimensions. *Acta Crystallogr D Biol Crystallogr* **60**, 2256-68 (2004).

53. Heier, C. et al. Identification of Yju3p as functional orthologue of mammalian monoglyceride
lipase in the yeast Saccharomycescerevisiae. *Biochim Biophys Acta* **1801**, 1063-71 (2010).

54. Lanéelle, M.A., Spina, L., Nigou, J., Lemassu, A. & Daffé, M. Lipid and Lipoarabinomannan
Isolation and Characterization. *Methods Mol Biol* **2314**, 109-150 (2021).

55. Kotov, V. et al. In-depth interrogation of protein thermal unfolding data with MoltenProt. *Protein*
Sci **30**, 201-217 (2021).

56. Provencher, S.W. & Glöckner, J. Estimation of globular protein secondary structure from circular
dichroism. *Biochemistry* **20**, 33-7 (1981).

ACKNOWLEDGEMENTS

We acknowledge the technical support by the Sample Preparation and Characterization facility of the

EMBL Hamburg Unit, the EMBL Proteomics Core Facility and the Proteomics Core Unit of the

University of Münster, Germany. We also acknowledge the contributions of Marina Royter during the

early phase of the project, as well as the technical contributions of Anke Peters and Barbara Klippel, all

from the Hamburg University of Technology, as well as Lucie Spina and Marie-Antoinette Laneelle from

the IPBS, University of Toulouse, CNRS, UPS, for contribution in methodology. We are grateful to Dr

Martin Walsh at the European Synchrotron Radiation Facility (ESRF), Grenoble, France, for providing

assistance in using the beamline BM14 for our X-ray data collection experiments. We thank Sihyun Sung,

Grzegorz Chojnowski and Barbara Ramsak from the EMBL Hamburg Unit for technical and scientific

support, and Chris Meier from the University of Hamburg for advice during the manuscript revision

process.

AUTHOR CONTRIBUTIONS

NP, AK, HM and MW designed the project. NP, AK, NT, SDC and CL performed the experiments. NP,

AK, SDC, SAM, HM and MW analyzed the data. NP, AK, SAM, HM and MW wrote the manuscript.

MD, AG and MW supported the project.

COMPETING INTERESTS

The authors declare no competing interests.

Fig. 1. Biophysical and structural properties of *Tth* MAG lipase. a, Size exclusion chromatograms

indicating dimeric (D) and monomeric (M) fractions and b, nano differential scanning fluorimetry profiles

of selected *Tth* MAG lipase variants. For further analysis, see **Supplementary Table 1**. c, topology

diagram of the molecular structure of the *Tth* MAG lipase protomer. CD, yellow; lid domain, pink.

Secondary structural elements are in proportion to their length and are labeled. The positions of key active

site residues (***cf*. Supplementary Fig. 5)** are highlighted. The active site glycerol interacting residue

Glu43 is shown in red. While in the text three-letter codes were used for residue specifications, for the

reason of clarity one-letter codes were used in all Figures. d, surface presentation of *Tth* MAG lipase

protomer, with the dimeric interface surface highlighted in stronger colors. Polar residues that contribute

to specific interactions to the dimeric interface are shown in red (glutamate, aspartate), blue (arginine,

lysine), and pale blue (glutamine), and are labeled. Glu72, which was mutated into an arginine for probing

the dimeric interface, is highlighted by an asterisk. The surface of the active site residue Ser113 is shown

in black, as point of reference for locating the active site. e, *Tth* MAG lipase dimer in surface

presentation. The catalytic domains (CD) and lid domains of each protomer are labeled and colored

separately. The two *Tth* MAG lipase protomers form a large joint substrate binding site, as indicated by

the bound C18 MAG ligands **(Fig. 2a)** and two asterisks in red, indicating the two active sites. In this

presentation, the proximal part of the C18 MAG ligands is visible (***cf*. Fig. 2)**. The orientation of *Tth*

MAG lipase protomer A is rotated by about 90 degrees around a vertical axis with respect to the *Tth*

MAG lipase monomer shown in panel d, as indicated. Source data are provided as a Source Data file.

Fig. 2: *Tth* MAG lipase active site and lid domain tunnel for long chain MAG esters. a, surface
presentation of the *Tth* MAG lipase long-chain MAG binding site, in part generated by a tunnel across the
lid domain. Bound C18 MAG intermediate is shown in green. Atom-specific colors are used for oxygen
(red) and nitrogen (blue). Every fifth methylene group of C18 MAG is numbered. Those parts of the C18
MAG that are sequestered by the lid domain tunnel (C11-C18) appear to be darker due to the semi-
transparent *Tth* MAG lipase surface. Side chains of key residues contributing to ligand binding and the
Tth active site are shown and labeled. Residues used for probing function are labeled with an asterisk. The
central part of the active site is zoomed in a box to the right, indicating specific polar interactions of the
glycerol product with *Tth* MAG lipase Glu43. b, (upper inlet), active site of the PMSF-inhibited *Tth* MAG
lipase structure for comparison. c, mesh surface representation of the C18 MAG lid domain tunnel from
the top of the lid domain, indicating that it is open at the most distal end (left panel), side view of the lid
tunnel (central panel) similar to the orientation in a. Right panel: for comparison, part of the surface
generated by residues Phe37 from the CD and Tyr154 from the lid domain is shown for the distinct
conformation of the PMSF-inhibited *Tth* MAG lipase structure, due a conformational change of Tyr154,
which is labeled in red. This change leads to blockage of the lid tunnel found in the *Tth* MAG lipase
structure without PMSF. d, superposition of *Tth* MAG lipase lid domain structures, without (violet) and
with PMSF (grey). e, surface presentation of *Tth* MAG lipase lid domain structures, without (violet, left
panel) and with PMSF (grey, right panel). Residues involved in the formation of the proximal lid
substrate tunnel opening and polar residues forming an outer half-ring wrapping the substrate tunnel exit
are labeled. The proximal end of the C18 MAG reaction intermediate bound to the *Tth* MAG lipase
structure without PMSF is shown in green. Comparison of the two conformations shows that the lid
tunnel is only opened in the *Tth* MAG lipase structure without PMSF, when C18 MAG ester ligand is
present. For reasons of optimizing illustration of the lid tunnel the orientation is different from the one in
panel b.

Fig. 3: *Tth* MAG lipase reaction scheme. Reaction intermediates are indicated. Color codes: acyl group,

pink; glycerol group, blue; reactive water, green; side chain of Glu43, red. The ligand density found C18

MAG ester reaction intermediate found in the *Tth* MAG lipase structure active site **(Supplementary Fig.**

3) is closest to loosely coordinated tetrahedral transition intermediate 1 (boxed).

Fig. 4: *Tth* MAG lipase activities for MAG esters. a, *Tth* MAG lipase activities for MAG esters of
variable length. Coloring scheme of esters characterized *in vitro* assays (**b-d**). MAG esters found in cell
extracts (**f**) and by structural data (**Fig. 2, Supplementary Fig. 3**) are also indicated, demonstrating
complementarity of different experimental approaches. b, left panel: *in vitro* turnover of C8 MAG, DAG
and TAG esters and C18:1 MAG. b, right panel: C8 MAG turnover of different *Tth* enzyme variants. c,
Tth MAG lipase activity for several *p*NP acyl esters of C2-C18 FAs and ferulate, indicating a broad range
of possible substrates. d, steady-state Michaelis-Menten analysis for MAG esters with quantifiable

activity profiles. Left panel, apparent substrate binding affinities for C2-C14 MAG esters measured by

Michaelis constants K_M, indicating increase of binding with increasing MAG ester length; central panel,

catalytic rates (k_{cat}) for C2-C14 MAG esters, showing an overall profile comparable to that measured for

overall substrate turnover (central panel, *cf*. panel b, left); right panel, catalytic efficiencies k_{cat}/K_M for the

same substrate spectrum, showing only minor variations for most of the MAG esters used (C4-C12). e,

left panel: experimental steady-state kinetic data of selected *Tth* variants (wt, Y154A, E43A) to estimate

Michaelis-Menten values for C6 MAG esters. f, Fatty Acid Methyl Esters (FAMEs) analysis by high-

performance thin-layer chromatography (HPTLC) of *E. coli* total lipid extracts (TLE) expressing *Tth*

MAG lipase. Left panel, 1, *Tth* MAG lipase (wt); 2, *Tth* MAG lipase (S113A); 3, *Tth* MAG lipase

(E43A). Right panel: FAMEs quantification for selected *Tth* MAG lipase mutants, normalized to *Tth*

MAG lipase (wt). Data statistics (b-e): Data are presented as mean values and individual data points,

$n = 3$-4 (technical replicates). Data statistics (f): Data are presented as means ± SD, and individual data

points. Error bars represent the standard deviation from biologically independent replicates (*n*=3 for

variants S113A, E43A, Y154R; *n*=4 for variants E43K, Y154A; *n*=6 for WT; and *n*=7 for ΔLid).

Unpaired t-test was used to evaluate statistical significance applying p values for a two-tailed confidence

interval as follows: $p < 0.001$ (***), $p < 0.01$ (**), $p < 0.05$ (*). Definition p values**:** p=0.0055 ΔLid,

p=<0.0001 S113A, p=0.0288 E43A, p=0.0013 E43K, p=0.4985 Y154A, p=0.1285 Y154R. Source data

and uncropped image are provided as a Source Data file.

a

7Q4J *T. thermohydrosulfuricus* **Monoacylglycerol lipase**
3S2Z *L. johnsonii* Cinnamoyl esterase
2WTM *B. proteoclasticus* Feruloyl esterase
5CML *R. marinus* Carboxyl esterase
3LLC *A. vitis* Hydroxymuconic semialdehyde hydrolase
6NY9 Mouse Palmitoyl-protein thioesterase A
6QE2 *P. ferrophilus* **Monoacylglycerol lipase**
4KE8 *B. sp. H257* **Monoacylglycerol lipase**
6EIC *M. tuberculosis* **Monoacylglycerol lipase**
7D79 *B. bassiana* Polyketide Cyclase
3HJU *Homo sapiens Monoacylglycerol lipase*

b

Fig. 5: Sequence, structural and functional relations of esterase/lipases with divergent HBH lid

domains. a, Scheme of HBH lid domain sequences related to *Tth* MAG lipase (**Fig. 1c**) with secondary

structural elements indicated proportional to sequence length. The two-stranded HBH lid β-hairpins are in

cyan, sequence and secondary structural elements inserted between strands βL1 and βL2 are in green. The

additional two-stranded β-hairpin structure hovering over the active site in the esterase from *L. johnsonii*

(PDB code: 3S2Z [https://doi.org/10.2210/pdb3S2Z/pdb]) is shown in dark blue. The related β-hairpin-

like structure in the esterase from *B. proteoclasticus* (PDB code: 2WTM

[https://doi.org/10.2210/pdb2WTM/pdb]) is shown in the same color with a diagonal stripe pattern. The

lid structure of the Osmotically inducible protein C from *R. marinus* (PDB code: 5CML

[https://doi.org/10.2210/pdb5CML/pdb]) comprises an additional third β-strand colored in dark violet,

which interacts with the first β-strand of the HBH β-hairpin. Human MAG lipase (3HJU

[https://doi.org/10.2210/pdb3HJU/pdb]), which does not have a HBH lid motif and presents a canonical

MAG lipase, is shown for comparison as well. Further details are summarized in **Supplementary Table**

8. b, HBH lid structures of three representative esterase/lipases with their PDB codes indicated

(**Supplementary Fig. 12b**), demonstrating common elements (N- and C-terminal α-helices, two-stranded

β-hairpin) and diversity of the inserted sequences in terms of length and structure. The CDs of these

structures are shown by surface presentation in light orange, and the HBH domains are illustrated by

cartoon presentation. The surface covered by the catalytic triad serine (Ser113 in *Tth* MAG lipase) is

shown in red, as a point of reference to locate the active site. Active site ligands are shown in sphere

presentation in atom-specific colors (carbon, green; oxygen, red; nitrogen, blue). Only in *Tth* MAG lipase,

the C18 ligand crosses the lid domain via a hydrophobic tunnel. The ligands in 3LLC

[https://doi.org/10.2210/pdb3LLC/pdb] (tetraethylene glycol) and 4KE8

[https://doi.org/10.2210/pdb4KE8/pdb] (tetradecyl hydrogen (R)-(3-azidopropyl)phosphonate, chain A)

are in similar positions next to the active site but they are too short to cross the respective lid domains

(Supplementary Fig. 4a).

Fig. 6: Structural comparison of the ligand bound HBH lid domains and active site area, indicating different substrate specificities. a, *Tth* MAG lipase in the presence of C18 MAG (left); b, esterase from *L. johnsonii* (PDB code: 3S2Z) in the presence of ferulate (right). Two residues critical for specific hydrogen bond interactions with ferulate in structure of the esterase from *L. johnsonii* (Asp138, Tyr169) are substituted by hydrophobic residues in *Tth* MAG lipase (equivalent residues Leu145, Phe175). Other key residues with critical roles for substrate binding are also shown. In the structure of the esterase from *L. johnsonii*, an additional N-terminal lid β-sheet is formed (colored in blue) that hovers over the CD active site. At the end of the short turn connecting the two β-strands, there is a highly exposed residue, which is Gln145 in this structure. Residues that are part of the conserved HGF motif (residues 35-37 in *Tth* MAG lipase, residues 32-34 in the *L. johnsonii* esterase) and less conserved FSE/D motif (residues 77-79 in *Tth* MAG lipase, residues 70-72 in in the *L. johnsonii* esterase) are labeled in red (**Supplementary Fig. 2).**