Spatial heterogeneity alters the trade-off between growth and dispersal during a range expansion

Patrizia Zamberletti[1,*]

Lionel Roques[1]

Florian Lavigne[2]

Julien Papaïx[1]

1. INRAE, BioSP, 84914, Avignon, France;

2. Université de Rouen Normandie, CNRS, Laboratoire de Mathématiques Raphaël Salem, France;

∗ Corresponding author; e-mail:patrizia.zamberletti@inrae.fr.

Keywords: Eco-evo model, Reaction-diffusion system, Evolutionary R-D trade-off, fragmented space, nonlocal competition.

Abstract

Individuals who invest more in the development of their dispersal-related traits often reduce their investment in reproduction. Thus, there are two possible eco-evolutionary strategies: grow faster or disperse faster ($R - D$ arbitrage). Here we explore, through a reaction-diffusion model, how spatial heterogeneity can shape the $R - D$ trade-off by studying the spreading dynamics of a consumer species exploiting a resource in a spatially fragmented environment. Based on numerical simulations and analytical solutions derived from simpler models, we show that the classical mathematical symmetry between the effects of growth and dispersal on the spatial spreading speed is broken in the presence of competition between phenotypes. At the back of the forefront, the dynamics is almost always driven by the R specialists. On the forefront, R-strategies are favored in spatially homogeneous environments, but the introduction of heterogeneity leads to a shift towards D-strategies. This effect is even stronger when spatial heterogeneity affects the diffusion term and when spatial fragmentation is lower. Introducing mutations between phenotypes produces an advantage towards the R-strategy and homogenizes the distribution of phenotypes, also leading to more polymorphism on the forefront.

1 Introduction

Rapid evolution in species traits can affect their ecological dynamics which in turn feedback on the evolutionary potential (Bonte and Bafort, 2019; Burton et al., 2010). Such interaction between ecological and evolutionary dynamics is crucial to understand the demography when species shift their range as in the case of evolutionary rescue (Anciaux et al., 2019), migrational meltdown (Ronce and Kirkpatrick, 2001), biological invasion (Szűcs et al., 2019). Population expansion is an ecological process mainly driven by traits related to reproduction and dispersal (Deforet et al., 2019; Turchin, 1998). Dispersal affects capabilities to exchange individuals and genes among different habitats (Legrand et al., 2017). Dispersal traits have been proven to be related to body dimension and condition (Duthie et al., 2015; Helms and Kaspari, 2015; Steenman et al., 2015), affecting competitive abilities, food web interactions (Bonte and de la Pena, 2009) or metabolic processes (Hirt et al., 2017). As a consequence, there are many examples where individuals who invest more in the development of their traits related to the dispersal strategy reduce the effort in foraging and reproduction (*e.g.*, reducing their mating period or with lower egg mass) (Baguette and Schtickzelle, 2006; Bonte and Bafort, 2019; Hanski et al., 2006). In such cases, two possible evolutionary strategies exist: dispersing faster or growing stronger (Deforet et al., 2019). This results in a species' trait trade-off that shapes the ecological and evolutionary dynamics of populations.

During invasion process, there is evidence that trait evolution can be very rapid altering demographic processes (Griette et al., 2015; Perkins et al., 2013). At the forefront, *i.e.*, in the frontmost part of the population range, studies highlight that spread rate jointly depends on population growth and dispersal, and that the evolution of these traits can

results in an accelerating spread (Fisher, 1937; Perkins et al., 2013). For example, Perkins et al. (2013) focused on how life-history or dispersal traits impact spread rates of the cane toad, *Rhinella marina*, in Australia by combining a stage-structured population dynamics model and an evolutionary quantitative genetic model. They pointed out that rapid evolution of life-history and dispersal traits at the forefront could have led to a more than twofold increase in the distance spread by cane toads across northern Australia. Indeed, spatial sorting of high-dispersal individuals drove dispersal evolution at the forefront and may have resulted in the accumulation of individuals with extreme dispersal abilities at its edge, accelerating invasion (Bouin et al., 2012; Perkins et al., 2013; Shine et al., 2011). However, the individuals leading the forefront should also face novel evolutionary pressures on reproduction, due to low population densities (Kelehear and Shine, 2020). An example of the interactions between dispersal and other key life history traits, such as reproduction, is wing polymorphism of various species of insects (Zera and Denno, 1997). The flight capability (defined by developed wings and flight muscles) is negatively correlated with age at first reproduction and fecundity (Denno, 1994). Thus, the energy efforts for flight and reproduction lead to a trade-off for internal resources (Zera and Denno, 1997).

Reaction-diffusion models are particularly well suited to the study of biological invasions and range expansions in general (Shigesada and Kawasaki, 1997; Turchin, 1998), and to mathematically formalize the relationship between species life-history traits and expansion speed. The first spatio-temporal models of this type considered a homogeneous environment and neglected adaptation (Skellam, 1951). In this case, if the population is initially concentrated in a bounded region, the organisms spread with a speed equal to $2\sqrt{RD}$ (Fisher, 1937; Kolmogorov et al., 1937), where R is the intrinsic growth

rate of the population and D is the diffusion coefficient which measures the dispersal capacities of the individuals. The population density tends to keep a constant profile: it converges to a traveling wave. The reaction-diffusion framework can be easily adapted to take into account spatial and/or temporal heterogeneities (Shigesada and Kawasaki, 1997). Several theoretical studies considered such models, and proposed a generalization of the notion of traveling wave to spatially-fragmented environments (Berestycki and Hamel, 2002, 2005; Berestycki et al., 2005; Weinberger, 2002). These studies, and other references that we mention in the following sections, have provided a detailed understanding of the dependence of the spreading speed on spatial fragmentation, according to the particular traits they affect (R, D or both), in the absence of adaptation. In particular, very different effects of fragmentation have been observed, depending on whether they affect R or D (Hamel et al., 2011).

Some recent works have proposed to take into account genetic adaptation in these spatio-temporal models, thanks to an additional variable, say y (interpreted as a phenotypic trait), a mutation term modeled with a Laplace diffusion operator, and a nonlocal selection term (Alfaro et al., 2017, 2013; Alfaro and Peltier, 2021; Peltier, 2020). These models describe adaptation along an environmental gradient, that is, a gradual change in various factors in space that determine the phenotypic traits that are favored by their growth rate $R(x, y)$. Here, each spatial position x is associated with a different optimal trait, *i.e.*, a trait which leads to a maximal growth rate. The value of this optimal trait may be proportional to the position (Alfaro et al., 2013; Peltier, 2020), may depend periodically on x (Alfaro and Peltier, 2021), or may change with time (Alfaro et al., 2017). Another important part of this literature has been interested in the case where the trait is the diffusion coefficient D (Benichou et al., 2012; Berestycki et al., 2015; Bouin and Calvez,

2014; Bouin et al., 2012), and mostly focused on the acceleration of the range expansion in this case, due to the selection of the individuals with enhanced dispersal abilities. The objective of these works was mainly to explain the acceleration of the range expansion of can toads since their introduction in Australia, and the corresponding model is often referred to as the "cane toad equation". Recently, this framework has been applied to other traits such as the Allee threshold (Alfaro et al., 2021). In all of these reaction-diffusion based models, the additional phenotypic variable only affects a single biological parameter, either directly when this variable is the trait itself such as the diffusion term D or the Allee threshold, or indirectly when the growth rate $R(x, y)$ depends on an abstract trait y. This means that trade-off between traits are not considered. Recently, Bouin et al. (2018) considered such a trade-off between dispersal and growth in the cane toad equation. They mainly focused on theoretical mathematical results and on the occurrence of acceleration, in a homogeneous environment, depending on the rate of increase of the mortality term when the diffusion term is increased.

In this work, we develop a reaction-diffusion model to describe the phenotype-space-time dynamics of a consumer species in a fragmented space during a range expansion. We focus on the trade-off between the growth rate $R(x, y)$ and dispersal rate $D(x, y)$, which are both defined as functions of the space variable x and the phenotype variable y. In a spatially homogeneous environment and in the absence of mutations and Allee effects, the standard formula $V = 2\sqrt{R\,D}$ clearly shows that growth and dispersal play a similar role on the spreading speed (Kolmogorov et al., 1937). We analyze here how this symmetry in the effects of R and D may be broken when facing spatial fragmentation, in the presence of competition between phenotypic traits or in the presence of mutations.

2   Model and methods

2.1   *Eco-evolutionary dynamics*

{model}

At time t and location x, the density of the consumer phenotype y is defined by $c(t, x, y)$. We describe the spatial dispersion in a one-dimensional environment with a Laplace diffusion operator, corresponding to random walk movements of the individuals, with a mobility parameter (also called diffusion coefficient) $D(x, y)$ (Shigesada and Kawasaki, 1997; Turchin, 1998). We assume a one-dimensional phenotype $y \in (y_{\min}, y_{\max})$. The mutations between phenotypes are also described with a Laplace diffusion approximation (Hamel et al., 2020; Tsimring et al., 1996) with constant mutation coefficient $\mu \geq 0$. The mutation coefficient μ is proportional to the mutation rate (per individual per generation) and to the average mutation effect on phenotype (Hamel et al., 2020). Finally, the population grows logistically with a spatially variable growth rate $R(x, y)$. Competition occurs locally on the geographical space but globally over phenotypes though a nonlocal term, and is modulated by a parameter γ. This leads to the following reaction-diffusion model for the phenotype-space-time dynamics of the consumer population:

$$\partial_t c(t, x, y) = \partial_{xx}(D(x, y) \, c(t, x, y)) + \mu \, \partial_{yy} c(t, x, y)$$
$$+ c(t, x, y) \left(R(x, y) - \gamma \int_{y_{\min}}^{y_{\max}} c(t, x, s) ds \right), \quad (1)$$

with $t > 0$, $x \in \mathbb{R}$ and $y \in (y_{\min}, y_{\max}) \subset \mathbb{R}$. In all cases, we assume an initial condition $c(0, x, y) = \mathbb{1}_{x<0}$, the characteristic function of the domain $(x, y) \in (-\infty, 0) \times (y_{\min}, y_{\max})$, and we focus on the spreading of the solution to the right, that is in the direction of positive x. In addition, we assume no-flux boundary conditions at the bound-

aries $y = y_{\min}$ and $y = y_{\max}$:

$$\partial_y c(t, x, y_{\min}) = \partial_y c(t, x, y_{\max}) = 0,$$

so that in the absence of demography (*i.e.*, if $R = \gamma = 0$), and with an integrable initial condition $c(0, x, y)$, the global population size $\mathbf{C}(t) = \int_{\mathbb{R} \times (y_{\min}, y_{\max})} c(t, x, y) dx dy$ would remain constant.

2.2 *Modeling genetic and spatial fragmentation in dispersal and growth*

Spatial fragmentation in environmental conditions are assumed to impact the consumer growth rate R and its mobility D. Genetic and environmental effects on R and D are assumed to be additive. The parameters

$R_0 > 0$ and $D_0 > 0$ are the basal values for growth and diffusion. These basal values are modified according to a genetic effect, R_g, respectively D_g, and a environmental effect, R_s, respectively D_s. The parameter L controls the spatial fragmentation: $R_s(x/L)$ and $D_s(x/L)$ are L-periodic. A small value of L corresponds to a highly fragmented (or rapidly varying) environment, and a large value corresponds to a low fragmented (or slowly varying) environment. Moreover, we also test the effect of introducing an amplitude effect to scale the fragmentation with respect to the scenario presented (see Appendix A).

Figure 1: **Genetic and environmental effects on growth (R) and dispersal (D)**. The panel A displays the curves representing the genetic effect for the dispersal rate $D_g(y)$ (red) and growth rate $R_g(y)$ (blue) expressed as a function of phenotypic traits $y \in (y_{min}, y_{max})$. The coefficient d is the distance between the optima for dispersal and growth rate. The panel B shows the environmental effect for the dispersal ($D_s(x/L)$)(dotted line) and ($R_s(x/L)$) (solid line). The panel C shows an example of resulting total population density $C(T_{sim}, x)$ (see Equation (5)) obtained from the solution of the Equation (1) at time $T_{sim} = 60$ along with the position x, with the parameter values: $\mu = 0$, $d = 2$, $R_0 = 1$, $D_0 = 1$, $\sigma = 1$ for in the Scenario R_{het}.

{fig:OdOr}

Genetic effect. Given its trait value y the genetic effect on the growth rate R and the diffusion coefficient D is assumed to be Gaussian (Figure 1):

$$R_g(y) = \exp(-(y + d/2)^2/(2\sigma^2)), \tag{2}$$

$$D_g(y) = \exp(-(y - d/2)^2/(2\sigma^2)) \tag{3}$$

where $d > 0$ corresponds to the distance between the two optima. The optimum trait for diffusion represents the consumer optimal dispersal strategy, and the optimum trait for the growth rate represents the consumer optimal resource exploitation strategy. Here, we assume that the optimum traits are symmetric with respect to 0, $O_R = -d/2$ and $O_D = +d/2$ for the growth rate and dispersal, respectively. The coefficient σ, fixed to 1 in the following, is the standard deviation of the Gaussian function and indicates the intensity of selection around the optimal trait value (smaller σ means higher intensity of selection).

Environmental effect. The terms $R_s(x/L)$ and $D_s(x/L)$ describe the periodic variations over the space x (Figure 1B). Here, R_s is a 1-periodic piecewise constant function of mean 0, with $R_s(x) = R_0$ on $[0, 1/2)$ and $R_s(x) = -R_0$ on $[1/2, 1)$. Equivalently, D_s is a smooth 1-periodic function, with mean value 0, and bounded from below by $-D_0$ (so that D is always positive). More precisely, we define the 1-periodic function $\delta_1(x)$ such that $\delta_1(x) = D_0$ in $[0, 1/2)$ and $\delta_1(x) = -D_0$ in $[1/2, 1)$.

Then, D_s is obtained by regularizing δ_1 with a convolution by a smooth function:

$$D_s(x) = \int_{\mathbb{R}} \delta_1(x - z)\, \phi(z)\, dz,$$

with ϕ a Gaussian function with small variance.

2.3 *Simulation scenario*

We define three scenarios, depending on the presence of spatial fragmentation, and on

 its effects on growth or dispersal:

- Scenario H: spatially homogeneous coefficients. In this case,

$$R(x,y) = R_h(y) := R_0 + R_g(y) \text{ and } D(x,y) = D_h(y) := D_0 + D_g(y).$$

- Scenario R_{het}: fragmented growth and homogeneous dispersal. In this case,

$$R(x,y) = R_0 + R_g(y) + R_s(x/L) \text{ and } D(x,y) = D_h(y) = D_0 + D_g(y).$$

- Scenario D_{het}: homogeneous growth and fragmented dispersal. In this case,

$$R(x,y) = R_h(y) = R_0 + R_g(y) \text{ and } D(x,y) = D_0 + D_g(y) + D_s(x/L).$$

Under these scenarios, we numerically simulated the model of Equation (1), to ex-

 plore the phenotypic trait composition in the population with different parameter value

combinations. Specifically, we focused on: i) the period of fragmentation L, considering

 rapidly (small period $L = 2$) and slowly (large period $L = 10$) varying environments; ii)

the distance d between the two optima, where we considered a short distance for weak

 trade-off ($d = 2$) and a large distance for strong trade-off ($d = 4$); iii) and, the mutation

parameter $\mu = 0$ (no mutations) or $\mu = 0.1$. The equations are solved numerically by

 transforming them into lattice dynamical systems (continuous time, discrete space with

small space step), and using a Runge-Kutta method over a fixed spatial domain (defined

 as $x \in [0; 250]$ by step 0.5). The phenotype space is defined between $y_{min} = -5$ and

$y_{max} = 5$ by step $\delta_x = 0.1$. The implementation is performed by using the software

 Matlab$^{\circledR}$ (code repository: DOI 10.17605/OSF.IO/V6N4M).

2.4   The spreading speed

The consumer species $R - D$ trade-off is investigated by focusing on the spreading properties and analyzing the population forefront under the defined scenarios. The spreading speed (to the right) V is the asymptotic rate at which a population, initially concentrated to the left of some point, expands its spatial range. It can be defined here as the smallest speed such that, if an observer travels to the right (*i.e.*, towards increasing x values) with this speed, he will observe that the population density vanishes. In mathematical terms, V is the only speed such that:

$$\begin{aligned}
\sup_{x \geq z} C(t, x + w\,t) &\xrightarrow[t \to +\infty]{} 0 \quad \text{for all } w > V \text{ and } z \in \mathbb{R}, \\
\inf_{x \leq z} C(t, x + w\,t) &\xarrownot[t \to +\infty]{} 0 \quad \text{for all } w < V \text{ and } z \in \mathbb{R},
\end{aligned} \tag{4}$$

with $C(t, x)$ the population density at spatial position x:

$$C(t, x) = \int_{y_{\min}}^{y_{\max}} c(t, x, y)\, dy. \tag{5}$$

For each phenotype y, the spreading speed $v(y)$ of the phenotype y can be defined as well by replacing $C(t, x + w\,t)$ with $c(t, x + w\,t, y)$ in the above expressions.

The existence of a spreading speed and analytical characterizations have been obtained for standard equations with spatially homogeneous coefficients and local competition terms (Aronson and Weinberger, 1975, 1978; Fife and McLeod, 1977; Kolmogorov et al., 1937). Comparable results have been obtained with a periodically varying coefficient as in Equation (1) and a local competition term (Berestycki and Hamel, 2002, 2005), namely for equations of the form:

$$\partial_t c(t, x, y) = \partial_{xx}(D(x, y)\, c(t, x, y)) + \mu\, \partial_{yy} c(t, x, y)$$

$$+ c(t, x, y)\, (R(x, y) - \gamma\, c(t, x, y)). \tag{6}$$

Here, the difference with Equation (1) is that the individuals with phenotype y only interact with individuals with the same phenotype. As we did not assume an Allee

effect in Equation (1), the solutions should be pulled by the individuals in the leading edge of the colonization (Roques et al., 2012; Stokes, 1976). Their speed should therefore

only depend on the growth term through its linearization around 0, here $R(x, y)c(t, x, y)$. We therefore conjecture that the spreading speeds V of the solutions of the nonlocal

Equation (1) and the local equation (6) are equal. This conjecture is supported by the results of Alfaro et al. (2013), which deal with a nonlocal equation of the form (1), with a

constant diffusion term D and with a growth term of the form $R(x, y) = r(y - B x)$, with $r(y) = r_{\max} - b y^2$ (to each position x is attached an optimal phenotype $B x$). This would

imply that the fastest phenotype,

$$y^* = \operatorname*{argmax}_{y \in (y_{\min}, y_{\max})} v(y),$$

222 has the same speed for the two Equations (1) and (6) with and without nonlocal interactions. As the fastest phenotype, y^* does not compete with other phenotypes, its

224 speed should indeed not be influenced by the competition term, and therefore be the same for the two equations. This conjecture is also supported by the results of Girardin

226 (2017) (Theorems 1.6 and 1.7), who studied an analogue of (1), but with a discrete phenotype space and a spatially homogeneous environment, leading to a system of reaction-

228 diffusion equations coupled by discrete Laplace mutation term.

For Equation (6), under our three scenarios (H, R_{het}, D_{het}), more or less explicit for-

mulas for the spreading speed are available. Thus, we compare these approximations of the spreading speeds with numerical results, using approached models and limiting

cases of rapidly and slowly varying environments and we compare these approximations with numerical results.

First, when the environment is spatially homogeneous (scenario H), *i.e.*, when $R(x, y) = R_h(y)$ and $D(x, y) = D_h(y)$, and in the absence of mutations ($\mu = 0$), the spreading speed

associated with a phenotype y is $v(y) = 2\sqrt{R_h(y)\, D_h(y)}$ (Kolmogorov et al., 1937). In that case, and according to the values of d and σ (see Equations (2) and (3)), the fastest

phenotypes can be the generalist, $y^* = 0$ or the two specialists, $y^* = O_R = -d/2$ and $y^* = O_D = d/2$, see Appendix B. The overall spreading speed defined by Equation (4) is

$V = 2\sqrt{R_h(y^*)\, D_h(y^*)}$. When the environmental fragmentation only impacts the growth rate $R(x, y)$ keeping the diffusion coefficient spatially homogeneous $D(x, y) = D_h(y)$

(scenario R_{het}), a general formula for the spreading speed has been obtained by Berestycki and Hamel (2005). Their results also encompass the case of a fragmented diffusion co-

efficient $D(x, y)$ but spatially homogeneous growth rate $R(x, y) = R_h(y)$ (scenario D_{het}). However, in this case, their result holds true for equations with "Fickian" spatial diffu-

sion term, *i.e.*, $\partial_x(D(x, y)\, \partial_x c)$ instead of the Fokker-Planck diffusion $\partial_{xx}(D(x, y)\, c)$ in (6) (Roques, 2013; Turchin, 1998).

In the spatially fragmented cases (scenarios R_{het} and D_{het}) the formulas rely on variational characterizations which make them hardly tractable, even numerically (see Ap-

pendix B). More tractable formulas for the phenotype spreading speeds can be obtained for rapidly varying (*i.e.*, when the period is small, $L \to 0$) and slowly varying (*i.e.*, when

the period is large, $L \to \infty$) environments in the absence of mutations (*i.e.*, $\mu = 0$) (Hamel et al., 2010, 2011; Smaily et al., 2009). These formulas are summarized in Table

1, see also Appendix B for more mathematical details. We check the accuracy of the analytical approximations in Table 1 and we compare them with numerical simulations

for the considered scenarios.

Table 1: Theoretical phenotype spreading speeds for Equation (6) with $\mu = 0$ (no mutation) for rapidly varying environments $L \to 0$ and slowly varying environments $L \to \infty$.

	Spatial fragmentation	$L \to 0$	$L \to \infty$
Scenario H	$R_h(y), D_h(y)$	$v(y) = 2\sqrt{R_h(y)\,D_h(y)}$	$v(y) = 2\sqrt{R_h(y)\,D_h(y)}.$
Scenario R_{het}	$R(x,y), D_h(y)$	$v_{R_{het},0}(y) = 2\sqrt{R_h(y)\,D_h(y)}^*$	$v_{R_{het},\infty}(y) = 4\sqrt{D_h(y)} \times \frac{(R^+(y))^2+(R^-(y))^2+(R^+(y)+R^-(y))\sqrt{\triangle(y)}}{(R^+(y)+R^-(y)+2\sqrt{\triangle(y)})^{\frac{3}{2}}}^{**}$
Scenario D_{het} (*Fickian diffusion*)	$R_h(y), D(x,y)$	$v_{D_{het},0}(y) = 2\sqrt{R_h(y)\,\langle D_1\rangle_H(y)}^{***}$	$v_{D_{het},\infty}(y) = 2\sqrt{R_h(y)}\,\langle\sqrt{D_1}\rangle_H(y)^{***}$

Where:

* $v_{Rh,0}(y) = 2\sqrt{\overline{R}_x(y)\,D_h(y)}$ with $\overline{R}_x(y) = R_0 + R_g(y) + \int_0^1 R_s(x)\,dx = R_h(y).$

** $R^+(y) = R_h(y) + R_0$, $R^-(y) = R_h(y) - R_0$, $\triangle(y) = (R^+(y))^2 + (R^-(y))^2 - R^+(y)\,R^-(y).$

*** $\langle F\rangle_H(y) = \left(\int_0^1 \frac{dx}{F(x,y)}\right)^{-1}$ the harmonic mean of F, and $D_1(x,y) = D_0 + D_g(y) + D_s(x).$

{tab:speed}

3   Results

{Res}

The forefront profiles highlight the role of the $R - D$ trade-off, the environmental heterogeneity and the mutation in influencing the spreading speed of the different phenotypes (Figures 2-5). In Section 3.1 we analyze these figures, then, in Section 3.2 we use the theoretical formulations to better understand the numerical simulations.

Hereafter, we identify the strategies favoring the selection of phenotypes with a behavior that increases dispersal capacities as D-strategy (namely, when $y^* > 0$), with a generalist behavior as G-strategy (namely, when $y^* = 0$) and with a behavior that increases growth rate as R-strategy (namely, when $y^* < 0$). As at the back of the front the R-strategy is always selected, our focus is entirely dedicated to the trade-off among R and D on the forefront, see Appendix C.

3.1 The $R − D$ trade-off selecting the fastest strategy on the forefront

In the absence of mutation ($\mu = 0$), when $d = 2$, the forefront is composed mostly by one phenotype with a low degree of specialization on R and D (Figure 2A, C and E; Figure 4A, C and E). For scenarios H and R_{het}, when fragmentation is high, numerical simulations show a small shift towards the R-strategy (Figure 2A and C, Table 2). Instead, when there is a slowly varying environment, the shift occurs towards D-strategy (Figure 2E, Table 2). Under scenario D_{het}, whatever environment fragmentation, there is a clear shift towards the D-strategy (Figure 4 C and E, Table 2).

When $d = 4$, the colonization is mostly driven by the D-strategy (Figures 2 and 4 B, D and F). A less fragmented habitat ($L = 10$), under the scenario R_{het}, increases the advantage of the D-specialist on the forefront, shifting the trade-off in favour of the D-strategy (see Figure 2 D *vs.* F). For the scenario D_{het}, the difference between weak and strong trade-off is even more remarkable as the advantage is completely shifted in favor of the strategy $y* = O_D$ (see Figure 4D), defining also a different forefront profile.

These outcomes are completely blurred when introducing a positive value for the mutation coefficient. In fact, the presence of mutations leads to a homogenization of the phenotypic distribution and therefore to a wider phenotype ensemble that leads the forefront: all of the phenotypes should theoretically spread with the same asymptotic speed (see Girardin, 2017, in a homogeneous case with discrete phenotype space). Yet, the population densities (Figures 3 and 5) indicate that the R-strategy becomes the preferred one almost in all cases when $d < d_{cr}$, except for the scenario D_{het} with a slowly varying environment (Figure 5 E). The D-specialist is still the fastest phenotype under the scenario D_{het} in case of strong trade-off ($d = 4$) (Figure 5D and F). However, we notice

that the shape of the solution at the leading part of the expansion is quite unusual. In all
cases, we observe a "bump" corresponding to a fraction of the population which adopts
the D-strategy. Thus, the expansion may take advantage of the larger diffusion coefficient
of the *D*-specialists and of the larger growth rate of the *R*-specialist by allowing more
polymorphism at the leading edge of the propagation.

Figure 2: **Population density contrasting scenario R_{het} (Panels C, D, E, F) with scenario H (Panels A and B) without mutation.** The forefront profiles show the population density with respect to the phenotypes and the space variables. Over lines the trade-off strength is showed: weak trade-off ($d = 2$) (Panel A, C, E) and strong trade-off ($d = 4$) (Panel B, D, F). Over columns the effect of the period L is showed: rapidly varying environment ($L = 2$) (Panel C and D) and slowly varying environment (Panel E and F). Solutions are obtained by numerically simulating the Equation (1) and results are reported at time $T_{sim} = 60$. White dashed lines highlight the optimum values (*i.e,* $O_D = d/2$ and $O_R = -d/2$).

{fig:FrontRh}

Figure 3: **Population density contrasting scenario R_{het} with scenario H with mutation.** Caption description is the same of Figure 2, but considering a positive value for mutation $(\mu > 0)$.

{fig:FrontRhmu}

Figure 4: **Population density contrasting scenario D_{het} with scenario H without mutation.** Caption description is the same of Figure 2, but for scenario D_{het}.

{fig:FrontDh}

Figure 5: **Population density contrasting scenario D_{het} with scenario H with mutation.** Caption description is the same of Figure 2, but for scenario D_{het} with a positive value for mutation $\mu > 0$.

{fig:FrontDhmu}

3.2 *Insight from the theoretical speeds*

{sec:theor_spee

In this section, we compare the numerical simulations of Equation (1) presented in Figures 2-5 with the analytical formulations presented in Figure 6 and Table 1. We first check if the outcomes that can be obtained from the theoretical speeds match with the numerical results. These outcomes are summarised in Table 2. Second, when there is a good match, we use the explicit formulas to explain the observed trends. We recall that the theoretical speeds of Table 1 were derived with the simpler local model (6) with $\mu = 0$ and therefore do not take the nonlocal competition and mutation effects into account.

Figure 6: **Theoretical phenotype spreading speeds.** The theoretical phenotype spreading speeds presented in Table 1 are showed in function of phenotypes $y \in [-5, 5]$ considering a weak trade-off ($d = 2$) (Panel A) and a strong trade-off ($d = 4$) (Panel B). Different colors refers to the formulations of the spreading speed highlighted in Table 1, the dots represent the fastest phenotype leading the forefront. Dashed lines highlight the positions of the optimum traits O_R and O_D.

{fig:v_th}

In absence of mutation ($\mu = 0$), there is a critical threshold d_{cr} on the distance d between the optima ($d_{cr} \approx 2.4$ with our parameter values, see Appendix B), such that the function $v(y) = 2\sqrt{R_h(y)\,D_h(y)}$, corresponding to the spreading speed, either admits one ($d < d_{cr}$) or two ($d > d_{cr}$) maxima. When $d < d_{cr}$ (*i.e*, $d = 2$) and with a rapidly varying environment, the generalist behavior is expected to be selected as the fastest trait following the theoretical formulations under scenarios H and R_{het}. However, numerical simulations show a small shift towards the R-strategy (Table 2). Instead, when there is a

slowly varying environment, or when heterogeneity impacts D, theoretical formulations consistently predict a shift towards the D-strategy (Table 2, Figure 6).

In the scenario D_{het}, the theoretical speed $v_{D_{het},0}(y) = 2\sqrt{R_h(y)\,\langle D_1\rangle_H(y)}$ in rapidly varying environments (small period $L = 2$ in the numerical results, $L \to 0$ in Table 1) involves the harmonic mean of the diffusion term. Contrarily to the arithmetic mean, the harmonic mean gives a higher weight to small values. Thus, small values of $D(x, y)$, even on a very small spatial interval, should lead to small speeds $v_{D_{het},0}(y)$, based on the results of Table 1. This leads to an imbalance in favor of the D-strategies (compare $v_{D_{het},0}(y)$ and $v(y)$ in Figure 6; Figure 4A and B *vs.* Figure 4C and D), which avoid very small values of $D(x, y)$. In the case of slowly varying environments (large period $L = 10$ in the numerical results, $L \to +\infty$ in Table 1), the theoretical speed $v_{D_{het},\infty}(y) = 2\sqrt{R_h(y)}\,\langle\sqrt{D_1}\rangle_H(y)$ again involves an harmonic mean of the diffusion term (here, its square root) which explains the advantage of the D-strategy, as in the case of rapidly varying environments.

When heterogeneity is introduced on R (scenario R_{het}), in rapidly varying environments, the theoretical formulas predict that the strategy remains unchanged (G when $d < d_{cr}$ or R+D when $d > d_{cr}$) compared to the homogeneous scenario (H). The theoretical spreading speed of each phenotype in the scenario R_{het}, $v_{R_{het},0}(y) = 2\sqrt{R_h(y)\,D_h(y)}$, is indeed the same as the speed $v(y)$ obtained in the homogeneous scenario H (both curves are superimposed in Figure 6): as the spatial arithmetic mean of the growth rate (noted $\overline{R}_x(y)$ in the legend of Table 1) is precisely equal to R_h, the homogenization results of Smaily et al. (2009) imply that the two speeds are equal. In the numerical simulations (see Table 2, lines A for numerical and C for theoretical), there are some discrepancies, as with a rapidly varying environment ($L = 2$), the homogenization limit is not reached. Hence, we observe a slight shift of the G-strategy towards the R-strategy when $d < d_{cr}$.

When $d > d_{cr}$, the two strategies are present on the forefront, with a slight advantage for the D-strategy (Figure 2A and B *vs.* Figure 2C and D). In slowly varying environments, the theoretical formulas and the numerical simulations consistently predict a shift towards the D-strategy. In this last case, the formula for $v_{R_{het},\infty}(y)$ can be written in the form $2\sqrt{F(y)\,D_h(y)}$, for some function F which satisfies $F(O_D) \approx (32/27)\,R_0$ (to be compared with $R_h(O_D) \approx R_0$) and $F(O_R) \sim R_0 + 1$ for small R_0 (to be compared with $R_h(O_R) = R_0 + 1$). Thus, compared to the homogeneous case $(v(y) = 2\sqrt{R_h(y)D_h(y)})$, the heterogeneity on R creates an asymmetry in favor of the D-strategy which manages to keep a growth rate larger than R_0.

When $\mu > 0$ (see Table 2 line B), as expected, there are some discrepancies between the numerical results and the theoretical predictions. However, the arguments above may explain some of the observations. First, in the scenario D_{het} in most cases (except when $d < d_{cr}$ in rapidly varying environments) we again observe a shift towards D-strategies which is most probably due to the "harmonic mean" effect described above. We note that this shift is stronger in slowly varying environments. In the scenario R_{het}, although the R-strategy is always selected when $\mu > 0$, the positive effect of the heterogeneity on the maintenance of the D-strategy which we noted above in the theoretical formulas is still visible on Figure 3F, which shows more polymorphism compared to the scenario H.

We note the positive effect of increasing the period L (equivalently, reducing the environmental fragmentation) on the spreading speeds: this effect, which is obvious in Figure 6 can also be observed in Figures 2-5.

		H	R_{het}		D_{het}	
			Rapidly varying	Slowly varying	Rapidly varying	Slowly varying
$d < d_{cr}$	A	-0.1	-0.1	0.1	0.5	0.3
	B	-0.7	-0.7	-0.7	-0.7	0.1
	C	0	0	0.2	0.6	0.5
$d > d_{cr}$	A	R+D	0.95	1	0.95	0.95
	B	-1.2	-1.3	-1.35	0.5	0.8
	C	R+D	R+D	1	1	1

Table 2: **Fastest phenotypes leading the forefront**. The table reports the values of the fastest phenotype leading the forefront with respect to the optimum value ($2y^*/d$). Results of numerical simulations of the Equation 1 are compared over the lines considering the presence and absence of mutation and the theoretical formulas: A) $y^* = \mathrm{argmax}(v_{sim}(y))$, with $\mu = 0$; B) $y^* = \mathrm{argmax}(v_{sim}(y))$, with $\mu > 0$; C) $y^* = \mathrm{argmax}(v_{th}(y))$, corresponding to the spreading speed v_{th} reported in Table 1. The corresponding strategy is highlighted: Blue cells correspond to R-strategies ($2y^*/d < 0$) and red cells to D-strategies ($2y^*/d > 0$). White cells correspond to generalists ($2y^*/d = 0$) or when both strategies lead to the same speed (noted R+D). $d < d_{cr}$ correspond to a weak trade-off ($d = 2$), $d > d_{cr}$ correspond to a strong trade-off ($d = 4$). Rapidly varying correspond to a period of heterogeneity $L = 2$ and slowly varying correspond to a period of heterogeneity $L = 10$.

{table:strategie

4 Discussion

Dispersing faster or growing stronger? In this work, we studied which of these strategies is selected in populations invading a heterogeneous environment. We gathered analytical solutions from the literature and performed numerical simulations of a reaction-diffusion model describing the demo-genetic dynamics of a population invading a one-dimensional environment. Results show that the symmetrical effects of growth and dispersal on the spreading speed is broken in the presence of competition between phenotypes, shrinking the population density around the optimum values. From here we observe that, at the back of the forefront, the dynamics is almost always carried out by the R-specialists, while, on the forefront, the selection of the fastest strategy is less obvious.

In this study, we identify the main following results: i) R-strategies are favored in spatially homogeneous environments, but the introduction of heterogeneity leads to a shift towards D-strategies, with at least more polymorphism at the forefront; ii) due to a "harmonic mean effect" that we have highlighted through analytical expressions obtained with a simpler model, this phenomenon is even stronger when spatial heterogeneity affects the diffusion term. In this case, the introduction of spatial heterogeneity can lead to a complete switch from an R-strategy to a D-strategy; iii) the spatial fragmentation does not affect a lot the $R - D$ trade-off, but tends to modulate the polymorphism: in situations where only R-strategists are present at the forefront when the level of fragmentation is high (small L), both R-strategists and D-strategists tend to be present at the forefront in low fragmented environments made of large patches (large L); iv) mutations produce an advantage towards the R-strategy, and homogenize the phenotype distribution, also

leading to more polymorphism on the forefront; v) these effects can be observed with a

weak trade-off (such that the generalist $y* = 0$ leads the population in a homogeneous

model without interactions), but become even stronger with a strong trade-off (such that

$R-$strategists and $D-$strategists have the same speed in a homogeneous model with-

out interactions) and vi) the comparison among theoretical and numerical simulations

allows checking when formulas obtained with a simpler model lead to results which are

consistent with of a more complex one.

Some of these results (points i,ii,iii) are in accordance with the ones of Burton et al.

(2010), who used an individual-based spatial model to study the evolution of three traits

in a population undergoing range expansion. When resources are highly fragmented, the

trade-off favors to the selection of an R-strategy on the forefront as high resource avail-

ability and fecundity facilitate expansion by increasing population growth. By contrast,

in a low fragmented environment, the faster dispersers take advantage of their mobility

to reach the most favorable habitats and lead the forefront. Evolution thus leads to the

selection of a greater capacity for dispersion. Conversely, when heterogeneity impacts

dispersal, only the D-specialists confer the maximal speed and persist on the forefront,

whatever the level of spatial fragmentation. Recently, given two species having growth

and dispersal coefficients R_1, D_1 and R_2, D_2 (for species 1 and 2 respectively), Deforet

et al. (2019) found that the evolutionary outcome mainly depends on the simple condi-

tion $v_1 = 2\sqrt{R_1 D_1} > v_2 = 2\sqrt{R_2 D_2}$, with success of the fastest (here, species 1). In our

work, the use of the theoretical formulas of Table 1 mainly relies on the same assumption,

that the fastest trait drives the expansion. In most cases, our results show that the obser-

vations of Deforet et al. (2019) remain true in a more general context with a continuum of

traits and possibly mutations between traits. However, we also observed some discrepan-

cies in the presence of mutations, which tend to advantage the R-strategy. Additionally, as recently observed by Keenan and Cornell (2021) in a homogeneous environment, in the presence of mutations, the furthest forward phenotypes are not necessarily those associated with the largest value of the product RD (see below).

Duputié and Massol (2013) argues that natural selection tends to favor dispersal to face spatio-temporal variation in local conditions. Consequently, more dispersive phenotypes are expected to predominate in unstable habitats, while less dispersive phenotypes are common in stable habitats and populations. Here, we found that in the bulk of the population, which corresponds to a saturated population, the R-strategy is always preferred, which is not always the case at the forefront. By definition, the expanding part of the population encounters a more variable environment, especially when the environment is itself highly heterogeneous. In such cases, we observed a shift towards the D-strategy. These findings are also consistent with the "Spatial sorting theory" which predicts that, at the forefront, dispersal may be strongly favored because of the accumulation of the best dispersers (Phillips et al., 2008; Shine et al., 2011; Travis and Dytham, 2002).

The presence of mutations homogenizes the spreading speed between morphs, even in presence of nonlocal competition. We also establish that polymorphism, caused by mutation, is maintained in the presence of spatial fragmentation impacting the R and D coefficients. Another possible effect of mutation is an increased spreading speed. Taking again a system with only two morphs (as in Deforet et al. (2019), but with a mutation term), typically an R-specialist and a D-specialist, Elliott and Cornell (2012) and Morris et al. (2019) investigated the effect of varying R and D on the spreading speed. They found that the system would spread faster in the presence of both phenotypes than

just one phenotype would spread in the absence of mutation for certain combination of R and D values. In a similar way, using the results of Girardin (2017), Keenan and Cornell (2021) considered the $R - D$ trades-off in the case of N phenotypes in a homogeneous environment, and obtained some conditions on the curvature of the trade-off curve $(D, R(D))$ such that this "anomalous" faster speed emerges. Here, although the trade-off curve $(D, R(D))$ has positive curvature, we did not observe this phenomenon: in all of our simulations of Figures 3-5, the speed is reduced when $\mu > 0$, compared to the speed of the fastest trait when $\mu = 0$. The theoretical results of Keenan and Cornell (2021) require a vanishing small mutation rate, and their numerical results use a mutation coefficient 10^{-6} (to be compared with $\mu/(\delta_x)^2 = (0.1)/(0.1)^2 = 10$ in our continuous framework), which may explain these differences.

Future works could consider a more detailed analysis of the lineages that pull the forefront, to determine for instance if the D-specialists in Figures 3 and 5 are produced by mutation from R-specialists or correspond to a self-sustaining fraction of the population. In that respect, one could reconstruct the genealogies of the fractions composing the population using the methods in (Roques et al., 2012). We recall that our results depend on the assumptions about the form of the dispersal and growth rate functions and the fragmentation definition. For instance, we do not take an Allee effect into account. It is demonstrated to have important consequences on the invasions dynamics and especially on the lineages that compose the forefront (Andrade-Restrepo et al., 2019; Chuang and Peterson, 2016; Roques et al., 2012).

Literature Cited

Alfaro, M., H. Berestycki, and G. Raoul (2017). The effect of climate shift on a species submitted to dispersion, evolution, growth, and nonlocal competition. SIAM Journal on Mathematical Analysis 49(1), 562–596.

Alfaro, M., J. Coville, and G. Raoul (2013). Travelling waves in a nonlocal reaction-diffusion equation as a model for a population structured by a space variable and a phenotypic trait. Communications in Partial Differential Equations 38(12), 2126–2154.

Alfaro, M., L. Girardin, F. Hamel, and L. Roques (2021). When the Allee threshold is an evolutionary trait: persistence vs. extinction. Journal de Mathématiques Pures et Appliquées 155, 155–191.

Alfaro, M. and G. Peltier (2021). Populations facing a nonlinear environmental gradient: steady states and pulsating fronts. arXiv preprint arXiv:2101.08078 32(02), 209–290.

Anciaux, Y., A. Lambert, O. Ronce, L. Roques, and G. Martin (2019). Population persistence under high mutation rate: from evolutionary rescue to lethal mutagenesis. Evolution 73(8), 1517–1532.

Andrade-Restrepo, M., N. Champagnat, and R. Ferrière (2019). Local adaptation, dispersal evolution, and the spatial eco-evolutionary dynamics of invasion. Ecology letters 22(5), 767–777.

Aronson, D. G. and H. F. Weinberger (1975). Nonlinear diffusion in population genetics, combustion, and nerve pulse propagation. In J. A. Goldstein (Ed.), Partial Differential

Equations and Related Topics, Berlin, Heidelberg, pp. 5–49. Springer Berlin Heidelberg.

Aronson, D. G. and H. G. Weinberger (1978). Multidimensional nonlinear diffusion arising in population genetics. Adv Math 30(1), 33–76.

Baguette, M. and N. Schtickzelle (2006). Negative relationship between dispersal distance and demography in butterfly metapopulations. Ecology 87(3), 648–654.

Benichou, O., V. Calvez, N. Meunier, and R. Voituriez (2012). Front acceleration by dynamic selection in fisher population waves. Physical Review E 86(4), 041908.

Berestycki, H. and F. Hamel (2002). Front propagation in periodic excitable media. Comm Pure Appl Math 55(8), 949–1032.

Berestycki, H. and F. Hamel (2005). Gradient estimates for elliptic regularizations of semilinear parabolic and degenerate elliptic equations. Communications in Partial Differential Equations 30(1-3), 139–156.

Berestycki, H., F. Hamel, and L. Roques (2005). Analysis of the periodically fragmented environment model: II - Biological invasions and pulsating travelling fronts. J Math Pures Appl 84(8), 1101–1146.

Berestycki, N., C. Mouhot, and G. Raoul (2015). Existence of self-accelerating fronts for a non-local reaction-diffusion equations. arXiv preprint arXiv:1512.00903.

Bonte, D. and Q. Bafort (2019). The importance and adaptive value of life-history evolution for metapopulation dynamics. Journal of Animal Ecology 88(1), 24–34.

Bonte, D. and E. de la Pena (2009). Evolution of body condition-dependent dispersal in metapopulations. Journal of Evolutionary Biology 22(6), 1242–1251.

Bouin, E. and V. Calvez (2014). Travelling waves for the cane toads equation with bounded traits. Nonlinearity 27(9), 2233.

Bouin, E., V. Calvez, N. Meunier, S. Mirrahimi, B. Perthame, G. Raoul, and R. Voituriez (2012). Invasion fronts with variable motility: phenotype selection, spatial sorting and wave acceleration. Comptes Rendus Mathematique 350(15-16), 761–766.

Bouin, E., M. H. Chan, C. Henderson, and P. S. Kim (2018). Influence of a mortality trade-off on the spreading rate of cane toads fronts. Communications in Partial Differential Equations 43(11), 1627–1671.

Burton, O. J., B. L. Phillips, and J. M. Travis (2010). Trade-offs and the evolution of life-histories during range expansion. Ecology letters 13(10), 1210–1220.

Chuang, A. and C. R. Peterson (2016). Expanding population edges: theories, traits, and trade-offs. Global change biology 22(2), 494–512.

Deforet, M., C. Carmona-Fontaine, K. S. Korolev, and J. B. Xavier (2019). Evolution at the edge of expanding populations. The American Naturalist 194(3), 291–305.

Denno, R. F. (1994). Life history variation in planthoppers. In Planthoppers, pp. 163–215. Springer.

Duputié, A. and F. Massol (2013). An empiricist's guide to theoretical predictions on the evolution of dispersal. Interface focus 3(6), 20130028.

Duthie, A. B., K. C. Abbott, and J. D. Nason (2015). Trade-offs and coexistence in fluctuating environments: evidence for a key dispersal-fecundity trade-off in five nonpollinating fig wasps. The American Naturalist 186(1), 151–158.

Elliott, E. C. and S. J. Cornell (2012). Dispersal polymorphism and the speed of biological invasions. PloS one 7(7), e40496.

Fife, P. C. and J. McLeod (1977). The approach of solutions of nonlinear diffusion equations to traveling front solutions. Arch Ration Mech Anal 65(1), 335–361.

Fisher, R. A. (1937). The wave of advance of advantageous genes. Ann Eugen 7, 335–369.

Girardin, L. (2017). Non-cooperative Fisher–KPP systems: traveling waves and long-time behavior. Nonlinearity 31(1), 108.

Griette, Q., G. Raoul, and S. Gandon (2015). Virulence evolution at the front line of spreading epidemics. Evolution 69(11), 2810–2819.

Hamel, F., J. Fayard, and L. Roques (2010). Spreading speeds in slowly oscillating environments. Bull Math Biol 72(5), 1166–1191.

Hamel, F., F. Lavigne, G. Martin, and L. Roques (2020). Dynamics of adaptation in an anisotropic phenotype-fitness landscape. Nonlinear Analysis: Real World Applications 54, 103107.

Hamel, F., G. Nadin, and L. Roques (2011). A viscosity solution method for the spreading speed formula in slowly varying media. Indiana Univ Math J 60, 1229–1247.

Hanski, I., M. Saastamoinen, and O. Ovaskainen (2006). Dispersal-related life-history trade-offs in a butterfly metapopulation. Journal of animal Ecology 75(1), 91–100.

Helms, J. and M. Kaspari (2015). Reproduction-dispersal tradeoffs in ant queens. Insectes sociaux 62(2), 171–181.

Hirt, M. R., T. Lauermann, U. Brose, L. P. Noldus, and A. I. Dell (2017). The little things that run: a general scaling of invertebrate exploratory speed with body mass. Ecology 98(11), 2751–2757.

Keenan, V. A. and S. J. Cornell (2021). Anomalous invasion dynamics due to dispersal polymorphism and dispersal–reproduction trade-offs. Proceedings of the Royal Society B 288(1942), 20202825.

Kelehear, C. and R. Shine (2020). Tradeoffs between dispersal and reproduction at an invasion front of cane toads in tropical australia. Scientific Reports 10(1), 1–7.

Kolmogorov, A. N., I. G. Petrovsky, and N. S. Piskunov (1937). Étude de l'équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique. Bull Univ État Moscou, Sér. Int. A 1, 1–26.

Legrand, D., J. Cote, E. A. Fronhofer, R. D. Holt, O. Ronce, N. Schtickzelle, J. M. Travis, and J. Clobert (2017). Eco-evolutionary dynamics in fragmented landscapes. Ecography 40(1), 9–25.

Morris, A., L. Börger, and E. Crooks (2019). Individual variability in dispersal and invasion speed. Mathematics 7(9), 795.

Peltier, G. (2020). Accelerating invasions along an environmental gradient. Journal of Differential Equations 268(7), 3299–3331.

Perkins, T., B. L. Phillips, M. L. Baskett, and A. Hastings (2013). Evolution of dispersal

and life history interact to drive accelerating spread of an invasive species. Ecology letters 16(8), 1079–1087.

Phillips, B. L., G. P. Brown, J. M. Travis, and R. Shine (2008). Reid's paradox revisited: the evolution of dispersal kernels during range expansion. the american naturalist 172(S1), S34–S48.

Ronce, O. and M. Kirkpatrick (2001). When sources become sinks: migrational meltdown in heterogeneous habitats. Evolution 55(8), 1520–1531.

Roques, L. (2013). Modèles de réaction-diffusion pour l'écologie spatiale. Editions Quae.

Roques, L., J. Garnier, F. Hamel, and E. K. Klein (2012). Allee effect promotes diversity in traveling waves of colonization. Proc Natl Acad Sci USA 109(23), 8828–8833.

Shigesada, N. and K. Kawasaki (1997). Biological Invasions: Theory and Practice. Oxford Series in Ecology and Evolution, Oxford: Oxford University Press.

Shine, R., G. P. Brown, and B. L. Phillips (2011). An evolutionary process that assembles phenotypes through space rather than through time. Proceedings of the National Academy of Sciences 108(14), 5708–5711.

Skellam, J. G. (1951). Random dispersal in theoretical populations. Biometrika 38, 196–218.

Smaily, M. E., F. Hamel, and L. Roques (2009). Homogenization and influence of fragmentation in a biological invasion model. arXiv preprint arXiv:0907.4951.

Steenman, A., A. Lehmann, and G. Lehmann (2015). Life-history trade-off between

macroptery and reproduction in the wing-dimorphic pygmy grasshopper Tetrix subu-lata (Orthoptera Tetrigidae). Ethology Ecology & Evolution 27(1), 93–100.

Stokes, A. N. (1976). On two types of moving front in quasilinear diffusion. Math Biosci 31, 307–315.

Szűcs, M., E. Vercken, E. V. Bitume, and R. A. Hufbauer (2019). The implications of rapid eco-evolutionary processes for biological control-a review. Entomologia Experimentalis et Applicata 167(7), 598–615.

Travis, J. M. and C. Dytham (2002). Dispersal evolution during invasions. Evolutionary Ecology Research 4(8), 1119–1129.

Tsimring, L. S., H. Levine, and D. A. Kessler (1996). RNA virus evolution via a fitness-space model. Physical review letters 76(23), 4440–4443.

Turchin, P. (1998). Quantitative Analysis of Movement: Measuring and Modeling Population Redistribution in Animals and Plants. Sinauer, Sunderland, MA.

Weinberger, H. F. (2002). On spreading speeds and traveling waves for growth and migration in periodic habitat. J Math Biol 45, 511–548.

Zera, A. J. and R. F. Denno (1997). Physiology and ecology of dispersal polymorphism in insects. Annual review of entomology 42(1), 207–230.

35