1

COP I and II dependent trafficking controls ER-associated degradation

Navit Ogen-Shtern[1,2] †‡, Chieh Chang[1,2] ‡, Haddas Saad[1,2] ‡, Niv Mazkereth[1,2], Chaitanya Patel[1,2], Marina Shenkman[1,2] and Gerardo Z. Lederkremer[1,2] *

[1]The Shmunis School of Biomedicine and Cancer Research, Cell Biology Division, George Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv 69978, Israel

[2]Sagol School of Neuroscience, Tel Aviv University, Tel Aviv 69978, Israel

†Present address: The Skin Research Institute, The Dead-Sea and Arava Science Centre, Masada 86910, Israel

‡Contributed equally

* corresponding author: Gerardo Z. Lederkremer

Email: gerardol@tauex.tau.ac.il

Author Contributions: Conceptualization: GZL. Investigation: NO-S, CC, HS, NM, CP, MS. Supervision: GZL. Writing: NO-S, GZL. Review and editing: GZL, NO-S, CC, HS, CP and MS.

Competing Interest Statement: The authors declare that they have no competing interests.

Keywords: endoplasmic reticulum/ ERAD/ vesicular trafficking/ retrotranslocation.

22
23
24

Abstract

Misfolded proteins and components of the endoplasmic reticulum (ER) quality control and ER associated degradation (ERAD) machineries concentrate in the pericentriolar ER-derived quality control compartment (ERQC), suggesting it as a staging ground for ERAD. By tracking the chaperone calreticulin and an established ERAD substrate, asialoglycoprotein receptor H2a, we have now determined that the trafficking to the ERQC is reversible and recycling back to the ER is slower than the movement in the ER periphery. The dynamics suggest vesicular trafficking rather than diffusion. Indeed, using dominant negative mutants of ARF1 and Sar1 or the drugs Brefeldin A and H89, we observed that COPI inhibition causes accumulation in the ERQC and increased retrotranslocation and ERAD, whereas COPII inhibition has the opposite effect. Our results suggest that targeting of misfolded proteins to ERAD involves COPII-dependent transport to the ERQC and that they can be retrieved to the peripheral ER in a COPI-dependent manner.

Introduction

During secretory protein biogenesis, after translocation into the ER, the quality control machinery facilitates protein folding and promotes segregation of misfolded proteins destined for ERAD. We had previously determined that misfolded proteins destined to ERAD accumulate in a specialized pericentriolar compartment, the ER-derived quality control compartment (ERQC), a staging ground for retrotranslocation and targeting to ERAD (1-5). The ERAD-associated luminal lectin OS-9 always appears localized mainly at the ERQC, whereas other quality control and ERAD components are recruited upon misfolded protein accumulation (ER stress). This compartmentalization depends on the PERK pathway of the unfolded protein response (UPR) and on the upregulation of Herp (4). PERK is activated by the initial accumulation of misfolded proteins and consequently upregulates Herp, resulting in a dynamic protein recruitment to the ERQC. Herp recruits the membrane-bound ubiquitin ligase HRD1 and additional quality control and ERAD machinery components, such as the luminal calreticulin (CRT), and the membrane proteins calnexin (CNX), Derlin-1, EDEM1, ERManI and cytosolic components, including the ubiquitin ligase SCFFbs2 and the AAA ATPase p97 (2, 6, 7). However, other machinery proteins such as BiP, PDI, UGGT and ERp57, do not accumulate in the ERQC(2-5, 8, 9). Here we show that compartmentalization of ERAD substrates in the ERQC, and subsequent targeting to ERAD, depend on a mechanism that is consistent with vesicular trafficking, involving coat protein complex II (COPII), and that recycling from the ERQC requires COPI. COPI and COPII coats had first been identified as complexes that take part in vesicular trafficking between the ER and the Golgi (10-12). Yet, COPII was recently found to participate in other processes taking place from the ER as well, including the formation of ER whorls in mammalian cells (13) and an autophagy process targeting the ER (ER-phagy) in yeast (14, 15).

Results

Dynamics of concentration of ER quality control and ERAD components at the ERQC upon proteasomal inhibition

We had previously shown that upon proteasomal inhibition, or under other conditions in which ERAD substrates accumulate (ER stress), several ER quality control and ERAD factors concentrate at the ERQC. This can be seen, for example, by comparing the localization of an established ERAD substrate, the uncleaved precursor of asialoglycoprotein receptor H2a, linked to monomeric red fluorescent protein (H2a-RFP), which serves as a marker of the ERQC (3, 4), with that of the soluble luminal chaperones BiP and CRT linked to GFP in NIH 3T3 cells. Whereas CRT-GFP was recruited to the ERQC under proteasomal inhibition with lactacystin (Lac), BiP-GFP was not (Fig. 1A, B), as we had previously seen (16). A proteasomal subunit linked to YFP, PSMD14-YFP, was also

partially recruited to the ERQC region upon proteasomal inhibition (Fig. 1C), similarly to what we
had seen with other cytosolic ERAD factors, p97 and SCFFbs2 (3, 7). Using another approach to
analyze the compartmentalization, iodixanol gradients optimized to separate the different ER-like
densities (4, 6), we observed H2a-RFP peaking in middle fractions (5-6), with a small shift to heavier
fractions, peaking in fraction 7, upon proteasomal inhibition (Fig. 1D). We had previously
characterized this shift as occurring upon protein concentration in the ERQC (4, 6). CRT-GFP had
a similar behavior and also endogenous CNX. In contrast, endogenous BiP appeared in the light
fractions and its migration did not change upon proteasomal inhibition. Most of the endogenous
proteasomes, likely ER membrane-bound, detected with antibodies against the 19S and 20S
proteasomal subunits also appeared in the middle fractions, although peaking at slightly lighter
densities in cells treated with the proteasome inhibitor bortezomib (Bz) (Fig. 1E).
To analyze the dynamics of protein concentration in the ERQC, we performed a live cell time-lapse
experiment with cells expressing H2a-RFP and CRT-GFP. In this case, we used CHO cells, which
are less mobile, to facilitate measurements for several hours. A stable cell line expressing CRT-
GFP was established, which was transfected with H2a-RFP and treated with Lac for 3 h, during
which we conducted live cell imaging (Fig. 2A). Both H2a-RFP and CRT-GFP concentrated in the
ERQC during this period, although a larger portion of CRT-GFP remained in the peripheral ER
compared to H2a-RFP (Fig. 2A, B). H2a-RFP concentrated in the ERQC at a rate of 13% of the
total cellular protein per minute, whereas the rate for CRT-GFP was 4.1% per minute, a 3.25-fold
slower rate (Fig. 2B).
The interaction of CRT with its glycoprotein substrates is dependent on a functional lectin domain.
We established a stable cell line expressing CRT(Y108F)-GFP, which carries a point mutation in
its lectin domain (17). A time lapse experiment similar to the one described above using WT CRT-
GFP was performed with this mutant (mutCRT-GFP). While H2a-RFP concentrated in the ERQC
of these cells as before, although at a slower rate (6.6% per min), mutCRT-GFP did not concentrate
to any significant extent within the 3h time frame (1.1% per min) (Fig. 2C, D).
We compared the colocalization by calculating the Pearson's coefficient. For WT CRT-GFP the
colocalization with H2a-RFP increased with time, whether only the ERQC region or the entire cell
were measured (light or dark blue trendlines respectively in Fig. 2E). The colocalization measured
for the entire cell area increased at a higher rate and to a higher extent than that measured for the
ERQC, suggesting that colocalization of CRT-GFP and H2a-RFP already occurs in peripheral ER
sites, before transport to the ERQC. In contrast, there was no colocalization and no increase with
time for mutCRT-GFP (Fig. 2E, orange and red trendlines) suggesting that the binding to substrates
through its lectin domain is necessary for the accumulation of CRT in the ERQC. These results
were not unique to the CHO cell lines, they were similar in NIH 3T3 cells, where after 3h of Lac
treatment mutCRT-GFP concentrated in the ERQC to a much lower extent than WT CRT-GFP (Fig.
2F, G).
115
Recycling from the ERQC to the peripheral ER
To study the reversibility of the process of protein concentration in the ERQC, we treated cells
expressing H2a-RFP and CRT-GFP with the reversible proteasome inhibitor MG-132 (as opposed
to the covalent inhibitor Lac) for 3h, after which the inhibitor was removed by changing the medium
(washout), and the cells were followed for up to another 2.5h. Both H2a-RFP and CRT-GFP
accumulated in the ERQC after treatment, and the ERQC accumulation decreased significantly
after washout (Fig. 3A, B). For H2a-RFP, this decrease could be due to recycling to the peripheral
ER or alternatively to its proteasomal degradation. However, in the case of CRT-GFP, as it is
neither an ERAD substrate nor a short-lived protein, the reduction in its concentration at the ERQC
must be ascribed to its recycling back to the peripheral ER.
To better understand the dynamics of recycling from the ERQC to the peripheral ER we performed
a fluorescence loss in photobleaching experiment (FLIP) in live NIH 3T3 cells expressing H2a-RFP
and CRT-GFP under continuous proteasomal inhibition. Cells were treated with Lac for 3h for
protein accumulation in the ERQC, after which both fluorescent proteins were repetitively bleached
in a peripheral region of the ER. H2a-RFP fluorescence intensity in the ERQC declined throughout
the time-lapse (Fig. 3C and Suppl. movie S1), though more slowly than in the ER periphery (Fig.

3D). This indicates that H2a-RFP cycles back to the ER periphery, but the nature of the trafficking
from the ERQC to the peripheral ER is slower than within the ER, suggesting that it is a vectorial
movement, likely vesicular trafficking rather than diffusion. In addition, transient increases in
fluorescence intensity in the ER periphery suggest uneven trafficking events, such as that occurring
at around 200 sec, with a concomitant decrease in the ERQC region (Fig. 3D). The fluorescence
intensity of CRT-GFP in the ERQC also declined more slowly than in the ER periphery (Fig. 3E).
Both CRT-GFP and H2a-RFP cycle back from the ERQC to the ER periphery in a non-diffusive
manner. To compare the trafficking dynamics of H2a-RFP and CRT-GFP we calculated the ratio
between the curve slopes (fluorescence loss rate) for each protein in the ER periphery and in the
ERQC (Fig. 3D, E). This ratio was more than 2-fold higher for H2a-RFP than for CRT-GFP (Fig.
3F), indicating that the movement out of the ERQC compared to its movement within the ER is
much slower for H2a-RFP than for CRT-GFP. This could be due to the different fates of the two
proteins. Whereas H2a-RFP interacts with ERAD components such as OS-9 at the ERQC (4, 9),
which would cause its retention, no such interactions are known or expected for CRT-GFP, which
would be actively shuttling between the ER periphery and the ERQC.
We then looked at another quality control component, ERManI. We had shown that ERManI resides
in quality control vesicles (QCVs) at the steady state, concentrating at the ERQC under ER stress
(6). We performed a FLIP experiment, similar to the one described, in NIH 3T3 cells expressing
H2a-RFP and ERManI-YFP under proteasomal inhibition. While H2a-RFP trafficking resembled
that of the previous experiment (Fig. 4A, B and Supplementary movie S2), the trafficking of
ERManI-YFP was very different than that of H2a-RFP and CRT-GFP. The fluorescence intensity
of ERManI-YFP, which appeared in a partially punctate vesicular pattern, decreased rapidly in the
ER periphery during the course of the time lapse (Fig. 4C) but barely changed in the ERQC, even
after 30 min, double the time of the experiment in Fig. 3C-F, suggesting that it does not recycle
back to the ER periphery.
157
Involvement of COPI and COPII coats in the trafficking to the ERQC and back
Given the results of the time lapse experiments, which indicate non-diffusive trafficking from the
ERQC to the peripheral ER, consistent with vesicular trafficking, we tested the involvement of
vesicular coats. We assessed the effect of overexpression of dominant-negative mutants of ARF1
(ARF1[T31N]) and Sar1 (Sar1[T39N]), which affect COPI and COPII coats respectively (18, 19),
on the localization of H2a-RFP. Similarly to treatment with Bz, ARF1[T31N] caused juxtanuclear
accumulation of H2a-RFP (Fig. 5A, B), which was confirmed to be in the ERQC, by comparing with
the ERQC localization of OS-9 (4) (Fig. 5C, D). In contrast, Sar1[T39N] reduced the number of cells
showing spontaneous juxtanuclear accumulation of H2a-RFP (Fig. 5A, B). A
coimmunoprecipitation experiment of H2a with OS-9 showed increased interaction in cells
overexpressing ARF1[T31N] and conversely, a reduced interaction in the presence of Sar1[T39N]
(Fig. 5E, F).
The effect on the compartmentalization by interference with COPII was also studied using iodixanol
gradients. Overexpression of Sar1[T39N] caused a shift of H2a from heavier fractions (peaking
around fractions 6-7) to lighter fractions (peaking around fractions 4-5) (Fig. 6A, B). There was a
reduction in the amount of H2a in the heavier fractions (6 to 9, which include the ERQC) from 92%
of the total protein in the absence of Sar1[T39N] to 41% in its presence (Fig. 6C). To discard the
possibility that this is an indirect effect due to the long-term (24h) interference by Sar1[T39N], a
similar experiment was done but using short-term (1h) incubation with the PKA inhibitor H89, which
inhibits COPII vesicular trafficking (20). Cells were incubated with or without Bz (1 M) for 3h with
the addition of H89 (50 M) for the last hour. The results were similar to those obtained with
Sar1[T39N]. There was a reduction of H2a in ERQC fractions, from close to 100% in untreated

cells to 34% in cells treated with H89, and in the presence of Bz from 100% to 60% with H89 (Fig.
6D-G).
It is important to note that in no case was there any detectable ERAD substrate in Golgi fractions
(peaking around fractions 1-2 as indicated by the Golgi markers Cab45 or GM130, Fig. 6 and
Supplementary Fig. 1).
Altogether these results suggest that COPII is necessary for trafficking of the ERAD substrate from
the peripheral ER to the ERQC and interaction with OS-9, whereas COPI appears to facilitate
recycling from the ERQC to the peripheral ER.
188
COPI and COPII are involved in the regulation of misfolded protein targeting to
retrotranslocation and ERAD.
The requirement of COPII function for efficient interaction of the ERAD substrate with the HRD1-
SEL1L-associated OS-9 suggested that it may also regulate its targeting to retrotranslocation and
proteasomal degradation. We tested the effect on retrotranslocation using the BAP/BirA system,
developed by the group of Burrone (21). This system utilizes the E. coli biotin ligase BirA expressed
in the cytosol of mammalian cells to specifically biotinylate a 15 amino acid biotin-acceptor peptide
(BAP). Secretory proteins with a BAP tag on their ER luminal side will be biotinylated only after
retrotranslocation. We made a construct to express H2a with BAP attached to its luminal C-
terminus (H2a is a type II membrane protein). H2a-BAP was biotinylated poorly when coexpressed
with cytosolic BirA and to a much higher extent when coexpressed with a luminally expressed
secBirA, which contains a signal peptide for translocation into the ER (21) (Fig. 7A). Biotinylated
H2a-BAP in the presence of cytosolic BirA represents retrotranslocated H2a-BAP molecules.
Interestingly, these biotinylated molecules concentrated at the ERQC fractions in iodixanol
gradients (especially upon proteasome inhibition), with only a minor fraction at light densities,
suggesting that H2a molecules remain associated to the cytosolic side of the ERQC membrane
prior to degradation (Fig. 7B).
We then tested the effect of ARF1[T31N]-HA and Sar1[T39N]-HA on retrotranslocation. The
fraction of retrotranslocated H2a-BAP increased approximately 2-fold under COPI interference
(+ARF1[T31N]-HA) and decreased to about half under COPII interference (+Sar1[T39N]-HA) (Fig.
7C). This suggests that COPII is required for efficient targeting of the ERAD substrate to
retrotranslocation. Concerning COPI, its role in recycling from the ERQC to the peripheral ER,
which we have observed in the previous experiments, could reduce the availability of molecules for
retrotranslocation. However, if COPI would be involved in events taking place after
retrotranslocation, this could also explain the increase seen by interfering with COPI.
As retrotranslocation is a rate-limiting step in ERAD, we reasoned that interference with COPI and
COPII may affect the degradation rate of the ERAD substrate. We tested this by pulse-chase
analysis, to examine H2a degradation over time under COPI/COPII interference. HEK293 cells
transiently expressing H2a were metabolically labeled with [^{35}S]cysteine. To minimize indirect
effects of long-term expression of the dominant negative constructs, we decided in this case to
incubate cells only during the chase period of up to 6h with brefeldin A (BFA) (5 µg/ml), a known
inhibitor of COPI vesicular transport, or H89 (100 µM). In untreated cells, H2a degraded with time,
and 28% of the labeled molecules remained after 6h of chase (Fig. 7D). COPII interference with
H89 inhibited the degradation, with 47% of the labeled protein remaining after 6h chase.
Interestingly, COPI interference with BFA accelerated the degradation, with 15% remaining after
6h, a reduction to almost half of the untreated control.
Altogether, these results suggest involvement of COPII in the targeting of the ERAD substrate to
retrotranslocation and degradation. In the case of COPI, the increased degradation seen by
interfering with its function suggests that it does not participate in events taking place after
retrotranslocation, but rather it functions in the recycling from the ERQC to the peripheral ER, which

reduces the availability of ERAD substrate molecules both for retrotranslocation and ultimately for
their degradation (Fig. 8).
231
232
233
234 **Discussion**
235
Previous work in our lab had characterized the ERQC as a distinct compartment, localized in the
centrosomal region of the cell and not colocalizing with the Golgi apparatus, ERGIC or other
organelles (2, 3). OS-9 resides constitutively in this compartment (4, 22), but ERAD substrates
and other machinery components are observed to concentrate at the ERQC only upon
accumulation of ERAD substrates, by inhibiting their proteasomal degradation (as done here), by
their overexpression or by induction of Herp, downstream of the activation of the PERK pathway
(3, 4). For example, recruitment to the ERQC can be achieved by overexpressing a
phosphomimetic mutant of eIF2 (S51D) (3), inhibiting eIF2 -P dephosphorylation or
overexpressing Herp (4). We had concluded from those results that even in unstressed
conditions, a transient accumulation of unfolded protein molecules causes a temporary activation
of the PERK pathway and an induction of Herp, leading to dynamic protein recruitment to the
ERQC. Therefore, ERAD machinery components are recruited according to the demand for
protein disposal and ERAD complexes might be transiently assembled only when needed,
followed by disassembly and dispersal in an "ERAD cycle". We have now observed that also
inhibition of COPI vesicular trafficking with a dominant-negative mutant of ARF1 causes an
accumulation of the ERAD substrate in the ERQC (Fig. 5), suggesting that the recycling or
dispersal is inhibited. Many quality control and ERAD components are recruited to the ERQC,
also from the cytosolic side, including proteasomes, but some are excluded, such as the
abundant chaperone BiP (Fig. 1) (5).
Both glycoproteins and non-glycosylated ERAD substrates concentrate at the ERQC, as they
share much of a common quality control machinery (1, 7). However, we have observed that the
trafficking of CRT to the ERQC is dependent on the presence of an active lectin domain (Fig. 2),
suggesting that it follows the accumulation of its glycoprotein clients. CRT appears to cycle
actively between the peripheral ER and the ERQC (Fig. 3). In contrast, another glycoprotein
quality control component, ERManI, does not recycle (Fig. 4) and seems to be effectively
sequestered in QCVs or in the ERQC.
We have described the QCVs as a new type of vesicles, where the mannosidases ERManI and
mannosidase IA (ManIA) are normally kept segregated from their substrate glycoproteins (6, 23).
Only occasionally do the glycoproteins interact with the mannosidases at the QCVs or at their
fusion sites with the ER. The QCVs are dependent on COPII, but coat components were not
observed on the vesicles. Future studies should determine whether the QCVs are the same
carriers for the trafficking of the ERAD substrates.
Interference with COPII inhibits ERQC localization of the ERAD substrate and reduces its
interaction with OS-9, retrotranslocation and degradation (Fig. 5-8). One possibility is that the
interference with COPII affects the QCVs, thus causing an inhibition of substrate interaction with
the mannosidases, which reduces mannose trimming and delivery to ERAD. However, the slow
dynamics of H2a-RFP to the ERQC (Fig. 2) and its slow recycling (Fig. 3-4) suggest vesicular
transport of the substrate and not an indirect effect of the mannose trimming. Trafficking of ERAD
substrates and recruitment of machinery components at the ERQC is also microtubule dependent
(2).
The COPII-dependent sorting to an ER subdomain, named ER-associated compartment (ERAC),
was found to be required for proteasomal degradation of an ERAD substrate, mutant CFTR, in *S.*
cerevisiae. Perhaps the ERAC is a functional homolog of the ERQC in yeast, although there are
some differences, such as the accumulation of the BiP homolog, Kar2 in the ERAC whereas BiP
does not accumulate in the ERQC (24, 25).
COPI and II vesicular trafficking is traditionally associated to movement between the ER and the
Golgi. However, COPII coats have recently been implicated in other pathways as well. For

example, in the formation of ER whorls, which appear in certain conditions and were found to be PERK and COPII dependent (13). COPII was also found to be involved in the initiation of the autophagic process in yeast (14, 15) and mammalian cells (26). Some interesting recent reports depart from the classical model, suggesting that COPII concentrates at the ER exit sites, creating a boundary, but does not coat the vesicular carriers (27, 28). COPII and also COPI involvement might be in the vesicle budding stage, with alternative factors possibly targeting the vesicles to different locations (29, 30).

There are reports of misfolded proteins that are sent back to the ER from the Golgi (31) or from the ERGIC (32). However, in the case of H2a that we have analyzed here, it is retained in the ER and does not traffic to the Golgi nor to the ERGIC (4, 5, 7).

Our results show that by interfering with misfolded protein delivery to the ERQC (Fig. 5, 6), interference with COPII surprisingly inhibits retrotranslocation and ERAD (Fig. 7, 8). COPII mutations are reported to cause ER stress, which has been interpreted as resulting from the accumulation of secretory proteins that cannot travel to the Golgi (33), but an additional cause might be an outcome of ERAD inhibition.

Materials and Methods

Materials
 Streptavidin-agarose-beads, ALLN, MG132, Bz, H89 and biotin were from Sigma. Lac was from Calbiochem. Protein A-Sepharose was from Repligen (Needham, MA). Other common reagents were from Sigma-Aldrich.

Antibodies
Rabbit polyclonal anti-H2a amino-terminal and carboxy-terminal antibodies were the one used in previous studies (34). Rabbit polyclonal anti-Cab45 was described in (35) and anti-BiP was the one used before (4). Rabbit polyclonal anti-GM130 was from BioLegend, anti-CNX from Sigma and anti-dsRED from MBL. Rabbit polyclonal anti-19S proteasome from Thermo scientific and anti-20S proteasome from Abcam and were kind gifts from Michal Sharon, Weizmann Institute. Mouse monoclonal anti-HA was from Sigma or from BioLegend, anti-actin and anti- -tubulin from Sigma, anti-S-tag from Novagen, anti-V5 from GenScript and anti-GFP from Santa Cruz Biotechnology. Strep-HRP, goat anti-mouse IgG-HRP, goat anti-rabbit IgG-HRP and goat anti-mouse IgG Dy649 were from Jackson-Immuno-Research Labs.

Plasmids and constructs
psmd14-YFP was a kind gift of A. Stanhill, (Open Univ., Israel)) (36). CRT-GFP and mutCRT-GFP: CRT and CRT(Y108F) expressed in pEGFP-C1 were kind gifts from Erik Snapp, (Janelia) (17). H2a in pcDNA1 was described before (2). BiP-GFP, H2a-RFP, S-tagged-OS-9.1 and S-tagged-OS-9.2 in pcDNA3.1(+) were used before (3, 4, 6, 23). ERManI-YFP was constructed with human ERManI cloned into peYFP-N1 (Clontech) using Hind III and Xma I yielding ERManI fused on its C-terminus to YFP through a 6 amino acid flexible linker (SGGGGS). Sar1[T39N]-HA in pTARGET or in IRES-GFP (Clontech) were used before (6). ARF1[T31N]-HA in pcDNA3.1(+)(37), ARF1[T31N]-GFP (38) and pBABE-puro were from Addgene. BirA and Sec-BirA plasmids were a kind gift from Gianlucca Petris and Oscar Burrone (ICGEB, Trieste, Italy) (21). H2a-BAP was cloned in pcDNA3.1 using restriction enzymes EcoR1 and XbaI for H2a(G78R) followed by a C-terminal BAP-SV5 tag using XbaI (39).

Cell culture, media and transfections
NIH 3T3, CHO and HEK293 cells were grown in DMEM supplemented with 10% bovine calf serum at 37 °C under 5% CO_2. Transfections of NIH 3T3 and CHO cells were carried out using a Neon MP-100 microporator system (Life Technologies, Carlsbad, CA). Transfection of HEK293 cells was performed using the calcium phosphate method. Stable CHO cell lines expressing CRT-GFP or CRT(Y108F)-GFP were obtained by co-transfection with pBABE-puro (Addgene) and selection with puromycin.

Microsome isolation and gradient fractionation
Cells were homogenized by passing through a 25-gauge needle 5 times followed by 30 strokes
with a Dounce homogenizer (Kontes glass co.) in an iso-osmotic buffer consisting of 10 mM
Hepes (pH 7.4) and 250 mM sucrose. Debris and nuclei were pelleted by centrifugation at
1,000xg at 4°C for 10 minutes. Then the supernatants were loaded on top of a iodixanol gradient
(10 to 34%) as previously described (23). The gradients were ultra-centrifuged at 24,000 rpm
(~98,500g, Beckman SW41 rotor) at 4ºC for 16 hours. Eleven fractions were collected from top to
bottom and subjected to gel electrophoresis followed by immunoblotting.

Metabolic labeling
Subconfluent (90%) cell monolayers in 60-mm dishes were labeled for 30 min with $[^{35}S]Cys$ and
chased for different periods of time with normal DMEM plus 10% FCS, lysed, and
immunoprecipitated with anti-H2a carboxy-terminal antibody as described previously (23, 34).
Quantitation was performed in a Fujifilm FLA 5100 phosphorimager (Tokyo, Japan).

Immunoprecipitation and Immunoblotting
Cell lysis and immunoprecipitation methods are described in (9). For precipitation with
streptavidin-agarose beads, cell lysates were incubated with the beads on a vertical rotating
mixer at 4°C overnight. After centrifugation, the beads were washed three times with 0.5% Triton
X-100, 0.25% NaDOC and 0.5% SDS in PBS and once with PBS. Samples were boiled in sample
buffer for 5 minutes and run on SDS-PAGE under reducing conditions. Transfer to a nitrocellulose
membrane, Immunoblotting, detection by ECL, and quantitation in a Bio-Rad ChemiDocXRS
Imaging System (Hercules, CA) were done as described previously (22). Quantity one or ImageJ
software were used for the quantitation.

BAP construct probing
BAP-tagged H2a (H2a-BAP) was transfected and probed as described before (39). Briefly, H2a-
BAP was cotransfected in HEK293 cells with an equal amount of plasmids carrying either BirA
(cytosolic biotin ligase) or Sec-BirA (ER luminal biotin ligase). Biotin $(100\mu M)$ was added 24 hours
post transfection for 30 to 180 min as indicated, cells were lysed, subjected to SDS-PAGE and
blots reacted with Strep-HRP in PBS-TWEEN (0.5%), followed by ECL detection. Total H2a-BAP
was detected with anti-H2a amino-terminal or anti-V5 antibodies.

Immunofluorescence Microscopy
Immunofluorescence was performed as described previously (2, 16, 40). Briefly, cells grown for
24 h after transfection on coverslips in 24-well plates were fixed with 3% paraformaldehyde for 30
min, incubated with 50mM glycine in PBS, and permeabilzed with 0.5% Triton X-100. After
blocking with normal goat IgG in PBS/2% BSA, they were exposed to primary antibody for 60
min, washed and incubated for 30 min with a secondary antibody, followed by washes and
staining of nuclei with DAPI. Specimens were observed with a Leica DMRBE microscope, or
confocal microscopy on a Zeiss laser scanning confocal microscope (LSM 510; Carl Zeiss, Jena,
Germany). ImageJ was used to quantify fluorescence intensity and to calculate Pearson's and
Mander's coefficients (using JACOP) for colocalization studies.

Live cell imaging
Cells were transfected and grown in 35mm culture dishes on 25mm coverslips. 24 hours post-
transfection images were captured using a Zeiss LSM 510 Meta confocal microscope. Cells were
kept in a stage incubator which provides tissue culture conditions (37°C, CO_2). For FLIP
experiments, GFP bleaching was achieved with an argon laser at a wavelength of 488nm, 100%
output. RFP bleaching was achieved with solid state laser at a wavelength of 561nm, 15% output.
Bleaching was done repetitively in an ER periphery region and scanned after each set of
bleaching (200 iterations at each time point). Analyses were performed in ImageJ software.

Statistical Analysis
The results are expressed as average ± SD or mean ± SEM as indicated. Student's t-test (two-
tailed) was used to compare the averages of two groups. Statistical significance was determined
at $P<0.05$ (*), $P<0.01$ (**), $P<0.001$ (***).

Acknowledgments

We would like to thank Erik Snapp, Ariel Stanhill, Michal Sharon, Gianlucca Petris and Oscar Burrone for plasmids and reagents. Work was supported by grants (1593/16 and 2577/20) from the Israel Science Foundation (GZL).

References

1. Benyair R, Ogen-Shtern N, & Lederkremer GZ (2015) Glycan regulation of ER-associated degradation through compartmentalization. *Seminars in cell & developmental biology* 41:99-109.

2. Kamhi-Nesher S*, et al.* (2001) A novel quality control compartment derived from the endoplasmic reticulum. *Mol Biol Cell* 12(6):1711-1723.

3. Kondratyev M, Avezov E, Shenkman M, Groisman B, & Lederkremer GZ (2007) PERK-dependent compartmentalization of ERAD and unfolded protein response machineries during ER stress. *Exp Cell Res* 313(16):3395-3407.

4. Leitman J*, et al.* (2014) Herp coordinates compartmentalization and recruitment of HRD1 and misfolded proteins for ERAD. *Mol Biol Cell* 25(7):1050-1060.

5. Shenkman M & Lederkremer GZ (2019) Compartmentalization and Selective Tagging for Disposal of Misfolded Glycoproteins. *Trends Biochem Sci* 44(10):827-836.

6. Benyair R*, et al.* (2015) Mammalian ER mannosidase I resides in quality control vesicles, where it encounters its glycoprotein substrates. *Mol Biol Cell* 26(2):172-184.

7. Shenkman M*, et al.* (2013) A Shared Endoplasmic Reticulum-associated Degradation Pathway Involving the EDEM1 Protein for Glycosylated and Nonglycosylated Proteins. *J Biol Chem* 288(4):2167-2178.

8. Frenkel Z, Shenkman M, Kondratyev M, & Lederkremer GZ (2004) Separate roles and different routing of calnexin and ERp57 in endoplasmic reticulum quality control revealed by interactions with asialoglycoprotein receptor chains. *Mol Biol Cell* 15(5):2133-2142.

9. Groisman B, Shenkman M, Ron E, & Lederkremer GZ (2011) Mannose trimming is required for delivery of a glycoprotein from EDEM1 to XTP3-B and to late endoplasmic reticulum-associated degradation steps. *J Biol Chem* 286(2):1292-1300.

10. Barlowe C*, et al.* (1994) COPII: a membrane coat formed by Sec proteins that drive vesicle budding from the endoplasmic reticulum. *Cell* 77(6):895-907.

11. Letourneur F*, et al.* (1994) Coatomer is essential for retrieval of dilysine-tagged proteins to the endoplasmic reticulum. *Cell* 79(7):1199-1207.

12. Orci L*, et al.* (1993) Budding from Golgi membranes requires the coatomer complex of non-clathrin coat proteins. *Nature* 362(6421):648-652.

13. Xu F*, et al.* (2021) COPII mitigates ER stress by promoting formation of ER whorls. *Cell Research* 31(2):141-156.

14. Cui Y, Parashar S, & Ferro-Novick S (2020) A new role for a COPII cargo adaptor in autophagy. *Autophagy* 16(2):376-378.

15. Cui Y*, et al.* (2019) A COPII subunit acts with an autophagy receptor to target endoplasmic reticulum for degradation. *Science* 365(6448):53-60.

16. Leitman J, Ulrich Hartl F, & Lederkremer GZ (2013) Soluble forms of polyQ-expanded
huntingtin rather than large aggregates cause endoplasmic reticulum stress. *Nature*
communications 4:2753.
17. Snapp EL, Sharma A, Lippincott-Schwartz J, & Hegde RS (2006) Monitoring chaperone
engagement of substrates in the endoplasmic reticulum of live cells. *Proceedings of the*
National Academy of Sciences 103(17):6536.
18. Dascher C & Balch WE (1994) Dominant inhibitory mutants of ARF1 block endoplasmic
reticulum to Golgi transport and trigger disassembly of the Golgi apparatus. *J Biol Chem*
269(2):1437-1448.
19. Weissman JT, Plutner H, & Balch WE (2001) The mammalian guanine nucleotide
exchange factor mSec12 is essential for activation of the Sar1 GTPase directing
endoplasmic reticulum export. *Traffic* 2(7):465-475.
20. Aridor M & Balch WE (2000) Kinase signaling initiates coat complex II (COPII)
recruitment and export from the mammalian endoplasmic reticulum. *J Biol Chem*
275(46):35673-35676.
21. Petris G, Vecchi L, Bestagno M, & Burrone OR (2011) Efficient detection of proteins
retro-translocated from the ER to the cytosol by in vivo biotinylation. *PLoS One*
6(8):e23712.
22. Ron E*, et al.* (2011) Bypass of glycan-dependent glycoprotein delivery to ERAD by up-
regulated EDEM1. *Mol Biol Cell* 22(21):3945-3954.
23. Ogen-Shtern N, Avezov E, Shenkman M, Benyair R, & Lederkremer GZ (2016)
Mannosidase IA is in Quality Control Vesicles and Participates in Glycoprotein Targeting
to ERAD. *Journal of molecular biology* 428(16):3194-3205.
24. Fu L & Sztul E (2003) Traffic-independent function of the Sar1p/COPII machinery in
proteasomal sorting of the cystic fibrosis transmembrane conductance regulator. *J Cell*
Biol 160(2):157-163.
25. Kakoi S, Yorimitsu T, & Sato K (2013) COPII machinery cooperates with ER-localized
Hsp40 to sequester misfolded membrane proteins into ER-associated compartments.
Mol Biol Cell 24(5):633-642.
26. Ge L, Wilz L, & Schekman R (2015) Biogenesis of autophagosomal precursors for LC3
lipidation from the ER-Golgi intermediate compartment. *Autophagy* 11(12):2372-2374.
27. Shomron O*, et al.* (2021) COPII collar defines the boundary between ER and ER exit site
and does not coat cargo containers. *Journal of Cell Biology* 220(6):e201907224.
28. Weigel AV*, et al.* (2021) ER-to-Golgi protein delivery through an interwoven, tubular
network extending from ER. *Cell* 184(9):2412-2429.e2416.
29. Phuyal S & Farhan H (2021) Want to leave the ER? We offer vesicles, tubules, and
tunnels. *J Cell Biol* 220(6).
30. Westrate LM, Hoyer MJ, Nash MJ, & Voeltz GK (2020) Vesicular and uncoated Rab1-
dependent cargo carriers facilitate ER to Golgi transport. *J Cell Sci* 133(14).
31. Hellerschmied D, Serebrenik YV, Shao L, Burslem GM, & Crews CM (2019) Protein
folding state-dependent sorting at the Golgi apparatus. *Mol Biol Cell* 30(17):2296-2308.
32. Sirkis DW, Aparicio RE, & Schekman R (2017) Neurodegeneration-associated mutant
TREM2 proteins abortively cycle between the ER and ER-Golgi intermediate
compartment. *Mol Biol Cell* 28(20):2723-2733.
33. Lu CL & Kim J (2020) Consequences of mutations in the genes of the ER export
machinery COPII in vertebrates. *Cell Stress Chaperones* 25(2):199-209.

34.   Tolchinsky S, Yuk MH, Ayalon M, Lodish HF, & Lederkremer GZ (1996) Membrane-bound
versus secreted forms of human asialoglycoprotein receptor subunits - Role of a
juxtamembrane pentapeptide. *J Biol Chem* 271(24):14496-14503.
35.   Scherer PE*, et al.* (1996) Cab45, a novel (Ca2+)-binding protein localized to the Golgi
lumen. *J Cell Biol* 133(2):257-268.
36.   Sukenik S, Braunstein I, & Stanhill A (2021) An Arsenite Relay between PSMD14 and
AIRAP Enables Revival of Proteasomal DUB Activity. *Biomolecules* 11(9):1317.
37.   Furman C, Short SM, Subramanian RR, Zetter BR, & Roberts TM (2002) DEF-1/ASAP1 is a
GTPase-activating protein (GAP) for ARF1 that enhances cell motility through a GAP-
dependent mechanism. *J Biol Chem* 277(10):7962-7969.
38.   Moorhead AM, Jung JY, Smirnov A, Kaufer S, & Scidmore MA (2010) Multiple host
proteins that function in phosphatidylinositol-4-phosphate metabolism are recruited to
the chlamydial inclusion. *Infect Immun* 78(5):1990-2007.
39.   Sharma N*, et al.* (2021) The Sigma-1 receptor is an ER-localized type II membrane
protein. *J Biol Chem* 297(5):101299.
40.   Avezov E, Frenkel Z, Ehrlich M, Herscovics A, & Lederkremer GZ (2008) Endoplasmic
reticulum (ER) mannosidase I is compartmentalized and required for N-glycan trimming
to Man5-6GlcNAc2 in glycoprotein ER-associated degradation. *Mol Biol Cell* 19(1):216-
225.

502
503
504

505   **Figures**
506

507
Figure 1. ERAD substrate H2a-RFP, CRT-GFP and members of the ERAD machinery
accumulate at the ERQC upon proteasomal inhibition. (A-C) Fluorescence microscopy of NIH
3T3 cells treated without or with the proteasomal inhibitor Lactacystin (Lac, 20μM, 3h), transiently
expressing H2a-RFP and CRT-GFP (A), BiP-GFP (B) or psMD14-YFP (C). H2a-RFP
concentrates and colocalizes with CRT-GFP (A) and with the proteasomal subunit psMD14-YFP
(C) at the juxtanuclear ERQC, especially in Lac-treated cells. In contrast, there is almost no
colocalization of H2a-RFP and BiP-GFP in Lac-treated cells (B). Bars= 10μm. **(D)** Immunoblots
of fractions of iodixanol density gradients from NIH 3T3 cells transiently expressing CRT-GFP
and H2a-RFP show comigration of CRT-GFP, H2a-RFP and endogenous CNX in mid-density
fractions (5-7), shifting to denser fractions (6-8) (ERQC) upon MG-132+ALLN (50μM, 3h)
treatment, compared to untreated cells. BiP appears in lighter fractions and does not shift upon
proteasomal inhibition. MW markers in kDa are indicated on the right. The graphs show
quantification of the percentage of the signal in each fraction relative to the sum of all fractions.
The experiment is representative of 3 independent repeat experiments. **(E)** Most of the
endogenous proteasome subunits 19S and 20S appear in similar mid-density fractions upon
iodixanol fractionation of Bz-treated cells.
524
525
526

528
Figure 2. Movement of H2a-RFP and CRT-GFP to the ERQC. CRT-GFP dependence on its
lectin activity. **(A)** CHO cells stably expressing CRT-GFP were transiently transfected with H2a-
RFP and treated with Lac (10μM). Live cell imaging shows the dynamic concentration of both
proteins in the juxtanuclear ERQC. Bar= 10μm.   **(B)** The percentage of H2a-RFP and CRT-GFP
concentrated in the ERQC (dashed circle in (A)) was calculated at the different time points. **(C-D)**
CHO cells stably expressing mutCRT-GFP, lacking a functional lectin domain, were transfected
with H2a-RFP and treated with 10μM Lac. Live cell imaging shows absence of mutCRT-GFP
accumulation at the ERQC. Bar= 10μm. **(E)** Pearson's coefficients of colocalization of CRT-GFP
or mutCRT-GFP with H2a-RFP in the entire cell or in the area of the ERQC were measured at
different time points in the experiments in (A) and (B). mutCRT-GFP showed no colocalization
with H2a-RFP. The results are representative of 3 independent experiments. **(F)** In Lac-treated
NIH 3T3 cells (20μM, 3h), CRT-GFP and H2a-RFP accumulate in the ERQC, whereas untreated
cells show an ER pattern. No colocalization appears in the ERQC (dashed circle) of NIH 3T3 cells
expressing mutCRT-GFP and H2a-RFP after incubation with Lac, compared to untreated cells.
(G) The percentage of GFP or RFP fluorescence in the ERQC compared to the entire cell was
measured in Lac-treated cells expressing CRT-GFP plus H2a-RFP or mutCRT-GFP plus H2a-

RFP. The graph shows mean +-SEM of cells from two independent experiments (~20 cells per
population). P value CRT-GFP vs. mutCRT-GFP = 0.00014.
547
548
549

Figure 3. Recycling from the ERQC to the ER is slower than movement within the ER periphery. (A) NIH 3T3 cells transiently expressing CRT-GFP and H2a-RFP treated with MG-132 (50μM, 3h), followed by drug washout. The accumulation in the ERQC after treatment reverts to an ER pattern after washout. Bar= 10μm. **(B)** Percent of cells with ERQC accumulation of H2a-RFP or CRT-GFP at the different times, average of 3 independent experiments +-SD. P values for H2a-RFP, +MG-132 vs. untreated = 0.0007, 90 min washout vs. +MG-132 =0.0002, 150 min washout vs. +MG-132 =0.0002. **(C)** Fluorescence loss in photobleaching (FLIP). Lac-treated NIH 3T3 cells (20μM, 3h) were repetitively bleached in an ER periphery region (yellow rectangles) and scanned after each set of bleaching. Fluorescence intensity was measured at the ERQC (white arrow), the ER periphery (yellow arrow) and the bleached region. Representative of 4 experiments. See Suppl. movie S1. **(D-E)** Loss of the fluorescence intensity during FLIP compared to the start point for H2a-RFP (D) and for CRT-GFP (E), in the ERQC, ER periphery, bleached region and in a control cell, during the first 400 sec. Recycling from the ERQC to the periphery is much slower than diffusion in the periphery. Fluorescence in the bleached region was lost quickly, while there was no non-specific bleaching of a nearby but non-irradiated control cell. **(F)** Ratio between the FLIP curve slopes (calculated from linear trendlines) for each protein in the ER periphery compared to the ERQC, average of 4 independent experiments +-SD. P value = 0.049. The relative recycling rate from the ERQC is much slower for H2a-RFP than for CRT-GFP.

Figure 4. No recycling of ER mannosidase I from the ERQC to the ER. (A) Lac-treated NIH
3T3 cells (20µM, 3h) transfected with H2a-RFP and ERManI-YFP expressing plasmids were
repetitively bleached in an ER periphery ROI (yellow rectangles) and scanned after each set of
bleaching as in Fig. 3C. Bar= 10µm. (**B-C**) The percentage of the fluorescence intensity
compared to the start (prior to bleaching) was calculated for each time point for H2a-RFP (B) and
for ERmanI-YFP (C), indicating no measurable recycling of ERmanI-YFP from the ERQC to the
periphery during the time of the experiment. The trendlines in the peripheral region are
polynomial (fourth order) and for H2a-RFP in the ERQC it is polynomial of the second order. See
Suppl. movie S2.

580
Fig. 5. Vesicular coats are involved in trafficking from the peripheral ER to the ERQC and
back. (A-B) NIH 3T3 cells transiently expressing H2a-RFP together with GFP, ARF1[T31N]-GFP
(dnARF1) or Sar1[T39N]-IRES-GFP (dnSar1). Where indicated, cells were treated for 3 h with Bz
(1µM). Bar= 10µm. The graph shows percent of cells with ERQC accumulation of H2a-RFP,
average of three independent experiments +-SD, which increased +Bz and +dnARF1, and
decreased + dnSar1. P values +Bz vs. control = 0.05, +dnARF1 vs. control = 0.005, +dnSar1 vs.
control = 0.05. **(C-D)** NIH 3T3 cells expressing H2a-RFP and S-tagged OS-9.1/2, with or without
ARF1[T31N]-GFP fixed and stained with anti-S-tag antibody and goat anti-mouse IgG Dy649.
Mander's coefficients indicate that dnARF1 increased H2a-RFP colocalization with OS-9. The
graph is a box plot with inclusive medians of 3 independent experiments (~30 cells per
experiment); box height=IQR; horizontal bar in the box=median; whiskers extend to the farthest
value smaller than 1.5xIQR. P value = 0.0003. **(E)** Immunoprecipitation with anti-H2a of lysates of
HEK293 cells expressing H2a together with combinations of OS-9.1/2-S, ARF1[T31N]-HA or
Sar1[T39N]-HA, as indicated. Immunoprecipitates and 10 % of the total lysates were
immunoblotted with anti-S-tag, anti-H2a, anti-HA and actin (loading control) antibodies. MW
markers in kDa are on the left. **(F)** The graph shows increased coimmunoprecipitation of OS-9
with H2a under COPI interference and decreased under COPII interference. Values are relative
to control (=100), normalized by total H2a and total OS-9, average of 3 independent experiments
+-SD. P value +dnSar1 vs. control = 0.002.

Fig. 6. Interference with COPII inhibits ERAD substrate accumulation in the ERQC. (A)
Homogenates from HEK293 cells expressing H2a with or without Sar1[T39N]-HA (dnSar1) were
loaded on an iodixanol gradient (10%-34%) as indicated in Methods. Eleven fractions were
collected from the top to the bottom, run on 12% SDS-PAGE and immunoblotted with anti-H2a,
anti-Cab45 (Golgi marker), anti-20S and anti-CNX antibodies (peak fractions indicated with
arrows). The results shown are representative of 3 independent experiments. **(B)** After
quantification, the percentage of total H2a in each fraction was plotted. **(C)** Comparison between
the percentage of total H2a in the sum of fractions 3 to 5 and that of fractions 6 to 9 (ERQC)
shows that most H2a shifts to lighter fractions under COPII interference by dnSar1. **(D)** Similar to
(A), but cells expressing H2a were incubated with 50μM H89 for 1h or 1μM Bz for 3h, or with 1μM
Bz for 3h adding 50μM H89 in the last hour of treatment. Anti-GM130 was used here as a Golgi
marker. The results shown are representative of 3 independent experiments. **(E-F)** The intensity
of each band was quantified and the percentage of total H2a in each fraction was plotted and
compared between the control and +H89 (E), or +Bz and +Bz+H89 (F). **(G)** Comparison between
the percentage of total H2a in the sum of fractions 3 to 5 and that of fractions 6 to 9 (ERQC)
shows that H2a shifts to lighter fractions under COPII interference by H89 treatment.

619
Fig. 7. COPII-dependent movement from the peripheral ER to the ERQC is required for
ERAD substrate retrotranslocation and degradation. (A) HEK293 cells transiently expressing
H2a-BAP with or without a cytosolic biotin ligase, BirA, or an ER luminal one, Sec-BirA. Cells
were incubated 24h post-transfection with biotin (100μM) for 30 min before lysis. Cytosolic BirA
biotinylates BAP exposed on the cytosolic side of the ER upon retrotranslocation of H2a-BAP,
while Sec-BirA biotinylates luminal H2a-BAP. Strep-HRP was used to detect biotinylated H2a-
BAP and anti-V5 to detect total H2a-BAP.MW markers (kDa) on the right. **(B)** Fractions of
iodixanol gradients from cells expressing H2a-BAP and BirA, were immunoblotted as in (A), (with
anti-H2a instead of anti-V5). Both retrotranslocated and total H2a-BAP peak in fractions 6-7,
which include the ERQC. **(C)** Biotinylated proteins from cells expressing H2a-BAP together with
BirA, ARF1[T31N]-HA or Sar1[T39N]-HA, (biotin for 3h), were precipitated with streptavidin-
agarose. Precipitates and 10% of total cell lysates were immunoblotted with anti-H2a, anti-HA-tag
or anti-tubulin antibodies. Retrotranslocated (biotinylated) H2a-BAP increased almost two-fold
under COPI interference and was reduced to half under COPII interference. Values are averages
of 4 independent experiments ±SD, relative to total H2a-BAP and standardized by the control
(=1). P value +dnARF1 vs. control = 0.007, +dnSar1 vs. control = 0.02. **(D)** Pulse-labeling for 20
min with [^{35}S]cysteine and chase for the indicated times with or without 5μg/ml BFA or 100μM
H89. H2a degradation (percentage remaining relative to pulse) increased under COPI
interference (+BFA) and was strongly inhibited under COPII interference (+H89). The graph
shows an average of 3 independent experiments ±SD. P value 6h +BFA vs. control = 0.03, 6h
+H89 vs. control = 0.01, 4h +H89 vs. control = 0.003.

Fig. 8. Model of COPI and COPII involvement. COPII is involved in the movement of ERAD substrates from the peripheral ER to the ERQC. Under COPII interference this movement is inhibited. The ERAD substrates are then mostly scattered in the peripheral ER instead of accumulating in the ERQC. The retrotranslocation is then decreased, followed by a slower degradation rate (light blue arrows). The recycling of ERAD substrates from the ERQC to the ER is dependent on COPI. COPI interference hinders this recycling, causing accumulation of ERAD substrates in the ERQC. The retrotranslocation is then increased and the degradation of the substrates is accelerated (orange arrows).