Informative title

Plant protection habits among Hungarian farmers and garden owners and their

implications for pollinator protection

Concise title

Pesticide use and pollinator support in Hungarian plant growers

Zsófia Varga-Szilay[1*], Gábor Pozsgai[2]

[1]Doctoral School of Biology, Institute of Biology, ELTE Eötvös Loránd University,

Budapest, Hungary

[2]cE3c – Centre for Ecology, Evolution and Environmental Changes/Azorean Biodiversity

Group, CHANGE – Global Change and Sustainability Institute, Departamento de

Ciências e Engenharia do Ambiente, Universidade dos Açores, Açores, Portugal

*Corresponding author: Zsófia Varga-Szilay, Doctoral School of Biology, Institute of

Biology, ELTE Eötvös Loránd University, 1117 Budapest, Hungary, email:

zsofia@vargaszilay.hu, https://orcid.org/0000-0001-9712-7654

Abstract

Pesticides are one of the most important anthropogenic-related stressors. In times of global pollinator decline, the role of integrated farming and that of urban gardens in supporting wild pollinators is becoming increasingly important. We circulated an online questionnaire to survey the plant protection practices among Hungarian farmers and garden owners with a particular emphasis on pollinator protection. We found that plant growers heavily rely on pesticide use, and pesticides are wildly used in otherwise pollinator-friendly gardens. Whether pesticide use practices were driven by expert opinion and the respondents' sex were the best predictors of pesticide use. Although most respondents supported pollinators, pesticides are also widely used among home garden owners, which can pose a non-evident ecological trap for pollinator populations in the gardens. Special attention should be paid to implementing measures to reduce pesticide, use not only in farmlands but also in home gardens. Environmental education and financial support through agroecological schemes could efficiently promote the transition. However, whereas farmers can be encouraged to reduce pesticide use mostly by expert advice, garden owners are likely to rely on more conventional information channels. The attitude of Hungarian plant growers can provide an insight into pesticide use practices of Central and Eastern European countries, but similar surveys are needed across Europe for a complete understanding of broad-scale processes. This work lays the foundations for similar studies which can inform and facilitate the transformation processes to pesticide-free farming and gardening.

Keywords: pest management, plant growing practices, synthetic pesticides, neonicotinoids, pollinator-friendly gardens, ecological trap

Introduction

In the Anthropocene, biodiversity declines at an alarming pace. One of the important groups, the insects, is among the most impacted (e.g. Hallmann et al. 2017; Zattara and Aizen 2021). Insect declines pose a major threat to a variety of ecosystem functions and the delivery of the derived ecosystem services, all of which are vital to humans (Dangles and Casas 2019; Vanbergen and the Insect Pollinators Initiative 2013). One important ecosystem service, mostly provided by insects, is pollination (Hanley et al. 2015). Insect pollinators suffer from habitat fragmentation, reduction of flower resources, lack of nesting space, as well as from exposure to pesticides from agricultural activities (Baldock 2020; Goulson and Nicholls 2022; Potts et al. 2016). Despite their long-known negative effects on human and environmental health (Rani et al. 2021, Zaller 2020), pesticides are widely used both in industrial-scale farming and urban green areas and their application has even increased with agricultural intensification in recent decades (FAO 2021). Indeed, the spillover of chemical insecticide residues from farmland can negatively affect wild insect pollinators in adjacent natural and semi-natural areas (Botías et al. 2015; Krupke et al. 2012) causing direct mortality, behavioural abnormalities, and reduced reproduction rates (Pisa et al. 2015). Furthermore, the concomitant use of agrochemicals (pesticides and fertilisers) can cause an even more detrimental 'cocktail effect' to insect pollinators (Raimets et al. 2018; Siviter et al. 2021; Wernecke and Castle 2020). In fact, a combination of over sixteen different agrochemicals was detected in flying insects in nature conservation areas adjacent to agricultural lands across Germany (Brühl et al. 2021) and the USA (Main et al. 2020). Thus, agrochemicals are suggested to play a major role in driving global insect declines (Sánchez-Bayo and Wyckhuys 2019; Tooker and Pearsons 2021), particularly on farmlands (Emmerson et al. 2016).

To address biodiversity loss on farmlands, particularly that of pollinators, the European Commission created a farm strategy to cut the use of chemical pesticides in European countries (European Commission 2020) and the reduction or complete elimination of pesticide use has been advised by the scientific community (e.g. Goulson and Nicholls 2022). A number of modern synthetic pesticides have been banned (e.g. in the European Union all neonicotinoids except acetamiprid) after they have been proven to harm non-target insects (like bees) in addition to the pest species targetted. In fact, the transition to alternative agricultural practices is possible without yield losses (EPRS 2019) whilst pest damage can be reduced and farm profitability maintained after lowering, but not completely abandoning, pesticide use (Lechenet et al. 2017; Pecenka et al. 2021). Despite the increasing number of organic farms (European Commission n.d. a; Willer and Lernoud 2018) in the European Union, the process leading to biodiversity-friendly farming can take years because the current conventional plant protection strategies employed on non-organic farms still require synthetic pesticide input. Nevertheless, evidence suggests that these integrated efforts may be a first step toward maintaining healthy ecosystems. For example, management that promotes ecosystem services (such as biological control or pollination) can support high insect diversity in areas of agricultural mosaics (Grass et al. 2019). Moreover, even in conventionally managed farms, increasing the proportion of semi-natural habitats, such as hedges or field-edge flower strips, can dramatically increase the diversity of insects that are beneficial to agriculture, including that of pollinators (Gurr et al. 2017; Pywell et al. 2015). However, although increasing natural habitat areas or employing other integrated pest management approaches can lead to increased pollinator and other insect diversity, unless these ecosystem-based approaches are combined with pollinator-friendly management, their

positive effect will be reduced or completely eliminated by the use of synthetic pesticides

(Brühl et al. 2022; Saqib et al. 2020; Tscharntke et al. 2021, 2022). Since socio-economic

factors can dictate how rapidly the transition to pesticide-free farming unfolds, knowing

farmers' approaches to these novel strategies is essential for future planning.

Whilst it may be difficult to achieve pesticide-free pest control within high-

intensity farming (especially in monocultures), it may be a more feasible approach in

small-scale farms and urban areas. Small-scale sustainable farming systems and well-

planned urban green areas, such as biodiversity-friendly parks and allotments

(community gardens), can mitigate pollinator declines (Baldock et al., 2019; Hülsmann

et al., 2015). In fact, in a landscape mosaic with a high proportion of urban areas,

organically managed parcels of land can maintain high biodiversity and serve as a source

of native pollinators within a landscape where most land is not managed with the

maintenance of biodiversity as a key goal (Holzschuh et al. 2008; Wray and Elle 2015).

Moreover, an increasing number of scientific papers support the premise that

urban and suburban gardens function as refuges and local hotspots for biodiversity (Hall

et al. 2017; Theodorou et al. 2020), and support diverse communities of insect pollinators,

even in highly urbanised areas (Sirohi et al. 2015). Near-natural gardens can even support

viable metapopulations of rare species (Baldock et al., 2015). However, the true

conservation potential of human-altered areas for pollinators depends on the available

floral resources, nesting and hiding spaces, and on the proportion of near natural areas

that can be found in the urban landscape. These factors also determine the abundance and

diversity of pollinator communities (Hülsmann et al. 2015; Wenzel et al. 2020).

Urban gardens may not be always beneficial for insect pollinators though. First,

there is a wide selection of pesticides in shops and supermarkets that are targeted at

domestic users and which may be applied in otherwise pollinator-friendly gardens.

Second, synthetic pesticides can also get into gardens unintentionally when ornamental

plants sold as 'bee-friendly' in horticultures are treated with various fungicides and

insecticides (Lentola et al. 2017). As a consequence, insects lured to supposedly bee-

friendly gardens can be exposed to a number of synthetic pesticides (especially

neonicotinoids) and their residues and this exposure, in turn, can lead to lethal and

sublethal effects (Botías et al. 2017; Main et al. 2020). The process of banning synthetic

pesticides for non-agricultural uses has already begun in some European countries (such

as France (Garden Organic n.d.; PAN Europe 2019; PAN Europe n.d.)) but others,

including Hungary, are lagging behind. Yet, we have no information on what proportion

of private gardens are treated with chemical plant protection products.

There is a large knowledge gap in our understanding of how efficiently farmlands

and urban and suburban gardens mitigate insect biodiversity loss at a country scale and

how farmers and garden owners approach the transition away from the use of pesticides.

Gaining insight into their management habits, motivations and willingness to change is

vital for developing further action.

Thus, we conducted a survey to measure plant growers' dependence on pesticides

(highlighting acetamiprid-containing insecticides), particularly to investigate the

pesticide application practices and the attitude towards protecting wild pollinators of

those who own less than one hectare of land (henceforth home gardens or gardens). We

focussed our work on Hungary, a typical, Central-Eastern European country with mainly

conventional agriculture in which chemical and more hazardous pesticide use trends are

likely to reflect those of general Europeans (European Commission n.d. b).

We aimed to investigate 1) what factors best predict pesticide use in agricultural areas and to what extent plant growers think their application is necessary, 2) to what extent plant growers think the use of insecticides (as a subset of pesticides) is necessary and what are the main considerations determining their selection, 3) how dependent plant growers are on the single currently allowed neonicotinoid (acetamiprid), and 4) if acetamiprid is used, what other pesticides are used simultaneously. Our additional aims were to specifically investigate the home garden owners' approach to pesticide use and pollinator support. We were interested in 1) how necessary garden owners think it is to use pesticides, 2) what they think about the threats to wild pollinators and how this affects their management practices and 3) what factors predict whether or not gardeners provide support for wild pollinators and what the most common such forms of support are.

We hypothesised that Hungarian plant growers are highly dependent on pesticide input and home garden owners have little awareness of linked environmental issues. Nevertheless, we predicted that home garden owners who predominantly produced for their own needs were more aware of the environmental hazards of pesticides than large-scale farmers and we also hypothesised that those who supported pollinators used pesticides less frequently.

Material and methods

Questionnaire Design

We circulated an online questionnaire that consisted of 61 closed-ended questions, all of which were mandatory to respond to. The questionnaire had eight sections to collect information about 1) sociodemographic factors, 2) type of farming, 3) use of plant protection products and 4-6) insecticides and their means of application, 7) protection of

wild pollinators, and the questionnaire included one question 8) about how the

questionnaire reached respondents (**Electronic Supplementary Material S1**). All

responses were recorded anonymously, however, respondents could provide their email

addresses at the end. The questionnaire was designed in Google Forms and circulated in

Hungarian language. The questionnaire was shared on social media platforms (such as

Facebook groups and Facebook pages, and agricultural websites) and on farming and

entomological mailing lists. The form was available from 26 April to 20 August in 2021.

Respondents who do not farm in Hungary were excluded from the analysis and

data from Pest county were merged with those from Budapest. The number of respondents

was standardised for 100,000 inhabitants in Hungary to improve representativeness.

In this study, we include both chemical and non-chemical pesticides in the group

of 'pesticides' and 'plant protection products'. We also use the word 'insecticide'

inclusively for synthetic insecticides and insecticides that can be used in organic farming.

Data processing and statistical analysis

The original categorical replies were on a few occasions re-categorised for analytical

purposes. Education categories were merged into 'elementary', 'middle', 'high', and

'postgraduate' levels. The most important sociodemographic parameters and the

categories used are listed in **Table 1**, and all other parameters can be found in **Electronic**

Supplementary Material S1. Although they were separated in the original questionnaire,

the two kinds of agricultural experts, 'plant doctors' and 'plant protection experts' were

later merged into a combined 'expert' category. When the additional plant protection

products which were used with acetamiprid were named, they were assigned into nine

categories or the combination of those, as 'adhesion promoter', 'insecticide(s)',

'insecticide(s) and acaricide(s)', 'insecticide(s) and fungicide(s)', 'insecticide(s),

fungicide(s) and fertiliser', 'fertiliser', 'fungicide(s)', 'fungicide(s) and acaricide(s)',

'fungicide(s) and fertiliser'. In the question about how respondents support pollinators

the textual responses for food and habitat provision-related answers may overlap although

when categorising these, we choose the one which was most strongly emphasised by the

respondent. In the same question, we did not create a separate category for 'outreach',

because it only occurred in a single response. When textual responses were given to the

types of support which could not have been categorised as direct action (e.g. 'I do not

harm them'), they were interpreted as 'no support'.

When the approach solely of garden owners to pesticide use and their attitude to

wild pollinators were investigated, only landowners with less than 1-hectare land were

included in the analysis.

We used Spearman correlation tests to examine if sociodemographic factors and

farming habits correlate with pesticide use. P-values were corrected according to Holm's

method. For calculating the correlation matrix the '*psych*' (version 2.1.9) (Revelle 2021),

and for visualising them the '*corrplot*' (version 0.92) (Wei and Simko 2021) R packages

were used. We used the chi-square test to compare plant protection habits of home garden

owners and large-scale farmers. We used machine learning techniques with Gradient

Boosting Machine (GBM) for generating our models to investigate 1) which

socioeconomic factors determine whether or not pesticides were used in farmlands, and

2) which socioeconomic factors determine whether or not pollinators were supported in

home gardens. The model fit was evaluated using the Area Under the Curve (AUC) score

and by examining the accuracy of the best fitting model. We used the '*gbm*' (version

2.1.8) (Greenwell et al. 2020), '*caret*' (version 6.0.90) (Kuhn 2021), and '*yardstick*'

(version 0.0.9) (Kuhn and Vaughan 2021) R packages for modelling. Likert Scales figures

were plotted using the '*likert*' (version 1.3.5) (Bryer and Speerschneider 2016) and the

map was created using the '*sf*' (version 1.0.3) (Pebesma 2018) R packages. All analyses

were done and figures were created using the R 4.1.1 statistical software (R Core Team

2021).

Results

218 Of the 463 people who completed the questionnaire, 246 were male, 214 were female,

219 and three did not state their gender. The willingness to respond was slightly unbalanced,

220 as more responses were received from the western than from the eastern counties (**Figure**

221 **1**). Pest country was the region that yielded the largest proportion of responses (22.0%).

 Figure 1. Distribution of respondents by gender ("M" = male, "F" = female, "O" =

unknown). Y-axis shows the percentage of the sexes and the numbers on the barplots

indicate the exact number of respondents in the 19 counties of Hungary. County names

are indicated in bold. The colour depth in the map indicates the number of respondents

per 100,000 inhabitants. Note that the sum of the numbers indicated on the barplots is

greater than the number of respondents because respondents who farmed in more than

one county were counted multiple times.

Among the respondents, the two middle-age categories (36–45 and 46–55 years old) were

the most frequent, and 60.3% of all respondents (n = 279) fall into the high-level (but not

Ph.D.) education group. Of all plant growers, 302 (65.2%) had less than one hectare of

farming area (**Table 1**). The most commonly grown crops were vegetables and fruits,

followed by grapes and root/tuberous plants (**Electronic Supplementary Material S2**).

Of the respondents, 181 (39.1%) used a pest forecasting system and 370 growers (79.9%)

supported natural enemies of pests (**Electronic Supplementary Material S3**).

Table 1. Sociodemographic characteristics of the study population (n = 463)

Variables		Total (n = 463)	
		n	%
Sex	Male	246	53.1
	Female	214	46.2
	Unknown	3	0.6
Age	18-25	27	5.8
	26-35	73	15.8
	36-45	126	27.2
	46-55	117	25.3
	56-65	62	13.4

	over 65	58	12.5
Education level	Elementary	6	1.3
	Middle	139	30.0
	High	279	60.3
	Postgraduate	39	8.4
Residence type	Farmland	20	4.3
	Countryside	174	37.6
	Town	136	29.4
	Major city	93	20.1
	Capital	40	8.6
Farming area size	< 1 ha	302	65.2
	1-9.9 ha	58	12.5
	10-29.9 ha	25	5.4
	30-49.9 ha	8	1.7
	50-99.9 ha	16	3.5
	100-299.9 ha	14	3.0
	300-499.9 ha	11	2.4
	500-999.9 ha	7	1.5
	≥ 1.000 ha	22	1.5

Plant protection habits of plant growers

The majority of plant growers in an area of less than one hectare were individual farmers

who produce exclusively for their own consumption (n = 251), while the majority of

farmers in an area larger than one hectare either produce for sale privately or as part of a

farmers' association (n = 107). Of these smallholders, over 95% used pesticides (**Table**

2). However, of all respondents, 311 (67.2%) used pesticides, and 212 of them (68.2%)

used them together with some additives. Among the pesticide users, 244 (78.5%) usually

did not spray during daytime in a flowering culture (**Electronic Supplementary**

Material 4). Of those plant growers who used pesticides, 243 (79.0 %) felt these products

were necessary for farming, with 150 (48.2%) of users considering pesticides as being

crucial, and 93 (29.9%) of them regarding them as important.

Table 2. Distribution of farming types among the study population (n = 463)

Area size	Type of farming	Uses pesticide		Pesticide-free	
		n	%	n	%
<1ha	Individual: production for oneself	134	53.4	117	46.6
	Individual: production for oneself and sale	27	69.2	12	30.8
	Individual: production for sale	9	81.8	2	18.2
	In farmers' association	1	100.0	-	-
>1ha	Individual: production for oneself	5	50.0	5	50.0
	Individual: production for oneself and sale	33	75.0	11	25.0
	Individual: production for sale	59	95.2	3	4.8
	In farmers' association	43	95.6	2	4.4

The sociodemographic and farming habit factors that were examined did not show a

strong correlation with pesticide use (and nor with each other); the highest significant

correlation ($p < 0.05$, Spearman's Rho = 0.35) was with what growers thought about the

risks of pesticide use (**Electronic Supplementary Material S5**). However, the GBM

model suggested that the best predictors for pesticide use in agricultural areas were if the

respondents had consulted with an expert or were themselves trained agricultural experts

(relative influence: 27.04, 84.4% of those who do versus 42.0% of those who do not

consult with experts, or are expert themselves, used pesticides) and the respondents'

gender (relative influence: 18.41, 50.5% of females versus 82.1% of males used

pesticides) (model accuracy: 0.79, AUC = 0.86, Sensitivity: 0.89, Specificity: 0.60)

(**Figure 2, Electronic Supplementary Material S6**).

Figure 2. The relative influence of factors generated from the Gradient Boosting

Machine (GBM) model for predicting pesticides used in farming areas.

Out of the 311 pesticide users, 243 (78.1%) felt that the use of insecticide was particularly

indispensable for them. The main aspects that determined the choice of an insecticide

were if they were harmless to humans, posed a low risk for bees and whether growers had

previous experience with the product. The techniques by which the insecticide can be

applied (such as if they can be used in flowering crops, if they can be used during daytime,

or if they can be sprayed from the air) were the least important aspects to users (**Figure**

3).

Figure 3. Relative frequency (%) of respondents' opinions on how important different

considerations are when choosing an insecticide. The response is colour-coded as

follows: dark grey – not important, light grey – negligible importance, light yellow –

moderately, light orange – important, dark orange – crucial.

Of those who used pesticides, 143 (46.0%) thought that banning neonicotinoids in the EU

impacted their management practices, and 218 (70.1%) used at a minimum one

acetamiprid-containing insecticides. Most users (55.2%) consider Mospilan (an

acetamiprid-containing insecticide) indispensable. Of those who used Mospilan, 124

(56.11%) use it together with other pesticides, mostly with fungicides.

Plant protection habits and the protection of pollinators of garden owners

Of all questionnaire respondents, 302 (65.2%) had less than one hectare of land and 171 (56.6%) of home garden owners used pesticides on their land. The use of pesticides was considered acceptable by most garden owners, and 34.5% and 31.6% of them even thought it was important or crucially important, respectively. However, a significantly lower proportion of garden owners than of larger-scale farmers used pesticides (Chi-squared = 42.455, p-value <0.001) and a significantly higher proportion of home garden owners than of large-scale farmers believed that pesticide-free farming is achievable (Chi-squared = 3.593, p-value = 0.029).

The home garden owners who responded to our questionnaire specified that widespread use of pesticides, habitat loss due to agriculture and intensive agricultural production were the three most likely threats for wild pollinators (**Figure 4**), whilst they thought the appearance of invasive species was the least significant. Nonetheless, this factor was labelled as crucial by over half of the respondents (Figure 4). Of these garden owners, 259 (85.8%) recognise that widespread use of pesticides is a crucial problem for wild pollinators and 87.7% have heard that certain pesticides that are considered safe may also harm these insects. A significantly higher proportion of home garden owners than of large-scale farmers assumed that the conversion of agricultural production can slow down the depletion of pollinator populations (Chi-squared = 10.998, p-value = <0.001).

Figure 4. Relative frequency (%) of garden owners' (with less than 1-hectare of land) opinions about the importance of factors that may threaten wild pollinators. The response is colour-coded as follows: grey – not at all, yellow – negligible, orange – crucial.

Of the garden owners, 81.1% carried out actions aimed at supporting wild pollinators. The examined sociodemographic and farming habits did not show a strong correlation with whether or not pollinators were supported (and neither did they with each other); yet the highest significant correlation with pollinator support was the growers' pesticide use (p < 0.05, Spearman's Rho = -0.22) (**Electronic Supplementary Material S7**). The GBM model suggested that the best predictors for supporting pollinators were whether garden owners had promoted biocontrol (relative influence: 38.85) and garden owners' age (relative influence: 26.34) (**Figure 5**). This model had relatively high accuracy (0.82),

and sensitivity (0.96) though only a moderate AUC (0.73), and very low specificity (0.18)

(**Electronic Supplementary Material S8**).

The proportion of those who supported pollinators was not significantly different

between home garden owners and large-scale farmers (Chi-squared = 0.856, p-value =

0.178). A significantly higher proportion of pesticide-free garden owners supported

pollinators than pesticide-using garden owners (Chi-squared = 13.159, p-value = <0.001).

Figure 5. The relative influence of factors generated from the Gradient Boosting Machine

(GBM) model for predicting whether or not garden owners support wild pollinators.

Among home garden owners, the most common activities to support wild pollinators were

to provide additional food sources (37.3%) (primarily by pollinator-friendly flowers) and

natural habitat improvement (35.7%) (e.g. wildflower strips). Providing artificial habitat

(19.9%) (e.g. bee hotels) and water (4.1%) were other forms of support. In some cases

(3.1%), growers claimed they support pollinators without providing additional

information. One respondent actively educated the neighbouring areas about the

importance of wild pollinators and how to protect them.

Discussion

In this study, we conducted an online survey in Hungary to investigate pesticide

application practices of plant growers, particularly of home garden owners and their

dependence on pesticides. Additionally, we also investigated the garden owners'

perspectives on environmental issues related to pesticides and their attitude to mitigating

pollinator declines.

Supporting our first hypothesis, we found that almost half of those who completed

our questionnaire claimed that general pesticide use is unavoidable in farming. This

proportion was even higher amongst those who actively used pesticides.

We found that expert knowledge was the best predictor of whether pesticides were

used in farming, and this was disproportionally important for large-scale farmers. Most

of the respondents who usually consult an expert, or who are experts themselves with

plant protection qualifications (e.g. plant doctor degree), use pesticides. Thus, farmers

strongly rely on (external) expert information for making decisions and embracing

alternative pest management practices this expert advice may be essential for encouraging

growers to move away from pesticide-based farming. The economic value of pollination

ecosystem services (Porto et al. 2020) and the yield losses related to pollinator decline

(Fijen et al. 2020) may be the most important points to raise in addition to emphasising

that maintaining pollinator populations requires drastic reduction or complete

abandonment of pesticide use (Geiger et al. 2010; Jacquet et al. 2022). However, farmers

who grow crops that are not dependent on insect pollination and do not face the negative

effects of their decline may be sceptical about the importance of this issue. Nonetheless,

in our study, 40% of large-scale farmers personally observed pollinator declines and over

70% of them believed that transitional agriculture can mitigate pollinator declines. These

results suggest that most growers are aware of the problem, yet their high level of

dependency on pesticides implies a distrust or lack of knowledge of alternative methods.

Also, for some crops no satisfactory management alternatives that protect yields are

available. Environmental education, subsidising ecological management (e.g. agri-

environment schemes supporting less intensive farming), and effective biodiversity offset

schemes can play an important role, especially when combined with expert advice.

However, the accessibility of this information varies from country to country (Wyckhuys

et al. 2019), and so increasing the ease with which stakeholders can access this

information is key in the transition process. Moreover, pressure from the agricultural

chemical lobby and the distrust among agricultural advisers of alternative plant protection

measures can strengthen market resistance (Villemaine et al. 2021), which can slow down

the dissemination of ecologically friendly practices.

The second-best predictor of whether respondents used pesticides or not was

gender. Despite sexes being evenly distributed among respondents, almost twice as many

men used pesticides as women. Indeed, in many respects, for instance, in eating habits

(such as food-selecting behaviour), women are more health-conscious than men (Wardle

et al. 2004), which can be reflected in differences in attitudes towards pesticide use (e.g.

Wang et al. 2017). Similar behavioural backgrounds may have created the emerging

between-gender imbalance in our study.

Besides showing patterns of general pesticide use, our survey showed that the

most important aspect for specifically choosing insecticides was the level of their effects

on humans and bees. This suggests that most users were aware that insecticides can cause

adverse, mostly sublethal, effects both in humans (Rani et al. 2021) and non-target insects

(Desneux et al. 2007; Müller 2018). This was further underpinned by the large proportion

of respondents (86.5%) who were aware that even insecticides labelled as harmless to

insect pollinators can nevertheless have negative effects. Previous experience with a

particular insecticide also influenced users' choices. Repeatedly using well-known

pesticides, however, may relax rigorous portioning habits which, in turn, may lead to

insecticide overuse (Huang et al. 2020). This fixed choice may also lead to brand fidelity,

which, consequently, may prevent experimenting with alternative, more environmentally

friendly, pesticides.

Indeed, despite scientific advice calling for the banning of all neonicotinoids

(Goulson and 232 Signatories 2018), this study showed that most respondents already

experience the effects of the present ban on neonicotinoids and that they heavily depend

on the use of acetamiprid, which is currently the only one freely available in the EU. This

may lead to a higher demand for acetamiprid-containing insecticides among plant

growers in the coming years (Varga-Szilay and Tóth 2022). Acetamiprid, like all

neonicotinoids, can persist in the tissues of treated plants (Lentola et al. 2017) and its

half-life can reach 450 days in soil (Goulson 2013) inducing sublethal effects in beneficial

organisms (Varga-Szilay and Tóth 2022), such as pollinators. On top of this, we also

found that many of those plant growers who used acetamiprid-containing pesticides co-

applied them in combination with other agrochemicals. Although concerns have been

raised about the negative effects of cocktails of pesticides on the fitness of non-target

insects (e.g. Gill et al. 2012; Williamson and Wright 2013), in our study the most extreme

example was one home garden owner who used Mospilan along with seven additional

fungicides. Based on our results, we can assume that a substantial proportion of

Hungarian growers have not yet attempted to reduce insecticides. Similarly to when

aiming to reduce pesticides at large, the publicising of relevant methodological advances

or alternative technologies is likely to be critical to achieving a reduction in insecticides

use and a transition to ecological farming.

Home gardens as ecological traps
421

Home gardens could be transformed into pesticide-free cultivation more quickly than

larger-scale farming areas, but, according to our study, Hungarian garden owners seem to

be reluctant of this conversion. Contrary to our expectations, pesticide use was

widespread among gardeners, and almost all respondents who considered that the issue

of pesticides causing harm to wild pollinators was unimportant themselves used

pesticides. Even those garden owners who acknowledged that the widespread use of

pesticides was a crucial problem for pollinators and have heard that certain pesticides

considered safe may also be harmful to wild pollinators kept using them. Hence, our

second hypothesis was not supported. Although a significantly greater proportion of home

gardeners than of large-scale farmers believed that pesticide-free farming is achievable.

Only 43.4% of the garden owners who completed our questionnaire grow plants

pesticide-free, and more than half of the garden owners who produce fruits and vegetables

for themselves and are not profit-oriented, use pesticides. These numbers are alarming

and suggest that despite the known negative effects of pesticide use and the potential

benefits of pesticide-free management, garden owners favour conventional approaches

including the use of pesticides. The proportion of pesticide-free gardeners is similar to

that found in Austria and Poland (pesticide-free: 41.0-51.7%) (Voigt et al. 2015) among

small-scale gardeners. In another survey conducted in the UK, only 30% of small

gardeners did not use pesticides (Resource Futures 2013). However, the comparability of

these results may be hampered by the differences in the definition of a 'home garden'

among surveys.

The majority of those who completed the questionnaire supported pollinators. Our

model indicated that whether or not one promoted biocontrol was the best predictor of

whether a garden owner also supported wild pollinators. However, due to the skew in

number towards pollinator-supporting garden owners, the model specificity was low,

making this prediction unreliable. A significantly higher proportion of pesticide-free

garden owners supported pollinators than of those who used pesticides, supporting our

third hypothesis. The most common means to support wild pollinators were to provide

pollinator-friendly flowers and many respondents provided bee hotels as a means of

support. These two approaches are probably widespread because in recent years

pollinators (particularly wild bees) have become an increasingly important part of

environmental education programs in the European Union (European Commission n.d.

c), including Hungary (e.g. the annual 'Pollinators day' event). Yet, Schmied et al. (2022)

demonstrated that urbanised areas, whilst being safe habitats in some cases (e.g.

Theodorou et al. 2020), can also act as ecological traps (e.g. Campioni et al. 2022;

Lehtonen et al. 2021) for insects in other cases. Indeed, although home gardens lure insect

pollinators, pesticides are used in many of those gardens, contaminating the nectar and

pollen of flowers (Nicholls et al. 2018) which, in turn, can have deleterious effects on

pollinators' fitness. Thus, these non-pesticide-free gardens act as ecological traps for

insect pollinators. For that reason, plant growers should be encouraged and motivated to

produce their vegetables and fruits pesticide-free. Garden owners should be aware that to

fully support pollinators in urban areas, pesticide use should be reduced or fully

abandoned. Realising the potential benefits of urban gardens as biodiversity refuges, and

the problems that pesticide use brings about for meeting this target, drive an increasing

number of European countries to aim to ban plant protection products in private areas in

addition to public areas (Goulson and Nicholls 2022). Home growers could also be

discouraged from buying and using these products if they are removed from being freely

available on supermarket shelves, as recently proposed in the UK (PAN UK n.d.).

Study limits and future perspectives

There are some limitations to our work. The respondents of the questionnaire were not

chosen randomly. Our population is a subsample of those who were aware of the

announcement and voluntarily took part in the study. The questionnaire could only be

completed online, therefore, it had a lower chance to reach the eastern part of the country

where there is a lower rate of internet access (KSH, n.d.). Most of the large-scale

agriculture takes place in Eastern Hungary, hence large-scale farmers may be

underrepresented. Nonetheless, our questionnaire was completed by a sufficiently large

number of people such that it should represent general plant protection habits and trends

in pesticide use among Hungarian growers.

Our study could have provided further insights if more landscape and biodiversity

variables had been available. However, asking for providing these may have been

demanding for farmers and so reduced response rates. Thus, unfortunately, we had to

compromise on this issue.

Furthermore, the generalisability of the result is limited by the fact that the survey

was conducted only among Hungarian plant growers, although it does provide a useful

case study in which results are not compounded by inter-country factors. Countries at a

similar development level and using similar agricultural practices should also be involved

in future studies to expand understanding of how pesticide use patterns and attitudes apply

across a wider geographic scale.

Conclusions and future perspectives

Additional questions in similar studies could provide deeper insight into farmers'

practices, for instance, the chemical structure of pesticides used, whether they were

synthetic or organic, or which organism were they used against.

One of the most pressing questions is whether home garden owners really want to

convert to a completely pesticide-free plant growing. Focused research is needed to

understand the willingness and motivations of garden owners for making this transition

and why (if so) they prefer to continue conventional practices. Additionally,

environmental education should establish the ecological foundations of insect-friendly

gardens and promote their local and global benefits. In particular, demonstration gardens

and demonstration farms should be set up which could demonstrate and teach pesticide-

free farming to plant growers. Alternatively, incentives and direct subsidies could be

provided to those who abandon the use of pesticides. Moreover, pesticide-free farming

could be advocated through mobile applications and social recognition, or through

granting 'pollination friendly' certification to home gardens. These gardens could also be

involved in biomonitoring programs to further strengthen links between nature and

garden owners.

However, at present, due to the unintentionally introduced pesticide pollution in

gardens (Lentola et al. 2017), not even the exact magnitude of exposure of pollinators to

chemical pesticides can be assessed. Therefore, future research should focus on this

invisible contamination and its effects on the assemblages of garden insects. From the

side of decision-makers, to deal with this issue, clear labelling practices should be

requested from suppliers to indicate whether or not products have come from pesticide-

free farms (e.g. Ecolabel Index, n.d.). The amount of freely available plant protection

products should be reduced, particularly those available in supermarkets, to discourage

direct and indirect pesticide pollution in the gardens.

Driving large-scale farmers and home garden owners toward pesticide-free

farming however may need different approaches. Whereas large scale farmers mostly rely

on expert advice, and therefore advisers should inform them about pesticide-free

practices, home gardeners may more heavily rely on conventional information channels

(such as social media and personal networks).

The attitude of Hungarian plant growers can provide a general insight into the

viewpoint of other Central and Eastern European countries and similar surveys would be

needed across Europe. Since survey approaches, similar to ours, through directed

questions about pesticide use habits, help us to better understand plant growers'

motivations, we hope this survey proves to be useful as an example for further online

questionnaires. The information gained then can help to find solutions towards a

pesticide-free future.

References

Baldock KCR (2020) Opportunities and threats for pollinator conservation in global towns and cities. Curr Opin Insect Sci 38:63–71. https://doi.org/10.1016/j.cois.2020.01.006

Baldock KCR, Goddard MA, Hicks DM, et al (2019) A systems approach reveals urban pollinator hotspots and conservation opportunities. Nat Ecol Evol 3:363–373. https://doi.org/10.1038/s41559-018-0769-y

Baldock KCR, Goddard MA, Hicks DM, et al (2015) Where is the UK's pollinator biodiversity? The importance of urban areas for flower-visiting insects. Proc R Soc B Biol Sci 282:20142849. https://doi.org/10.1098/rspb.2014.2849

Botías C, David A, Hill EM, Goulson D (2017) Quantifying exposure of wild bumblebees to mixtures of agrochemicals in agricultural and urban landscapes. Environ Pollut 222:73–82. https://doi.org/10.1016/j.envpol.2017.01.001

Botías C, David A, Horwood J, et al (2015) Neonicotinoid residues in wildflowers, a potential route of chronic exposure for bees. Environ Sci Technol 49:12731–12740. https://doi.org/10/8bk

Brühl CA, Bakanov N, Köthe S, et al (2021) Direct pesticide exposure of insects in nature conservation areas in Germany. Sci Rep 11:24144. https://doi.org/10.1038/s41598-021-03366-w

Brühl CA, Zaller JG, Liess M, Wogram J (2022) The rejection of synthetic pesticides in organic farming has multiple benefits. Trends Ecol Evol 37:113–114. https://doi.org/10.1016/j.tree.2021.11.001

Bryer J, Speerschneider K (2016) likert: Analysis and visualization likert items

Campioni L, Marengo I, Román J, D'Amico M (2022) Mud-puddling on roadsides: a potential ecological trap for butterflies. J Insect Conserv 26:131–134. https://doi.org/10.1007/s10841-021-00367-y

Dangles O, Casas J (2019) Ecosystem services provided by insects for achieving sustainable development goals. Ecosyst Serv 35:109–115. https://doi.org/10.1016/j.ecoser.2018.12.002

Desneux N, Decourtye A, Delpuech J-M (2007) The sublethal effects of pesticides on
     beneficial arthropods. Annu Rev Entomol 52:81–106.
     https://doi.org/10.1146/annurev.ento.52.110405.091440

Emmerson M, Morales MB, Oñate JJ, et al (2016) Chapter two - How agricultural
     intensification affects biodiversity and ecosystem services. In: Dumbrell AJ,
     Kordas RL, Woodward G (eds) Advances in Ecological Research. Academic
     Press, pp 43–97

EPRS (2019) Farming without plant protection products. Can we grow without using
     herbicides, fungicides and insecticides? In: Service EEPR (ed). Publications
     Office, Panel for the Future of Science and Technology, PE 634.416

European Commission (2020) Communication from the Commission to the European
     Parliament, the Council, the European Economic and Social Committee and the
     Committee of the Regions: A Farm to Fork Strategy for a fair, healthy and
     environmentally-friendly food system COM/2020/381 final. https://eur-
     lex.europa.eu/resource.html?uri=cellar:ea0f9f73-9ab2-11ea-9d2d-
     01aa75ed71a1.0001.02/DOC_1&format=PDF. Accessed 20 Jan 2022

European Commission Organic farming statistics.
     https://ec.europa.eu/eurostat/statistics-
     explained/index.php?title=Organic_farming_statistics. Accessed 18 Feb 2022a

European Commission Member States: Trends. Trend in use and risk of chemical and
     more hazardous pesticides.
     https://ec.europa.eu/food/plants/pesticides/sustainable-use-pesticides/farm-fork-
     targets-progress/member-states-trends_en. Accessed 20 Jan 2022b

European Commission EU Pollinators Initiative.
     https://ec.europa.eu/environment/nature/conservation/species/pollinators/policy_
     en.htm. Accessed 14 Feb 2022c

FAO (2021) Pesticides use, pesticides trade and pesticides indicators. Global, regional
     and country trends, 1990–2019. FAOSTAT Anal Brief Ser No 29 Rome 22

Fijen TPM, Scheper JA, Vogel C, et al (2020) Insect pollination is the weakest link in
     the production of a hybrid seed crop. Agric Ecosyst Environ 290:106743.
     https://doi.org/10.1016/j.agee.2019.106743

Garden Organic France declares public spaces pesticide free - private gardens will
follow. https://www.gardenorganic.org.uk/news/france-declares-public-spaces-
pesticide-free. Accessed 1 Mar 2022

Geiger F, Bengtsson J, Berendse F, et al (2010) Persistent negative effects of pesticides
on biodiversity and biological control potential on European farmland. Basic
Appl Ecol 11:97–105. https://doi.org/10.1016/j.baae.2009.12.001

Gill RJ, Ramos-Rodriguez O, Raine NE (2012) Combined pesticide exposure severely
affects individual- and colony-level traits in bees. Nature 491:105–108.
https://doi.org/10/f4cg7z

Goulson D (2013) REVIEW: An overview of the environmental risks posed by
neonicotinoid insecticides. J Appl Ecol 50:977–987. https://doi.org/10/mv6

Goulson D, 232 Signatories (2018) Call to restrict neonicotinoids. Science 360:973.
https://doi.org/10.1126/science.aau0432

Goulson D, Nicholls E (2022) Anthropogenic influences on bee foraging. Science
375:970–972. https://doi.org/10.1126/science.abn0185

Grass I, Loos J, Baensch S, et al (2019) Land-sharing/-sparing connectivity landscapes
for ecosystem services and biodiversity conservation. People Nat 1:262–272.
https://doi.org/10.1002/pan3.21

Greenwell B, Boehmke B, Cunningham J, Developers G (2020) gbm: Generalized
boosted regression models

Gurr GM, Wratten SD, Landis DA, You M (2017) Habitat management to suppress pest
populations: progress and prospects. Annu Rev Entomol 62:91–109.
https://doi.org/10.1146/annurev-ento-031616-035050

Hall DM, Camilo GR, Tonietto RK, et al (2017) The city as a refuge for insect
pollinators. Conserv Biol 31:24–29. https://doi.org/10.1111/cobi.12840

Hallmann CA, Sorg M, Jongejans E, et al (2017) More than 75 percent decline over 27
years in total flying insect biomass in protected areas. PLOS ONE 12:e0185809.
https://doi.org/10.1371/journal.pone.0185809

Hanley N, Breeze TD, Ellis C, Goulson D (2015) Measuring the economic value of
pollination services: Principles, evidence and knowledge gaps. Ecosyst Serv
14:124–132. https://doi.org/10.1016/j.ecoser.2014.09.013

Holzschuh A, Steffan-Dewenter I, Tscharntke T (2008) Agricultural landscapes with
organic crops support higher pollinator diversity. Oikos 117:354–361.
https://doi.org/10.1111/j.2007.0030-1299.16303.x

Huang Y, Luo X, Tang L, Yu W (2020) The power of habit: does production experience
lead to pesticide overuse? Environ Sci Pollut Res 27:25287–25296.
https://doi.org/10.1007/s11356-020-08961-4

Hülsmann M, von Wehrden H, Klein A-M, Leonhardt SD (2015) Plant diversity and
composition compensate for negative effects of urbanization on foraging bumble
bees. Apidologie 46:760–770. https://doi.org/10.1007/s13592-015-0366-x

Jacquet F, Jeuffroy M-H, Jouan J, et al (2022) Pesticide-free agriculture as a new
paradigm for research. Agron Sustain Dev 42:8. https://doi.org/10.1007/s13593-
021-00742-8

KSH Központi Statisztikai Hivatal A háztartások információs- és
kommunikációseszköz-használatának főbb jellemzői.
https://www.ksh.hu/docs/hun/xftp/idoszaki/ikt/2020/01/index.html. Accessed 5
Apr 2022

Krupke CH, Hunt GJ, Eitzer BD, et al (2012) Multiple routes of pesticide exposure for
honey bees living near agricultural fields. PLOS ONE 7:e29268.
https://doi.org/10.1371/journal.pone.0029268

Kuhn M (2021) caret: Classification and regression training

Kuhn M, Vaughan D (2021) yardstick: Tidy characterizations of model performance

Lechenet M, Dessaint F, Py G, et al (2017) Reducing pesticide use while preserving
crop productivity and profitability on arable farms. Nat Plants 3:1–6.
https://doi.org/10.1038/nplants.2017.8

Lehtonen TK, Babic NL, Piepponen T, et al (2021) High road mortality during female-
biased larval dispersal in an iconic beetle. Behav Ecol Sociobiol 75:26.
https://doi.org/10.1007/s00265-020-02962-6

Lentola A, David A, Abdul-Sada A, et al (2017) Ornamental plants on sale to the public
are a significant source of pesticide residues with implications for the health of
pollinating insects. Environ Pollut 228:297–304.
https://doi.org/10.1016/j.envpol.2017.03.084

Main AR, Hladik ML, Webb EB, et al (2020) Beyond neonicotinoids – Wild pollinators
654   are exposed to a range of pesticides while foraging in agroecosystems. Sci Total
655   Environ 742:140436. https://doi.org/10/ghp9xk

656   Müller C (2018) Impacts of sublethal insecticide exposure on insects — Facts and
657   knowledge gaps. Basic Appl Ecol 30:1–10.
658   https://doi.org/10.1016/j.baae.2018.05.001

659   Nicholls E, Botías C, Rotheray EL, et al (2018) Monitoring neonicotinoid exposure for
660   bees in rural and peri-urban areas of the U.K. during the transition from pre- to
661   post-moratorium. Environ Sci Technol 52:9391–9402. https://doi.org/10/gh6hqh

662   PAN Europe (2019) Pesticide Action Week 2019: No more pesticides in non-
663   agricultural areas! https://www.pan-europe.info/sites/pan-
664   europe.info/files/PANEurope_PR_PesticideActionWeek2019_Final.pdf.
665   Accessed 7 Mar 2022

666   PAN Europe Pesticide Free Towns. https://www.pan-europe.info/campaigns/pesticide-
667   free-towns. Accessed 7 Mar 2022

668   PAN UK Pesticide Action Network UK. https://www.pan-uk.org/. Accessed 28 Feb
669   2022

670   Pebesma E (2018) Simple features for r: Standardized support for spatial vector data. R
671   J 10:439–446. https://doi.org/10.32614/RJ-2018-009

672   Pecenka JR, Ingwell LL, Foster RE, et al (2021) IPM reduces insecticide applications
673   by 95% while maintaining or enhancing crop yields through wild pollinator
674   conservation. Proc Natl Acad Sci 118:e2108429118.
675   https://doi.org/10.1073/pnas.2108429118

676   Pisa LW, Amaral-Rogers V, Belzunces LP, et al (2015) Effects of neonicotinoids and
677   fipronil on non-target invertebrates. Environ Sci Pollut Res 22:68–102.
678   https://doi.org/10.1007/s11356-014-3471-x

679   Porto RG, de Almeida RF, Cruz-Neto O, et al (2020) Pollination ecosystem services: A
680   comprehensive review of economic values, research funding and policy actions.
681   Food Secur 12:1425–1442. https://doi.org/10.1007/s12571-020-01043-w

682   Potts SG, Imperatriz-Fonseca V, Ngo HT, et al (2016) Safeguarding pollinators and
683   their values to human well-being. Nature 540:220–229.
684   https://doi.org/10.1038/nature20588

685 Pywell RF, Heard MS, Woodcock BA, et al (2015) Wildlife-friendly farming increases
686   crop yield: evidence for ecological intensification. Proc R Soc B Biol Sci
687   282:20151740. https://doi.org/10.1098/rspb.2015.1740

688 R Core Team (2021) R: A language and environment for statistical computing. Vienna,
689   Austria

690 Raimets R, Karise R, Mänd M, et al (2018) Synergistic interactions between a variety of
691   insecticides and an ergosterol biosynthesis inhibitor fungicide in dietary
692   exposures of bumble bees (*Bombus terrestris* L.). Pest Manag Sci 74:541–546.
693   https://doi.org/10.1002/ps.4756

694 Rani L, Thapa K, Kanojia N, et al (2021) An extensive review on the consequences of
695   chemical pesticides on human health and environment. J Clean Prod
696   283:124657. https://doi.org/10.1016/j.jclepro.2020.124657

697 Resource Futures (2013) Pesticide user habits survey 2013: public purchasing, use,
698   storage, and disposal of pesticides in plant protection products

699 Revelle W (2021) psych: Procedures for psychological, psychometric, and personality
700   research. Evanston, Illinois

701 Sánchez-Bayo F, Wyckhuys KAG (2019) Worldwide decline of the entomofauna: A
702   review of its drivers. Biol Conserv 232:8–27. https://doi.org/10/gfvb6r

703 Saqib HSA, Chen J, Chen W, et al (2020) Local management and landscape structure
704   determine the assemblage patterns of spiders in vegetable fields. Sci Rep
705   10:15130. https://doi.org/10.1038/s41598-020-71888-w

706 Schmied H, Getrost L, Diestelhorst O, et al (2022) Between perfect habitat and
707   ecological trap: even wildflower strips mulched annually increase pollinating
708   insect numbers in intensively used agricultural landscapes. J Insect Conserv.
709   https://doi.org/10.1007/s10841-022-00383-6

710 Sirohi MH, Jackson J, Edwards M, Ollerton J (2015) Diversity and abundance of
711   solitary and primitively eusocial bees in an urban centre: a case study from
712   Northampton (England). J Insect Conserv 19:487–500.
713   https://doi.org/10.1007/s10841-015-9769-2

714 Siviter H, Bailes EJ, Martin CD, et al (2021) Agrochemicals interact synergistically to
715   increase bee mortality. Nature 596:389–392. https://doi.org/10.1038/s41586-
716   021-03787-7

Theodorou P, Radzevičiūtė R, Lentendu G, et al (2020) Urban areas as hotspots for bees

and pollination but not a panacea for all insects. Nat Commun 11:576.

https://doi.org/10.1038/s41467-020-14496-6

Tooker JF, Pearsons KA (2021) Newer characters, same story: neonicotinoid

insecticides disrupt food webs through direct and indirect effects. Curr Opin

Insect Sci 46:50–56. https://doi.org/10.1016/j.cois.2021.02.013

Tscharntke T, Grass I, Wanger TC, et al (2021) Beyond organic farming – harnessing

biodiversity-friendly landscapes. Trends Ecol Evol 36:919–930.

https://doi.org/10.1016/j.tree.2021.06.010

Tscharntke T, Grass I, Wanger TC, et al (2022) Restoring biodiversity needs more than

reducing pesticides. Trends Ecol Evol 37:115–116.

https://doi.org/10.1016/j.tree.2021.11.009

Vanbergen AJ, the Insect Pollinators Initiative (2013) Threats to an ecosystem service:

pressures on pollinators. Front Ecol Environ 11:251–259.

https://doi.org/10.1890/120126

Varga-Szilay Z, Tóth Z (2022) Is acetamiprid really not that harmful to bumblebees

(Apidae: *Bombus* spp.)? Apidologie 53:2. https://doi.org/10.1007/s13592-022-

00909-6

Villemaine R, Compagnone C, Falconnet C (2021) The social construction of

alternatives to pesticide use: A study of biocontrol in Burgundian viticulture.

Sociol Rural 61:74–95. https://doi.org/10.1111/soru.12320

Voigt A, Latkowska M, Rutecka A, et al (2015) Environmental behavior of urban

allotment gardeners in Europe. Landsc Flux Est Univ Life Sci Est 78–82

Wang W, Jin J, He R, Gong H (2017) Gender differences in pesticide use knowledge,

risk awareness and practices in Chinese farmers. Sci Total Environ 590–591:22–

28. https://doi.org/10.1016/j.scitotenv.2017.03.053

Wardle J, Haase AM, Steptoe A, et al (2004) Gender differences in food choice: The

contribution of health beliefs and dieting. Ann Behav Med 27:107–116.

https://doi.org/10.1207/s15324796abm2702_5

Wei T, Simko V (2021) R package "corrplot": Visualization of a correlation matrix

Wenzel A, Grass I, Belavadi VV, Tscharntke T (2020) How urbanization is driving

pollinator diversity and pollination – A systematic review. Biol Conserv

241:108321. https://doi.org/10.1016/j.biocon.2019.108321

Wernecke A, Castle D (2020) Effects of tank mixtures of plant protection products on

honey bees and possible physiological interactions. J Für Kult 154-161 Seiten.

https://doi.org/10.5073/JFK.2020.05.05

Willer H, Lernoud J (2018) The world of organic agriculture. Statistics and emerging

trends 2018. https://orgprints.org/id/eprint/34669/1/WILLER-LERNOUD-2018-

final-PDF-low.pdf. Accessed 14 Feb 2022

Williamson SM, Wright GA (2013) Exposure to multiple cholinergic pesticides impairs

olfactory learning and memory in honeybees. J Exp Biol 216:1799–1807.

https://doi.org/10.1242/jeb.083931

Wray JC, Elle E (2015) Flowering phenology and nesting resources influence pollinator

community composition in a fragmented ecosystem. Landsc Ecol 30:261–272.

https://doi.org/10.1007/s10980-014-0121-0

Wyckhuys KAG, Pozsgai G, Lovei GL, et al (2019) Global disparity in public

awareness of the biological control potential of invertebrates. Sci Total Environ

660:799–806. https://doi.org/10.1016/j.scitotenv.2019.01.077

Zaller JG (2020) Pesticide Impacts on the Environment and Humans. In: Zaller JG (ed)

Daily Poison : Pesticides - an Underestimated Danger. Springer International

Publishing, Cham, pp 127–221

Zattara EE, Aizen MA (2021) Worldwide occurrence records suggest a global decline in

bee species richness. One Earth 4:114–123.

https://doi.org/10.1016/j.oneear.2020.12.005

Ecolabel Index - All ecolabels on food.

https://www.ecolabelindex.com/ecolabels/?st=category,food. Accessed 14 Feb

2022

Acknowledgements

We are grateful to Zoltán Tóth for pointing us towards conducting a questionnaire-based study. We also thank Marco Ferrante and Nick Littlewood for their comments on the manuscript.

Author information

Affiliations

Doctoral School of Biology, Institute of Biology, ELTE Eötvös Loránd University, Budapest, Hungary

Zsófia Varga-Szilay

cE3c – Centre for Ecology, Evolution and Environmental Changes/Azorean Biodiversity Group, Departamento de Ciências e Engenharia do Ambiente, Universidade dos Açores, Açores, Portugal

Gábor Pozsgai

Corresponding author

Correspondence to Zsófia Varga-Szilay

ORCID number of authors

Zsófia Varga-Szilay: https://orcid.org/0000-0001-9712-7654

Gábor Pozsgai: https://orcid.org/0000-0002-2300-6558

Author contributions

Zsófia Varga-Szilay conceived and designed the study. The questionnaire was created and data were collected by Zsófia Varga-Szilay. Analysis was performed by Zsófia Varga-Szilay and Gábor Pozsgai. The manuscript was written by Zsófia Varga-Szilay and Gábor Pozsgai. Both authors read and approved the final manuscript.

Declarations

Conflict of interest

The authors declare no competing interests.

Code availability

The underlying computer code is available in the GitHub repository

https://github.com/zsvargaszilay/pesticide_questionnaire