Antibiotic-producing Micrococcales govern the microbiome that inhabits the fur of two- and three-toed sloths

Diego Rojas-Gätjens[1], Katherine S. Valverde-Madrigal[1], Keilor Rojas-Jimenez[2], Reinaldo Pereira[3] Judy Avey-Arroyo[4] & Max Chavarría[1,5,6*]

[1]Centro Nacional de Innovaciones Biotecnológicas (CENIBiot), CeNAT-CONARE, 1174-1200, San José, Costa Rica, [2]Escuela de Biología, Universidad de Costa Rica, 11501-2060, San José, Costa Rica; [3]Laboratorio Nacional de Nanotecnología (LANOTEC), CeNAT-CONARE, 1174-1200 San José, Costa Rica. [4]The Sloth Sanctuary of Costa Rica, Limon, Costa Rica, [5]Escuela de Química, Universidad de Costa Rica, 11501-2060, San José, Costa Rica, [6]Centro de Investigaciones en Productos Naturales (CIPRONA), Universidad de Costa Rica, 11501-2060, San José, Costa Rica.

Running title: Antibiotic-producing Micrococcales in sloth`s fur

Keywords: Sloths, Micrococcales, Antibiotic-producing bacteria, *Kocuria, Brevibacterium*, *Rothia*, Fur microbiota

* Correspondence to: Max Chavarría

Escuela de Química & Centro de Investigaciones en Productos Naturales (CIPRONA)

Universidad de Costa Rica

Sede Central, San Pedro de Montes de Oca, San José, 11501-2060, Costa Rica

Phone (+506) 2511 8520. Fax (+506) 2253 5020

E-mail: max.chavarria@ucr.ac.cr; ORCID: https://orcid.org/0000-0001-5901-3576

Summary

Sloths have a dense coat on which insects, algae, and fungi coexist in a symbiotic relationship. This

complex ecosystem requires different levels of control, however, most of these mechanisms remain

unknown. We investigated the bacterial communities inhabiting the hair of two- (*Choloepus Hoffmani*)

and three-toed (*Bradypus variegatus)* sloths and evaluated their potential for producing antibiotic

molecules capable of exerting control over the hair microbiota. The analysis of 16S rRNA amplicon

sequence variants (ASVs) revealed that the communities in both host species are dominated by

Actinobacteriota and Firmicutes. The most abundant genera were *Brevibacterium, Kocuria/Rothia,*

Staphylococcus, Rubrobacter, Nesterenkonia, and Janibacter. In addition, we isolated nine strains of

Brevibacterium and *Rothia* able to produce substances that inhibited the growth of common

mammalian pathogens. The analysis of the biosynthetic gene clusters (BCGs) of these nine isolates

suggests that the pathogen-inhibitory activity could be mediated by the presence of siderophores,

terpenes, beta-lactones, Type III polyketide synthases (T3PKS), ribosomally synthesized, and post-

translationally modified peptides (RiPPs), non-alpha poly-amino acids (NAPAA) like e-Polylysin,

ectoine or nonribosomal peptides (NRPs). Our data suggest that Micrococcales inhabiting sloth hair

could have a role in controlling microbial populations in that habitat, improving our understanding of

this highly complex ecosystem.

____

Introduction

Almost all mammals have fur or other hair-related structures that help them adapt to their habitats.

Hair usually provides thermal regulation and camouflage (Dawson et al., 2014) also it has been

related to some communication and mating processes (Buffoli et al., 2014). Beyond their functional

role for animals, hair constitutes a habitat for multiple microorganisms on which they coexist and

associate in complex ways with their host (Chen et al., 2018). In mammals, due to their anatomical

proximity, it is difficult to separate the microbiome associated with skin and fur, and the vast majority

of studies refer to the skin microbiome (Belheouane et al., 2020; Chen et al., 2018; Council et al.,

2016; Ross et al., 2019). For example, a high microbial diversity has been reported in human skin

(Byrd et al., 2018), and it has been shown that it presents a high proportion of symbionts that control

the population of pathogenic bacteria through the secretion of multiple compounds such as

bacteriocins or fatty acids (Byrd et al., 2018; O'Sullivan et al., 2019). For its part, the hair microbiota

in humans has been shown to have a similar composition to skin samples (Polak-Witka et al., 2020;

Saxena et al., 2021). The most abundant genera found in hair follicles correspond to *Cutibacterium*

acnes (previously known as *Propionibacterium acnes*), *Staphylococcus* sp., *Corynebacterium*,

Streptococcus, *Acinetobacter*, *Prevotella* and *Malassezia* (Lousada et al., 2021; Polak-Witka et al.,

2020; Watanabe et al., 2021), however, it is important to mention that this microbiota is influenced by

multiple factors including demography, use of cosmetics products or oils and genetics (Brinkac et al.,

2018).

The human skin (and fur) microbiome is the most studied; the microbiome present in the skin and

hair of other mammals and their possible functions are poorly understood or remain completely

unknown. Understanding these microbe-mammalian hair relationships is important not only from an

ecological perspective but also because we could reveal microorganisms, enzymes or small

molecules of biotechnological interest. A global analysis of multiple animals' skin microbiome, based

on 16S rRNA amplicon sequencing revealed that the skin of most mammals differs significantly from

the human skin microbiome and resembles soil microbiome in most cases, which is probably a

consequence of the constant environmental exposure of their fur. Also, a phylosymbiosis of the

microbiota was found among the different mammals' clades. Bacteria belonging to *Staphylococcus,*

Corynebacterium, and *Micrococcus* genera are reported as recurrent members on the skin and fur of

mammals (Kloss et al., 1976; Ross et al., 2019). In non-human primates, it was revealed that even

though members of the *Staphylococcus* and *Corynebacterium* genera were present, other phylotypes

such as *Anaerococcus* and *Prevotella* are an abundant part of the microbial community (Council et

al., 2016). Furthermore, on livestock animals, it has been shown that the skin microbiome has an

important prevalence of Bacteoidetes, Chloroflexi, and Proteobacteria (Ross et al., 2018; Ross et al.,

2019). The role of these phylotypes in the animal's physiology is unclear. Nevertheless, it has been

postulated they can be involved in protection from pathogenic bacteria via competition or maturation

of the immune system (Bradley et al., 2016; Rodrigues Hoffman, 2017).

Sloths are mammals whose skin and hair microbiota is partially known (Higginbotham et al., 2014;

Pauli et al., 2014). Sloths are mammals adapted to live in the canopy layer of the rainforest. These

animals are characterized as being solitary with parsimonious movements and a relatively low

metabolic rate compared to other mammals (Falconi et al., 2015; Taube et al., 2008). They have a

low body temperature, making them intolerant to sudden variations in temperature; for this reason,

they spend most of their time sunbathing and feeding (De Oliveira et al., 2014). There are a total of

six different sloth species divided into two genera: *Bradypus* (two-toed sloths), with four species, and

Choloepus (three-toed sloths), with two species (Dill-McFarland et al., 2016; Mendoza et al., 2015).

In Costa Rica, only two species are present *B. variegatus* and *C. hoffmanni* (Vaughan et al., 2007).

These two species were declared in August 2021 as a national symbol in the country due to their

importance in touristic activities and their ecological roles (see Figure 1).

Sloth fur is a dense coat on which macro- and microorganisms coexist (see Figure 1). It is usually

covered with green algae from the genus *Trichophilus,* which helps sloths camouflage from predators

(Suutari et al., 2010). Also, a moth classified as *Cryptoses* lives on sloths' hair and is in charge of the

nitrification process that supports the growth of *Trichophilus* (Waager and Montgomery, 1976).

Previously, a study that focused on culturable fungi that inhabit two-toed sloth's hair revealed that

these fungi secrete multiple compounds that inhibit the growth of numerous bacteria and parasites

such as *T. cruzi* (Higginbotham et al., 2014). Even though sloths fur has been proved to be a

complex habitat with various organisms occupying it, little information exists about its bacterial

community, functions, and possible roles as symbionts of these mammals. In this work, we

characterize the bacterial community inhabiting the hair of both sloth species from Costa Rica

(*Choloepus hoffmani* and *Bradypus variegatus*) (see Figure 1). Additionally, we intend to explore the

possible roles of these bacteria in the pathogen control of the microbial community on sloth hair.

Experimental Procedures

Sampling and permits

All samples of sloth hair were obtained from the Sloth Sanctuary (http://www.slothsanctuary.com) in

Cahuita, Limon (9°47'58.5"N 82°54'53.1"W) in February 2020. The sanctuary shelters around 250

individuals of *Bradypus variegatus* and *Choloepus hoffmani* (see Figure 1). The Sloth Sanctuary has

been operating since 1992 and is dedicated to the rescue, rehabilitation, and research of sloths and

conserving their rainforest habitat. All the sloths have come to the sanctuary because they were

abandoned by their mothers, or have suffered some kind of accident. In cases where it is possible,

the animals are reinserted into their natural habitats, however, many do not have that capacity and

are therefore kept in the refuge. All sloths are kept in cages which can be either alone or with

partners of the same species. In the case of shared cages, only one sloth per cage was sampled.

Hair samples were taken from the back of the sloth usinors and then stored in sterile 50 mL conical

tubes. We obtained two subsamples from 13 individuals of *Bradypus* and 15 individuals of

Choloepus. The names, code, years in the refuge, and other related characteristics of each sampled

animal are shown in Table S1. Samples for total DNA extraction were stored at -20 °C until

processed. The samples for bacterial isolation were immediately transported to the laboratory and

processed as described below. Permits to sample were obtained from the Institutional Commission of

Biodiversity of the University of Costa Rica (resolution Nº 253) and Judy Avey-Arroyo, the property

owner.

Scanning electron microscopy

Hairs of *B. variegatus* and *C. hoffmani* ~5 mm long were fixed with osmium tetroxide vapor (4%, in

0.2 M sodium phosphate buffer, pH 7.2). The hair fragments were placed in 2-cm-diameter plastic

lids, which were kept inside a 15-cm-diameter glass Petri dish containing filter paper moistened with

the osmium solution for 48 hours in a hood of extraction. After this time, the material was removed

from the Petri dish and left to dry in the fume hood for 12 hours. The hairs were placed on an

aluminum base containing double-sided carbon conductive tape and plated with gold in a Denton

Vaccum Desk V ion spreader (20mA, 3 minutes). Subsequently, the hairs were observed in scanning

electron microscopy (JEOL JSM 6390LV) at 20 KV.

Total DNA extraction and sequencing

For total DNA extraction of sloth hair, the samples were proceeded as described by Rojas-Gätjens

(2021) with some modifications. Briefly, hair samples were cut with sterile scissors into smaller

pieces. A total of 300-500 mg of hair per sample was extracted with a DNA isolation kit (PowerSoil®,

MoBio, Carlsbad, CA, USA) as described by the manufacturer. Cell lysis was accomplished by two

steps of bead beating (FastPrep-24, MP Biomedicals, Santa Ana, CA, USA) for 30 s at 6.0 m s^{-1}. For

the construction of the microbial amplicon library (16S rRNA), the V4 hypervariable region was PCR-

amplified with universal primers 515F (5'-GTGCCAGCMGCCGCGGTAA-3') and 806R (5'-

GGACTACHVGGGTWTCTAAT-3') (Carporaso et al., 2010). The PCR-generated products were

subjected to a 250 nt paired-end sequencing (Novoseq, Novogene).

Bioinformatic and statistical analysis

Amplicon sequencing data was analyzed using the DADA2 pipeline (version 1.12), as described by

Rojas-Gätjens et al. (2021). Briefly, reads were quality-filtered, deduplicated, and amplicon sequence

variant (ASV), a higher-resolution analog of the traditional operational taxonomic unit (OTU), was

inferred. The chimeric sequences were removed. This process resulted in an ASV table, which

records the number of times each exact amplicon sequence variant was observed in a sample. ASVs

were taxonomically assigned by comparing sequences against the SILVA reference database v138

(Quast et al., 2013) and then curated by comparing sequences against the GenBank database of

NCBI. Global singletons, doubletons, and sequences classified as Eukarya, Chloroplast, and

Mithocondria were removed. The average number of sequences per sample obtained was 132265,

ranging from 91415 to 170677. For estimation of the core microbiome of both sloths, we use the

package R package microbiome (Lahti and Shetty, 2018). The core microbiome was defined with a

prevalence of 50% and abundance of 1% according to Lahti and Shetty (2018). The raw sequencing

data were deposited in the sequence-read archive (SRA) of Genbank under the Bioproject ID

PRJNA804868

The statistical analysis and their visualization were performed with R statistical program (R-core-

Team 2019) and the Project Jupyter notebook interface. The package Vegan v2.5-6 (Oksanen et al.

2019) was used to calculate alpha diversity estimators (Observed abundance, Shannon index, and

Simpson index) and Non-metric multidimensional scaling analysis (NMDS). Wilcoxon test was

performed for comparison of the alpha diversity estimators between hosts (*Choloepus hoffmani* and

Bradypus variegatus).   Data tables with the ASV abundances were normalized into relative

abundances and then converted into a Bray–Curtis similarity matrix. The effects of the hosts and sex

on the bacterial community composition were evaluated with non-parametric multivariate analysis of

variance (PERMANOVA) and pairwise PERMANOVA (adonis2 function with 999 permutations). The

effect of the time they present in captivity was evaluated using the envfit function developed on the

package Vegan v2.5-6 (Oksanen et al. 2019).

Isolation of antibiotic-producing bacteria

Samples were processed as described by Cambronero-Heinrichs (2019). Briefly, samples were

diluted in PBS 1X (Na_2HPO_4 0.01 mol L^{-1}, KH_2PO_4 0.0018 mol L^{-1}, NaCl 0.137 mol L^{-1}, KCl 0.0027

mol L^{-1}) employing a 1:1 sample weight/volume ratio and then serially diluted until 10^{-3}. Dilutions were

pre-incubated (60°C for 10 min) for avoiding the growth of Enterobacteria. Dilutions were plated on

oatmeal agar (Jensen et al., 2005) supplemented with nalidixic acid (25 mg ml^{-1}) and

cycloheximide/clotrimazole (50 mg ml^{-1}). Plates were incubated for three months (30°C) and

examined every week. All bacteria were transferred to ISP2 medium (Pridham et al., 1957). All

isolates were preserved (−80°C, 15 % glycerol).

Screening of antimicrobial activity

The ability of isolates to inhibit bacterial growth was evaluated by the agar disc diffusion method

described by Cambronero-Heinrichs (2019). Since there are no known possible pathogens on sloth

skin, we decided to perform antibiotic activity tests against ATCC model microorganisms such:

Escherichia coli ATCC25922, *Pseudomonas aeruginosa* PAO1, *Staphylococcus aureus* ATCC25923,

Candida albicans ATCC10231, and *Bacillus subtilis subsp. spizizenii* ATCC6336. In addition, we

included in the test set other available microorganisms known to be pathogenic in mammals (i.e.

Streptococcus agalactiae, *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, *Kleibsella*

pneumonie, *Serratia marcescens)*, as well as some genera of fungi that were isolated from the hair of

sloths previously (i.e. *Cytospora sp., Fusarium sp., Colletrichum sp., Hypocrea sp., Trichoderma sp)*

(Higginbotham et al., 2014).

Bacteria and yeast pathogens strains were individually inoculated in either Luria–Bertani agar or

potato dextrose agar (OD = 0.1, from overnight cultures). A piece of agar (diameter 6mm) with

confluent growth of each actinobacterial isolate (2-week-old cultures in ISP2) was cut and placed on

the surface of the medium, with bacterial growth uppermost. A Whatman® cellulose filter paper

inoculated with kanamycin (5 uL, 5 mg/mL; 50 mg/mL for *P. aeruginosa*) or clotrimazole (5 mg/mL)

was used as a positive control. A filter without any substance was used as the negative control. The

plates were incubated for 24h at 37 °C for *E. coli, P. aeruginosa, S. aureus, Streptococcus*

agalactiae, Staphylococcus epidermidis, Staphylococcus saprophyticus, Kleibsella pneumonie,

Serratia marcescens, and *C. albicans,* and 30° C for all others. Every experiment was performed in

duplicate. Inhibition halos were measured in centimeters, and the results were ranked by size (see

Table 1).

Identification of bacterial isolates

For identification of the antibiotic-producing isolates, each one was grown individually (ISP2 agar, 1-4

weeks). Then, DNA was extracted according to the protocol described by Chun and Goodfellow

(1995). DNA samples were quantified using a NanoDrop 2000 spectrophotometer (Thermo Fisher,

USA). Polymerase chain reaction (PCR) protocols described by Cambronero-Heinrichs et al. (2019)

to amplify 16S rRNA with primers 27F and 1492R were used. PCR products were visualized on

agarose gels (1%) stained with Gel Red (Biotium Inc., USA). The PCR products were sequenced

using BigDye Terminator v3.1 cycle sequencing kit and cleaned with BigDye XTerminator purification

kit (Applied Biosystems, USA). Products were analyzed on a 3130xl Genetic Analyzer (Applied

Biosystems, USA). Forward and reverse sequences were assembled using Bioedit. The obtained

consensus sequences were compared against 16S rRNA bacterial/archaeal database of the NCBI

using BlastN. All sequences were deposited on the Genbank under accessions OM638436-

OM638444.

Biosynthetic gene clusters (BGCs) analyses of antibiotic-producing isolates

For biosynthetic gene clusters (BCGs) analyses, the nine bioactive isolates were grown on ISP2

media (5 days at 30 ºC). Specimens classified within the same genera were combined in sterile

microcentrifuge tubes with a difference in OD600 of 1:2 to achieve a good binning of the genomes

according to their abundance. DNA was extracted using a DNA isolation kit (PowerSoil®, MoBio,

Carlsbad, CA, USA) following the manufacturer's instructions. DNA was quality checked and

subjected to a 250 nt paired-end sequencing (Novoseq, Novogene). The raw sequences from this

study were deposited in the GenBank database BioProject ID PRJNA804868

For the construction of the Metagenome Assembled Genomes (MAGs) sequences were quality

filtered using bbmaptools (function bbduck) and then quality checked with FastQC. Sequences were

assembled with MEGAHIT (Li et al., 2016). Only contigs larger than 1500 bp were further analyzed.

The open reading frames were mapped using Prodigal v2.6.3 (Hyatt et al., 2010) and the single-copy

core genes were identified with HMMER v3.3 (Johnson et al., 2010). The taxonomy of the SCG

genes was estimated using DIAMOND (Buchfink et al., 2015). Manual binning was performed using

Anvio v6.2. A bin was considered a MAG if it presented 100% completeness and >5% redundancy.

Four bins were obtained from which two were classified as MAGs. Phylogenetic analysis of the four

bins with closely related genomes (see table 2 for Genbank accessions) was performed by using 71

single-copy core genes (a curated list of genes available in Anvi'o phylogenomic workflow

(https://merenlab.org/tutorials/infant-gut/#chapter-iii-phylogenomics) shared among the different

genomes. Sequences were concatenated and further aligned using MUSCLE v3.8.1551 (Edgar,

2004). Phylogenetic relationship was estimated using FasTree v2.1 (Price et al., 2009).

BGC analyses were conducted employing antiSMASH 5.0 (Blin et al., 2019) with default relaxed

detection parameters that identify well-defined complete clusters and partial clusters missing one or

more functional parts. We applied the following criteria for determining a known BGC: 100% of its

genes have a significant BLAST hit to the corresponding genes of a known compound in the MIBiG

database.

Results and Discussion

SEM reveals a high density of cocci-and bacilli-shaped microorganisms in sloths hair.

Similar to the results obtained by Wujek and Cocuzza (1986), we observed through a scanning electron microscope (SEM) that the hair of sloths of the genus *Choloepus* has a series of longitudinal furrows and an average width of 138 ± 13 μm. On the other hand, the hair of sloths of the genus *Bradypus* is smoother, with irregular fissures throughout the entire hair and an average width of 261 ± 36 μm. Hair samples from both *Choloepus* and *Bradypus* (Fig. 2), showed a high density of cocci- and bacilli-shaped microorganisms. This high density of microorganisms was more noticeable in the hair of the sloths of the *Choloepus* species, which could be explained by the morphology. As observed in the SEM micrographs, the microbial communities are preferentially established within the furrows of the hair of the genus *Choloepus*. Presumably, this location can be favored by a lower exposure to mechanical stress, and greater availability of water and nutrients with respect to the ridges of the hair. Most of the microorganisms observed in those habitats correspond to coccoid bacteria, nevertheless, some hyphae-like structures and rod-shaped bacteria were present in less abundance. In *Bradypus* hair, a lower microbial density was observed and its location was not necessarily associated with the fissures throughout the entire hair. In *Bradypus*, it is also possible to observe coccoid bacteria, as on the hair of *Choloepus*.

Since the sloths live in the refuge, the development of the characteristic greenish color in their fur was not observed, typically associated with the presence of the green algae *Trichophilus*. In the SEM

photos, we were not able to determine the presence of cells of this algae either. Its absence is not

rare, since the appearance of this algae occurs in wild sloths as a camouflage mechanism, which is

not required in individuals living in captivity. Despite the absence of green algae *Trichophilus*, these

first observations show that sloth hair is an environment rich in microorganisms. To reveal the identity

of these microbial communities we decided to perform an analysis of 16S rRNA gene amplicons.

Analysis of 16S rRNA gene amplicons indicates that sloth hair is governed by members of the

Micrococcales order and shaped by the environment

The bacterial community inhabiting the hair of both sloth genera showed high diversity values. The

average estimations of the Shannon-index were 3.06 and 2.72 for *C. hoffmani* and *B. variegatus*,

respectively (see Figure S1). Wilcoxon test revealed a statistically significant difference (Shannon-

index, $p = 0.045$) in the diversity of the microbial communities between the two sloths, in which

Choloepus showed higher diversity than *Bradypus*; although multiple phylotypes were identified in all

samples. In total, 4892 amplicon sequence variants (ASV) were identified in the 28 samples. The

most abundant bacterial phyla were Actinobacteriota (10.82-96.44%), Firmicutes (1.60-65.56%),

Proteobacteria (0.53-36.75%), and Bacteroidota (0.53-32.81%). Other phyla were detected in lower

proportions, such as Acidobacteriota (0-1.91%), Chloroflexi (0.59-2.87%), Cyanobacteria (0-0.59%),

Desulfobacteria (0-0.62%), Gemmatimonadota (0.84-3.46%) (see Figure 3A and Table S2).

At the order level, we observed that the bacterial community inhabiting the hair of both species of

sloths was mainly dominated by members of the Micrococcales (see Figure 3B). Within this order, the

families Brevibacteriaceae (0.70-75.30%) and Micrococcaceae (2.61-71.23%) were the most

abundant (see Figure S2). Members of the Brevibacteriaceae family have been previously isolated

from a variety of environments including dairy products (Forquin et al., 2009), poultry skin (Forquin-

Gomez et al., 2014), insects (Katı et al., 2010), mural paintings (Heyrman et al., 2004), clinical

samples (Roux and Raoult, 2009; Talento et al., 2009), saline sediments (Chen et al., 2016),

mammal skin (Collins et al., 1983) and human microbiota (Forquin-Gomez et al., 2014). Most of the

organisms found in this family are characterized as being halotolerant with the ability to produce

sulfur volatile compounds (Arfi et al., 2003). Members of the Micrococcaceae family are usually

associated with the skin microbiota of mammals (Collins et al., 2000), blood cultures (Hansen, 1985),

and in various soil samples including marine sediments (Kim et al., 2004). These microorganisms

have demonstrated a vigorous metabolism and an exceptional ability to adapt to harsh conditions

(Dastager et al., 2014). Additionally, even though some members of both families ie.

Brevibacteriaceae and Micrococcaceae have been associated with bacteremia and other infections

(Asai et al., 2019; Bal et al., 2015; Kandi et al., 2016; Ramanan et al., 2014), most of the

microorganisms belonging to this group are commensals frequently found on the skin and oral

cavities of mammals (Collins et al., 1983; Dastager et al., 2014).

Furthermore, a significant proportion of members of the Staphylococcales (1.08-45.39%) and

Rubrobacterales (0.17-44.57%) were found in the hair samples of *Bradypus* and *Choloepus*

specimens (Figure 4B). Members of the family Staphylococcaceae are commonly found in the skin of

multiple mammals (Lory et al., 2014). Particularly *Staphylococcus aureus* is a widely-known pathogen

associated with multiple infections and has become more deadly due to the increase of antibiotic

resistance (Hassoun et al., 2017). The prevalence of members of the Staphylococcaceae family in

skin samples has been previously reported in multiple mammals, particularly in those mammals that

have a strong anthropogenic influence (Ross et al., 2018) which is the case of the sloths sampled,

that dwell in a human-controlled refugee. Contrarily to Staphylococcaceae, Rubrobacteriaceae

members are characterized for being extremophilic bacteria, which have shown to survive extreme

radiation, including gamma radiation (Ferreira et al., 1999). Members of this family are pink-

pigmented bacteria commonly isolated from hot springs (Carreto et al., 1996), biodeteriorated

monuments (Jurado et al., 2012), marine sponges (Kämpfer et al., 2014) and deep-sea sediments

(Albuquerque et al., 2014). The presence of this taxon in sloth hair is probably associated with the

high temperatures (the Sloth Sanctuary is located in Cahuita, with an annual temperature range of 21

to 31 ºC) and the constant exposure to UV radiation that the sloths endure in their niche (Duan et al.,

2017; Gohli et al., 2019; Sun et al., 2018). Nevertheless, further study is required to comprehend the

association between these extremophilic microorganisms and their hosts.

As seen in Figure 3C, abundant ASVs detected in sloth's hair encompass members of the genera

Brevibacterium (0.70-75.30%), *Kocuria* (0.94-47.17%), *Staphylococcus* (1.01-45.09%),

Brachybacterium (0.60-31.07%), *Nesterenkonia* (0.04-69.61%) and *Rubrobacter* (0.17-44.57%).

These genera are characterized for inhabiting sediments usually with a high salt concentration

(Chander et al., 2017; Dastager et al., 2014; Duan et al., 2017; Forquin-Gomez et al., 2014;

Stackebrandt, 2014). The Sloth Sanctuary in Cahuita is located just 1.7 km from the Atlantic coast of

Costa Rica, where high concentrations of salt in the environment are expected to occur. Therefore, it

is not surprising that the microorganisms that inhabit the hair of the sloths in this refuge have a high

content of bacteria associated with habitats with high concentrations of salt, thus being a factor that

shapes the microbial community. It has also been shown that compounds secreted by the glands of

the host can shape the microbial community through the variation of pH, salinity, and osmotic

pressure (Byrd et al., 2018; Grice and Segre, 2011; Sanford and Gallo, 2013). The presence of

microorganisms associated with saline sediments on our hair samples agrees with previous studies

that showed that mammals' skin microbiome resembles the community from the environment,

including soils and sediments (Ross et al., 2018).

Specifically, sequences classified as *Brevibacterium* ASV1 and ASV6 were abundant ASVs

particularly in *Bradypus* samples (see Figure 3C). *Brevibacterium* is a highly heterogeneous genus

with multiple metabolic capabilities (Forquin-Gomez et al., 2014). It has been shown that this genus

usually prefers to metabolize lactate as a source of carbon; lactate has been reported as a secreted

compound by sebaceous mammals' glands (Downey and Kealey, 2004). Additionally, this genus can

also metabolize amino acids as carbon and nitrogen sources, probably allowing them to adapt well to

the highly competitive hair niche on which algae, insects, protozoa, and other bacteria compete for

resources (Mounier et al., 2007). Similarly, multiple sequences identified as *Kocuria* were found in

samples belonging to both genera (*Choloepus* and *Bradypus*) (ASV3, ASV12, ASV16, and ASV17).

Kocuria is characterized for mainly being aerobic chemoheterotroph microorganisms (Dastager et al.,

2014). It usually expresses polymer degrading enzymes such as esterases (Kaur et al., 2011), and

lipases (Reddy et al., 2003), suggesting that they might have the ability to degrade biopolymers found

in the skin. A keratinase has also been found in *Kocuria* specimens (Bernald et al., 2006), which

probably allow them to degrade the hair in which they inhabit to obtain energy.

Additionally, various abundant ASVs were classified as *Rubrobacter*. As mentioned above, the

genus *Rubrobacter* is widely known for its ability to survive radiation and desiccation (Chen et al.,

2004). Its resistance to these conditions probably allows members of this taxon to live on the sloth

hair which is constantly exposed to UV radiation and desiccation due to the high sodium chloride

content. Additionally, ASV classified as *Staphylococcus* were highly abundant, similar to what has

been reported in human skin, where *Staphylococcus* and *Propionibacterium* are the main members

of the community (Byrd et al., 2018; Cogen et al., 2008). It has been previously shown that

Staphyloccocus is part of the normal hair microbiota in humans (Lousada et al., 2021). Even though

some species such as *S. aureus* are important pathogens, most of the genotypes belonging to this

group are commensal microorganisms that accomplish important processes for maturation of the

immune system or secreted bacteriocins-like proteins that inhibit the proliferation of pathogenic

bacteria (Nguyen et al., 2017; Newstead et al., 2020; Carson et al., 2017).

Finally, other sequence variants were assigned to the Actinobacteriota genera *Brachybacterium*

(ASV5, ASV10, ASV13) and *Nesterenkonia* (ASV4, ASV9, ASV32, ASV29, ASV15). Both of these

genera are poorly studied bacteria which, similar to *Brevibacterium* and *Kocuria*, show a great

diversity in their niches, which range from plant growth promoters to human opportunistic pathogens

(Edouard et al., 2014; Tak et al., 2018; Singh et al., 2016).

The sampled two- and three-toed sloths share almost 50% of the core microbiome in their hair

Further Non-Metric Multidimensional and PERMANOVA analysis revealed that even though similar

bacterial phyla were found in *Choloepus* and *Bradypus* hair, a significant difference between the

bacterial communities was found (PERMANOVA, p = 0.001) (see Figure 4A). No significant

difference was found between the sex (p = 0.18) or time in captivity (p = 0.143) for either species

(data not shown). Previous analyses of the skin microbiome of multiple mammals found a

specialization along mammal orders (Ross et al., 2018). It is reasonable to think that this

specialization may also occur between sloth species. The difference in the microbial community

observed could be associated with the physiology of each sloth as the compounds secreted by

multiple glands in the skin would shape the microbial community.

Even though there is a significant difference in the microbial community between sloth species (see

Figure 6A), our data revealed that the core microbiome of both sloths is similar (see Figure 4B). The

core microbiome (according to the definition of Lahti and Shati (2017)) of both sloths consists mainly

of members of the Actinobacteriota phylum and bacteria from the *Staphylococcus* genus, particularly

of ASVs classified as *Brevibacterium*, *Staphylococcus*, *Kocuria*, *Nesterenkonia*, *Brachybacterium*,

Rubrobacter, *Janibacter*, *Dietzia*, and *Saccharopolyspora*. The core microbiome of both sloth species

in the Sanctuary shares almost 50% of the ASVs (see Figure 6B) which could suggest that these

microorganisms exert a functional role in this habitat. Although the sampled sloths are in separate

cages, not mixed between species, and the Sanctuary employees avoid handling the animals, the

presence of a core microbiome that is shared between the two species must be validated with

samples from other regions and with free-living animals. Despite this, our work provides a base and

gives the first light of the microbiota that establishes itself in the fur of the sloths.

In an attempt to elucidate the role of the members of the microbiome in both sloth species and

considering that many of these microorganisms belong to the phylum Actinobacteriota, we decided to

evaluate the presence of bacteria that produce antibiotic molecules. Members of the phylum

Actinobacteriota are known for their ability to produce antibiotic molecules, and many of them are

found in association with higher organisms establishing symbiotic relationships (Matarrita-Carranza et

al., 2021).

Several members of the core microbiome are antibiotic-producing bacteria

To evaluate the capacity of the bacteria inhabiting sloth's hair to produce antibiotic compounds, we

isolated Actinobacteriota and evaluated their anti-microbial activity against multiple pathogenic

microorganisms (see Experimental procedures). We were able to isolate 54 bacteria of which nine

were antibiotic-producing. From these, five were isolated from *Choloepus* hair and four from

Bradypus hair.

The antibiotic-producing bacteria were classified as either *Brevibacterium sediminis*, *Rothia*

koreensis, or *Rothia kristinae*. Interestingly these genera were classified as part of the core

microbiome in the microbial community analysis. *Rothia kristinae* and *Rothia koreensis* were only

recently classified within the *Rothia* genus through phylogenomic analysis (Nouioui et al., 2018),

before 2018 these bacteria were classified within the *Kocuria* genus according to its 16S rRNA

sequence (Stackebrandt et al., 1995) suggesting that the absence of *Rothia* genus is a consequence

of the close phylogenic relationship between *Rothia* and *Kocuria* that do not allow a clear separation

through methodologies. Blastn of abundant ASV classified as Kocuria showed a 100% identity with

Rothia kristinae or *Rothia koreensis.*

Biological activity was observed against *S. aureus, E. coli, P. aeruginosa, B. Subtilis. S. agalatiae*

and *Cytospora* sp., however, the antimicrobial activity presented a wide variation even among strains

of the same genus (see Table 1). Strains that showed activity against *S. aureus* (*R. koreensis* 10C1,

R. kristanae 13C8, 9C1, 11C2, 3B2, and 3B6) showed no activity against *S. epidermidis* which is

known to be a common symbiont in mammal's skin (Liu et al., 2020). The ability of these isolates to

inhibit well-known pathogens growing but not affect closely related symbionts may indicate that these

Micrococcales may act as a protective mechanism principally against pathogens on sloth skin.

Nevertheless, no inhibition was observed against other well-known pathogens such as *K. pneumonie,*

S. agalactiae, and *S. saprohyticus.* Notably, *S. agalactiae* presented growth inhibition with isolate

3B2 an inhibition on its capacity to perform hemolysis when exposed to 9C1, 11C2, 3B2, and 13C9

(data not shown). Additionally, multiple strains presented activity against *Bacillus subtillis* which have

been associated with food poisoning (13C8, 9C1, 11C2, 3B2, and 3B6).

Since many fungi have demonstrated the ability to infect the skin, we attempt to evaluate the

antifungal activity of our isolates against fungi closely related to the ones previously found in

Bradypus fur (Higginbotham et al., 2014) (see Experimental procedures).   Three isolates

(11C2,13C9, and 13C8) showed antibiosis against *Cytospora* sp. Despite *Cytospora* sp. is not a

recognized mammal pathogen, its presence in sloth hair may indicate that these Microccocales have

evolved to compete with other microorganisms that occupy a similar niche.

Furthermore, multiple isolates showed the ability to inhibit other Gram-negative bacteria such as

Pseudomonas aeruginosa and *Escherichia coli* (see Table 1). *P. aeruginosa* is an opportunistic

pathogen that has shown an intrinsic resistance to a variety of antibiotics (Gellatly et al., 2013; Qiu et

al., 2009; Thi et al., 2020). Nowadays, *P. aeruginosa* strains resistant to almost all antibiotics (Molina-

Mora et al., 2020) have been isolated and represent a significant health problem; therefore, isolates

that show the ability to inhibit its growth are interesting for future research to identify the compounds

responsible for this activity.

BGCs found in antibiotic-producing bacteria are poorly studied but well conserved among strains of

Micrococcales.

When analyzing the Biosynthetic Gene Clusters (BGCs) in the assembled genomes, we were able to

manually bin the sequences into four bins (see Table 2). Bins were classified as *Rothia koreensis*

(Bin 3), *Rothia kristinae* (Bin 1 and Bin 2), and *Brevibacterium* (Bin 4). Furthermore, a comparison

with closely related genomes obtained from the Genbank (see Figure 5) showed that our Bins have a

very similar BGC composition in contrast with reference genomes of the same genus (see Figure 5)

revealing that the BGCs are well conserved among multiple strains of the same genus.

Bin 4 is the MAG with more BGCS with a total of eight BGCS classified as siderophores, terpenes,

T3PKS, RiPP-like, NAPAA, ectoine, and NRPS (Figure 5). However, none of the BGCs found in this

genome showed a 100% similarity with any known cluster on the MiBiG database therefore, we

cannot ensure that any specific metabolite is produced (Table 3). Nevertheless, BGC 22.1 was

classified as a RIPP-like (see Table 2) and present a high similarity with Linocin M18 (81%),

suggesting that a similar bacteriocin may be produced. Linoncin M18 is a bacteriocin produced by

Brevibacterium linens; this peptide has been shown to be active against Gram-positive bacteria,

particularly *Bacillus subtilis, Staphylococcus aureus,* and *Listeria monocytogenes* (Valdés-Stauber et

al., 1994). Furthermore, other bacteriocins have been shown to be responsible for the antibiotic

activity of different *Brevibacterium* strains; specifically Linencin A (Forquin-Gomez et al., 2014) and

Linenscin OC2 (Maisnier-Patin et al., 1995). Linencin A is a 95 KDa peptide that inhibits the growth of

only other *B. linus* strains. On the other hand, Linenscin OC2 showed to inhibit the growth of

Arthrobacter, Microccus, and *Corynebacterium.*

Furthermore, Bin 1 and Bin 2 classified as *Rothia kristinae* showed the presence of NRPS (three

BGCs), terpenes (two BGCs), beta lactone (two BGCs), and a T3PKS. Finally, Bin 3 showed six

BGCs classified as terpene (one BGCs), beta lactone (one BGCs), NRPS-like (one BGCs), NRPS

(one BGCs), T3PKS (one BGCs), and a siderophore (one BGCs). No BGCs found in the MAGs

present a 100% similarity with the ones reported on the MIBIG database, therefore we can not

ensure completely that any cluster produces a known secondary metabolite molecule. The low

identity of the clusters found in these bins may indicate that novel metabolites are being produced,

ensuring the potential of these microorganisms for the bioprospecting of novel antibiotics. Members

of the *Rothia* genus have been poorly described as antibiotic-producing bacteria with only a few

reports found (Uranga et al., 2020). Particularly, a porcine commensal classified as *Rothia*

nasimurium showed the genetic capacity to produce the ionophore antibiotic valinomycin (Roiser et al

2017), but no identification of the compound was performed.

In summary, this work characterizes the bacterial community inhabiting the hair of *Choloepus*

hoffmani and *Bradypus variegatus* in a group of animals in captivity. Analysis of the V4 region of the

16S rRNA gene showed that Actinobacteriota, Proteobacteria, and Firmicutes were the most

dominant phyla, and *Brevibacterium*, *Janibacter*, *Rubrobacter*, *Brachybacterium*, *Kocuria*, and

Staphylococcus the most dominant genera across all samples. Additionally, further analysis revealed

that even though a significant difference exists between the community inhabiting both sloths' sloths'

species, the phylotypes that conform to the core microbiome were mostly shared. Isolation of bacteria

revealed that phylotypes closely related to the core microbiome exert an antibiotic activity against a

wide variety of pathogens. Furthermore, metagenomics made from the antibiotic-producing strains

revealed that multiple BGCs were found in these isolates including siderophores, terpenes,

betalactone, Type III polyketide synthases (T3PKS), ribosomally synthesized and post-translationally

modified peptides (RiPPs) similar to bacteriocin Linocin M18, non-alpha poly-amino acids (NAPAA)

like e-Polylysin, ectoine and nonribosomal peptides (NRPs).

Our results suggest the presence of antibiotic-producing bacteria in the fur of sloths that could

modulate the microbial community in that ecosystem. This modulation makes it possible to control the

proliferation of potentially pathogenic bacteria for the host or inhibit other competitors (e.g. fungi) in

the niche. On the other hand, our results point to sloth fur as a new source of antibiotic-producing

bacteria that may open the opportunity for the discovery of new bioactive molecules.

Funding

This work was supported by The Vice-rectory of Research of Universidad de Costa Rica (project

number VI 809-C1-009) and the National Center of Biotechnological Innovations (CENIBiot).

Acknowledgements

We thank Sofía Vieto, Roberto Avendaño and Felipe Vasquez-Castro for their help in collecting

samples or processing them in the laboratory.

Conflicts of Interest

The authors declare no conflict of interest.

Author Contribution

MC and DRG conceived and designed the experiments; DRG, KVM, RP performed the experiments;

DRG, MC and KRJ analyzed the data; MC, JA-A and RP contributed reagents or materials; DRG and

MC wrote the paper. All authors reviewed and approved the final version of the manuscript.

References

Albuquerque, L., Johnson, M.M., Schumann, P., Rainey, F.A., & da Costa, M.S. (2014). Description

of two new thermophilic species of the genus *Rubrobacter*, *Rubrobacter calidifluminis* sp. nov. and

Rubrobacter naiadicus sp. nov., and emended description of the genus *Rubrobacter* and the species

Rubrobacter bracarensis. *Syst Appl Microbiol* **37**: 235–243.

Amiri H., Azarbaijani R., Parsa Yeganeh, L., Shahzadeh Fazeli, A., Tabatabaei, M., Salekdeh, G.H.,

and Karimi, K. (2016) *Nesterenkonia* sp. strain F, a halophilic bacterium producing acetone, butanol,

and ethanol under aerobic conditions. *Sci Rep*, **6**: e18408.

Arfi, K., Amárita, F., Spinnler, H. E., and Bonnarme, P. (2003) Catabolism of volatile sulfur

compounds precursors by *Brevibacterium linens* and *Geotrichum candidum*, two microorganisms of

the cheese ecosystem. *J Biotech* **105**: 245–253.

Asai, N., Suematsu, H., Yamada, A., Watanabe, H., Nishiyama, N., Sakanashi, D., Kato, H., Shiota,

A., Hagihara, M., Koizumi, Y., Yamagishi, Y., and Mikamo, H. (2019) *Brevibacterium paucivorans*

bacteremia: case report and review of the literature. *BMC Infect Dis* **19**: e344.

Bal, Z.S., Sen, S., Karapinar, D.Y., Aydemir, S., and Vardar, F. (2015) The first reported catheter-

related *Brevibacterium casei* bloodstream infection in a child with acute leukemia and review of the

literature. *Braz J Infect Dis* **19**: 213–215.

Belheouane, M., Vallier, M., Čepić, Chung, C.J., Ibrahim, S., and Baines, J.F. (2020) Assessing

similarities and disparities in the skin microbiota between wild and laboratory populations of house

mice. *ISME J* **14**: 2367–2380.

Bernal, C., Cairó, J., Coello, N. (2006) Purification and characterization of a novel exocellular

keratinase from Kocuria rosea. *Enzyme Microb Technol* **38**: 49-54.

Blin, K., Shaw, S., Steinke, K., Villebro, R., Ziemert, N., Lee, S. Y., Medema, M. H., and Weber, T.

(2019) antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. *Nucleic Acids*

Res **47**: 81–87.

Bradley, C.W., Morris, D.O., Rankin, S.C., Cain, C.L., Misic, A.M., Houser, T., Mauldin, E.A., and

Grice, E.A. (2016) Longitudinal evaluation of the skin microbiome and association with

microenvironment and treatment in canine atopic dermatitis. *J Invest Dermatol* **136**: 1182–1190.

Brinkac, L., Clarke, T.H., Singh, H., Greco, C., Gomez, A., Torralba, M.G., Frank, B., and Nelson,

K.E. (2018) Spatial and Environmental Variation of the Human Hair Microbiota. *Sci Rep* **8**: e9017.

Buchfink, B., Xie, C., and Huson, D.H. (2015) Fast and sensitive protein alignment using DIAMOND.

Nat Methods **12**: 59–60.

Buffoli, B., Rinaldi, F., Labanca, M., Sorbellini, E., Trink, A., Guanziroli, E., Rezzani, R., and Rodella,

L.F. (2014) The human hair: from anatomy to physiology. *Int J Dermatol* **53**: 331–341.

Byrd, A.L., Belkaid, Y., and Segre, J.A. (2018) The human skin microbiome. *Nat Rev Microbiol* **16**:

143–155.

Cambronero-Heinrichs, J.C., Matarrita-Carranza, B., Murillo-Cruz, C., Araya-Valverde, E., Chavarría,

M., and Pinto-Tomás, A.A. (2019) Phylogenetic analyses of antibiotic-producing *Streptomyces* sp.

isolates obtained from the stingless-bee *Tetragonisca angustula* (Apidae: Meliponini). *Microbiology*

165: 292–301.

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J.,

Fierer, N., Knight, R. (2010) Global patterns of 16S rRNA diversity at a depth of millions of sequences

per sample. *Proc Natl Acad Sci USA* **108**: 4516-4522.

Carreto, L., Moore, E., Nobre, M.F., Wait, R., Riley P.W., Sharp, R.J. and Da Costa M.S. (1996)

Rubrobacter xylanophilus sp. nov., a new thermophilic species isolated from a thermally polluted

effluent. *Int J Syst Evol Microbiol* **46**: 460-465.

Carson, D.A., Barkema, H.W., Naushad, S., and De Buck, J. (2017) Bacteriocins of non-aureus

Staphylococci isolated from bovine milk. *Appl Environ Microbiol* **8**: e01015-17.

Chander, A.M., Nair, R.G., Kaur, G., Kochhar, R., Dhawan, D.K., Bhadada, S.K., and Mayilraj, S.

(2017) Genome insight and comparative pathogenomic analysis of *Nesterenkonia jeotgali* strain

CD08_7 isolated from duodenal mucosa of celiac disease patient. *Front Microbiol* **8**: 129.

Chen, M.Y., Wu, S.H., Lin, G.H., Lu, C.P., Lin, Y.T., Chang, W.C., and Tsay, S.S. (2004) *Rubrobacter*

taiwanensis sp. nov., a novel thermophilic, radiation-resistant species isolated from hot springs. *Int J*

Syst Evol Microbiol **54**: 1849–1855.

Chen, P., Zhang, L., Wang, J., Ruan, J., Han, X., and Huang, Y. (2016) *Brevibacterium sediminis* sp.

nov., isolated from deep-sea sediments from the Carlsberg and Southwest Indian Ridges. *Int J Syst*

Evol Microbiol **66**: 5268–5274.

Chen, Y.E., Fischbach, M.A., and Belkaid, Y. (2018) Skin microbiota-host interactions. *Nature* **553**:

427–436.

Chun, J., Goodfellow, M. (1995) A phylogenetic analysis of the genus *Nocardia* with 16S rRNA gene

sequences. *Int J Syst Evol Microbiol* **45**: 240–245.

Cogen, A.L., Nizet, V., and Gallo, R.L. (2008) Skin microbiota: a source of disease or defense? *Brit J*

Dermatol **158**: 442–455.

Collins, M.D., Farrow, J.A., Goodfellow, M., and Minnikin, D.E. (1983) *Brevibacterium casei* sp. nov.

and *Brevibacterium epidermidis* sp. nov. *Syst Appl Microbiol* **4**: 388–395.

Collins, M.D., Hutson, R.A., Båverud, V., and Falsen, E. (2000). Characterization of a *Rothia*-like

organism from a mouse: description of *Rothia nasimurium* sp. nov. and reclassification of

Stomatococcus mucilaginosus as *Rothia mucilaginosa* comb. nov. *Int J Syst Evol Microbiol* **50**: 1247–

1251.

Council, S.E., Savage, A.M., Urban, J.M., Ehlers, M.E., Skene, J.H., Platt, M.L., Dunn, R.R., and

Horvath, J.E. (2016) Diversity and evolution of the primate skin microbiome. *Proc Biol Sci* **283**: e

20152586.

Dastager, S.G., Krishnamurthi, S., Rameshkumar, N., Dharne, M. (2014). The family

Micrococcaceae. In: Rosenberg, E., DeLong, E.F., Lory, S., Stackebrandt, E., Thompson, F. (eds)

The Prokaryotes. Springer, Berlin, Heidelberg.

Dawson, T.J., Webster, K.N., and Maloney, S.K. (2014) The fur of mammals in exposed

environments; do crypsis and thermal needs necessarily conflict? The polar bear and marsupial koala

compared. *J Comp Physiol B* **184**: 273–284.

De Oliveira Moreira, D., Leite, G.R., de Siqueira, M.F., Coutinho, B.R., Zanon, M.S., and Mendes,

S.L. (2014) The distributional ecology of the maned sloth: environmental influences on its distribution

and gaps in knowledge. PLoS ONE **9**: e110929.

Dill-McFarland, K.A., Weimer, P.J., Pauli, J.N., Peery, M.Z., and Suen, G. (2016) Diet specialization

selects for an unusual and simplified gut microbiota in two-and three-toed sloths. *Environ Microbiol*

18: 1391-1402.

Downie, M.M., and Kealey, T. (2004) Human sebaceous glands engage in aerobic glycolysis and

glutaminolysis. *Brit J Dermatol* **151**: 320–327.

Duan, Y., Wu, F., Wang, W., He, D., Gu, J.D., Feng, H., et al. (2017) The microbial community

characteristics of ancient painted sculptures in Maijishan Grottoes, China. PLoS ONE **12**: e0183598.

Edgar R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput.

Nucleic Acids Res **32**: 1792–1797

Edouard, S., Sankar, S., Dangui, N.P., Lagier, J.C., Michelle, C., Raoult, D., and Fournier, P.E.

(2014) Genome sequence and description of *Nesterenkonia massiliensis* sp. nov. strain

NP1(T.). *Stand Genomic Sci* **9**: 866–882.

Falconi, N., Vieira, E.M., Baumgarten, J., Faria, D., and Fernandez Giné, G.A. (2015) The home

range and multi-scale habitat selection of the threatened maned three-toed sloth (*Bradypus*

torquatus). *Mamm Biol* **80**: 431–439.

Ferreira, A.C., Nobre, M.F., Moore, E., Rainey, F.A., Battista, J.R., and da Costa, M.S. (1999)

Characterization and radiation resistance of new isolates of *Rubrobacter radiotolerans* and

Rubrobacter xylanophilus. *Extremophiles* **3**: 235–238.

Forquin, M.P., Duvergey, H., Proux, C., Loux, V., Mounier, J., Landaud, S., et al. (2009). Identification

of brevibacteriaceae by multilocus sequence typing and comparative genomic hybridization analyses.

Appl Environ Microbiol **75**: 6406–6409.

Forquin-Gomez M.P., Weimer B.C., Sorieul L., Kalinowski J., Vallaeys T. (2014) The Family

Brevibacteriaceae. In: Rosenberg E., DeLong E.F., Lory S., Stackebrandt E., Thompson F. (eds) The

Prokaryotes. Springer, Berlin, Heidelberg.

Gaiser R.A., Medema, M.H., Kleerebezem, M., van Baarlen, P., Wells J.M. (2017) Draft genome

sequence of a porcine commensal, *Rothia nasimurium*, encoding a nonribosomal peptide synthetase

predicted to produce the ionophore antibiotic valinomycin. *Genome Announc* **5**: e00453-17.

Gellatly, S.L., and Hancock, R.E. (2013) *Pseudomonas aeruginosa*: new insights into pathogenesis

and host defenses. *Pathog Dis* **67**:159–173.

Gohli, J., Bøifot, K.O., Moen, L.V., Pastuszek, P., Skogan, G., Udekwu, K.I., and Dybwad, M. (2019)

The subway microbiome: seasonal dynamics and direct comparison of air and surface bacterial

communities. *Microbiome* **7**: e160.

Grice, E.A., and Segre, J.A. (2011) The skin microbiome. *Nat Rev Microbiol* **9**: 244–253.

Hansen B.G. (1985) Etiologic importance of coagulase-negative Micrococcaceae isolated from blood

cultures. *Acta Pathol Microbiol Immunol Scand B* **93**: 1–6.

Hassoun, A., Linden, P.K., and Friedman, B. (2017). Incidence, prevalence, and management of

MRSA bacteremia across patient populations-a review of recent developments in MRSA

management and treatment. *Critical care* **21**: e211.

Heyrman, J., Verbeeren, J., Schumann, P., Devos, J., Swings, J., and De Vos, P. (2004)

Brevibacterium picturae sp. nov., isolated from a damaged mural painting at the Saint-Catherine

chapel (Castle Herberstein, Austria). *Int J Syst and Evol Microbiol* **54**: 1537–1541.

Higginbotham, S., Wong, W.R., Linington, R.G., Spadafora, C., Iturrado, L., Spadafora, C., Iturrado,

L., Arnold A.E. (2014) Sloth hair as a novel source of fungi with potent anti-parasitic, anti-cancer and

anti-bacterial bioactivity. *PLoS ONE* **9**(1): e84549.

Hyatt, D., Chen, G.L., Locascio, P.F., Land, M.L., Larimer, F.W., and Hauser, L.J. (2010) Prodigal:

prokaryotic gene recognition and translation initiation site identification. *BMC bioinformatics* **11**: e119.

Jensen, P.R., Gontang, E., Mafnas, C., Mincer, T.J., and Fenical, W. (2005) Culturable marine

actinomycete diversity from tropical Pacific Ocean sediments. *Environ Microbiol* **7**:1039–1048.

Jurado, V., Miller, A.Z., Alias-Villegas, C., Laiz, Saiz-Jimenez, C. (2012) *Rubrobacter bracarensis* sp.

nov., a novel member of the genus *Rubrobacter* isolated from a biodeteriorated monument. *Syst Appl*

Microbiol **35**: 306-309.

Kämpfer, P., Glaeser, S.P., Busse, H.J., Abdelmohsen, U.R., and Hentschel, U. (2014) *Rubrobacter*

aplysinae sp. nov., isolated from the marine sponge *Aplysina aerophoba*. *Int J Syst Evol Microbiol* **64**:

705–709

Kandi, V., Palange, P., Vaish, R., Bhatti, A.B., Kale, V., Kandi, M. R., and Bhoomagiri, M.R. (2016).

Emerging bacterial infection: identification and clinical significance of *Kocuria* species. *Cureus* **8**:

e731.

Katı, H., İnce, İ. A., Demir, İ., and Demirbağ, Z. (2010) *Brevibacterium pityocampae* sp. nov., isolated

from caterpillars of *Thaumetopoea pityocampa* (Lepidoptera, Thaumetopoeidae). *Int J Syst Evol*

Microbiol **60**: 312–316.

Kaur, C., Kaur, I., Raichand, R., Bora, T.C., and Mayilraj, S. (2011) Description of a novel

actinobacterium *Kocuria assamensis* sp. nov., isolated from a water sample collected from the river

Brahmaputra, Assam, India. *Antonie van Leeuwenhoek* **99**: 721–726.

Kim, S.B., Nedashkovskaya, O.I., Mikhailov, V.V., Han, S.K., Kim, K.O., Rhee, M.S., and Bae, K.S.

(2004). *Kocuria marina* sp. nov., a novel actinobacterium isolated from marine sediment. *Int J Syst*

Evol Microbiol **54**:1617–1620.

Kloos, W.E., Zimmerman, R.J., and Smith, R.F. (1976) Preliminary studies on the characterization

and distribution of *Staphylococcus* and *Micrococcus* species on animal skin. *Appl Environ Microbiol*

31: 53–59.

Lahti L., and Shetty S. (2018) Introduction to the microbiome R package.URL

http://bioconductor.statistik.tu-

dortmund.de/packages/3.6/bioc/vignettes/microbiome/inst/doc/vignette.html

Li, D., Luo, R., Liu, C.M., Leung, C.M., Ting, H.F., Sadakane, K., Yamashita, H., and Lam, T.W.

(2016) MEGAHIT v1.0: A fast and scalable metagenome assembler driven by advanced

methodologies and community practices. *Methods* **102**: 3–11.

Liu, Q., Liu, Q., Meng, H., Lv, H., Liu, Y., Liu, J., et al. (2020). *Staphylococcus epidermidis* contributes

to healthy maturation of the nasal microbiome by stimulating antimicrobial peptide production. *Cell*

Host Microbe **27**: 68–78.

Lory S. (2014) The Family Staphylococcaceae. In: Rosenberg E., DeLong E.F., Lory S.,

Stackebrandt E., Thompson F. (eds) The Prokaryotes. Springer, Berlin, Heidelberg.

Lousada, M.B., Lachnit, T., Edelkamp, J., Rouillé, T., Ajdic, D., Uchida, Y., Di Nardo, A., Bosch, T.,

and Paus, R. (2021) Exploring the human hair follicle microbiome. *Brit J Dermatol* **184**: 802–815.

Maisnier-Patin, S., and Richard, J. (1995). Activity and purification of linenscin OC2, an antibacterial

substance produced by *Brevibacterium linens* OC2, an orange cheese coryneform bacterium. *Appl*

Environ Microbiol **61**: 1847–1852.

Matarrita-Carranza, B., Murillo-Cruz, C., Avendaño, R., Ríos, M. I., Chavarría, M., Gómez-Calvo, M.

L., Tamayo-Castillo, G., Araya, J.J., et al. (2021) *Streptomyces* sp. M54: an actinobacteria associated

with a neotropical social wasp with high potential for antibiotic production. *Antonie van Leeuwenhoek*

114: 379–398.

Mendoza, J.E., Peery, M.Z., Gutiérrez, G.A., Herrera, G., and Pauli, J.N. (2015) Resource use by the

two-toed sloth (*Choloepus hoffmanni*) and the three-toed sloth (*Bradypus variegatus*) differs in a

shade-grown agro-ecosystem. *J Trop Ecol* **31**: 49-55.

Molina-Mora, J.A., Chinchilla-Montero, D., Chavarría-Azofeifa, M., Ulloa-Morales, A.J., Campos-

Sánchez, R., Mora-Rodríguez, R., Shi, L., and García, F. (2020) Transcriptomic determinants of the

response of ST-111 *Pseudomonas aeruginosa* AG1 to ciprofloxacin identified by a top-down systems

biology approach. *Sci Rep* **10**: e13717.

Mounier, J., Rea, M.C., O'Connor, P.M., Fitzgerald, G.F., and Cogan, T.M. (2007). Growth

characteristics of *Brevibacterium*, *Corynebacterium*, *Microbacterium*, and *Staphylococcus* spp.

isolated from surface-ripened cheese. *Appl Environ Microbiol* **73**: 7732–7739.

Newstead, L.L., Varjonen, K., Nuttall, T., and Paterson, G.K. (2020) Staphylococcal-produced

bacteriocins and antimicrobial peptides: their potential as alternative treatments for *Staphylococcus*

aureus infections. *Antibiotics* **9**: e40.

Nguyen, T.H., Park, M.D., and Otto, M. (2017) Host response to *Staphylococcus epidermidis*

colonization and infections. *Front Cell Infect Microbiol* **7**: e90.

Nouioui, I., Carro, L., García-López, M., Meier-Kolthoff, J. P., Woyke, T., Kyrpides, N. C., Pukall, R.,

Klenk, H. P., Goodfellow, M., & Göker, M. (2018). Genome-based taxonomic classification of the

Phylum Actinobacteria. *Front Microbiol* **9**: e2007.

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R.,

O'Hara, R.B., Simpson, G.L., Solymos, P. M., Stevens, M.H.H., Szoecs, E., Wagner, H. (2019)

Vegan: community ecology package. R package version 2:5–6. URL https://cran.r-

project.org/web/packages/vegan/index.html

O'Sullivan, J.N., Rea, M.C., O'Connor, P.M., Hill, C., and Ross, R.P. (2019) Human skin microbiota is

a rich source of bacteriocin-producing staphylococci that kill human pathogens. *FEMS Microbiol Ecol*

95: fiy241.

Pauli, J.N., Mendoza, J.E., Steffan, S.A., Carey, C.C., Weimer, P.J., and Peery M.Z. (2014) A

syndrome of mutualism reinforces the lifestyle of a sloth. *Proc R Soc B* **281**: e20133006.

Polak-Witka, K., Rudnicka, L., Blume-Peytavi, U., and Vogt, A. (2020) The role of the microbiome in

scalp hair follicle biology and disease. *Exp Dermatol* **29**: 286–294.

Price, M.N., Dehal, P. S., & Arkin, A. P. (2009) FastTree: computing large minimum evolution trees

with profiles instead of a distance matrix. *Mol Biol Evol* **26**: 1641–1650.

Pridham, T.G., Anderson, P., Foley, C., Lindenfelser, L.A., Hesseltine, C.W., and Benedict, R.G.

(1957) A selection of media for maintenance and taxonomic study of *streptomycetes*. *Antibiot Annu*

1957: 947–953.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O.

(2013) The SILVA ribosomal RNA gene database project: improved data processing and web-based

tools. *Nucleic Acids Res* **41**: 590–596.

Qiu, X., Kulasekara, B. R., and Lory, S. (2009). Role of horizontal gene transfer in the evolution of

Pseudomonas aeruginosa virulence. *Genome Dyn* **6**: 126–139.

Ramanan, P., Barreto, J.N., Osmon, D.R., and Tosh, P.K. (2014) *Rothia* bacteremia: a 10-year

experience at Mayo Clinic, Rochester, Minnesota. *J Clin Microbiol* **52**: 3184–3189.

Reddy, G., Prakash, J., Prabahar, V., Matsumoto, G. I., Stackebrandt, E., and Shivaji, S. (2003)

Kocuria polaris sp. nov., an orange-pigmented psychrophilic bacterium isolated from an Antarctic

cyanobacterial mat sample. *Int J Syst Evol Microbiol* **53**: 183–187.

R-Core-Team (2019) R: A language and environment for statistical computing. R foundation for

statistical computing, Vienna, Austria. URL https://www.r-project.org/

Rodrigues Hoffmann A. (2017) The cutaneous ecosystem: the roles of the skin microbiome in health

and its association with inflammatory skin conditions in humans and animals. *Vet Dermatol* **28**: 60–

e15.

Rojas-Gätjens, D., Fuentes-Schweizer, P., Rojas-Jiménez, K., Pérez-Pantoja, D., Avendaño, R.,

Alpízar, R., Coronado-Ruíz, C., and Chavarría, M. (2022). Methylotrophs and hydrocarbon-degrading

bacteria are key players in the microbial community of an abandoned century-old oil exploration well.

Microb Ecol **83**: 83–99.

Ross, A.A., Müller, K.M., Weese, J.S., and Neufeld, J.D. (2018) Comprehensive skin microbiome

analysis reveals the uniqueness of human skin and evidence for phylosymbiosis within the class

Mammalia. *Proc Natl Acad Sci USA* **115**: 5786–5795.

Ross, A.A., Rodrigues Hoffmann, A., and Neufeld, J.D. (2019) The skin microbiome of vertebrates.

Microbiome **7**: e79.

Roux, V., and Raoult, D. (2009). *Brevibacterium massiliense* sp. nov., isolated from a human ankle

discharge. *Int J Syst Evol Microbiol* **59**: 1960–1964.

Sanford, J.A., and Gallo, R.L. (2013) Functions of the skin microbiota in health and disease. *Semin

Immunol* **25**: 370–377.

Saxena, R., Mittal, P., Clavaud, C., Dhakan, D.B., Roy, N., Breton, L., Misra, N., and Sharma, V.K.

(2021) Longitudinal study of the scalp microbiome suggests coconut oil to enrich healthy scalp

commensals. *Sci Rep* **11**: e7220

Singh, H., Du, J., Yang, J. E., Shik Yin, C., Kook, M., & Yi, T. H. (2016) *Brachybacterium horti* sp.

nov., isolated from garden soil. *Int J Syst Evol Microbiol* **66**: e189–195.

Stackebrandt E. (2014) The Family Dermabacteraceae. In: Rosenberg E., DeLong E.F., Lory S.,

Stackebrandt E., Thompson F. (eds) The Prokaryotes. Springer, Berlin, Heidelberg.

Sun, Y., Shi, Y.L., Wang, H., Zhang, T., Yu, L. Y., Sun, H., and Zhang, Y.Q. (2018) Diversity of

bacteria and the characteristics of Actinobacteria community structure in Badain Jaran desert and

Tengger desert of China. *Front Microbiol* **9**: e1068.

Suutari, M., Majaneva, M., Fewer, D.P., Voirin, B., Aiello, A., Friedl, T., Chiarello, A.G., and Blomster,

J. (2010) Molecular evidence for a diverse green algal community growing in the hair of sloths and a

specific association with *Trichophilus welckeri* (Chlorophyta, Ulvophyceae). *BMC Evol Biol* **10**: e86.

Stackebrandt, E., Koch, C., Gvozdiak, O., and Schumann, P. (1995) Taxonomic dissection of the

genus *Micrococcus*: *Kocuria* gen. nov., *Nesterenkonia* gen. nov., *Kytococcus* gen. nov.,

Dermacoccus gen. nov., and *Micrococcus* Cohn 1872 gen. emend. *Int J Syst Bacteriol* **45**: 682–692.

Tak, E.J., Kim, P.S., Hyun, D.W., Kim, H.S., Lee, J.Y., Kang, W., et al. (2018) Phenotypic and

genomic properties of *Brachybacterium vulturis* sp. nov. and *Brachybacterium avium* sp. nov. *Front*

Microbiol, **9**: e1809.

Talento, A.F., Malnick, H., Cotter, M., Brady, A., McGowan, D., Smyth, E., and Fitzpatrick, F. (2013)

Brevibacterium otitidis: an elusive cause of neurosurgical infection. *J Med Microbiol* **62**: 486–488.

Taube, E., Keravec, J., Vié, J.C., and Duplantier, J.M. (2008) Reproductive biology and postnatal

development in sloths, *Bradypus* and *Choloepus*: review with original data from the field (French

Guiana) and from captivity. *Mamm Rev* **31**: 173–188.

Thi, M., Wibowo, D., and Rehm, B. (2020) *Pseudomonas aeruginosa* Biofilms. *Int J Mol Sci* **21**:

e8671.

Uranga, C.C., Arroyo, P., Jr, Duggan, B.M., Gerwick, W.H., and Edlund, A. (2020) Commensal oral

Rothia mucilaginosa produces enterobactin, a metal-chelating siderophore. *mSystems*, **5**: e00161-20.

Valdés-Stauber, N., and Scherer, S. (1994). Isolation and characterization of Linocin M18, a

bacteriocin produced by *Brevibacterium linens*. *Appl Environ Microbiol* **60**: 3809–3814.

Vaughan, C., Ramírez, O., Herrera, G., Guries, R. (2007) Spatial ecology and conservation of two

sloth species in a cacao landscape in limón, Costa Rica. *Biodivers Conserv* **16**: 2293–2310.

Waage J.F. and Montgomery G. (1976) *Cryptoses choloepi*: A coprophagous moth that lives on a

Sloth. *Science* **193**: 157-158.

Watanabe, K., Yamada, A., Nishi, Y., Tashiro, Y., and Sakai, K. (2021) Host factors that shape the

bacterial community structure on scalp hair shaft. *Sci Rep* **11**: e17711.

Wujek, D.E., and Cocuzza, J.M. (1985) Morphology of hair of two- and three-toe sloths (Edentata:

Bradypodidae). *Revista de biologia tropical* **34**: 243–246.

Tables

Table 1. Identity and antimicrobial activity of bacteria isolated from sloths' hair.

Strain	Host	Activity against pathogen														
		SA	EC	PA	BS	CA	SAg	SS	SE	KP	SM	Cy	Co	Fu	Hy	Tr
Rothia koreensis 10C1	*Choloepus*	++	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Rothia kristinae 13C9	*Choloepus*	-	-	+	-	-	-	-	-	-	-	+	-	-	-	-
Rothia kristinae 13C8	*Choloepus*	+	-	+	+++	-	-	-	-	-	-	+	-	-	-	-
Rothia sp. 9C1	*Choloepus*	+	-	+	+++	-	-	-	-	-	-	-	-	-	-	-
Rothia kristinae 11C2	*Choloepus*	+	-	+	+	-	-	-	-	-	-	+	-	-	-	-
Rothia kristinae 6B2	*Bradypus*	-	++	++	-	-	-	-	-	-	-	-	-	-	-	-
Rothia kristinae 3B2	*Bradypus*	+	-	+	+++	-	+	-	-	-	-	-	-	-	-	-
Brevibacterium sediminis 11B2	*Bradypus*	-	++	-	-	-	-	-	-	-	-	-	-	-	-	-
Rothia kristinae 3B6	*Bradypus*	+	-	+	+	-	-	-	-	-	-	-	-	-	-	-

SA = *S. aureus*; EC = *E. coli*; PA = *P. aeruginosa*; BS = *B. subtilis*; CA = *C. albicans*; Sag = *S. agalatiae*; SS =

S. saprophyticus; SE = *S. epidermidis*; KP = *K. pneumonie*; SM = *S. marcenses*; Cy = *Cytospora* sp.; Co =

Colleotrichum sp.; Fu = *Fusarium* sp.; Hy = *Hypocrea* sp; Tr = *Trichoderma* sp.

Table 2. Putative biosynthetic gene clusters in *Rothia koreensis* (Bin 3), *Rothia kristinae* (Bin 1

and Bin 2) and *Brevibacterium* (Bin 4). BGCs were identified using Antismash 5.0 and classified

agains the MiBiG database as described in "Experimental procedures".

Bin	Region	Type	Most Similar known BGC	Similarity (%)
Rothia kristinae Bin_1	1	NRPS		
	2	NRPS		
	3	T3PKS		
Rothia kristinae Bin_2	1	Terpene	Carotenoid	35
	2	Betalactone	Microansamycin	7
	3	Betalactone	Microansamycin	7
	4	Terpene	Carotenoid	35
	5	NRPS	Salinichelins	23
Rothia koreensis 3	1	Betalactone	Microansamycin	7
	2	NRPS-like	Kistamicin	12
	3	T3PKS	Endophenazine A1	17
	4	Siderophore	Desferrioxamin B	60
	5	NRPS		
	6	Terpene	Pimaricin	11
Brevibacterium 4	1	Siderophore	Desferrioxamine E	75
	2	Terpene	Carotenoid	71
	3	T3PKS	Endophenazine A1	21
	4	RiPP-like		
	5	NAPAA		
	6	Ectoine	Ectoine	75
	7	NRPS	Pepticinnamin E	10
	8	NAPAA		

Figure legends

Figure 1. Sloths that inhabit in the Sloth Sanctuary of Costa Rica located in Cahuita, Costa Rica. A) *Choloepus hoffmani*, B) *Bradypus variegatus.*

Figure 2. SEM images of the hair of *Choloepus hoffmani* and *Bradypus variegatus*. A) *Choloepus* hair are filament-like structures B) Filaments present a width of 138 ± 13 μm. C) The hair present multiple cavities along its longitude D) These cavities seem to be the niche for the microbial community inhabiting sloth hair. E) The higher abundance of microorganisms in these furrows is probably associated to the lack of mechanical stress they suffer F) Most microorganisms presented a coccoid-shape morphology, also some rod-shaped microorganisms were present. G) *Bradypus* hair is smoother H) It have average width of 261 ± 36 μm. I) It present irregular fissures throughout the entire hair longitude. J) Microorganisms survive in hair surface K) Coccoid and bacilli shaped bacteria are observed across the hair. L) The density of microorganisms is lower probably as a consequence of the hair morphology.

Figure 3. Taxonomic composition of prokaryotic community inhabiting the hair of *Bradypus variegatus* and *Choloepus hoffmani*. Relative abundance of bacterial and archaeal organisms at the A) phylum, B) Order levels. C) Relative percentage of 16S rRNA gene sequences assigned to each ASV (y axis) across the 28 samples analyzed (x axis).The ASV were taxonomically classified using SILVA reference database v138, as described in "Experimental procedures". *Bradypus* samples are identified as B1 to B13 and *Choloepus* samples are identified as C1-C15.

956 **Figure 4. Non-metric multidimensional scaling (NMDS) analysis and Venn diagram of the**

957 **prokaryotic communities in the hair of both sloth's species**. A) The NMDS analysis revealed a

958 clustering of communities according to the host is shown. The NMDS and the permanova analyses

959 were performed with Package Vegan v2.5- 6. B) Venn diagram presenting the number of ASV of the

960 core microbiome shared between *Bradypus variegatus* and *Choloepus hoffmani*. Both sloths share

961 almost 50% of the core microbiome. Core microbiome was calculated using the R package

962 microbiome as described in Experimental procedures.

964 **Figure 5. Heat map of BGCs types from the MAGs assembly and references genomes**

965 **identified in antiSMASH and BiG-SCAPE.**