Natural and human-mediated drivers of microevolution in Neotropical palms: a

historical genomics approach

Louise Brousseau[1,2]*, Sylvain Santoni[3], Audrey Weber[3], Guillaume Odonne[4]

*corresponding author: louise.brousseau@ird.fr (ORCID 0000-0003-2769-2767)

[1]UMR AMAP, IRD, Univ. Montpellier, CIRAD, CNRS, INRAE, Montpellier, France

[2]UMR DIADE, IRD, Univ. Montpellier, CIRAD, Montpellier, France

[3]UMR AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, Montpellier, France

[4]UAR LEEISA, CNRS, Cayenne, French Guiana

Abstract

Abstract

•   Microevolution in Amazonia has been far less documented than in temperate regions

and still constitutes a major knowledge gap. Moreover, the actual influence of pre-

Columbian societies through the Holocene remains unclear, although it could provide

interesting insights on the relationship between natural and human-mediated

evolutionary processes in driving microevolution.

•   Because they are widespread and traditionally managed since pre-Columbian times,

Neotropical palms are choice models to investigate the drivers of microevolution in

Amazonia. In this study, we carried out a preliminary exploration of the genomic

diversity in two pairs of congeneric palm species in French Guiana (*Astrocaryum* spp.

and *Oenocarpus* spp.).

•   We built upon an original sampling design, taking into account both regional climate

variations and local-scale pre-Columbian occupation, and designed a new target capture

kit of 20,000 molecular probes scattered across exonic regions of more than 5,000

nuclear genes in Arecoïdeae ("ARECO5000+"). Hundreds of palm libraries were

sequenced through Illumina sequencing, providing a unique – high-coverage – genomic

dataset in these non-model species.

•   We explored patterns of genomic diversity and differentiation within and across

populations, bringing state-of-the-art knowledge about the roles of climate adaptation

and pre-Columbian domestication. By documenting original cases of "incipient"

domestication, these pioneer results open new avenues toward a better understanding of

microevolution in Amazonia and of the impact of pre-Columbian societies on present-

day biodiversity patterns.

Key word: *Oenocarpus* spp., *Astrocaryum* spp., targeted capture, Illumina sequencing, climate

adaptation, pre-Columbian societies, ring ditch, incipient domestication.

Introduction

Evolution in Amazonia has been far less documented than in temperate regions and still constitutes a major knowledge gap. Evolutionary studies in Amazonia also largely concentrated on macroevolution through long-term geological periods, providing interesting insights onto speciation and extinction dynamics in relation to landscape changes associated with the Andean uplift and the formation of the Amazon river (Hoorn et al. 2010), to species dispersal abilities (B. T. Smith et al. 2014) and to biotic interactions (Onstein et al. 2018). On the contrary, less attention has been paid to the microevolution of plants and animals during the Quaternary (Cavers and Dick 2013), and even less during the Holocene. This period, which started about 11,000 years ago, experienced both a global warming subsequent to the last glacial period resulting in the present climate, and the establishment of dense networks small-scale human societies subsequent to the human colonization of South America in late Pleistocene (Llamas et al. 2016; Rothhammer and Dillehay 2009). A better understanding of the drivers of plant and animal evolution during the Holocene could thus bring interesting insights on the interactions between natural and human-mediated evolutionary processes and their role shaping present-day patterns of biodiversity.

Neotropical palms of the Arecoideae subfamily are choice models to investigate the drivers of microevolution in Amazonia. Dated phylogenies revealed that Arecoideae crown node diversified in mid-Cretaceous in South America, suggesting a Neotropical origin (Baker and Couvreur 2013). Today, Neotropical palms are widespread in Amazonia ; Arecaceae standing at the second position of the most abundant plant families in forest tree inventories with some "hyperdominant" species (ter Steege et al. 2013). Neotropical palms also constitute a key resource to human societies since pre-Columbian times, whose multiple uses – including

subsistence food and building materials – have been extensively documented at least since the 1850's (Wallace 1853; N. Smith 2015). Surprisingly, they are also particularly abundant at archaeological sites with evidence of long-term human occupation, and some species (in particular *Astrocaryum* spp. and *Oenocarpus* spp.) are commonly credited as 'biological indicators' of ancient settlements (Odonne et al. 2019).

This is notably the case at the so-called "ring ditches" that are abundant in interfluvial terra-firme forests in lowland Amazonia, from Bolivia (John F Carson et al. 2015) to French Guiana (Petitjean Roget 1991). These pre-Columbian earthworks provide valuable evidence of anthropogenic disturbances, with belowground enrichments in pre-Columbian ceramics, organic matter and charcoals characterizing "anthropogenic soils" that are commonly attributed to slash-and-burn agriculture (Bodin et al. 2020; Brancier et al. 2014). Even if the actual magnitude of land use at these sites remain controversial – with hypotheses varying from extensive forest clearance to the exploitation of naturally open savanna (Watling et al. 2017; John Francis Carson et al. 2014) – there is no more doubt that pre-Columbian societies have strongly impacted forest landscapes, thus contrasting with previous depictions of Amazonia as an undisturbed, primary forest (Levis et al. 2017).

These repeated and converging observations feed the long-standing hypothesis that many plant species, including Neotropical palms, may have been "incipiently-" or "semi-" domesticated through pre-Columbian times (Clement 1999; Clement et al. 2010, 2006). Whether pre-Columbian societies have actually shaped patterns of palm diversity through domestication, and how human-mediated processes interacted with natural ones, remain however largely unknown. This knowledge gap is primarily underpinned by a profound lack of observational genomic data in non-model Neotropical species that we intend to fill in this study.

We carried out a preliminary exploration of the genomic diversity in two pairs of congeneric

Neotropical palms in coastal French Guiana: *Astrocaryum paramaca* and *A. sciophilum* on one

side, *Oenocarpus bacaba* and *O. bataua* on the other side. We sampled populations following

an original sampling design taking into account both regional and local-scale evolutionary

processes related to climate variations and pre-Columbian occupation, respectively. At the

regional scale, we sampled populations in study sites with contrasted rainfall to catch climate

adaptation. At the local scale, we sampled palms both on the top of ring ditches and in the

surrounding forests to catch the impact of domestication associated with local anthropogenic

disturbances. We designed a new capture kit composed of 20,000 molecular probes to target

exonic regions within thousands of nuclear genes, and carried out an extensive targeted-capture

experiment and next-generation sequencing (NGS) of hundreds of libraries.

Here, we introduce this new capture kit ("ARECO5000+") and provide new NGS data on the

state and distribution of genomic diversity in Neotropical palms in coastal French Guiana. We

investigate intra-specific patterns of diversity and differentiation driven by both neutral

evolutionary processes (gene flow and admixture) and selective ones, disentangling the roles

of (natural) climate adaptation and of (human-mediated) pre-Columbian domestication that

operated at the different spatial scales.

Materials and Methods

1. Palm species

We focused on four palm species that belong to the Arecoideae sub-family and that are widespread in forest landscapes of French Guiana: two species of the genus *Astrocaryum* (*A. paramaca* and *A. sciophilum*) and two species of the genus *Oenocarpus* (*O. bacaba* and *O. bataua*). *Oenocarpus bacaba* and *O. paramaca* are pan-Amazonian species that are broadly distributed from Colombia to north Brazil. *Oenocarpus bataua* stands at the 7[th] position amongst the 20 most abundant species in the Amazonian tree flora, while *O. bacaba* stands at the 33[th] position (ter Steege et al. 2013). The distribution of *A. paramaca* and *A. sciophilum* is restricted to northeast Amazonia (Suriname, French Guiana and Brazil, GBIF Secretariat: GBIF Backbone Taxonomy. https://doi.org/10.15468/39omei). Despite this, *A. sciophilum* stands at the 51[th] position amongst the most abundant species in forest tree inventories. Moreover, *O. bacaba* and *A. sciophilum* are two biological indicators of pre-Columbian settlements in French Guiana, as their abundance is significantly higher in forest plots with evidence of strong pre-Columbian anthropogenic disturbances than in natural plots (Odonne et al. 2019). *Astrocaryum* spp. and *Oenocarpus* spp. are major sources of food and of building and handicraft materials in Amazonia (Sosnowska et al. 2015; Cummings and Read 2016). Their leaves are traditionally used in basketry and to build houses (see for example maloca's walls and roof, García et al. 2015; Davy 2007; Fadiman 2008; Coomes 2004; Gutierrez 2020; Kahn 1993), while their fruits are used to make oil (*e.g.* 'seje' oil) and beverages (*e.g.* so called 'comou' and 'wassaï' in French Guiana and 'bacaba wine' in Northwest Amazonia, Cunha et al. 2019; Castro et al. 2014; Puerari et al. 2015).

2. Sampling strategy

Palm populations of the four palms species were sampled in five study sites in the coastal

French Guiana with different rainfall ranging from about 2400 to 3500 mm.year^{-1} (**Figure 1**):

three archaeological sites with ring ditches (SPA, MC87 and NOU), plus two additional

undisturbed sites without evidence of pre-Columbian occupation (LAU and PAR). In the three

study sites with ring ditches, we sampled palm populations both at the top of the ring ditches

and in the surrounding (undisturbed) forest (< 5 km from the ring ditch), **Table 1**. When the

species were present, we collected a piece of leaf for 20 to 100 adult palms per species in each

sampling site and local condition (pre-Columbian occupation "+" and "–").

Figure 1. Localization of the study sites and annual precipitation (in mm.year^{-1}) in French

Guiana. Current precipitations were extracted from WorldClim version 1 using R (libraires

"raster" and "sp") and are representative of the period 1960-1990

(http://www.worldclim.com/version1).

Table 1. Localization of populations in the different sampling sites and local conditions of pre-Columbian occupation. "+" refers the top of the ring ditch with evidence of pre-Columbian occupation, and "–" refers the surrounding sampling forest area without evidence of pre-Columbian occupation. "Nouragues North" and "Nouragues South" indicate the localization of two ring ditches in the north and in the south of the Nouragues Ecological research station, respectively. Coordinates are provided in decimal degrees (WGS84).

Study site	Site ID	Local condition	Latitude (N)	Longitude (E)	Rainfall (mm.year⁻¹)
Sparouine	SPA	–	5.272861999	-54.22715197	2529
	SPA	+	5.253569037	-54.253331	2491
Laussat	LAU	–	5.479205009	-53.59530601	2391
Paracou	PAR	–	5.279138032	-52.92425998	2818
Nouragues South	NOU_S	–	4.038217971	-52.68241298	3246
	NOU_S	+	4.029483022	-52.681869	3246
Nouragues North	NOU_N	–	4.081238005	-52.67338903	3299
	NOU_N	+	4.079981977	-52.67142003	3299
Montagne Couronnée 87	MC87	–	4.053646959	-52.09517601	3517
	MC87	–	4.05699201	-52.086698	3517

3. Probes design, targeted capture and sequencing

We took advantage of the availability of the chromosome-level reference genome for the oil palm (*Elaeis guineensis*, Arecaceae, Arecoideae) to design a custom of capture kit of 20,000 molecular probes located within 5,209 orthologous nuclear genes scattered across all 16 chromosomes ("ARECO5000+"). Details on probes design is provided in **Supplementary material 1**. Probe sequences are available upon request from the corresponding author. The probes were then synthetized by Daicel Arbor Biosciences (myBaits Custom DNA-Seq©). Molecular experiments were further performed at the AGAP genotyping platform (Montpellier, France). Plant DNA was extracted from 15 mg of dried leaves. Ten to fifteen best-quality DNAs

per population were retained for library preparation, targeted-capture and sequencing, resulting

in the sampling design described in **Table 2** and totaling 405 libraries (one per palm). Genomic

library preparation for multiplexed individuals and targeted capture by bait hybridization

followed published protocols (Rohland and Reich 2012; Mascher et al. 2013) with some

modifications. The molecular protocol is provided in **Supplementary material 2**. Briefly, for

each genotype a genomic barcoded library was built; 48 barcoded libraries were mixed together

for the capture by hybridization using the myBaits Custom DNA-Seq© kit "ARECO5000+".

Libraries of *Oenocarpus bataua* were sequenced (paired-end, $2 \times 150bp$) on an Illumina MiSeq

at the AGAP genotyping platform (Montpellier, France), while libraries of *Astrocaryum* spp.

and *Oenocarpus bacaba* were sequenced by the Get-PlaGe core facility (GenoToul platform,

INRA Toulouse, France http://www.genotoul.fr) on an Illumina HiSeq3000. Sequence data are

available on EMBL-ENA under accession PRJEB51800.

Table 2. Sampling design for library preparation and sequencing. "NA" indicates that the

species was absent at this sites and/or local condition.

Site ID	Local condition	*Astrocaryum paramaca*	*Astrocaryum sciophilum*	*Oenocarpus bacaba*	*Oenocarpus bataua*
SPA	–	12	12	12	12
SPA	+	12	12	12	12
LAU	–	12	12	12	12
PAR	–	12	12	12	12
NOU_S	–	12	NA	NA	NA
NOU_S	+	12	NA	12	12
NOU_N	–	15	10	NA	12
NOU_N	+	14	15	15	12
MC87	–	12	12	NA	12
MC87	+	12	12	NA	12
TOTAL		**125**	**97**	**75**	**108**

4. Bioinformatics

Reads demultiplexing and pre-processing. Reads were demultiplexed based on their 6

nucleotide barcodes with `fastq-multx` of the `ea-utils-2.7` suit of tools (Aronesty

2013) with options `-b -x -m 0 -d 1`, and split into individual .fastq files (two files per

library, corresponding to "R1" and "R2" paired reads). Read sequences that failed to pass the

chastity filter (identified "Y" in the "@" line in CASAVA 1.8) were discarded using the

command:

`zcat $lib_name.R1.fastq.gz | grep -A 3 '^@.* [^:]*:N:[^:]*:' |`

`grep -v "^--$" > $lib_name_pf.R1.fastq`.

Reads were further cleaned using `FASTX-Toolkit (RRID:SCR_005534) v.0.0.14`:

they were trimmed with `fastx_trimmer` with option `-Q 33 -f 10 -l 140`, resulting

in reads of 140-nucleotide long. Individual bases of low quality (phred-score < 25) were then

masked with `fastq_masker` with options `-q 25 -Q 33`. At last, read 3'-end and 5'-end

were cut and orphan reads were discarded using `sickle-1.33` (Joshi and Fass 2011) with

options `pe -l 100` (*i.e.* default quality threshold of 20 and length threshold of 100bp). Once

cleaned, reads were quality controlled with `FastQC-0.11.5` (Andrews 2010).

Mapping. Cleaned reads of *Astrocaryum* spp. and *Oenocarpus* spp. libraries were mapped

separately against *Elaeis guineensis* reference genome (EG5) using `bwa` (H. Li and Durbin

2010, 2009): the reference was indexed with the command `bwa index` and paired-reads were

mapped using `bwa mem` with option `-t 10`.

Mapping were post-processed using `samtools-1.9` (H. Li 2011; H. Li et al. 2009).

Multiple mapped reads were discarded from .bam alignments using `samtools view` with

the command:

```
200   samtools view -h mapping.bam | awk '$17 !~ /XA:/ || $1 ~ /^@/'
201   |    /usr/local/samtools-1.9/bin/samtools    view    -bS    >
202   mapping_unique.bam
```

Alignments were then sorted and indexed using `samtools sort` and `samtools index`,

respectively. Mapping statistics were computed using `samtools stats`, `samtools`

`flagstat` and `samtools bedcov`. Sequencing and mapping statistics are provided in

Supplementary materials 3 and 4.

Following a first mapping against *Elaeis guineensis* reference genome, the reference was

updated by editing genus-specific consensus (one for *Astrocaryum* spp. and one for *Oenocarpus*

spp.) to take into account the genomic divergence between *Elaeis guinensis* and the genera

under study. This step aimed to avoid detecting polymorphisms caused by inter-genus variation

in the following. This was achieved by calling variants using `samtools mpileup` with

options `-uf` for each *Astrocaryum* spp. and *Oenocarpus* spp. alignments, respectively. The two

consensus references were edited with `bcftools index` (v.1.3) and `bcftools`

`consensus` (Danecek et al. 2021). Then, the libraries of each species (*Astrocaryum*

paramaca, *A. sciophilum*, *Oenocarpus bacaba* and *O. bataua*) were mapped separately against

the corresponding genus-specific consensus reference following the mapping procedure

described above.

SNP and genotype calling. Variants were called using `samtools-1.9` and `bcftools-`

`1.3`: genus-specific consensus were indexed with `samtools faidx` and variant were called

with `samtools mpileup` with options `-g -t DP` followed by `bcftools call` with

options `-vc`. The resulting .vcf file was filtered with `grep` to discard monomorphic sites,

variants detected on unassembled scaffolds or chloroplast as well as indels to retain nuclear

single-nucleotide polymorphisms (SNPs) only. SNPs were post-filtered with `R version`

`3.5.1` (R Core Team 2021) to retain SNPs and genotypes with a quality threshold of 20, and

a per-library coverage threshold of 6X. SNPs and genotypes that did not meet the criteria were

edited as missing ("NA"), and SNPs that became monomorphic or contained more than 20% of

missing values were discarded. This resulted in 178,358 and 23,252 high-quality SNPs in

Astrocaryum spp. and *Oenocarpus* spp., respectively. Last, palm samples with more than 20%

of missing values were discarded. The list of samples retained for genetic analyses is provided

in **Supplementary material 5**.

5. Population genetics

Genome-wide patterns of genetic structure and diversity were computed on a random subset

of 5,000 SNPs in `R version 3.5.1` (command `sample` with argument `replace =`

`F`). Genotype tables were converted from tabular (data frame) to `genind` objects using the

function `df2genind` (package `adegenet`).

Genetic differentiation. Pairwise genetic differentiation (F_{ST}) was estimated both between

congeneric species and within each species. Weir and Cockerham's F_{ST} were estimated using

the function `genet.dist` (package `hierfstat`) with argument `method="WC84"`.

Individual ancestry. Genome-wide genetic structure was assessed for each species by

estimating individual ancestry (Frichot et al. 2014) with the function `snmf` (package `LEA`) with

arguments `ploidy = 2`, `entropy = TRUE` and `CPU = 8`. The genetic structure was

explored from `K=1` to `K=10` ancestral populations (*A. paramaca*), to `K=8` (*A. sciophilum*), to

`K=6` (*O. bacaba*) or to `K=9` (*O. bataua*). The most probable number of clusters was retained

based on cross-entropy criterion (lower values indicating better fit to the predictions) and each

palm sample was assigned to a given group with an ancestry threshold of 0.5.

Isolation-by-distance. Isolation-by-distance was tested across populations within each species

through Mantel tests between genetic and geographic distances. Genetic distance were

estimated with the function `dist.genpop` (package `adegenet`) with argument `method =`

`3` (*i.e.* coancestry coefficient or Reynolds' distance, Jombart 2008; Jombart and Ahmed 2011).

Geographic distances (euclidean) were computed from geographic coordinates using the

function `dist` with default arguments. Mantel tests between geographic and genetic distance

matrices were computed with the function `mantel.randtest` (package `ade4`, Mantel

1967; Thioulouse et al. 1997).

Admixture tests. Admixture between populations was evaluated through *f3* admixture tests for

all combinations of three populations (Patterson et al. 2012; Peter 2016; Reich et al. 2009):

$f_3(P_X ;P_A,P_B)$, where P_X is the target population and P_A and P_B are two source populations.

Admixture tests were encoded with `R version 3.5.1`. f_3 statistic was estimated following

the equation by Reich et al. (2009):

$$\mathbb{E}(p_X - p_a)(p_X - p_b),$$

where p_x, p_a and p_b are the allele frequencies in the populations P_X, P_A and P_B, respectively, and

$\mathbb{E}$ the expected value across all SNPs. Negative values indicate admixture between A and B.

The significance of f_3 was assessed by bootstrap (function `sample` with argument `replace`

`= T`) through 500 iterations. Z-scores were converted into p-values (left-tailed) with the

function `pnorm`.

Intra-population diversity. Observed (*Ho*) and expected heterozygosity (*He*) within each

species and populations was estimated with `adegenet` from `genind` objects. `Genind`

objects containing multiple populations were split into separate populations using the function

`seppop` and summary statistics were exported with the function `summary`. Whether *Ho* was

higher than *He* was assessed for each population through non-parametric Wilcoxon tests

(because of non-homogeneity of variances between *Ho* and *He*) with the function

`wilcox.test`. Inbreeding coefficients (*Fis*) were estimated following the formula: $Fis =$

$1 - \frac{Ho}{He}$. Hardy Weinberg equilibrium was tested for each SNP through Chi-squared tests with

the function `hw.test` (package `pegas`), and the proportion of SNPs showing significant

departure from HW equilibrium at a 1% threshold was assessed.

Selection test. We carried out a genome-wide scan for adaptive differentiation within each

species using `BayPass v.2.1`. Genotype data were converted into allele counts and

`BayPass` core model was run a first time using observed SNP counts as input. The number of

populations in each dataset was specified with option `-npop` and MCMC parameters were

specified with options `-nthreads 8 -thin 15 -burnin 2500 -npilot 15 -`

`pilotlength 5000`. The estimated covariance matrix (Ω), which provides an accurate

estimate of genome-wide allele frequency covariance across populations, was converted into a

correlation matrix in `R version 3.5.1` using the function `cov2cor`. The singular values

of the covariance matrix was decomposed using `svd` and left singular vectors (U) were

extracted to plot the genome-wide structure across populations. Pseudo-observed data (PODs)

of 10,000 SNPs (for *Oenocarpus* spp.) or 100,000 SNPs (for *Astrocaryum* spp.) were simulated

with the `simulate.baypass` function (`BayPass` miscellaneous R function), specifying

the total number of allele counts for each SNP (*i.e.* sample size) and the estimated

hyperparameters a_π and b_π (*i.e.* posterior mean and standard deviation of the Beta distribution

expressing the mean reference allele frequency, ~ancestral reference allele frequency) as input.

These PODs correspond to a simulated set of SNP counts that take into account the genome-

wide (neutral) genetic structure empirically inferred from observed SNPs counts. The core

model was run again using PODs as input under the same parameters to calibrate SNP-specific

statistics (standardized allele frequencies α^{std}, *XtX* and derived statistics) under neutral

hypothesis. `BayPass` outputs were post-processed with `R version 3.5.1`. The

robustness of the core model was assessed by evaluating the FMD distance (Förstner and

Moonen 2003) between covariance matrices estimated on observed data (Ω^{OBS}) and on

simulated data (Ω^{POD}) using `fmd.dist` function (`BayPass` miscellaneous R function). SNP-

specific departures from genome-wide (*i.e.* neutral) differentiation were detected based on the

joint distribution of:

• SNP-specific XtX_i (Günther and Coop 2013) estimated by `BayPass` (which describes

the differentiation across all populations for each SNP i explicitly corrected for genome-

wide covariance, whatever their regional and local environmental conditions), and

• two additional statistics (δ_i and Δ_i) derived from estimated standardized allele

frequencies ($\alpha_{i,j}^{STD}$, which describes allele frequencies for each SNP i within each

population j explicitly corrected for the genome-wide covariance matrix) and designed

to quantify variations in α^{STD} driven by climate adaptation and pre-Columbian

domestication, respectively.

Because α^{std} are unbounded, α^{std} were rescaled between 0 and 1 through min-max

normalization:

$$\alpha_{i,j}^{STD\,(OBS)} = \frac{\alpha_{i,j}^{STD\,(OBS)} - \min(\alpha^{STD})}{\max(\alpha^{STD}) - \min(\alpha^{STD})} \text{ and } \alpha_{i,j}^{STD\,(POD)} = \frac{\alpha_{i,j}^{STD\,(POD)} - \min(\alpha^{STD})}{\max(\alpha^{STD}) - \min(\alpha^{STD})},$$

where:

• $\alpha_{i,j}^{STD\,(OBS)}$ and $\alpha_{i,j}^{STD\,(POD)}$ are the standardized allele frequency at SNP i in population

j estimated from observed and simulated data respectively ;

• $\min(\alpha^{STD}) = \min(\alpha^{STD\,(OBS)}, \alpha^{STD\,(POD)})$;

• $\max(\alpha^{STD}) = \max(\alpha^{STD\,(OBS)}, \alpha^{STD\,(POD)})$.

The statistic δ quantifies clines in α^{STD} along the regional precipitation gradient:

$$\delta_i = |\beta_i|.r_i^2 \,,$$

where $|\beta_i|$ and r_i^2 are the absolute regression coefficient and the r-squared of the linear model

between allele frequency at SNP i and precipitation: $\alpha_{i,j}^{STD} = \beta_i.prec_j$, with $prec_j$ the

precipitation in population j (scaled between 0 and 1). Linear models were fitted for each SNP

with the function lm.

The statistic Δ quantifies absolute differences in α^{STD} between contrasted local conditions of

pre-Columbian occupation (+ *versus* –) within ($\Delta_{i,s}$) and across study sites (Δ_i):

$$\Delta_{i,s} = \left| \alpha_{i,(j+)}^{STD} - \alpha_{i,(j-)}^{STD} \right| ,$$

327   and

$$\Delta_i = \sum_s (\Delta_{i,s})$$

where $j+$ and $j–$ refer to populations located on the ring ditch and in the surrounding natural

forest for each study site s (*i.e.* single-site contrast), respectively, and $\Delta_{i,s}$ is the sum across

study sites s (*i.e.* multi-site contrast).

δ and Δ statistics where separately computed on observed data (δ_i^{OBS}, $\Delta_{i,s}^{OBS}$, Δ_i^{OBS}) and on PODs

(δ_i^{POD}, $\Delta_{i,s}^{POD}$, Δ_i^{POD}). A SNP was retained as being shaped by climate adaptation if it was both

over-differentiated across all populations whatever their local environment based on XtX at a

95% PODs threshold (*i.e.* condition 1), and if it displayed clinal allele frequency variation with

regional precipitation based on δ at a 95% PODs threshold (*i.e.* condition 2). Similarly, a SNP

was retained as being shaped by domestication if both condition 1 was met, and if it displayed

local differences in allele frequencies in relation to pre-Columbian occupation based on Δ at a

95% PODs threshold (*i.e.* condition 3).

Climate adaptation SNP	**Domestication SNP**
Condition 1: $XtX_i^{OBS} > q_{(95)}(XtX_i^{POD})$	
Condition 2: $\delta_i^{OBS} > q_{(95)}(\delta_i^{POD})$	Condition 3: $\Delta_i^{OBS} > q_{(95)}(\Delta_i^{POD})$

where:

• XtX_i^{OBS} is the SNP-specific differentiation estimated on observed data, and

$q_{(95)}(XtX_i^{POD})$ the 95-quantile of the distribution (across all SNPs) of SNP-specific

differentiation estimated on simulated data ;

• δ_i^{OBS} is the SNP-specific cline in allele frequency along the regional precipitation

gradient estimated on observed data, and $q_{(95)}(\delta_i^{POD})$ the 95-quantile of the distribution

(across all SNPs) of SNP-specific clines in allele frequency along the regional

precipitation gradient estimated on simulated data ;

• Δ_i^{OBS} is the SNP-specific difference in allele frequency between local conditions of pre-

Columbian occupation (+ *versus* –) estimated on observed data, and $q_{(95)}(\Delta_i^{POD})$ the

95-quantile of the distribution (across all SNPs) of SNP-specific differences in allele

frequency between local conditions of pre-Columbian occupation estimated on

simulated data.

Outlier blastn. Last, we retrieved the identity of genes containing precipitation and adaptation

SNPs. For each species, we extracted the sequences of the genomic regions surrounding each

outlier SNP (+/- 200 bp) from consensus genomic sequences using `bedtools getfasta`

(`bedtools-2.27.1`). Each SNP-surrounding region was then blasted against the NCBI

nucleotide database using `blastn` (`NCBI BLAST+ v. 2.10.1+`), limiting blast search by

taxonomy (*Elaeis guineensis* taxid = 51953 and *Phoenix dactylifera* taxid = 42345) and

returning the first good hit to optimize computation time.

Results

1. **Genome-wide genetic structure and differentiation**

Genetic differentiation. At the inter-specific level, pairwise F_{ST} across populations ranged

between 0.30 and 0.55 in *Astrocaryum* spp. and between 0.23 and 0.30 in *Oenocarpus* spp.

Intra-specific F_{ST} ranged between 0.003 and 0.4 in *A. paramaca*, between 0.01 and 0.11 in *A.*

sciophilum, between 0 and 0.06 in *O. bacaba* and between 0 and 0.01 in *O. bataua*, **Figure 2**

and **Supplementary material 6 (Tables S5.1 & S5.2)**. In *A. paramaca*, the population

originating from the undisturbed area close to the ring ditch in MC87 (i.e "MC87−") showed a

surprisingly high differentiation with all other population of this species ranging between 0.39

and 0.40, while the differentiation between all other populations ranged between 0 and 0.28.

Figure 2. Correlation plots showing patterns of genome-wide genetic differentiation across

populations of *Astrocaryum* spp. (left) and *Oenocarpus* spp. (right). "AP" refers to *A.*

paramaca, "AS" to *A. sciophilum*, "OBc" to *O. bacaba* and "OBt" to *O. bataua*.

Individual ancestry. Cross-entropy criterion of individual ancestry was lower for K=3 and

K=4 in *A. paramaca* and *A. sciophilum*, respectively, and for K=4 and K=2 in *O. bacaba* and

O.bataua, respectively, thus indicating a genome-wide genetic structuring at a grain lower than

the species level (**Supplementary material 6, Figures S6.1 & S6.2**). Within each species, the

genetic structure was consistent with the regional location of the palm populations from west

to east, **Figure 3**.

In *A. paramaca*, we detected one cluster in the west (in red) including all individuals from the

sites of Sparouine (SPA), Laussat (LAU) and Paracou (PAR), and two genetic clusters in the

east (sites of NOU and MC87), with all individuals associated to the blue group originating

from MC87.

In *A. sciophilum*, we detected two genetic clusters in the west: one corresponding to the study

site SPA (in green) and one (in purple) corresponding to the study sites LAU and PAR. In the

east (NOU and MC87), most individuals were assigned to the light blue group. Few individuals

originating from MC87 were assigned to a fourth group (in red). In addition, one individual was

assigned to the purple group and one ambiguous individual was not assigned to any group.

In *O. bacaba*, we detected three genetic groups in the west, and one in the east. The palms from

SPA were in majority assigned to the light blue group, while the populations from LAU and

PAR included palms assigned to both light blue and red groups. The third group (green)

corresponded to two individuals originating from the ring ditch (+) of SPA. The palms from

NOU were all assigned to the fourth group (purple).

In *O. bataua*, we detected two groups with one group (purple) including palms from SPA and

PAR, and another group (blue) including palms from NOU and MC87 except one palm in the

latter that clustered with the purple group with an ancestry coefficient slightly above 0.5. In

LAU, quite all palms clustered with the blue group except two palms clustered with the purple

group.

Figure 3. Geographic genetic structure. The pies indicate the proportion of individuals assigned to genetic each cluster within each study site and local condition of pre-Columbian occupation. Grey slices indicate ambiguous individuals that cannot be assigned to a given group with an ancestry coefficient > 0.5.

Isolation-by-distance. Patterns of isolation by distance were significant in *A. sciophilum* (p-value = 0.001) and *O.bataua* (p-value = 0.002), but not in *A. paramaca* (p-value = 0.20) nor in *O. bacaba* (p-value = 0.14). In *A. paramaca*, however, IBD became significant (p-value = 0.003) after exclusion of populations from MC87 (MC87 + and MC87–), **Supplementary material 6 (Figure S6.3)**.

Admixture. Significantly negative f_3 statistics were detected at a threshold of 5% for four target

populations of *Astrocaryum* spp. (*A. paramaca* from MC87 +, *A. sciophilum* from MC87 + and

MC87 – and *A. paramaca* from NOU_N –), and for one target population *of Oenocarpus* spp.

(*O.bacaba* in SPA +), **Supplementary material 6 (Figure S6.4)**. Populations of *A. paramaca*

from MC87 + appeared to be admixed between populations of *A. paramaca* from NOU (N and

S, + and –) and populations of *A. paramaca* from MC87 –, while *A. paramaca* from NOU_N –

appeared to be admixed between *A. paramaca* from NOU_S – and *A. sciophilum* from NOU_N

(+ and –), LAU – and/or PAR, thus suggesting patterns of inter-specific hybridization between

A. paramaca and *A. sciophilum* in the study site of NOU. Similar patterns of inter-specific

admixture also appeared in *A. sciophilum* from MC87 (+ and –), revealing admixture between

A. sciophilum from NOU_N (+ and –) and *A. paramaca* from MC87 –. *Oenocarpus bacaba*

from SPA + also appeared to be admixed between *O. bacaba* from SPA – and *O. bataua* from

different sites.

Intra-population diversity. In *Astrocaryum* spp., observed heterozygosity (*Ho*) ranged

between 0.136 (*A. sciophilum* in MC87 +) and 0.163 (*A. paramaca* in NOU_S –). In

Oenocarpus spp., *Ho* ranged between 0.351 (*O. bacaba* SPA – and LAU – ; *O.bataua* NOU_S

+, SPA – and PAR –) and 0.372 (*O. bacaba* SPA+). *Fis* was negative, indicating outcrossing.

Fis was slightly higher in *A. paramaca* than in *A. sciophilum* although non-significant

(Wilcoxon p-value = 0.1), and higher in *O. bacaba* than in *O. bataua* (Wilcoxon p-value =

0.0004), **Table 3** and **Supplementary material 6 (Figure S6.5)**. In *Astrocaryum* spp., *Fis* was

higher in the populations from MC 87 where it became non-significantly negative. *Fis* also tend

to be slightly higher in populations on ring ditches (+) than in surrounding natural populations

(–) in *Astrocaryum* spp. and *O. bataua* while the trend was opposite in *O. bacaba*, **Table 3** and

Supplementary material 6 (Figure S6.5). A large proportion of SNPs shew significant

departure from HW equilibrium with large variations across populations ranging from 7% to

12% in *A. paramaca*, from 0 % and 14% in *A. sciophilum*, from 18% to 27% in *O. bacaba* and

between 0 and 25% in *O.bataua*.

Table 3. Intra-population diversity statistics (*He*, *Ho* and *Fis*) in *Astrocaryum* spp. with

Wilcoxon tests for differences between *Ho* and *He* (** : p-value <0.01 ; * : p-value<0.05).

	Ho	*He*	*Fis*
A. paramaca			
SPA –	0.156 (sd=0.292)	0.105	-0.333 (sd=0.381) **
SPA +	0.156 (sd=0.294)	0.105	-0.337 (sd=0.396) **
LAU –	0.160 (sd=0.301)	0.107	-0.365 (sd=0.401) **
PAR –	0.157 (sd=0.294)	0.106	-0.345 (sd=0.389) **
NOU_N –	0.165 (sd=0.290)	0.114	-0.273 (sd=0.362) **
NOU_N +	0.162 (sd=0.289)	0.112	-0.288 (sd=0.372) **
NOU_S –	0.163 (sd=0.296)	0.110	-0.340 (sd=0.371) **
NOU_S +	0.162 (sd=0.292)	0.111	-0.312 (sd=0.371) **
MC87 –	0.140 (sd=0.285)	0.104	-0.141 (sd=0.587)
MC87 +	0.154 (sd=0.278)	0.128	-0.101 (sd=0.556)
A. sciophilum			
SPA –	0.143 (sd=0.303)	0.089	-0.496 (sd=0.408) **
SPA +	0.139 (sd=0.289)	0.092	-0.384 (sd=0.407) *
LAU –	0.137 (sd=0.287)	0.090	-0.403 (sd=0.400) *
PAR –	0.137 (sd=0.285)	0.091	-0.363 (sd=0.402) *
NOU_N –	0.144 (sd=0.292)	0.094	-0.383 (sd=0.389) **
NOU_N +	0.137 (sd=0.285)	0.091	-0.366 (sd=0.405) *
MC87 –	0.137 (sd=0.257)	0.142	0.030 (sd=0.614) *
MC87 +	0.136 (sd=0.272)	0.110	-0.004 (sd=0.625)

444   **Table 4.** Intra-population diversity statistics (*He*, *Ho* and *Fis*) in *Oenocarpus* spp. with

445   Wilcoxon tests for differences between *Ho* and *He* (** : p-value <0.01 ; * : p-value<0.05).

	Ho	*He*	*Fis*
O. bacaba			
SPA –	0.352 (sd=0.415)	0.205	-0.575 (sd=0.389) **
SPA +	0.373 (sd=0.393)	0.228	-0.438 (sd=0.408) **
LAU –	0.351 (sd=0.381)	0.226	-0.427 (sd=0.436) **
PAR –	0.356 (sd=0.385)	0.228	-0.453 (sd=0.442) **
NOU_N +	0.353 (sd=0.411)	0.207	-0.535 (sd=0.400) **
NOU_S +	0.351 (sd=0.413)	0.206	-0.554 (sd=0.398) **
O. bataua			
SPA –	0.351 (sd=0.428)	0.198	-0.659 (sd=0.367) **
SPA +	0.353 (sd=0.427)	0.200	-0.639 (sd=0.376) **

LAU –	0.352 (sd=0.430)	0.198	-0.700 (sd=0.347) **
PAR –	0.352 (sd=0.427)	0.200	-0.686 (sd=0.353) **
NOU_N –	0.358 (sd=0.427)	0.203	-0.644 (sd=0.375) **
NOU_N +	0.361 (sd=0.423)	0.206	-0.614 (sd=0.384) **
NOU_S +	0.357 (sd=0.424)	0.204	-0.631 (sd=0.378) **
MC87 –	0.355 (sd=0.421)	0.204	-0.617 (sd=0.376) **
MC87 +	0.353 (sd=0.422)	0.202	-0.642 (sd=0.365) **

2. SNP-specific differentiation

BayPass core model provided accurate estimates of genome-wide genetic structure, SNP-specific differentiation (XtX_i) and derived statistic designed to disentangle footprints of divergent selection caused by precipitation (δ_i) at the regional scale, and pre-Columbian occupation (*i.e.* domestication) at the local scale (Δ_i). Covariance matrices estimated on observed data (Ω^{OBS}) were close to covariance matrices estimated on pseudo-observed data (Ω^{POD}), with $\overline{FMD}$ = 1.06 in *A. paramaca*, $\overline{FMD}$ = 1.25 in *A. sciophilum*, $\overline{FMD}$ = 2.07 in *O. bacaba* and $\overline{FMD}$ = 2.9 in *O. bataua*, **Supplementary material 6 (Figures S6.6 to S6.9)**. The genome-wide genetic structure estimated by BayPass through covariance matrices was also consistent with the genetic structure estimated by snmf through individual ancestry, confirming the robustness of BayPass to empirically infer the genomic structure across populations **Supplementary material 6 (Figures S6.6 to S6.10)**. We further detected SNPs displaying patterns of differentiation related to precipitation and/or pre-Columbian occupation based on the joint distribution of XtX_i and derived statistics (δ_i and Δ_i), **Figure 4** and **Supplementary material 6 (Figure S6.11)**.

Figure 4. Example of SNP detection in relation to precipitation in *Oenocarpus bataua* based on the joint distribution of XtX_i and δ_i.

Between 0.74% (*A. sciophilum*) and 1.42% (*O. bataua*) climate adaptation SNPs were detected in the different species, and between 0.20% (*O. bataua*) and 1.25% (*A. paramaca*) domestication SNPs across all study sites. Moreover, the proportion of domestication SNPs was higher when considering all sites together (i.e. multi-site contrast) than within each site (single-site contrast), with many SNPs detected in one or two study sites only. Nevertheless, many domestication SNPs were shared between study sites: in *A. paramaca*, 1,252 domestication SNPs were detected in two sites, 285 were detected in three sites, and 28 in four sites ; in *A. sciophilum*, 564 SNPs were detected in two sites and 105 in three sites ; in *O. bataua*, only 10 SNPs were detected in two sites. The number of shared SNPs between pairs of study sites is provided in **Supplementary material 6 (Tables S6.3 to S6.6).**

476   **Table 5.** Number of over-differentiated SNPs detected in the four species in relation to climate

477   adaptation and domestication (out of 178,358 SNPs in *Astrocaryum* spp. and 23,252 SNPs in

478   *Oenocarpus* spp.). Because *O. bacaba* was only present on the ring ditches in NOU_N and

479   NOU_S and was absent from MC87, the detection of domestication SNP was based on a single

480   contrast in SPA.

	A. paramaca	*A. sciophilum*	*O. bacaba*	*O. bataua*
Climate adaptation SNPs				
Across all populations and sites	1345 (0.75%)	1320 (0.74%)	297 (1.25%)	331 (1.42%)
Domestication SNPs				
Across sites (multi-site contrast)	2238 (1.25%)	386 (0.22%)	NA	47 (0.20%)
Within sites (single-site contrast)				
• *SPA*	624 (0.35%)	640 (0.36%)	284 (1.22%)	2 (0.01%)
• *NOU N*	679 (0.38%)	810 (0.45%)	NA	70 (0.30%)
• *NOU S*	623 (0.35%)	NA	NA	NA
• *MC87*	2212 (1.24%)	374 (0.21%)	NA	16 (0.07%)

The median of SNP-specific differentiation (XtX_i) for neutral SNPs was 9.8 (Q90 = [8.6 ; 12.0])

in *A. paramaca*, 8.0 ([6.7 ; 9.9]) in *A. sciophilum*, 5.6 ([4.5 ; 8.1]) in *O. bacaba* and 8.7 ([7.6 ;

11.0]) in *O. bataua*. XtX_i ranged between 9.6 ([8.2 ; 14.6]) in *O. bacaba* and 13.9 ([12.5 ; 18.1])

in *A. paramaca* for climate adaptation SNP, and between 10.48 ([8.2 ; 13.8]) in *O. bacaba* and

13.1 ([12.5 ; 16.0]) in *A. paramaca* without differences in the extent of genetic differentiation

between climate adaptation and domestication SNPs, **Figure 6**.

Figure 6. Distribution of SNP-specific differentiation (XtX_i) in neutral SNPs, domestication (DOM) and climate adaptation SNPs (PREC).

Interestingly, several SNPs shew variations in allele frequency in relation to both precipitation and pre-Columbian occupation (**Supplementary material 6**, **Tables S6.3 to S6.6**)), allowing to distinguish four groups of SNPs within each species: (i) neutral SNPs, (ii) climate adaptation SNPs showing (clinal) variations in allele frequency in relation to regional precipitation, (iii) domestication SNPs showing differences in allele frequency in relation to local pre-Columbian occupation, and (iv) climate and domestication SNPs showing allele frequency variations in relation to both, **Figure 7**.

Figure 7. Principal Component Analysis representing the multivariate projection of SNPs based on SNP-specific estimates of XtX_i, δ_i, Δ_i and $\Delta_{i,s}$ in *Oenocarpus bataua*. Green: neutral SNPs (*i.e.* non-outlier); blue: climate adaptation SNPs ("PREC") ; pink: domestication SNPs ("DOM") ; violin: climate and domestication SNPs ("PREC+DOM"). Heatmaps on the right show standardized allele frequencies in the different populations and for the different SNPs of each group.

Discussion

This study opens a new research framework to document microevolution in non-model Amazonian palms. We provide original observational genomic data describing the state and distribution of genomic diversity in French Guiana, and we bring new knowledge about the microevolutionary processes at play, disentangling neutral and selective processes on one side, and natural and human-mediated ones on the other.

First of all, we must recall that all the four species were not present in all study sites and local conditions of pre-Columbian occupation. This was particularly the case for *Oenocarpus* spp. that were more difficult to find and to collect in natural forest areas than on ring ditches were these species are frequent. As a result, we did not collect any *Oenocarpus* spp. in NOU_S –, while *O. bacaba* was also absent from NOU_N – and MC87 (+ and –). This is consistent with previous observations at the community level by Odonne et al. 2019 who reported an over-abundance of palms in sites with strong evidence of pre-Columbian occupation compared to natural forest landscapes. This constrained our study, as we had to deal with incomplete sampling designs. Nevertheless, the amount of population-scale NGS data acquired was largely sufficient to explore microevolutionary processes in the studied palm species, and to disentangle natural and human mediated ones.

Neutral microevolution driven by gene flow and admixture.

Patterns of genome-wide diversity revealed a west-east genetic structuring at the regional scale, with similar trends between the four species. Genome-wide structure and differentiation are commonly driven by neutral evolutionary processes, in particular gene flow. This is comforted by significant isolation by distance we detected in three species, indicating that regional patterns of genomic diversity are primarily driven by gene flow, which gradually decreases with the

geographic distances between populations. We also detected intriguing patterns of genome-

wide genetic structure in *Astrocaryum* spp. in the study site of MC87. In particular, the natural

population (–) of *A. paramaca* revealed high levels of differentiation (about 40%) with

populations from other study sites, while the population on the ring ditch was closer to

populations from other study sites than to the population in the surrounding natural forest. A

similar pattern was also observed in the congeneric species *A. sciophilum*, although to a lesser

extent, suggesting a site-specific process probably involving inter-specific hybridization. This

hypothesis was corroborated by admixture tests, which suggested inter-specific admixture

between *A. paramaca* and *A. sciophilum* in the study site of MC87. Similarly, the genetic

structure of *Oenocarpus bataua* revealed a site-specific structure in SPA+ corroborated by

admixture tests suggesting inter-specific hybridization between *O. bacaba* and *O. bataua*.

These site-specific patterns of inter-specific hybridization were neither expected nor anticipated

when designing this study and carrying out the sampling. Most interestingly, *Astrocaryum*

putative inter-specific hybrids were identified either as *A. paramaca* or as *A. sciophilum*,

suggesting that they alternatively exhibit the morphological characteristics of one or the other

species. The two putative *Oenocarpus* hybrids, on their side, were both identified as

Oenocarpus bacaba. While there is no other species of *Oenocarpus* known in French Guiana,

there is at least fifteen different species of *Astrocaryum* recorded on *GBIF* (GBIF Secretariat:

GBIF   Backbone   Taxonomy.   https://doi.org/10.15468/39omei   Accessed   via

https://www.gbif.org/species/2738048 [30 March 2022]), raising the possibility of complex

inter-specific hybridization with possible cryptic species. Even though we cannot conclude

about the exact species involved in the pattern observed here, this preliminary observation

would deserve in-depth investigation of inter-specific hybridization and introgression in the

Astrocaryum complex.

Independently from site-specific hybridization patterns, estimates of intra-population genetic

diversity revealed high heterozygosity (H_o) and significantly negative F_{is}, indicating the

absence of inbreeding. These estimates are in line with previous knowledge about the breeding

system in Arecoideae that are known to be monoecious and predominantly outcrossed due to

protandry and/or genetic incompatibility (Henderson 1986; Eguiarte et al. 1992; Consiglio and

Bourne 2001; Picanço-Rodrigues et al. 2015; Gaiotto et al. 2003; Ottewell et al. 2012). The

genetic structure and diversity of populations on ring ditches was further close to populations

in the surrounding natural forests, indicating the absence of pre-Columbian impacts on

population genome-wide diversity and divergence. This result is not surprising, considering the

geographic proximity between "+" and "–" populations within study sites, which are probably

connected by extensive gene flow. The absence of genome-wide differentiation, however, does

not preclude the existence of SNP-specific selective processes driven by adaptation and/or

domestication.

Climate adaptation and pre-Columbian domestication.

Adaptation and domestication are acknowledged as important drivers of microevolution

worldwide, but they have been poorly documented in lowland Amazonia. While Central and

South America are known to be important centers of plant domestication, most studies have

focused on major crops such as maize (Kistler et al. 2018), papaya (Chávez-Pesqueira and

Núñez-Farfán 2017), manioc (Mühlen et al. 2019; Rival and McKey 2008; Alves-Pereira et al.

2020) or common bean (Rendón-Anaya et al. 2017), which provide the most obvious cases of

domestication. On the opposite, studies on minor crops are scarce although they are receiving

increasing attention since the last ten years (*e.g.* Brazil nut, Shepard and Ramirez 2011 ; piquiá,

Alves et al. 2016 ; peach palm, Clement et al. 2017 ; treegourd, Moreira et al. 2017). For its

part, the lack of interest in adaptation can be explained by the popularity of the neutral theory

under which the distribution of alleles within species (Kimura 1983) or species within

communities (Hubbell 2001) is assumed to be independent from fitness differences between

genotypes or species, being thus driven by stochastic mutation, demography and dispersal. In

hyperdiverse tropical rainforests, neutral models are powerful to predict global patterns of

diversity, as illustrated in the above section, and very useful to overcome the complexity of

niche differentiation modelling. Neutral and niche models are however highly complementary

(Chave 2004), and the neutral theory provides an elegant neutral hypothesis to test for niche

differentiation and adaptation (Leigh 2007). This is precisely the theoretical assumption of most

population genetics selection tests, which allows detecting gene- or SNP-specific departures

from the genome-wide, supposedly neutral, background of diversity and differentiation

(Beaumont and Nichols 1996).

By detecting many over-differentiated SNPs in relation to precipitation and pre-Colombian

occupation, our results bring new evidence of the coaction of climate adaptation and ancient

domestication in shaping patterns of genomic diversity across multiple spatial scales. As

expected, divergent selection related to adaptation and domestication affected a very small

fraction of the studied SNPs (always below 1.5%). Both the proportion of SNPs and the extent

of differentiation (XtX_i) were in the same order of magnitude between climate adaptation and

domestication SNPs, in spite of the different geographical scales involved (*i.e.* hundreds of

kilometers between study sites with different rainfall *versus* hundreds of meters between local

populations in contrasting conditions of pre-Columbian occupation). Regarding climate

adaptation, the present results are in line with a previous study in the hyperdominant tree *Eperua*

falcata which provided another evidence of climate adaptation along the sharp rainfall gradient

in coastal French Guiana (Brousseau et al. 2021).

Climate and domestication SNPs were located within or in the neighborhood (< 200bp) of many

genes whose exhaustive list is provided in **Supplementary material 7**. Although documenting

the functions of these genes in *Astrocaryum* spp. and *Oenocarpus* spp. is beyond the scope of

this study, we can nevertheless pick a few examples to illustrate the variety of genes involved.

For example, we detected climate adaptation SNPs in several genes of major importance for

plant development and response to stresses, including ABC transporters (Yazaki 2006; Do et

al. 2018), leaf rust disease resistance loci (Lim et al. 2015), Apetala2-like ethylene responsive

transcription factor (Licausi et al. 2013), aquaporins (G. Li et al. 2014; Maurel et al. 2015), and

several peroxidases (Blokhina et al. 2003; Yoshida et al. 2003). Several domestication SNPs

were detected in genes involved in lipid and sugar metabolism that could be of importance for

fruit development, including phospholipase D (Sun et al. 2011; Mao et al. 2007; Pinhero et al.

2003), acyl-transferases, fructose-bisphosphate aldolase (Feng et al. 2020; Yin et al. 2012),

phospholipid-transporting ATPases, and UDP-glucose 6-dehydrogenase 4.

Interestingly, many SNPs showed patterns of differentiation in relation to both climate

adaptation and domestication, indicating that these two processes are closely intertwined. In

other words, pre-Columbian societies had either domesticated "*in situ*" locally pre-adapted

populations or had to deal with adaptation during domestication. The absence of genome-wide

differentiation between palm populations on ring ditches and surrounding natural populations

tend to suggest that palm populations were collected in the forest near to the ring ditches.

Moreover, extensive gene flow between wild and domesticated populations had probably

continuously homogenized populations, thus preventing the genome-wide differentiation and

the fixation of the so-called "domestication syndrome" in domesticated populations (see next

section). Instead, SNP-specific patterns of domestication observed here provide a new evidence

of the subtle influence of pre-Columbian societies and depicts what is commonly referred to as

"incipient" domestication (Lins Neto et al. 2014; Casas et al. 2016; Clement 1999). Gene flow

since land abandonment may also have partly erased patterns of differentiation caused by

domestication (*i.e.* de-domestication, Qiu et al. 2017). Moreover, many SNPs were detected in

one study site only, supporting that pre-Columbian impacts were partly site-specific.

Nevertheless, many SNPs were also detected in more than one study site, indicating some

common trends in the different study sites, probably caused by similar management practices.

Whether observed domestication patterns result from "conscious" domestication through direct

selection for desired traits or "unconscious" domestication through indirect local adaptation to

anthropogenic landscapes (Gepts 2010; Engelbrecht 1916; Darlington 1973) cannot however

be solved yet, and would require in-depth investigation at the phenotypic level to assess which

traits have been selected for or against in palm populations established on ring ditches.

Moving beyond the domestication syndrome paradigm.

The domestication of plants is a long-term process, with multiple possible outcomes. However,

most studies have focused on major crops exhibiting clear phenotypic differences with wild

populations, often underpinned by a sharp – genome-wide – genetic differentiation. Studies on

major crops led to the elaboration of an earlier model of domestication through which the

domestication is initiated by the isolation of a small population with reduced diversity.

Domesticated populations are then expected to accumulate phenotypic differences from their

wild ancestors under the action of strong selection (*i.e.* artificial selection for desired traits and

natural selection exerted by new environmental conditions) reinforced by inbreeding, both

resulting in the fast decrease in genetic diversity and in the fixation of phenotypic differences

in domesticated populations: the so-called "domestication syndrome" (Hammer 1984; Harlan

et al. 1973; Allaby 2014). The domestication syndrome has been extensively documented, and

quite over-emphasized (Kantar et al. 2017), leading to a narrow view of the domestication

process and to a misleading dichotomous distinction between domesticated varieties with a

clear domestication syndrome, and other (supposedly wild) populations. The domestication

process is, however, gradual through time and variable across populations (Meyer et al. 2012;

O. Smith et al. 2019; Kantar et al. 2017), and it is not necessarily expected to result in a sharp

– genome-wide – differentiation, in particular in cases of domestication "*in situ*" with extensive

and continuous gene flow with wild populations (Meyer and Purugganan 2013). In a broader

sense, the domestication process is better depicted as the modification of the evolutionary

trajectory of a population through human-mediated evolutionary processes (including human-

mediated dispersal, artificial selection for useful traits and/or indirect adaptation to

anthropogenic landscapes) that can outcome – in most extreme cases – in the fixation of the

domestication syndrome.

Because lowland Amazonia offers a great diversity of domestication histories and cases, from

"incipiently-" or "semi-" domesticated populations to stable varieties (Lins Neto et al. 2014;

Clement 1999), it offers an ideal study system to reconsider the domestication process beyond

the domestication syndrome paradigm. Integrating the diversity of evolutionary histories

underlying domestication further opens a new perspective toward a better understanding of

microevolution in Amazonia, and the present study argues in favor of a paradigm shift to better

integrate "incipiently-" and "semi-" domesticated minor crops in domestication studies.

Supplementary materials

Supplementary material 1 (S1). ARECO5000+ Custom probe kit for targeted capture in Arecoideae.

Supplementary material 2 (S2). Molecular methods – library preparation, capture, sequencing and adapters.

Supplementary material 3 (S3). Sequencing and bioinformatics (405 libraries) – supplementary figures, tables and information.

Supplementary material 4 (S4). Fastq quality check.

Supplementary material 5 (S5). List of palm samples retained for genetic analyses.

Supplementary material 6 (S6). Population genetics – supplementary figures and tables.

Supplementary material 7 (S7). Genes containing climate adaptation and domestication SNPs.

Ethics statement

This study was carried out in compliance with the Nagoya protocol and French regulations on access to genetic resources with no direct commercial purposes. Palm samples were collected for fundamental research following declaration to the French Ministry of the Ecological Transition registered under the Internationally Recognized Certificate of Compliance ref. ABSCH-IRCC-FR-245923-1 of 23 May 2018. The authors declare no commercial use of the data generated and that no monetary benefit was derived from this study. Genomic data are made available according to shared values and principles for Open Science.

Data and codes availability

Sequencing data (i.e. paired-end reads) are available on the European Nucleotide Archive under accession PRJEB51800. Bioinformatics codes are available on GitLab under Project ID

696 34990830 (https://gitlab.com/LouiseBrousseau/palm-capture-bioinformatics). Data and code

697 remains under embargo until formal acceptance of the present article.

Competing interests

700 The authors declared no conflict of interest.

Authors contribution

703 LB and GO imagined the study and carried out palm sampling. LB designed the capture kit,

704 implemented the bioinformatics pipeline, and carried out bioinformatics and population

705 genetics. SS and AW carried out the molecular experiments (gDNA extraction and purification,

706 library preparation, targeted capture and sequencing of *O. bataua* librairies). All authors wrote

707 and approved the article.

Acknowledgements

This study has been conducted as part of the research program *PalmOmix* coordinated by L.

Brousseau. *PalmOmix* (ID 1702-01) was publicly funded by the French National Research

Agency (Agence Nationale de la Recherche, ANR) under the "*Investissement d'avenir*"

program with reference ANR-10-LABX-001-01 LabEx Agro and coordinated by Agropolis

Foundation under the frame of I-SITE MUSE (ANR-16-IDEX-0006). Sampling in the

Nouragues research Station has been supported by the LongTIme project coordinated by G.

Odonne and J-F. Molino and funded by an "*Investissement d'avenir*" grant from the French

National Research Agency (Agence Nationale de la Recherche, ANR) with reference CEBA:

ANR-10-LABX-25-01.

We are grateful to the Nouragues Ecological Research Station (CNRS) and the Réserve

Naturelle des Nouragues, to the Paracou Research Station (CIRAD) and to the National Forests

Office (ONF) in French Guiana for access to the study sites. We are also grateful to GenoToul

("GeT" and bioinformatics) facilities for next-generation sequencing of *Astrocaryum* spp. and

O. bacaba libraries, and for access to computing facilities to carry out bioinformatics analyses.

We thank Thierry Joët, Stéphane Dussert and James Tregear for their help in pre-filtering

candidate proteins ("bottom-up approach") to design the custom capture kit. We also thank T.

Joët for his help in palm sampling and Anaïs Prud'homme who explored *O. bataua* NGS data

as part of her internship at UMR DIADE under the supervision of L. Brousseau.

References

Allaby, Robin G. 2014. "Domestication Syndrome in Plants." In *Encyclopedia of Global Archaeology*, 2182–84. Springer New York. https://doi.org/10.1007/978-1-4419-0465-2_2416.

Alves-Pereira, Alessandro, Charles R. Clement, Doriane Picanço-Rodrigues, Elizabeth Ann Veasey, Gabriel Dequigiovanni, Santiago Linorio Ferreyra Ramos, José Baldin Pinheiro, Anete Pereira de Souza, and Maria Imaculada Zucchi. 2020. "A Population Genomics Appraisal Suggests Independent Dispersals for Bitter and Sweet Manioc in Brazilian Amazonia." *Evolutionary Applications* 13 (2): 342–61. https://doi.org/10.1111/eva.12873.

Alves, Rubana P., Carolina Levis, and Charles R. Clement. 2016. "Use and Management of Piquiá Suggest *in Situ* Domestication along the Lower Tapajós River, Brazilian Amazonia." *Economic Botany* 70 (2): 198–202. https://doi.org/10.1007/s12231-016-9340-4.

Andrews, Simon. 2010. "FastQC: A Quality Control Tool for High Throughput Sequence Data."

Aronesty, Erik. 2013. "Comparison of Sequencing Utility Programs." *The Open Bioinformatics Journal* 7 (1): 1–8. https://doi.org/10.2174/1875036201307010001.

Baker, William J., and Thomas L.P. Couvreur. 2013. "Global Biogeography and Diversification of Palms Sheds Light on the Evolution of Tropical Lineages. I. Historical Biogeography." Edited by W. Daniel Kissling. *Journal of Biogeography* 40 (2): 274–85. https://doi.org/10.1111/j.1365-2699.2012.02795.x.

Beaumont, Mark A, and Richard A Nichols. 1996. "Evaluating Loci for Use in the Genetic Analysis of Population Structure." *Proceedings of The Royal Society B: Biological*

Sciences 263: 1619–26.

Blokhina, Olga, Eija Virolainen, and Kurt V. Fagerstedt. 2003. "Antioxidants, Oxidative

Damage and Oxygen Deprivation Stress: A Review." *Annals of Botany* 91 (2): 179–94.

https://doi.org/10.1093/aob/mcf118.

Bodin, Stephanie C., Jean François Molino, Guillaume Odonne, and Laurent Bremond. 2020.

"Unraveling Pre-Columbian Occupation Patterns in the Tropical Forests of French

Guiana Using an Anthracological Approach." *Vegetation History and Archaeobotany* 29

(5): 567–80. https://doi.org/10.1007/s00334-019-00767-w.

Brancier, Jeanne, Cecilia Cammas, Dominique Todisco, and Eric Fouache. 2014. "A

Micromorphological Assessment of Anthropogenic Features in Pre-Columbian French

Guiana Dark Soils (FGDS): First Results." *Zeitschrift Fur Geomorphologie* 58

(October): 109–39. https://doi.org/10.1127/0372-8854/2014/S-00159.

Brousseau, Louise, Paul V.A. Fine, Erwin Dreyer, Giovanni G. Vendramin, and Ivan Scotti.

2021. "Genomic and Phenotypic Divergence Unveil Microgeographic Adaptation in the

Amazonian Hyperdominant Tree Eperua Falcata Aubl. (Fabaceae)." *Molecular Ecology*

30 (5): 1136–54.

Carson, John F, Jennifer Watling, Francis E Mayle, Bronwen S Whitney, José Iriarte, Heiko

Prümers, and J Daniel Soto. 2015. "Pre-Columbian Land Use in the Ring-Ditch Region

of the Bolivian Amazon." *The Holocene* 25 (8): 1285–1300.

https://doi.org/10.1177/0959683615581204.

Carson, John Francis, Bronwen S. Whitney, Francis E. Mayle, José Iriarte, Heiko Prům̈ers, J.

Daniel Soto, and Jennifer Watling. 2014. "Environmental Impact of Geometric

Earthwork Construction in Pre-Columbian Amazonia." *Proceedings of the National*

Academy of Sciences of the United States of America 111 (29): 10497–502.

776 https://doi.org/10.1073/pnas.1321770111.

777 Casas, Alejandro, José Blancas, Adriana Otero-Arnaiz, Jeniffer Cruse-Sanders, Rafael Lira,

778  Aidé Avendaño, Fabiola Parra, et al. 2016. "Evolutionary Ethnobotanical Studies of

779  Incipient Domestication of Plants in Mesoamerica." In , 257–85. Springer, New York,

780  NY. https://doi.org/10.1007/978-1-4614-6669-7_11.

781 Castro, J. W., M. S. Hernández, and R. H. Gutiérrez. 2014. "Uses for Amazonian Seje Oil

782  (*Oenocarpus Bataua*) Extracted by Mechanical and Manual Methods." In *Acta*

783  *Horticulturae*, 1047:335–40. International Society for Horticultural Science.

784  https://doi.org/10.17660/actahortic.2014.1047.43.

785 Cavers, Stephen, and Christopher W. Dick. 2013. "Phylogeography of Neotropical Trees."

786  *Journal of Biogeography* 40 (4): 615–17. https://doi.org/10.1111/jbi.12097.

787 Chave, Jerome. 2004. "Neutral Theory and Community Ecology." *Ecology Letters* 7 (august):

788  241–53. https://doi.org/10.1111/j.1461-0248.2003.00566.x.

789 Chávez-Pesqueira, Mariana, and Juan Núñez-Farfán. 2017. "Domestication and Genetics of

790  Papaya: A Review." *Frontiers in Ecology and Evolution* 5: 155.

791  https://www.frontiersin.org/article/10.3389/fevo.2017.00155.

792 Clement, Charles R. 1999. "1942 and the Loss of Amazonian Crop Genetic Resources. I. The

793  Relation between Domestication and Human Population Decline." *Economic Botany*.

794  SpringerNew York Botanical Garden Press. https://doi.org/10.2307/4256179.

795 Clement, Charles R., Joseph M. McCann, and Nigel J. H. Smith. 2006. "Agrobiodiversity in

796  Amazônia and Its Relationship with Dark Earths." In *Amazonian Dark Earths*, 159–78.

797  Kluwer Academic Publishers. https://doi.org/10.1007/1-4020-2597-1_9.

798 Clement, Charles R, Michelly de Cristo-Araújo, Geo Coppens d'Eeckenbrugge, Vanessa

799  Maciel dos Reis, Romain Lehnebach, and Doriane Picanço-Rodrigues. 2017. "Origin and

Dispersal of Domesticated Peach Palm." *Frontiers in Ecology and Evolution* 5: 148. https://www.frontiersin.org/article/10.3389/fevo.2017.00148.

Clement, Charles R, Michelly de Cristo-Araújo, Geo Coppens D'Eeckenbrugge, Alessandro Alves Pereira, and Doriane Picanço-Rodrigues. 2010. "Origin and Domestication of Native Amazonian Crops." *Diversity* 2 (1): 72–106. https://doi.org/10.3390/d2010072.

Consiglio, Trisha K., and Godfrey R. Bourne. 2001. "Pollination and Breeding System of a Neotropical Palm *Astrocaryum Vulgare* in Guyana: A Test of the Predictability of Syndromes." *Journal of Tropical Ecology* 17 (4): 577–92. https://doi.org/10.1017/S0266467401001432.

Coomes, Oliver T. 2004. "Rain Forest 'Conservation-through-Use'? Chambira Palm Fibre Extraction and Handicraft Production in a Land-Constrained Community, Peruvian Amazon." *Biodiversity and Conservation* 13 (2): 351–60. https://link.springer.com/article/10.1023/B:BIOC.0000006503.90980.e8.

Cummings, Anthony R., and Jane M. Read. 2016. "Drawing on Traditional Knowledge to Identify and Describe Ecosystem Services Associated with Northern Amazon's Multiple-Use Plants." *International Journal of Biodiversity Science, Ecosystem Services & Management* 12 (1–2): 39–56. https://doi.org/10.1080/21513732.2015.1136841.

Cunha, Vânia Maria Borges, Marcilene Paiva da Silva, Sérgio Henrique Brabo de Sousa, Priscila do Nascimento Bezerra, Eduardo Gama Ortiz Menezes, Nayara Janaina Neves da Silva, Débora Ariane Dornelas da Silva Banna, Marilena Emmi Araújo, and Raul Nunes de Carvalho Junior. 2019. "Bacaba-de-Leque (*Oenocarpus Distichus* Mart.) Oil Extraction Using Supercritical CO_2 and Bioactive Compounds Determination in the Residual Pulp." *Journal of Supercritical Fluids* 144 (February): 81–90. https://doi.org/10.1016/j.supflu.2018.10.010.

Danecek, Petr, James K. Bonfield, Jennifer Liddle, John Marshall, Valeriu Ohan, Martin O.

Pollard, Andrew Whitwham, et al. 2021. "Twelve Years of SAMtools and BCFtools."

GigaScience 10 (2). https://doi.org/10.1093/gigascience/giab008.

Darlington, Cyril Dean. 1973. *Chromosome Botany and the Origins of Cultivated Plants*.

London: Allen & Unwin.

Davy, Damien. 2007. *"Vannerie et Vanniers" Approche Ethnologique d'une Activité*

Artisanale En Guyane Française. Université d'Orleans.

Do, Thanh Ha Thi, Enrico Martinoia, and Youngsook Lee. 2018. "Functions of ABC

Transporters in Plant Growth and Development." *Current Opinion in Plant Biology*.

Elsevier Ltd. https://doi.org/10.1016/j.pbi.2017.08.003.

Eguiarte, Luis E., Nidia Perez-Nasser, and Daniel Piñero. 1992. "Genetic Structure,

Outcrossing Rate and Heterosis in *Astrocaryum Mexicanum* (Tropical Palm):

Implications for Evolution and Conservation." *Heredity* 69 (3): 217–28.

https://doi.org/10.1038/hdy.1992.119.

Engelbrecht, Th. H. 1916. "Über Die Entstehung Einiger Feldmäßig Angebauter

Kulturpflanzen." *Geographische Zeitschrift* 22 (6): 328–34.

http://www.jstor.org/stable/27809914.

Fadiman, Maria G. 2008. "Use of Mocora, *Astrocaryum Standleyanum* (Arecaceae), by Three

Ethnic Groups in Ecuador: Differences, Similarities and Market Potential." *Journal of*

Ethnobiology 28 (1): 92–109. https://doi.org/10.2993/0278-

0771(2008)28[92:UOMASA]2.0.CO;2.

Feng, Dan, Li Gao, Yusheng Zheng, Dongdong Li, and Peng Zhou. 2020. "Molecular Cloning

and Function Characterisation of a Cytoplasmic Fructose-1, 6-Bisphosphate Aldolase

Gene from Coconut (*Cocos Nucifera* L.)." *Journal of Horticultural Science and*

848   *Biotechnology* 95 (6): 703–11. https://doi.org/10.1080/14620316.2020.1749139.

849 Förstner, Wolfgang, and Boudewijn Moonen. 2003. "A Metric for Covariance Matrices." In

850   *Geodesy-The Challenge of the 3rd Millennium*, 299–309. Berlin, Heidelberg: Springer

851   Berlin Heidelberg. https://doi.org/10.1007/978-3-662-05296-9_31.

852 Frichot, Eric, François Mathieu, Théo Trouillon, Guillaume Bouchard, and Olivier François.

853   2014. "Fast and Efficient Estimation of Individual Ancestry Coefficients." *Genetics* 196:

854   973–983. https://doi.org/10.1534/genetics.113.160572.

855 Gaiotto, Fernanda A, Dario Grattapaglia, and Roland Vencovsky. 2003. "Genetic Structure,

856   Mating System, and Long-Distance Gene Flow in Heart of Palm (Euterpe Edulis Mart.)."

857   *Journal of Heredity* 94 (5): 399–406. https://doi.org/10.1093/jhered/esg087.

858 García, Néstor, Gloria Galeano, Laura Mesa, Nicolás Castaño, Henrik Balslev, Rodrigo

859   Bernal, Néstor García, et al. 2015. "Management of the Palm *Astrocaryum Chambira*

860   Burret (Arecaceae) in Northwest Amazon." *Acta Botanica Brasilica* 29 (1): 45–57.

861   https://doi.org/10.1590/0102-33062014abb3415.

862 Gepts, Paul. 2010. "Crop Domestication as a Long-Term Selection Experiment." In *Plant*

863   *Breeding Reviews*, 1–44. Oxford: John Wiley & Sons, Inc.

864   https://doi.org/10.1002/9780470650288.ch1.

865 Günther, Torsten, and Graham Coop. 2013. "Robust Identification of Local Adaptation from

866   Allele Frequencies." *Genetics* 195 (1): 205–20.

867   https://doi.org/10.1534/genetics.113.152462.

868 Gutierrez, Maria Paz. 2020. "The Northwestern Amazon Malocas: Craft Now and Then."

869   *Journal of Material Culture* 25 (1): 3–35. https://doi.org/10.1177/1359183519836141.

870 Hammer, Karl. 1984. "Das Domestikationssyndrom (The Domestication Syndrome)." *Die*

871   *Kulturpflanze* 32 (1): 11–34. https://doi.org/10.1007/BF02098682.

Harlan, Jack R, J M J de Wet, and E Glen Price. 1973. "Comparative Evolution of Cereals."

Evolution 27 (2): 311–25. https://doi.org/10.2307/2406971.

Henderson, Andrew. 1986. "A Review of Pollination Studies in the Palmae." *The Botanical*

Review 52 (3): 221–59. https://doi.org/10.1007/BF02860996.

Hoorn, Carina, Frank P. Wesselingh, Hans Ter Steege, M. A. Bermudez, A. Mora, J. Sevink,

I. Sanmartín, et al. 2010. "Amazonia through Time: Andean Uplift, Climate Change,

Landscape Evolution, and Biodiversity." *Science* 330 (6006): 927–31.

https://doi.org/10.1126/science.1194585.

Hubbell, Stephen P. 2001. *The Unified Neutral Theory of Biodiversity and Biogeography*.

Princeton University Press.

Jombart, Thibaut. 2008. "Adegenet: A R Package for the Multivariate Analysis of Genetic

Markers." *Bioinformatics* 24 (11): 1403–5.

https://doi.org/10.1093/bioinformatics/btn129.

Jombart, Thibaut, and Ismaïl Ahmed. 2011. "Adegenet 1.3-1: New Tools for the Analysis of

Genome-Wide SNP Data." *Bioinformatics* 27 (21): 3070–71.

https://doi.org/10.1093/bioinformatics/btr521.

Joshi, N A, and Joseph Fass. 2011. "Sickle: A Sliding-Window, Adaptive, Quality-Based

Trimming Tool for FastQ Files (Version 1.33)."

Kahn, Francis. 1993. "Amazonian Palms: Food Resources for the Management of Forest

Ecosystems." *Tropical Forests, People and Food: Biocultural Interactions and*

Applications to Development 13: 153.

Kantar, Michael B, Amber R Nashoba, Justin E Anderson, Benjamin K Blackman, and Loren

H Rieseberg. 2017. "The Genetics and Genomics of Plant Domestication." *BioScience*

67 (11): 971–82. https://doi.org/10.1093/biosci/bix114.

Kimura, Motoo. 1983. *The Neutral Theory of Molecular Evolution*. Cambridge: Cambridge

University Press. https://doi.org/DOI: 10.1017/CBO9780511623486.

Kistler, Logan, S. Yoshi Maezumi, Jonas Gregorio De Souza, Natalia A. S. Przelomska,

Flaviane Malaquias Costa, Oliver Smith, Hope Loiselle, et al. 2018. "Multiproxy

Evidence Highlights a Complex Evolutionary Legacy of Maize in South America."

Science 362 (6420): 1309–13. https://doi.org/10.1126/science.aav0207.

Leigh, E. G. 2007. "Neutral Theory: A Historical Perspective." *Journal of Evolutionary*

Biology 20 (6): 2075–91. https://doi.org/10.1111/j.1420-9101.2007.01410.x.

Levis, Carolina, Flávia R C Costa, Frans Bongers, Marielos Peña-Claros, Charles R Clement,

André B Junqueira, Eduardo G Neves, Eduardo K Tamanaha, Fernando O G Figueiredo,

and Rafael P Salomão. 2017. "Persistent Effects of Pre-Columbian Plant Domestication

on Amazonian Forest Composition." *Science* 355 (6328): 925–31.

https://doi.org/10.1126/science.aal0157.

Li, Guowei, Véronique Santoni, and Christophe Maurel. 2014. "Plant Aquaporins: Roles in

Plant Physiology." *Biochimica et Biophysica Acta - General Subjects*. Elsevier.

https://doi.org/10.1016/j.bbagen.2013.11.004.

Li, Heng. 2011. "A Statistical Framework for SNP Calling, Mutation Discovery, Association

Mapping and Population Genetical Parameter Estimation from Sequencing Data."

Bioinformatics 27 (21): 2987–93. https://doi.org/10.1093/bioinformatics/btr509.

Li, Heng, and Richard Durbin. 2009. "Fast and Accurate Short Read Alignment with

Burrows–Wheeler Transform." *Bioinformatics* 25 (14): 1754–60.

https://doi.org/10.1093/bioinformatics/btp324.

———. 2010. "Fast and Accurate Long-Read Alignment with Burrows–Wheeler Transform."

Bioinformatics 26 (5): 589–95. https://doi.org/10.1093/bioinformatics/btp698.

Li, Heng, Bob Handsaker, Alec Wysoker, Tim Fennell, Jue Ruan, Nils Homer, Gabor Marth,

Goncalo Abecasis, Richard Durbin, and Genome Project Data Processing Subgroup.

2009. "The Sequence Alignment/Map Format and SAMtools." *Bioinformatics* 25 (16):

2078–79. https://doi.org/10.1093/bioinformatics/btp352.

Licausi, Francesco, Masaru Ohme-Takagi, and Pierdomenico Perata. 2013.

"APETALA2/Ethylene Responsive Factor (AP2/ERF) Transcription Factors: Mediators

of Stress Responses and Developmental Programs." *New Phytologist* 199 (3): 639–49.

https://doi.org/10.1111/nph.12291.

Lim, Chae Woo, Seung Hwan Yang, Ki Hun Shin, Sung Chul Lee, and Sang Hyon Kim.

2015. "The AtLRK10L1.2, Arabidopsis Ortholog of Wheat LRK10, Is Involved in ABA-

Mediated Signaling and Drought Resistance." *Plant Cell Reports* 34 (3): 447–55.

https://doi.org/10.1007/s00299-014-1724-2.

Lins Neto, Ernani M., Nivaldo Peroni, Alejandro Casas, Fabiola Parra, Xitlali Aguirre, Susana

Guillén, and Ulysses P. Albuquerque. 2014. "Brazilian and Mexican Experiences in the

Study of Incipient Domestication." *Journal of Ethnobiology and Ethnomedicine*. BioMed

Central Ltd. https://doi.org/10.1186/1746-4269-10-33.

Llamas, Bastien, Lars Fehren-Schmitz, Guido Valverde, Julien Soubrier, Swapan Mallick,

Nadin Rohland, Susanne Nordenfelt, et al. 2016. "Ancient Mitochondrial DNA Provides

High-Resolution Time Scale of the Peopling of the Americas." *Science Advances* 2 (4):

e1501385. https://doi.org/10.1126/sciadv.1501385.

Mantel, N. 1967. "The Detection of Disease Clustering and a Generalized Regression

Approach." *Cancer Research*.

Mao, Linchun, Huaqing Pang, Guoze Wang, and Chenggang Zhu. 2007. "Phospholipase D

and Lipoxygenase Activity of Cucumber Fruit in Response to Chilling Stress."

Postharvest Biology and Technology 44 (1): 42–47.

https://doi.org/10.1016/j.postharvbio.2006.11.009.

Mascher, Martin, Todd A. Richmond, Daniel J. Gerhardt, Axel Himmelbach, Leah Clissold,

Dharanya Sampath, Sarah Ayling, et al. 2013. "Barley Whole Exome Capture: A Tool

for Genomic Research in the Genus Hordeum and Beyond." *Plant Journal* 76 (3): 494–

505. https://doi.org/10.1111/tpj.12294.

Maurel, Christophe, Yann Boursiac, Doan Trung Luu, Véronique Santoni, Zaigham Shahzad,

and Lionel Verdoucq. 2015. "Aquaporins in Plants." *Physiological Reviews* 95 (4):

1321–58. https://doi.org/10.1152/physrev.00008.2015.

Meyer, Rachel S., Ashley E. DuVal, and Helen R. Jensen. 2012. "Patterns and Processes in

Crop Domestication: An Historical Review and Quantitative Analysis of 203 Global

Food Crops." *New Phytologist* 196 (1): 29–48. https://doi.org/10.1111/j.1469-

8137.2012.04253.x.

Meyer, Rachel S., and Michael D. Purugganan. 2013. "Evolution of Crop Species: Genetics of

Domestication and Diversification." *Nature Reviews Genetics* 14 (12): 840–52.

https://doi.org/10.1038/nrg3605.

Moreira, Priscila Ambrósio, Cédric Mariac, Leila Zekraoui, Marie Couderc, Doriane Picanço

Rodrigues, Charles R. Clement, and Yves Vigouroux. 2017. "Human Management and

Hybridization Shape Treegourd Fruits in the Brazilian Amazon Basin." *Evolutionary*

Applications 10 (6): 577–89. https://doi.org/10.1111/eva.12474.

Mühlen, Gilda Santos, Alessandro Alves-Pereira, Cássia Regina Limonta Carvalho, André

Braga Junqueira, Charles R. Clement, and Teresa Losada Valle. 2019. "Genetic

Diversity and Population Structure Show Different Patterns of Diffusion for Bitter and

Sweet Manioc in Brazil." *Genetic Resources and Crop Evolution* 66 (8): 1773–90.

https://doi.org/10.1007/s10722-019-00842-1.

Odonne, Guillaume, Martijn Bel, Maxime Burst, Olivier Brunaux, Miléna Bruno, Etienne

Dambrine, Damien Davy, et al. 2019. "Long-term Influence of Early Human

Occupations on Current Forests of the Guiana Shield." *Ecology* 100 (10): e02806.

https://doi.org/10.1002/ecy.2806.

Onstein, Renske E, William J Baker, Thomas L P Couvreur, Søren Faurby, Leonel Herrera-

Alsina, Jens-Christian Svenning, and W Daniel Kissling. 2018. "To Adapt or Go

Extinct? The Fate of Megafaunal Palm Fruits under Past Global Change." *Proceedings.*

Biological Sciences 285 (1880): 20180882. https://doi.org/10.1098/rspb.2018.0882.

Ottewell, K, E Grey, F Castillo, and J Karubian. 2012. "The Pollen Dispersal Kernel and

Mating System of an Insect-Pollinated Tropical Palm, *Oenocarpus Bataua*." *Heredity*

109 (6): 332–39. https://doi.org/10.1038/hdy.2012.40.

Patterson, Nick, Priya Moorjani, Yontao Luo, Swapan Mallick, Nadin Rohland, Yiping Zhan,

Teri Genschoreck, Teresa Webster, and David Reich. 2012. "Ancient Admixture in

Human History." *Genetics* 192 (3): 1065–93.

https://doi.org/10.1534/genetics.112.145037.

Peter, Benjamin M. 2016. "Admixture, Population Structure and F-Statistics." *Genetics* 4:

1485–1501. https://doi.org/10.1534/genetics.115.183913.

Petitjean Roget, Hugues. 1991. "50 Sites de Montagnes En Guyane Française: Contribution A

L'inventaire Des Sites Archeologiques D'Emile Abonnenc." In *Proceedings of the*

Twelfth Congress of the International Association for Caribbean Archaeology, 241–58.

Picanço-Rodrigues, Doriane, Spartaco Astolfi-Filho, Maristerra R Lemes, Rogerio Gribel,

Alexandre M Sebbenn, and Charles R Clement. 2015. "Conservation Implications of the

Mating System of the Pampa Hermosa Landrace of Peach Palm Analyzed with

Microsatellite Markers." *Genetics and Molecular Biology*. scielo.

Pinhero, Reena Grittle, Kurt C. Almquist, Zuzana Novotna, and Gopinadhan Paliyath. 2003.

"Developmental Regulation of Phospholipase D in Tomato Fruits." *Plant Physiology and*

Biochemistry 41 (3): 223–40. https://doi.org/10.1016/S0981-9428(03)00014-7.

Puerari, Cláudia, Karina Teixeira Magalhães-Guedes, and Rosane Freitas Schwan. 2015.

"Bacaba Beverage Produced by Umutina Brazilian Amerindians: Microbiological and

Chemical Characterization." *Brazilian Journal of Microbiology* 46 (4): 1207–16.

https://doi.org/10.1590/S1517-838246420140964.

Qiu, Jie, Yongjun Zhou, Lingfeng Mao, Chuyu Ye, Weidi Wang, Jianping Zhang, Yongyi Yu,

et al. 2017. "Genomic Variation Associated with Local Adaptation of Weedy Rice

during De-Domestication." *Nature Communications* 8 (May): 15323.

https://doi.org/10.1038/ncomms15323.

R Core Team, R Core Team. 2021. "R: A Language and Environment for Statistical

Computing." In *R Foundation for Statistical Computing, Vienna, Austria*. https://www.r-

project.org/.

Reich, David, Kumarasamy Thangaraj, Nick Patterson, Alkes L. Price, and Lalji Singh. 2009.

"Reconstructing Indian Population History." *Nature* 461 (7263): 489–94.

https://doi.org/10.1038/nature08365.

Rendón-Anaya, Martha, Josaphat M. Montero-Vargas, Soledad Saburido-Álvarez, Anna

Vlasova, Salvador Capella-Gutierrez, José Juan Ordaz-Ortiz, O. Mario Aguilar, et al.

2017. "Genomic History of the Origin and Domestication of Common Bean Unveils Its

Closest Sister Species." *Genome Biology* 18 (1): 60. https://doi.org/10.1186/s13059-017-

1190-6.

Rival, Laura, and Doyle McKey. 2008. "Domestication and Diversity in Manioc (*Manihot*

Esculenta Crantz Ssp. *Esculenta*, Euphorbiaceae).” *Current Anthropology* 49 (6): 1119–

28. https://doi.org/10.1086/593119.

Rohland, Nadin, and David Reich. 2012. “Cost-Effective, High-Throughput DNA Sequencing

Libraries for Multiplexed Target Capture.” *Genome Research* 22 (5): 939–46.

https://doi.org/10.1101/gr.128124.111.

Rothhammer, Francisco, and Tom D. Dillehay. 2009. “The Late Pleistocene Colonization of

South America: An Interdisciplinary Perspective.” *Annals of Human Genetics*. Ann Hum

Genet. https://doi.org/10.1111/j.1469-1809.2009.00537.x.

Shepard, Glenn H., and Henri Ramirez. 2011. “‘Made in Brazil’: Human Dispersal of the

Brazil Nut (*Bertholletia Excelsa*, Lecythidaceae) in Ancient Amazonia.” *Economic

Botany* 65 (1): 44–65. https://doi.org/10.1007/s12231-011-9151-6.

Smith, Brian Tilston, John E. McCormack, Andrés M. Cuervo, Michael. J. Hickerson,

Alexandre Aleixo, Carlos Daniel Cadena, Jorge Pérez-Emán, et al. 2014. “The Drivers of

Tropical Speciation.” *Nature* 515 (7527): 406–9. https://doi.org/10.1038/nature13687.

Smith, Nigel. 2015. *Palms and People in the Amazon*. Geobotany Studies. Cham: Springer

International Publishing. https://doi.org/10.1007/978-3-319-05509-1.

Smith, Oliver, William V. Nicholson, Logan Kistler, Emma Mace, Alan Clapham, Pamela

Rose, Chris Stevens, et al. 2019. “A Domestication History of Dynamic Adaptation and

Genomic Deterioration in *Sorghum*.” *Nature Plants* 5 (4): 369–79.

https://doi.org/10.1038/s41477-019-0397-9.

Sosnowska, Joanna, Adam Walanus, and Henrik Balslev. 2015. “Asháninka Palm

Management and Domestication in the Peruvian Amazon.” *Human Ecology* 43 (3): 451–

66. https://doi.org/10.1007/s10745-015-9745-1.

Steege, Hans ter, Nigel C A Pitman, Daniel Sabatier, Christopher Baraloto, Rafael P Salomão,

Juan Ernesto Guevara, Oliver L Phillips, et al. 2013. "Hyperdominance in the

Amazonian Tree Flora." *Science* 342 (6156).

Sun, Jian, Xiangrong You, Li Li, Hongxiang Peng, Weiqiang Su, Changbao Li, Quanguang

He, and Fen Liao. 2011. "Effects of a Phospholipase D Inhibitor on Postharvest

Enzymatic Browning and Oxidative Stress of Litchi Fruit." *Postharvest Biology and*

Technology 62 (3): 288–94. https://doi.org/10.1016/j.postharvbio.2011.07.001.

Thioulouse, Jean, Daniel Chessel, Sylvain Dolédec, and Jean Michel Olivier. 1997. "ADE-4:

A Multivariate Analysis and Graphical Display Software." *Statistics and Computing* 7

(1): 75–83. https://doi.org/10.1023/A:1018513530268.

Wallace, Alfred Russel. 1853. *Palm Trees of the Amazon and Their Uses*. J. van Voorst.

Watling, Jennifer, José Iriarte, Francis E. Mayle, Denise Schaan, Luiz C.R. Pessenda, Neil J.

Loader, F. Alayne Street-Perrott, Ruth E. Dickau, Antonia Damasceno, and Alceu Ranzi.

2017. "Impact of Pre-Columbian Geoglyph Builders on Amazonian Forests."

Proceedings of the National Academy of Sciences of the United States of America 114

(8): 1868–73. https://doi.org/10.1073/pnas.1614359114.

Yazaki, Kazufumi. 2006. "ABC Transporters Involved in the Transport of Plant Secondary

Metabolites." *FEBS Letters*. No longer published by Elsevier.

https://doi.org/10.1016/j.febslet.2005.12.009.

Yin, Yuxin, Xiaowei Zhang, Yongjun Fang, Linlin Pan, Gaoyuan Sun, Chengqi Xin,

Mohammed M.Ba Abdullah, et al. 2012. "High-Throughput Sequencing-Based Gene

Profiling on Multi-Staged Fruit Development of Date Palm (*Phoenix Dactylifera*, L.)."

Plant Molecular Biology 78 (6): 617–26. https://doi.org/10.1007/s11103-012-9890-5.

Yoshida, K., P. Kaothien, T. Matsui, A. Kawaoka, and A. Shinmyo. 2003. "Molecular

Biology and Application of Plant Peroxidase Genes." *Applied Microbiology and*

Biotechnology. Springer Verlag. https://doi.org/10.1007/s00253-002-1157-7.

1065