Oncogenic metabolic rewiring independent of proliferative control in human mammary epithelial cells

Wentao Dong[1,2], Mark A. Keibler[1], Sun Jin Moon[1], Patricia Cho[3], Nian Liu[1], Christian J. Berrios[4,5], Joanne K. Kelleher[1], Hadley D. Sikes[1], Othon Iliopoulos[6,7], Jonathan L. Coloff[3,8], Matthew G. Vander Heiden[9] and Gregory Stephanopoulos[1,*]

1. Department of Chemical Engineering, Massachusetts Institute of Technology, Cambridge, MA, 02139, USA
2. Current address: Department of Chemical Engineering and the Institute for Chemistry, Engineering & Medicine for Human Health (ChEM-H), Stanford University, Stanford, CA, 94205, USA
3. Department of Cell Biology, Harvard Medical School, Boston, MA, 02115, USA
4. Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA, 02115, USA
5. Program in Virology, Harvard University, Boston, MA, 02115, USA
6. Center for Cancer Research, Massachusetts General Hospital Cancer Center, Charlestown, MA, 02129, USA
7. Division of Hematology/Oncology, Department of Medicine, Massachusetts General Hospital, Boston, MA, 02114, USA

8. Current address: Department of Physiology and Biophysics, College of Medicine, University

of Illinois at Chicago, Chicago, IL, 60612, USA

9. Koch Institute for Integrative Cancer Research and Department of Biology, Massachusetts

Institute of Technology, Cambridge, MA, 02139, USA

[*] Corresponding author:

77 Massachusetts Avenue, Room 56-469C

Cambridge, MA, 02139, USA

Phone: (617)-253-4583

Fax: (617)-253-3122

Email: gregstep@mit.edu

Summary

The use of isotopic tracers and metabolic flux analysis (MFA) has unveiled a number of

metabolic pathways differentially activated in cancer cells. To support efforts to design effective

metabolic therapies for cancer, we sought to distinguish metabolic behavior in cancer versus

normal cells growing at the same rate. To this end, we performed ^{13}C-isotope tracing and MFA

in human mammary epithelial cells (HMECs) harboring different combinations of oncogenes. By

introducing a new quantity termed metabolic flux intensity, defined as pathway flux divided by

specific growth rate, we showed that metabolism is dually controlled by proliferation and

oncogenotypes. ^{13}C-MFA further revealed that oxidative pentose phosphate pathway (oxPPP),

malate dehydrogenase (MDH) and isocitrate dehydrogenase (IDH) were most enhanced in

cancerous HMECs. Drug targeting of these pathways selectively reduced growth in the

tumorigenic HMEC line. Our study provides direct evidence that metabolism of cancer cells is

different than that of normal proliferating cells.

Introduction

Metabolic rewiring is one of the hallmarks of cancer (Hanahan and Weinberg, 2011). Our

knowledge of such changes has now extended beyond the Warburg effect (Bensinger and

Christofk, 2012; DeBerardinis et al., 2007; Hsu and Sabatini, 2008; Keibler et al., 2021;

Koppenol et al., 2011; Liberti and Locasale, 2016; Tran et al., 2020), which refers to the

paradoxical phenomenon that cancer cells preferentially metabolize glucose to lactate regardless

of oxygen availability (Warburg, 1956). Over the past decades, various types of metabolic

alterations in cancer cells have been documented (Ahn et al., 2018; Diaz-Moralli et al., 2016;

Man et al., 2021; Metallo et al., 2012; Pacold et al., 2016; Tran et al., 2020), prompting extensive

investigation of the link between reprogrammed cell signaling pathways and rewired cellular

metabolism (Dang et al., 2008; Forcina et al., 2022; Gaglio et al., 2011; Isidor et al., 2021; Liu et

al., 2012; Son et al., 2013). In addition, drug targeting of certain metabolic pathways has been

demonstrated to be a promising cancer treatment strategy (Vander Heiden, 2011; Méndez-lucas

et al., 2020; Ngoi et al., 2020). Despite this progress, a fundamental question remains

unanswered: whether there is a difference in the metabolism of cancer cells and their normal

proliferating counterparts. This deficiency has hindered our understanding of cancer metabolism

and the efforts to develop effective cancer therapies targeting metabolism with reduced side

effects.

This question has not been directly addressed before. It has been speculated that rewired

metabolic behavior associated with cancer may be a simple manifestation of reprogrammed

cellular energetics required to accommodate deregulated cell growth (Hanahan and Weinberg,

2011; Schulze and Harris, 2012; Ward and Thompson, 2012). For example, the Warburg effect is

observed in both tumor and highly proliferative normal cells (Hanahan and Weinberg, 2011;

Vander Heiden and DeBerardinis, 2017; Vander Heiden et al., 2009). Several types of

proliferating immune cells and stem cells rely on the glycolytic phenotype, a metabolic hallmark

of cancer, to maintain their respective functionality and lineage (Agathocleous and Harris, 2013;

Ananieva, 2015; Chen et al., 2017; Folmes et al., 2012; Vacanti and Metallo, 2013). Furthermore,

both cancer and normal proliferating cells may selectively express the M2 isoform of pyruvate

kinase, which is capable of promoting anabolism and cellular growth (Christofk et al., 2008;

Vander Heiden et al., 2011). Reductive carboxylation of glutamine has also been reported in both

proliferating endothelial cells (Kim et al., 2017) and several types of cancer cells (Metallo et al.,

2012; Mullen et al., 2012), suggesting that rewired glutamine metabolism is not unique to cancer.

However, the notion that metabolic behavior in cancer and normal proliferative cells is

equivalent is still questionable. For example, comparison between normal proliferative and

tumorigenic liver cells revealed that the latter exhibited upregulated glutaminase and

transaminase activities (Méndez-lucas et al., 2020). In addition, copy number amplification of

phosphoglycerate dehydrogenase was reported in melanoma relative to normal tissue

counterparts (Beroukhim et al., 2010; Locasale et al., 2011). Moreover, increased expression of

proline oxidase has been observed in pancreatic ductal adenocarcinoma compared to normal

pancreatic cells (Olivares et al., 2017). Although these findings suggest that cancer metabolism

may not be simply regarded as rewired proliferative energetics, the evidence is still inconclusive:

the proliferation rate of cancer cells in these studies was different than that of the normal cells.

Therefore, the metabolic alteration due to proliferation introduces a confounding factor in the

observed metabolic shifts. This challenge limits our ability to conclusively determine whether

metabolism of cancer cells is truly different than that of normal proliferating cells.

In order to address this question, we developed a panel of HMECs from the same genetic

background yet harboring different combinations of oncogenic manipulations. Through the use

of ^{13}C-MFA and the introduction of a new quantity referred to as metabolic flux intensity (MFI),

we decoupled proliferative control of metabolism and removed cellular growth as a factor

affecting metabolic rewiring. In addition, ^{13}C-isotopic labeling analysis and MFA also revealed a

distinct metabolic pattern within the tricarboxylic acid (TCA) cycle and lipogenesis. Our results

suggest that metabolism is directly controlled by both oncogenotypes and proliferation. The

proliferation-independent metabolic shifts identified in our work support the effort to design

cancer therapies that target specific metabolic pathways related with tumors only and not

affecting growth of healthy cells.

Results

Construction of HMECs with different combinations of oncogenes

To compare cancer and proliferative metabolism, we developed a panel of HMECs harboring

different oncogenes. The resulting HMECs share the same genetic background from the parental

cell: an immortalized, yet non-cancerous, HMEC 184A1 (ATCC), the growth of which is

dependent on the amount of epidermal growth factor (EGF) present in media. We then

overexpressed defined oncogenes and dominant negative mutant forms of tumor suppressors in

the original HMEC 184A1.

Based on the characteristic properties of cancer cells (Hanahan and Weinberg, 2000), we

considered three key attributes relevant to our specific *in vitro* system during cell line

development: growth factor independence, apoptotic evasion and limitless replicative potential.

We then generated several genetically modified HMECs carrying different oncogenes (Figure

1a). The constitutively active mutant form of EGF receptor (EGFR L858R) and the oncogenic K-

ras (KRas) were stably integrated into HMECs to confer growth factor-independent proliferation

(Schlaeth et al., 2010), and the resulting modified cell lines are referred to as HMEC-EGFR and

HMEC-KRas, respectively. To promote apoptotic evasion (Polyak et al., 1997), we introduced a

dominant negative mutant form of p53 (p53DD) to HMEC-KRas and generated HMEC-p53DD-

KRas. Moreover, we developed another cell line harboring Simian Virus 40 Early Region

(SV40ER), which also attenuates apoptosis due to the inhibitory role of SV40 large T antigen

(LT) on p53. We refer to the resulting cell line developed from HMEC-KRas as HMEC-

SV40ER-KRas. The successful expression of these genetic elements was validated by Western

blot (Figure S1). In addition, we also obtained an HMEC line carrying Simian Virus 40 large T

antigen (SV40-LT), oncogenic H-ras (HRas) and human telomerase reverse transcriptase

(hTERT) (Elenbaas et al., 2001). In this cell line, HRas and SV40-LT are able to induce growth

factor-independent proliferation and evasion of apoptosis (Shen and Shenk, 1998), respectively.

The introduction of hTERT further enhances cellular replicative potential, so the resulting cell

line HMEC-hTERT-LT-HRas possesses all three characteristic properties of cancer (Hanahan

and Weinberg, 2011) relevant to our *in vitro* system (Figure 1a).

Before characterizing the metabolic behavior of these cell lines, we first performed

proliferation assays. Parental and empty vector-expressing HMECs exhibited exponential growth

and were sensitive to EGF (Figure 1b). Based on exponential cell growth, we calculated specific

growth rates at different EGF concentrations. EV cells proliferated faster at higher EGF

concentrations, and the maximum specific growth rate was achieved at 5 ng/mL EGF (Figure 1c).

We next performed proliferation assays in oncogene-containing HMECs with 5 ng/mL or no

EGF (+/- EGF). Upon the integration of EGFR, cells began to acquire a small extent of growth

factor independence, which means that they were able to actively proliferate in the absence of

EGF. The uncontrolled proliferation was further enhanced in HMECs carrying KRas, p53DD-

KRas, SV40ER-KRas and hTERT-LT-HRas. Accordingly, the panel of HMECs was ranked by

increasing growth factor independence in Figure 1d. In addition, HMECs in this order also

exhibited increasing oncogene multiplicity (including downregulation of tumor suppressor

genes), with EV carrying no oncogene, EGFR and KRas harboring one, p53DD-KRas and

SV40ER-KRas carrying two, and hTERT-LT-HRas possessing three oncogenic elements.

Furthermore, previous work (Elenbaas et al., 2001) has validated that HMEC-hTERT-LT-HRas

is capable of undergoing anchorage-independent growth and forming tumors in immunodeficient

mice, demonstrating the greatest level of tumorigenicity among all HMECs (Figure 1d). These

HMECs with the same genetic background and defined oncogenic modifications enabled us to

investigate whether cancer metabolism differs from normal proliferative metabolism in a well-

controlled manner.

Quantification of extracellular fluxes suggests dual control of metabolism by proliferation and

oncogenotypes

We examined extracellular fluxes of glucose, lactate, glutamine and glutamate in our HMECs

(Murphy and Young, 2013). The extracellular fluxes followed a similar trend as the proliferation

rate: the difference between +/- EGF conditions was smaller in cell lines harboring more

oncogenes (Figure 2a). We further calculated lactate yield on glucose and glutamate yield on

glutamine by dividing the respective extracellular fluxes and accounting for the stoichiometry of

converting glucose to lactate and glutamine to glutamate (Figure S2a). Interestingly, although the

glutamate yield from glutamine exhibited a similar pattern as the extracellular fluxes, the lactate

yield from glucose was not significantly different at varying oncogenotypes or in the presence or

absence of EGF (Figure S2a). The only exception was the HMEC-EV -EGF condition, which

showed a relatively small lactate yield. The invariant yield of lactate suggests the strong presence

of the Warburg effect while cells are actively proliferating (Hanahan and Weinberg, 2011;

Vander Heiden and DeBerardinis, 2017; Vander Heiden et al., 2009).

Since the pattern of differential extracellular fluxes between +/- EGF conditions (Figure

2a) parallels that of growth (Figure 1 d), we sought to investigate whether growth rate regulates

extracellular fluxes. To this end, we plotted extracellular fluxes against specific growth rates. All

major extracellular fluxes (Figure 2b) and glutamate yield (Figure S2b) were higher at elevated

growth, except for lactate yield, which stayed constant again presumably because of the Warburg

effect (Figure S2b). Remarkably, the extracellular fluxes from both +/- EGF conditions tend to

collapse onto a single regression line, which suggests the presence of direct control of

extracellular fluxes by proliferation (Figure 2b). However, although the entire data set can be

described by regression lines to some extent, a closer examination of the data points with

comparable specific growth rates (0.02-0.025 h^{-1}) suggests that extracellular fluxes are not solely

determined by proliferation (Figure 2b). In fact, we constructed separate plots for HMECs grown

at the +EGF condition (Figure 2c). The apparent proliferative control of extracellular fluxes, as

indicated by the R^2 and regression p-values, were reduced and increased respectively when

considering only the +EGF condition (Figure 2c) compared to those in both +/-EGF conditions

(Figure 2b and S3). Collectively, these data suggest that metabolism, at least manifested by

extracellular fluxes, may be dually controlled by proliferation and specific oncogenotypes. In

addition, proliferative control of metabolism was reduced in +EGF HMECs that shared similar

growth rates.

<u>^{13}C-isotopic labeling reveals distinct substrate utilization patterns in the TCA cycle and *de novo*</u>

<u>lipogenesis</u>

To determine further metabolic changes, we performed ^{13}C-isotopic labeling experiments by

culturing HMECs in media separately containing three different tracers: uniformly ^{13}C-labeled

glucose (U-^{13}C$_6$-glucose); 1,2-^{13}C-labeled glucose (1,2-^{13}C$_2$-glucose); and uniformly ^{13}C-labeled

glutamine (U-^{13}C$_5$-glutamine). Previous studies have shown that these three tracers give the best

resolution of fluxes for glycolysis, PPP and the TCA cycle (Metallo et al., 2009). Measurement

of isotope enrichment of TCA cycle intermediates and amino acids revealed a distinct substrate

utilization pattern in all HMECs used in our study. The canonical view of anaplerosis predicts

strong enrichment of all TCA cycle intermediates by U-^{13}C$_6$-glucose. However, we observed a

low level of enrichment for all U-^{13}C$_6$-glucose-derived metabolites of the TCA cycle. In fact, the

majority of TCA cycle metabolites were weakly labeled by U-^{13}C$_6$-glucose, except for citrate,

which was mainly labeled by two ^{13}C atoms (M+2). This labeling pattern indicated that the ^{13}C

influx from glucose stopped at the point of citrate (Figure 3a). This conclusion was further

corroborated by 1,2-^{13}C$_2$-glucose labeling, which generated similar profiles compared to those

from U-^{13}C$_6$-glucose, consistent with the notion that M+2 citrate was the only metabolite

moderately labeled by glucose tracers (Figure S4).

Intrigued by this stark distribution pattern of glucose-derived carbons, we further labeled

cells with U-^{13}C$_5$-glutamine, hypothesizing that glutamine may be the primary carbon substrate

fueling the TCA cycle. The crucial role of glutamine anaplerosis has been reported in various

studies. In several types of cancer cells grown under hypoxia, glutamine provides carbon sources

for *de novo* lipogenesis through the reversal of isocitrate dehydrogenase (IDH) (Gameiro et al.,

2013; Metallo et al., 2012; Yoo et al., 2008). In contrast to reductive metabolism, in our HMECs,

glutamine was primarily incorporated into the oxidative TCA cycle. Mass isotopomer

distribution (MID) data showed that TCA intermediates were labeled much more strongly by U-

$^{13}C_5$-glutamine than U-$^{13}C_6$-glucose. Across all cell lines, TCA cycle metabolites were

approximately 50% enriched to the highest isotopomer states (M+4/5) when cultures were

labeled by U-$^{13}C_5$-glutamine. Moreover, M+4 abundance was much greater than that of M+5 for

citrate, indicating that the primary route of glutamine anaplerosis was oxidative cycling rather

than reductive carboxylation (Dong et al., 2017). This data suggests that glutamine, but not

glucose, is the primary carbon source that fuels the oxidative TCA cycle (Figure 3b).

In addition to glutamine anaplerosis, we also investigated the metabolic fate of glucose-

derived carbons. Aside from citrate, TCA cycle metabolites were minimally labeled by U-$^{13}C_6$-

glucose (Figure 3a). This labeling pattern suggests that glucose-derived fluxes are diverted away

from the TCA cycle at the point of citrate. Because citrate initiates *de novo* lipogenesis, we

analyzed the nonpolar metabolites from cells cultured in ^{13}C-labeled substrates to identify the

metabolic fate of citrate. The MIDs of palmitate and stearate suggest that glucose, rather than

glutamine, was the main contributor to *de novo* fatty acid synthesis (Figure 3c-d). To further

quantify the fractional contributions of glucose and glutamine to acetyl-CoA, a two-carbon unit

recruited for *de novo* lipogenesis, we performed isotopomer spectral analysis (ISA) (Kelleher

and Nickol, 2015). ISA confirmed the lipogenic role of glucose relative to glutamine (Figure 3e

and S5). The labeling and ISA results suggest that glucose-derived carbons were mostly utilized

for *de novo* lipogenesis. Taken together, our [13]C-isotopic labeling results identified a distinct

substrate utilization pattern regarding the TCA cycle and *de novo* lipogenesis in HMECs. In all

cell lines that we studied, the TCA cycle functions in a truncated fashion: glutamine supplies the

majority of the carbon substrates for TCA cycle intermediates, while lipogenesis is maintained

primarily by glucose rather than glutamine (Figure 3f).

<u>Quantification of intracellular metabolism via [13]C-metabolic flux analysis ([13]C-MFA)</u>

<u>corroborates dual control of metabolism by proliferation and oncogenotypes</u>

In order to quantitatively investigate intracellular metabolism in HMECs, we performed [13]C-

metabolic flux analysis ([13]C-MFA). Briefly, [13]C-MFA estimates metabolic fluxes by fitting them

such as to most closely reproduce the measured metabolite isotopic enrichment patterns

(Antoniewicz et al., 2007) (Figure 4a). The metabolic network model constructed for our [13]C-

MFA consists of major biochemical reactions within central carbon metabolism (Figure S6 and

Table S1). [13]C-MFA was performed under pseudo-steady state hypothesis, which was difficult to

achieve in slowly growing cells. Accordingly, we report converged [13]C-MFA flux results for all

+EGF HMECs (Table S6-11) and those -EGF HMECs that still actively proliferated (KRas,

p53DD-KRas, SV40ER-KRas and hTERT-LT-HRas, Table S2-5).

We also performed several consistency tests to examine if model-estimated fluxes from

the totality of labeling data were in agreement with local labeling results generated from

specialized isotopic tracers. For example, culturing cells with $1,2$-[13]C_2-glucose allows estimation

of relative fluxes between glycolysis and oxPPP. In brief, the glycolytic pathway retains both [13]C

atoms from $1,2$-[13]C_2-glucose and therefore generates M+2 labeled lactate as the main product.

The oxidative branch of the PPP, on the other hand, involves a decarboxylation reaction which

results in a loss of a ^{13}C atom in the form of carbon dioxide, yielding M+1 labeled lactate

(Carpenter et al., 2014; Dong et al., 2020). Relative magnitude of glycolysis and oxPPP can

therefore be deduced by comparing the MIDs of M+2 against M+1 lactate. This can be compared

with the same ratio calculated from the MFA-estimated fluxes using different isotopic labels. In

accordance with the ^{13}C-labeling pattern indicating a significantly lower fraction of M+1 (0.03)

as opposed to M+2 lactate (0.45), ^{13}C-MFA results also suggested a minor diversion of fluxes

into oxPPP (50 pmol·h^{-1}·µg^{-1}) from glycolysis (2813 pmol·h^{-1}·µg^{-1}). Moreover, we cultured

HMECs with U-^{13}C$_5$-glutamine and U-^{13}C$_6$-glucose, and calculated fractional enrichments of

major TCA intermediates. The labeling results indicated that the TCA cycle was primarily

maintained by glutamine. Consistently, MFA also suggested stronger anaplerotic flux by

glutamine than that by glucose (Figure 4b).

It has been hypothesized that metabolic behavior in cancer and normal proliferative cells

can be regarded similarly (Hanahan and Weinberg, 2011; Schulze and Harris, 2012; Sun et al.,

2010; Ward and Thompson, 2012). To test this notion and investigate how proliferation affects

metabolism, we plotted major intracellular fluxes against specific growth rates in all HMECs

with converged results. The regression trend lines suggested that glycolysis and lactate excretion

increased as cells grew faster, consistent with the Warburg Effect (Figure 5). As expected, the

malic enzyme (ME) reaction was also enhanced, potentially regenerating NADPH for cellular

redox needs (DeBerardinis and Chandel, 2016; Fan et al., 2014; Jiang et al., 2013) (Figure 5).

Moreover, glutamine anaplerosis was enhanced when cells grew faster. This metabolic pattern

further pinpoints the vital role of glutamine metabolism in maintaining the TCA cycle (Figure 3f).

Additionally, ^{13}C-MFA results suggested that PDH, rather than pyruvate carboxylase (PC),

catalyzed the primary route of pyruvate entry into the TCA cycle (Figure S8), consistent with

what has been reported in other cell lines (DeBerardinis et al., 2007). Importantly, although the

overall trend lines suggested that metabolic fluxes were controlled by growth rate at some level,

we also noticed non-negligible deviations of certain fluxes from the regression lines (Figure 5).

Such deviations indicated that growth was not the only factor determining fluxes. In other words,

metabolism is affected by both proliferation and oncogenotypes.

<u>Normalizing fluxes against growth by introducing metabolic flux intensity (MFI)</u>

Our ^{13}C-MFA results (Figure 5) and extracellular fluxes (Figure 2) suggest that metabolism may

be dually controlled by both proliferation and oncogenotypes. Indeed, there may be two modes

of action through which metabolism is regulated: an indirect route through which oncogene-

altered proliferation affects metabolism and a direct oncogenic effect independent of

proliferative control (Figure S10).

The two modes of control were uncoupled by dividing metabolic fluxes by the specific

growth rate (Figure S10). We term the resulting quantity as Metabolic Flux Intensity (MFI).

When the MFI of a pathway increases under some conditions, this suggests that cells require a

higher flux along the pathway to sustain the same level of growth, whereas lower MFI suggests

that the corresponding metabolic pathway plays a less essential role in sustaining cellular growth.

Therefore, the MFI of a pathway is an indicator of how strongly cells rely on the pathway to

proliferate. By defining this new quantity, we can potentially assess the direct impact of

oncogenotypes on metabolism independent of proliferation.

We thus calculated MFIs for all major metabolic pathways and constructed the plots of

MFIs against specific growth rates (Figure 6). To isolate the impact of oncogenotype alone, we

chose to focus on the +EGF HMECs. In addition, MFIs were averaged for oncogene-containing

yet non-cancerous HMECs (EGFR, KRas, p53DD-KRas and SV40ER-KRas) to underscore the

overall trend. We noticed that MFIs of certain metabolic pathways were increased in more

cancerous lines, while other pathways remained unchanged (Figure 6). Interestingly, the

glycolytic pathway and the lactate dehydrogenase (LDH) reaction exhibited enhanced MFIs,

suggesting that the Warburg effect extends beyond the simple notion of proliferative

upregulation of aerobic glycolysis (Figure 6). Instead, cells with higher oncogenic potential may

need even greater glycolytic and LDH fluxes per growth rate to maintain proliferation.

Importantly, pathways with the greatest MFIs were oxPPP, malate dehydrogenase (MDH) and

isocitrate dehydrogenase (IDH). Based on the definition of MFI, these may be the most critical

reactions for sustaining growth in HMEC-hTERT-LT-HRas relative to the control cell line

(Figure 6).

<u>Assessing the therapeutic potential of targeting oxPPP, MDH and IDH</u>

Our MFI analysis identified oxPPP, MDH and IDH as the most enhanced reactions in HMECs

harboring hTERT-LT-HRas. Interestingly, these three pathways are essential sources

regenerating NADH and NADPH. OxPPP is one of the primary routes of NADPH synthesis,

which can also be replenished by ME (Figure 7a). However, ME exhibited statistically invariant

MFIs across cell lines, whereas the reliance on oxPPP was higher in more cancerous HMECs

(Figure 6). Additionally, MDH and IDH play an important role in maintaining the integrity of the

TCA cycle, which functions as a central hub for supplying ATP, NADH and amino-anabolic

precursors (Figure 7b). Therefore, inhibition of oxPPP, MDH and IDH may be selectively toxic

in cancerous HMECs.

To test these two strategies, we performed drug inhibition experiments for the

tumorigenic HMEC-hTERT-LT-HRas against the control line HMEC-EV. We used 6-

aminonicotinamide (6AN), LW6 and GSK864 to inhibit oxPPP (Tsouko et al., 2014), MDH

(Zhang et al., 2020) and IDH (Calvert et al., 2017), respectively. The fold changes of cell count

for drug-treated compared to DMSO-treated cells at different drug concentrations are shown in

Figure 7c. We found that these inhibitors were selectively more toxic in the tumorigenic HMEC.

Specifically, 0.5 μM 6AN, 100 μM LW6 or 10 μM GSK864 significantly reduced the

proliferation of HMEC-hTERT-LT-HRas compared to the control cell line. It is worth noting

that drug targeting of the metabolic pathways exhibiting statistically invariant MFIs might not be

able to achieve selective toxicity (Figure S13). Therefore, pathways with the greatest MFIs in the

tumorigenic line may serve as better drug targets. These experiments underscore the therapeutic

potentials of targeting oxPPP, MDH and IDH for selectively killing cancerous HMECs.

<u>Discussion</u>

We have demonstrated the use of ^{13}C-isotopic labeling and MFA to quantitatively resolve the

metabolic fluxes of HMECs with different combinations of oncogenes. Our data and analysis

provide direct evidence that the metabolisms of cancer and normal proliferative cells differ and

this can be the basis for identifying targets for selective cancer treatment. In addition, we found a

distinct substrate utilization pattern involving a truncated TCA cycle and glucose-supported

lipogenesis. Moreover, we introduced the concept of MFI to unveil proliferation-independent

oncogenic metabolic rewiring that helped identify pathways like oxPPP, MDH and IDH as new

drug targets with improved selective toxicity.

Our work further underscores the metabolic significance of glutamine anaplerosis. The

fact that glutamine is metabolized as one of the major respiratory substrates in cancer cells has

been previously reported (Moreadith and Lehninger, 1984; Parlo and Coleman, 1984; Piva and

McEvoy-Bowe, 1998). More recently, the important role of glutamine in supporting *de novo*

lipogenesis has been elucidated (Gameiro et al., 2013; Metallo et al., 2012; Mullen et al., 2012).

In contrast to these works, here we revealed another type of glutamine metabolism: a truncated

TCA cycle for glutamine oxidation with glucose mainly supporting the lipogenic pathway. It is

worth mentioning that a truncated TCA cycle has been reported in previous investigations. For

example, glutamine can be utilized to fuel the TCA cycle in cancer cell lines harboring

heterozygous IDH-1 mutations under hypoxia or mitochondrial inhibition (Grassian et al., 2014).

Moreover, cell lines exhibiting a truncated TCA cycle exhibited reduced proliferation due to

their inability to perform a fully functional reductive glutamine metabolism along IDH (Grassian

et al., 2014). In contrast, our results show that the presence of a truncated TCA cycle does not

undermine cellular growth in HMECs (Figure 1 b-d). Instead, further truncation was observed in

the cell lines that proliferated faster (Figure 3e), suggesting that this utilization pattern may be

regulated by proliferation. In line with our results, the truncated TCA cycle and glucose-

supported lipogenesis have also been reported in glioblastomas, and such rewired metabolism

does not impair cancer cell proliferation either (DeBerardinis et al., 2007). Importantly, our work

shows that the truncated TCA cycle is present in all HMECs being studied, indicating that this

metabolic pattern is not unique to cancer cells. Furthermore, we employed [13]C-MFA to fully

resolve all major metabolic pathways and quantified fluxes within the truncated TCA cycle and

de novo lipogenesis. This quantitative depiction provides opportunities to further investigate this

metabolic pattern in depth. For example, the mechanism by which this metabolic phenomenon

occurs is still unclear at this point and remains to be elucidated by future work. These rewired

energetics may expose potential metabolic vulnerabilities that can be exploited for therapeutic

intervention to treat cancer. This idea is in line with the therapeutic efforts to target both glucose

and glutamine metabolism for enhanced overall effectiveness of chemotherapy (Jin et al., 2016;

Tennant et al., 2010; Wise and Thompson, 2011).

Our quantitative flux results enabled us to address the question of whether cancer and

normal proliferative metabolisms are different. One of the hallmarks of cancer – the Warburg

effect – has been proposed to be shared by both cancer and proliferating cells (Hanahan and

Weinberg, 2011; Vander Heiden and DeBerardinis, 2017; Vander Heiden et al., 2009). However,

there has been no study directly investigating whether and how cancer metabolism is different

from proliferative metabolism beyond the Warburg effect. In fact, it would be difficult to

eliminate confounding factors such as different cell types and genetic backgrounds if common

existing cancer cell lines were used in such studies. These concerns prompted us to develop a

new panel of HMECs that shared the same original genetic background by modifying defined

genetic elements. More importantly, we believe that the main challenge to study the difference

between cancer and normal proliferative metabolism is to decouple the growth-independent

impact by oncogenotypes from that by altered growth phenotypes (Fritz and Fajas, 2010).

To decouple these two modes through which cancer metabolism is manifested, we

introduced a new quantity called metabolic flux intensity (MFI) obtained by dividing metabolic

fluxes by the specific growth rate. In this way, the effect of growth on metabolism is normalized

when one compares MFIs across different cell lines. MFI essentially serves as an indicator

gauging the importance of a certain metabolic pathway in sustaining cellular growth. MFIs are

different across our HMEC variants, which suggests that proliferation is not the only factor

governing metabolic behavior.

MFI analysis helps in selecting targets for treating cancer. Based on the definition of MFI,

metabolic pathways with higher MFIs are relied upon by the cell more heavily than those with

lower MFIs for supplying metabolites and energy for growth. Therefore, pathways with higher

MFIs can be therapeutically more relevant in exerting selective toxicity on cancer cells. MFI

analysis emphasizes the notion of selective toxicity because it compares flux changes

independent of proliferative control of metabolism. Therefore, MFI may be more suitable for

identifying potential drug targets.

Since oxPPP, MDH and IDH are responsible for producing NADPH and NADH (Moon

et al., 2020), cofactor regeneration may be one of the key constraints limiting cancer cell growth.

In line with this idea, inhibition of these reactions rendered a more drastic reduction of

proliferation in the tumorigenic HMEC line. Previous studies also reported similar promising

results (Fujii et al., 2016; Lee et al., 2013; Levis, 2013; Ramos-Montoya et al., 2006). We

believe that MFI analysis can be helpful in identifying potential drug targets in other types of

cancer.

Significance

In order to distinguish the metabolism of cancer and normal growing cells, we performed stable

isotope tracing and metabolic flux analysis in a panel of human mammary epithelial cells

(HMECs) harboring different oncogenes. We identified oxidative pentose phosphate pathway,

malate dehydrogenase and isocitrate dehydrogenase as the most activated metabolic reactions

and potential drug targets in the tumorigenic HMEC. Moreover, we introduced a new quantity

termed metabolic flux intensity to describe metabolic rewiring independent of proliferative

control. Our quantitative metabolic analysis provides direct evidence that metabolism is dually

controlled by proliferation and oncogenes, thus supporting efforts to develop effective cancer

treatment by selectively targeting metabolic pathways.

Acknowledgements

We thank W. Hahn, O. Gjoerup, J. Deangelo and W. Kim for generously granting us one of the

cell lines to study. We also thank J. Park, B. Woolston, B. Pereira, Z. Li, E. Lien, K. Abbott and

A. Muir for discussions. We would like to further thank X. Lian and X. Bao for the suggestions

on manuscript revision. The authors are supported by NIH R01 CA160458.

Author contributions

Conceptualization, W.D., M.A.K. and G.S.; Methodology, W.D., M.A.K. and G.S.; Software,

W.D., M.A.K. and N.L.; Validation, W.D., P.C. and J.L.C.; Formal analysis, W.D., M.A.K.,

S.J.M. and N.L.; Investigation, W.D. M.A.K. and S.J.M.; Resources, W.D., P.C., J.L.C., C.B.

and G.S.; Writing – Original Draft, W.D. M.A.K. and P.C.; Writing – Review and Editing, W.D.,

M.A.K., S.J.M., P.C., J.L.C., N.L., J.K.K., M.G.V.H., O.I., H.D.S. and G.S., Supervision, J.K.K.,

M.G.V.H., O.I., H.D.S. and G.S.; Project Administration, G.S.; Funding Acquisition, G.S.

Declaration of interests

<u>Figure titles and legends</u>

Figure 1. Development of a panel of HMECs with different combinations of oncogenes.

a, Overexpression of defined oncogenes and dominant negative mutant forms of tumor

suppressors in HMECs. Ovals indicate HEMC lines. Rounded rectangles indicate cancer-

associated characteristics achieved during cell line development. Pointed and blunt-end arrows

within the signaling pathway denote activation and inhibition, respectively.

b, Proliferation of the normal HMECs in response to different EGF concentrations.

c, Specific growth rate of the normal HMECs as a function of EGF concentration.

d, Growth factor independence and other properties observed in genetically modified HMECs.

Data are represented as mean ± SEM (n=3). **P<0.005. ***P<0.001.

Figure 2. Extracellular fluxes were determined by both specific growth rates and

oncogenotypes.

a, Extracellular fluxes of glucose, lactate, glutamine and glutamate for HMECs at +/- EGF

conditions.

b, Extracellular fluxes of +/- EGF HMECs plotted against specific growth rates.

c, Extracellular fluxes of only the +EGF HMECs plotted against specific growth rates.

Data are represented as mean ± SEM (n=3). *P<0.05. **P<0.005. ***P<0.001.

Figure 3. ^{13}C-isotopic labeling analysis revealed a truncated TCA cycle.

a, U-^{13}C$_6$-glucose labeling pattern suggests weak incorporations of glucose-derived carbons into

the TCA cycle except for citrate, the majority of which stays at M+2.

b, U-^{13}C$_5$-glutamine labeling pattern suggests much stronger incorporations of glutamine-derived

carbons into the TCA cycle.

c, U-^{13}C$_5$-glutamine labeling pattern suggests weak incorporations of glutamine-derived carbons

into *de novo* fatty acid synthesis.

d, U-^{13}C$_6$-glucose labeling pattern suggests stronger incorporations of glucose-derived carbons

into *de novo* fatty acid synthesis.

e, Isotopomer spectral analysis (ISA) confirmed that glucose dominated lipogenesis over

glutamine.

f, ^{13}C-isotopic labeling analysis revealed a distinct substrate utilization pattern within the TCA

cycle and *de novo* lipogenesis in contrast to the canonical view.

Columns along the axis of abscissas in each bar graph represent ^{13}C-isotopic labeling data in the

following order: EV -EGF, EGFR -EGF, KRas -EGF, p53DD-KRas -EGF, SV40ER-KRas -EGF,

hTERT-LT-HRas -EGF; EV +EGF, EGFR +EGF, KRas +EGF, p53DD-KRas +EGF, SV40ER-

KRas +EGF, hTERT-LT-HRas +EGF. Mass isotopomer distribution (MID) is shown in **a** and **b**.

Fractional enrichment is shown in **c** and **d**. Open and filled circles in the schematics indicate ^{12}C

an ^{13}C atoms within the metabolites, respectively. Acetyl-CoA.c and acetyl-CoA.m refer to

cytosolic and mitochondrial acetyl-CoA, respectively. Data are represented as mean ± SEM

(n=3). Abbreviations: UGC, U-^{13}C$_6$-glucose; UGN: U-^{13}C$_5$-glutamine; Glu, glutamate.

Figure 4. Introduction to ^{13}C-MFA and consistency tests against the labeling data.

a, Essential components and steps for ^{13}C-metabolic flux analysis (^{13}C-MFA). Based on a

detailed metabolic network model, ^{13}C-MFA determines absolute flux scales from extracellular

flux data. ^{13}C-isotopic labeling data is used to resolve intracellular flux distributions. The

numerical algorithm works through iterations, during which ^{13}C-MFA frequently checks the fit

of experimentally determined extracellular fluxes and ^{13}C-isotopic labeling data by values

simulated by hypothesized flux distributions. The converged results are shown at the center of

the schematic.

b, ^{13}C-MFA generated consistent results with the labeling data regarding the partition of fluxes

between glycolysis and oxPPP and the anaplerotic contributions from glucose and glutamine.

Flux results of +EGF HMEC-EV are shown here as an example. Unit of metabolic fluxes,

pmol·h^{-1}·μg^{-1}. Uncertainties are shown as 95% confidence intervals.

Data are represented as mean ± SEM (n=3). ***P<0.001.

Abbreviations: oxPPP, oxidative pentose phosphate pathway; LDH, lactate dehydrogenase; ME:

malic enzyme; MDH, malate dehydrogenase; PDH, pyruvate dehydrogenase; CS, citrate

synthase; IDH, isocitrate dehydrogenase; TCA cycle, tricarboxylic acid cycle; Glu, glutamate.

Figure 5. ^{13}C-MFA suggested that both intracellular and extracellular metabolisms may be

dually controlled by proliferation and oncogenotypes.

Data are represented as mean ± 95% confidence intervals. Abbreviations: oxPPP, oxidative

pentose phosphate pathway; LDH, lactate dehydrogenase; ME: malic enzyme; MDH, malate

dehydrogenase; PDH, pyruvate dehydrogenase; CS, citrate synthase; IDH, isocitrate

dehydrogenase; TCA cycle, tricarboxylic acid cycle; Glu, glutamate.

Figure 6. Quantitative ^{13}C-MFI analysis validated that oncogenotypes directly impact

metabolism independent of proliferative control.

Normalized MFI results for all +EGF HMECs. Data are represented as mean ± 95% confidence

intervals. *P<0.05. **P<0.005. ***P<0.001. Abbreviations: oxPPP, oxidative pentose phosphate

pathway; MDH, malate dehydrogenase; IDH, isocitrate dehydrogenase; LDH, lactate

dehydrogenase; ME: malic enzyme; PDH, pyruvate dehydrogenase; CS, citrate synthase; TCA

cycle, tricarboxylic acid cycle.

Figure 7. Therapeutic potentials of targeting oxPPP, MDH and IDH to selectively kill the tumorigenic HMEC as opposed to the normal proliferating counterpart.

a, ^{13}C-MFI analysis identified oxPPP as the most enhanced reaction in HMEC-hTERT-LT-HRas. Targeting oxPPP is expected to impair NADPH regeneration.

b, ^{13}C-MFI analysis identified MDH and IDH as one of the most enhanced reactions in HMEC-hTERT-LT-HRas. Targeting MDH and IDH is expected to impair NADH regeneration and the TCA cycle.

c, Drug targeting results for oxPPP, MDH and IDH.

Data are represented as mean ± 95% confidence intervals (**a**, **b**), and mean ± SEM (n=3) (**c**).

*P<0.05. **P<0.005. ***P<0.001.

Abbreviations: oxPPP, oxidative pentose phosphate pathway; ME: malic enzyme; MDH, malate dehydrogenase; IDH, isocitrate dehydrogenase; G6PD, glucose-6-phosphate dehydrogenase; 6AN, 6-aminonicotinamide.

<u>Materials and methods</u>

Cell culture conditions

All cell lines were obtained from ATCC, tested for mycoplasma and cultured at 37 ºC and 5%

CO_2. The MCDB 170 medium was used for regular subculture on 10-cm tissue-culture treated

polystyrene dishes (Corning). The complete formulation of the MCDB 170 medium includes a

mammary epithelial basal medium (MEBM) (Lonza) supplemented with 5 µg/mL insulin, 0.5

µg/ml hydrocortisone, 5 µg/mL transferrin, 0.07 mg/mL bovine pituitary extract (BPE)

(Hammond), 10 uL isoproterenol and 5 ng/mL EGF (Peprotech). Trypsin-EDTA (0.25%)

(Thermo Fisher) was used as the dissociation reagent to passage cells upon 70-80% confluence

for up to 4 passages. Fresh media (15 mL per plate) was replaced every 2-3 days. Cells were

plated and tested on 6-well plates (Corning) for the experiments determining growth rates and

extracellular fluxes. Dulbecco's Modified Eagle's Medium (DMEM) (without glucose,

glutamine and sodium pyruvate, Corning) was used as the basal medium (2.5 mL per well)

supplemented with other MCDB 170 components, 10 mM glucose, 4 mM glutamine, 0.1 mM

ethanolamine, 0.1 mM phosphoethanolamine and 10 mM HEPEs. All components were obtained

from Sigma unless otherwise noted.

Transfection/infection and drug selection for HMECs

HMEC 184A1 was used as the parental cell line to develop all HMEC variants, except for

HMEC-hTERT-LT-HRas, which was kindly provided by Dr. W. Hahn (Dana-Farber Cancer

Institute). An empty vector, K-ras G12V, EGFR L858R, p53DD and SV40ER (Addgene) were

stably integrated into HMEC 184A1 through γ-retroviral infections under the CMV promoter.

530 HEK 293 T cells at 40-60% confluence were transfected with 5.25 µg pBABE-puro EV,

531 pBABE-puro K-ras G12V, pBABE-puro EGFR L858R, pBABE-neo p53DD or pBABE-neo

532 SV40ER, 4.725 µg gag/pol vector, 0.525 µg VSV-G and 31.5 µL X-tremeGENE HP DNA

533 Transfection Reagent (Roche). All components were premixed and equilibrated for 15 min at 25

534 ºC before being transferred in a dropwise manner into HEK 293 T cells grown in a 10-cm dish

535 with 10 mL DMEM and 10 vol % fetal bovine serum (FBS) (Sigma SAFC). Media containing γ-

536 retroviruses was harvested after 36-48 h incubation at 37 ºC and 5% CO_2 and was passed through

537 0.45 µm filters (Pall). The retentate (~10 mL) containing γ-retroviruses was then mixed with 8

538 µg/mL Polybrene (Millipore) and added into HMECs at 20-30% confluence grown in a 10-cm

539 dish. After 24-36 hours, the spent media with retroviruses was aspirated and the plate was

540 washed by 10 mL PBS (Corning) for three times.

541  HMEC-EV, HMEC-EGFR and HMEC-KRas with successful integrations of retroviral

542 vectors were selected with 0.5 µg/mL puromycin. HMEC-p53DD-KRas and HMEC-SV40ER-

543 KRas were selected with 0.5 µg/mL puromycin and subsequently with 400 µg/mL G418. The

544 concentrations of these drugs were verified to be able to kill all mock-infected HMECs. To avoid

545 bias due to position effect, the final population of each cell line was composed of at least 25

546 independent clones. HMEC lines after drug selection were then cryopreserved by mixing 1.5-

547 $3×10^6$ cells with 1mL freezing media containing 75 vol % MEBM, 15 vol % FBS and 10 vol %

548 glycerol (Sigma).

Western blots

Cells were lysed in RIPA buffer (Thermo) containing protease and phosphatase inhibitor cocktails (Bimake B14001 and B15001-B). Protein concentration was quantified by BCA assay (Thermo Fisher). Equal amounts of protein were run on 4-20% Tris-Glycine gels (Invitrogen) and transferred to PVDF membranes. Membranes were blocked with 5% bovine serum albumin and probed with the following primary antibodies: β-actin (Sigma, Cat. No. A1978), EGFR (Thermo Fisher, Cat. No. MS-400-P1), ERK1/2 (Cell Signaling, Cat. No. 9102) phospho-ERK1/2 (Cell Signaling, Cat. No. 4370), p53 (BD Biosciences, Cat. No. 554293), SV40 T Ag (Santa Cruz, Cat. No. SC-147), α-Tubulin (Cell Signaling, Cat. No. 3873). Blots were imaged using Luminata Western HRP substrate (EMD Millipore).

Determinations of specific growth rates and extracellular fluxes

Protein concentrations for cells within 6-well plates were quantified by BCA assay (Thermo Pierce). Specific growth rates (μ) were determined by the following equation:

$$\mu = \frac{1}{t} ln \left(\frac{X_t}{X_0} \right)$$

where X_t and X_0 are the protein concentrations at time 0 and time t, respectively. The initial cell seeding densities were 80×10^3 and 35×10^3 per well for -EGF and +EGF conditions, respectively. MCDB 170 medium was used during seeding (D-3), and DMEM with MCDB 170 components at +/-EGF conditions was used after 24 hours during media change (D-2), preceded by a two-time wash with PBS. Spent media was collected and new media was replaced after another 48 hours following a one-time wash with 2mL/well PBS (D0). Another 48 hours later, the spent

media was harvested, and the cells (D2) were first washed once by 2 mL/well PBS. Then 500

µL/well of cold RIPA buffer (Thermo) was pipetted into each well, and the 6-well plates were

shaken briefly and placed at 4 ℃ for overnight incubation. Cell lysates were then harvested,

vortexed and centrifuged for the retrieval of supernatant. Protein concentration within each well

was then measured in 96-well plates through colorimetric detection and quantification according

to the standard protocol of BCA assay.

Concentrations of extracellular glucose, lactate, glutamine and glutamate in spent media

were then measured by Yellow Spring Instruments (YSI) 2950. Based on the equation governing

extracellular fluxes during the exponential growth phase, the extracellular fluxes for glucose,

lactate and glutamate were determined according to the following equation (Murphy and Young,

2013):

584
$$v = \frac{C_t - C_0}{e^{\mu t} - 1} \frac{\mu}{X_0}$$

where v is the extracellular flux of a metabolite, C_t the concentration of that metabolite at time t,

C_0 the concentration of that metabolite at time 0, μ the specific growth rate and X_0 the initial

protein content at time 0.

Due to the fact that glutamine is unstable and undergoes spontaneous degradation in

normal cell culture condition, the extracellular fluxes for glutamine were determined based on

the following equation with an additional term of decay constant k (Murphy and Young, 2013):

591
$$v = \frac{C_t e^{kt} - C_0}{e^{(\mu+k)t} - 1} \frac{\mu+k}{X_0}$$

592   where the first-order decay constant k was determined based on the following equation:

593
$$k = \frac{1}{t} \ln \left(\frac{C_{0,d}}{C_{t,d}} \right)$$

where $C_{t,d}$ and $C_{0,d}$ are the glutamine concentrations in media without cells at time t and time 0,

respectively.

^{13}C-isotopic labeling experiments and intracellular metabolite extraction

^{13}C-isotopic tracers (U-^{13}C$_6$-glucose, 1,2-^{13}C$_2$-glucose and U-^{13}C$_5$-glutamine, Cambridge Isotope

Laboratories) were administered at the same concentrations as the unlabeled counterparts (10

mM for the glucose tracers and 4 mM for the glutamine tracer). The tracers were used one at a

time at full enrichment. For example, 10 mM of U-^{13}C$_6$-glucose was used with 4 mM unlabeled

glutamine in the U-^{13}C$_6$-glucose tracing experiment. The media formulation is exactly the same

as that used in the growth and extracellular flux experiments. The experimental procedures are

the same as that for the growth and extracellular flux experiments until D2. Instead of harvesting

at D2, we prolonged the ^{13}C-istopic labeling duration to 72 hours (D3) after the second media

change (D0) to allow for more sufficient time to reach metabolic and isotopic steady states.

To harvest metabolites with minimal metabolic perturbations, we immediately placed the

6-well plates on ice once the plates were out of the incubator. Spent media was then quickly

aspirated, and each well was washed once by cold (0-4 ℃) saline, followed by addition of 500

μL freezing cold (-20 ℃) methanol. Next, 300 μL cold (0-4 ℃) milliQ water containing 2 μg

norvaline (Sigma) as an internal standard was pipetted to each well. Cells were then quickly

scrapped off from the wells by pipette tips in the presence of the liquid mixture, and the mixture

was transferred to microcentrifuge tubes. These steps were performed one at a time for each well.

Metabolite extraction was then performed by adding 600 μL freezing cold (-20 ℃) chloroform to

each microcentrifuge tube and then quickly vortexing the mixture at 4 ℃ for 10 min, followed by

a centrifugation at 21000×g for 10 min. After these steps, the mixture was separated into two

phases with the top and bottom layers containing polar and nonpolar metabolites, respectively.

The two layers were then retrieved and placed into separate microcentrifuge tubes, dried by air

and then stored at -80 ℃ for less than 1-2 weeks before being derivatized and loaded onto

GC/MS.

Metabolite derivatization and GC/MS analysis

Polar metabolites were derivatized by incubating dried samples in each microcentrifuge tube

with 15 µL methoxyamine in pyridine (MOX) (Thermo) at 40 ℃ for 1.5 hours, followed by

another incubation with 20 µL N-(tert-butyldimethylsilyl)-N-methyl-trifluoroacetamide with 1%

tert-Butyldimethylchlorosilane (TBDMS) (Sigma) at 60 ℃ for 1 hour. The derivatized mixture

was then briefly vortexed, centrifuged and transferred into polypropylene GC/MS vials (Agilent).

Nonpolar metabolites were treated by incubating dried samples in each microcentrifuge tube

with 500 µL methanol with 2 vol % sulfuric acid at 60 ℃ for 3 hours. Next, 600 µL hexane and

175 µL saturated sodium chloride solution were added, and the mixture was vortexed at room

temperature for 30 min and subsequently centrifuged at 21000×g for 1 min. The resulting

mixture was separated into two phases, and the top phase containing nonpolar metabolites was

retrieved into a separate microcentrifuge tube, dried by air, reconstituted by 30-50 µL hexane

and transferred into amber glass GC/MS vials with glass inserts (Agilent).

Agilent 6890N GC and 5975B Inert XL MS were used for polar metabolite analysis. The

column for the 6890N GC is Agilent J&W DB-35ms (35%-phenyl-methylpolysiloxane, mid-

polarity) and the electron ionization mode with 70 eV was used for the 5975B MS.

Chromatography grade helium (Airgas) at 1mL/min flow rate was used as the carrier gas. The

inlet temperature for the 6890N GC was set to 270 ºC, and the oven temperature was first

maintained at 100 ºC, and ramped to 300 ºC at a speed of 2.5 ºC/min. Samples of either 1 or 2 µL

were injected into the instrument with either split or splitless mode based on sample abundances.

The scan mode with a detection range of 150-625 m/z was used for all measurements. Mass

isotopomer distributions (MIDs) have been corrected for natural abundance.

Agilent 7890B GC and 5977B MS were used for nonpolar metabolite analysis. The

column for the 7890B GC is Agilent J&W HP-5ms (5%-phenyl-methylpolysiloxane, nonpolar)

and the electron ionization mode with 70 eV was used for the 5977B MS. Ultra high purity grade

helium (Airgas) at 3 mL/min was used as the carrier gas. The inlet temperature for the 7890B GC

was set to 280 ºC, and the oven temperature was first maintained at 165 ºC and then ramped to

226 °C at a speed of 2 ºC/min. Samples of 1 µL were injected into the instrument with the

splitless mode. The scan mode with a detection range of 200-400 m/z was used for all

measurements. MIDs have been corrected for natural abundance.

[13]C-metabolic flux analysis ([13]C-MFA) and isotopomer spectral analysis (ISA)

An elementary metabolite unit (EMU)-based software Metran coded within MATLAB

(MathWorks) was used to perform [13]C-MFA (Antoniewicz et al., 2007; Young et al., 2008) and

ISA (Kelleher and Nickol, 2015; Metallo et al., 2012). Experimentally determined extracellular

fluxes of glucose, lactate, glutamine and glutamate, as well as the [13]C-labeling data were used in

conjunction with a metabolic reaction model (Figure S3) to generate [13]C-MFA results. The 95%

confidence intervals for metabolic fluxes were obtained by performing parameter continuation

on converged flux results (Antoniewicz et al., 2006). The critical steps of ISA are explained in

Figure S5.

663   Inhibition of oxPPP, MDH and IDH

664   HMEC-hTERT-LT-HRas and HMEC-EV were seeded at 35×10^{3} per well with 5ng/mL EGF in

665   6-well plates. The small molecule drugs 6-aminonicotinamide (6AN), LW6 and GSK864 at

666   different concentrations were administered in the usual MEBM-based MCDB 170 medium.

667   DMSO was used to prepare the initial concentrated drug solutions. Cell counts were measured by

668   Cellometer (Nexcelom) 48 hours after seeding.

670   Definition of quantities

671   Fractional enrichments from ^{13}C-isotopic tracers were calculated using MID data corrected for

672   natural abundance based on the following equation:

$$673 \qquad\qquad \text{Fractional enrichment} = \sum_{i=0}^{n}(i \times m_i) / (n \times \sum_{i=0}^{n} m_i)$$

674   where n is the number of carbon atoms in a metabolite, m_i the abundance of a mass isotopomer

675   and i the labeling state (M+i) of a mass isotopomer.

676   Metabolic flux intensities (MFIs) were defined and calculated based on the following

677   equation:

$$678 \qquad\qquad \text{MFI} = \frac{v}{\mu}$$

where v and µ refer to metabolic flux (pmol·h^{-1}·µg^{-1}) and specific growth rate (h^{-1}), respectively.

The unit of MFI is therefore pmol·µg^{-1}.

References

Agathocleous, M., and Harris, W.A. (2013). Metabolism in physiological cell proliferation and

differentiation. Trends Cell Biol. *23*, 484–492.

Ahn, W.S., Dong, W., Zhang, Z., Cantor, J.R., Sabatini, D.M., Iliopoulos, O., and

Stephanopoulos, G. (2018). Glyceraldehyde 3-phosphate dehydrogenase modulates nonoxidative

pentose phosphate pathway to provide anabolic precursors in hypoxic tumor cells. AIChE J. *64*,

4289–4296.

Ananieva, E. (2015). Targeting amino acid metabolism in cancer growth and anti-tumor immune

response. World J. Biol. Chem. *6*, 281–289.

Antoniewicz, M.R., Kelleher, J.K., and Stephanopoulos, G. (2006). Determination of confidence

intervals of metabolic fluxes estimated from stable isotope measurements. Metab. Eng. *8*, 324–

337.

Antoniewicz, M.R., Kelleher, J.K., and Stephanopoulos, G. (2007). Elementary metabolite units

(EMU): A novel framework for modeling isotopic distributions. Metab. Eng. *9*, 68–86.

Bensinger, S.J., and Christofk, H.R. (2012). New aspects of the Warburg effect in cancer cell

biology. Semin. Cell Dev. Biol. *23*, 352–361.

Beroukhim, R., Mermel, C.H., Porter, D., Wei, G., Raychaudhuri, S., Donovan, J., Barretina, J.,

Boehm, J.S., Dobson, J., Urashima, M., et al. (2010). The landscape of somatic copy-number

alteration across human cancers. Nature *463*, 899–905.

Calvert, A.E., Chalastanis, A., Wu, Y., Hurley, L.A., Kouri, F.M., Bi, Y., Kachman, M., May,

J.L., Bartom, E., Hua, Y., et al. (2017). Cancer-Associated IDH1 Promotes Growth and

Resistance to Targeted Therapies in the Absence of Mutation. Cell Rep. *19*, 1858–1873.

Carpenter, K.L.H., Jalloh, I., Gallagher, C.N., Grice, P., Howe, D.J., Mason, A., Timofeev, I.,

Helmy, A., Murphy, M.P., Menon, D.K., et al. (2014). 13C-labelled microdialysis studies of

cerebral metabolism in TBI patients. Eur. J. Pharm. Sci. *57*, 87–97.

Chen, Z., Liu, M., Li, L., and Chen, L. (2017). Involvement of the Warburg effect in non-tumor

diseases processes. J. Cell. Physiol. *233*, 2839–2849.

Christofk, H.R., Vander Heiden, M.G., Harris, M.H., Ramanathan, A., Gerszten, R.E., Wei, R.,

Fleming, M.D., Schreiber, S.L., and Cantley, L.C. (2008). The M2 splice isoform of pyruvate

kinase is important for cancer metabolism and tumour growth. Nature *452*, 230–233.

Dang, C. V, Kim, J., Gao, P., and Yustein, J. (2008). The interplay between MYC and HIF in

cancer. Nat. Rev. Cancer *8*, 51–56.

DeBerardinis, R.J., and Chandel, N.S. (2016). Fundamentals of cancer metabolism. Sci. Adv. *2*,

1–18.

DeBerardinis, R.J., Mancuso, A., Daikhin, E., Nissim, I., Yudkoff, M., Wehrli, S., and

Thompson, C.B. (2007). Beyond aerobic glycolysis: transformed cells can engage in glutamine

metabolism that exceeds the requirement for protein and nucleotide synthesis. Proc. Natl. Acad.

Sci. U. S. A. *104*, 19345–19350.

Diaz-Moralli, S., Aguilar, E., Marin, S., Coy, J.F., Dewerchin, M., Antoniewicz, M.R., Meca-

Cortés, O., Notebaert, L., Ghesquière, B., Eelen, G., et al. (2016). A key role for transketolase-

like 1 in tumor metabolic reprogramming. Oncotarget *7*, 51875–51897.

Dong, W., Keibler, M.A., and Stephanopoulos, G. (2017). Review of metabolic pathways

activated in cancer cells as determined through isotopic labeling and network analysis. Metab.

Eng. *43*, 113–124.

Dong, W., Moon, S.J., Kelleher, J.K., and Stephanopoulos, G. (2020). Dissecting Mammalian

Cell Metabolism through 13C- And 2H-Isotope Tracing: Interpretations at the Molecular and

Systems Levels. Ind. Eng. Chem. Res. *59*, 2593–2610.

Elenbaas, B., Spirio, L., Koerner, F., Fleming, M.D., Zimonjic, D.B., Donaher, J.L., Popescu,

N.C., Hahn, W.C., and Weinberg, R.A. (2001). Human breast cancer cells generated by

oncogenic transformation of primary mammary epithelial cells. Genes Dev. *15*, 50–65.

Fan, J., Ye, J., Kamphorst, J.J., Shlomi, T., Thompson, C.B., and Rabinowitz, J.D. (2014).

Quantitative flux analysis reveals folate-dependent NADPH production. Nature *10*, 298–302.

Folmes, C.D.L., Dzeja, P.P., Nelson, T.J., and Terzic, A. (2012). Metabolic plasticity in stem cell

homeostasis and differentiation. Cell Stem Cell *11*, 596–606.

Forcina, G.C., Pope, L., Murray, M., Dong, W., Abu-Remaileh, M., Bertozzi, C.R., and Dixon,

S.J. (2022). Ferroptosis Regulation by the NGLY1/NFE2L1 Pathway. Proc. Natl. Acad. Sci. U. S.

A. *119*, 1–11.

Fritz, V., and Fajas, L. (2010). Metabolism and proliferation share common regulatory pathways

in cancer cells. Oncogene *29*, 4369–4377.

Fujii, T., Khawaja, M.R., DiNardo, C.D., Atkins, J.T., and Janku, F. (2016). Targeting Isocitrate

Dehydrogenase (IDH) in cancer. Discov. Med. *21*, 373–380.

Gaglio, D., Metallo, C.M., Gameiro, P.A., Hiller, K., Danna, L.S., Balestrieri, C., Alberghina, L.,

Stephanopoulos, G., and Chiaradonna, F. (2011). Oncogenic K-Ras decouples glucose and

glutamine metabolism to support cancer cell growth. Mol. Syst. Biol. *7*, 1–15.

Gameiro, P.A., Yang, J., Metelo, A.M., Pérez-Carro, R., Baker, R., Wang, Z., Arreola, A.,

Rathmell, W.K., Olumi, A., López-Larrubia, P., et al. (2013). In vivo HIF-mediated reductive

carboxylation is regulated by citrate levels and sensitizes VHL-deficient cells to glutamine

deprivation. Cell Metab. *17*, 372–385.

Grassian, A.R., Parker, S.J., Davidson, S.M., Divakaruni, A.S., Green, C.R., Zhang, X., Slocum,

K.L., Pu, M., Lin, F., Vickers, C., et al. (2014). IDH1 mutations alter citric acid cycle

metabolism and increase dependence on oxidative mitochondrial metabolism. Cancer Res. *74*,

3317–3331.

Hanahan, D., and Weinberg, R.A. (2000). The hallmarks of cancer. Cell *100*, 57–70.

Hanahan, D., and Weinberg, R.A. (2011). Hallmarks of cancer: The next generation. Cell *144*,

646–674.

Vander Heiden, M.G. (2011). Targeting cancer metabolism: A therapeutic window opens. Nat.

Rev. Drug Discov. *10*, 671–684.

Vander Heiden, M.G., and DeBerardinis, R.J. (2017). Understanding the Intersections between

Metabolism and Cancer Biology. Cell *168*, 657–669.

Vander Heiden, M.G., Cantley, L.C., Thompson, C.B., Mammalian, P., Exhibit, C., and

Metabolism, A. (2009). Understanding the Warburg Effect : Cell Proliferation. Science *324*,

1029–1034.

Vander Heiden, M.G., Lunt, S.Y., Dayton, T.L., Fiske, B.P., Israelsen, W.J., Mattaini, K.R.,

Vokes, N.I., Stephanopoulos, G., Cantley, L.C., Metallo, C.M., et al. (2011). Metabolic Pathway

Alterations that Support Cell Proliferation. Cold Spring Harb. Symp. Quant. Biol. *76*, 325–334.

Hsu, P.P., and Sabatini, D.M. (2008). Cancer cell metabolism: Warburg and beyond. Cell *134*,

703–707.

Isidor, M.S., Dong, W., Servin-Uribe, R.I., Villarroel, J., Altıntaş, A., Ayala-Sumuano, J.T.,

Varela-Echavarría, A., Barrès, R., Stephanopoulos, G., Macotela, Y., et al. (2021). Insulin

resistance rewires the metabolic gene program and glucose utilization in human white adipocytes.

Int. J. Obes. 1–9.

Jiang, P., Du, W., Mancuso, A., Wellen, K.E., and Yang, X. (2013). Reciprocal regulation of p53

and malic enzymes modulates metabolism and senescence. Nature *493*, 689–693.

Jin, L., Alesi, G.N., and Kang, S. (2016). Glutaminolysis as a target for cancer therapy.

Oncogene *35*, 3619–3625.

Keibler, M.A., Dong, W., Korthauer, K.D., Hosios, A.M., Moon, S.J., Sullivan, L.B., Liu, N.,

Abbott, K.L., Arevalo, O.D., Ho, K., et al. (2021). Differential substrate use in EGF- and

oncogenic KRAS-stimulated human mammary epithelial cells. FEBS J. 1–21.

Kelleher, J.K., and Nickol, G.B. (2015). Isotopomer Spectral Analysis: Utilizing Nonlinear

Models in Isotopic Flux Studies. Methods Enzymol. *561*, 303–330.

Kim, B., Li, J., Jang, C., and Arany, Z. (2017). Glutamine fuels proliferation but not migration of

endothelial cells. EMBO J. *36*, 2321–2333.

Koppenol, W.H., Bounds, P.L., and Dang, C. V. (2011). Otto Warburg's contributions to current

concepts of cancer metabolism. Nat. Rev. Cancer *11*, 325–337.

Lee, K., Ban, H.S., Naik, R., Hong, Y.S., Son, S., Kim, B.K., Xia, Y., Song, K. Bin, Lee, H.S.,

and Won, M. (2013). Identification of malate dehydrogenase 2 as a target protein of the HIF-1

inhibitor LW6 using chemical probes. Angew. Chem. Int. Ed. Engl. *52*, 10286–10289.

Levis, M. (2013). Targeting IDH: The next big thing in AML. Blood *122*, 2770–2771.

Liberti, M. V., and Locasale, J.W. (2016). The Warburg Effect: How Does it Benefit Cancer

Cells? Trends Biochem. Sci. *41*, 211–218.

Liu, W., Le, a., Hancock, C., Lane, a. N., Dang, C. V., Fan, T.W.-M., and Phang, J.M. (2012).

Reprogramming of proline and glutamine metabolism contributes to the proliferative and

metabolic responses regulated by oncogenic transcription factor c-MYC. Proc. Natl. Acad. Sci.

109, 8983–8988.

Locasale, J.W., Grassian, A.R., Melman, T., Lyssiotis, C.A., Mattaini, K.R., Bass, A.J., Heffron,

G., Metallo, C.M., Muranen, T., Sharfi, H., et al. (2011). Phosphoglycerate dehydrogenase

diverts glycolytic flux and contributes to oncogenesis. Nat. Genet. *43*, 869–874.

Man, C.H., Mercier, F.E., Liu, N., Dong, W., Stephanopoulos, G., Jiang, L., Jung, Y., Lin, C.,

Leung, A.Y., and Scadden, D.T. (2021). Proton export alkalinizes intracellular pH and

reprograms carbon metabolism to drive hematopoietic progenitor growth. Blood.

doi:10.1182/blood.2021011563.

Méndez-lucas, A., Lin, W., Driscoll, P.C., Legrave, N., Novellasdemunt, L., Xie, C., Charles, M.,

Wilson, Z., Jones, N.P., Rayport, S., et al. (2020). Identifying strategies to target the metabolic

flexibility of tumours. Nat. Metab. *2*, 335–350.

Metallo, C.M., Walther, J.L., and Stephanopoulos, G. (2009). Evaluation of 13C isotopic tracers

for metabolic flux analysis in mammalian cells. J. Biotechnol. *144*, 167–174.

Metallo, C.M., Gameiro, P.A., Bell, E.L., Mattaini, K.R., Yang, J., Hiller, K., Jewell, C.M.,

Johnson, Z.R., Irvine, D.J., Guarente, L., et al. (2012). Reductive glutamine metabolism by IDH1

mediates lipogenesis under hypoxia. Nature *481*, 380–384.

Moon, S.J., Dong, W., Stephanopoulos, G.N., and Sikes, H.D. (2020). Oxidative pentose

phosphate pathway and glucose anaplerosis support maintenance of mitochondrial NADPH pool

under mitochondrial oxidative stress. Bioeng. Transl. Med. *5*, 1–18.

Moreadith, R.W., and Lehninger, A.L. (1984). Purification, kinetic behavior, and regulation of

NAD(P)+ malic enzyme of tumor mitochondria. J. Biol. Chem. *259*, 6222–6227.

Mullen, A.R., Wheaton, W.W., Jin, E.S., Chen, P.-H., Sullivan, L.B., Cheng, T., Yang, Y.,

Linehan, W.M., Chandel, N.S., and DeBerardinis, R.J. (2012). Reductive carboxylation supports

growth in tumour cells with defective mitochondria. Nature *481*, 385–388.

Murphy, T.A., and Young, J.D. (2013). ETA: Robust software for determination of cell specific

rates from extracellular time courses. Biotechnol. Bioeng. *110,* 1748–1758.

Ngoi, N.Y.L., Eu, J.Q., Hirpara, J., Wang, L., Lim, J.S.J., Lee, S.C., Lim, Y.C., Pervaiz, S., Goh,

B.C., and Wong, A.L.A. (2020). Targeting Cell Metabolism as Cancer Therapy. Antioxidants

Redox Signal. *32*, 285–305.

Olivares, O., Mayers, J.R., Gouirand, V., Torrence, M.E., Gicquel, T., Borge, L., Lac, S., Roques,

J., Lavaut, M.N., Berthezène, P., et al. (2017). Collagen-derived proline promotes pancreatic

ductal adenocarcinoma cell survival under nutrient limited conditions. Nat. Commun. *8*, 1–14.

Pacold, M.E., Brimacombe, K.R., Chan, S.H., Rohde, J.M., Lewis, C.A., Swier, L.J.Y.M.,

Possemato, R., Chen, W.W., Sullivan, L.B., Fiske, B.P., et al. (2016). A PHGDH inhibitor

reveals coordination of serine synthesis and one-carbon unit fate. Nat. Chem. Biol. *12*, 452–458.

Parlo, R.A., and Coleman, P.S. (1984). Enhanced rate of citrate export from cholesterol-rich

hepatoma mitochondria. The truncated Krebs cycle and other metabolic ramifications of

mitochondrial membrane cholesterol. J. Biol. Chem. *259*, 9997–10003.

Piva, T.J., and McEvoy-Bowe, E. (1998). Oxidation of glutamine in hela cells: Role and control

of truncated TCA cycles in tumour mitochondria. J. Cell. Biochem. *68*, 213–225.

Polyak, K., Xia, Y., Zweier, J.L., Kinzler, K.W., and Vogelstein, B. (1997). A model for p53-

induced apoptosis. Nature *389*, 300–305.

Ramos-Montoya, A., Lee, W.N.P., Bassilian, S., Lim, S., Trebukhina, R. V., Kazhyna, M. V.,

Ciudad, C.J., Noé, V., Centelles, J.J., and Cascante, M. (2006). Pentose phosphate cycle

oxidative and nonoxidative balance: A new vulnerable target for overcoming drug resistance in

cancer. Int. J. Cancer *119*, 2733–2741.

Schlaeth, M., Berger, S., Derer, S., Klausz, K., Lohse, S., Dechant, M., Lazar, G.A., Schneider-

Merck, T., Peipp, M., and Valerius, T. (2010). Fc-engineered EGF-R antibodies mediate

improved antibody-dependent cellular cytotoxity (ADCC) against KRAS-mutated tumor cells.

Cancer Sci. *101*, 1080–1088.

Schulze, A., and Harris, A.L. (2012). How cancer metabolism is tuned for proliferation and

vulnerable to disruption. Nature *491*, 364–373.

Shen, Y., and Shenk, T.E. (1998). Viruses and apoptosis. Curr. Opin. Genet. Dev. *5*, 105–111.

Son, J., Lyssiotis, C. a, Ying, H., Wang, X., Hua, S., Ligorio, M., Perera, R.M., Ferrone, C.R.,

Mullarky, E., Shyh-Chang, N., et al. (2013). Glutamine supports pancreatic cancer growth

through a KRAS-regulated metabolic pathway. Nature *496*, 101–105.

Sun, W., Liu, Y., Glazer, C.A., Shao, C., Bhan, S., Demokan, S., Zhao, M., Rudek, M.A., Ha,

P.K., and Califano, J.A. (2010). TKTL1 is activated by promoter hypomethylation and

contributes to head and neck squamous cell carcinoma carcinogenesis through increased aerobic

glycolysis and HIF1α stabilization. Clin. Cancer Res. *16*, 857–866.

Tennant, D.A., Durán, R. V., and Gottlieb, E. (2010). Targeting metabolic transformation for

cancer therapy. Nat. Rev. Cancer *10*, 267–277.

Tran, T.Q., Hanse, E.A., Habowski, A.N., Li, H., Ishak Gabra, M.B., Yang, Y., Lowman, X.H.,

Ooi, A.M., Liao, S.Y., Edwards, R.A., et al. (2020). α-Ketoglutarate attenuates Wnt signaling

and drives differentiation in colorectal cancer. Nat. Cancer *1*, 345–358.

Tsouko, E., Khan, A.S., White, M.A., Han, J.J., Shi, Y., Merchant, F.A., Sharpe, M.A., Xin, L.,

and Frigo, D.E. (2014). Regulation of the pentose phosphate pathway by an androgen receptor-

mTOR-mediated mechanism and its role in prostate cancer cell growth. Oncogenesis *3*, 1–10.

Vacanti, N.M., and Metallo, C.M. (2013). Exploring metabolic pathways that contribute to the

stem cell phenotype. Biochim. Biophys. Acta *1830*, 2361–2369.

Warburg, O. (1956). On the origin of cancer cells. Science *123*, 309–314.

Ward, P.S., and Thompson, C.B. (2012). Metabolic Reprogramming: A Cancer Hallmark Even

Warburg Did Not Anticipate. Cancer Cell *21*, 297–308.

Wise, D.R., and Thompson, C.B. (2011). Glutamine Addiction: A New Therapeutic Target in

Cancer. Trends Biochem Sci. *35*, 427–433.

Yoo, H., Antoniewicz, M.R., Stephanopoulos, G., and Kelleher, J.K. (2008). Quantifying

reductive carboxylation flux of glutamine to lipid in a brown adipocyte cell line. J. Biol. Chem.

283, 20621–20627.

Young, J.D., Walther, J.L., Antoniewicz, M.R., Yoo, H., and Stephanopoulos, G. (2008). An

elementary metabolite unit (EMU) based method of isotopically nonstationary flux analysis.

Biotechnol. Bioeng. *99*, 686–699.

Zhang, X., Liu, P., Shang, Y., Kerndl, H., Kumstel, S., Gong, P., Vollmar, B., and Zechner, D.

(2020). Metformin and LW6 impairs pancreatic cancer cells and reduces nuclear localization of

YAP1. J. Cancer *11*, 479–487.

Figure 1. Development of a panel of HMECs with different combinations of oncogenes.

Figure 2. Extracellular fluxes were determined by both specific growth rates and oncogenotypes.

Figure 3. ¹³C-isotopic labeling analysis revealed a truncated TCA cycle.

Figure 4. Introduction to ¹³C-MFA and consistency tests against the labeling data.

Figure 5. ¹³C-MFA suggested that both intracellular and extracellular metabolisms may be dually controlled by proliferation and oncogenotypes.

Figure 6. Quantitative ¹³C-MFI analysis validated that oncogenotypes directly impact metabolism independent of proliferative control.

a

b

c

Figure 7. Therapeutic potentials of targeting oxPPP, MDH and IDH to selectively kill the tumorigenic HMEC as opposed to the normal proliferating counterpart.