NEUROeSTIMator: Using Deep Learning to Quantify Neuronal Activation from Single-Cell and Spatial Transcriptomic Data

Ethan Bahl[1,2], Snehajyoti Chatterjee[3,4], Utsav Mukherjee[3,4,8], Muhammad Elsadany[1,2], Yann Vanrobaeys[2,3], Li-Chun Lin[3,4], Miriam McDonough[4,9], Jon Resch[4], K Peter Giese [5], Ted Abel[3,4], and Jacob J. Michaelson[1,3,6,7*]

[1]Department of Psychiatry, University of Iowa.
[2]Interdisciplinary Graduate Program in Genetics, University of Iowa.
[3]Iowa Neuroscience Institute, University of Iowa.
[4]Department of Neuroscience & Pharmacology, University of Iowa.
[5]Department of Basic and Clinical Neuroscience, King's College London, London, UK.
[6]Department of Biomedical Engineering, University of Iowa.
[7]Department of Communication Sciences & Disorders, University of Iowa.
[8]Interdisciplinary Graduate Program in Neuroscience, University of Iowa.
[9]Interdisciplinary Graduate Program in Molecular Medicine, University of Iowa.
* correspondence to jacob-michaelson@uiowa.edu

ABSTRACT

Neuronal activity-dependent transcription directs molecular processes that regulate synaptic plasticity, brain circuit development, behavioral adaptation, and long-term memory. Single cell RNA-sequencing technologies (scRNAseq) are rapidly developing and allow for the interrogation of activity-dependent transcription at cellular resolution. Here, we present NEUROeSTIMator, a deep learning model that integrates transcriptomic signals to estimate neuronal activation in a way that we demonstrate is associated with Patch-seq electrophysiological features and that is robust against differences in species, cell type, and brain region. We demonstrate this method's ability to accurately detect neuronal activity in previously published studies of single cell activity-induced gene expression. Further, we applied our model in a spatial transcriptomic study to identify unique patterns of learning-induced activity across different brain regions. Altogether, our findings establish NEUROeSTIMator as a powerful and broadly applicable tool for measuring neuronal activation, whether as a critical covariate or a primary readout of interest.

INTRODUCTION

Transcriptional regulation is a key component in the downstream response to neuronal stimulation, and the dysregulation of this activity-dependent gene expression has been linked to several nervous system disorders[1-6]. Estimating neuronal activation based on gene expression data is vital for unraveling the intricate processes governing neuronal function and dysfunction, and decades of research has been devoted to characterizing these processes[1,4,7]. Accurate estimation of activation could provide insights into neural circuitry, advance our understanding of neurological disorders, and ultimately contribute to the development of targeted therapeutic interventions. With advances in single-cell RNA-sequencing (scRNAseq) technologies, researchers can now study activity-dependent transcription at the individual neuron level[8,9]. However, scRNAseq has persistent limitations such as sparsity and stochasticity in gene expression measurements, which challenge our ability to use these tools to understand neuronal activation[10-13].

To address these challenges, we present an innovative tool that estimates neuronal activation using transcriptome-scale data as input. Our method combines several technical innovations such as a carefully sampled, multi-species training dataset of neurons, data-driven selection of immediate early genes (IEGs) as model targets, a deep learning for neuron activity estimation, including an architecture designed to protect the latent space from being influenced by cell type biases.

Distilling transcriptome-scale signatures into a 1-dimensional information bottleneck, our approach reconstructs the expression profiles of 22 well-established neuronal activity markers. We show that the resulting activity score is significantly associated with electrophysiological features of increased excitability, demonstrating its applicability in classifying neurons along this essential characteristic. We further show its application in secondary analyses of several single-cell data sets, as well as in new spatial transcriptomic data that captures learning-related circuit activation in the murine brain.

To facilitate community usage, we developed NEUROeSTIMator, an R package with a tutorial that demonstrates our model's application to single-cell data. This innovative method offers a powerful solution to estimate neuronal activation based on de-noised latent patterns in single-cell gene expression, outperforming alternative approaches and providing significant associations with electrophysiological features at the single-cell level.

RESULTS

Multi-species training, test, and validation data sets

We developed a neural network to quantify signatures of transcriptional activation by training an autoencoder to reconstruct expression of 22 coexpressed activity-dependent genes from a 1-dimensional latent space. We trained, validated, and tested

the model using publicly available datasets, including single cell datasets from the Allen
Institute of Brain Science[14] (AIBS) totaling over one million combined samples, as well
as GEO datasets and a novel spatial transcriptomics dataset (**Supplementary Table 1**).
Activity-dependent genes were identified by intersecting differential expression results
of three studies of experimental manipulation of neuron activity[15-17]. All three differential
expression studies examined different brain regions, used different methods of neuronal
stimulation, and were published from independent groups. Intersecting these three lists
yielded a list of 29 candidate target genes (*i.e.*, genes to be reconstructed in
autoencoder output) Principal component analysis of the candidate genes revealed a
relatively low dimensionality of expression profiles, with PC1 explaining a majority of the
variance. (**Supplementary Fig. 1a**). Using the Allen datasets, we then performed
bootstrap permutations of principal component analysis on candidate target genes and
found a subset of 22 genes robustly exhibiting greater PC1 loadings than expected
under a null PCA with randomly shuffled data (**Supplementary Fig. 1b**). These 22
genes were selected as the final set of target genes to be reconstructed by the
autoencoder during training. The Allen Cell Type Databases were downsampled with
hierarchical weighting to increase balance of cell types, datasets, technical variation,
and biological variation (**Supplementary Fig. 2a,b**). Five hundred thousand neurons
were selected and partitioned into cell type-balanced training and testing sets, and the
training set was further split into 5 folds for cross validation. We trained the neural
network to predict expression of the 22 activity-dependent target genes through a 1-
dimensional hidden bottleneck layer with sigmoidal activation (see methods for a
detailed description of model architecture and training). To evaluate model
performance, we applied it to a diverse test set of 164,658 neurons held out from the
training process, including 86,650 samples from the Allen and 78,108 samples from
additional labeled datasets. We selected our final model based on classification
accuracy of labeled test set data through cross validation.
111
112   For further analyses, model output at the bottleneck activation layer was
113   extracted to index activity level for each neuron passed through the model. Hereafter,
114   we refer to this output as the predicted activity or activity score.
115

Genes informing model predictions

117
To identify genes whose expression levels influence model predictions, we
calculated integrated gradients[18] for all input genes with respect to predicted activity
using data from the held out test set. Integrated gradients attribute model predictions to
input features for each cell. We first compared the attributions of target and non-target
genes on predicted activity. The most influential genes were *Fos, Egr1*, *Fosl2, Junb,*
Fbxo33, and *Nr4a1*. We further explored gene importance for all non-target input genes
and found varying degrees of influence throughout the transcriptome (**Supplementary**
Table 2). The most influential non-target genes, such as *Fosb*, *Npas4*, *Egr2*, and *Ntrk2*
tended to have higher initial support for consideration as a target gene and were more
often differentially expressed in multiple transcriptomic studies of neuronal activity-
dependent transcription (**Supplementary Fig. 3a).** Other known activity response
genes such as *Homer1, Egr4, Dusp6,* and *Bdnf* were also among the top fifty genes

with the highest feature attribution. To evaluate whether influential genes were enriched
for specific annotated biological mechanisms, we performed gene set enrichment
analysis on gene ontology and pathway annotation sets (**Supplementary Fig. 3b,**
Supplementary Table 3). Among the most highly enriched gene sets were DNA
binding transcription factor activator activity ($p_{adjusted}$ = 3.97x10^{-9}, NES = 2.29), signaling
by NTRKs ($p_{adjusted}$ = 2.07x10^{-8}, NES = 2.40), cognition ($p_{adjusted}$ = 5.35x10^{-3}, NES =
2.01), MAPK signaling pathway ($p_{adjusted}$ = 2.02x10^{-4}, NES = 2.01), circadian rhythm
($p_{adjusted}$ = 1.60x10^{-3} , NES = 2.03), and cellular response to extracellular stimulus
($p_{adjusted}$ = 0.033, NES = 1.89). To identify gene sets with higher attribution in non-target
genes, we removed target genes, repeated the analysis, and identified gene sets that
remained, or became significant. Of the 45 pathways significantly enriched when
examining all genes, seven sets remained significant when excluding target genes, and
seven new sets became significant, including positive regulation of neurogenesis
($p_{adjusted}$ = 0.019, NES = 1.48), regulation of synapse structure or activity ($p_{adjusted}$ =
0.018, NES = 1.49), and PI3/AKT signaling ($p_{adjusted}$ = 0.019, NES = 1.37)
(**Supplementary Fig. 3b**).

146
147 **Electrophysiological features of individual neurons predicted by transcription via**
148 **NEUROeSTIMator**

149
As an initial application of our model, we examined neurons subjected to direct
electrical stimulation. We integrated single cell RNA assays of a Patch-seq dataset[19]
with a SMART-seq dataset[14] of mouse GABAergic neurons, both generated by the Allen
Institute for Brain Science (AIBS) (**Supplementary Fig 4a**). Using the standard Seurat
workflow[20], we identified seven clusters corresponding to AIBS-defined neuron types
including Sst, Sst Chodl, Pvalb, Vip, Lamp5, Sncg, and Meis2 neuron clusters. We
applied NEUROeSTIMator to both datasets and observed elevated activity score
predictions in the Patch-seq dataset relative to the SMART-seq dataset for seven
neuron types (**Fig. 2a**). We obtained empirical p-value estimates of the Kolmogorov-
Smirnov (KS) test by permuting expression values in each cell and recalculating activity
score for one thousand permutations. Six of the seven neuron types showed significant
differences in activity score distributions between datasets (**Fig. 2b**), of which Lamp5
neurons showed the greatest difference (p < 0.001, D = 0.79, KS test) and Vip neurons
showing the smallest difference (p = 0.033, D = 0.41, KS test). We also noted that
several immediate early genes, which serve as predictive targets for NEUROeSTIMator,
were among the top genes with higher mean expression in the Patch-seq dataset
compared to the SMART-seq dataset (**Supplementary Fig. 4b**). Together, these results
suggest that the Patch-seq protocol induces significantly greater transcriptional
upregulation of activity-dependent genes as compared to the standard SMART-seq
protocol, and our model is sensitive to these differences. Notably, this effect is
consistent for nearly all neuronal types tested.

171
We established that our model detects transcriptomic signals that separate the
electrically stimulated Patch-seq dataset from the unstimulated SMART-seq dataset.
However, it is not clear whether elevated activity was due to electrical stimulation and
neuron depolarization, or general features of the Patch-seq protocol. Patch-seq neurons

showed variability in predicted activity and, because all neurons underwent the
stimulation protocol, we hypothesized that variability in activity score predictions were
linked to individual neuron differences in electrophysiological features, either intrinsic
properties, or responses to stimulation.
180
To determine if NEUROeSTIMator predictions of activity could be explained by
electrophysiology, we fit a lasso-regularized generalized linear model with 10-fold cross
validation to predict NEUROeSTIMator output using only electrophysiological features
derived from Patch-seq recordings. We used 41 electrophysiological features, including
features measured at baseline and in response to the current ramp stimulus. As
substantial differences in electrophysiology exist between neuron types, we residualized
activity score and derived features for neuron type in order to force the model to learn
cell type-agnostic combinations of features that explain variability in activity score. For
further analyses, we focus on the four most abundant neuron types: Lamp5, Pvalb, Sst,
and Vip neurons. A PCA on these 41 features revealed 90% of variance is explained by
fifteen principal components, three of which were associated with activity score (p <
0.05) (**Supplementary Fig. 4c**). The cross-validated lasso model predictions were
significantly correlated with NEUROeSTIMator output (R = 0.234, $p < 2.2 \times 10^{-16}$). To
evaluate feature importance for each electrophysiological feature, we obtained
bootstrap coefficient estimates and standard errors by permuting sample weights for
each of 10,000 model fits (**Fig. 2c**). Three features showed robust non-zero coefficients;
membrane time constant (tau) (coefficient = -0.154, standard error = 0.037), input
resistance (coefficient: 0.088, standard error = 0.041), and the threshold current to
depolarization (coefficient = -0.080, standard error = 0.038). The combination of these
features and their direction of effect suggest that the most stimulus-responsive neurons,
those which exhibit a transcriptional response, are more excitable than other neurons of
the same type (**Fig. 2d**). We then examined specifically the relationship between input
resistance and NEUROeSTIMator predictions in an independent Patch-seq dataset of
human excitatory neurons and, again, found a significant positive correlation (r = 0.497,
p = 0.011) (**Fig. 2e**). Despite the fundamental differences between Patch-seq datasets
in species and major neuron class, this successful validation emphasizes the
robustness of the relationship between neuronal excitability and transcriptional
response to stimulation, as revealed through NEUROeSTIMator predictions.
209
To better understand how NEUROeSTIMator connects gene expression to
electrophysiology, we performed generalized canonical correlation analysis using three
input matrices: the gene expression matrix, a matrix containing the three significant
electrophysiology features, and a one column matrix containing the NEUROeSTIMator
output. We computed a single component and examined gene expression loadings to
understand genes explaining covariance of both predicted activity and the
electrophysiological features of excitability. We used GSEA to identify gene sets
enriched for the genes correlated to both predicted activity and excitability
(**Supplementary Table 4**). We found significant enrichment of several gene sets
including DNA binding transcription activator activity ($p_{adjusted} = 2.74 \times 10^{-5}$, NES = 1.91),
MAPK signaling ($p_{adjusted} = 2.54 \times 10^{-4}$, NES = 1.78), and RAS signaling ($p_{adjusted} =$
3.80×10^{-2}, NES = 1.48) (**Supplementary Fig. 4d**). Genes common to these pathways

with high loadings included *Dusp6, Dusp10, Map3k4, Ntrk2, Rasgrp1*, and *Prkca*. These
pathways constitute an established molecular signaling cascade that functions to link
neuron depolarization and related calcium influx to transcriptional response[21,22].
225
Taken together, these results demonstrate that NEUROeSTIMator captures
transcriptomic signatures of activity that are associated with electrophysiological
properties measured in response to electrical stimulation of individual neurons. Building
upon this successful application in electrophysiology, we next extended our analyses to
demonstrate NEUROeSTIMator's utility across a range of applications (**Supplementary**
Table 1).
232
233 **Detecting pharmacological activation of neurons**
234
We applied our model to three single-cell datasets containing neurons from
rodents and human cell lines treated with powerful stimulating, pharmacological agents.
From the first dataset, we computed activity score for neurons of mice treated with
either saline or pentylenetetrazol (PTZ), a depolarizing agent used to model status
epilepticus and induce seizures[8,23,24]. As expected, we observed significant differences
in distributions of predicted activity (KS test) for 11/14 neuronal subclasses tested,
including excitatory cortical layer 2/3 (L2/3) neurons, Lamp5 interneurons, and Sst
interneurons (L2/3 IT CTX: $D = 0.40$, $p = 9.67\times10^{-24}$), (L2/3 IT ENT: $D = 0.5$, $p =$
4.64×10^{-28}), (Lamp5: $D = 0.49$, $p = 1.73\times10^{-4}$), (Sst: $D = 0.42$, $p = 3.81\times10^{-6}$) (**Fig. 3a,**
Supplementary Table 5).
245
Next, we compared activity score between neurons from the nucleus accumbens
(NAc) of rats treated with either saline or cocaine, a stimulant acting on dopaminergic
neurotransmission[25 26]. We found neuron subtype-specific increases in activity score.
Two subclasses of dopaminergic medium spiny neurons (MSNs) and both Pvalb
interneurons exhibited significant differences in predicted activity (Drd1-MSN: $D = 0.15$,
$p = 5.92\times10^{-7}$), (Drd3-MSN: $D = 0.19$, $p = 0.049$), (Pvalb: $D = 0.22$, $p = 0.028$ (**Fig. 3b,**
Supplementary Table 5). Because our model was not trained using any rat or
dopaminergic neurons, these findings provide further endorsement for robust estimation
of neuronal activity induced by potent pharmacological agents of stimulation.
255
To further evaluate whether our model could be successfully applied to human
data, we examined a dataset of human induced pluripotent stem cell-derived neurons[27].
Neuron cultures were either unstimulated or treated with KCl depolarization[28] buffer for
1, 2, or 4 hours. Our model predicted low neuronal activity for the unstimulated group,
which was consistent among cell types and biological replicates. Cells treated with KCl
for 1 hour demonstrated substantial and significant increases in predicted activity.
Although significant shifts in predicted activity were observed for all cell types ($p < 0.05$),
the activity score of post-mitotic neurons displayed a stronger response to KCl
compared to mitotic progenitors. The most responsive neuron types were the mature
striatal neurons and pallial neurons (striatal: $D_{1hr} = 0.69$, $D_{2hr} = 0.58$, $D_{4hr} = 0.50$,
$p_{1hr/2hr/4hr} < 2.22\times10^{-16}$), (pallial: $D_{1hr} = 0.75$, $D_{2hr} = 0.67$, $D_{4hr} = 0.64$, $p_{1hr/2hr/4hr} < 2.22\times10^{-16}$
^{16}) (**Fig. 3c, Supplementary Table 5**). Notably, all cell clusters except neural

progenitors and septal neurons followed a similar temporal pattern of predicted activity
modestly declining at 2 hours relative to peak activity at 1 hour, with a further decline at
4 hours (*i.e.*, $D_{1hr} > D_{2hr} > D_{4h}$). Despite the steady decline in predicted activity after 1
hour, none of the neuron clusters completely returned to basal activity by 4 hours, the
final time point in the experiment. The temporal trends of predicted activity are indicative
of synaptic depression[29], suggesting NEUROeSTIMator is sensitive to detecting short
term plasticity mechanisms.
275
276 Together, these analyses suggest our model can robustly assign higher
277 estimates of activity to cells subjected to chemical exposures that are expected to elicit
278 strong and generally ubiquitous transcriptional responses to stimulation.
279
280 **Activity score as a generalizable classifier of neuronal stimulation**
281
Next, we asked if our model could detect neuronal activation by more subtle
forms of stimuli, such as sensory experience. We applied our model to a dataset
containing visual cortex neurons from mice housed in darkness for seven days prior to
light exposure for 0, 1, or 4 hours[30] (**Fig. 4a**). We observed a significant increase in
predicted activity for neurons from mice exposed to light, relative to controls. To
elucidate temporal patterns of activity, we tested differences in activity score
distributions between pairs of each time point. Activity score was significantly increased
at 1 hour for all cell types, in particular for excitatory cortical neurons (layer 2/3: D =
0.56, $p < 2.22 \times 10^{-16}$), (layer 4: D = 0.60, $p < 2.22 \times 10^{-16}$), (layer 6: D = 0.54, $p < 2.22 \times 10^{-16}$
). At 4 hours of light exposure, predicted activity began to show diverging trends which
were foreshadowed by predicted activity at 1 hour. Although trending towards a return
to baseline, activity scores of neurons at 4 hours were not significantly different from
neurons at 1 hour, while several glial cell types returned to baseline (**Fig. 4a,
Supplementary Table 6**).
296
As we observed similar trends in temporal activity predictions between the
unstimulated (0h) and 1h group, we investigated the degree to which the activity score
derived from our model could be used as a classifier of experimental group. The degree
to which activity score is predictive of experimental group in a particular cell type is
expected to represent the robustness of the response in that cell type. Using the visual
cortex dataset (VIS) mentioned above, we constructed receiver-operator curve (ROC)
plots for neuronal and non-neuronal cell types (**Fig. 4b**). For both excitatory neuron and
interneuron subtypes, the activity score demonstrated varying degrees of predictive
power. For example, the activity score alone was able to almost perfectly separate
stimulation groups when considering excitatory cortical layer cell types (layer 2/3: AUC
= 0.84), (layer 4: AUC = 0.89), (layer 6: AUC = 0.823), as well as certain interneuron
cell types (Sst-2: AUC: 0.97), (Vip: AUC = 0.90).)Despite the model being trained on
purely neuronal cell type populations, the activity score was able to separate stimulation
groups for astrocytes (AUC = 0.84) and endothelial cells (AUC = 0.72) with comparable
accuracy to neurons.
312
Utility in data modalities beyond scRNA-seq

Next, we asked whether our model could identify spatial signatures of learning in
brain slices of mice following spatial object recognition (SOR) training - a widely used
behavioral paradigm to investigate spatial memory mechanisms[31]. Using spatial
transcriptomic data from brain slices of SOR-trained and homecage control (HC) mice,
we applied our model to predict activity for each spot (**Supplementary Fig. 5a**) and
clustered all spots into anatomical regions. The 23 resulting clusters were annotated
with brain region naming conventions from the Allen Mouse Brain Atlas (**Fig. 5a**). In
baseline homecage samples, we noted weak activation signatures, primarily covering
regions of the striatum and hypothalamus as well as subregions of the hippocampus
(**Fig. 5b, left**). Following SOR training, we noted region-specific increases in predicted
activity (**Fig. 5b, right**). To identify a spatial activation signature of SOR for the entire
brain slice, we tested for differences in predicted activity for each brain region cluster
(**Fig. 5c**). We observed significant increases in predicted activity for several cortical and
subcortical regions. Multiple layers of the isocortex and the retrosplenial area showed
increases in activity following SOR (**Fig. 5d**), with the greatest increase observed in
layers 2/3 ($\beta = 0.17$, $p < 2.22 \times 10^{-16}$) and the retrosplenial cortex ($\beta = 0.17$, $p < 2.22 \times 10^{-16}$
). The caudoputamen area of the dorsal striatum, subregions of the hippocampus, and
the olfactory/piriform areas also showed significant increases in activity, of comparable
magnitude (caudoputamen: $\beta = 0.11$, $p < 2.22 \times 10^{-16}$). Subregions of the hippocampus
were variably activated following SOR, with the strongest induction observed in the CA1
region ($\beta = 0.13$, $p < 2.22 \times 10^{-16}$). Regions predicted to be least activated by SOR
include the thalamus, hypothalamus, dentate gyrus of the hippocampus, the lateral
ventricle, and fiber tracts (**Supplementary Table 7**).
Visium samples contain signal from multiple neuronal and non-neuronal cells.
Cortical layers and CA1 showed profound increases in activity but are also known to
have relatively high neuron density. Therefore, we considered the extent to which
regional neuron density influences coefficient estimates in **Fig. 5d.** We performed cell
type deconvolution on each Visium spot, using neuronal and non-neuronal cell type
expression profiles from the Allen Cell Types Database. For a subset of genes with high
variability across cell classes, we fit a linear model to each Visium spot as a function of
GABAergic, glutamatergic, and non-neuronal expression profiles. Coefficient estimates
for each cell class were rescaled across samples to a range of zero to one, and then
adjusted for each sample to sum to one, thus providing a proxy measure of cell class
density (**Supplementary Fig. 6a**). We found a significant positive relationship between
estimated regional neuron abundance and $\log_{10}$-estimated regional effect of learning on
activity ($\beta = 0.86$, $p = 0.01$), (Spearman correlation = 0.52, p = 0.01) (**Supplementary.**
Fig. 6b). We used the residuals of the linear model in **Supplementary Fig. 6b** as a
neuron density-corrected view of the effect of learning (**Supplementary Fig. 6c**).
Cortical layers, retrosplenial cortex, CA1, and caudoputamen regions show greater
learning-induced increases in activity score than expected based on neuron density
estimates alone, suggesting that neurons in these regions exhibit a more potent
transcriptional response to learning, or that there are other cell types contributing to the
activity signature. The dentate gyrus and regions of the hypothalamus, amygdala,
thalamus showed lower activity induction than expected.

To validate the finding that the CA1 region was preferentially activated by spatial
learning over the dentate gyrus, we performed RNAscope[32] on coronal brain slices from
SOR-trained mice (1 hour post-training) and a homecage control mouse
(**Supplementary Fig. 5b**). We selected three NEUROeSTIMator target genes, *Egr1*,
Egr3, and *Nr4a1* as probes. For each brain slice, we quantified mean fluorescent
intensity (MFI) of each probe in CA1 and dentate gyrus. A two-way ANOVA revealed a
significant interaction between hippocampus subregion and training group (F = 8.015,
p=0.0299). In line with the spatial transcriptomic data (**Supplementary Fig. 5c**), we
found that CA1 showed a preferential increase in average probe intensity following
spatial learning (Sidak's test, CA1: $p_{adjusted}$ = 0.0075, dentate gyrus: $p_{adjusted}$ = 0.5553)
(**Supplementary Fig. 5d**).

371
372

373 **NEUROeSTIMator outperforms competing methods in discriminating**
374 **experimental stimulation groups**
375

The NEUROeSTIMator approach to quantifying activity uses an encoding neural
network to infer transcriptome-wide coexpression with a set of 22 activity-dependent
target genes. We investigated whether the added complexities of our model are
necessary to produce a metric that successfully discerns stimulated samples from
unstimulated. We compared NEUROeSTIMator to three alternative models. The first
competing method is a scaled additive model, a simple and easily implementable
approach that scales normalized expression of NEUROeSTIMator's 22 target genes
across samples to range of [0,1], and then sums scaled values within samples. The
second method, the PC1 model, scores samples with target gene loadings derived from
the first component of a PCA on the NEUROeSTIMator training dataset. The third
method, the transcriptome-naive model is a neural network trained with identical
parameters to NEUROeSTIMator, but without access to non-target gene expression
inputs. We evaluated each model by scoring activity in stimulated and unstimulated
samples from six datasets (**Supplementary Table 1**) and determined separability of the
experimental groups using area under the curve (AUC). This approach was applied to
datasets featured in Figures 3, 4, and 5, as well as a novel single nuclei dataset from
mouse dorsal hippocampus following SOR. For each method, dataset and neuron type
we computed AUC and summarized the performance differences between
NEUROeSTIMator and competing methods (ΔAUC). (**Supplementary Fig. 7,**
Supplementary Table 8). NEUROeSTIMator demonstrated higher performance than
each of the competing methods ($AUC_{NEUROeSTIMator}$ = 0.724, $AUC_{scaled.additive}$ = 0.541,
AUC_{PC1} = 0.658, $AUC_{transcriptome.naive}$ = 0.666), (mean $\Delta AUC_{scaled.additive}$ = +0.183, ΔAUC_{PC1}
= +0.066, $\Delta AUC_{transcriptome.naive}$ = +0.058). Across all competing methods, datasets, and
sample groupings, NEUROeSTIMator increased AUC in 88.1% of 336 comparisons. For
comparisons where NEUROeSTIMator outperformed the competing method, the
average increase in AUC was +0.11. For the remaining 11.9% of comparisons where
NEUROeSTIMator did worse than the competing method, the average decrease in AUC
was -0.031. Together, these results suggest that the neural network architecture and
NEUROeSTIMator's use of broader transcriptomic signals underlie its enhanced
performance compared to alternative approaches.

DISCUSSION

We introduce NEUROeSTIMator, a versatile tool that estimates neuronal activity from transcriptome-wide single-cell gene expression data. Our method generates an activity score that reflects transcriptional response to stimulation, facilitating the rapid identification and prioritization of neurons displaying stimulus response. NEUROeSTIMator's performance and generalizability are demonstrated through capturing electrophysiological (e-phys) features and detecting activation signatures across various stimulation protocols, neuron subtypes, species, and sequencing technologies, including spatial transcriptomics.

Our analysis reveals that activity-predictive signals are distributed broadly across the transcriptome and are enriched for genes related to external stimulus response, synaptic plasticity, MAP kinase activity, and circadian rhythm. Moreover, we demonstrate a significant link between NEUROeSTIMator's predictions and increased neuronal excitability, utilizing a unique single-cell Patch-seq dataset containing gene expression and e-phys readouts from individual GABAergic murine neurons[19]. These results were further confirmed through a significant association between the activity score and measures of excitability in human excitatory neurons[33].

We showcase NEUROeSTIMator's flexibility and generalizability through secondary analyses of several diverse published datasets, detecting potent pharmacologically induced neuronal activity and generalizing across species and cell types. Importantly, our model also detects physiologically relevant activation in response to sensory experiences and uncovers unexpected activity in non-neuronal cells (e.g., astrocytes), suggesting potential applicability to other cell types and stimulus responses.

In a novel application, we use NEUROeSTIMator to analyze spatial transcriptomic data from mice subjected to a spatial learning task, revealing a brain-wide map of learning-induced activation. This highlights the method's adaptability to spatial transcriptomics (and other varieties of bulk RNA-seq data) and suggests potential for uncovering distinct spatial signatures of transcriptional activation related to cognitive processes.

While our model demonstrates a significant link between transcriptional response and e-phys features in murine interneurons and human excitatory neurons, its applicability to e-phys features in broader cell types is limited by the few available Patch-seq datasets. Additionally, our model does not inherently differentiate stimulation types (e.g., pharmacological vs. naturalistic), and we observed some sensitivity to activity signatures in non-neuronal cell types (a potentially desirable feature, depending on the application). Despite these limitations, NEUROeSTIMator delivers key insights across several neuroscience research domains.

451 As the first robust and generalizable method for estimating neuronal activation from
452 transcriptomic data, NEUROeSTIMator has broad implications for molecular
453 neuroscience research. Our method enables the estimation of cellular activity state, a
454 fundamental variable in gene expression research, paving the way for its integration into
455 single-cell and spatial transcriptomics analyses.
456
457
458 **METHODS**
459
Dataset for model training and evaluation
To train the model, we utilized publicly available datasets provided by the Allen Institute
for Brain Science, including a single-cell RNA-sequencing (scRNAseq) dataset of over a
million cells isolated from mouse cortical and hippocampal tissue[14], and a single-nuclei
RNA-sequencing (snRNAseq) dataset of 76,000 nuclei isolated from human cortical
tissue. Hereafter, these datasets will be referred to as the Allen Mouse and Allen
Human datasets. Both datasets used the 10X Genomics Chromium system for droplet
capture. The Allen Mouse dataset was prepared using the Chromium Next GEM Single
Cell 3' v3 reagent kit, while the Allen Human dataset used v2.
469
470 **Datasets for model application – publicly available**
471 We downloaded multiple datasets from Gene Expression Omnibus (GEO) to
472 demonstrate the utility of our model. The following GEO accessions were included in
473 analyses: GSE102827[34], GSE106678[9], GSE136656[27], and GSE137763[25].
474
Gene identifier mapping
We used the R package biomaRt to map gene identifiers from various annotations used
in public datasets, and between species, to a common set of reliably mapped
genes[35,36]. Ensembl gene identifiers (Ensembl IDs) were used as the primary identifier
for mapping genes, and gene symbols were used as secondary identifiers in cases of
ambiguous mapping. The Ensembl release 93 archive (July 2018 release) was used for
cross-species gene mapping[37]. Genes with one-to-one orthology between mouse and
human, as well as mouse and rat, were selected to facilitate cross species utility. All
datasets lacking Ensembl ID annotation contained gene symbols, which were then
queried against multiple Ensembl archives to determine which archive maximized
identifier mapping rate. For instances when a gene symbol mapped to multiple Ensembl
IDs, identifiers present in the cross-species mapping table were preferentially selected.
We provide a helper function for mapping gene identifiers to the feature set used by our
model, and we further demonstrate usage in the associated tutorial.
489
Choice of neural network target genes
To identify robust markers of neuronal activity for use as targets of the neural network,
we intersected lists of stimulus-responsive genes from three published RNA-sequencing
experiments. Each publication was from a different group of authors, focused on
different brain regions, and used different forms of neuronal stimulation (see Table 1).
One publication categorized stimulus-responsive genes into three groups – rapid
primary response genes (rPRGs), delayed primary response genes (dPRGs), and

secondary response genes (SRGs)[17]. As SRGs are thought to demonstrate higher
celltype-specificity relative to PRGs[1], only rPRGs and dPRGs were considered from this
publication. In another publication, approximately 600 genes upregulated in response to
kainic acid treatment in the hippocampus were considered[16].

In a third publication, two sets of KCl-responsive genes were available, one from
a brain region enriched in glutamatergic neurons and one enriched for GABAergic
neurons[15]. For this study, we sought to intersect the results of both the glutamatergic
and GABAergic analyses into one list of genes. Published p-value distributions
suggested differences in statistical power between these two analyses, and only
statistically significant results were published. To expand the list of genes overlapping
between these analyses, we reanalyzed the data using GEO2R to obtain two sets of
transcriptome-wide statistics. Using significance rankings from the GEO2R reanalysis,
we jointly determined p-value thresholds for each analysis based on rank-rank
hypergeometric overlap (RRHO, see **Supplementary Fig. 5**) and identified genes with
p-values below these thresholds in both sets with concordant direction of effect[38].
Because this approach used unconventional p-value thresholding, we additionally
required intersecting genes to have an estimated fold change greater than or equal to
0.5 in both analyses.

Intersecting the three lists of stimulus response genes provided 41 candidate
target genes. From these 41, four genes were excluded due to ambiguous or failed
cross-species gene identifier mapping. We also excluded an additional eight genes that
were detected in less than 1% of cells in more than one out of the four species-by-class
groups (i.e., mouse-GABA, mouse-glut, human-GABA, human-glut) to mitigate the
potential for the model to learn spurious target correlations to cell class markers or
species markers. We performed principal component analysis (PCA) on the reaming 29
candidate target genes to assess dimensionality and found that PCs beyond PC1 added
very little to the variance explained (**Supplementary Fig. 1a**). All but one gene had a
positive loading on PC1, and several genes had a loading near zero. We identified
genes that had extreme positive loadings by comparing them to an outlier threshold for
PC1 loadings from a shuffled, null PCA. A scaling factor, the ratio of variance from the
observed PC1 to the null PC1, was used to make the loadings comparable. The outlier
threshold was defined as the median null loading plus or minus three times the median
absolute deviation of the null loadings. We found that 22 of 29 genes had loadings
considered to be outliers under the null expectation (**Supplementary Fig. 1b**). We then
did 1000 bootstrap permutations of this analysis, each using random samples of 10,000
cells, and counted the number of permutations where each gene's loading surpassed
the outlier threshold. The same 22 genes surpassed the threshold in at least 5% of the
permutations, while the other 7 genes did not (**Supplementary Fig. 1c**). With evidence
of robust coexpression, those 22 genes were selected as the final set of targets for our
neural network: *Arc*, *Btg2*, *Coq10b*, *Crem*, *Dusp1*, *Dusp5*, *Egr1*, *Egr3*, *Fbxo33*, *Fos*,
Fosl2, *Gadd45g*, *Gmeb2*, *Grasp*, *Junb*, *Nr4a1*, *Nr4a2*, *Nr4a3*, *Per1*, *Rgs2*, *Sertad1*, and
Tiparp.

Sample filtering, downsampling, and partitioning
Cells were filtered out if they had less than 5,000 total counts, greater than 50,000 total
counts, less than 1,000 genes detected, more than 10% reads originating from

mitochondrial genes, or more than 10% reads originating from the most highly
expressed gene. Non-neuronal cells and ultra-rare neuron subclasses with less than
1,000 cells were removed, and imbalances among species, neuron type, quality control
metrics and naively-estimated activity were alleviated using weighted downsampling
with hierarchically readjusted weights. Cells were grouped hierarchically by species,
class, brain region neighborhood, subclass, and finally, by two-dimensional binning of
one source of technical variation ($\log_{10}$ of total counts), and one source of biological
variation (PC1 of target genes). Each group at the leaf nodes of the hierarchy were
assigned equal weights summing to one. At each level of the hierarchy, moving from
leaves to root, weights were readjusted to sum to one for each group. Using these
weights, we downsampled the dataset and calculated Shannon's entropy to measure
imbalance at different values of N (**Supplementary Fig. 2a**) We chose to downsample
to 500,000 cells to strike a balance between dataset size and diversity. The effects of
downsampling are demonstrated for four subclasses in **Supplementary Fig. 2b**. The R
package *groupdata2* was used to create five training folds (91.3%, 456,725 samples)
and one test split (8.7%, 43,275 samples). We used the *groupdata2* function 'fold' with
the parameters 'cat_col' set to the species-class group and 'num_col' set to the target
gene PC1 values in order to keep these balanced across folds, but also set the 'id_col'
parameter to the subclass label so each subclass was present in some folds but not
others. This was done to discourage the undesirable learning of subclass-specific bias
in cross-validation.
564
Dataset augmentation
The Allen Cell Types Database has higher average total counts per cell compared to
many publicly available datasets. To increase representation of lower quality cells, we
created augmented samples by synthetically downsampling raw counts using the R
package scater[39]. For each combination of species and neuron subclass, an equal
number of cells were randomly assigned a value of either 10^3, $10^{3.25}$, or $10^{3.5}$ total
counts to be downsampled to. All genes were considered for downsampling.
572
Feature normalization and preprocessing
Log-normalization, as implemented in Seurat, was used to normalize input gene
expression. Total counts for each cell were calculated by summing only the 10,017
features used by the model. Normalized expression levels for each gene were centered
and scaled based on mean and standard deviation estimated from the training data. For
cross validation, mean and standard deviation were estimated without the held-out fold.
579
Model architecture
The architecture of the model was adapted from DCA[40]. Briefly, input gene expression
is supplied to encoder branches; a series of three fully connected dense layers with
ELU activations and batch normalization. ELU activations were chosen over RELU to
increase learning speed and performance. The first three hidden layers of the encoders
contained 16, 8, and 4 units. The encoder then connects to the information bottleneck,
a single-unit dense layer with sigmoid activation. To restrict the bottleneck from
capturing biases in the training data, we directly supplied neuron subclass, dataset, and
quality control variables to a small encoder with two hidden layers and an output of six

units, which was then concatenated with the bottleneck activation as well as the output
branch for dispersion and dropout parameters to form penultimate decoder layers. The
decoder outputs the estimated mean parameter **μ** of the zero-inflated negative binomial
(ZINB) model. A separate encoder branch outputs estimates of the dispersion and
dropout parameters. The ZINB loss function was used as implemented in DCA, with the
only difference being that our implantation only computes loss for the target genes
rather than all inputs to the model. This reconstruction loss is based on the maximum
likelihood estimate of the ZINB count model, a common model assumed to underlie the
count distributions of single cell RNA-seq datasets. A binary cross entropy loss was
added to the bottleneck layer to monitor classification accuracy. For model applications,
the model outputs are not used, but the sigmoidal bottleneck activation value is the
metric extracted to index neuronal activity (see **Supplementary Fig. 10**).
601
Model training
We trained the model using keras, as implemented in keras R package, version 2.3.0.0.
Training proceeded for 10 epochs using the ADAM optimizer. Gaussian dropout was
applied to input expression to simulate uncertainty in measurements. Augmented
samples were given the same output as the original data to curtail the learning of depth-
dependent information, and augmented samples appeared in the same training batches
as the samples they were derived from. We used basic hyperparameter tuning to
evaluate the impact of training epochs, learning rate, and regularization parameters
such as dropout, L1, and L2 penalty. The final model was selected based on the
minimum cross-validation bottleneck layer loss for labeled data (binary cross entropy).
Although samples from labeled datasets contributed to the ZINB reconstruction loss, the
bottleneck loss did not contribute to weight updates during training to curtail the
possibility of information leakage between the training and test splits. Only test split
samples were included in any downstream analysis presented in this paper.
616
Evaluating feature importance
To evaluate relative importance of each gene on predicted activity, we implemented a
strategy adapted from the integrated gradients[18], a popular feature attribution approach
used in image classification. Briefly, model gradients were calculated from input genes
to bottleneck activation for the test set. For each gene we fit a generalized additive
model (GAM) with smoothing between the sample gradients and bottleneck activation.
We estimated global feature importance by approximating the integral of the fit. We took
the summation of the GAM output for the full range of possible bottleneck activation
values [0,1], with a step size of 1×10^{-3}. Gene set enrichment analysis was run using the
R package fgsea. We calculated enrichment for gene ontology, Reactome, KEGG, and
Wikipathways terms of size between 100-200.
628
Comparison of Patch-seq and SMART-seq datasets
The SMART-seq and Patch-seq datasets were integrated, clustered, and visualized
using the R package Seurat[20]. The sctransform 'v2' method was used for
normalization[41]. Cell clustering results were cross-referenced with provided neuron type
annotations to identify neuron types common to both datasets and harmonize labels.
We used the Kolmogorov-Smirnov test (KS test) to assess distributional differences in

predicted activity of samples from the SMART-seq and Patch-seq datasets at the
neuron type level. Only neuron types with at least one hundred samples in each dataset
were considered. To derive empirical p-values and determine whether dataset-specific
biases is transcriptome-wide expression levels contributed to observed differences in
predicted activity, we ran 1,000 permutations of the KS test using activity score
predictions from within-sample shuffled gene expression values. Empirical p-values
were calculated from the frequency of permutation D statistics that were more extreme
than those observed in the actual data.
643
Modeling activity score with electrophysiological features
The whole-cell Patch-seq electrophysiology recordings from neurons of mouse visual
cortex provided by the Allen Institute for Brain Sciences were downloaded as NWB files
from the DANDI (ID: 000020, September 2021 update)[19]. The dataset included
electrophysiological recordings of 4,284 specimens from 1,040 mouse subjects. The
Python package IPFX was used to perform quality control and compute
electrophysiological features from the NWB files. We built a generalized linear model
with lasso regularization using the R package "glmnet" to predict output of
NEUROeSTIMator using only electrophysiological features. Cells included in the glmnet
training belonged to major neuron types (*e.g.*, Vip, Sst) with at least 200 cells, and
minor neuron types (*e.g.*, Sst Rxfp1 Prdm8) with at least 10 cells. Features with missing
values in five hundred or more cells were removed, and cells with any missing values
for the remaining features were discarded. We engineered additional features from the
IPFX output, including time and voltage changes between various action potential
landmarks (*e.g.*, voltage change between action potential threshold and peak). Minor
neuron type was regressed out of the activity score and electrophysiological features.
Cells with feature values more extreme than three standard deviations of the mean
were considered outliers and removed. Significance of non-zero coefficients from the
glmnet model was evaluated with 10,000 bootstrap permutations. In each permutation,
a vector of weights is shuffled and used in fitting the glmnet model. For each input
feature, we computed the mean bootstrap coefficient. The confidence interval was
defined as the mean estimate plus or minus 1.96 times the standard deviation of the
bootstrapped coefficients. Feature significance is indicated by the confidence intervals
not reaching or overlapping zero. To validate the electrophysiology findings, we did a
Pearson's correlation test between NEUROeSTIMator output and input resistance for
Patch-seq neurons from the AIBS dataset 'Multimodal Analysis of Human Layer 5
Pyramidal Neurons'[33]. Canonical correlation analysis (CCA) was used to explore gene
expression profiles with shared relationships to predicted activity and three
electrophysiological features. Inputs to the generalizable CCA included the
NEUROeSTIMator output matrix, the matrix of three significant electrophysiological
features, and the gene expression matrix. One canonical component was calculated for
each input matrix, with the tau parameter set to 0 to maximize correlations between all
sets of variables. Gene set enrichment analysis was performed as described above
using the gene loadings for the gene expression canonical component.
678
Preparation of application datasets

Application datasets were downloaded from the Gene Expression Omnibus (GEO). In
datasets without provided labels, cell types were identified by running the standard
Seurat pipeline, identifying cell clusters, and cross-referencing cluster marker genes to
those used in the source publications or the Allen Cell Types Database. To prepare
datasets for input to NEUROeSTIMator, gene expression data was log-normalized
sample-wise and scaled gene-wise using mean and standard deviation values from the
original training dataset. Samples with at least 1000 counts were then input to the model
and activity score was the 1-unit bottleneck activation layer output.
688
Testing differences in predicted activity
For all datasets analyzed in figures 2 and 3, we used the KS test to evaluate differences
in activity score distributions. A linear model was used to test for differences (i.e., using
t-statistic of the regression slope) in predicted activity of the spatial transcriptomic
clusters in figure 5. In **Supplementary Figure 5b** we adjusted the coefficient estimates
of learning on predicted activity for neuron density. We estimated neuron density using
a cell type deconvolution technique. We derived gene signatures for GABAergic
neurons, glutamatergic neurons, and non-neuronal cell classes by averaging the
median gene expression for their respective clusters and subclasses from the Allen Cell
Types Database.

For each sample in the Visium dataset, we constructed a linear model, correlating the
gene expression vector with a three-column matrix that contained class-averaged
expression signatures. We then normalized the coefficient estimates across all Visium
samples for each cell class by scaling them to a 0-1 range and then adjusting their sum
to one within samples. This process provided an estimate of abundance for each cell
class. We summed the estimated abundances of glutamatergic and GABAergic neurons
to determine neuron abundance or density for a specific Visium spot, Subsequently, we
created a linear model of brain region log coefficient estimates as a function of brain
region averaged neuron density estimates. The residuals of this model represent the
neuron density-corrected coefficient estimates.

709
Performance benchmarking
We compared the performance of NEUROeSTIMator to three alternative approaches to
quantify activity. Performance was defined by the ability to separate labeled cells based
on the stimulation status (*i.e.*, experimental group) using area under the curve (AUC).
The three alternative approaches were calculated as follows. The scaled additive
measure of activity is calculated by first scaling log-normalized target expression values
to a range between 0 and 1, and then summing the scaled targets in each cell or
sample. To calculate the PC1 measure, we first performed PCA on target genes of the
training set used to train NEUROeSTIMator. The loadings for PC1 are then applied to
the test set. The transcriptome-naïve model is a version of NEUROeSTIMator trained
with the same parameters, but information from the non-target input genes is blocked by
setting their input values and weights to the first hidden layer to zero. To benchmark
NEUROeSTIMator against alternative models for binary outcome prediction of
stimulation group status, we evaluated model ability to separate stimulated from

unstimulated samples. We computed the Area Under the Receiver Operating Characteristic Curve (AUC-ROC) for each model across all combinations of source dataset and sample groupings. We then determined the AUC differences between NEUROeSTIMator and the alternative models for each sample grouping. To summarize the average AUC difference within each dataset, we calculated the mean of the AUC differences for the respective sample groups. This yielded a single value representing the mean AUC difference for each source dataset in the test set. Lastly, we derived the final performance metric by computing the mean of these mean AUC differences across all data sources, offering a comprehensive evaluation of the NEUROeSTIMator's performance relative to the alternative models.

Animals

All experiments were performed using male C57BL/6J mice obtained from Jackson Laboratory (#000664), aged between 2 and 4 months. The mice were housed in groups of up to five individuals per cage, which contained soft bedding material. They had access to food (NIH-31 irradiated modified mouse diet #7913) and water ad libitum. A 12-hour light-dark schedule was maintained, with the start of the lights-on period designated as Zeitgeber time zero (ZT 0). The experimental procedures followed the guidelines for animal care and use set forth by the National Institutes of Health, and they were approved by the Institutional Animal Care and Use Committee (IACUC) at the University of Iowa.

Behavioral training for spatial transcriptomics

We generated a novel spatial transcriptomic dataset examining the effects of spatial learning in mice. The dataset contains spatial RNA-sequencing of whole brain slices from 1 hour after SOR training or home cage controls (HC). 10-12 weeks old C57BL/6J male mice were trained in SOR task as described previously[31]. Briefly, mice were handled for 5 consecutive days prior to the SOR training. On the day of the training, animals were individually placed in an open field with a spatial cue and was allowed to habituate for 6 mins. They were returned to their home cage and the arena was wiped clean with ethanol. The animals were again placed back into the open field with three objects at specific spatial locations. The animals were allowed to freely explore the objects in the arena for 6 mins. This training with objects comprised of three trials of 6 mins each. One hour after the completion of the training, mice were euthanized and whole brains were flash frozen in isopentane placed on dry ice.

Spatial transcriptomics

Mouse brain section per mouse was cut at 10 μm thickness and mounted onto each Visium slide capture area. After H&E staining, each bright-field image was taken as described in the spatial transcriptomics protocol. Tissue permeabilization was performed for 18 minutes, as established in the tissue optimization assay. The Visium Spatial Gene Expression Slide & Reagent kit (10x Genomics) was used to generate sequencing libraries for Visium samples. Libraries were constructed according to the 10x Visium library construction protocol and sequenced by Illumina NovaSeq6000. Raw data was then processed using the 10x Genomics Space Ranger analysis pipeline. See **Supplementary Figure 5** for images of predicted activity for each replicate.

In situ hybridization

In situ hybridization was performed on 20 μm coronal brain sections from OR-trained and HC mice using RNAscope reagents (Advanced Cell Diagnostics). Briefly, fixed frozen brains were sectioned on a microtome and mounted on Superfrost™ Plus microscope slides (Fisherbrand). Slides then underwent serial dehydration in Ethanol, followed by Hydrogen Peroxide treatment, target retrieval, and Protease III treatment – all of which were done according to the manufacturer's protocol. Hybridization of probes was done at 40°C for 2 hours in an HybEZ oven using a 50:1:1 cocktail of Egr1, Nr4a1, and Egr3 respectively. The probe signals were amplified with Pre-amplifier 1 (Amp 1-FL) and counterstained with OPAL dyes (Akoya Biosciences) corresponding to different excitation-emission wavelengths. Finally, the slides were mounted with Vectashield® Antifade Mounting Medium with DAPI (Vector Laboratories). Slides were stored in 4^0C until they were imaged.

Confocal imaging and analysis

After performing the RNAscope assay, brain sections were imaged using Olympus FV3000 confocal microscope with a 10X NA = 0.4 objective at 800 × 800-pixel resolution. High magnification images of the hippocampus were obtained using a 100X NA = 1.45 oil immersion objective at 800 × 800-pixel resolution and 1.25 optical zoom. All images (8 bit) were acquired with identical settings for laser power, detector gain and pinhole diameter for each experiment and between experiments. Images from the different channels were stacked and projected at maximum intensity using ImageJ (NIH). Hippocampal subregion-specific Mean Fluorescence Intensity (MFI) of each RNAscope probe was evaluated using plugins in ImageJ. The average MFI for all the 3 probes corresponding to each hippocampal subregion was compared between the two experimental groups. Two-way ANOVA was used to assess significance of subregion-by-training interaction (n = 4 for each subregion/training combination). A post-hoc Sidak's multiple comparisons test was applied to assess significance of individual subregion differences.

DATA AVAILABILITY

Spatial RNA-sequencing data, including gene expression measurements, tissue images, spot coordinates, and raw FASTQ files have been deposited in the Gene Expression Omnibus repository under the reference series ID GSE201610.

CODE AVAILABILITY

NEUROeSTIMator is available at https://research-git.uiowa.edu/michaelson-lab-public/neuroestimator/ as a free R package with installation instructions and a tutorial.

ACKNOWLEDGEMENTS

Funding: This work was supported by NIH grant R01 MH 087463 to T.A., NIH grant R01 DC014489 to J.J.M., NIH grant K99 AG 068306 and the Nellie Ball Trust to S.C., and The University of Iowa Hawkeye Intellectual and Developmental Disabilities

815 Research Center (HAWK-IDDRC) P50 HD103556 to T.A. and Lane Strathearn. T.A.
816 and J.J.M. are also supported by the Roy J. Carver Charitable Trust.
817

AUTHOR CONTRIBUTIONS
EB, SC, TA, and JJM conceived the study. EB performed data curation, model training,
analyses, and visualizations. SC and TA designed the molecular biology experiments
with input from JJM and KPG. SC performed behavioral experiments, stereotactic
surgeries, and single nuclei sequencing. LC performed Visium spatial transcriptomic
sequencing experiments. YV and ME processed Visium data. JR and MM performed
RNAscope experiments. UM performed statistical analysis and created visualization
pertaining to the RNAscope data. EB and JJM wrote the article.
826

Data Sources
We thank the Allen Institute for Brain Sciences for their valuable datasets we used to
train our model. We thank the creators and authors of DCA, whose work inspired the
approach we implemented in this paper. Finally, we thank the investigators who make
their data publicly available on repositories like GEO. Specifically, we would like to
thank the authors providing data for GEO series GSE111899, GSE125068, GSE55591,
GSE106678, GSE137763, GSE136656, and GSE102827.
834

835 **Competing interests:** The authors declare that they have no competing interests.
836

837 We also thank Mahesh Shetty for his valuable contributions to discussions about this
838 project.
839

840
841
842   References
843
1   Yap, E. L. & Greenberg, M. E. Activity-Regulated Transcription: Bridging the Gap
between Neural Activity and Behavior. *Neuron* **100**, 330-348 (2018).
https://doi.org:10.1016/j.neuron.2018.10.013
2   Cohen, S. & Greenberg, M. E. Communication between the synapse and the
nucleus in neuronal development, plasticity, and disease. *Annu Rev Cell Dev Biol*
24, 183-209 (2008). https://doi.org:10.1146/annurev.cellbio.24.110707.175235
3   West, A. E. & Greenberg, M. E. Neuronal activity-regulated gene transcription in
synapse development and cognitive function. *Cold Spring Harb Perspect Biol* **3**
(2011). https://doi.org:10.1101/cshperspect.a005744
4   Gallo, F. T., Katche, C., Morici, J. F., Medina, J. H. & Weisstaub, N. V. Immediate
Early Genes, Memory and Psychiatric Disorders: Focus on c-Fos, Egr1 and Arc.
Front Behav Neurosci **12**, 79 (2018). https://doi.org:10.3389/fnbeh.2018.00079
5   Mews, P. *et al.* From Circuits to Chromatin: The Emerging Role of Epigenetics in
Mental Health. *J Neurosci* **41**, 873-882 (2021).
https://doi.org:10.1523/JNEUROSCI.1649-20.2020
6   Nido, G. S., Ryan, M. M., Benuskova, L. & Williams, J. M. Dynamical properties
of gene regulatory networks involved in long-term potentiation. *Front Mol*
Neurosci **8**, 42 (2015). https://doi.org:10.3389/fnmol.2015.00042
7   Sheng, M. & Greenberg, M. E. The regulation and function of c-fos and other
immediate early genes in the nervous system. *Neuron* **4**, 477-485 (1990).
https://doi.org:10.1016/0896-6273(90)90106-p
8   Wu, Y. E., Pan, L., Zuo, Y., Li, X. & Hong, W. Detecting Activated Cell
Populations Using Single-Cell RNA-Seq. *Neuron* **96**, 313-329 e316 (2017).
https://doi.org:10.1016/j.neuron.2017.09.026
9   Hu, P. *et al.* Dissecting Cell-Type Composition and Activity-Dependent
Transcriptional State in Mammalian Brains by Massively Parallel Single-Nucleus
RNA-Seq. *Mol Cell* **68**, 1006-1015 e1007 (2017).
https://doi.org:10.1016/j.molcel.2017.11.017
10   Vallejos, C. A., Risso, D., Scialdone, A., Dudoit, S. & Marioni, J. C. Normalizing
single-cell RNA sequencing data: challenges and opportunities. *Nat Methods* **14**,
565-571 (2017). https://doi.org:10.1038/nmeth.4292
11   Hicks, S. C., Townes, F. W., Teng, M. & Irizarry, R. A. Missing data and technical
variability in single-cell RNA-sequencing experiments. *Biostatistics* **19**, 562-578
(2018). https://doi.org:10.1093/biostatistics/kxx053
12   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell
RNA-seq data using regularized negative binomial regression. *Genome Biol* **20**,
296 (2019). https://doi.org:10.1186/s13059-019-1874-1
13   Lahnemann, D. *et al.* Eleven grand challenges in single-cell data science.
Genome Biol **21**, 31 (2020). https://doi.org:10.1186/s13059-020-1926-6
14   Yao, Z. *et al.* A taxonomy of transcriptomic cell types across the isocortex and
hippocampal formation. *Cell* **184**, 3222-3241 e3226 (2021).
https://doi.org:10.1016/j.cell.2021.04.021

15   Spiegel, I. *et al.* Npas4 regulates excitatory-inhibitory balance within neural
circuits through cell-type-specific gene programs. *Cell* **157**, 1216-1229 (2014).
https://doi.org:10.1016/j.cell.2014.03.058
16   Fernandez-Albert, J. *et al.* Immediate and deferred epigenomic signatures of in
vivo neuronal activation in mouse hippocampus. *Nat Neurosci* **22**, 1718-1730
(2019). https://doi.org:10.1038/s41593-019-0476-2
17   Tyssowski, K. M. *et al.* Different Neuronal Activity Patterns Induce Different Gene
Expression Programs. *Neuron* **98**, 530-546 e511 (2018).
https://doi.org:10.1016/j.neuron.2018.04.001
18   Sundararajan, M., Taly, A. & Yan, Q. Axiomatic Attribution for Deep Networks.
arXiv:1703.01365 (2017).
<https://ui.adsabs.harvard.edu/abs/2017arXiv170301365S>.
19   Gouwens, N. W. *et al.* Integrated Morphoelectric and Transcriptomic
Classification of Cortical GABAergic Cells. *Cell* **183**, 935-953 e919 (2020).
https://doi.org:10.1016/j.cell.2020.09.057
20   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-
3587 e3529 (2021). https://doi.org:10.1016/j.cell.2021.04.048
21   Thomas, G. M. & Huganir, R. L. MAPK cascade signalling and synaptic plasticity.
Nat Rev Neurosci **5**, 173-183 (2004). https://doi.org:10.1038/nrn1346
22   Rosen, L. B., Ginty, D. D., Weber, M. J. & Greenberg, M. E. Membrane
depolarization and calcium influx stimulate MEK and MAP kinase via activation of
Ras. *Neuron* **12**, 1207-1221 (1994). https://doi.org:10.1016/0896-6273(94)90438-
3
23   Baraban, S. C., Taylor, M. R., Castro, P. A. & Baier, H. Pentylenetetrazole
induced changes in zebrafish behavior, neural activity and c-fos expression.
Neuroscience **131**, 759-768 (2005).
https://doi.org:10.1016/j.neuroscience.2004.11.031
24   Dhir, A. Pentylenetetrazol (PTZ) kindling model of epilepsy. *Curr Protoc Neurosci*
Chapter 9, Unit9 37 (2012). https://doi.org:10.1002/0471142301.ns0937s58
25   Savell, K. E. *et al.* A dopamine-induced gene expression signature regulates
neuronal function and cocaine response. *Sci Adv* **6**, eaba4221 (2020).
https://doi.org:10.1126/sciadv.aba4221
26   Luo, Z., Volkow, N. D., Heintz, N., Pan, Y. & Du, C. Acute cocaine induces fast
activation of D1 receptor and progressive deactivation of D2 receptor striatal
neurons: in vivo optical microprobe [Ca2+]i imaging. *J Neurosci* **31**, 13180-13190
(2011). https://doi.org:10.1523/JNEUROSCI.2369-11.2011
27   Boulting, G. L. *et al.* Activity-dependent regulome of human GABAergic neurons
reveals new patterns of gene regulation and neurological disease heritability. *Nat
Neurosci* **24**, 437-448 (2021). https://doi.org:10.1038/s41593-020-00786-1
28   Rienecker, K. D. A., Poston, R. G. & Saha, R. N. Merits and Limitations of
Studying Neuronal Depolarization-Dependent Processes Using Elevated
External Potassium. *ASN Neuro* **12**, 1759091420974807 (2020).
https://doi.org:10.1177/1759091420974807
29   Rienecker, K. D. A. *et al.* Mild membrane depolarization in neurons induces
immediate early gene transcription and acutely subdues responses to a

successive stimulus. *J Biol Chem* **298**, 102278 (2022).
https://doi.org/10.1016/j.jbc.2022.102278
30 Hrvatin, S. *et al.* Single-cell analysis of experience-dependent transcriptomic
states in the mouse visual cortex. *Nat Neurosci* **21**, 120-129 (2018).
https://doi.org/10.1038/s41593-017-0029-5
31 Chatterjee, S. *et al.* Endoplasmic reticulum chaperone genes encode effectors of
long-term memory. *Sci Adv* **8**, eabm6063 (2022).
https://doi.org/10.1126/sciadv.abm6063
32 Wang, F. *et al.* RNAscope: a novel in situ RNA analysis platform for formalin-
fixed, paraffin-embedded tissues. *J Mol Diagn* **14**, 22-29 (2012).
https://doi.org/10.1016/j.jmoldx.2011.08.002
33 Kalmbach, B. E. *et al.* Signature morpho-electric, transcriptomic, and dendritic
properties of human layer 5 neocortical pyramidal neurons. *Neuron* **109**, 2914-
2927 e2915 (2021). https://doi.org/10.1016/j.neuron.2021.08.030
34 Hrvatin, S. *et al.* Publisher Correction: Single-cell analysis of experience-
dependent transcriptomic states in the mouse visual cortex. *Nat Neurosci* **21**,
1017 (2018). https://doi.org/10.1038/s41593-018-0112-6
35 Durinck, S. *et al.* BioMart and Bioconductor: a powerful link between biological
databases and microarray data analysis. *Bioinformatics* **21**, 3439-3440 (2005).
https://doi.org/10.1093/bioinformatics/bti525
36 Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the
integration of genomic datasets with the R/Bioconductor package biomaRt. *Nat*
Protoc **4**, 1184-1191 (2009). https://doi.org/10.1038/nprot.2009.97
37 Zerbino, D. R. *et al.* Ensembl 2018. *Nucleic Acids Res* **46**, D754-D761 (2018).
https://doi.org/10.1093/nar/gkx1098
38 Plaisier, S. B., Taschereau, R., Wong, J. A. & Graeber, T. G. Rank-rank
hypergeometric overlap: identification of statistically significant overlap between
gene-expression signatures. *Nucleic Acids Res* **38**, e169 (2010).
https://doi.org/10.1093/nar/gkq636
39 McCarthy, D. J., Campbell, K. R., Lun, A. T. & Wills, Q. F. Scater: pre-
processing, quality control, normalization and visualization of single-cell RNA-seq
data in R. *Bioinformatics* **33**, 1179-1186 (2017).
https://doi.org/10.1093/bioinformatics/btw777
40 Eraslan, G., Simon, L. M., Mircea, M., Mueller, N. S. & Theis, F. J. Single-cell
RNA-seq denoising using a deep count autoencoder. *Nat Commun* **10**, 390
(2019). https://doi.org/10.1038/s41467-018-07931-2
41 Choudhary, S. & Satija, R. Comparison and evaluation of statistical error models
for scRNA-seq. *Genome Biol* **23**, 27 (2022). https://doi.org/10.1186/s13059-021-
02584-9
970

Figure 1 - Schematic Overview of NEUROeSTIMator. The graphical abstract illustrates the application of NEUROeSTIMator, a deep learning model, to estimate neuronal activity. Transcriptome-wide gene expression data is input to an autoencoder, which reconstructs expression of immediate early genes from a single unit latent space called the 'activity score'. The activity score can be applied to gene expression data, including single cell RNA-sequencing, Patch-seq, or spatial transcriptomics studies for detecting transcriptomic signatures of activity.

Figure 2 – Electrophysiological features of neuronal stimulation captured by transcriptome-based activity score.
(A) Activity score distributions compared between stimulated Patch-seq neurons (gold) and unstimulated SMART-seq neurons (violet). (B) Observed D statistics (red line) and empirical distribution from Kolmogorov-Smirnov tests between activity score distributions of Patch-seq and SMART-seq datasets, using one thousand permutations of shuffled expression values . (C) Associations between electrophysiological features and predicted activity. Heatmap shows feature value columns ordered by the bootstrap coefficient, and individual neuron rows ordered by lasso model prediction (top). Summary of lasso bootstrap permutations. Dots represent the mean coefficient estimate across bootstraps, lines indicate standard error. Significant features depicted by colored dots (bottom). (D) Spike train comparison of two Vip Ptprt Pkp2 neurons exhibiting representative electrophysiological features of high (red) and low (blue) NEUROeSTIMator predictions (ramp current injected - top panel, membrane potential - bottom panel) (E) Patch-seq validation of the positive association between input resistance and NEUROeSTIMator predictions in a novel set of human excitatory neurons. Pearson's correlation test statistics annotated; linear fit (orange line).

974
975

A

detecting activation of mouse neurons
response to seizure induced by pentylenetetrazol (PTZ)

☐ saline ■ seizure-affected

B

detecting activation of rat nucleus accumbens neurons
response to cocaine injection

☐ saline ■ cocaine

C

detecting activation of human cultured neurons
temporal response to potassium chloride depolarization

☐ KCl_0hr ■ KCl_1hr ☐ KCl_2hr ☐ KCl_4hr

Figure 3 – Multi-species generalization of neuronal activity score applied to previously published chemical induction studies. Predicted activity for various neuron types responding to treatment with potent chemical modulators of neuron activity. **(A)** Response to PTZ-induced seizure (red) and controls (white) in mouse cortical neurons. **(B)** Cocaine treatment (red) or controls (white) in rat nucleus accumbens neurons. **(C)** Time series of predicted activity for human iPSCs treated with depolarizing potassium chloride at 0 hours (control), 1 hour, 2 hours, and 4 hours. Annotated statistics from Kolmogorov-Smirnov test are provided in **Supplementary Table 5**.

976
977

Figure 4 - Temporal patterns and classification of *in vivo* sensory activation. (**A**) Cell type activity predictions of visual cortex neurons in freely behaving mice exposed to light for 0, 1, or 4 hours. Activity predictions summarized by cell type and experimental group, and Kolmogorov-Smirnov test results are provided in **Supplementary Table 6**. (**B**) ROC plots indicate the ability of predicted activity to distinguish between the 0h and 1h experimental groups. Diagonals from bottom left to top right indicate an accuracy similar to random chance, while lines moving straight vertically, then straight horizontally indicate perfect separation.

Figure 5 – Spatial transcriptomic patterns of neuronal activation after spatial learning.
(**A**) Spatial anatomical clustering of RNA-sequencing spots. (**B**) Activity score per spot, averaged within experimental groups of homecage controls (left) and 1 hour after SOR training (right). Using the spatial anatomical clustering derived from the expression data, we were able to group individual spots into clusters to. (**C**) Brain region-specific induction of activity following SOR training. Differences tested using linear models, colored by the SOR coefficient estimate. Cluster-wise differential activity statistics are summarized in (**D**) and provided in **Supplementary Table 7**. Bar length represents estimated coefficient of SOR on activity score. Brackets indicate standard error and circle size represents the number of spots per comparison, an index of statistical power.

Figure S1 - Selection of target genes to be reconstructed by autoencoder. (**A**) A principle component analysis (PCA) of 29 candidate target genes reveals the dominance of the first component (PC1) over the other components, suggesting PC1 captures the majority of the underlying structure in the data. (**B**) Gene loadings onto PC1. The red line marks the outlier threshold for extreme loadings, as defined by a shuffled null PCA. (**C**) Number of bootstrap permutations where feature loadings of genes surpassed the outlier threshold in random subsets of samples. The red line indicates where loadings surpassed thresholds in 5% of permutations. Genes surpassing this threshold are colored blue, and these genes were selected as the final target genes used by the autoencoder. These findings show that the expression pattern of these genes is not overly complex and is suitable to be modeled as a single dimension.

Figure S2 - Weighted downsampling improves balance and preserves heterogeneity. (A) Shannon entropy values to measure balance of cell groupings across multiple downsampling thresholds. Downsampling to 500,000 cells (dashed line) offers a balance of sample numbers and group balance, particularly for species (red), cell class (gold), and cell subclass (light blue). (B) Hierarchical downsampling visualized. Hexagon bin colors represents the number of cells in a given x,y coordinate range before downsampling (left) and after downsampling (right). The y-axis represents a cell's target gene PC1 score, and the x-axis shows total unique molecular identifier (UMI) counts per cell. Downsampling with hierarchical weighting preserves diversity in both biological and technical sources of variation.

984
985

Figure S3 - Feature attribution of model predictions. (**A**) We compared the mean feature attribution for target genes (blue) to non-target genes (gray) split by the number of stimulus-responsive differential expression lists they appeared in from the target selection analysis. Target genes have higher average importance in model predictions, but mean feature attribution for non-target genes is highest for genes appearing in two or three differential expression lists. (**B**) Gene set enrichment results for feature attributions. We examined enrichment scores for several gene sets using feature attributions of all input genes (light bar) or non-target genes (dark bar). Asterisks indicate the gene set remained significantly enriched when target genes were removed from the analysis (see **Supplemental Table 3**).

Figure S4 - Analysis of Patch-seq data with electrophysiological and transcriptional perspectives.
(**A**) Seurat integration of Patch-seq and Smart-seq datasets. (**B**) Comparison of mean gene expression between Patch-seq and Smart-seq, with NEUROeSTIMator target genes colored pink. (**C**) Principle component analysis of electrophysiological features from the Patch-seq dataset. Component variances (top), their correlation with NEUROeSTimator output (middle), and contribution to total variance explained (bottom). (**D**) Gene set enrichment analysis of gene loadings from canonical correlation analysis for GO: Biological Processes (GOBP), GO: Molecular Function (GOMF), Reactome, and Wikipathways (WP) (Supplementary data 1).

Figure S5 - Spatial transcriptomic techniques reveal region-specific activity in response to spatial learning.
(**A**) Predicted activity of individual Visium spots for each sample from a homecage control group (top) and an SOR-trained group (bottom). (**B**) RNAScope validation of region-specific activation *Egr1*, *Nr4a1*, and *Egr3*. Scale bar represents 1000 um in length. (**C**) Image comparison of learning induced activity as measured by RNAScope (top) and Visium (bottom). Scale bar depicts 500 um in length. (**D**) Dorsal hippocampus subregion-specific induction of transcriptional activity following SOR.

Figure S6 - Cell Type Deconvolution of Spatial Transcriptomic Data. (A) Results of cell type deconvolution in Visium samples, colored by estimated proportion of cell types. (B) Quadratic relationship between estimated proportion of neuronal cell types and estimated effect of learning (C). Estimated effects of learning, adjusted for estimated proportion of neurons. Adjusted estimates are provided in **Supplementary Table 7**.

992
993

Figure S7 - Generalized Improvement in Classification Accuracy Over Competing Methods.
Comparison of classification accuracy (AUC) of experimental stimulation groups for the NEUROeSTIMator model and three competing methods. Cells are grouped by datasets, cell types, and time points when applicable. Number of samples per comparison is shown on the left of the plot and further columns correspond to the method being compared. Positive x-axis values indicate NEUROeSTIMator classified the experimental groups with greater accuracy than the method being compared.

Figure S8 - Benchmarking performance across technical and biological variables. Comparing classification accuracy across technical and biological groups. Positive values (red) indicate NEUROeSTIMator classified the experimental groups with greater accuracy than the method being compared. Column titles indicate the method being compared, x-axis groups correspond to time passed since onset of stimulation, and y-axis groups are binned by $\log_{10}$ sum of UMI counts.

Figure S9 - Candidate Target Gene Selection. (**A**) Rank-rank hypergeometric overlap (RRHO) was used to identify p-value thresholds of maximum enrichment between differential expression lists for potassium chloride depolarization in glutamatergic-enriched and GABAergic-enriched neuron samples. Log fold change thresholds were set at one (**B**) to select the final set of candidate targets from genes below RRHO p-value thresholds in both lists (**C**).

Figure S10 - Graphical representation of model architecture. The architecture NEUROeSTIMator was trained with includes an input for expression of target genes and another for the rest of the transcriptome. Inputs are concatenated before being passed to an encoder layer consisting of a dense fully-connected layer, a batch-normalization layer, and an activation layer. Diverging branches with additional encoder units lead to dispersion and dropout estimates of the zero-inflated negative binomial model (right branch), or a single unit bottleneck which estimates the mean parameter (left branch). The parameter estimates are provided with contextual information about cell type, dataset, and quality control metrics from an additional series of encoder layers (middle branch). Importantly, in application, NEUROeSTIMator utilizes only the left branch through the bottleneck activation layer.