1
2
3 **A novel lineage of the *Capra* genus discovered in the Taurus Mountains of**
4 **Turkey using ancient genomics.**
5

6 Kevin G. Daly[1], Benjamin S. Arbuckle[2], Conor Rossi[1], Valeria Mattiangeli[1], Phoebe A. Lawlor[1],
7 Marjan Mashkour[3,4], Eberhard Sauer[5], Joséphine Lesur[3], Levent Atici[6], Cevdet Merih Erek[7],
8 Daniel G. Bradley[1]
9

[1]Smurfit Institute of Genetics, Trinity College Dublin, Dublin 2, Ireland
[2]Department of Anthropology, University of North Carolina at Chapel Hill, 301 Alumni Building,
Chapel Hill, NC 27599 USA
[3]Centre National de Recherche Scientifique, / Muséum national d'Histoire naturelle,
Archéozoologie, Archéobotanique, (AASPE)-UMR 7209, CP 56, 55 rue Buffon, 75005 Paris,
France
[4]University of Tehran, Bioarchaeology Laboratory, (Central Laboratory), Archaeozoology
section, Engelab av. Qods av., Poursina av., Jalalieh str. N° 6, CP1417634934, Tehran, Iran
[5]School of History, Classics and Archaeology, University of Edinburgh, Edinburgh EH8 9AG, UK
[6]Department of Anthropology, University of Nevada, Las Vegas, Las Vegas, Nevada, USA
[7]Ankara Hacı Bayram Veli University, Faculty of Letters, Department of Archeology, Department
of Prehistoric Archeology, Emniyet Mah. Abant 1 cad. 10/2D, Kat 3, Oda no 320, Yenimahalle,
Ankara, Turkey
23
Corresponding author: Kevin G. Daly, dalyk1@tcd.ie, ORCID: 0000-0002-5579-6144

Abstract

Direkli Cave, located in the Taurus Mountains of southern Turkey, was occupied by Late

Epipaleolithic hunters-gatherers for the seasonal hunting and processing of game including

large numbers of wild goats. We report genomic data from new and published *Capra* specimens

from Direkli Cave and, supplemented with historic genomes from multiple *Capra* species, find a

novel lineage best represented by a ~14,000 year old 2.59X genome sequenced from specimen

Direkli4. This newly discovered *Capra* lineage is a sister clade to the Caucasian tur species

(*Capra cylindricornis* and *Capra caucasica*), both now limited to the Caucasus region. West

Eurasian domestic goats in the past, but not those today, appear enriched for Direkli4-specific

alleles, and we further identify genomic regions introgressed in domestic goats with high affinity

to Direkli4. This forgotten "Taurasian tur" likely survived Late Pleistocene climatic change in a

Taurus Mountain refugia and its genomic fate is unknown.

Introduction

The genus *Capra* includes the domestic goat (*Capra hircus*) as well as a variety of wild

mountain-dwelling goat/ibex species distributed across Eurasia and North Africa including

several listed as endangered or vulnerable (Shackleton 1997; Pidancier et al. 2006). Nine

species are currently recognized by the IUCN; however, taxonomic relationships are still under

revision (Pidancier et al. 2006; Zheng et al. 2020). Among these, the status of the two species

endemic to the Caucasus Mountains has been debated (Parrini, Cain, and Krausman 2009;

Groves and Grubb 2011). The East Caucasian tur (*Capra cylindricornis*) has been considered

either a species distinct from the West Caucasian tur (*Capra caucasica*) or they comprise a

single species of two potentially-hybridizing populations (Heptner, Nasimovich, and Bannikov

1961). Moreover the bezoar (*Capra aegagrus*), progenitor of domestic goat, has also been

reported to hybridize with both tur varieties with which it shares seasonal grazing territories in

the Caucasus region (Pfitzenmayer 1915; Sarkisov 1953; P. J. Weinberg 2002). Interspecies

Capra gene flow is well known (Manceau et al. 1999; Pidancier et al. 2006; Kazanskaia,

Kuznetsova, and Danilkin 2007), and may explain discordant phylogenies across loci (Pidancier

et al. 2006; Ropiquet and Hassanin 2006). Such admixture may have shaped the evolution of

domestic goat; for example, the tur has been identified as a putative source of a *MUC6* allele

driven to fixation in domestic populations and likely selected for gastrointestinal parasite

resistance (Zheng et al. 2020; Grossen et al. 2020). Tur additionally shows differing affinity to

domestic and wild goat genomes indicating a complex evolutionary history of the genus.

Although tur are currently restricted to the Caucasus region, ancient wild goat specimens

recovered from Direkli Cave, a camp site used by Late Pleistocene hunters in the Central

Taurus Mountains of southern Turkey (Figures 1A, S1 and S2) (B. S. Arbuckle and Erek 2012),

were found to carry a tur-like mitochondrial lineage, designated T (Daly et al. 2018). Three of

these four reported ancient *Capra* specimen fall within the bezoar autosomal diversity, but a

fourth - Direkli4, dated to 12,164-11,864 cal BCE (Ramsey 2009; Reimer et al. 2020) and

sequenced here to 2.59X mean genome coverage (Table 1) - shows an excess of ancestral

alleles in *D* statistic tests (Green et al. 2010) when paired with domestic/bezoar goat (Figure S3,

Table S2) implying Direkli4 carries ancestry basal to that clade. To explore this signal further we

generated low coverage genomes from historic and rare *Capra* samples, including a 20th

century CE zoo-born East Caucasian tur (Tur2), tur specimens from the Dariali-Tamara Fort

archaeological site at Kazbegi, Georgia (Mashkour et al. 2020), a zoo-born Walia ibex, and

supplemented with published modern and ancient *Capra* genomes (Table S1, S3, Figure S4)

(Zheng et al. 2020; Grossen et al. 2020). Surprisingly, a neighbour joining tree from nuclear

genome identity-by-state (IBS) information places Direkli4 as sister to a clade of both Caucasian

tur taxa, a signal obtained using either goat- or sheep-aligned data (Figures 1B and S5). The

Direkli4 genome thus suggests a previously-unrecognized *Capra* lineage sister to both

Caucasian tur inhabited the Taurus Mountains ~14,000 years ago.

Materials and methods

DNA from 7 postcranial bone elements from Direkli Cave and 13 historic *Capra* specimen was

extracted via standard aDNA protocols (Yang et al. 1998), with a 0.5% sodium hypochlorite pre-

wash (Korlević et al. 2015) performed for the Direkli material. Following uracil excision (Rohland

et al. 2015) and dsDNA library construction (Meyer and Kircher 2010), libraries were subject to

shotgun sequencing (Illumina HiSeq 2000 and NovaSeq 6000) or RNA-bait enrichment of

mtDNA reads prior to shotgun sequencing. Additional sequencing data was also generated for

specimen Direkli4.

Using bwa aln (Li, Ruan, and Durbin 2008) a relaxed alignment (Meyer al. 2012) was performed

against the goat reference ARS1 (Bickhart et al. 2017) or an outgroup genome

(Oar_rambouillet_v1.0). Subsequent analyses were primarily performed in the ANGSD

environment (Korneliussen, Albrechtsen, and Nielsen 2014) using single read sampling. A more

detailed methodology is provided in the Supplementary Material, available at OSF (DOI

10.17605/OSF.IO/3ECQD).

Table 1: Sample providence and sequencing summary.

Sample	Morphological Species	Origin	Age	Sex[b]	Nuclear Cov.[c]	mtDNA Cov.
Direkli4	*Capra spc.*	E4/8A, Direkli Cave, Turkey	12,164-11,864 cal BCE (2σ)	M	2.59	642.23
Direkli9	*Capra spc.*	B6/5, Direkli Cave, Turkey	Est. 12,100-8,900 BCE[a]	M	0.0003	9.5
Direkli12	*Capra spc.*	B13/4A, Direkli Cave, Turkey	Est. 12,100-8,900 BCE	M	0.07	44.06
Direki13	*Capra spc.*	B13/7B, Direkli Cave, Turkey	Est. 12,100-8,900 BCE	M	0.0055	76.99
Direkli14	*Capra spc.*	D3/7, Direkli Cave, Turkey	Est. 12,100-8,900 BCE	M	0.0005	13.24
Direkli15	*Capra spc.*	B6/5, Direkli Cave, Turkey	Est. 12,100-8,900 BCE	F	0.0002	24.21
Direkli16	*Capra spc.*	B8/7, Direkli Cave, Turkey	Est. 12,100-8,900 BCE	F	0.01	15.11
Direkli17	*Capra spc.*	E5/5, Direkli Cave, Turkey	Est. 12,100-8,900 BCE	F	0.0001	0.0845
Caucasus1	*Capra caucasica*	Tamara Fort, Kazbegi, Georgia	4th-21st c. CE, probably 5th-15th c. CE	M	0.55	64.63
Caucasus2	*Capra caucasica*	Tamara Fort, Kazbegi, Georgia	4th-21st c. CE, probably 5th-15th c. CE	M	0.0021	379.71
Caucasus3	*Capra caucasica*	Tamara Fort, Kazbegi, Georgia	4th-21st c. CE, probably 5th-15th c. CE	M	0.004	464.51
Falconeri1	*Capra falconeri hepteneri*[d]	Unknown, via Parc de la Haute-Touche	20th Century CE	M	0.58	72.11
Falconeri2	*Capra falconeri*	Born at MNHN Zoo, Paris	20th Century CE	M	0.06	45.78
Ibex1	*Capra ibex*	Unknown	20th Century CE	F	3.93	179.03
Ibex2	*Capra ibex*	Pointe de Calabre, Savoie	20th Century CE	M	0.05	21.4
Sibirica1	*Capra sibirica*	Born at MNHN Zoo, Paris	20th Century CE	M	0.04	163.16
Sibirica2	*Capra sibirica*	Born at MNHN Zoo, Paris	20th Century CE	F	1.48	205.78
Tur2	*Capra cylindricornis*	Unknown, via Vincennes Zoo	20th Century CE	F	0.02	4.96
Walie1	*Capra walie*	Born at MNHN Zoo, Paris	20th Century CE	M	0.75	103.12
Pyrenica2	*Capra pyrenica*	Unknown	20th Century CE	M	0.16	51.2
Nubiana1	*Capra nubiana*	Unknown	20th Century CE	M	1.25	211.32

97 [a]Estimated ages for Direkli material is based on calibrated ages from the cave stratigraphy.

98 [b]M=Male, F=Female

99 [c]Cov. = coverage.

100 [d.]Falconeri1, a likely Barbary sheep (see Supplementary Methods).

Figure 1: A) Elevation map of southwest Asia. Key sites are indicated, with *C. caucasica* and *C. cylindricornis* distributions from (Gavashelishvili et al. 2018) displayed. MSL=metres above mean sea level. B) Neighbour joining phylogeny of genomes >0.5X using pairwise IBS and transversions, rooted on Sheep (not shown, as well as a likely Barbary Sheep sample Falconeri1, see Supplementary Methods). Node support from 100 replicates using 50 5Mb regions sampled without replacement shown when <100. Pink=Direkli4, green=other genomes first reported here.

Results

Our additional screening of Direkli Cave *Capra* remains identified seven with surviving DNA (Table S4); two genomes show greater affinity to Direkli4 than bezoar from the same site (Figure 2A). A total of three out of the eleven Direkli Cave *Capra* specimens therefore are assigned to the tur-related clade, implying that while less numerous than bezoar, members of this clade were not rare in the region in the Late Pleistocene. Nuclear genome types (tur-like or bezoar) do not necessarily co-associate with mitochondrial lineages (tur T and bezoar F), with all combinations except "tur-like genome, tur-like mitochondria" observed (Figure 2B, Figures S6 and S7, Table S5), establishing that there was gene flow between these lineages. Additionally, there is little variation among Direkli T mtDNA (average 4.07 pairwise differences, compared to 67 among Direkli F mtDNA), suggesting a recent, single introgression origin for this Direkli tur-like matrilineage.

A tur-like population in the Taurus Mountains is consistent with the high variability in body size of the *Capra* material at Direkli Cave where extremely large *Capra* remains have been reported alongside smaller bezoar-size individuals (Figure S8). Extant tur exhibit body weights 20-50% larger than bezoar (Masseti 2009; Castelló, Huffman, and Groves 2016) and it is plausible that the 'large' *Capra* from Direkli represent tur-lineage animals. Although there are clear differences between bezoar and tur horn morphologies (Pidancier et al. 2006), unfortunately, diagnostic horncore remains have not been recovered from Direkli. The cave was initially inferred to be occupied primarily during summer months (B. S. Arbuckle and Erek 2012), with subsequent discoveries of architectural remains and zooarchaeological analyses indicating more intensive use (Benjamin S. Arbuckle 2019). The presence of tur-related goats may reflect use of the cave in the winter months when, based on Caucasian analogs (Gavashelishvili 2009), tur would be expected to descend from the higher elevations surrounding the cave (>2000m above sea level).

Given the Direkli4 genome was recovered together with bezoar specimens, the two lineages of *Capra* likely had proximate ranges and hybridised. We use *D* statistics (Green et al. 2010) to measure Direkli4 derived allele sharing relative to either a likely-hunted (Table S6) or likely-herded (Table S7, Pearson's *r* >0.99) ~10,000 year old goat from the Zagros Mountains. A Late Pleistocene wild goat from the Armenian Lesser Caucasus, Hovk1, shows highest affinity with Direkli4 (Figure S9). Bezoar goats from Direkli Cave also show high Direkli4 allele sharing,

Figure 2: A) *D* statistic test of affinity using specimens from Direkli Cave and a historic *C. caucasica* individual for reference. Positive values indicate sample X has greater affinity with the Tur-like Direkli4 genome; negative values indicate greater affinity with the *C. aegagrus* Direkli1-2. Error bars represent 3 standard errors. B) ML phylogeny of mtDNA, abbreviated. Bootstrap node support values (100 replicates) are displayed when <100. The complete phylogeny including likely Barbary sheep Falconeri1 is displayed in Figure S6. T haplogroup is as defined by (Daly et al. 2018) Low coverage sample Direkli17 is displayed in a highly reduced phylogeny Figure S7, C=collapsed. *=Direkli sample with discordant mtDNA and nuclear genome affinity.

mirroring affinity measures with west Caucasian tur (Zheng et al. 2020). While directionality is

uncertain, these statistics imply gene flow between the tur-like lineage and wild bezoar.

Examining domestic goats we find that Neolithic genomes from Europe show greater affinity to

Direkli4 (Figure S9), but Neolithic Iranian goats do not, echoing the distribution of Direkli bezoar-

related ancestry in West Eurasian populations (Daly et al. 2018). We account for possible gene

flow from Caucasian tur into modern European goat using the statistic D(Tur1, Direkli4; X,

Sheep) to compare relative affinity with Tur1 and Direkli4 (Table S8). With the exception of two

other tur samples, all examined domestic/bezoar goats show either a bias towards Direkli4 or

gave a non-significant result, consistent with Direkli4-related admixture or a more complicated

genetic history.

Genetic exchange between bezoar and the ancestors of Direkli4 could confound these

measures of shared variation among domestic populations. We identified variants specific to

Direkli4, conditioned on ancestral allele fixation in a range of defined groups (Figure 3A,

Supplementary). Using this we calculate a statistic analogous to the D statistic, here termed the

extended D or D_{ex}. Relative to Neolithic Zagros goats, ancient domestic genomes from western

Eurasia have an excess of Direkli4-specific variants (Figure 3B, Figure S10, Table S9). This

"Direkli4-specific" allele sharing signal is absent in ancient goats from Iran-eastwards, and in all

tested modern goat (Figure 3C). To control for possible reference biases, we calculated D_{ex}

ascertaining on variants segregating in sheep (Table S10) and recovered similar results

(Pearson's r = 0.9935). Repeating the analysis using other ancient/historic *Capra* "specific"

alleles shows somewhat correlated results (Table S11), but the distinct patterns of allele sharing

(Figure S11, Table S12) imply that Direkli4 ancestry in domestic goat varies temporally and

geographically.

We next identify alleles derived in Direkli4 also at a low frequency (>0%, ≤10%) in other *Capra*

and bezoar, and then measure their abundance in domestic goats. The west Caucasian tur

(Tur1) most frequently shares derived alleles with Direkli4 and domestic goat (Figure S12),

consistent with their cladal relationship (although this measure is sensitive to genome depth,

Figure 3: A) Extended D statistic. To control for gene flow with the *Capra* genus, we condition on variants derived in Direkli4 (H3) and a genome X (H2), but ancestral in other populations (here: Sheep, non-bezoar *Capra* genomes, bezoar from Direkli Cave, and other bezoar). Values are calculated relative to a set of Neolithic goat from the Zagros Mountains, and normalized similarly to the D statistic. B) Extended D statistic values for Direkli4 using transversions, C) plotted through time. Each symbol is a test genome, with shape denoting time period. Black borders indicates a |Z| score ≥ 3, using 1000 bootstrap replicates and 5Mb blocks.

Supplementary Methods). Ancient European domestic goats share a higher proportion of alleles with both Direkli4 and the high-coverage bezoar from Direkli Cave, Direkli1-2. In comparison, Modern European and African goats carry variation present in Direkli4 plus one of the two Caucasian tur (Tur1 and Caucasus1). This discrepancy could be explained by either gene flow from domestic goats into tur during the last 8,000 years, or alternatively an increase in tur-Direkli4 related ancestry in European populations over time.

Investigating gene flow events within *Capra*, automated tree-based model exploration (Pickrell

and Pritchard 2012) detects admixture between the Direkli4/Tur lineage and the ancestors of

the Late Pleistocene bezoar Hovk-1 (Figure S13). Residuals of this graph point to unmodelled

affinity between Direkli4 and both Direkli Cave bezoar and with Neolithic Serbian domestic goat

(Figure S14). Modern European goat do not show unmodeled Direkli4 affinity, supporting the

interpretation that Direkli4-related ancestry has declined with time in west Eurasian goats. A

reduced set of populations explored using ML network orientation (Molloy, Durvasula, and

Sankararaman 2021) reiterates the Tur1/Direkli4 and Direkli bezoar lineages admixture, and

also between Direkli bezoar and domestic goat (Figure S15). Investigating admixture graph

space (Supplementary Methods) reiterates gene flow between Direkli bezoar and Direkli4, as

well as genetic differentiation of Neolithic and modern European goats (Figure S16). However

our data presently cannot distinguish between 1-3 gene flow events or gene flow directionality

(Table S13).

We finally identify 3 out of 112 regions introgressed from other *Capra* species to domestic goat

(Zheng et al. 2020) which show high affinity with Direkli4 (Figures S17-19, Supplementary Data

Files 1 and 2). A further 7 regions appear to have most affinity with the Direkli4-tur clade

(Figures S20-28), including a locus encompassing *MUC6*, a target of selection in domestic

goats during the last 10,000 years (Zheng et al. 2020), implicating the Direkli4 lineage in the

makeup of domestic goat gene pool.

Discussion

Our results indicate that a lineage related to the Caucasian tur existed in the Taurus Mountains

during the Late Pleistocene, as late as the 12th millennium cal BCE. We designate this novel

taxa as *Capra taurensis*, sister species of the tur *C. caucasica* and *C. cylindricornis* following

IUCN convention (P. Weinberg and Lortkipanidze 2020) or *Capra caucasica taurensis* under a

subspecies classification (Wilson and Reeder 2005). This "Taurasian tur" may have diverged

from the Caucasian lineages 130-200kya based on mtDNA coalescent estimates (Daly et al.

2018; Bouckaert et al. 2014). The current distribution of *Capra* species is mostly discontinuous

and is suggestive of climate-induced fragmentation (Shackleton 1997). The ancestors of

Caucasian tur likely extended over a broader range in Eurasia during the Late Pleistocene but

may have been poorly captured by the fossil record (Uerpmann 1987; Crégut-Bonnoure 1991;

P. J. Weinberg 2002). The large variability and high upper size range of *Capra* remains are

consistent with both smaller-bodied bezoar and larger-bodied tur-relatives being present within

the faunal assemblage at Direkli as well as other sites in the central Taurus and Lebanese

mountains (Üçagızlı cave, Ksar Akil, Saaide II), but not in the western Taurus where only bezoar

are evident ([Figure S8](#)) (B. S. Arbuckle and Erek 2012). *C. taurensis* could have survived the

Last Glacial Maximum within the central Taurus Mountains, a plausible refugia for *Capra*

species in addition to the Pontic and Anti-Taurus ranges (Gavashelishvili et al. 2018) while

experiencing a severe matrilineal bottleneck (Figure 2B). *C. taurensis* appears to have produced

fertile offspring with other members of the *Capra* genus; the traces of shared ancestry in ancient

bezoar and likely managed goat (Figure 3B) may be the consequence of direct gene flow or

secondarily via admixed bezoar. Gene flow between early managed goats and Anatolian bezoar

carrying *C. taurensis* ancestry could partially explain the divergence between Zagros and more

westerly herds.

Given the tremendous pressure on *Capra* species via Anthropocene over-hunting and habitat

disruption (Shackleton 1997), it is assumed that *C. taurensis* is extinct, with its existence only

now revealed via palaeogenomics. The Caucasian tur's preference for snowier habitats

(Gavashelishvili et al. 2018) combined with the lower altitude of the Taurus Mountains relative to

the Caucasus (Figure 1A) may have rendered the lineage vulnerable to climatic change via

Holocene warming and interspecific competition with bezoar, which are still found in the Taurus

mountains (Naderi et al. 2008, Gavashelishvili 2009), leading to its hypothesised extinction. As

the history of *C. taurensis* following the Late Pleistocene is still unknown, further genomic

surveys of Holocene *Capra* remains and present-day populations from this and adjacent regions

may illuminate its genetic legacy.

Acknowledgements

We thank Matthew Teasdale for his advice on interpretation of results. Excavations at Direkli

Cave are sponsored by T.C. Kültür ve Turizm Bakanlığı. Permission to export samples from

Direkli Cave provided by T.C. Kültür Varlıkları ve Müzeler Genel Müdürlüğü and T.C. Ankara

Valılığı, Il Kültür ve Turizm Müdürlüğü, Anadolu Medeniyetleri Müzesi Müdürlüğü (#70583208-

160.99(06)-899).

Funding

Zooarchaeological work at Direkli has been supported by grants from the Office of Vice Provost

for Research, Baylor University and a URC grant from the Office of the Vice Chancellor for

Research at the University of North Carolina at Chapel Hill. This work was supported by the

European Research Council under the European Union's Horizon 2020 research and innovation

programme (grant numbers 885729-AncestralWeave, 295729-CodeX, 295375-Persia and its

Neighbours); and supported in part by a Grant from Science Foundation Ireland under grant

number 21/PATH-S/9515.

276 Conflict of interest disclosure

277 The authors declare they have no conflict of interest relating to the content of this article.

279 Data, script and code availability

280 *Capra taurensis* has been registered under the Zoobank LSID

281 urn:lsid:zoobank.org:act:1261A42B-B0C0-4571- 87F4-8EC3B5381A88. Raw sequencing reads,

282 aligned QCed final bam files, and mitochondrial fasta files have been deposited in ENA under

283 the project accession PRJEB51668. Scripts for extended D calculation/disentangling derived

284 allele sharing are available at https://github.com/Xevkin/direkli_caprid_extended_ds/tree/main.

286 <u>References</u>

Arbuckle, Benjamin S. 2019. "Zooarchaeology at Epipaleolithic Direkli Cave, Kahramanmaras,
Turkey." *Journal of Anatolian Prehistoric Research/Anadolu Prehistorya Araştırmaları*
Dergisi 5: 1–14.
Arbuckle, B. S., and C. M. Erek. 2012. "Late Epipaleolithic Hunters of the Central Taurus:
Faunal Remains from Direkli Cave, Kahramanmaraş, Turkey." *International Journal of*
Osteoarchaeology 22 (6): 694–707.
Bickhart, Derek M., Benjamin D. Rosen, Sergey Koren, Brian L. Sayre, Alex R. Hastie, Saki
Chan, Joyce Lee, et al. 2017. "Single-Molecule Sequencing and Chromatin Conformation
Capture Enable de Novo Reference Assembly of the Domestic Goat Genome." *Nature*
Genetics 49 (4): 643–50.
Bouckaert, Remco, Joseph Heled, Denise Kühnert, Tim Vaughan, Chieh-Hsi Wu, Dong Xie,
Marc A. Suchard, Andrew Rambaut, and Alexei J. Drummond. 2014. "BEAST 2: A Software
Platform for Bayesian Evolutionary Analysis." *PLoS Computational Biology* 10 (4):
e1003537.
Castelló, José R., Brent Huffman, and Colin Groves. 2016. *Bovids of the World: Antelopes,*
Gazelles, Cattle, Goats, Sheep, and Relatives. Vol. 104. Princeton Field Guides. Princeton
University Press.
Crégut-Bonnoure, Evelyne. 1991. "Intérêt Biostratigraphique de La Morphologie Dentaire de
Capra (Mammalia, Bovidae)." *Annales Zoologici Fennici* 28 (3/4): 273–90.
Daly, Kevin G., Pierpaolo Maisano Delser, Victoria E. Mullin, Amelie Scheu, Valeria Mattiangeli,
Matthew D. Teasdale, Andrew J. Hare, et al. 2018. "Ancient Goat Genomes Reveal Mosaic
Domestication in the Fertile Crescent." *Science* 361 (6397): 85–88.
Gavashelishvili, Alexander. 2009. "GIS-Based Habitat Modeling of Mountain Ungulate Species
in the Caucasus Hotspot." In *Status and Protection of Globally Threatened Species in the*
Caucasus, edited by N. Zazanashvili and D. Mallon, 74–82. Tbilisi: CEPF, WWF.
Gavashelishvili, Alexander, Yuriy A. Yarovenko, Elmar A. Babayev, Giorgi Mikeladze, Zurab
Gurielidze, Davit Dekanoidze, Niko Kerdikoshvili, Levan Ninua, and Nika Paposhvili. 2018.

"Modeling the Distribution and Abundance of Eastern Tur (Capra Cylindricornis) in the Caucasus." *Journal of Mammalogy*, June. https://doi.org/10.1093/jmammal/gyy056.

Green, Richard E., Johannes Krause, Adrian W. Briggs, Tomislav Maricic, Udo Stenzel, Martin Kircher, Nick Patterson, et al. 2010. "A Draft Sequence of the Neandertal Genome." *Science* 328 (5979): 710–22.

Grossen, Christine, Frédéric Guillaume, Lukas F. Keller, and Daniel Croll. 2020. "Purging of Highly Deleterious Mutations through Severe Bottlenecks in Alpine Ibex." *Nature Communications* 11 (1): 1001.

Groves, Colin, and Peter Grubb. 2011. *Ungulate Taxonomy*. Baltimore: JHU Press.

Heptner, V. G., A. A. Nasimovich, and A. G. Bannikov. 1961. *Mammals of the Soviet Union*. Edited by V. G. Heptner and N. P. Naumov. Moscow: Vysshaya Shkola.

Kazanskaia, E. Iu, M. V. Kuznetsova, and A. A. Danilkin. 2007. "[Phylogenetic reconstructions in the genus Capra (Bovidae, Artiodactyla) based on the mitochondrial DNA analysis]." *Genetika* 43 (2): 245–53.

Korneliussen, Thorfinn Sand, Anders Albrechtsen, and Rasmus Nielsen. 2014. "ANGSD: Analysis of Next Generation Sequencing Data." *BMC Bioinformatics* 15 (November): 356.

Li, Heng, Jue Ruan, and Richard Durbin. 2008. "Mapping Short DNA Sequencing Reads and Calling Variants Using Mapping Quality Scores." *Genome Research* 18 (11): 1851–58.

Manceau, V., L. Després, J. Bouvet, and P. Taberlet. 1999. "Systematics of the Genus Capra Inferred from Mitochondrial DNA Sequence Data." *Molecular Phylogenetics and Evolution* 13: 504–10.

Mashkour, Marjan, Sarieh Amiri, Homa Fathi, Roya Khazaeli, Karyne Debue, Delphine Decruyenaere, Sanaz Beizaee Doost, Safoora Kamjan, and Eberhard W. Sauer. 2020. "Herding and Hunting in the Highlands from the Sasanian to Late Medieval Periods: The Archaeozoology of the Dariali Gorge." In *Dariali: The "Caspian Gates" in the Caucasus from Antiquity to the Age of the Huns and the Middle Ages: The Joint Georgian-British Dariali Gorge Excavations and Surveys of 2013–2016*, edited by Eberhard W. Sauer, 6:729–80. British Institute of Persian Studies Archaeological Monograph Series. Oxbow.

Masseti, Marco. 2009. "The Wild goatsCapra aegagrusErxleben, 1777 of the Mediterranean Sea and the Eastern Atlantic Ocean Islands." *Mammal Review* 39 (2): 141–57.

Meyer, Matthias, and Martin Kircher. 2010. "Illumina Sequencing Library Preparation for Highly Multiplexed Target Capture and Sequencing." *Cold Spring Harbor Protocols* 2010 (6): db.prot5448.

Meyer, Matthias, Martin Kircher, Marie-Theres Gansauge, Heng Li, Fernando Racimo, Swapan Mallick, Joshua G. Schraiber, et al. 2012. "A High-Coverage Genome Sequence from an Archaic Denisovan Individual." *Science* 338 (6104): 222–26.

Molloy, Erin K., Arun Durvasula, and Sriram Sankararaman. 2021. "Advancing Admixture Graph Estimation via Maximum Likelihood Network Orientation." *Bioinformatics* 37 (Suppl_1): i142–50.

Naderi, Saeid, Hamid-Reza Rezaei, François Pompanon, Michael G. B. Blum, Riccardo Negrini, Hamid-Reza Naghash, Ozge Balkiz, et al. 2008. "The Goat Domestication Process Inferred from Large-Scale Mitochondrial DNA Analysis of Wild and Domestic Individuals." *Proceedings of the National Academy of Sciences of the United States of America* 105: 17659–64.

Parrini, Francesca, James W. Cain, and Paul R. Krausman. 2009. "Capra Ibex (Artiodactyla: Bovidae)." *Mammalian Species*, no. 830 (May): 1–12.

Pfitzenmayer, E. V. 1915. "Nekotoryye Interesnyye Ublyudki Semeistva Polorogikh Iz Zakavkaz'ya [Some Interesting Bovid Mongrels from Transcaucasia]." *Izvestiya Kavkazskogo Muzeya* 8: 251–66.

Pickrell, Joseph K., and Jonathan K. Pritchard. 2012. "Inference of Population Splits and Mixtures from Genome-Wide Allele Frequency Data." *PLoS Genetics* 8 (11): e1002967.

Pidancier, Nathalie, Steve Jordan, Gordon Luikart, and Pierre Taberlet. 2006. "Evolutionary
History of the Genus Capra (Mammalia, Artiodactyla): Discordance between Mitochondrial
DNA and Y-Chromosome Phylogenies." *Molecular Phylogenetics and Evolution* 40 (3):
739–49.
Ramsey, Christopher Bronk. 2009. "Bayesian Analysis of Radiocarbon Dates." *Radiocarbon* 51
(1): 337–60.
Reimer, Paula J., William E. N. Austin, Edouard Bard, Alex Bayliss, Paul G. Blackwell,
Christopher Bronk Ramsey, Martin Butzin, et al. 2020. "THE INTCAL20 NORTHERN
HEMISPHERE RADIOCARBON AGE CALIBRATION CURVE (0–55 CAL kBP)."
Radiocarbon, August, 1–33.
Rohland, Nadin, Eadaoin Harney, Swapan Mallick, Susanne Nordenfelt, and David Reich. 2015.
"Partial Uracil-DNA-Glycosylase Treatment for Screening of Ancient DNA." *Philosophical
Transactions of the Royal Society of London. Series B, Biological Sciences* 370 (1660):
20130624.
Ropiquet, Anne, and Alexandre Hassanin. 2006. "Hybrid Origin of the Pliocene Ancestor of Wild
Goats." *Molecular Phylogenetics and Evolution* 41 (2): 395–404.
Sarkisov, A. A. 1953. "0 Pomesyakh Polorogikh [On Bovid Crossbreeds]." *Priroda* 42: 113.
Shackleton, David M. 1997. "Wild Sheep and Goats and Their Relatives: Status Survey and
Conservation Action Plan for Caprinae." IUCN.
https://portals.iucn.org/library/sites/library/files/documents/1997-006.pdf.
Uerpmann, Hans-Peter. 1987. *The Ancient Distribution of Ungulate Mammals in the Middle
East: Fauna and Archaeological Sites in Southwest Asia and Northeast Africa.* Vol. Nr. 27.
Beihefte Zum Tübinger Atlas Des Vorderen Orients. Wiesbaden: L. Reichert Verlag.
Weinberg, Paul J. 2002. "Capra Cylindricornis." *Mammalian Species*, July, 1–9.
Weinberg, Paul, and Bejan Lortkipanidze. 2020. "IUCN Red List of Threatened Species: Capra
Cylindricornis." *IUCN Red List of Threatened Species.*
https://www.iucnredlist.org/species/3795/91287260.
Wilson, Don E., and Deeann M. Reeder. 2005. *Mammal Species of the World: A Taxonomic
and Geographic Reference.* JHU Press.
Yang, D. Y., B. Eng, J. S. Waye, J. C. Dudar, and S. R. Saunders. 1998. "Technical Note:
Improved DNA Extraction from Ancient Bones Using Silica-Based Spin Columns."
American Journal of Physical Anthropology 105 (4): 539–43.
Zheng, Zhuqing, Xihong Wang, Ming Li, Yunjia Li, Zhirui Yang, Xiaolong Wang, Xiangyu Pan, et
al. 2020. "The Origin of Domestication Genes in Goats." *Science Advances* 6 (21):
eaaz5216.

400