Multilevel selection analysis reveals diversity in selfish mitochondrial genome dynamics

Bryan L. Gitschlag[1], Claudia V. Pereira[1], James P. Held[1], & Maulik R. Patel[1,2,3] *

[1]Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA

[2]Department of Cell and Developmental Biology, Vanderbilt University School of Medicine,

Nashville, TN, USA

[3]Diabetes Research and Training Center, Vanderbilt University School of Medicine, Nashville,

TN, USA

Corresponding author: maulik.r.patel@vanderbilt.edu

Abstract

Mutant copies of mitochondrial DNA (mtDNA) can behave as selfish entities, propagating within

organisms at the expense of host fitness. We previously developed experiments to separately

measure the within-host proliferation, and host fitness cost, of selfish mtDNA. By focusing on a

single mutant genome, the *uaDf5* variant, our previous work raises the question of

generalizability: do selfish genomes uniformly overcome a heavy fitness cost by proliferating to

high frequency, or do their population dynamics vary? By applying a standardized multilevel

selection analysis across a collection of mitochondrial mutants, we uncover variation in cheating

strategies. Mutants that impose a heavy fitness cost nevertheless persist by outcompeting

wildtype mtDNA within hosts. Conversely, some mutants fail to outcompete wildtype mtDNA but

nevertheless persist, maintaining a frequency range at which host fitness cost is negligible.

These findings suggest that the population dynamics of mutant mtDNA depend on the loci

affected.

Introduction

Cooperation and cheating represent conflicting adaptive strategies understood to shape the evolution and behavior of complex biological systems (Fisher and Regenberg, 2019, Gulli et al., 2019, Michod et al., 2006, West et al., 2015, Hammerschmidt et al., 2014). In particular, transitions toward larger, more complex forms of life occur when natural selection favors cooperation and loss of conflict between individual life forms, enabling them to become integrated into a collective entity that replicates and undergoes selection as a unified whole (Michod et al., 2006, West et al., 2015, Queller and Strassmann, 2009). However, because cooperators incur the near-term cost of contributing to the fitness of partners, albeit for mutual long-term benefit, cooperation creates conditions that can select for the emergence of selfish, "cheater" entities. Cheating can be defined as occurring when a cooperative partner provides a benefit to the fitness of another entity (the "cheater"), at the expense of its own fitness (Ghoul et al., 2014). Cheaters reap the benefit of belonging to a cooperative relationship without incurring the cost of reciprocating. Examples of cheating can be found across the various levels of scale in the hierarchy of biological organization, including the level of genes (Bravo Nunez et al., 2018, Hammond et al., 2012, Hu et al., 2017, Larracuente and Presgraves, 2012, Schimenti, 2000), cells (Aktipis et al., 2015), and social animals (Riehl and Frederickson, 2016). By exploiting the costly contributions of others without reciprocating, cheaters maintain an advantage within a group. Competition between groups, however, maintains a selection pressure that can select for cooperation and against cheating (Aktipis et al., 2015, de Vargas Roditi et al., 2013, Fiegna and Velicer, 2003, Moreno-Fenoll et al., 2017, Rainey and Rainey, 2003, Wenseleers and Ratnieks, 2004). The combination of conflicting selection pressures across these different levels of scale, termed conflicting multilevel selection, thus resolve an apparent paradox, namely the coexistence between selfish and cooperative entities within a hierarchically structured population.

The eukaryotic cell was first recognized as the result of a cooperative symbiosis more than fifty

years ago (Sagan, 1967). Moreover, mitochondria were hypothesized to experience conflicting

evolutionary pressures (Leigh, 1983), even before pathogenic mitochondrial mutations were first

identified (Holt et al., 1988). Specifically, natural selection was proposed to occur within

individual host organisms in a manner that favors selfish genetic elements, while selection for

host fitness was proposed as a form of "group selection," favoring the faithful transmission of a

metabolically competent population of mitochondrial genomes (Leigh, 1983). Consistent with

this notion, multiple studies in recent years have identified mitochondrial mutations across a

wide diversity of life that propagate while inflicting a fitness cost on their hosts (Havird et al.,

2019, Klucnika and Ma, 2019). These observations have inspired a widespread interest among

mitochondrial geneticists and microbial ecologists in the phenomenon of multilevel selection

(Clark et al., 2012, Dubie et al., 2020, Gitschlag et al., 2020, Havird et al., 2019, Klucnika and

Ma, 2019, Shou, 2015).

Consistent with the predictions of conflicting multilevel selection, a deleteriously mutated

mitochondrial DNA (mtDNA) variant—the *uaDf5* variant in *Caenorhabditis elegans*—has been

previously reported to stably persist in a heteroplasmic state at approximately 60 percent

frequency across hundreds of generations (Tsang and Lemire, 2002), despite being deleterious

at the level of host fitness (Liau et al., 2007). More recent follow-up studies have used the *uaDf5*

heteroplasmy as a model to identify mechanisms underlying the selfish propagation of

deleterious mtDNA (Gitschlag et al., 2016, Gitschlag et al., 2020, Lin et al., 2016, Yang et al.,

2022). To what extent can the results of such work be generalized? Mutant mitochondrial

genomes that propagate selfishly at the expense of host fitness have been observed to arise

spontaneously in laboratory populations of yeast (Bernardi, 2005) and *C. elegans* (Dubie et al.,

2020, Konrad et al., 2017), suggesting that cheating may represent a relatively common theme

in the population dynamics of mitochondrial mutations. On the other hand, while *uaDf5* persists

at approximately 60 percent heteroplasmic frequency, another heteroplasmy in *C. elegans*, a

spontaneously-arising mutant mtDNA variant labeled the Δ*ctb-1* mitotype, was found to maintain

upwards of 90 percent heteroplasmy frequency (Dubie et al., 2020). These findings suggest that

mitochondrial mutations differ from one another in their heteroplasmy dynamics, consistent with

the observed diversity of phenotypic manifestations among mtDNA variants (Taylor and

Turnbull, 2005). Given that selfish mtDNA proliferation has been observed to arise as a result of

interactions between the mitochondrial and host genomes (Gitschlag et al., 2016, Gitschlag et

al., 2020, Lin et al., 2016, Yang et al., 2022), variation among mtDNA variants in average

frequency and in phenotypic effects raise the possibility that the propensity for cheating

behavior likely differs between deleterious mtDNA variants.

To address this possibility, we sought to develop a standardized method for quantifying the

cheating behavior of different mutant mtDNA (ΔmtDNA) variants. We recently developed an

experimental strategy to isolate and measure the strength of natural selection at each

respective level (Gitschlag et al., 2020), namely within hosts (sub-organismal selection) and at

the level of overall host fitness (organismal selection). Accordingly, we were able to identify

effects of nutrient status on both the selfish proliferation and the host fitness effects of *uaDf5*

(Gitschlag et al., 2020). Building on this prior work, here we describe an approach for integrating

empirical measurement with theory, in order to model the population dynamics of selfish

mitochondrial genomes. Using this approach, we sought to expand the scope of our

investigation to include other mtDNA variants, representing mutations in enzyme complexes I,

III, and IV of the electron transport chain (ETC). By characterizing their within-host and

between-host population dynamics, we find that although cheating is evidently a common

feature among deleterious mitochondrial mutations, the specific nature of the cheating "strategy"

varies considerably among different mtDNA variants.

Results

An experimental strategy to isolate selection on a selfish mitochondrial genome at different levels

We sought to identify broad themes and notable differences between different stably propagating heteroplasmic ΔmtDNA variants. We elected to use the mutant mtDNA variants *mpt1*, *mpt2*, *mptDf3*, *mptDf2,* and *mpt4* (Figure 1A). These encompass mutations in ETC complex I (*mpt1* and *mpt2*), complex III (*mptDf3*), and complex IV (*mpt4* and *mptDf2*). Mutations in *mpt1*, *mpt2*, and *mpt4* are each confined to a single protein-coding gene. The *mptDf2* deletion also deletes transfer RNA (tRNA) genes and the mutation in *mptDf3* deletes the five most 3' nucleotides of the tRNA gene for isoleucine suggesting that these mutations may also disrupt ETC function more broadly by interfering with mitochondrial protein synthesis. Importantly, these mutant variants persist at substantially different heteroplasmic frequencies (Figure 1B), suggesting that they differ with respect to within-host proliferation, host-level selection, or both. We therefore sought to develop a standardized approach for quantifying differences in cheating behavior among ΔmtDNA variants; that is, differences in their within-host proliferation and host fitness consequences.

We previously measured the impact of sub-organismal selection on the propagation of the ΔmtDNA variant *uaDf5* (Figure 1A, red bar) by quantifying *uaDf5* frequency longitudinally between individual parents and their own respective adult progeny (Figure 2A) (Gitschlag et al., 2020). These data revealed negative frequency-dependent selection, in which *uaDf5* outcompetes wildtype mtDNA for replication across organismal development, but its ability to do

so declines as a function of increasing starting (parental) *uaDf5* frequency. Selection against

uaDf5 that occurs strictly at the level of host fitness was measured by competing *uaDf5*-carrying

heteroplasmic animals against their homoplasmic (wildtype) counterparts on the same food

plate (Figure 2B) (Gitschlag et al., 2020). Non-competing control lines were maintained, which

consisted of exclusively heteroplasmic animals, with no wildtype animals to compete against. To

control for variation in *uaDf5* frequency due to reasons other than the organismal fitness cost of

uaDf5, the *uaDf5* frequencies of competing lines were normalized to that of non-competing lines

at each generation; any resulting change in the *uaDf5* frequency of the competing lines relative

to the non-competing lines can therefore be attributed to an organismal fitness effect of *uaDf5*,

since the presence of animals lacking *uaDf5* represents the only distinguishing condition

between competing and non-competing lines. Accordingly, we have developed an approach to

separately measure the effects of selection on ∆mtDNA at the sub-organismal and organismal

levels.

Multilevel selection analysis of the *uaDf5* heteroplasmy

Here, we sought to integrate the measurements of sub-organismal and organismal selection

into a single theoretical framework describing the overall population dynamics of ∆mtDNA. The

Price Equation, which provides such a framework, describes evolution as the covariance

between a trait and its reproductive fitness (Frank, 1997, Price, 1972). Importantly, since this

equation can accommodate more than one source of covariance, including non-linear

interactions (Frank, 2012), it can be used to describe selection at different levels

simultaneously:

$$w\Delta z = cov(w,z)_{org} + E[cov(w,z)_{sub}]$$

In this form of the equation, *z* refers to a heritable trait that can be quantitatively measured.

Since mitochondrial genotype consists of a heteroplasmic mix of mtDNA variants, trait-value *z*

can be defined in this case as *uaDf5* frequency. Fitness *w* refers to the net replicative success

of *uaDf5* relative to wildtype mtDNA, defined as the ratio of new *uaDf5* copies to new wildtype

mtDNA copies per generation. Subscripts $_{org}$ and $_{sub}$ label the covariance between *w* and *z* at the

organismal and sub-organismal levels, respectively, with covariance defined by the following

formula:

$$cov(w,z) = \frac{\sum(w_i - w_{average})(z_i - z_{average})}{n-1}$$

The term *E* in front of the sub-organismal covariance term denotes expected, or average, sub-

organismal covariance between *w* and *z*; note that the total sub-organismal change in *uaDf5*

frequency is the average number of replicated copies per *uaDf5* template compared to that of

wildtype mtDNA. Moreover, since the strength or intensity of selection can be expressed as the

selection coefficient *S* (equal to 1 – *w*), the change in *uaDf5* frequency at each respective level

of selection can be expressed using the covariance term that relates selection coefficient to

uaDf5 frequency (see Figure 2C for a sample illustration):

$$cov(S, \%\Delta mtDNA) = \frac{\sum(S_i - S_{average})(\%uaDf5_i - \%uaDf5_{average})}{n-1}$$

167   Although the above formula for covariance merely describes how variation in *uaDf5* frequency

168   relates to variation in selection acting on *uaDf5*, this relation can be applied to the empirically

169   measured shifts in *uaDf5* frequency reported previously (Gitschlag et al., 2020) to derive a

170   function that relates selection to *uaDf5* frequency, taking both levels of selection into account

171   (for a more detailed derivation of this equation, see Figure 3—Figure Supplement 1):

$$S_{overall} = S_{sub-organismal} + S_{organismal} = (0.01154 * \%uaDf5 - 0.923) + (10^{0.0181 * \%uaDf5 - 1.81})$$

The term shown in the parentheses on the left defines the relationship between *uaDf5*

frequency and sub-organismal selection (plotted in Figure 3A). The term shown in the

parentheses on the right defines the relationship between *uaDf5* frequency and organismal

selection (plotted in Figure 3B, solid red line), relying on two additional assumptions. The first

assumption is that selection coefficient would be zero when *uaDf5* frequency is 0 percent, since

there is no *uaDf5* to select against. Second, the selection coefficient is assumed to reach 1—

fitness reaches 0, reflecting elimination of *uaDf5* from the population within a single

generation—as *uaDf5* frequency reaches 100 percent. This assumption rests on the fact that

the mutation deletes essential genes and should therefore be lethal if present at 100 percent

frequency. Summing the organismal and sub-organismal regressions yields the overall,

population-wide relationship between heteroplasmic *uaDf5* frequency and the strength of

selection acting on *uaDf5* (Figure 3C).

The Price Equation framework shows that selection at the sub-organismal and organismal

levels balance—the overall selection coefficient approaches 0, reflecting no net difference in

fitness between *uaDf5* and wildtype mtDNA—when *uaDf5* is near 60 percent heteroplasmic

frequency (Figure 3C). Note that this is the approximate frequency at which *uaDf5* persists in

the non-competing populations, in which the conflicting levels of selection are both in effect

(Gitschlag et al., 2020). Importantly, this conclusion is compatible with a range of alternate

assumptions regarding the impact of rising *uaDf5* frequency on organismal fitness (Figure 3B-

C). The mutant genome may become lethal before reaching fixation, at 90 or 95 percent

frequency for example. Consistent with this idea, *uaDf5* is rarely observed at or above 90

percent frequency (Gitschlag et al., 2016, Gitschlag et al., 2020, Lin et al., 2016, Tsang and

Lemire, 2002). Alternatively, the fitness cost of *uaDf5* might not rise smoothly, as depicted in the

non-linear regression in Figure 3B. For example, the organismal selection coefficient may rise

linearly with increasing *uaDf5* frequency up to and beyond the mean of 60 percent

heteroplasmic frequency, perhaps until reaching a critical threshold before imposing a heavy

fitness cost, resulting in an abrupt upward shift in the strength of selection. However, taking

these alternative assumptions into account had little effect on the overall conclusions (Figure

3B-C). In conclusion, we have integrated measurements of selection on *uaDf5* at different levels

into a unified theoretical framework that robustly predicts the overall population-level dynamics

observed for *uaDf5*.

Quantification of tracking sub-organismal frequency and organismal selection amongst different heteroplasmies

By measuring sub-organismal and organismal selection using the approach initially developed for the study of *uaDf5*, we were able to isolate changes in mutant mtDNA frequency in a manner specific to each level of selection (Figure 4 and Figure 4—Figure Supplements 1-2). Since each mtDNA variant stably persists in a heteroplasmic population, despite harboring a deleterious mutation in at least one essential gene, they could all be predicted to undergo conflicting multilevel selection similarly to *uaDf5* (Figure 3). Surprisingly, however, the capacity for within-host proliferation, as well as the host-level fitness costs, varies considerably between mutant mitochondrial genomes.

Neither of the ETC complex I deletions show a significant effect on host fitness (Figure 4B and 4D). It should be noted, however, that the two complex I mutations stably persist at different frequencies (Figures 1B, 4A, 4C, and Figure 4—Figure Supplement 2A-B). Consistent with a lack of host fitness effects, these mutations appear to undergo neither within-host proliferation nor negative organismal selection, remaining at similar frequencies across generations in both sub-organismal and organismal experiments (Figure 4A-D). The same is true for the complex III mutation in *mptDf3* (Figure 4E-F). Interestingly, the population-wide frequency of *mptDf3* rose dramatically, albeit in only one of the competition lines (Figure 4F), consistent with random drift toward a higher population-wide fraction of heteroplasmic animals. The complex IV mutations, on the other hand, inflict dramatic fitness effects, being almost entirely outcompeted by homoplasmic (wildtype) animals within six generations. This was true for *mptDf2* (Figure 4H), which affects both complex IV and tRNA genes, as well as *mpt4* (Figure 4J), which only affects the complex IV gene *CO1*. Consistent with these findings, mutations in human *CO1* are associated with several pathologies (Brown et al., 1992, Namslauer and Brzezinski, 2009,

Varlamov et al., 2002). These severe host fitness costs predict that for these mutations to

persist in a population, they must undergo significant within-host proliferation, sufficient to

overcome the strong negative selection at the organismal level. Indeed, this prediction is

remarkably consistent with our observation (Figure 4G and 4I). We therefore have identified

mutations that behave similarly to *uaDf5*, and mutations that differ, from a collection of five

additional heteroplasmic mtDNA variants.

Multilevel selection analysis of a panel of different heteroplasmies

Finally, we sought to integrate these findings using the same theoretical framework developed

for *uaDf5* (Figure 3), by summing the regressions between mutant mtDNA frequency and the

strength of selection at the sub-organismal and organismal levels (Figure 5). Not surprisingly,

when the levels of selection are combined into a single overarching description of their

respective population dynamics, there is no overall (population-level) covariance between

mutant frequency and selection, at least for *mpt1*, *mpt2*, and *mptDf3* (Figure 5A-I). In other

words, the frequency of these mutations does not seem to correlate with their fitness effects, at

least within the frequency ranges at which these heteroplasmies were observed to persist. In

contrast, the regressions for the mutant genome *mptDf2* (Figure 5J-L) predicts that this genome

stably persists at very high frequency—near 80 percent—due to a balance between a strong

sub-organismal proliferative advantage and a strong host fitness cost. As with *uaDf5*, this

prediction is compatible with alternate assumptions regarding the relationship between

heteroplasmy frequency and host fitness cost (Figure 5—Figure Supplement 1); moreover,

these findings are highly consistent with the empirical observation that *mptDf2* maintains very

high frequency in non-competing lines, where the forces of sub-organismal and organismal

selection are both in effect (Figure 1B and Figure 4—Figure Supplement 2D). Although *mpt4*

undergoes robust sub-organismal proliferation (Figure 4I and Figure 4—Figure Supplement 4E),

the magnitude of proliferation does not appear to co-vary with *mpt4* frequency (Figure 5M),

indicating that the combined population-level covariance fails to predict the level at which this

mutant genome persists (Figure 5O). However, the substantial host fitness cost of this genome

(Figures 4J and 5N) implies a strong selection pressure that acts in opposition to the sub-

organismal proliferation, similarly to *uaDf5* and *mptDf2*. In conclusion, we have identified

notable similarities and differences amongst the population dynamics of mitochondrial

mutations.

Discussion

Multilevel selection theory provides a powerful explanatory account of cooperator-cheater

dynamics. We developed an experimental approach to empirically quantify selection for selfish

mitochondrial genomes at the sub-organismal (within-host) level by tracking their frequency

longitudinally within isolated parent-progeny lineages. At the organismal (between-host) level,

we measure selection against selfish mitochondrial genomes by competing heteroplasmic and

wildtype animals on the same food plate and tracking the population-wide frequency of mutant

mtDNA. By enabling the empirical measurement of selection at different levels, this approach

can be applied to explore cooperator-cheater dynamics broadly. Integrating these empirical

measurements into a theoretical model (Figure 3) reveals a more complete, higher-resolution

depiction of the overall population dynamics of selfish mtDNA. Apart from confirming that the

mutant persists within a given frequency range due to balancing selection among the different

levels, measuring selection at each level makes it possible to account for their respective

contributions toward experimentally induced shifts in heteroplasmy level, providing a framework

for detailed mechanistic follow-up.

The mutant mtDNA variant *uaDf5* is an excellent model for the study of selfish mitochondrial

genome dynamics, for a number of reasons. For example, this mutant rises in frequency over

time within hosts (Figure 3A) at the expense of inflicting a fitness cost (Figure 3B). The

proliferation of *uaDf5* is also frequency-dependent, a common feature of cheater entities

(Dobata and Tsuji, 2013, Dugatkin et al., 2005, Pruitt and Riechert, 2009, Riehl and

Frederickson, 2016, Ross-Gillespie et al., 2007). Moreover, *uaDf5* proliferates by exploiting

interactions with the host genome that exist to minimize the harmful effects of metabolic and

nutrient stress (Gitschlag et al., 2016, Gitschlag et al., 2020, Lin et al., 2016, Yang et al., 2022)*,*

or which reason it can be viewed as a bona fide cheater entity. However, mitochondrial

mutations are not all alike with respect to their fitness effects. How broadly generalizable, then,

are the findings associated with the study of *uaDf5*? We sought to address this question here by

identifying noteworthy similarities and differences in the population dynamics of different

mitochondrial DNA mutations/deletions.

Variation in selfish mitochondrial genome dynamics

By quantitatively measuring the strength of selection acting on mutant mtDNA at the sub-

organismal and organismal levels, across a collection of mitochondrial mutations, we identify

two noteworthy themes. On one hand, like *uaDf5*, some mutant mtDNA variants propagate in

spite of inflicting a heavy fitness burden on their hosts. As expected, their seemingly paradoxical

persistence—in other words, their ability to avoid extinction in the face of strong negative

organismal selection—appears to coincide with strong positive sub-organismal selection (Figure

6, left side). These conflicting multilevel selection dynamics can be observed in three of the six

heteroplasmies featured here: *uaDf5*, *mptDf2*, and *mpt4*. Although the molecular details

underlying these dynamics are not clear, the only thing that all three mutations have in common

is a disruption to ETC complex IV. Consistent with these findings, mutations in the complex IV

gene *CO1* are associated with a number of human clinical pathologies (Brown et al., 1992,

Namslauer and Brzezinski, 2009, Varlamov et al., 2002). In any case, selfish propagation in

which mutant genomes survive negative organismal selection by outcompeting the wildtype

mtDNA within individual hosts, appears to be a relatively common, albeit not universal, theme

among stably persisting deleterious heteroplasmies.

In addition to these conflicting multilevel selection dynamics, a second general theme emerges

from tracking the levels of some of the mutant genomes featured here. In contrast to *uaDf5*,

mptDf2, and *mpt4*, not all mutant mitochondrial genomes persist by rising in frequency across

development at the expense of a strong host fitness cost. On the contrary, despite carrying

deletions in essential metabolic genes, some mutant genomes appear to undergo neutral drift—

that is, fluctuations in frequency of existing alleles not attributable to natural selection—even

when the heteroplasmic animals compete for several generations against their wildtype

counterparts on the same food. This is true for both mutations in ETC complex I—the *mpt1*

deletion in *ND1* and the *mpt2* deletion in *ND5*—as well as for *mptDf3*, which affects the complex

III gene for cytochrome b (Figure 6, right side). Taken together, these observations suggest that

mtDNA variants can persist in a manner consistent with neutral drift, despite carrying mutations

that deleteriously affect essential ETC components.

Comparison with previous findings

Interestingly, *mptDf3*, which carries a deletion affecting the complex III mtDNA-encoded protein,

namely cytochrome b, does not appear to inflict a significant host fitness cost, at least not within

the frequency range at which *mptDf3* is observed to persist. This may be difficult to reconcile

with previous work showing that an mtDNA variant labeled Δ*ctb-1*—owing to a cytochrome b

mutation—behaves much more like *uaDf5*, *mptDf2*, and *mpt4*, in particular by proliferating within

hosts while imposing a host fitness cost (Dubie et al., 2020). In fact, in contrast to *uaDf5*, Δ*ctb-1*

was reported to persist at upwards of 90 percent heteroplasmic frequency for dozens of

generations (Dubie et al., 2020). One possible explanation for this apparent discrepancy

between the dynamics of Δ*ctb-1* and *mptDf3* is that Δ*ctb-1* contains other mutations in addition

to the deletion in cytochrome b, which may contribute to its observed selfish proliferation, or

host fitness consequences, or both (Dubie et al., 2020).

Hypothesis and future directions

The sub-organismal selection dynamics described here are remarkably consistent with recent

mammalian work. In particular, a tRNA point mutation in mouse mtDNA, and three point

mutations in human mtDNA, were recently found to undergo an upward shift in frequency from

parents to offspring, especially among parents with lower starting mutant levels (Zhang et al.,

2021, compare Figure 1C and Figure 8B, D and F to Figure 2B in (Gitschlag et al., 2020). These

shifts occur in spite of deleterious phenotypic effects associated with this heteroplasmy (Zhang

et al., 2021), suggesting a cheating behavior strikingly similar to mtDNA variants *uaDf5*, *mptDf2*,

and *mpt4* in *C. elegans*. Moreover, we previously applied the experiments described in this

study to identify conditions and host genotypes that modulate both the sub-organismal

proliferation and host fitness cost of *uaDf5* (Gitschlag et al., 2020). Taken together, in addition to

an apparent wide prevalence of cheating behavior among animal mitochondrial genomes, these

findings further suggest an overlap between the mechanisms underlying cheating and those

underlying disease. By applying the multilevel selection analysis described here to a diversity of

conditions and host genotypes, we anticipate that it will be possible to identify avenues for

mitigating the risk and severity of mitochondrial disease.

How can a mitochondrial genome lacking an essential gene manage to stably persist with no

measurable fitness effect at either level of selection (organismal or sub-organismal)? Although

the explanation is not known, these observations raise the interesting possibility that some

mutant genomes propagate, not in spite of a deleterious host fitness cost, but rather because of

what could be considered a "flying under the radar" strategy. Some mutant genomes seem to

successfully persist by maintaining a frequency range that enables them to avoid exposure to

negative organismal selection. One possible basis for this phenomenon would invoke a balance

between positive and negative selection at the sub-organismal level. For example, these mutant

genomes might affect mitochondrial function in a manner that elicits a combination of cellular

mechanisms that simultaneously favor and disfavor the mutant genome within individual hosts.

If this balance is achieved within a frequency range that is too low to significantly impact host

fitness, the mutant mtDNA could undergo neutral drift at the organismal level while maintaining

a steady-state heteroplasmy.

This possibility could be explored in future research by testing host genes for their effect on

heteroplasmy dynamics within hosts. For example, previous work has shown that mtDNA

replication can be biased in favor of either a mutant or wildtype genome depending on the

expression level of the mtDNA polymerase, POLG (Chiang et al., 2019, Lin et al., 2016). This

raises the possibility that physiological stress-response mechanisms might bias replication in

favor of the mutant genome by influencing the expression or localization of mtDNA replication

machinery. For example, the gene *daf-16* encodes a Forkhead box O (FoxO) transcription factor

that functions as a master regulator of metabolic homeostasis. Although *daf-16* does not appear

to directly regulate POLG expression (Tepper et al., 2013), it nevertheless regulates other

aspects of mitochondrial function, including the expression of proteins involved in energy

metabolism and stress response (Depuydt et al., 2014, Tepper et al., 2013, Webb et al., 2016).

If its attempt to rescue metabolic integrity influences mtDNA replication, even indirectly, then a

compensatory effect could result that favors underperforming regions of the mitochondrial

network, inadvertently promoting replication of mutant genomes. Consistent with this possibility,

daf-16 mediates rising *uaDf5* frequency across organismal development (Gitschlag et al., 2020).

Like *daf-16*, the gene *atfs-1* also regulates a transcriptional response to metabolic stress, in this

case by activating the mitochondrial unfolded protein response (UPR^{mt}) (Nargund et al., 2012).

Importantly, *atfs-1* gain-of-function mutations are associated with elevations in both *uaDf5*

frequency (Gitschlag et al., 2016, Lin et al., 2016) and expression of *polg-1*, the gene encoding

POLG (Lin et al., 2016). Together, these considerations highlight *daf-16* and *atfs-1* as attractive

candidate genes underlying positive sub-organismal selection for the mutant mitochondrial

genomes featured in this study. Conversely, *drp-1* is involved in mitochondrial fission, while

pink-1 and *pdr-1* play central roles in PINK1/Parkin-dependent mitophagy, and both of these

processes have been implicated in selection against mutant mtDNA at the sub-organismal level

(Gitschlag et al., 2016, Kandul et al., 2016, Lieber et al., 2019, Lin et al., 2016, Suen et al.,

2010, Valenci et al., 2015). Does the combined activity, and hence influence, of each of these

pathways vary amongst different selfish mitochondrial genomes? This question could be

addressed by future research, for example by comparing UPR^{mt} activation and levels of

mitophagy, across different heteroplasmies. It would also be interesting to determine how

differently, stably-propagating heteroplasmies compare in their sensitivity to the disruption of

these pathways.

Materials and Methods

Nematode husbandry

C. elegans strains used in this study were maintained on 60-mm standard nematode growth

medium (NGM) plates (for measuring sub-organismal selection), or 100-mm NGM plates (for

measuring organismal selection), seeded with live OP50-strain *E. coli* bacteria as a food source.

Nematode strains were incubated at 20°C. In addition to the Bristol wildtype strain and the

heteroplasmic *uaDf5* strain featured in prior work (Gitschlag et al., 2020), five additional *C.*

elegans strains were used in this study. These consisted of heteroplasmic mutant genomes

mpt1, *mpt2*, *mptDf3*, *mptDf2*, and *mpt4*, each crossed into the nuclear background of the Bristol

strain.

Nuclear genome exchange in heteroplasmic strains.

To ensure that our analysis was not confounded by variation in the nuclear genome, we

employed a novel unigametic inheritance method to completely exchange the nuclear genome

of each heteroplasmic strain with the nuclear genome of wildtype (Bristol strain) *C. elegans*

(method from Artiles *et al.* 2019). Thus, all heteroplasmic strains used in this study have

identical nuclear backgrounds. Briefly, each heteroplasmic strain was crossed to the *gpr-1* over-

expression strain PD2220 following Mendelian genetics. Next, hermaphrodites of the stable *gpr-*

1 over-expression heteroplasmy strains were crossed to wildtype males. Non-Mendelian

hermaphrodite progeny from these crosses in which the paternal nuclear background is

unigametically inherited in the germline cell lineage (determined by pharyngeal mosaic

patterning) were individually propagated. Stock strains were established from the progeny of

these animals as they have a complete wildtype nuclear genomic background and retain the

given heteroplasmy.

Lysate preparation

To prepare animals for quantification of mutant mtDNA frequency, nematodes were transferred

to sterile PCR tubes or 96-well PCR plates containing lysis buffer with 100 µg/mL proteinase K.

Lysis buffer consisted of 50 mM KCl, 10 mM Tris pH 8.3, 2.5 mM $MgCl_2$, 0.45% Tween 20,

0.45% NP-40 (IGEPAL), and 0.01% gelatin, in deionized water. Volume of lysis buffer varied by

worm count: 10 µL for individual adults of the parent generation and 20 µL for pooled adult

progeny for measuring sub-organismal selection, and 50 µL for pooled animal lysates from the

competition experiments for measuring organismal selection. After transferring worms to lysis

buffer, each tube or plate was immediately sealed and incubated at -80˚C for 10 minutes to

rupture nematode cuticles, followed by lysis incubation at 60˚C for 60 minutes (90 minutes for

pooled nematodes), and then at 95˚C for 15 minutes to inactivate the proteinase K. Nematode

lysates were then kept at -20˚C for stable long-term storage until being used for genotyping and

quantification.

Quantifying mutant mtDNA levels

Levels of each mutant mtDNA variant were quantified as described previously for *uaDf5*

(Gitschlag et al., 2020), using droplet digital PCR (ddPCR). Nematodes were lysed as described

above, then diluted in nuclease-free water, with a dilution factor varying depending on nematode

concentration: 200x for single adults, 1000x for pooled adults (sub-organismal selection

experiment), and 20,000x for pooled nematodes of mixed age (organismal selection

experiment). Next, 5 µL of each dilute nematode lysate was combined with 0.25 µL of a 10-µM

dilution of each primer needed for amplifying each mtDNA variant. The following primers were

used for ddPCR amplification of each mutant variant as necessary.

mpt1:

Forward primer 1: 5'-TTCTGAAGGTGAAAGGGAGTTAG-3'

Reverse primer 1: 5'-ACTGACCTTAATGGTAAAGTATTTG-3'

Forward primer 2: 5'-GGGTACTTTCTTCTGCTATAT-3'

Reverse primer 2: 5'-CGATAACGAGGGTATGAACTACG-3'

mpt2:

Forward primer 1: 5'-GAAGAAGGTGGTAGCCTTGAGGAC-3'

Reverse primer 1: 5'-CGTATAAGAAAAGTCTTGGGATGTTAAG-3'

Forward primer 2: 5'-GGATTAATTTTCTCAAGGGGTGCTG-3'

Reverse primer 2: 5'-CTTTTTCAAAGACGAAAACTGTAACC-3'

mptDf3:

Forward primer 1: 5'-CCCTGAAGAGGCTAAGAATATTAGG-3'

Reverse primer 1: 5'-GGCAATGTCACCAACATCC-3'

Reverse primer 2: 5'-CCCAATACAATAACTAGAATAGCTCACG-3'

mptDf2:

Forward primer 1: 5'-GGATTGGCAGTTTGATTAGAGAG-3'

Reverse primer 1: 5'-AAGTAACAAACACTAAAACTCCCAAC-3'

Forward primer 2: 5'-CGTGCTTATTTTTCGGCTGC-3'

Reverse primer 2: 5'-CTTTAACACCTGTTGGCACTG-3'

mpt4:

Forward primer 1: 5'-CGGTGGTTTTGGTAACTG-3'

Reverse primer 1: 5'-TCATAGTGTAACACCCGTGAAAATCC-3'

Forward primer 2: 5'-TGATCCAAGAACTGGAGGTAATC-3'

Reverse primer 2: 5'-CCTGTTGGCACTGCAATAAC-3'

Mixtures of dilute nematode lysate and primer were combined with 12.5 μL of Bio-Rad QX200^{TM}

ddPCR EvaGreen Supermix and nuclease-free water to a volume of 25 μL in Eppendorf^{TM} 96-

well twin.tec^{TM} PCR plates. Droplet generation and PCR amplification were performed according

to manufacturer's protocol. Wildtype and mutant-specific primers were combined in the same

reaction, and each droplet was scored as containing either wildtype or mutant mtDNA using the

2-dimensional (518 nm and 554 nm dual-wavelength) clustering plot option in the Bio-Rad

QuantaSoft^{TM} program.

Sub-organismal selection assay

The strength of sub-organismal (within-host) selection on mutant mtDNA was measured

longitudinally across isolated parent-progeny lineages, as previously described (Gitschlag et al.,

2020). Briefly, multiple L4-stage (late larval) heteroplasmic animals were picked at random

under a dissecting microscope from stock populations carrying each heteroplasmy that had

been crossed in the Bristol strain (wildtype) nuclear background. These larvae were transferred

to fresh NGM plates seeded with OP50 *E. coli* bacteria as a food source and incubated for 2

days at 20˚C. The day-2 adults were then segregated by transfer onto individual plates, one

adult per plate, and incubated for 4 hours at 20˚C to produce embryos that are age-

synchronized to within a four-hour window. Each parent was then individually lysed. After 4 days

of continued incubation at 20˚C, the progeny had progressed from embryos to day-2 adults,

reaching the same age at which their respective parents were lysed. Adult progeny were lysed

at this point to obtain progeny that are age-matched to their parents, to control for age-

dependent differences in mutant mtDNA levels. Progeny from each parent were lysed in pools

of 3 to minimize the confounding effect of random drift. Each parent-progeny lineage was

individually segregated from the rest, to ensure that mutant mtDNA frequency from each

progeny lysate was being compared with that of its own respective parent, thereby minimizing

the confounding effect of competition between lineages (organismal selection). Mutant mtDNA

frequency of parents and progeny was determined for each heteroplasmy using ddPCR as

described above, across multiple replicate parent-progeny lineages for statistical power.

Experimental evolution (organismal selection)

Selection against mutant mtDNA that occurs strictly at the level of host fitness was measured

using an organismal competition experiment similarly to the experiment previously described

(Gitschlag et al., 2020). Briefly, for each mutant mtDNA variant, heteroplasmic nematodes

carrying mutant mtDNA in the Bristol nuclear background were combined with Bristol-strain

nematodes on 10-cm NGM plates seeded with live OP50 *E. coli* bacteria as a food source.

Approximately 500 nematodes were transferred to each plate. In addition to 4 replicate

competition lines for each heteroplasmy, 4 non-competing control lines were established by

transferring only heteroplasmic animals onto their own food plates, with no homoplasmic-

wildtype animals to compete against. Every 3 days, the generation for each experimental line

was reset; nematodes were washed off the plates using M9 buffer into a sterile 1.7 mL

collection tube. Approximately 500 animals of mixed age from each line were transferred to a

fresh food plate. Another 500 nematodes were lysed together in a single pooled lysate. This

experiment was continued for 6 consecutive generations.

Statistical analysis

The effect of initial mutant mtDNA frequency on sub-organismal heteroplasmy dynamics was

observed in three previously reported experiments involving *uaDf5* (Gitschlag et al., 2020), and

one experiment each for the five additional mutant mtDNA variants featured in this study. Each

data point represents a biological replicate, consisting of a parent-progeny pair (sub-organismal

selection) or a population of nematodes confined to the same food plate (organismal selection).

For each experiment, sample sizes (number of biological replicates) are provided in Figure 3—

Source Data (for *uaDf5*) and the Figure 4 figure legend (for the remaining mutant mtDNA

variants). Number of animals per replicate is provided in sub-sections **Sub-organismal**

selection assay and **Experimental evolution (organismal selection)** of the **Materials and**

Methods. Sample sizes were chosen based on prior observation of sample sizes sufficient for

drawing statistical inferences (Gitschlag et al., 2020). Where appropriate, statistical analyses

are indicated in the figure legend (Figures 3-5, and Figure 4—Figure Supplement 1); for more

information on analysis of data, see Figure 3—Source Data 1-2 and Figure 3—Figure

Supplement 1. The Wilcoxon matched-pairs sign-ranks was used for the longitudinal measure of

sub-organismal mutant mtDNA dynamics (Figure 4A, 4C, 4E, 4G, and 4I); this was due to the

pairing of parent and progeny inherent in the experimental design. Moreover, the Wilcoxon test

was chosen because it is non-parametric; that is, it does not require assuming the distribution of

mutant mtDNA frequency. Linear regression analysis was used for comparing sub-organismal

selection versus starting (parental) mutant mtDNA frequency and for tracking the multi-

generational decline in mutant mtDNA frequency due to organismal selection (Figures 3, 5,

Figure 4—Figure Supplement 1, Figure 4—Figure Supplement 2, and Figure 5—Figure

Supplement 1). No statistical testing was performed to identify or remove outliers. For the sub-

organismal selection assay (Figure 4A, 4C, 4E, for which N<24), biological replicates were

omitted from final analysis if the ddPCR amplification failed to generate a clear, unambiguous

measurement of mutant mtDNA levels in the corresponding plate-well.

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Genetic reagent (*Caenorhabditis elegans*)	*him-8*(e1489); ΔmtDNA(*uaDf5*)/+	Caenorhabditis Genetics Center	RRID:WB-STRAIN:WBStrain00024106	LB138
Genetic reagent (*Caenorhabditis elegans*)	ccTi1594 [mex-5p::GFP::gpr-1::smu-1 3'UTR + Cbr-unc-119(+), III: 680195] III. umnIs27 [myo-2::GFP + NeoR, III: 8856215 (intergenic)] III	Artiles et al. 2019		PD2220

Genetic reagent (*Caenorhabditis elegans*)	ΔmtDNA(*uaDf5*)/+ in Bristol nuclear background	Gitschlag et al., 2020		See Materials and Methods: Genetic crosses and genotyping
Genetic reagent (*Caenorhabditis elegans*)	ΔmtDNA(*mpt1*)/+ in Bristol nuclear background	This study		See Materials and Methods: Nuclear genome exchange in heteroplasmic strains
Genetic reagent (*Caenorhabditis elegans*)	ΔmtDNA(*mpt2*)/+ in Bristol nuclear background	This study		See Materials and Methods: Nuclear genome exchange in heteroplasmic strains
Genetic reagent (*Caenorhabditis elegans*)	ΔmtDNA(*mptDf3*)/+ in Bristol nuclear background	This study		See Materials and Methods: Nuclear genome exchange in heteroplasmic strains
Genetic reagent (*Caenorhabditis elegans*)	ΔmtDNA(*mptDf2*)/+ in Bristol nuclear background	This study		See Materials and Methods: Nuclear genome exchange in heteroplasmic strains
Genetic reagent (*Caenorhabditis elegans*)	ΔmtDNA(*mpt4*)/+ in Bristol nuclear background	This study		See Materials and Methods: Nuclear genome exchange in heteroplasmic strains
Genetic reagent (*Escherichia coli*)	OP50 strain	Caenorhabditis Genetics Center	RRID: WB-STRAIN:WBStrain00041971	
Sequence-based reagent	PCR primers used in this study	This study		See Materials and Methods: Quantifying mutant mtDNA levels
Other	Eppendorf™ 96-well twin.tec™ PCR plates	Fisher Scientific	Cat#951020303	
Other	QX200 ddPCR EvaGreen Supermix	Bio-Rad	Cat#1864034	

Other	Automated Droplet Generation Oil for EvaGreen	Bio-Rad	Cat#1864112	
Other	DG32 Automated Droplet Generator Cartridges	Bio-Rad	Cat#1864108	
Other	Droplet Reader Oil for ddPCR	Bio-Rad	Cat#1863004	
Software, algorithm	QuantaSoft™	Bio-Rad	Cat#1864011	
Software, algorithm	Prism 8 for macOS	GraphPad Software, Inc.	RRID:SCR_002798	Version 8.1.2

Acknowledgments

The *uaDf5* heteroplasmic strain was kindly provided by the Caenorhabditis Genetics Center

(CGC). We thank O. Thompson and R. Waterston at University of Washington for identifying the

mpt1, *mpt2*, *mptDf3*, *mptDf2*, and *mpt4* deletions from the Million Mutation Project worm

collection. We thank Patrick Abbot, Ann T. Tate and Allison Hilbun of Vanderbilt University for

valuable guidance and assistance with data analysis. This work was supported by the NIH

research project grants (R01 GM123260 and GM145378), the Discovery Award (PR170792)

from the Department of Defense's Congressionally Directed Medical Research Program, the

Vanderbilt University Medical Center Diabetes Research and Training Center Pilot & Feasibility

Grant, the NIH-sponsored Cellular, Biochemical and Molecular Sciences Training Program

(5T32GM008554-18), and the Ruth L. Kirschstein National Research Service Award Individual

Predoctoral Fellowship grant (1F31GM125344).

579   **References**

AKTIPIS, C. A., BODDY, A. M., JANSEN, G., HIBNER, U., HOCHBERG, M. E., MALEY, C. C. &
WILKINSON, G. S. 2015. Cancer across the tree of life: cooperation and cheating in
multicellularity. *Philos Trans R Soc Lond B Biol Sci,* 370.
BERNARDI, G. 2005. Lessons from a small, dispensable genome: the mitochondrial genome of
yeast. *Gene,* 354**,** 189-200.
BRAVO NUNEZ, M. A., NUCKOLLS, N. L. & ZANDERS, S. E. 2018. Genetic Villains: Killer Meiotic
Drivers. *Trends Genet,* 34**,** 424-433.
BROWN, M. D., YANG, C. C., TROUNCE, I., TORRONI, A., LOTT, M. T. & WALLACE, D. C. 1992. A
mitochondrial DNA variant, identified in Leber hereditary optic neuropathy patients,
which extends the amino acid sequence of cytochrome c oxidase subunit I. *Am J Hum*
Genet, 51**,** 378-85.
CHIANG, A. C., MCCARTNEY, E., O'FARRELL, P. H. & MA, H. 2019. A Genome-wide Screen
Reveals that Reducing Mitochondrial DNA Polymerase Can Promote Elimination of
Deleterious Mitochondrial Mutations. *Curr Biol,* 29**,** 4330-4336 e3.
CLARK, K. A., HOWE, D. K., GAFNER, K., KUSUMA, D., PING, S., ESTES, S. & DENVER, D. R. 2012.
Selfish little circles: transmission bias and evolution of large deletion-bearing
mitochondrial DNA in Caenorhabditis briggsae nematodes. *PLoS One,* 7**,** e41433.
DE VARGAS RODITI, L., BOYLE, K. E. & XAVIER, J. B. 2013. Multilevel selection analysis of a
microbial social trait. *Mol Syst Biol,* 9**,** 684.

DEPUYDT, G., XIE, F., PETYUK, V. A., SMOLDERS, A., BREWER, H. M., CAMP, D. G., 2ND, SMITH,
R. D. & BRAECKMAN, B. P. 2014. LC-MS proteomics analysis of the insulin/IGF-1-
deficient Caenorhabditis elegans daf-2(e1370) mutant reveals extensive restructuring of
intermediary metabolism. *J Proteome Res,* 13**,** 1938-56.
DOBATA, S. & TSUJI, K. 2013. Public goods dilemma in asexual ant societies. *Proc Natl Acad Sci*
U S A, 110**,** 16056-60.
DUBIE, J. J., CARAWAY, A. R., STOUT, M. M., KATJU, V. & BERGTHORSSON, U. 2020. The conflict
within: origin, proliferation and persistence of a spontaneously arising selfish
mitochondrial genome. *Philos Trans R Soc Lond B Biol Sci,* 375**,** 20190174.
DUGATKIN, L. A., PERLIN, M., LUCAS, J. S. & ATLAS, R. 2005. Group-beneficial traits, frequency-
dependent selection and genotypic diversity: an antibiotic resistance paradigm. *Proc Biol*
Sci, 272**,** 79-83.
FIEGNA, F. & VELICER, G. J. 2003. Competitive fates of bacterial social parasites: persistence and
self-induced extinction of Myxococcus xanthus cheaters. *Proc Biol Sci,* 270**,** 1527-34.
FISHER, R. M. & REGENBERG, B. 2019. Multicellular group formation in Saccharomyces
cerevisiae. *Proc Biol Sci,* 286**,** 20191098.
FRANK, S. A. 1997. The Price Equation, Fisher's fundamental theorem, kin selection, and causal
analysis. *Evolution,* 51**,** 1712-1729.
FRANK, S. A. 2012. Natural selection. IV. The Price equation. *J Evol Biol,* 25**,** 1002-19.
GHOUL, M., GRIFFIN, A. S. & WEST, S. A. 2014. Toward an evolutionary definition of cheating.
Evolution, 68**,** 318-31.
GITSCHLAG, B. L., KIRBY, C. S., SAMUELS, D. C., GANGULA, R. D., MALLAL, S. A. & PATEL, M. R.
2016. Homeostatic Responses Regulate Selfish Mitochondrial Genome Dynamics in C.
elegans. *Cell Metab,* 24**,** 91-103.
GITSCHLAG, B. L., TATE, A. T. & PATEL, M. R. 2020. Nutrient status shapes selfish mitochondrial
genome dynamics across different levels of selection. *Elife,* 9.
GULLI, J. G., HERRON, M. D. & RATCLIFF, W. C. 2019. Evolution of altruistic cooperation among
nascent multicellular organisms. *Evolution,* 73**,** 1012-1024.
HAMMERSCHMIDT, K., ROSE, C. J., KERR, B. & RAINEY, P. B. 2014. Life cycles, fitness decoupling
and the evolution of multicellularity. *Nature,* 515**,** 75-9.
HAMMOND, T. M., REHARD, D. G., XIAO, H. & SHIU, P. K. 2012. Molecular dissection of
Neurospora Spore killer meiotic drive elements. *Proc Natl Acad Sci U S A,* 109**,** 12093-8.
HAVIRD, J. C., FORSYTHE, E. S., WILLIAMS, A. M., WERREN, J. H., DOWLING, D. K. & SLOAN, D. B.
2019. Selfish Mitonuclear Conflict. *Curr Biol,* 29**,** R496-R511.
HOLT, I. J., HARDING, A. E. & MORGAN-HUGHES, J. A. 1988. Deletions of muscle mitochondrial
DNA in patients with mitochondrial myopathies. *Nature,* 331**,** 717-9.
HU, W., JIANG, Z. D., SUO, F., ZHENG, J. X., HE, W. Z. & DU, L. L. 2017. A large gene family in
fission yeast encodes spore killers that subvert Mendel's law. *Elife,* 6.
KANDUL, N. P., ZHANG, T., HAY, B. A. & GUO, M. 2016. Selective removal of deletion-bearing
mitochondrial DNA in heteroplasmic Drosophila. *Nat Commun,* 7**,** 13100.
KLUCNIKA, A. & MA, H. 2019. A battle for transmission: the cooperative and selfish animal
mitochondrial genomes. *Open Biol,* 9**,** 180267.
KONRAD, A., THOMPSON, O., WATERSTON, R. H., MOERMAN, D. G., KEIGHTLEY, P. D.,
BERGTHORSSON, U. & KATJU, V. 2017. Mitochondrial Mutation Rate, Spectrum and

Heteroplasmy in Caenorhabditis elegans Spontaneous Mutation Accumulation Lines of Differing Population Size. *Mol Biol Evol,* 34**,** 1319-1334.

LARRACUENTE, A. M. & PRESGRAVES, D. C. 2012. The selfish Segregation Distorter gene complex of Drosophila melanogaster. *Genetics,* 192**,** 33-53.

LEIGH, E. G. 1983. When does the good of the group override the advantage of the individual? *Proc Natl Acad Sci U S A,* 80**,** 2985-9.

LIAU, W. S., GONZALEZ-SERRICCHIO, A. S., DESHOMMES, C., CHIN, K. & LAMUNYON, C. W. 2007. A persistent mitochondrial deletion reduces fitness and sperm performance in heteroplasmic populations of C. elegans. *BMC Genet,* 8**,** 8.

LIEBER, T., JEEDIGUNTA, S. P., PALOZZI, J. M., LEHMANN, R. & HURD, T. R. 2019. Mitochondrial fragmentation drives selective removal of deleterious mtDNA in the germline. *Nature,* 570**,** 380-384.

LIN, Y. F., SCHULZ, A. M., PELLEGRINO, M. W., LU, Y., SHAHAM, S. & HAYNES, C. M. 2016. Maintenance and propagation of a deleterious mitochondrial genome by the mitochondrial unfolded protein response. *Nature,* 533**,** 416-9.

MICHOD, R. E., VIOSSAT, Y., SOLARI, C. A., HURAND, M. & NEDELCU, A. M. 2006. Life-history evolution and the origin of multicellularity. *J Theor Biol,* 239**,** 257-72.

MORENO-FENOLL, C., CAVALIERE, M., MARTINEZ-GARCIA, E. & POYATOS, J. F. 2017. Eco-evolutionary feedbacks can rescue cooperation in microbial populations. *Sci Rep,* 7**,** 42561.

NAMSLAUER, I. & BRZEZINSKI, P. 2009. A mitochondrial DNA mutation linked to colon cancer results in proton leaks in cytochrome c oxidase. *Proc Natl Acad Sci U S A,* 106**,** 3402-7.

NARGUND, A. M., PELLEGRINO, M. W., FIORESE, C. J., BAKER, B. M. & HAYNES, C. M. 2012. Mitochondrial import efficiency of ATFS-1 regulates mitochondrial UPR activation. *Science,* 337**,** 587-90.

PRICE, G. R. 1972. Extension of Covariance Selection Mathematics. *Annals of Human Genetics,* 35**,** 485-&.

PRUITT, J. N. & RIECHERT, S. E. 2009. Frequency-dependent success of cheaters during foraging bouts might limit their spread within colonies of a socially polymorphic spider. *Evolution,* 63**,** 2966-73.

QUELLER, D. C. & STRASSMANN, J. E. 2009. Beyond society: the evolution of organismality. *Philos Trans R Soc Lond B Biol Sci,* 364**,** 3143-55.

RAINEY, P. B. & RAINEY, K. 2003. Evolution of cooperation and conflict in experimental bacterial populations. *Nature,* 425**,** 72-4.

RIEHL, C. & FREDERICKSON, M. E. 2016. Cheating and punishment in cooperative animal societies. *Philos Trans R Soc Lond B Biol Sci,* 371**,** 20150090.

ROSS-GILLESPIE, A., GARDNER, A., WEST, S. A. & GRIFFIN, A. S. 2007. Frequency dependence and cooperation: theory and a test with bacteria. *Am Nat,* 170**,** 331-42.

SAGAN, L. 1967. On the origin of mitosing cells. *J Theor Biol,* 14**,** 255-74.

SCHIMENTI, J. 2000. Segregation distortion of mouse t haplotypes the molecular basis emerges. *Trends Genet,* 16**,** 240-3.

SHOU, W. 2015. Acknowledging selection at sub-organismal levels resolves controversy on pro-cooperation mechanisms. *Elife,* 4.

686   SUEN, D. F., NARENDRA, D. P., TANAKA, A., MANFREDI, G. & YOULE, R. J. 2010. Parkin
687   overexpression selects against a deleterious mtDNA mutation in heteroplasmic cybrid
688   cells. *Proc Natl Acad Sci U S A,* 107**,** 11835-40.
689   TAYLOR, R. W. & TURNBULL, D. M. 2005. Mitochondrial DNA mutations in human disease. *Nat*
690   *Rev Genet,* 6**,** 389-402.
691   TEPPER, R. G., ASHRAF, J., KALETSKY, R., KLEEMANN, G., MURPHY, C. T. & BUSSEMAKER, H. J.
692   2013. PQM-1 complements DAF-16 as a key transcriptional regulator of DAF-2-mediated
693   development and longevity. *Cell,* 154**,** 676-690.
694   TSANG, W. Y. & LEMIRE, B. D. 2002. Stable heteroplasmy but differential inheritance of a large
695   mitochondrial DNA deletion in nematodes. *Biochem Cell Biol,* 80**,** 645-54.
696   VALENCI, I., YONAI, L., BAR-YAACOV, D., MISHMAR, D. & BEN-ZVI, A. 2015. Parkin modulates
697   heteroplasmy of truncated mtDNA in Caenorhabditis elegans. *Mitochondrion,* 20**,** 64-70.
698   VARLAMOV, D. A., KUDIN, A. P., VIELHABER, S., SCHRODER, R., SASSEN, R., BECKER, A., KUNZ,
699   D., HAUG, K., REBSTOCK, J., HEILS, A., ELGER, C. E. & KUNZ, W. S. 2002. Metabolic
700   consequences of a novel missense mutation of the mtDNA CO I gene. *Hum Mol Genet,*
701   11**,** 1797-805.
702   WEBB, A. E., KUNDAJE, A. & BRUNET, A. 2016. Characterization of the direct targets of FOXO
703   transcription factors throughout evolution. *Aging Cell,* 15**,** 673-85.
704   WENSELEERS, T. & RATNIEKS, F. L. 2004. Tragedy of the commons in Melipona bees. *Proc Biol*
705   *Sci,* 271 Suppl 5**,** S310-2.
706   WEST, S. A., FISHER, R. M., GARDNER, A. & KIERS, E. T. 2015. Major evolutionary transitions in
707   individuality. *Proc Natl Acad Sci U S A,* 112**,** 10112-9.
708   YANG, Q., LIU, P., ANDERSON, N. S., SHPILKA, T., DU, Y., NARESH, N. U., LI, R., ZHU, L. J., LUK, K.,
709   LAVELLE, J., ZEINERT, R. D., CHIEN, P., WOLFE, S. A. & HAYNES, C. M. 2022. LONP-1 and
710   ATFS-1 sustain deleterious heteroplasmy by promoting mtDNA replication in
711   dysfunctional mitochondria. *Nat Cell Biol,* 24**,** 181-193.
712   ZHANG, H., ESPOSITO, M., PEZET, M. G., ARYAMAN, J., WEI, W., KLIMM, F., CALABRESE, C.,
713   BURR, S. P., MACABELLI, C. H., VISCOMI, C., SAITOU, M., CHIARATTI, M. R., STEWART, J.
714   B., JONES, N. & CHINNERY, P. F. 2021. Mitochondrial DNA heteroplasmy is modulated
715   during oocyte development propagating mutation transmission. *Sci Adv,* 7**,** eabi5657.
716
717

Figure 1. Mitochondrial mutations featured in this study. (**A**) Map of the mitochondrial genome in *C. elegans* showing location and size of each heteroplasmic mutation featured in this study. The relative sizes and locations of protein and ribosomal RNA genes are represented by the gray arrows; thick black bars represent tRNA genes; thin black lines represent noncoding regions. The *uaDf5* deletion featured in previous work (Gitschlag et al., 2020) is shown in dark red. Colored bars around the perimeter of the genome represent approximate locations and sizes of the remaining mtDNA deletions used in this study. Mutations are color-coded corresponding to the visualization of data in panel (**B**) and in subsequent figures in this study. (**B**) Heteroplasmic mutant mtDNA frequency sampled from N=4 lines stably maintaining each mutation featured in (**A**), each crossed into the wildtype (Bristol strain) nuclear background, and quantified using droplet digital PCR.

Figure 2. Experimental approach to characterize multilevel selection dynamics of mutant mtDNA. (**A**) Longitudinal experiment, featuring measurement of mutant mtDNA between individual parents and their own respective adult progeny, to quantify sub-organismal selection. (**B**) Competition experiment designed to quantify organismal selection against mutant mtDNA. (**C**) Graphical depiction of covariance between a quantifiable trait value and selection coefficient. Covariance implies that variation in a measured trait value *z*, such as mutant mtDNA frequency, corresponds to variation in the selection coefficient *S*, a measure of the strength of the selection acting on *z*. A slope of zero indicates that variation in the trait does not correspond to variation in the fitness effects of the trait.

Figure 3. Integration of the multilevel selection dynamics of *uaDf5* using the Price Equation. (**A**) Linear regression between *uaDf5* frequency and the strength of sub-organismal selection, obtained from converting the previously measured intergenerational shifts in *uaDf5* frequency (Gitschlag et al., 2020) to selection coefficients (red circles) and plotting them as a function of parental *uaDf5* frequency. Red shaded regions represents 95% confidence interval. F-test for non-zero slope (sample size and raw data shown in Figure 3—Source Data 1). (**B**) Non-linear regression between heteroplasmic (not population-wide) *uaDf5* frequency and the strength of organismal selection, obtained from converting the previously measured declines in *uaDf5* frequency from the organismal competition experiment (Gitschlag et al., 2020) to selection coefficients (red circles). Solid regression and shaded region represents the best fit and 95% confidence interval, respectively, assuming the organismal selection coefficient reaches 1 (lethality) when *uaDf5* frequency is 100%. For comparison, dashed line and horizontally-striped region represents the best fit and 95% confidence interval, respectively, assuming the organismal selection coefficient reaches 1 when *uaDf5* frequency is 90%. Dotted line and vertically-striped region represents the best fit and 95% confidence interval, respectively, assuming the selection coefficient increases linearly with *uaDf5* frequency (raw data shown in Figure 3—Source Data 2). (**C**) Overall population-level

regression (summation of sub-organismal and organismal regressions in panels **A**-**B**), showing the

population-level consequences of each organismal covariance model assumed in panel (**B**). For each

graph, the black horizontal dotted line represents no net selection. Above the dotted horizontal line:

selection favors wildtype mtDNA. Below the dotted horizontal line: selection favors *uaDf5*.

Figure 4. Data from experiments designed to measure influence of sub-organismal and organismal levels

of selection on mtDNA heteroplasmies. (**A-B**) Intergenerational shifts in *mpt1* frequency at the levels of

sub-organismal (**A**) and organismal (**B**) selection. Wilcoxon matched-pairs signed-rank test, N=21 (**A**).

Linear regression analysis (**B**). (**C-D**) Same as above but for the *mpt2* heteroplasmy. N=23 (**C**). (**E-F**)

Same as above but for the *mptDf3* heteroplasmy. N=23 (**E**). (**G-H**) Same as above but for the *mptDf2*

heteroplasmy. N=24 (**G**). (**I-J**) Same as above but for the *mpt4* heteroplasmy. N=24 (**I**). Raw data shown

in Figure 4—Source Data 1 and Figure 4—Source Data 2.

Figure 5. Integration of the multilevel selection dynamics of all five heteroplasmies using the Price

Equation framework. For each graph shown above, the horizontal dotted line represents no net selection.

Above the dotted horizontal line: selection favors wildtype mtDNA. Below the dotted horizontal line:

selection favors ΔmtDNA. (**A-C**) The plotted regressions between heteroplasmic frequency and selection

coefficient at sub-organismal, organismal, and overall (combined) levels, using data from Figure 4A-B.

Same analysis as in Figure 3 but applied to the *mpt1* heteroplasmy. (**D-F**) Same as above but for *mpt2*

(data from Figure 4C-D). (**G-I**) Same as above but for *mptDf3* (data from Figure 4E-F). (**J-L**) Same as

above but for *mptDf2* (data from Figure 4G-H). (**M-O**) Same as above but for *mpt4* (data from Figure 4I-J).

F-test for non-zero slope (panels **A**, **D**, **G**, **J**, and **M**; sample sizes and mutant frequency data shown in

Figure 4 and Figure 4—Figure Supplement 1). As before (see Figure 3), the organismal and combined

("overall") plots respectively represent the 95% confidence ranges that result from assuming the selection

coefficient reaches 1 (lethality) when the mutant is at 100 percent frequency. For visual simplicity, only

this assumption is shown here; see Figure 5—Figure Supplement 1 for consequences of alternately

assuming the organismal selection coefficient reaches 1 when mutant mtDNA frequency is 90% or

increases linearly with mutant mtDNA frequency. Note that confidence ranges capable of accommodating

a horizontal regression indicate a slope that is not significantly non-zero, implying no significant

relationship between mutant mtDNA frequency and selection coefficient, at least within the mutant

frequency range in which the data was collected.

Figure 6. Alternate population dynamics depending on mitochondrial genotype. **Left**: sub-organismal or

within-host dynamics favor the preferential proliferation of mutant over wildtype mtDNA across individual

parent-progeny lineages, when the effect of organismal selection is minimized. Simultaneously, selection

for host fitness strongly favors wildtype over mutant mtDNA. These dynamics are observed for

heteroplasmic mtDNA variants *uaDf5*, *mptDf2*, and *mpt4*. In contrast to *uaDf5* and *mptDf2*, however, the

sub-organismal proliferation of *mpt4* does not appear to be frequency-dependent (Figure 5M). **Right**: for

some heteroplasmic variants (*mpt1*, *mpt2*, and *mptDf3*), neither sub-organismal dynamics nor selection at

the organismal level favor a significant shift in mutant mtDNA frequency, at least at the frequencies at

which these mutant genomes were observed to stably persist (Figure 1B).

851   **Figure 3—Figure Supplement 1.** Mathematical framework for multilevel selection analysis of *uaDf5*.
852
853   Here, the regressions shown in Figure 3 are derived from the data obtained from the measurements of
854   organismal and sub-organismal selection on *uaDf5* (Gitschlag et al., 2020), using the general framework
855   of the Price Equation. Hereafter, *uaDf5* is referred to as ΔmtDNA to denote a deletion-bearing genome, to

856 emphasize that this framework can be applied to other mutant mtDNA (Figure 5). We begin with the Price
857 Equation:
858

$$w\Delta z = cov(w,z)_{org} + E[cov(w,z)_{sub}]$$

859
Where:
w = reproductive fitness, or number of replication events per generation, for one trait relative to another
z = measurable value of a trait (in this case, mutant mtDNA frequency)
$cov(w,z)_{org}$ = covariance between w and z contributed by selection at the organismal level
$E[cov(w,z)_{sub}]$ = expected or average covariance between w and z contributed by sub-organismal
selection
866
867 Note that the version of the Price Equation shown above is equivalent to the form shown as equation A17
868 in Price, 1972. At each level of selection, organismal and sub-organismal, the term for the covariance
869 between w and z can be defined as follows:
870

$$cov(w,z) = \frac{\sum(w_i - w_{average})(z_i - z_{average})}{n-1}$$

871
Where:
w_i = fitness w of trait z in the ith replicate parent-progeny lineage from the sample or population
$w_{average}$ = average w across the sample or population
z_i = value of trait z in the parent of the ith replicate parent-progeny lineage from the sample or population
$z_{average}$ = average value of z across the sample or population
n = number of parent-progeny lineages sampled
878
879 Since the selection coefficient, which measures the strength of selection acting on a trait, is directly to
880 fitness, the Price Equation can be expressed using covariance terms between selection coefficient, S,
881 and the value of a trait, z. Defining trait-value as the heteroplasmic (sub-organismal or within-host)
882 frequency of mutant mtDNA ($\%\Delta mtDNA$) yields the following form of the Price Equation:
883

$$Change\ in\ \%\Delta mtDNA = cov(S, \%\Delta mtDNA)_{org} + E[cov(S, \%\Delta mtDNA)_{sub}]$$

884
885 Replacing w and z in the Price Equation with S and $\%\Delta mtDNA$ yields the following formula for covariance:
886

$$cov(S, \%\Delta mtDNA) = \frac{\sum(S_i - S_{average})(\%\Delta mtDNA_i - \%\Delta mtDNA_{average})}{n-1}$$

887
Where:
S_i = selection coefficient of ΔmtDNA in the ith replicate of each experiment (sub-organismal and
organismal)
$S_{average}$ = average S across the sample or population of each experiment
$\%\Delta mtDNA_i$ = heteroplasmic ΔmtDNA frequency from the ith parent (sub-organismal) or line (organismal)
$\%\Delta mtDNA_{average}$ = average heteroplasmic ΔmtDNA frequency across all replicates
894
895 In the sub-sections that follow, the individual regressions are derived for the strength of sub-organismal
896 and organismal selection as a function of heteroplasmic ΔmtDNA frequency, shown using the data from
897 (Gitschlag et al., 2020). (see also Figure 3—Source data files). The multilevel selection analyses in Figure
898 5 are performed using the same approach and data shown in Figure 4.
899
900
901
902
903 **Sub-organismal regression (Figure 3A)**
904

For selection acting on ΔmtDNA at the sub-organismal level, selection coefficient, S, is calculated for a given change in ΔmtDNA frequency between parent and progeny using the following equation:

$$S = 1 - w = 1 - \frac{\%\Delta mtDNA_{g+1}/\%\Delta mtDNA_g}{\%wildtype\ mtDNA_{g+1}/\%wildtype\ mtDNA_g}$$

Where:
w = average replicative fitness of ΔmtDNA relative to wildtype mtDNA
g = parental generation

The following regression, relating selection coefficient to heteroplasmic ΔmtDNA frequency, can be derived using the covariance relationship defined on the previous page and the data shown in Figure 2—Source data 1:

$$S = slope(S, \%\Delta mtDNA) * \%\Delta mtDNA + intercept(S, \%\Delta mtDNA)$$

Note that this equation takes the form of the slope-intercept equation, $y=mx+b$, in which y corresponds to selection coefficient, S, and x corresponds to heteroplasmic ΔmtDNA frequency, denoted *%ΔmtDNA*. The slope, m, can be calculated from its relationship to covariance and variance. In particular, the covariance between S and *%ΔmtDNA* is equal to the variance of *%ΔmtDNA* multiplied by the slope of the regression (Price, 1970):

$$cov(S, \%\Delta mtDNA) = var(\%\Delta mtDNA) * slope(S, \%\Delta mtDNA)$$

Here, the term *var(S,%ΔmtDNA)* represents the variance of *%ΔmtDNA*, calculated as follows:

$$var(\%\Delta mtDNA) = \frac{\sum(\%\Delta mtDNA_i - \%\Delta mtDNA_{average})^2}{n-1}$$

Note that the formula for variance is similar to the formula for covariance, with one exception: the errors or residuals in S (S_i - $S_{average}$) are replaced by the errors in *%ΔmtDNA* (*%ΔmtDNA$_i$* - *%ΔmtDNA$_{average}$*). In other words, in contrast to the covariance of S and *%ΔmtDNA*, the variance of *%ΔmtDNA* is equivalent to the covariance of *%ΔmtDNA* with itself. The term *slope(S,%ΔmtDNA)*, shown above, represents the slope of the regression obtained by plotting S as a function of *%ΔmtDNA*. The slope can be calculated by rearranging the relation between covariance, variance, and slope, as follows:

$$slope(S, \%\Delta mtDNA) = \frac{cov(S, \%\Delta mtDNA)}{var(\%\Delta mtDNA)} = \frac{\sum(S_i - S_{average})(\%\Delta mtDNA_i - \%\Delta mtDNA_{average})}{\sum(\%\Delta mtDNA_i - \%\Delta mtDNA_{average})^2}$$

Entering the empirically derived values for S and *%ΔmtDNA* (Figure 3—Source data 1) yields a slope of 0.01154. Next, the slope-intercept equation can be rearranged to solve for the intercept, as follows:

$$intercept(S, \%\Delta mtDNA) = S - slope(S, \%\Delta mtDNA) * \%\Delta mtDNA$$

By entering the slope, together with any pair of corresponding values of S and *%ΔmtDNA* that fall on the regression line (such as $S_{average}$ and *%ΔmtDNA$_{average}$*, taken from Figure 3—Source data 1), the intercept is calculated to be -0.923 as follows:

$$intercept(S, \%\Delta mtDNA) = S - slope(S, \%\Delta mtDNA) * \%\Delta mtDNA = -0.289 - 0.01154 * 54.96 = -0.923$$

Accordingly, the covariance between selection coefficient and ΔmtDNA frequency at the sub-organismal level yields the following relation:

$$S = 0.01154 * \%\Delta mtDNA - 0.923$$

950
951 This equation defines the regression line shown in Figure 3A. The confidence interval surrounding the
952 regression (defining the shaded region) is calculated for each value of ΔmtDNA frequency, from 0 to 100,
953 using the following formula:
954

$$95\% \; confidence \; interval = t_{n-2} * \sigma_R \sqrt{\frac{1}{n} + \frac{(\%\Delta mtDNA^* - \%\Delta mtDNA_{ave})^2}{\sigma_{RS}^2(n-1)}}$$

955
<u>Where:</u>
t_{n-2} = critical value from Student's t-table for *n-2*, two-tailed, α=0.05
σ_R = standard deviation of the residuals for selection coefficient *S*
n = sample size
%ΔmtDNA* = each value of ΔmtDNA frequency at which confidence interval is being calculated
%ΔmtDNA$_{ave}$ = average ΔmtDNA frequency across all samples
σ_{RS} = standard deviation of the residuals of *S*, with each residual defined as $S_{actual} - S_{predicted}$
963
964
965 **Organismal regression (Figure 3B)**
966
For selection acting on ΔmtDNA at the organismal level, in other words the selection against ΔmtDNA that
arises strictly from the cost that ΔmtDNA imposes on host fitness, the selection coefficient, *S*, is
calculated from the change in ΔmtDNA frequency across a population consisting of heteroplasmic hosts
(carrying ΔmtDNA) competed against their wildtype counterparts on the same food plate. In contrast to
the sub-organismal selection experiment, where *S* is calculated from the change in ΔmtDNA frequency
between parents and their immediate progeny, the organismal competition experiment involved
measuring the change in ΔmtDNA frequency across multiple successive generations. Accordingly, the
selection coefficient can be calculated as follows:
975

$$S = 1 - w = 1 - e^{slope[g,LN(\%\Delta mtDNA/\%WT \, mtDNA)]}$$

976
<u>Where:</u>
LN = natural logarithm
%ΔmtDNA = population-wide (not heteroplasmic) ΔmtDNA frequency
%WT mtDNA = population-wide wildtype mtDNA frequency (equal to 100 – *%ΔmtDNA*)
g = time, measured in number of generations
LN(%ΔmtDNA / %WT mtDNA)] = natural logarithm of the ratio of *%ΔmtDNA* to *%WT mtDNA*
slope[*g,LN(%ΔmtDNA / %WT mtDNA)*] = slope of *LN(%ΔmtDNA / %WT mtDNA)*] versus time, as follows:
984

$$slope[g, LN(\%\Delta mtDNA/\%WT \, mtDNA)] = \frac{\sum \left[LN\left(\frac{\Delta}{W}\right)_i - LN\left(\frac{\Delta}{W}\right)_{average} \right](g_i - g_{average})}{\sum(g_i - g_{average})^2}$$

985
986 <u>Where:</u>
987 $LN(\Delta/W)_i$ = *LN(%ΔmtDNA / %WT mtDNA)* for the *i*th generation
988 $LN(\Delta/W)_{average}$ = average value of $LN(\%\Delta mtDNA_I \, / \, \%WT \, mtDNA)$ across all generations
989 g_i = the *i*th generation
990 $g_{average}$ = average g_i value across all generations (equal to 5.5 for generations 1 to 10)
991
The selection coefficient of ΔmtDNA is calculated for both the competed and non-competed populations
in the organismal competition experiment. Although ΔmtDNA persists within a stable frequency range in
non-competed populations, ΔmtDNA undergoes slight variation in frequency, perhaps due to drift,
variations in sub-organismal selection, and population structure (during population expansion at the onset
of each new generation, adults become rapidly outnumbered by embryos and larvae, which tend to carry

lower ∆mtDNA than adults, see "Non-competing" section in Figure 3—Source data 2). In any case, to
control for sources other than organismal selection (that is, variation in ∆mtDNA frequency occurring for
reasons other than the fitness differences between heteroplasmic animals and their wildtype
counterparts), the average selection coefficient across all replicate non-competed populations (containing
exclusively heteroplasmic animals) is subtracted from the selection coefficient of each replicate competed
population, yielding the selection coefficient for organismal selection against ∆mtDNA:
1003

$$S_{organismal} = S_{competed} - \bar{S}_{non-competed}$$

1004
1005 <u>Where:</u>
1006 $S_{organismal}$ = selection coefficient, S, strictly at the level of organismal selection
1007 $S_{competed}$ = S across a competed line
1008 $\bar{S}_{non-competed}$ = average S across all non-competed lines
1009
Deriving a regression between selection coefficient and heteroplasmic ∆mtDNA frequency is trickier at the
organismal level, since the relationship is non-linear. In particular, a large deletion such as in ∆mtDNA
results in the loss of essential genes and is therefore expected to be lethal when present at very high
frequency (fitness w = 0, or S = 1). However, the organismal selection coefficient is approximately only
0.1892 at a heteroplasmic frequency of 60% ∆mtDNA. Moreover, S is expected to be 0 when ∆mtDNA
frequency is zero, since no selection is occurring against ∆mtDNA among hosts that lack ∆mtDNA. The
appropriate regression will therefore constitute a non-linear trend-line that rises sharply at progressively
higher ∆mtDNA frequency, passing through S=0 when %∆mtDNA=0, S=0.189 when %∆mtDNA=60, and
S=1 when %∆mtDNA=100. An ideal candidate regression would therefore be logarithmic, in which rather
than being directly proportional to ∆mtDNA frequency, S is equal to some quantity raised to the power of
an exponent (which itself is proportional to ∆mtDNA frequency). The least-squares regression method
can be used to fit a non-linear regression. As the name suggests, this method seeks to minimize the
square of the errors, or residuals, along the dependent variable (in this case S), which can be calculated
as shown:
1024

$$SSQ = \sum [S_i - f(\%\Delta mtDNA)]^2$$

1025
1026 <u>Where:</u>
1027 SSQ = sum of the squares of residuals
1028 S_i = organismal selection coefficient for replicate line i in the organismal competition experiment
1029 $f(\%\Delta mtDNA)$ = predicted value of S, defined as a function of %∆mtDNA
1030 $S_i - f(\%\Delta mtDNA)$ = error, or residual, between actual and predicted values of S
1031
1032 The least-squares method, computed here using the GraphPad Prism software package, fits a regression
1033 by finding the curve with a shape that minimizes the value of SSQ. When assuming S=0 at 0% ∆mtDNA
1034 and S=1 at 100% ∆mtDNA, together with the empirically derived value of S=0.1892 at 60% ∆mtDNA, the
1035 least-squares method identifies the following regression to minimize SSQ:
1036

$$S = 10^{0.0181 * \%\Delta mtDNA - 1.81}$$

1037
1038 However, S may approach 1 at a ∆mtDNA frequency lower than 100%. Consistent with this possibility,
1039 hosts are almost never observed to carry ∆mtDNA at or above 90% frequency. To account for the
1040 uncertainty in how the fitness cost of ∆mtDNA scales with ∆mtDNA frequency, the following regression
1041 was obtained using the least-squares method when assuming that S=1 at 90% instead of 100% ∆mtDNA:
1042

$$S = 10^{0.0241 * \%\Delta mtDNA - 2.17}$$

1043
1044 Finally, S might rise modestly until reaching a particular (albeit unknown) ∆mtDNA frequency threshold,
1045 followed by an abrupt shift to higher S in a sort of stepwise fashion. To account for this uncertainty, the
1046 assumption that S reaches 1 at a given ∆mtDNA frequency was eliminated, enabling a standard linear
1047 regression between S=0 at 0% ∆mtDNA and S=0.1892 at 60% ∆mtDNA:

1048

$$S = 0.003153 * \%\Delta mtDNA$$

1049
1050   The confidence interval for the non-linear regressions can be calculated using the following formula:
1051

$$95\% \; confidence \; interval = t_{n-1}\sqrt{CM * dS^2}$$

1052
1053   Where:
1054 t_{n-1} = critical value from student's t-table for *n-1*
1055   *CM* = covariance matrix value
1056   *dS* = derivative of the non-linear regression *S* = *f(%ΔmtDNA)* (Figure 3—Source data 2)
1057   *n* = sample size
1058
1059   The confidence interval for the linear regressions can be calculated in a similar manner as in the case of
1060   the linear regressions between *S* and *%ΔmtDNA* at the sub-organismal level (see the previous sub-
1061   section):
1062

$$95\% \; confidence \; interval = t_{n-2} * \sigma_R \sqrt{\frac{1}{n} + \frac{(\%\Delta mtDNA^* - \%\Delta mtDNA_{ave})^2}{\sigma_{\%\Delta mtDNA}{}^2(n-1)}}$$

1063
Where:
t_{n-2} = critical value from student's t-table for *n-2*
σ_R = standard deviation of the residuals for selection coefficient *S* (Figure 3—Source data 2)
%ΔmtDNA* = each value of ΔmtDNA frequency at which the confidence interval is being calculated
%ΔmtDNA$_{ave}$ = average ΔmtDNA frequency across input values (*S*=0.1892 at 60% and *S*=0 at 0%
ΔmtDNA)
$\sigma_{\%\Delta mtDNA}$ = standard deviation of ΔmtDNA frequency
1071
1072
1073   **Overall (combined) regression (Figure 3C)**
1074
1075   According to the general framework of the Price Equation, the overall relationship between ΔmtDNA
1076   frequency and the strength of selection—when both the sub-organismal and organismal levels are taken
1077   into account—results from summing the regressions as well as their confidence intervals. The overall
1078   relationship is therefore defined as the following (assuming $S_{organismal}$=1 at 100% ΔmtDNA):
1079

$$S_{overall} = S_{sub-organismal} + S_{organismal} = (0.01154 * \%\Delta mtDNA - 0.923) + (10^{0.0181 * \%\Delta mtDNA - 1.81})$$

1080
1081   Assuming $S_{organismal}$=1 at 90% ΔmtDNA:
1082

$$S_{overall} = S_{sub-organismal} + S_{organismal} = (0.01154 * \%\Delta mtDNA - 0.923) + (10^{0.0241 * \%\Delta mtDNA - 2.17})$$

1083
1084   Assuming $S_{organismal}$ rises linearly from 0% ΔmtDNA to 60% ΔmtDNA and beyond:
1085

$$S_{overall} = S_{sub-organismal} + S_{organismal} = (0.01154 * \%\Delta mtDNA - 0.923) + (0.003153 * \%\Delta mtDNA)$$

1086
1087   Finally, for each regression, the combined confidence interval can be calculated as follows:
1088

$$95\% \; confidence \; interval_{overall} = \frac{1}{2} * \sqrt{(2 * 95\% CI_{sub-org})^2 + (2 * 95\% CI_{org})^2}$$

1089
1090   Where:
1091   *95%CI$_{sub-org}$* = confidence interval for the regression of *S* and *%ΔmtDNA* at the sub-organismal level
1092   *95%CI$_{org}$* = confidence interval for the regression of *S* and *%ΔmtDNA* at the organismal level

1093

1094

Figure 4—Figure Supplement 1. Intergenerational shifts in mutant mtDNA frequency at the sub-

organismal level, similar to Figure 2B in (Gitschlag et al., 2020) and Figures 1C and 8B,D,F in (Zhang et

al., 2021). (**A**) Shift in *mpt1* frequency per generation, obtained by subtracting *mpt1* frequency of mature

adult progeny in Figure 4A from *mpt1* frequency of the respective parent, plotted as a function of parental

mpt1 frequency. (**B**) Shift in *mpt2* frequency per generation, obtained by subtracting *mpt2* frequency of

mature adult progeny in Figure 4C from *mpt2* frequency of the respective parent, plotted as a function of

parental *mpt2* frequency. (**C**) Shift in *mptDf3* frequency per generation, obtained by subtracting *mptDf3*

frequency of mature adult progeny in Figure 4E from *mptDf3* frequency of the respective parent, plotted

as a function of parental *mptDf3* frequency. (**D**) Shift in *mptDf2* frequency per generation, obtained by

subtracting *mptDf2* frequency of mature adult progeny in Figure 4G from *mptDf2* frequency of the

respective parent, plotted as a function of parental *mptDf2* frequency. (**E**) Shift in *mpt4* frequency per

generation, obtained by subtracting *mpt4* frequency of mature adult progeny in Figure 4I from *mpt4*

frequency of the respective parent, plotted as a function of parental *mpt4* frequency.

1108

1109

Figure 4—Figure Supplement 2. Raw measurements of decline in mutant frequency upon competition

with homoplasmic (wildtype) counterparts, across the collection of heteroplasmies featured in Figure 3.

(**A**) Same experiment as shown in Figure 4B but with y-axis expressed in raw (non-normalized) *mpt1*

frequency. (**B**) Same experiment as shown in Figure 4D but with y-axis expressed in raw *mpt2* frequency.

(**C**) Same experiment as shown in Figure 4F but with y-axis expressed in raw *mptDf3* frequency. (**D**)

Same experiment as shown in Figure 4H but with y-axis expressed in raw *mptDf2* frequency. (**E**) Same

experiment as shown in Figure 4J but with y-axis expressed in raw *mpt4* frequency. Straight lines

represent best-fit regression across all replicates.

1118

1119

Figure 5—Figure Supplement 1. Integration of the multilevel selection dynamics of all five

heteroplasmies using the Price Equation framework with alternate organismal covariance assumptions.

The same empirical data and analysis as in Figure 5 but with the following model adjustments: dashed

line and horizontally-striped region represents the best fit and 95% confidence interval, respectively,

assuming the organismal selection coefficient reaches 1 when mutant mtDNA frequency is 90%; dotted

line and vertically-striped region represents the best fit and 95% confidence interval, respectively,

assuming the selection coefficient changes linearly with mutant mtDNA frequency. For each graph shown

above, the horizontal dotted line represents no net selection. Above the dotted horizontal line: selection

favors wildtype mtDNA. Below the dotted horizontal line: selection favors mutant mtDNA. Note that

confidence ranges capable of accommodating a horizontal regression indicate a slope that is not

significantly non-zero, implying no significant relationship between mutant mtDNA frequency and

selection coefficient, at least within the mutant frequency range in which the data was collected.

1132