1 Capped nascent RNA sequencing reveals novel therapy-responsive enhancers in prostate
2 cancer

Kellie A. Cotter[1,#], Sagar R. Shah[2,3,#], Mauricio I. Paramo[2,3,#], Shaoke Lou[4,#], Li Yao[3,5], Philip D.
Rubin[1], You Chen[2,3], Mark Gerstein[4,*], Mark A. Rubin[1,6*], Haiyuan Yu[3,5,*]

[1]Department of Biomedical Research, University of Bern, Bern, 3008, CH
[2]Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853, US
[3]Weill Institute for Cell and Molecular Biology, Cornell University, Ithaca, NY 14853, US
[4]Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, CT 06520,
US
[5]Department of Computational Biology, Cornell University, Ithaca, NY 14853, US
[6]Bern Center for Precision Medicine, University of Bern and Inselspital, Bern, 3008, CH

[#]Contributed equally
*Correspondence to: mark.gerstein@yale.edu (M.G), mark.rubin@dbmr.unibe.ch (M.A.R) and
haiyuan.yu@cornell.edu (H.Y)

Abstract

Mounting evidence suggests that enhancer RNA (eRNA) transcription start sites (TSSs) provide higher sensitivity and specificity for enhancer identification than histone modifications and chromatin accessibility. The extent to which changes in eRNA transcription correspond to changes in enhancer activity, however, remains unclear. Here, we used precision run-on and capped RNA sequencing (PRO-cap) to assess changes in enhancer activity in response to treatment with the androgen receptor signaling inhibitor, enzalutamide (ENZ). We identified 6,189 high-confidence candidate enhancers in the human prostate cancer cell line, LNCaP; 853 of which demonstrated significant changes in activity in response to drug treatment. Notably, we found that 67% and 54% of drug-responsive enhancers did not show similar changes in activity in previous studies that utilized ChIP-seq and ATAC-seq, respectively. Strikingly, 79% of regions with increased eRNA transcription showed no other biochemical alterations, implying that PRO-cap can capture a set of precise changes in enhancer activity that classical approaches lack the sensitivity to detect. We performed *in vivo* functional validations of candidate enhancers and found that CRISPRi targeting of PRO-cap-specific drug-responsive enhancers impaired ENZ regulation of downstream target genes, suggesting that changes in eRNA TSSs mark true biological changes in enhancer activity with high sensitivity. Our study highlights the utility of using PRO-cap as a complementary approach to canonical biochemical methods for detecting precise changes in enhancer activity and, in particular, for better understanding disease progression and responses to treatment.

Main

First-line treatment for advanced metastatic prostate cancer (PCa) generally involves androgen deprivation therapy (ADT) to reduce the activity of androgen receptor (AR)(Heinlein and Chang, 2004). Of particular clinical concern is metastases of castration-resistant PCa (CRPC), where the disease has developed resistance to both first-line and second-generation AR signaling inhibitors (ARSi, e.g., enzalutamide)(Scher and Sawyers, 2005). Thus, determining both the mechanisms behind ADT resistance and the distinct signaling pathways activated in CRPC are essential to improving existing therapies and finding new potential drug targets.

Given the enrichment of genomic alterations in metastatic PCa(Armenia et al., 2018), there are increased efforts to understand the complexity of the PCa genome. Recent studies using whole genome sequencing have revealed several examples of alterations in gene regulatory regions. For example, upwards of 80% of samples were found to have a duplication of a region upstream of *AR,* which was then shown to be a previously unidentified enhancer(Quigley et al., 2018; Takeda et al., 2018; Viswanathan et al., 2018). This enhancer duplication was shown to increase the expression of *AR* and in doing so decrease sensitivity to ARSi. Therefore, it is important to precisely identify and characterize enhancer dynamics in response to therapeutic intervention in PCa. However, despite the exhaustive amount of sequencing information captured by WGS, our interpretation and understanding of the data is limited due to the incomplete annotation of the non-coding genome, including transcriptional regulatory elements such as enhancers.

In general, enhancer regions are currently defined by biochemical features such as chromatin accessibility (DNase I hypersensitivity or transposase accessibility) and histone modification marks (H3K27ac and H3K4me1) along with transcription factor binding profiles as determined by ChIP-seq(Gasperini et al., 2020). Large-scale reporter assays (e.g., STARR-seq or MPRA) have also been used to evaluate the enhancer potential of candidate DNA regions(Arnold et al., 2013; Kheradpour et al., 2013). However, these reporter assays have

consistently shown that less than 50% of regions with these biochemical annotations exert

enhancer activity(Kwasnieski et al., 2014; Vanhille et al., 2015). Likewise, similar techniques have

been used to identify thousands of potential enhancer regions in PCa by combining ChIP-seq and

whole genome STARR-seq(Kron et al., 2017; Liu et al., 2017; Sharma et al., 2013; Stelloo et al.,

2018). In addition, chromosome capture has been used to elucidate chromatin interactions

genome-wide and those loci specifically associated with AR and RNA Polymerase II using ChIA-

PET(Ramanand et al., 2020; Rhie et al., 2019; Taberlay et al., 2016; Zhang et al., 2019).

Interestingly, a recent study demonstrated that the majority of AR binding sites are not active

enhancers(Huang et al., 2021). Moreover, despite increased recruitment of AR to these regions,

dihydrotestosterone (DHT) stimulation did not increase enhancer activity in over half of the AR-

bound active enhancers. Perhaps most intriguing, however, CRISPR interference of some

"inactive" enhancers altered the expression of the enhancer-regulated gene, at times at a similar

level of interference to nearby "active" enhancers. Overall, this highlights the limitations of the

datasets produced thus far and the need for alternative methods for enhancer identification given

that epigenomic-mark-based approaches identify enhancers with a high false-positive rate, while

reporter assays cannot fully reproduce much of the biological complexity of large enhancers

regions in their native genomic context, in particular with regards to multiple enhancers acting on

a single gene.

More recently, widespread RNA polymerase II-mediated bidirectional transcription has

been observed in enhancer regions, which produce biochemically unstable transcripts known as

enhancer RNAs (eRNAs)(Core et al., 2014; Kim et al., 2010; Tippens et al., 2018). Although the

functional significance of eRNAs remains unclear, evidence suggests that enhancer transcription

corresponds with activation(Chen et al., 2018; Chen and Liang, 2020), with close to 50% of short

capped nascent RNAs that map to previously unannotated TSSs overlapping with episomal

reporter-validated enhancers(Henriques et al., 2018). Moreover, it has been found that ~95% of

putative active enhancers found within accessible chromatin drive local transcription and do so

using factors and mechanisms overwhelmingly similar to those of promoters(Core et al., 2014).

Recently, we performed systematic interrogation of enhancer elements and showed that active

enhancer units are precisely marked by divergent eRNA TSSs genome-wide. Moreover, although

eRNA transcription is closely correlated with histone marks, we saw that these epigenomic marks

offer lower resolution and specificity for enhancer activation than transcription initiation(Tippens

et al., 2020). Thus, these data support a model whereby transcription is required for distal

enhancer function, challenging classical methods that rely on chromatin accessibility and histone

modifications to identify active enhancers.

On average, enhancers transcribe at 5% the level of promoters(Core et al., 2014;

Henriques et al., 2018). Thus, due to their low abundance and instability, eRNA detection requires

highly sensitive alternative methods to standard RNA sequencing approaches. Recently, we

performed systematic comparisons of genome-wide RNA sequencing assays suitable for the

identification of active enhancers and found that the nuclear run-on followed by cap-selection

assays (namely, Global/Precision Run-On and capped RNA sequencing GRO/PRO-cap) provide

the highest sensitivity and specificity for eRNA detection and active enhancer identification across

the whole genome(Yao et al., 2022). Importantly, PRO-cap libraries undergo a series of cap state

selection reactions to modify the 5′ ends of transcripts and allow for the accurate identification of

transcription initiation sites (Figure 1A). As a result, PRO-cap is highly sensitive for capturing

eRNA transcription and therefore, a powerful tool for enhancer identification.

Here, we utilized changes in eRNA expression to assess changes in enhancer activity in

response to short-term treatment with the ARSi enzalutamide (ENZ). We applied PRO-cap to

enrich and sequence only nascent RNAs associated with engaged RNA polymerase and to

identify divergent transcription start sites marking active enhancers at base pair-resolution(Mahat

et al., 2016; Tippens et al., 2020). We identified over 6,000 candidate enhancers in LNCaP cells;

853 of which demonstrated significant changes in enhancer activity in response to ENZ treatment.

Importantly, these results identified a large percentage of therapy-responsive enhancers, which

were not previously shown to be responsive in studies utilizing other biochemical marks

(H3K27ac-ChIP-seq/ATAC-seq) or reporter assays (i.e., STARR-seq). Our study highlights the

utility of using eRNA transcription, in particular PRO-cap, as a generalizable and complementary

approach to canonical biochemical methods for detecting precise changes in enhancer activity,

specifically for applications in disease prognosis, progression, and treatment.

Results
Identification of active enhancers in a prostate cancer cell line model using PRO-cap

To measure enhancer activity in their native genomic and cellular context, we utilized PRO-cap

to preferentially sequence nascent transcriptional start sites including those generating eRNAs

(Figure 1A). Overall, this analysis in LNCaP cells identified 91,705 statistically significant peaks

(n = 2 replicates), 68.4% of which were in gene distal regions. 6,189 of these gene distal regions

contained divergent significant peaks on both the plus and minus strands and were extended by

200 bp in both directions and called as high confidence enhancer regions (Data S1). As expected,

the majority of these regions contained canonical biochemical marks delineating enhancers

including open chromatin (DNase I hypersensitivity, ATAC-seq), H3K27 acetylation, and H3K4

monomethylation, and are bound by prostate-enriched transcription factors (AR and FOXA1)

(Figure 1B-C). However, 11% (n = 674) of these regions did not exhibit those classical

biochemical features (at least two of the three: DNase I HS, H3K27ac, or H3K4me1),

demonstrating that PRO-cap identifies a novel set of enhancer regions that would otherwise be

missed (Figure 1C).

Using publicly available H3K27ac HiChIP data, we estimated the number of chromatin

connections of each candidate enhancer with other genomic regions(Giambartolomei et al.,

2021). We found that the regions delineated by PRO-cap have a significantly greater number of

loops (P < 0.0001, Kolmogorov-Smirnov test) than H3K27ac or DNase I peaks (Figure 1D). Next,

we inquired whether these enhancers identified in LNCaP cells were expressed in patient-derived

PCa samples. To that end, we analyzed published ChIP-seq data from clinical specimens of non-

neoplastic prostate, primary PCa, and metastatic PCa(Pomerantz et al., 2020). As seen in Figure

1E, these tissues recapitulate H3K27ac and the binding pattern of prostate-enriched transcription

factors (AR, FOXA1, and HOXB13) in enhancers seen in LNCaP cells. Furthermore, given that

LNCaP cells are derived from an androgen-responsive metastatic lymph node

lesion(Horoszewicz et al., 1983), the enhancer expression profiles are indeed strongest in

metastatic PCa tissues. Overall, these results demonstrate that, unlike traditional epigenomic-

based methods, PRO-cap can uncover previously unknown enhancer loci with hallmarks of PCa-

relevant TF-binding patterns observed in primary and metastatic lesions.

Measurement of changes in enhancer activity in response to androgen deprivation therapy

Given our confidence in detecting enhancers by PRO-cap, we investigated whether this method

could be utilized to assess changes in enhancer activity in response to treatment. To do so, we

treated LNCaP cells with the ARSi enzalutamide (ENZ) for 4 or 24 hours. Using the same analysis

as described above, we identified 6769 and 8513 high confidence enhancers using PRO-cap in

the 4- and 24-hour time point conditions, respectively (Data S2-3). Approximately 4,479 of these

enhancers were identified in both ENZ timepoints and 2,922 were identified irrespective of

treatment (Figure S1A).

In addition to detecting enhancers that are expressed at only a single treatment time point,

we were interested in identifying those which were active at more than one time point, but with

differential levels of expression. Thus, we compiled the total list of enhancers found in any of the

two treatment timepoints and vehicle and assessed their differential expression upon treatment

(versus vehicle) on at least one strand. This analysis identified 853 enhancers that were

significantly activated or repressed after 24 hours of ENZ treatment (Figure 2A, Figure S1B-D,

Data S4-5). Furthermore, 73 of these regions were significantly activated or repressed on both

the plus and minus strands.

To predict putative transcription factors (TF) regulating these enhancer regions, we utilized

GIGGLE(Layer et al., 2018) analysis (Figure 2B) to probe the CISTROME(Mei et al., 2017) ChIP-

seq database containing published datasets with overlapping genomic regions. Of the TFs

predicted to bind to the ENZ repressed elements, AR was the top candidate with the highest

GIGGLE score. Interestingly, other significant TF predicted to bind the ENZ repressed enhancers

included glucocorticoid receptor (*NR3C1*), HNF4G (a TF which drives ARSi resistance(Shukla et

al., 2017)), and the co-activator EP300, the SWI/SNF family chromatin remodeler SMARCA4, and

the pioneering factor FOXA1. In addition to AR, FOXA1, and NR3C1, other TF predicted to bind

the ENZ-activated enhancers included the histone methyltransferase ASHL2, the AR co-

regulators GRHL2 and HOXB13, the AR-regulated TF NKX3-1, the pioneering factor GATA2,

ARID1A another SWI/SNF family member, and the histone deacetylase HDAC3. Motif analysis

further supported these predictions. Motifs for FOXA1, AR, and MAFA were significantly enriched

in ENZ-repressed enhancers, while FOXA1, HOXB13, and HOXA13 motifs were significantly

enriched in the ENZ-activated enhancers (Figure 2C).

We further mined published ChIP-seq data from LNCaP cells to determine how the binding

of the identified enhancer-associated TF is changing in response to ENZ treatment or androgen

stimulation with dihydrotestosterone (DHT) or the synthetic androgen, R1881) (Figure 2D). Both

ENZ-activated and -repressed enhancers lose AR binding in response to ENZ (AR inhibition) or

the absence of hormones (charcoal-stripped, CS FCS), and binding is regained with androgen

stimulation. Strikingly, FOXA1 binding at the ENZ repressed enhancers is lost with ENZ or CS

FCS treatments and regained with androgen stimulation, while the binding is unaffected at ENZ

activated enhancers. Similar patterns were seen with NKX3.1 and ASH2L with dynamic changes

in binding seen at the ENZ repressed enhancers with trivial changes at the activated ones. In

contrast, both GRHL2 and HOXB13 show minimal binding at the ENZ repressed enhancers as

compared to the activated enhancers irrespective of treatment. Similar patterns were seen

between the activated and repressed groups with ChIP-seq analysis of metastatic patient

samples(Pomerantz et al., 2020) (Figure S1E).

Using a previously described method(Wang et al., 2018), we predicted which genes are

regulated by each ENZ-responsive enhancer. We generated predictions for 690 of the 853

regions with an average of 6.3 genes per candidate enhancer (Figure S1F, Data S6-7). GO

analysis of these gene lists demonstrated that the majority of genes are related to steroid hormone

signaling (in particular androgen response) or the cell cycle (Figure 2E).

Altogether, these findings demonstrate that PRO-cap can detect a large set of precise

changes in enhancer activity that other approaches lack the sensitivity to capture.

Non-coding mutation analysis in PRO-cap-detected ENZ-responsive candidate enhancers

We next queried if PRO-cap-detected enhancer regions could help prioritize somatic genomic

variation in non-coding regions. To that end, using 286 PCa whole genomes available through

ICGC, we searched for single nucleotide variants (SNVs) in the identified 853 ENZ-regulated

enhancers. In total 137 variants were discovered in these regions; 20 of which were recurrent in

more than one patient (Figure S2A). Interestingly, eight of these enhancers also had more than

one SNV within the enhancer region. An example of this is shown in Figure S2B in which two

patients have a recurrent SNV which disrupts an ESRRA motif within the enhancer, and two

different patients have a recurrent SNV which creates an HNF4 motif within the enhancer.

Next, we investigated whether the PRO-cap-identified enhancers may harbor potential

PCa-associated germline single nucleotide polymorphisms (SNPs) (Conti et al., 2021). Of the 269

known PCa risk variants, none overlap with our 853 ENZ-regulated enhancers. However, 10

SNPs did overlap with our larger list of 6,189 total high confidence PCa enhancers ($P < 0.001$). A

representative example of a significantly enriched SNP at a PCa enhancer is shown in Figure

S2C. It is noteworthy that none of these SNPs have been previously identified as residing in gene

regulatory regions. Thus, these results highlight the power of PRO-cap in identifying and

delimiting the non-coding regulatory genome to prioritize enhancer-associated mutations.

***in vivo* functional validation of PRO-cap-detected ENZ-responsive candidate enhancers**

Next, we sought to determine how PRO-cap compares with other methods at measuring enhancer

activity changes. We first surveyed published H3K27ac ChIP-seq and ATAC-seq data from

LNCaP cells treated with ENZ. Many of the regions with altered enhancer activity as measured

by PRO-cap could also be detected by changes in the ChIP- or ATAC-seq data (Figure 4A).

Surprisingly, however, 67% and 54% of the regions did not show a similar change in the ChIP- or

ATAC-seq data, respectively (highlighted in navy and burgundy). Particularly striking was that 78-

79% of the regions with an increase in eRNA transcription with ENZ showed no biochemical

alterations.

223   Given our observations, we sought to validate the enhancer activity of our candidates

identified by PRO-cap using an *in vivo* approach via CRISPR interference (CRISPRi) of these

enhancers. To that end, we genetically targeted two of the previously tested candidates (#1 and

#3) that were repressed by ENZ (Figure 4B). Enhancer candidate #1 is located within in intronic

region of *CACNG4*, while enhancer candidate #3 is found within the intron of *KCNMA1*, (Figures

S3-4). Both of these genes are known to be repressed by ENZ treatment. We designed 3 sgRNAs

each against these regions centered around the TSS of the eRNAs. We then transfected dCas9-

KRAB-stably expressing LNCaP cells with these sgRNAs. CRISPRi-mediated repression of the

two candidate enhancers reduced the expression of both the eRNA and the predicted target

genes (Figure 3C-D).

233   We next sought to determine whether targeting the TSS altered the ability of the candidate

enhancer to regulate gene expression in response to ENZ treatment. To do so we again

transfected dCas9-KRAB-expressing LNCaP cells with the chosen sgRNAs followed by 24-hour

treatment with ENZ. Again, for both genes we demonstrated significant downregulation with ENZ

(*KCNMA1, P* < 0.0001; *CACNG4, P* < 0.0001), with the sgRNA (*KCNMA1, P* < 0.0001; *CACNG4,*

P < 0.0001), and a significant reduction in the ENZ downregulation in combination with the sgRNA

(*KCNMA1, P* = n.s., 1.42 vs 2.05-fold change; *CACNG4, P* = 0.0007, 2.35 vs 4.48-fold change)

(Figure 4A-B).

Next, we functionally validated the ability of PRO-cap to identify changes in enhancer

activity that are not detected by ATAC-seq or H3K27ac ChIP-seq assays (Figure 3A). Given that

the two above candidates both show concomitant changes in both of those biochemical marks

(Figure S3-4), we pursued enhancer candidates which: (1) show significant changes in PRO-cap

upon ENZ treatment, (2) show no changes in ATAC-seq or H3K27ac ChIP-seq after ENZ

treatment, and (3) are predicted to regulate known ENZ-responsive genes.

The first enhancer candidate (#8) selected resides within an intronic region of *DENND1B*

(Figure S5A), a gene that has been shown to be upregulated with ENZ (Figure S5B). While

DENND1B was not significantly upregulated upon ENZ treatment in control sgRNA-expressing

cells, a dramatic increase in its expression was seen in cells expressing sgRNAs against the

candidate target #8 with ENZ treatment (P < 0.0001, 2.06 vs 0.946-fold change) (Figure S5C).

The next enhancer candidate selected (#9) is ~80 kb upstream of *RASD1* (Figure 4C, Figure

S6A), which is a known ENZ-repressed gene (Figure S6B). *RASD1* was significantly

downregulated with ENZ treatment in control sgRNA-expressing cells (P = 0.0084). Likewise, a

significant decrease in RASD1 expression was observed in cells transfected with the candidate

#9 targeting sgRNAs than in control sgRNA-expressing cells (P = 0.0214). and there was a

significant reduction of ENZ downregulation in combination with the sgRNA (P = n.s., 1.15 vs

1.44-fold change) (Figure 5D).

Altogether, these *in vivo* functional results confirm the utility of our PRO-cap assay as a

highly-sensitive approach to identify putative enhancers and detect changes in their activities

genome-wide.

Discussion

This study highlights the generalizable utility of using eRNA transcription patterns obtained from

PRO-cap to detect precise changes in enhancer activity, which has broad applications in human

genetics spanning development and disease. Despite previous efforts in generating chromatin

accessibility and histone modification landscapes to indirectly map transcriptional regulatory

networks, our knowledge of the key regulatory mechanisms that orchestrate the complexity of

cellular differentiation, human development, and disease pathogenesis is still limited by our

incomplete understanding and characterization of the non-coding genome, in particular with the

annotation of cell-state specific transcriptional regulatory elements. Thus, the ability to delineate

more precise maps of enhancer activity dynamics will facilitate the systematic examination of the

transcriptional programs of developing cells across cell transition states and different

differentiation lineages, including disease progression. With a significant fraction of disease-

associated risk variants harbored within non-coding regions of the genome, detailed

characterization of these regulatory networks that coordinate cell- and tissue-type specificity could

provide insights into the molecular mechanisms that underlie dysregulation in numerous disorders

and assist in the mapping of variants functional only at specific cellular states. Similarly, precise

mapping of enhancer dynamics genome-wide in clinical specimens by PRO-cap can also help

better understand disease mechanisms and heterogeneity across patients as well as response

and resistance to treatment.

Hence, our study underscores the clinical value of identifying and delineating aberrant

distal regulatory elements in cancer to identify potential therapeutic vulnerabilities.

Materials and Methods
Cell lines

LNCaP cells (male, ATCC, RRID: CVCL_1379) were maintained in RPMI medium (Gibco,

A1049101), supplemented with 10% FBS (Gibco, 10270106), and 1% penicillin-streptomycin

(Gibco, 11548876) on poly-L-lysine coated plates. All cell lines were grown at 37 °C with 5% CO_2.

All cell lines were authenticated by STR analysis and regularly tested for mycoplasma.

ChIP-seq data analysis

For Figure 1B, previously aligned and normalized bigwig files were downloaded from the

Cistrome Data Browser allowing for consistent and standardized analysis of data from multiple

studies. For other plots previously aligned and normalized bigwig files were downloaded and

analyzed in their published format from GEO. Heatmaps and summary plots were generated

using the deepTools suite(Ramirez et al., 2016).

PRO-cap

LNCaP cells were treated with 10 µM enzalutamide or DMSO for 4 or 24 hrs. For PRO-cap,

approximately 10 to 30 million cells were processed per sample. Library preparations for two

biological replicates each consisting of two technical replicates per condition were processed

separately. Cells were permeabilized and run-on reactions were carried out as previously

described(Mahat et al., 2016). Following RNA isolation, two adaptor ligations using T4 RNA

Ligase 1 (catalog no. M0204; NEB) and reverse transcription using SuperScript III Reverse

Transcriptase (catalog no. 18080044; Invitrogen) were performed, with custom adaptors

detailed in Table S1. Between adaptor ligations, cap state selection reactions were performed

by treating the samples with CIP (catalog no. M0290; NEB) to reduce uncapped RNAs to 5′

hydroxyls and make them incapable of ligating to 5′ adaptor and Cap-Clip (catalog no. C-

CC15011H; Cambio) to remove the 5′ cap of transcripts that had undergone guanylation and

allow them to be incorporated into the library through 5′ adapter ligation. RNA washes,

phenol:chloroform extractions and ethanol precipitations were carried out between reactions.

All steps were performed under RNase-free conditions and following manufacturer protocols.

Libraries were sequenced on an Illumina HiSeq 3000 following PCR amplification and library

clean-up.

ChIA-PET and Hi-C data processing for downstream gene predictions

Similar to a previously described method(Wang et al., 2018), LNCaP RNA Pol II ChIA-PET

interactions(Ramanand et al., 2020) (GSM3423998) were lifted over from GRCh37 to GRCh38.

LNCaP Hi-C data (ENCFF676WJO) was downloaded from ENCODE and interactions were lifted

over from GRCh37 to GRCh38. ChIA-PET and Hi-C interactions were merged and filtered to

remove duplicate interactions. The merged dataset was used as the final 3D interaction set.

Active enhancers and active promoters were inferred based on a merged set of

unidirectional and bidirectional PRO-cap peaks from LNCaP treated with either ENZ or DMSO

for 4 or 24 hrs. PRO-cap peak regions were intersected with GENCODE (V28) annotated

promoters to get a list of active promoter regions. Active enhancers and promoters were

overlapped with the final 3D interaction set to identify potential enhancer-gene pairs. A maximum

of 1 Mb distance was allowed.

Analysis of whole genome patient data

Prostate adenocarcinoma whole genome variants were downloaded from TCGA. This included

SNV data from 286 samples and indel data from 200 samples after filtering and overlapping with

enhancer regions. The effects of these variants were predicted using Funseq2(Fu et al., 2014)

(V2.1.4). The motif break and gain events of the variants were extracted from Funseq2

annotations.

Analysis of PRO-cap data

Differential transcription at eRNA peaks was quantified using EdgeR(Robinson et al., 2010)

analysis of the total read counts in the core promoter (-35 to 60 bp from the peak TSS)(Tippens

et al., 2020) of the plus and minus strands separately. Induced enhancers were defined as FDR

adj p-value < 0.05 in at least one peak direction.

CRISPRi

Lenti-dCas9-KRAB-blast was a gift from Gary Hon (Addgene plasmid # 89567). Lentivirus was

produced in HEK293T cells, and subsequent virus-containing media was used to transduce

LNCaP cells, followed by blasticidin selection.

gRNAs against the candidate enhancer regions were designed using CRISPick.

(https://portals.broadinstitute.org/gppx/crispick/public)(Doench et al., 2016; Sanson et al., 2018)

(Table S1). Custom crRNAs were ordered from IDT and were annealed with Alt-R® CRISPR-

Cas9 tracrRNA (IDT, 1072532) to generate sgRNAs according to the manufacturer's instructions.

For each experiment LNCaP-dCas9-KRAB cells were transfected with 30 nM sgRNA complex

using Lipofectamine RNAiMAX (Invitrogen, 13778030). 48 hours post-transfection cells were

treated with media containing 10 µM enzalutamide or DMSO for a further 24 hours.

qPCR

RNA was extracted directly from cells using the RNeasy Mini Kit (Qiagen, 74106), and genomic

DNA was removed using the DNA-free kit (Ambion, AM1906). RNA was reverse transcribed using

random primers and SuperScript IV Reverse Transcriptase (Invitrogen, 18090010). Quantitative

real-time PCR was performed on the ViiA 7 system (Applied Biosystems) using HOT FIREPol

EvaGreen qPCR mix (Solis Biodyne, 08-24-00020) following the manufacturer's instruction.

Primer sequences are listed in Table S1. All quantitative real-time PCR assays were carried out

using three technical replicates using HMBS as the housekeeping gene.

Data availability
PRO-cap enhancer calls are available in Data S1-7. Raw PRO-cap data and bigWig files are
available through GEO under accession no. GSE198268. Other datasets used in this study are
publicly available under the following accession nos. LNCaP ATAC-seq (accession no.
GSE105116). LNCaP DNase-seq (accession no. GSM816637). LNCaP H3K27ac ChIP-seq
(accession no. GSE107780). LNCaP H3K4me1 ChIP-seq (accession no. GSE73783). LNCaP
RNA Pol II ChIA-PET (accession no. GSM3423998). LNCaP Hi-C (accession no.
ENCFF676WJO). LNCaP AR ChIP-seq, LNCaP FOXA1 ChIP-seq, and LNCaP H3K27ac ChIP-
seq (accession no. GSE85558). ATAC-seq, AR ChIP-seq, FOXA1 ChIP-seq, and H3K27ac
ChIP-seq (accession no. GSE137775) of LNCaP treated with ENZ. ASH2L ChIP-seq (accession
no. GSE60841) of VCaP treated with R1881 or cultured in media containing CS FCS. ChIP-seq
(accession nos. GSE137775, GSE125245, GSE94682, GSE70079, GSE40269, GSE28264, and
GSE80256) of LNCaP treated with ENZ, R1881, DHT, or cultured in media containing CS FCS.
ChIP-seq (accession no. GSE130408) of normal prostate tissue, primary prostate cancer, and
metastatic prostate cancer. LNCaP STARR-seq (GSE151064) of AR binding regions treated with
DHT or EtOH.

Code availability

PRO-cap enhancers were called using Peak Identifier for Nascent Transcript Starts (PINTS)(Yao
et al., 2022). All analysis was performed using common publicly available tools.

Competing interests
The authors declare that there are no competing interests.

375 **References**

Armenia, J., Wankowicz, S.A.M., Liu, D., Gao, J., Kundra, R., Reznik, E., Chatila, W.K., Chakravarty, D., Han, G.C., Coleman, I.*, et al.* (2018). The long tail of oncogenic drivers in prostate cancer. Nat Genet *50*, 645-651.

Arnold, C.D., Gerlach, D., Stelzer, C., Boryn, L.M., Rath, M., and Stark, A. (2013). Genome-wide quantitative enhancer activity maps identified by STARR-seq. Science *339*, 1074-1077.

Chen, H., Li, C., Peng, X., Zhou, Z., Weinstein, J.N., Cancer Genome Atlas Research, N., and Liang, H. (2018). A Pan-Cancer Analysis of Enhancer Expression in Nearly 9000 Patient Samples. Cell *173*, 386-399 e312.

Chen, H., and Liang, H. (2020). A High-Resolution Map of Human Enhancer RNA Loci Characterizes Super-enhancer Activities in Cancer. Cancer Cell *38*, 701-715 e705.

Conti, D.V., Darst, B.F., Moss, L.C., Saunders, E.J., Sheng, X., Chou, A., Schumacher, F.R., Olama, A.A.A., Benlloch, S., Dadaev, T.*, et al.* (2021). Trans-ancestry genome-wide association meta-analysis of prostate cancer identifies new susceptibility loci and informs genetic risk prediction. Nat Genet *53*, 65-75.

Core, L.J., Martins, A.L., Danko, C.G., Waters, C.T., Siepel, A., and Lis, J.T. (2014). Analysis of nascent RNA identifies a unified architecture of initiation regions at mammalian promoters and enhancers. Nat Genet *46*, 1311-1320.

Doench, J.G., Fusi, N., Sullender, M., Hegde, M., Vaimberg, E.W., Donovan, K.F., Smith, I., Tothova, Z., Wilen, C., Orchard, R.*, et al.* (2016). Optimized sgRNA design to maximize activity and minimize off-target effects of CRISPR-Cas9. Nat Biotechnol *34*, 184-191.

Fu, Y., Liu, Z., Lou, S., Bedford, J., Mu, X.J., Yip, K.Y., Khurana, E., and Gerstein, M. (2014). FunSeq2: a framework for prioritizing noncoding regulatory variants in cancer. Genome Biol *15*, 480.

Gasperini, M., Tome, J.M., and Shendure, J. (2020). Towards a comprehensive catalogue of validated and target-linked human enhancers. Nat Rev Genet *21*, 292-310.

Ge, S.X., Jung, D., and Yao, R. (2020). ShinyGO: a graphical gene-set enrichment tool for animals and plants. Bioinformatics *36*, 2628-2629.

Giambartolomei, C., Seo, J.H., Schwarz, T., Freund, M.K., Johnson, R.D., Spisak, S., Baca, S.C., Gusev, A., Mancuso, N., Pasaniuc, B.*, et al.* (2021). H3K27ac HiChIP in prostate cell lines identifies risk genes for prostate cancer susceptibility. Am J Hum Genet *108*, 2284-2300.

Heinlein, C.A., and Chang, C. (2004). Androgen receptor in prostate cancer. Endocr Rev *25*, 276-308.

Henriques, T., Scruggs, B.S., Inouye, M.O., Muse, G.W., Williams, L.H., Burkholder, A.B., Lavender, C.A., Fargo, D.C., and Adelman, K. (2018). Widespread transcriptional pausing and elongation control at enhancers. Genes Dev *32*, 26-41.

Horoszewicz, J.S., Leong, S.S., Kawinski, E., Karr, J.P., Rosenthal, H., Chu, T.M., Mirand, E.A., and Murphy, G.P. (1983). LNCaP model of human prostatic carcinoma. Cancer Res *43*, 1809-1818.

Huang, C.F., Lingadahalli, S., Morova, T., Ozturan, D., Hu, E., Yu, I.P.L., Linder, S., Hoogstraat, M., Stelloo, S., Sar, F.*, et al.* (2021). Functional mapping of androgen receptor enhancer activity. Genome Biol *22*, 149.

Hwang, J.H., Seo, J.H., Beshiri, M.L., Wankowicz, S., Liu, D., Cheung, A., Li, J., Qiu, X., Hong, A.L., Botta, G.*, et al.* (2019). CREB5 Promotes Resistance to Androgen-Receptor Antagonists and Androgen Deprivation in Prostate Cancer. Cell Rep *29*, 2355-2370 e2356.

Kheradpour, P., Ernst, J., Melnikov, A., Rogov, P., Wang, L., Zhang, X., Alston, J., Mikkelsen, T.S., and Kellis, M. (2013). Systematic dissection of regulatory motifs in 2000 predicted human enhancers using a massively parallel reporter assay. Genome Res *23*, 800-811.

Kim, J., Lee, Y., Lu, X., Song, B., Fong, K.W., Cao, Q., Licht, J.D., Zhao, J.C., and Yu, J. (2018). Polycomb- and Methylation-Independent Roles of EZH2 as a Transcription Activator. Cell Rep *25*, 2808-2820 e2804.

Kim, T.K., Hemberg, M., Gray, J.M., Costa, A.M., Bear, D.M., Wu, J., Harmin, D.A., Laptewicz, M., Barbara-Haley, K., Kuersten, S.*, et al.* (2010). Widespread transcription at neuronal activity-regulated enhancers. Nature *465*, 182-187.

Kron, K.J., Murison, A., Zhou, S., Huang, V., Yamaguchi, T.N., Shiah, Y.J., Fraser, M., van der Kwast, T., Boutros, P.C., Bristow, R.G.*, et al.* (2017). TMPRSS2-ERG fusion co-opts master transcription factors and activates NOTCH signaling in primary prostate cancer. Nat Genet *49*, 1336-1345.

Kwasnieski, J.C., Fiore, C., Chaudhari, H.G., and Cohen, B.A. (2014). High-throughput functional testing of ENCODE segmentation predictions. Genome Res *24*, 1595-1602.

Layer, R.M., Pedersen, B.S., DiSera, T., Marth, G.T., Gertz, J., and Quinlan, A.R. (2018). GIGGLE: a search engine for large-scale integrated genome analysis. Nat Methods *15*, 123-126.

Liu, Y., Yu, S., Dhiman, V.K., Brunetti, T., Eckart, H., and White, K.P. (2017). Functional assessment of human enhancer activities using whole-genome STARR-sequencing. Genome Biol *18*, 219.

Mahat, D.B., Kwak, H., Booth, G.T., Jonkers, I.H., Danko, C.G., Patel, R.K., Waters, C.T., Munson, K., Core, L.J., and Lis, J.T. (2016). Base-pair-resolution genome-wide mapping of active RNA polymerases using precision nuclear run-on (PRO-seq). Nat Protoc *11*, 1455-1476.

Malik, R., Khan, A.P., Asangani, I.A., Cieslik, M., Prensner, J.R., Wang, X., Iyer, M.K., Jiang, X., Borkin, D., Escara-Wilke, J.*, et al.* (2015). Targeting the MLL complex in castration-resistant prostate cancer. Nat Med *21*, 344-352.

McNair, C., Xu, K., Mandigo, A.C., Benelli, M., Leiby, B., Rodrigues, D., Lindberg, J., Gronberg, H., Crespo, M., De Laere, B.*, et al.* (2018). Differential impact of RB status on E2F1 reprogramming in human cancer. J Clin Invest *128*, 341-358.

Mei, S., Qin, Q., Wu, Q., Sun, H., Zheng, R., Zang, C., Zhu, M., Wu, J., Shi, X., Taing, L.*, et al.* (2017). Cistrome Data Browser: a data portal for ChIP-Seq and chromatin accessibility data in human and mouse. Nucleic Acids Res *45*, D658-D662.

Meuleman, W., Muratov, A., Rynes, E., Halow, J., Lee, K., Bates, D., Diegel, M., Dunn, D., Neri, F., Teodosiadis, A.*, et al.* (2020). Index and biological spectrum of human DNase I hypersensitive sites. Nature *584*, 244-251.

Palit, S.A., Vis, D., Stelloo, S., Lieftink, C., Prekovic, S., Bekers, E., Hofland, I., Sustic, T., Wolters, L., Beijersbergen, R.*, et al.* (2019). TLE3 loss confers AR inhibitor resistance by facilitating GR-mediated human prostate cancer cell growth. Elife *8*.

Paltoglou, S., Das, R., Townley, S.L., Hickey, T.E., Tarulli, G.A., Coutinho, I., Fernandes, R., Hanson, A.R., Denis, I., Carroll, J.S.*, et al.* (2017). Novel Androgen Receptor Coregulator GRHL2 Exerts Both Oncogenic and Antimetastatic Functions in Prostate Cancer. Cancer Res *77*, 3417-3430.

Pomerantz, M.M., Li, F., Takeda, D.Y., Lenci, R., Chonkar, A., Chabot, M., Cejas, P., Vazquez, F., Cook, J., Shivdasani, R.A.*, et al.* (2015). The androgen receptor cistrome is extensively reprogrammed in human prostate tumorigenesis. Nat Genet *47*, 1346-1351.

Pomerantz, M.M., Qiu, X., Zhu, Y., Takeda, D.Y., Pan, W., Baca, S.C., Gusev, A., Korthauer, K.D., Severson, T.M., Ha, G.*, et al.* (2020). Prostate cancer reactivates developmental epigenomic programs during metastatic progression. Nat Genet *52*, 790-799.

Quigley, D.A., Dang, H.X., Zhao, S.G., Lloyd, P., Aggarwal, R., Alumkal, J.J., Foye, A., Kothari, V., Perry, M.D., Bailey, A.M.*, et al.* (2018). Genomic Hallmarks and Structural Variation in Metastatic Prostate Cancer. Cell *174*, 758-769 e759.

Ramanand, S.G., Chen, Y., Yuan, J., Daescu, K., Lambros, M.B., Houlahan, K.E., Carreira, S., Yuan, W., Baek, G., Sharp, A.*, et al.* (2020). The landscape of RNA polymerase II-associated chromatin interactions in prostate cancer. J Clin Invest *130*, 3987-4005.

Ramirez, F., Ryan, D.P., Gruning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S., Dundar, F., and Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing data analysis. Nucleic Acids Res *44*, W160-165.

Rasool, R.U., Natesan, R., Deng, Q., Aras, S., Lal, P., Sander Effron, S., Mitchell-Velasquez, E., Posimo, J.M., Carskadon, S., Baca, S.C.*, et al.* (2019). CDK7 Inhibition Suppresses Castration-Resistant Prostate Cancer through MED1 Inactivation. Cancer Discov *9*, 1538-1555.

Rhie, S.K., Perez, A.A., Lay, F.D., Schreiner, S., Shi, J., Polin, J., and Farnham, P.J. (2019). A high-resolution 3D epigenomic map reveals insights into the creation of the prostate cancer transcriptome. Nat Commun *10*, 4154.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics *26*, 139-140.

Sanson, K.R., Hanna, R.E., Hegde, M., Donovan, K.F., Strand, C., Sullender, M.E., Vaimberg, E.W., Goodale, A., Root, D.E., Piccioni, F., *et al.* (2018). Optimized libraries for CRISPR-Cas9 genetic screens with multiple modalities. Nat Commun *9*, 5416.

Scher, H.I., and Sawyers, C.L. (2005). Biology of progressive, castration-resistant prostate cancer: directed therapies targeting the androgen-receptor signaling axis. J Clin Oncol *23*, 8253-8261.

Sharma, N.L., Massie, C.E., Ramos-Montoya, A., Zecchini, V., Scott, H.E., Lamb, A.D., MacArthur, S., Stark, R., Warren, A.Y., Mills, I.G., *et al.* (2013). The androgen receptor induces a distinct transcriptional program in castration-resistant prostate cancer in man. Cancer Cell *23*, 35-47.

Shukla, S., Cyrta, J., Murphy, D.A., Walczak, E.G., Ran, L., Agrawal, P., Xie, Y., Chen, Y., Wang, S., Zhan, Y., *et al.* (2017). Aberrant Activation of a Gastrointestinal Transcriptional Circuit in Prostate Cancer Mediates Castration Resistance. Cancer Cell *32*, 792-806 e797.

Stelloo, S., Nevedomskaya, E., Kim, Y., Schuurman, K., Valle-Encinas, E., Lobo, J., Krijgsman, O., Peeper, D.S., Chang, S.L., Feng, F.Y., *et al.* (2018). Integrative epigenetic taxonomy of primary prostate cancer. Nat Commun *9*, 4900.

Taberlay, P.C., Achinger-Kawecka, J., Lun, A.T., Buske, F.A., Sabir, K., Gould, C.M., Zotenko, E., Bert, S.A., Giles, K.A., Bauer, D.C., *et al.* (2016). Three-dimensional disorganization of the cancer genome occurs coincident with long-range genetic and epigenetic alterations. Genome Res *26*, 719-731.

Takeda, D.Y., Spisak, S., Seo, J.H., Bell, C., O'Connor, E., Korthauer, K., Ribli, D., Csabai, I., Solymosi, N., Szallasi, Z., *et al.* (2018). A Somatically Acquired Enhancer of the Androgen Receptor Is a Noncoding Driver in Advanced Prostate Cancer. Cell *174*, 422-432 e413.

Tan, P.Y., Chang, C.W., Chng, K.R., Wansa, K.D., Sung, W.K., and Cheung, E. (2012). Integration of regulatory networks by NKX3-1 promotes androgen-dependent prostate cancer survival. Mol Cell Biol *32*, 399-414.

Tippens, N.D., Liang, J., Leung, A.K., Wierbowski, S.D., Ozer, A., Booth, J.G., Lis, J.T., and Yu, H. (2020). Transcription imparts architecture, function and logic to enhancer units. Nat Genet *52*, 1067-1075.

Tippens, N.D., Vihervaara, A., and Lis, J.T. (2018). Enhancer transcription: what, where, when, and why? Genes Dev *32*, 1-3.

Vanhille, L., Griffon, A., Maqbool, M.A., Zacarias-Cabeza, J., Dao, L.T., Fernandez, N., Ballester, B., Andrau, J.C., and Spicuglia, S. (2015). High-throughput and quantitative assessment of enhancer activity in mammals by CapStarr-seq. Nat Commun *6*, 6905.

Viswanathan, S.R., Ha, G., Hoff, A.M., Wala, J.A., Carrot-Zhang, J., Whelan, C.W., Haradhvala, N.J., Freeman, S.S., Reed, S.C., Rhoades, J., *et al.* (2018). Structural Alterations Driving Castration-Resistant Prostate Cancer Revealed by Linked-Read Genome Sequencing. Cell *174*, 433-447 e419.

Wang, D., Liu, S., Warrell, J., Won, H., Shi, X., Navarro, F.C.P., Clarke, D., Gu, M., Emani, P., Yang, Y.T., *et al.* (2018). Comprehensive functional genomic resource and integrative model for the human brain. Science *362*.

Yao, L., Liang, J., Ozer, A., Leung, A.K., Lis, J.T., and Yu, H. (2022). A comparison of experimental assays and analytical methods for genome-wide identification of active enhancers. Nat Biotechnol.

Zhang, Z., Chng, K.R., Lingadahalli, S., Chen, Z., Liu, M.H., Do, H.H., Cai, S., Rinaldi, N., Poh, H.M., Li, G., *et al.* (2019). An AR-ERG transcriptional signature defined by long-range chromatin interactomes in prostate cancer cells. Genome Res *29*, 223-235.

Figures

Figure 1. PRO-cap sequencing of eRNAs to map active enhancers in prostate cancer. A)

Overview of experimental pipeline implemented in this study. **B)** Enhancer candidates

demonstrate divergent transcription of eRNAs. Shown is the upstream enhancer for androgen

receptor (AR) along with published H3K27Ac HiChIP loops(Giambartolomei et al., 2021) and read

tracks for DNase (GSM816637) and H3K27Ac (GSE85558) ChIP-seq(Meuleman et al., 2020;

Shukla et al., 2017) of LNCaP cells. **C)** Putative enhancers identified by PRO-cap (n = 6,189) are

bound by canonical biochemical marks and known important prostate transcription factors in

LNCaP cells. Heatmaps show read density from publicly available ChIP-seq data(Kim et al., 2018;

McNair et al., 2018; Meuleman et al., 2020; Shukla et al., 2017; Taberlay et al., 2016) (DNase:

GSM816637, AR/FOXA1: GSE85558, ATAC-seq: GSE105116, H3K4me1: GSE73783, and

H3K27Ac: GSE107780) 1kb up- and downstream of the center of the enhancer. **D)** H3K27Ac loop

anchors(Giambartolomei et al., 2021) are significantly more likely ($P < 0.0001$, Kolmogorov-

Smirnov test) to overlap with PRO-cap peaks versus H3K27Ac or DNase I peaks. **E)** Putative

enhancers identified by PRO-cap are also bound by biochemical marks and prostate transcription

factors in patient samples. Heatmaps show average read density from publicly available ChIP-

seq(Pomerantz et al., 2020) data (GSE130408) from normal prostate tissue, primary PCa, and

metastatic PCa (n = 8 samples for each).

A

Plus Strand — DMSO | ENZ 4h | ENZ 24h
Minus Strand — DMSO | ENZ 4h | ENZ 24h
Repressed
Activated
log2 FC

B

ENZ Activated

GIGGLE Score
ASH2L, GRHL2, HOXB13, FOXA1, NKX3-1, AR, GATA2, NR3C1, ARID1A, HDAC3

ENZ Repressed

AR, NR3C1, EP300, SMARCA4, HNF4G, FOXA1

C

ENZ Activated

	FOXA1	1 × 10^49
	HOXB13	1 × 10^20
	HOXA13	1 × 10^15

ENZ Repressed   *P*-val

	FOXA1	1 × 10^38
	AR	1 × 10^30
	MAFA	1 × 10^19

D

AR — Vehicle | ENZ | DHT | CS FCS
FOXA1 — Vehicle | ENZ | R1881 | CS FCS
Repressed / Activated

HOXB13 — CS FCS | R1881
NKX3.1 — CS FCS | DHT
ASH2L — R1881 | CS FCS
GRHL2

E

TNFA SIGNALING VIA NFKB
ANDROGEN RESPONSE
G2M CHECKPOINT
ESTROGEN RESPONSE EARLY
MYC TARGETS V1
E2F TARGETS
P53 PATHWAY
UV RESPONSE UP
DNA REPAIR
PI3K AKT MTOR SIGNALING
FATTY ACID METABOLISM
OXIDATIVE PHOSPHORYLATION
HYPOXIA
PROTEIN SECRETION
UNFOLDED PROTEIN RESPONSE
HEME METABOLISM
XENOBIOTIC METABOLISM
CHOLESTEROL HOMEOSTASIS
APICAL JUNCTION
MYC TARGETS V2
MTORC1 SIGNALING
NOTCH SIGNALING
IL2 STAT5 SIGNALING

ESTROGEN RESPONSE EARLY
G2M CHECKPOINT
APOPTOSIS
ESTROGEN RESPONSE LATE
P53 PATHWAY
MITOTIC SPINDLE
MYC TARGETS V1
TNFA SIGNALING VIA NFKB
GLYCOLYSIS

-log (*P*-value)

Figure 2. PRO-cap sequencing of eRNAs identifies enhancers affected by enzalutamide. A)

PRO-cap analysis of LNCaP cells treated with 10 µM enzalutamide (ENZ) for 24 hours identified

853 putative enhancers which demonstrated significant differential activation or repression on the

plus or minus strands as determined by edgeR analysis (FDR < 0.05). Data is displayed as a

heatmap of the fold-change in mapped reads with 4 or 24 hour ENZ treatments. For each

treatment condition two independent biological replicates are shown. **B)** GIGGLE(Layer et al.,

2018) analysis demonstrates overlap of the ENZ regulated enhancers with regions identified in

published ChIP-seq datasets. Each dot represents a single study, and the GIGGLE score

incorporates both enrichment and significance. **C)** Overrepresented TF motifs as determined by

Cistrome MDSeqPos(Mei et al., 2017) for the ENZ regulated enhancers. **D)** Heatmaps and

summary plots show read density 1kb up- and downstream of the center of the ENZ regulated

enhancers. Data is from publicly available ChIP-seq data(Hwang et al., 2019; Palit et al., 2019;

Paltoglou et al., 2017; Pomerantz et al., 2015; Rasool et al., 2019; Tan et al., 2012) (GSE137775,

GSE125245, GSE94682, GSE70079, GSE40269, GSE28264, and GSE80256) for LNCaP cells

treated with ENZ, R1881, DHT or cultured in media containing charcoal stripped (CS) FCS.

ASH2L data(Malik et al., 2015) (GSE60841) is from VCaP cells treated with R1881 or cultured in

media containing CS FCS. **E)** Gene ontology analysis of the genes predicted to be regulated by

these enhancers generated using ShinyGO(Ge et al., 2020).

Figure 3. Functional validation of PRO-cap enhancers. A) Comparing changes in eRNA

expression with ENZ treatment as detected by PRO-cap with changes in H3K27Ac ChIP-seq and

ATAC-seq. XY charts and heatmaps show read density 1kb up- and downstream of the center of

the ENZ regulated enhancers. Data is from publicly available data(Hwang et al., 2019)

(GSE137775). **B)** Selected candidate enhancers for functional analysis. **C)** CRISPRi targeting

PRO-cap candidate enhancer #3 with three different sgRNAs significantly reduces expression of

the eRNA itself ($P < 0.0001$), and of the downstream target gene *KCNMA1* ($P = 0.0083$). **D)**

CRISPRi targeting PRO-cap candidate enhancer #1 with three different sgRNAs significantly

reduces expression of the eRNA itself ($P = 0.0002$), and of the downstream target gene *CACNG4*

($P = 0.0011$).

A

Mean Normalized Expression

****

****

ns

DMSO ENZ DMSO ENZ

Control sgRNA Repressed #3 sgRNA

B

Mean Normalized Expression

****

****

***

DMSO ENZ DMSO ENZ

Control sgRNA Repressed #1 sgRNA

D

Mean Normalized Expression

*

**

ns

DMSO ENZ DMSO ENZ

Control sgRNA Repressed #9 sgRNA

C

Enhancer #9

DNAse HS

RASD1

17,420 17,440 17,460 17,480 17,500 Kb

chr17

3.2 / -0.3 PRO-cap DMSO rep 1

3.2 / -0.3 PRO-cap DMSO rep 2

3.2 / -0.3 PRO-cap ENZ rep 1

3.2 / -0.3 PRO-cap ENZ rep 2

1.4 DNAse HS

2.5 H3K27Ac DMSO

2.5 H3K27Ac ENZ

1.2 ATAC DMSO

1.2 ATAC ENZ

0.8 AR DMSO

0.8 AR ENZ

1.5 FOXA1 DMSO

1.5 FOXA1 ENZ

Figure 4. CRISPRi interrogation of PRO-cap enhancer candidates. A) CRISPRi targeting

PRO-cap candidate enhancer #3 with three different sgRNAs impairs the ENZ-regulation of the

downstream target gene *KCNMA1* (2.05-fold change, *P* < 0.0001 vs. 1.42-fold change). **B)**

CRISPRi targeting PRO-cap candidate enhancer #1 with three different sgRNAs impairs the ENZ-

regulation of the downstream target gene *CACNG4* (4.48-fold change, *P* < 0.0001 vs. 2.35-fold

change, *P* = 0.0007). **C)** Shown is the candidate enhancer PRO-cap signal along with published

H3K27Ac HiChIP loops(Giambartolomei et al., 2021) and read tracks for DNase (GSM816637),

ATAC-seq, and ChIP-seq with and without ENZ for H3K27Ac, AR, and FOXA1

(GSE137775)(Hwang et al., 2019). **D)** CRISPRi targeting PRO-cap candidate enhancer #9 with

two different sgRNAs impairs the ENZ-regulation of the downstream target gene *RASD1* (1.44-

fold change, *P* = 0.0084 vs. 1.15-fold change).

Figure S1. PRO-cap sequencing of eRNAs identifies enhancers affected by enzalutamide.

A) Venn diagram of the overlap of enhancer regions between the three treatments paradigms **B)**

Heatmap of the PRO-cap signal 250 bp up- and downstream of ENZ activated or repressed peaks

from two replicates of LNCaP cells treated with DMSO or 10 μM ENZ for 24 hours. **C)** PRO-cap

analysis of LNCaP cells treated with 10 μM enzalutamide (ENZ) for 24 hours identified 853

putative enhancers which demonstrated significant differential activation or repression on the plus

or minus strands as determined by edgeR analysis (FDR < 0.05). **D)** PRO-cap analysis of LNCaP

cells treated with 10 μM enzalutamide (ENZ) for 4 hours identified 44 putative enhancers which

demonstrated significant differential activation or repression on the plus or minus strands as

determined by edgeR analysis (FDR < 0.05). **E)** Heatmaps and summary plots show read density

1kb up- and downstream of the center of the ENZ regulated enhancers in metastatic PCa. Data

is from publicly available ChIP-seq data(Pomerantz et al., 2020) (GSE130408) and is displayed

as the mean signal from n = 8 metastatic PCa samples. **F)** Histogram demonstrating the number

of downstream target gene predictions for the 853 ENZ-regulated enhancers identified with PRO-

cap.

Figure S2. PRO-cap helps prioritize candidate enhancer regions to search for cancer-

associated germline and somatic variants. A) Analysis of 286 PCa whole genomes identified

137 single nucleotide variants (SNV) in the ENZ-regulated enhancers identified via PRO-cap, 20

of which were recurrent in more than one patient. **B)** Example of an enhancer downregulated by

ENZ which demonstrates two separate recurrent SNVs, one in two patients that breaks an

ESRRA motif, and another in two different patients that creates an HNF4 motif. **C)** Example of a

PCa risk variant(Conti et al., 2021) which overlaps with an enhancer identified via PRO-cap

A

H3K27Ac HiChIP

DNAse HS

KCNMA1

77,200   77,300   77,400   77,500   77,600 Kb   77,700

chr10

1.1

-6.1   PRO-cap DMSO rep 1

1.1

-6.1   PRO-cap DMSO rep 2

1.1

-6.1   PRO-cap ENZ rep 1

1.1

-6.1   PRO-cap ENZ rep 2

13.8   DNAse HS

7   H3K27Ac DMSO

7   H3K27Ac ENZ

12   AR DMSO

12   AR ENZ

10   FOXA1 DMSO

10   FOXA1 ENZ

sgRNA

B

KCNMA1

Figure S3. Candidate Enhancer #3. A) Shown is the candidate enhancer PRO-cap signal along

with published H3K27Ac HiChIP loops(Giambartolomei et al., 2021) and read tracks for DNase

(GSM816637), and ChIP-seq with and without ENZ for H3K27Ac, AR, and FOXA1

(GSE137775)(Hwang et al., 2019). **B)** Expression of the candidate downstream target gene

KCNMA1 in response to ENZ from multiple published RNA-seq data sets, accession number as

indicated.

A

DNAse HS

CACNG4

66,900 66,920 66,940 66,960 66,980 67,000 67,020 67,040 Kb

chr17

0.9

-0.5 PRO-cap DMSO rep 1

0.9

-0.5 PRO-cap DMSO rep 2

0.9

-0.5 PRO-cap ENZ rep 1

0.9

-0.5 PRO-cap ENZ rep 2

4.7 DNAse HS

4.5 H3K27Ac DMSO

4.5 H3K27Ac ENZ

10 AR DMSO

10 AR ENZ

6.5 FOXA1 DMSO

6.5 FOXA1 ENZ

sgRNA

B

Figure S4. Candidate Enhancer #1. A) Shown is the candidate enhancer PRO-cap signal along

with published H3K27Ac HiChIP loops(Giambartolomei et al., 2021) and read tracks for DNase

(GSM816637), and ChIP-seq with and without ENZ for H3K27Ac, AR, and FOXA1

(GSE137775)(Hwang et al., 2019). **B)** Expression of the candidate downstream target gene

CACNG4 in response to ENZ from multiple published RNA-seq data sets, accession number as

indicated.

A

DNAse HS

DENND1B

197.5   197.6   197.7   197.8   197.9 Mbp
chr1

0.6

-0.5   PRO-cap DMSO rep 1
0.6

-0.5   PRO-cap DMSO rep 2
0.6

-0.5   PRO-cap ENZ rep 1
0.6

-0.5   PRO-cap ENZ rep 2
1.8

DNAse HS
3

H3K27Ac DMSO
3

H3K27Ac ENZ
2.5

ATAC DMSO
2.5

ATAC ENZ
2

AR DMSO
2

AR ENZ
9

FOXA1 DMSO
9

FOXA1 ENZ

B

C

Figure S5. Candidate Enhancer #8. A) Shown is the candidate enhancer PRO-cap signal along

with published H3K27Ac HiChIP loops(Giambartolomei et al., 2021) and read tracks for DNase

(GSM816637), ATAC-seq, and ChIP-seq with and without ENZ for H3K27Ac, AR, and FOXA1

(GSE137775)(Hwang et al., 2019). **B)** Expression of the candidate downstream target gene

DENND1B in response to ENZ from multiple published RNA-seq data sets, accession number as

indicated. **C)** CRISPRi targeting PRO-cap candidate enhancer #8 with three different sgRNAs

impairs the ENZ-regulation of the downstream target gene DENND1B (0.946-fold change vs.

2.06-fold change P < 0.0001).

A

B

RASD1

Figure S6. Candidate Enhancer #9. A) Shown is the candidate enhancer PRO-cap signal along

with published H3K27Ac HiChIP loops(Giambartolomei et al., 2021) and read tracks for DNase

(GSM816637), ATAC-seq, and ChIP-seq with and without ENZ for H3K27Ac, AR, and FOXA1

(GSE137775)(Hwang et al., 2019). **B)** Expression of the candidate downstream target gene

RASD1 in response to ENZ from multiple published RNA-seq data sets, accession number as

indicated.