Disrupting ACE2 Dimerization Mitigates the Infection by SARS-COV-2

Jiaqi Zhu, Yue Su, Young Tang*

Address: Department of Animal Science, Institute of Systems Genetics, University of Connecticut, Storrs, CT 06269, U.S.A.

*Correspondence: yong.tang@uconn.edu

Abstract

The coronavirus disease 2019 (COVID-19) pandemic has caused over 6 million death and 460 million reported cases globally. More effective antiviral medications are needed to curb the continued spread of this disease. The infection by SARS-COV-2 virus is initiated via the interaction between the receptor binding domain (RBD) of the viral glycoprotein Spike (S protein) and the N-term peptidase domain (PD) of the angiotensin-converting enzyme 2 (ACE2) expressed on host cell membrane. ACE2 forms protein homodimer primarily through its ferredoxin-like fold domain (aka. Neck-domain). We investigated whether the dimerization of ACE2 receptor plays a role in SARS-COV-2 virus infection. We report here that the ACE2 receptor dimerization enhances the recognition of SARS-COV-2 S protein. A 43 amino acid peptide based on the N-term of Neck-domain could block the ACE2 dimerization and the interaction between RBD and ACE2, and mitigate the SARS-COV-2/host cell interaction. Our study illustrated a new route to develop potential therapeutics for the prevention and treatment of SARS-COV-2 viral infection.

Introduction

The COVID-19 pandemic causes severe respiratory syndrome and mortality especially in adult and elderly human beings, and has paralyzed the global societal and economic development and interactions. The entry of SARS-COV-2 virus in host cells starts with the recognition of its S protein by the angiotensin-converting enzyme 2 (ACE2) on the cell membrane, which triggers the cleavage of Spike (S protein) into two subunits (S1 and S2, Fig. 1) by host cell proteases(Hoffmann et al., 2020, Shang et al., 2020), with subsequent membrane fusion between virus and host cells driven by the S2 subunit. In addition to the development of vaccines, many efforts have focused on blocking the viral entry by targeting the receptor binding domain (RBD) of S protein or PD of ACE2 with neutralizing antibodies, recombinant or soluble ACE2, peptides based on the N-term peptidase domain (PD) of ACE2 and RBD region of S protein, and small molecules targeting these domains(Datta et al., 2020, Seyedpour et al., 2020, Tay et al., 2020), with additional focuses on development of peptides and protease inhibitors blocking

47 viral fusion and replication, and host cell endosomal and some other
48 pathways(Seyedpour et al., 2020, Datta et al., 2020).
49
The S protein in SARS-COV-2 viral envelop forms homotrimer with the rotation of one of
the three RBDs from "down" to "up" position to be accessible by the PD of ACE2(Wrapp
et al., 2020, Walls et al., 2020). The ectodomain (ECT) region of ACE2 consists of PD
and the ferredoxin-like fold domain (aka. Neck-domain)(Yan et al., 2020). Crystal
structure analysis revealed that ACE2 receptor naturally forms homodimer between
their Neck-domains, with additional "closed" or "open" conformation developed via either
formation or break of another weak interaction in PD region of the protein dimer(Yan et
al., 2020). However, the significance of this dimerization for the ACE2/S protein
interaction is unclear. In this study, we found that the ACE2 dimerization plays a
previously unrecognized role for enhancing SARS-COV-2 infection, while inhibiting this
dimerization mitigated viral infection. Our study will have important impact for the
development of new therapeutics to modify the sensibility of host cells to SARS-COV-2
virus infection.
63
64
65   **Materials and Methods**
66
DNA Constructs, Cell Culture and Virus Production
For BiFC assay, the RBD domain in S protein of SARS-COV-2 virus was inserted into
pBiFC-VN155 (I152L) vector, the ACE2 and its ECT/PD domains were cloned into
pBiFC-VN155 (I152L) and pBiFC-VC155 vector(Kodama and Hu, 2010) (Addgene, MA,
USA). The RBD, Neck-domain and different Neck-domain fragments followed directly
with a stop codon were cloned into pBiFC-VN155 (I152L) vector. For protein expression,
the cell membrane penetrating peptide (TAT), red fluorescence protein (DsRed) and
NK-NT or NKN1 fragments were cloned into pET6xHN-N Vector (Takara, CA, USA).
HEK293T cells were cultured in FP medium (DMEM containing 10% FBS, 2 mM
GlutaMAX™ Supplement, 0.1 mM MEM Non-Essential Amino Acids, 50 U/mL and 50
µg/mL Penicillin-Streptomycin). For ACE2-expressing lentivirus packaging,
pscALPSpuro-HsACE2 (human) (Addgene, MA, USA) were co-transfected with psPAX2
and pCMV-VSV-G packaging plasmids into HEK293T cells using FuGENE 6 (Promega,
WI, USA). Supernatants containing virus were collected at 48h and 72h after
transfection. For doxycycline (Dox) inducible, Spike protein pseudotyped luciferase-
expressing lentivirus preparation, HEK293T cells were transfected with FUW-RLuc-
T2A-PuroR(Kanarek et al., 2018) (Addgene, MA), psPAX2 and pUNO1-SARS2-S
(D614G) (InvivoGen, CA) packaging plasmids using FuGENE 6. Supernatants
containing virus were collected at 48h and 72h after transfection. To establish ACE2-
expressed cell line (ACE2-HEK293T cells), HEK293T cells were infected with ACE2-
expressing lentivirus and ACE2-positive cells were selected by 2 ug/mL of puromycin.
88
Bimolecular Fluorescence Complementation (BiFC) Assay
For BiFC assay, HEK293T cells were cultured in 12-well plates and were co-transfected
with 0.5 µg of each construct expressed in pBiFC-VN155 (I152L) and 0.5 µg pBiFC-
VC155 vectors per well using FuGENE 6. In competition BiFC Assay, HEK293T cells

were co-transfected with 0.5 µg of each construct expressed in pBiFC-VN155 (I152L)
and 0.5 µg pBiFC-VC155 vectors, together with and 5 µg competitor constructs with
stop codon in pBiFC-VN155 (I152L) vector. For the protein competition BiFC Assay,
HEK293T cells were co-transfected with 0.25 µg of each construct expressed in pBiFC-
VN155 (I152L) and 0.5 µg pBiFC-VC155 vectors in 12-well plates. FP medium
containing 0.1 µg, 1 µg or 2 µg of recombinant TAT-Dsred-GFP/NK-NT/NKN1 proteins
were added to the cells at 4 h after plasmid transfection. Fluorescence images were
taken at 24 h and 48 h after transfection using a Nikon fluorescence microscope and
fluorescence intensity was quantified by Image J.
102
Protein Expression and Purification
The recombinant pET6xHN-N constructs were transformed into *Escherichia coli* strain
BL21(DE3) (New England Biolabs, MA, USA). The transformed clones were cultured at
37 ℃ in LB medium with 100 µg/mL Carbenicillin and were induced by adding 1 mM
isopropyl β-D-1-thiogalactopyranoside (IPTG) at OD 0.6-0.8 and incubated at 23 ℃ for
overnight. For protein purification, cells were harvested by centrifugation at 5,000 rpm
for 15 min and resuspended in xTractor Buffer containing DNAse I, Lysozyme solution
and Protease Inhibitor Cocktail (Takara Bio USA, Inc., CA). The suspension was
sonicated 10 sec for 3 times with a 30 s pause, and centrifuged at 10,000 rpm for 20
min. Supernatant was incubated with equilibrated TALON Metal Affinity Resin
(Takara Bio USA, Inc., CA) for 20 min on ice. The proteins were eluted with the Elution
Buffer (pH 7.0, 150 mM Imidazole, 50 mM NaH2PO4, 300 mM NaCl). And the eluted
proteins were concentrated and with buffer exchanged to PBS (pH 7.4) using Protein
Concentrator PES, 3K MWCO (Pierce Biotechnology, PA, USA).
117
Immunoprecipitation and Western Blotting
HEK293T cells in 6-well-plate were transfected with 1 µg Myc-tagged ECT-expressing
plasmids per well, and total proteins were extracted from the cells by adding Cell Lysis
Buffer (Cell Signaling Technology, MA, USA) containing 500 mM NaCl, 1.5% Triton X-
100 and 1 mM PMFS at 48 h after transfection. Supernatant was collected after
centrifugation at 14,000 x g for 10 min and was incubated with 5 µg of recombinant
GFP, NK-NT or NKN-1 proteins, and Myc-Tag (9B11) Mouse mAb (1:1000, Cell
Signaling Technology, MA, USA) at 4 ℃ for overnight. 30 µL of protein G Agarose
Beads (Cell Signaling Technology, MA, USA) slurry was added to the protein mixture
and rotated at 4 ℃ for 3 h. Agarose beads were washed with Cell Lysis Buffer for 5
times and then resuspended in 30 µL of NuPAGE™ LDS Sample Buffer (4X) and 12 µL
NuPAGE™ Sample Reducing Agent (10X) (Invitrogen, MA, USA).
130
Samples collected from Immunoprecipitation were separated by 10% SDS-PAGE gel
and transfected on an Immobilon-P PVDF Membrane. The membrane was blocked by
5% skim milk and incubated with ACE-2 Antibody (1:2,000, Novus Biologicals, CO,
USA), 6xHN Polyclonal Antibody (1:2000, Takara, CA, USA) at 4℃ for overnight. The
membrane was then incubated with secondary HRP-linked, Anti-rabbit IgG (1:10,000,
Cell Signaling Technology, MA, USA) for 1 h at room temperature. For BiFC assay,
whole proteins were extracted from HEK293T cells, separated by 10% SDS-PAGE gel
and transferred onto a Immobilon-P PVDF Membrane. The membrane was incubated

with ACE-2 Antibody (1:2,000, Novus Biologicals, CO, USA), Myc-Tag (9B11) Mouse
mAb (1:1,000, Cell Signaling Technology, MA) and GAPDH (D16H11) XP® Rabbit mAb
(1:1,000, Cell Signaling Technology, MA, USA) at 4℃ overnight. The membrane was
then incubated with secondary HRP-linked, Anti-rabbit IgG (1:10,000, Cell Signaling
Technology, MA, USA) and Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary
Antibody, HRP (1:2,000, Thermo Fisher Scientific, MA, USA) for 1 h at room
temperature. Images were developed with Clarity™ Western ECL Substrate (Bio-Rad
Laboratories, CA, USA) and visualized under the ChemiDox XRS Image System (Bio-
Rad Laboratories, CA, USA).
148
mRNA Transfection and S Protein Pseudovirus Luciferase Assay
For mRNA preparation, GFP, NK-NT and NTN1 DNA fragments harboring T7 promoter
sequence were amplified by PCR, purified, and transcribed to mRNAs *in vitro* using
mMESSAGE mMACHINE™ T7 ULTRA Transcription Kit (Invitrogen, CA, USA). For the
S protein pseudovirus infection and luciferase assay, ACE2-expressing HEK293T cells
were seeded to 24-well plates and transfected with 2 µg of each mRNAs/well using
Lipofectamine™ MessengerMAX™ Transfection Reagent (Invitrogen, CA, USA). After
24 h, cells were infected with Spike-RLuc pseudotyped lentivirus, M2rtTA lentivirus and
10 ug/mL Polybrene for 1 h. The luciferase expression was induced by adding 1 µg/mL
Doxycycline at 24 h after viral infection. Luciferase activity in cells was measured using
Renilla Luciferase Glow Assay Kit (Pierce Biotechnology, PA, USA) and the
CLARIOstar Plus plate reader (BMG LABTECH, NC, USA) at 24 h later.

Statistial Analysis
One way–ANOVA with Tukey's multiple comparison post hoc test or Student's t-test
was used for data analysis. The figures were presented as mean ± standard deviation
(sd). A *p*-value < 0.05 was considered statistically significant.
165
166
167   **Results**
168
Neck-Domain Is Needed for ACE2 Receptor Dimerization in Living Cells.
The ACE2 ECT region consists of PD, the Neck-domain, and a long linker before the
transmembrane helix (TM, excluded from ECT) (Fig. 1). The intensive polar interaction
between Neck-domains is believed to primarily mediate ACE2 dimerization based on
cryo-EM structural analysis(Yan et al., 2020). The bimolecular fluorescence
complementation (BiFC) assay based on reconstitution of the N- and C-term fragments
of the YFP protein Venus (VN and VC) has been widely applied for protein-protein
interaction (PPI) study under physiological conditions(Hu et al., 2002, Kodama and Hu,
2010), allowing direct visualization of PPI in living cells with high sensitivity without
exogenous reagent and complicated data process(Bhat et al., 2006, Miller et al., 2015).
To verify the function of Neck-domain in living cells, we designed BiFC assays to
evaluate the Neck-domain mediated ACE2 dimerization (Figs. 2A, 2B). ACE2 PD
(without Neck-domain) fused with VN or VC were co-expressed in HEK293T cells.
Alternatively, ACE2 ECT (with Neck-domain) fused with VN or VC were co-expressed.
As expected, co-expression of PD-VN and PD-VC yielded no fluorescence, while co-

expression of ECT-VN and ECT-VC produced strong fluorescence (Figs. 2C, 2D). Also,
co-expression of ECT-VN with PD-VC, or ECT-VC with PD-VN did not yield any
fluorescence (Figs. 2C, 2D). These data in living cells support the notion that Neck-
domain is critical for ACE2 protein dimerization(Yan et al., 2020).
188
Neck-Domain Is Needed for the ACE2/S Protein Interaction
To investigate the significance of Neck-domain for the PPI between human ACE2 and
SARS-COV-2 S protein, we developed BiFC assay constructs by fusing S protein RBD
with VN, and ACE2 PD or ECT with VC (Fig. 3A). Western blotting (WB) confirmed co-
expression of S protein RBD-VN and ACE2 PD-VC or ECT-VC fusion proteins in
HEK293T cells (Fig. 3B). The YFP fluorescence was then quantified at 24 h after
plasmids transfection. Intriguingly, the BiFC assay revealed that ACE2 PD (w/o Neck-
domain) is incapable of binding to S protein RBD in living cells, with background
fluorescence signal similar as a pair of previously reported non-interactive
proteins(Huang et al., 2020) - CD163 scavenger receptor cysteine-rich domain 2
(SRCR2) and porcine reproductive and respiratory virus glycoprotein GP2a (Fig. 4). On
the contrary, ACE2 ECT which contains Neck-domain showed strong interaction with S
protein RBD (Fig. 4). These results strongly indicate that the presence of Neck-domain
is necessary for the recognition of viral S protein by ACE2. Based on the fact that Neck-
domain is critical for ACE2 dimerization, our results here support the hypothesis that
Neck-domain is needed for the ACE2 recognition of S protein, most likely through
mediating the dimerization of ACE2.
206
Screening Parts of Neck-Domain Critical for ACE2 Dimerization and S protein
Recognition
To investigate which part of Neck-domain is critical for ACE2 dimerization, we designed
a competition assay by co-expressing different segments of Neck-domain as the
competitor (Fig. 5A) in the ECT-VN/ECT-VC BiFC system (Fig. 2B). The competitor
plasmids were added at 5-fold more than ECT-VN/ECT-VC vectors, with data collected
at 72 h after plasmid transfection to allow the proper expression of ECT-VN/ECT-VC
vectors. As expected, PD without Neck-domain failed to interfere with the interaction
between ECT-VN and ECT-VC (Fig. 5B). The competition BiFC screening revealed that
N-term residues 616-671 of Neck-domain (NK-NT, Fig. 5A) strongly inhibited the PPI
between ECT-VN and ECT-VC (Fig. 5B). Interestingly, overexpression of NK-MT
caused aggregation and death of cells (Fig. 5B). To investigate how the disturbance of
ECT dimerization could impact the PPI between S protein RBD and ECT, we co-
expressed these segments of Neck-domain as competitors in RBD-VN/ECT-VC BiFC
system (Fig. 3A). Over-expression of PD as a competitor could only slightly inhibit the
recognition of RBD by ECT (Fig. 5C). However, NK-NT fragment exhibited the most
significant inhibitory effect on the PPI between RBD and ECT (Figs. 5C, 5D), similar as
in the ECT dimerization BiFC assay (Fig. 5B). Further screening of different parts of NK-
NT identified that residues 616-658 of Neck-domain (NKN1, Fig. 5A) shares similar
potency as NK-NT in blocking the RBD/ECT interaction (Fig. 5E). This correlates well
with the comparable efficiency between NK-NT and NKN1 on the inhibition of full length
ACE2 protein dimerization in BiFC assay (Fig. 5F).
229

In order to double verify the results of the competition BiFC assays, we engineered
histidine and asparagine (HN)-tagged protein constructs with fusion of cell penetrating
peptide TAT(Frankel and Pabo, 1988), red fluorescence protein DsRed, and NK-NT (Fig.
6A, Upper). The TAT-DsRed(Tang et al., 2011) and TAT-DsRed-NK-NT recombinant
proteins were purified and quantified, and added directly to culture medium of 293T
cells transfected with ECT-VN/ECT-VC or ACE2-VN/ACE2-VC plasmids (Fig. 6A,
Lower). Compared with the TAT-DsRed control, one µg/mL and two µg/mL TAT-DsRed-
NK-NT significantly, though not completely inhibited the dimerization between ECT (Fig.
6B) or ACE2 (Fig. 6C). We also expressed HN-tagged DsRed-NKN1, DsRed-NK-NT
fusion proteins (Fig. 6D) and conducted Co-immunoprecipitation (Co-IP) experiment
using total lysates of 293T cells over-expressing Myc-tagged ECT. We found that both
NKN1 and NK-NT recombinant proteins interacted with ECT (Fig. 6E). Taken together,
our results revealed a 43-a.a. peptide (NKN1) as the critical segment in Neck-domain
necessary for the PPI between ACE2/S protein by mediating ACE2 dimerization.
244
NK-NT and NKN1 mRNAs Inhibit SARS-COV-2 Viral Infection
Previous cryl-EM structure study mapped extensive polar interactions for ACE2
dimerization in the second (a.a. 636-658, NKA-2) and fourth (a.a. 708-717, NKA-3)
helices of Neck-domain(Yan et al., 2020) (Fig. 7A), which may explain the partial
activities we observed for NKA-2 and NKA-3 segments in blocking ECT dimerization
(Fig. 5B). However, NK-NT and NKN1 fragments exhibited the greatest potency in
blocking ECT dimerization (Fig. 5B), indicating that in addition to the second helix,
additional residue(s) before it (residues 616-636) also has critical function for ACE2
dimerization (Fig. 7A). One such could be Y633 that forms cation-π interaction with
R710 in the fourth helix(Yan et al., 2020) (Fig. 7A). In order to test whether the
expression of NK-NT and NKN1 fragments can interfere with the SARS-COV-2 viral
infection, we in vitro transcribed and purified the NK-NT and NKN1 mRNAs with the
GFP mRNA as negative control (Fig. 7B). Two µg of mRNA per well were transfected to
293T cells stably expressing human ACE2 in 24-well plates. After 24 h of transfection,
the cells were infected with Renilla luciferase-expressing FUW-lentivirus pseudotyped
with SARS-COV-2 S protein (D614G) as the envelope protein, and the luciferase
activity were evaluated at 48 h after viral infection. We found that compared with the
GFP mRNA, the transfection of both NK-NT and NKN1 mRNAs significantly inhibited S
protein pseudotyped lentivirus infection of the cells (Fig. 7C).
264
265
266   **Discussion**
267
The COVID-19 pandemic has caused nearly 500 million infection cases and over 6
million death globally. Numerous research has been done to better understand the
mechanism of the SARS-COV-2 viral infection and its pathological effect in order to
combat this disease. In this study we tried to understand the fundamental aspects that
determine this virus/host cell interaction to aid in the design of effective therapeutics,
and used uniquely designed serial BiFC assays to verify the essential role of Neck-
domain for the ACE2/S protein PPI. We found that only the ACE2 fragment capable of
self-dimerization can effectively bind to S protein of the SARS-COV-2. This is consistent

with the previous observation from cryl-EM structural study that the interaction with S protein RBD was only observed in ACE2 dimer with "closed" conformation(Yan et al., 2020). These data indicate that conformational changes caused by dimerization could affect the binding affinity of ACE2 to S protein. We further identified that the presence of Neck-domain is critical for the successful ACE2/S protein recognition. This indicates that the disruption of Neck-domain mediated ACE2 dimerization in host cells might have significant impact on S protein recognition.

We further identified the minimal peptide sequence from residues 616-658 (NKN1) in Neck-domain is necessary for blocking ACE2 dimerization and the PPI between ACE2 and S protein. It is interesting that NKA-1 fragment (Fig. 5A) containing both the second and forth helices could not effectively inhibit the PPI between ACE2 or ACE2/S protein. One reason could be the potential self-association of NKA-1 fragments caused by extensive hydrogen bond and Cation-π interaction among residues within these two helices (Fig. 7A). By expressing NKN1 or NK-NT peptides in ACE2-expressing cells, we confirmed that these peptides could inhibit the S protein pseudotyped lentivirus infection. These results correlate well with a previous study showing that artificial dimerization of ACE2 PD by fusion with the mouse IgG Fc has much lower efficiency than that of ACE2-ECT in blocking SARS-COV-2 pseudovirus infection(Li et al., 2020). Our study lets us to predict that in host cell surface, ACE2 receptors form homodimer to enable S protein recognition, and the dimerization mediated by Neck-domain is necessary for ACE2/S protein interaction and the SARS-COV-2 viral infection. Thus, drugs interfering with Neck-domain to disrupt ACE2 homodimerization would have the potential to minimize the susceptibility of host cells to SARS-COV-2 virus.

We therefore expect that future research would identify refined protein peptides based on Neck-domain and small molecules that can effectively block the ACE2 dimerization, to enhance the host cell "immunity" to SARS-COV-2 virus. These could be applied to either protect the host from viral infection or curb subsequent viral amplification and viremia in infected hosts. These drugs could also have extended effect on preventing the infection by other members of the coronavirus family that utilize the recognition of ACE2 receptor to enter the cells.

Data Availability Statement

All data presented in the study are included in the article, further inquiries can be directed to the corresponding author.

Author Contributions

YT and JZ conceived and designed the study and wrote the manuscript. JZ and YS performed the experiments, conducted data analysis, and revision of the manuscript.

322
323 **Acknowledgement**
324
325 This study was supported by the University of Connecticut OVPR COVID-19 Rapid Start
326 Funding.
327

328 **Confliction of Interest**
329
330 The authors declare that they have no conflict of interest.
331
332
333 **Figure Legends:**
334

335 **Fig. 1: Structures of the SARS-COV-2 S Protein and Human ACE2 with Relevant**
336 **Amino Acid Residues.** NTD: N-term Domain. FP: Fusion Peptide. HR: Heptad Repeat.
337 TM: Transmembrane helix.
338

Fig. 2: BiFC Assays to Verify the Neck-domain Function on ACE2 Dimerization.
A) The PD is fused with the N-term or C-term fragment of YFP protein Venus (VN or
VC). B) The ECT is fused with VN or VC. C) Representative fluorescence images of
HEK293T cells co-expressing different combinations of the ACE2-ECT and ACE2-PD
constructs at 24 and 48 h after the plasmid transfection. Bar = 250 μm. D) Quantitation
of the fluorescence intensity at 24 (upper) and 48 h (lower) after plasmid transfection. p
< 0.001, N = 3.
346

347 **Fig. 3: Design of BiFC Assay to Detect the ACE2/S Protein PPI.**
348 A) The RBD of S protein was fused with VN, and PD (a.a. 19-615) or ECT (a.a. 19-740)
349 of ACE2 was fused with VC. B) Western Blot detection of the recombinant proteins co-
350 expressed in HEK293T cells for BiFC assay using specific antibodies. GAPDH served
351 as the loading control.
352

Fig. 4: ACE2 Neck-domain Is Critical for ACE2/S Protein Interaction.
Left: Representative BiFC fluorescence images of HEK293T cells co-transfected with
ACE2-PD-VC or ACE2-ECT-VC and S protein RBD-VN plasmids at 24 and 48 h,
respectively. Bar = 250 μm. Right: Quantitation of the relative fluorescence intensity. p <
0.001, N = 3. NC: negative control using a pair of non-interactive proteins (CD163-
SRCR2 and GP2a) as described in the main text.
359

Fig. 5: Screening Parts of Neck-domain Critical for ACE2 Dimerization and
Recognition of S Protein.
A) Schematic diagram of different Neck-domain fragments for BiFC screening. B)
Competition ECT-VN/ECT-VC BiFC results with the co-expression of different Neck-
domain fragments as competitors. Left: Representative BiFC fluorescence images. Bar
= 250 μm. Right: Quantitation of the relative fluorescence intensity. p <0.01, N = 3. NC:
Negative control of ECT-VN/ECT-VC BiFC with no competitors. C) Competition RBD-
VN/ECT-VC BiFC results with the co-expression of different Neck domain fragments as

competitors. Left: Representative BiFC fluorescence images. Bar = 250 μm. Right:
Quantitation of the relative fluorescence intensity. p <0.001, N = 3. NC: Negative control
of RBD-VN/ECT-VC BiFC with no competitors. D) Competition RBD-VN/ECT-VC BiFC
results with RBD, Neck-domain, and NK-NT as competitors. Left: Representative BiFC
fluorescence images. Bar = 250 μm. Right: Quantitation of the relative fluorescence
intensity. p <0.001, N = 3. E) Comparing competition RBD-VN/ECT-VC BiFC results
with additional fragments of NK-NT as competitors. Left: Representative BiFC
fluorescence images. Bar = 250 μm. Right: Quantitation of the relative fluorescence
intensity. p <0.001, N = 3. F) Comparing competition full length ACE2-VN/ACE2-VC
BiFC results with PD, NK-NT, and NKN1 as competitors. Left: Representative BiFC
fluorescence images. Bar = 250 μm. Right: Quantitation of the relative fluorescence
intensity. p <0.001, N = 3.
380
Fig. 6: Recombinant NK-NT and NKN1 Proteins Bind to ECT and Block ACE2
Dimerization.
A) Upper: Schematic diagram of histidine and asparagine (HN)-tagged recombinant
protein constructs with cell penetrating peptide TAT. Lower: Representative image
showing the HN-TAT-NK-NT recombinant protein transfected into 293T cells. Bar = 120
μm. B) Competition ECT-VN/ECT-VC BiFC results with the TAT-DsRed control or TAT-
DsRed-NK-NT as competitors. Left: Representative BiFC assay fluorescence images.
Bar = 250 μm. Right: Quantitation of the relative fluorescence intensity. p <0.001, N = 3.
C) Competition ACE2-VN/ACE2-VC BiFC assay results with the TAT-DsRed control or
TAT-DsRed-NK-NT as competitors. Left: Representative BiFC fluorescence images.
Bar = 250 μm. Right: Quantitation of the relative fluorescence intensity. p <0.001, N = 3.
D) Schematic diagram of HN-tagged recombinant NK-NT and NKN1 protein constructs.
E) Co-IP results of purified HN-tagged recombinant DsRed, NK-NT and NKN1 proteins
incubated with lysates of the 293T cells expressing Myc-tagged ECT.
395
Fig. 7: NK-NT and NKN1 mRNAs Mitigate SARS-COV-2 Pseudovirus Infection.
A) Residues in NK-NT, NKN1, and Neck-domain critical for ACE2 dimerization. The
second and forth helices are highlighted by yellow and green, respectively. B) Upper:
GFP, NK-NT and NKN1 mRNAs transcribed in vitro via DNA constructs with a T7
promoter. Non-tail: Purified mRNAs transcribed in vitro. Tailing: Purified mRNAs with
addition of poly-A tail. Lower: Expression of GFP mRNA after transfection into 293T
cells. Bar = 250 μm. C) Relative luminescence intensity in 293T cells in 24-well plates
transfected with 2 μg of GFP, NK-NT or NKN1 mRNAs per well and followed by
infection of S protein pseudotyoped lentivirus expressing Renilla luciferase. p<0.01, N =
3.

414
415
416
417 **References**
418
BHAT, R. A., LAHAYE, T. & PANSTRUGA, R. 2006. The visible touch: in planta visualization of
protein-protein interactions by fluorophore-based methods. *Plant Methods,* 2**,** 12.
DATTA, P. K., LIU, F., FISCHER, T., RAPPAPORT, J. & QIN, X. 2020. SARS-CoV-2 pandemic and
research gaps: Understanding SARS-CoV-2 interaction with the ACE2 receptor and
implications for therapy. *Theranostics,* 10**,** 7448-7464.
FRANKEL, A. D. & PABO, C. O. 1988. Cellular uptake of the tat protein from human
immunodeficiency virus. *Cell,* 55**,** 1189-93.
HOFFMANN, M., KLEINE-WEBER, H., SCHROEDER, S., KRUGER, N., HERRLER, T., ERICHSEN, S.,
SCHIERGENS, T. S., HERRLER, G., WU, N. H., NITSCHE, A., MULLER, M. A., DROSTEN, C. &
POHLMANN, S. 2020. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is
Blocked by a Clinically Proven Protease Inhibitor. *Cell,* 181**,** 271-280 e8.
HU, C. D., CHINENOV, Y. & KERPPOLA, T. K. 2002. Visualization of interactions among bZIP and
Rel family proteins in living cells using bimolecular fluorescence complementation. *Mol*
Cell, 9**,** 789-98.
HUANG, C., BERNARD, D., ZHU, J., DASH, R. C., CHU, A., KNUPP, A., HAKEY, A., HADDEN, M. K.,
GARMENDIA, A. & TANG, Y. 2020. Small molecules block the interaction between
porcine reproductive and respiratory syndrome virus and CD163 receptor and the
infection of pig cells. *Virol J,* 17**,** 116.
KANAREK, N., KEYS, H. R., CANTOR, J. R., LEWIS, C. A., CHAN, S. H., KUNCHOK, T., ABU-
REMAILEH, M., FREINKMAN, E., SCHWEITZER, L. D. & SABATINI, D. M. 2018. Histidine
catabolism is a major determinant of methotrexate sensitivity. *Nature,* 559**,** 632-636.
KODAMA, Y. & HU, C. D. 2010. An improved bimolecular fluorescence complementation assay
with a high signal-to-noise ratio. *Biotechniques,* 49**,** 793-805.
LI, Y., WANG, H., TANG, X., FANG, S., MA, D., DU, C., WANG, Y., PAN, H., YAO, W., ZHANG, R.,
ZOU, X., ZHENG, J., XU, L., FARZAN, M. & ZHONG, G. 2020. SARS-CoV-2 and three related
coronaviruses utilize multiple ACE2 orthologs and are potently blocked by an improved
ACE2-Ig. *J Virol.*
MILLER, K. E., KIM, Y., HUH, W. K. & PARK, H. O. 2015. Bimolecular Fluorescence
Complementation (BiFC) Analysis: Advances and Recent Applications for Genome-Wide
Interaction Studies. *J Mol Biol,* 427**,** 2039-2055.
SEYEDPOUR, S., KHODAEI, B., LOGHMAN, A. H., SEYEDPOUR, N., KISOMI, M. F., BALIBEGLOO, M.,
NEZAMABADI, S. S., GHOLAMI, B., SAGHAZADEH, A. & REZAEI, N. 2020. Targeted
therapy strategies against SARS-CoV-2 cell entry mechanisms: A systematic review of in
vitro and in vivo studies. *J Cell Physiol.*
SHANG, J., YE, G., SHI, K., WAN, Y., LUO, C., AIHARA, H., GENG, Q., AUERBACH, A. & LI, F. 2020.
Structural basis of receptor recognition by SARS-CoV-2. *Nature,* 581**,** 221-224.
TANG, Y., LIN, C. J. & TIAN, X. C. 2011. Functionality and transduction condition evaluation of
recombinant Klf4 for improved reprogramming of iPS cells. *Cell Reprogram,* 13**,** 99-112.

TAY, M. Z., POH, C. M., RENIA, L., MACARY, P. A. & NG, L. F. P. 2020. The trinity of COVID-19:
immunity, inflammation and intervention. *Nat Rev Immunol,* 20**,** 363-374.
WALLS, A. C., PARK, Y. J., TORTORICI, M. A., WALL, A., MCGUIRE, A. T. & VEESLER, D. 2020.
Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell,* 181**,**
281-292 e6.
WRAPP, D., WANG, N., CORBETT, K. S., GOLDSMITH, J. A., HSIEH, C. L., ABIONA, O., GRAHAM, B.
S. & MCLELLAN, J. S. 2020. Cryo-EM structure of the 2019-nCoV spike in the prefusion
conformation. *Science,* 367**,** 1260-1263.
YAN, R., ZHANG, Y., LI, Y., XIA, L., GUO, Y. & ZHOU, Q. 2020. Structural basis for the recognition
of SARS-CoV-2 by full-length human ACE2. *Science,* 367**,** 1444-1448.
467

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7