A Leak K[+] Channel TWK-40 Regulates Rhythmic Motor Program in *C. elegans*

Zhongpu Yue [1#], Bin Yu [1#], Yi Li [1#], Yueqing Xu [2#], Lili Chen [1], Jyothsna Chitturi [4], Jun Meng [4], Wesley Hung [4], Yuhang Tian [3], Sonia El Mouridi [5,6], Cuntai Zhang [3], Thomas Boulin [5], Mei Zhen [4], Shangbang Gao [1,2]*

1. Key Laboratory of Molecular Biophysics of the Ministry of Education, College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan 430074, China

2. College of Biomedical Engineering, South-Central University for Nationalities, Wuhan 430074, China

3. Department of Geriatrics, Tongji Hospital of Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

4. Lunenfeld-Tanenbaum Research Institute, Mount Sinai Hospital, University of Toronto, Toronto, Ontario M5G 1X5, Canada

5. Univ Lyon, Université Claude Bernard Lyon 1, MeLiS, CNRS UMR 5284, INSERM U1314, Institut NeuroMyoGène, Lyon 69008, France

6. King Abdullah University of Science and Technology (KAUST), Biological and Environmental Sciences and Engineering Division (BESE), Thuwal 23955–6900, Kingdom of Saudi Arabia

[#] **Equal contributions**

* **Correspondence**: sgao@hust.edu.cn

Running title: TWK-40 regulates the defecation rhythm

Abstract

Leak potassium (K^+) currents, conducted by two-pore domain K^+ (K_{2P}) channels, are critical for the stabilization of the membrane potential. The effect of K_{2P} channels on motor rhythm remains enigmatic. We show here that the K_{2P} TWK-40 regulates the rhythmic defecation motor program (DMP) in *Caenorhabditis elegans*. Disrupting TWK-40 suppresses the expulsion defects of *nlp-40* and *aex-2* mutants. By contrast, a gain-of-function (*gf*) mutant of *twk-40* significantly reduces the expulsion frequency per DMP cycle. *In situ* whole-cell patch clamping demonstrates that TWK-40 forms an outward current that hyperpolarize the resting membrane potential of DVB neuron. In addition, TWK-40 substantially contributes to the rhythmic activity of DVB. Specifically, DVB Ca^{2+} oscillations exhibit obvious defects in *twk-40* mutants. Expression of TWK-40(*gf*) in DVB recapitulates the expulsion deficiency of the *twk-40(gf)* mutant, and inhibits DVB Ca^{2+} oscillations in both wild-type and *twk-40(lf)* animals. Moreover, DVB innervated enteric muscles also exhibit rhythmic Ca^{2+} defects. Taken together, these results demonstrate that TWK-40 is an essential neuronal inhibitor of DMP, thus revealing a cellular mechanism linking leak K_{2P} channels with rhythmic motor activity.

Introduction

Two-pore domain potassium (K_{2P}) channels conduct K^+ leak currents. In contrast to voltage-gated K^+ channels, K_{2P} channels are mostly voltage-independent and non-
45   inactivating channels, which stabilize the cell's resting membrane potential (Enyedi and Czirjak, 2010). Along with most K_{2P} channels profiling by heterologous expression systems or in cultured primary cells, the channel associated behaviors and the intrinsic physiological characterization of specific K_{2P} channel have remained largely unknown.

    The human genome encodes 15 K_{2P} channels, which are grouped into six families
50   based on their functional resemblance and structural similarity (Goldstein et al., 2005; Lesage and Lazdunski, 2000). Aberrant functions of K_{2P} channels have been implicated in various disorders associated with genetic variation. For example, mutations of human TASK-3 (KCNK9 p.Gly236Arg) cause *KCNK9* Imprinting Syndrome, a pediatric neurodevelopmental disease with severe feeding difficulties, delayed development and
55   intellectual disability (Graham et al., 2016). With predicted conservation of amino acid sequences across species, K_{2P} channels associated genetic and functional investigations in animal models are emerging (Buckingham et al., 2005; Lalevee et al., 2006). For instance, cardiac-specific inactivation or overexpression of Ork1, a *Drosophila* two-pore domain potassium channel, led to an increase or a complete arrest of fly heart beating,
60   respectively (Lalevee et al., 2006).

    The nematode *Caenorhabditis elegans* (or *C. elegans*) genome contains a large family with at least 47 K_{2P} genes (Bargmann, 1998; Buckingham *et al.*, 2005). This large set of K_{2P} channels may allow for exceptionally fine 'tuning' of the firing activity of individual cells within the very compact nematode nervous system (Salkoff et al., 2001;

65   Witvliet et al., 2021). Encouraged by the increasing understanding of the molecular and

cellular wiring of the worm's neural network (Bargmann, 1998; Cook et al., 2019; White

et al., 1986; Witvliet *et al.*, 2021), we use *C. elegans* to explore the functional complexity

of K_{2P} channels as potential regulators of rhythmic motor behaviors *in vivo* (Ben Soussia

et al., 2019; Wang et al., 1999).

70   The *C. elegans* defecation motor program (DMP) exhibits a highly coordinated

ultradian rhythm. It is achieved by periodically activating a stereotyped sequence of

muscle contractions, including the initial contraction of posterior body wall muscles

(pBoc), followed by the second contraction of anterior body wall muscles (aBoc), and the

final contraction of enteric muscles (EMC) (Thomas, 1990). Driven by the synchronizing

75   activity of electrically-coupled DVB and AVL motor neurons (Choi et al., 2021), EMC

leads to the robust expulsion (Exp) of gut contents (**Figure 1A**) (Dal Santo et al., 1999;

Liu and Thomas, 1994). Unlike most rhythmic motor circuits that are composed of a

central pattern generator (CPG) neural network or self-oscillating pacemaker neurons

(Marder and Calabrese, 1996), the expulsion rhythm is precisely timed by endogenous

80   calcium oscillation signals from the intestine (Branicky and Hekimi, 2006; McIntire et

al., 1993a). The intestinal IP_3 receptor/*itr-1*-driven pacemaker Ca^{2+} oscillations trigger a

Ca^{2+}-dependent secretion of neuropeptide NLP-40, which activates the neuronal G

protein-coupled receptor (GPCR) AEX-2 in DVB and AVL (Mahoney et al., 2008;

Nehrke et al., 2008; Teramoto and Iwasaki, 2006; Wang et al., 2013). GABA release

85   from DVB (McIntire et al., 1993b), cyclically initiated by NLP-40 (Wang *et al.*, 2013),

activates the muscular excitatory GABA-gated cation channel EXP-1 to drive the final

contraction of enteric muscles (**Figure 1B**) (Beg and Jorgensen, 2003). Deficiency of

either NLP-40, AEX-2 or EXP-1 results in serious expulsion defects in *C. elegans*. Although the intestinal Ca^{2+} oscillation sets the instructive timing signal of the expulsion

90 rhythm (Branicky and Hekimi, 2006; McIntire *et al.*, 1993a), the cellular processes that control the expulsion rhythm in the intestine-neuron-muscle circuit are still unclear.

We show here that loss of function of *twk-40* – a two-pore domain potassium channel subunit – specifically suppresses the expulsion defects of *nlp-40* and *aex-2* mutants, but not of *exp-1* mutant. TWK-40 exhibits the common structural features of K_{2P}

95 channel based on sequence analyses. The single subunit of TWK-40 has four transmembrane regions and two pore-forming domains (**Figure 2A**), which exhibit sequence similarity with the mammalian TASK (tandem-pore-acid-sensing K^+) subfamily (**Supplementary Figure 1A**). A gain of function mutant of *twk-40(bln336)*, generated by CRISPR/Cas9 gene editing (Ben Soussia *et al.*, 2019), induces an expulsion defect by

100 reducing the frequency of expulsion per DMP cycle. Moreover, *in-situ* whole-cell patch clamping revealed a *twk-40*-dependent outward K^+ current that regulates the resting membrane potential (RMP) in DVB neurons, in which the RMP was depolarized in *twk-40(lf)* and hyperpolarized in *twk-40(gf)* mutant, respectively. In combination with real-time Ca^{2+} imaging, we provide several lines of evidence to demonstrate that TWK-40

105 regulates rhythmic motor behaviors – including defecation – by cell-autonomously regulating DVB's resting membrane potential.

Remarkably, we also found recently that TWK-40 modulates global locomotor activity by regulating the activity of the AVA premotor interneuron (Meng et al., 2022). Briefly, *twk-40(lf)* mutant animals exhibit exaggerated body bends during both forward

110 and backward movement, and more frequent and prolonged reversals. *twk-40(gf)* mutant

animals, by contrast, exhibit strongly altered locomotion including obvious decreased body bending and velocity in both forward and backward undulation. Heterologous expression of the TWK-40 in HEK293T cells produced a voltage-insensitive K^+ selective K_{2P}-like ion channel. Therefore, the K_{2P} channel TWK-40 is required for the generation

115 of multiple motor rhythms, including the defecation and locomotion motor programs.

Results

twk-40(lf) suppresses the expulsion defect of nlp-40 and aex-2 mutants

To determine a potential role of the TWK-40 K_{2P} channel in the cellular mechanisms controlling rhythmic DMP activity, we took advantage of three *exp* mutants that disrupt

120 DMP in different tissues within the expulsion motor circuit (**Figure 1B**). First, *nlp-40(tm4085)*, an instructive neuropeptide from the intestinal pacemaker that delivers temporal information to the DVB neuron (Wang *et al.*, 2013). Second, *aex-2(sa3)*, a G-protein-coupled receptor that functions as the DVB receptor for NLP-40 (Mahoney *et al.*, 2008; Wang *et al.*, 2013). And the third, *exp-1(sa6)*, which encodes an excitatory GABA

125 receptor expressed in enteric muscles (Beg and Jorgensen, 2003; Lee et al., 2005). Loss-of-function mutants of these genes induced severe expulsion defects (**Figure 1C, E, G**) by disrupting the normal function of the intestine, DVB neuron and enteric muscles, respectively.

We reasoned that a loss of the potassium conductance *twk-40* would increase the

130 excitability of cells within the circuit and compensate the inhibitory effect of the *nlp-40*, *aex-2*, or *exp-1* mutations. Hence, we combined the *twk-40* allele *hp834* (**Figure 2A**) (Meng *et al.*, 2022) with each DMP mutation. *hp834* contains a 7 bp deletion of the sequence (GTTCGAG) at the base of 127-133 exon before the first pore-forming domain

of T28A8.1a (**Figure 2A**), causing a frameshift and subsequent premature stop codon of

135   *twk-40a* (Meng *et al.*, 2022), one of three annotated isoforms of *twk-40*

(www.wormbase.org). We found that *twk-40(hp834)* significantly improved the

expulsion defect in *nlp-40* and *aex-2* mutants (**Figure 1C-F**). Specifically, the expulsion

frequency per DMP was increased in *nlp-40(tm4085); twk-40(hp834)* (0.18±0.02 Exp per

cycle) or *aex-2(sa3); twk-40(hp834)* (0.15±0.02 Exp per cycle), when compared to the

140   single mutants of *nlp-40(tm4085)* (0.08±0.01 Exp per cycle) and *aex-2(sa3)* (0.08±0.01

Exp per cycle), respectively (**Figure 1C-F**). Similar expulsion defect suppression was

observed in another *twk-40(lf)* allele *hp733* (**Supplementary Figure 1B**), which harbors

a missense mutation that changes an amino acid at the end of the first transmembrane

domain (E58K) (Meng *et al.*, 2022). By contrast, no substantial suppression of the

145   expulsion defect of *exp-1* mutants was observed for either loss-of-function allele of *twk-*

40. Indeed, no difference in expulsion frequency was observed between *exp-1(sa6)*

(0.17±0.01 Exp per cycle) and *exp-1(sa6); twk-40(hp834)* (0.17±0.01 Exp per cycle) or

exp-1(sa6); twk-40(hp733) (0.16±0.01 Exp per cycle) double mutants, respectively

(**Figure 1G, H, Supplementary Figure 1B**). These results demonstrated that *twk-40* is

150   selectively required for rhythmic defecation regulation, upstream of *exp-1*.

A twk-40 gain-of-function mutation reduces expulsion frequency

       The selective suppression of *exp* defects in distinct *exp* mutants suggested that *twk-*

40, per se, may regulate the DMP. Indeed, in *twk-40(hp834, lf)* single mutants, the

expulsion number was modestly but significantly increased (13.87±0.21/10 min)

155   compared to wild-type animals (12.97±0.19/10 min) (**Figure 2C, D**). Similar results were

also observed in *twk-40(hp733)* mutants. These results confirm that *twk-40* regulates the DMP negatively.

To further confirm the involvement of *twk-40* in the defecation motor program, we examined the expulsion step in a gain-of-function mutant allele of *twk-40, bln336* (**Figure 2A**). These mutants harbor a pan-K_{2P} activating mutation (Ben Soussia *et al.*, 2019) that promotes the gating of vertebrate and invertebrate K_{2P} channels. Indeed, heterologous expression of the corresponding TWK-40(L159N) channel in HEK293T cells showed ~5 fold current increase from the wild-type TWK-40 (Meng *et al.*, 2022). Consistent with an inhibitory function of *twk-40*, we observed a strongly reduced expulsion frequency (4.8±0.25/10 min) in *twk-40(bln336, L159N)* animals, i.e. approximately 25% of wild type (12.97±0.19/10 min) (**Figure 2C, D**). Thus, gain-of-function of *twk-40* inhibits the expulsion rhythm, which encouraged us to further investigate the functional effects of *twk-40* loss- and gain-of-function mutants.

twk-40 regulates expulsion in the DVB neuron

To pinpoint the exact role of *twk-40* in the DMP, we first assessed its cellular focus of action. Loss of *twk-40* suppressed the *exp* defect of *nlp-40* and *aex-2* mutants but not of *exp-1*, indicating that *twk-40* regulates the expulsion frequency from the intestine and/or expulsion neurons. Indeed, functional TWK-40 driven by a small fragment of its upstream region (P*twk-40*::TWK-40::sl2dGFP, Methods) revealed the expression exclusively in the nervous system and intestine (**Figure 2B**). The neuronal expression pattern includes the excitatory ventral motor neurons (A- and B-types), but no expression was observed in inhibitory D-motor neurons. Meanwhile, sparse labeling of head and tail

neurons was also observed, including the DVC interneuron (strong) and the excitatory GABAergic DVB motor neuron/interneuron (moderate) (**Figure 2B**).

180   We then systematically tested the tissue or cell requirements of *twk-40* for the expulsion behavior. A wild-type TWK-40 cDNA driven by different promoters was competitively expressed in *twk-40(bln336, gf)* mutant (0.26±0.02 Exp per cycle), in an attempt to reduce its severe *exp* defect. Expression of wild-type TWK-40 (i.e., TWK-40(WT)) by the short promoter P*twk-40* (Methods) rescued the animal's expulsion

185   frequency to (0.55±0.04 Exp per cycle) (**Figure 2E**). Expression of TWK-40(WT) in GABAergic DVB/AVL/D-motor neurons (P*unc-47*), also significantly restored the DMP (0.67±0.06 Exp per cycle). More importantly, when we restored TWK-40(WT) expression exclusively in the DVB neuron (P*flp-10*)(Choi *et al.*, 2021), the expulsion defect of *twk-40(bln336, gf)* mutant was rescued (0.7±0.06 Exp per cycle). In contrast,

190   expression of the TWK-40(WT) in the intestine (P*ges-1*, 33.2±1.9%) (**Supplementary Figure S1C**) or D-motor neurons (P*unc-25*s, 33.8±5.6%) (Jin et al., 1999), however, did not restore a normal expulsion frequency (**Figure 2E**). Therefore, consistent with the expression pattern, these results suggest that regulation of the expulsion rhythm by *twk-40* requires DVB neurons.

195   To further confirm this notion, we ectopically expressed the TWK-40(L159N) gain-of-function mutant in DVB neurons. Indeed, in the wild-type N2 background, TWK-40(L159N) expression in DVB (P*twk-40*, P*unc-47* or P*flp-10*) resembled the expulsion defect as found in *twk-40(bln336)* mutant animals, exhibiting strongly reduced expulsion frequency per cycle (**Figure 2F**). TWK-40(L159N) expression only in D-MNs (P*unc-*

200   25s), however, could not replicate this defect. Similar phenotypes were also observed by

expressing TWK-40(L159N) in a *twk-40(lf)* mutant context (**Figure 2G**).

Taken together, these results demonstrate that *twk-40* is sufficient and necessary in

DVB neurons for the regulation of the expulsion rhythm.

Rhythmic Ca^{2+} oscillation of the DVB neuron is disrupted in twk-40 mutants

205   To determine whether *twk-40* directly regulates DVB's neuronal activity, we

performed Ca^{2+} imaging in animals expressing the genetically encoded Ca^{2+} sensor

GCaMP6s in the DVB neuron (**Figure 3A**) (Li et al., 2023). We conducted the Ca^{2+}

imaging in a liquid environment on restricted animals, in which DVB exhibited tight

correlation of rhythmic Ca^{2+} oscillations with cyclic expulsion behavior (**Figure 3B-D**).

210   In wild-type animals, DVB fired periodic Ca^{2+} spikes (or Ca^{2+} oscillations) with a

frequency of 4.5±0.3 Hz/300s (**Figure 3B, E, F**). To verify these Ca^{2+} spikes represent

pacemaker-activated DVB activation, we examined their dependence on NLP-40, the

instructive neuropeptide from the intestinal pacemaker that delivers temporal information

to the DVB neuron (Wang *et al.*, 2013). Consistent with previous reports (Choi *et al.*,

215   2021; Wang *et al.*, 2013), we observed severe decrease in both DVB Ca^{2+} and expulsion

frequency in the *nlp-40(tm4085)* mutant (**Supplementary Figure S2A-C**). These results

demonstrate that the activation dynamics of DVB are indeed dependent on the NLP-40-

initiated pacemaker signal.

In *twk-40(bln336, gf)* mutants, the frequency of Ca^{2+} spikes was significantly

220   reduced to 3.2±0.3 Hz/300s (**Figure 3E, F**). Interestingly, the Ca^{2+} amplitude was also

diminished in *twk-40(bln336, gf)* mutants (**Figure 3G**). Conversely, in *twk-40(hp834, lf)*

mutants, the Ca^{2+} frequency was significantly increased to 5.9±0.5 Hz/300s. Yet, the Ca^{2+}

amplitude of *twk-40(hp834)* was unchanged (**Figure 3G**), and the individual Ca^{2+} spike

kinetics – including the rise and decay time – were not modified either (**Supplementary**

Figure S3A-C), suggesting that *twk-40* contributes to the basal-activity of the DVB

neuron. Taken together, our experiments demonstrate that the intrinsic DVB Ca^{2+}

oscillation activity was substantially regulated by *twk-40*.

To further decipher *twk-40*'s cellular specificity, we firstly measured neuronal Ca^{2+}

activity by expressing wild-type TWK-40 cDNA under different exogenous promoters in

a *twk-40(bln336, gf)* mutant background. Consistent with our behavioral data, the Ca^{2+}

frequency and amplitude were partially rescued when TWK-40(WT) was expressed in

DVB in *twk-40(bln336, gf)* mutants (**Figure 4A-C**). By contrast, neither frequency nor

amplitude was restored when TWK-40(WT) was expressed in D-motor neurons (**Figure**

4C). In addition, we then expressed TWK-40(WT) in *twk-40(hp834, lf)* mutant and found

that the increased Ca^{2+} frequency of these mutants could also be rescued to wild-type

levels when TWK-40(WT) was expressed in DVB (**Figure 4D-F**). Thus, TWK-40(WT)

is sufficient to restore the neuronal Ca^{2+} activity of the DVB neurons.

Collectively, these results support the notion that *twk-40* regulates neuronal Ca^{2+}

oscillations of the DVB neuron.

twk-40 cell-autonomously regulates DVB Ca²⁺ oscillations

The broad neuronal expression pattern of *twk-40* indicates that TWK-40 may

regulate the DMP not limited in the DVB neuron (White *et al.*, 1986). To examine

whether synaptic input upstream of DVB may play a role, we examined the DVB Ca^{2+}

oscillations in an *unc-13(lf)* background (Wang *et al.*, 2013). UNC-13 is a conserved and

essential presynaptic Ca^{2+} effector that triggers exocytosis and neurotransmitter release

(Brose et al., 1995; Gao and Zhen, 2011; Richmond and Jorgensen, 1999). Loss of

function of *unc-13* strongly impairs global neurotransmission with minor effect on NLP-

40 release (Wang *et al.*, 2013). Interestingly, in the *unc-13(lf)* mutant background, we

found that Ca^{2+} oscillation defects were retained in *twk-40* mutants, including the

250   decreased frequency and amplitude in *twk-40(bln336, gf)* mutants and increased

frequency in *twk-40(hp834, lf)* mutants (**Supplementary Figure S4A-C**). These results

demonstrate that potential presynaptic inputs to DVB don't interfere with the regulation

of DVB's Ca^{2+} activities by TWK-40, and that *twk-40* most likely regulates the cellular

excitability of DVB neuron in a cell-autonomous manner.

255   To reinforce this notion, we further tested TWK-40's inhibition of DVB by

measuring the Ca^{2+} oscillation in different transgenic lines that expressed the *twk-

40(L159N, gf)* cDNA. When TWK-40(L159N) was expressed in DVB in wild-type

animals (P*twk-40*, P*unc-47* or P*flp-10*), the frequency and amplitude of Ca^{2+} spikes were

significantly reduced (**Supplementary Figure S5A**), similar to *twk-40(bln336, gf)*

260   mutants (**Supplementary Figure S5B, C**). By contrast, expression of TWK-40(L159N)

in D-motor neurons had no effect. Furthermore, reduced Ca^{2+} oscillations were also

observed when TWK-40(L159N) was expressed in a *twk-40(hp834, lf)* mutant

background (**Supplementary Figure S5E-F**). These results reveal that ectopic

expression of TWK-40(L159N) is sufficient to silence the Ca^{2+} oscillation of DVB

265   neurons. Namely, TWK-40 execute a dominant inhibition of the intrinsic activity

regulation in DVB neuron.

DVB activity regulates Ca^{2+} dynamics in enteric muscles

The expulsion behavior ultimately relies on the coordinated contraction of a group of muscles, including the anal depressor and sphincter, and two enteric muscles (EM),

270 which wrap around the posterior gut to further pressurize the intestinal contents (McIntire *et al.*, 1993b). EMs are innervated by DVB (**Figure 5A**) and AVL neurons (McIntire *et al.*, 1993b; White *et al.*, 1986), and the contraction of EMs is driven by intracellular Ca^{2+} transients (LeBoeuf and Garcia, 2017). We then ask whether EM Ca^{2+} activity is also regulated in *twk-40* mutants.

275 We thus performed Ca^{2+} imaging of animals expressing the calcium indicator GCaMP6s in the enteric muscles (EM::GCaMP6s, **Figure 5B**) (Li *et al.*, 2023). Similar to activation dynamics in DVB neuron, enteric muscles displayed rhythmic Ca^{2+} spikes closely correlated with expulsion action (**Figure 5C, D**). In our experimental conditions, the EM Ca^{2+} spikes exhibited a frequency of 6.4±0.5 Hz/300s (n = 20) in wild-type

280 animals (**Figure 5E**), which was also eliminated in *nlp-40(tm4085)* mutant (**Supplementary Figure S2D-E**). In *twk-40* mutants, the periodicity of EM Ca^{2+} spikes was affected. Specifically, the EM Ca^{2+} frequency was significantly reduced in *twk-40(bln336, gf)* mutants (3.0±0.2 Hz/300s, n = 20), and increased to 8.6±0.6 Hz/300s (n = 20) in *twk-40(hp834, lf)* mutants (**Figure 5E, F**), reminiscent of *twk-40*'s effects for DVB

285 Ca^{2+} frequency. EM Ca^{2+} amplitudes, however, were unchanged in *twk-40(bln336, gf)* animals (**Figure 5G**), which differs from the reduction observed for DVB Ca^{2+} spike amplitudes. Given that no *twk-40* expression was observed in enteric muscles, we propose that the effects of *twk-40* mutations on the Ca^{2+} activity of enteric muscles are a consequence of the dysregulation of presynaptic DVB neurons.

290   To confirm this idea, we performed additional experiments by expressing wild-type

TWK-40 in EM and DVB neurons, respectively. We found that the reduced EM Ca^{2+}

frequency in *twk-40(bln336, gf)* mutant was almost fully rescued by expressing TWK-

40(WT) in DVB (P*twk-40*, P*unc-47* and P*flp-10*), but not in enteric muscles (P*exp-1*) or

in D-motor neurons (P*unc-25s*) (**Figure 6A, B**). Consistently, the increased frequency of

295   EM Ca^{2+} spikes in *twk-40(hp834, lf)* mutants was also restored to wild-type by TWK-

40(WT) expression in DVB neurons (**Figure 6C, D**). We also ectopically expressed the

gain-of-function TWK-40(L159N) in *twk-40(hp834)* animals to inhibit DVB's activity

and recorded the EM Ca^{2+} spikes. Consistently, the frequency of EM Ca^{2+} spikes was

significantly diminished in this experiment. This inhibition was however not observed in

300   animals expressing TWK-40(L159N) in the enteric muscles or D-motor neurons (**Figure**

6E, F). Collectively, our experiments argue that *twk-40* regulates Ca^{2+} activities of enteric

muscles by modulating DVB activity.

TWK-40 hyperpolarizes the resting membrane potential of DVB neuron

        The functional inhibition of DVB neuron by *twk-40* is consistent with an inhibitory

305   function of this potassium channel. To directly investigate the functional effect of *twk-40*,

we dissected and recorded GFP-labeled DVB neurons *in vivo* in the whole-cell patch

clamp configuration (Methods, **Figure 7A**). In wild-type animals, stable outward K^+

currents were recorded at stepwise holding voltages from -60 mV to +80 mV (Jiang et al.,

2022; Xie et al., 2013). These currents were significantly diminished in *twk-40(hp834)*

310   loss-of-function mutants (**Figure 7B, C**). Conversely, in *twk-40(bln336)* gain-of-function

mutants, outward currents were dramatically increased (**Figure 7B, C**). Thus, these *in-*

situ neuronal recordings demonstrate that *twk-40* supports a functional K^+ current in DVB neuron.

Furthermore, the resting membrane potential of DVB was also analyzed. While wild-type DVB neurons had an RMP of -38.9±3.7 mV (n = 6), loss of *twk-40* depolarized the RMP to -22.1±2.9 mV (n = 7). Conversely, in *twk-40(bln336)* mutants, the RMP of DVB neuron was dramatically hyperpolarized to -63.1±5.8 mV (n = 6) (**Figure 7D, E**). Taken together, these results demonstrate that TWK-40 constitutes a K^+ channel that stabilizes the RMP of DVB neuron.

In summary, we show here that TWK-40 forms a *bona fide* potassium channel that cell-autonomously maintains the RMP of DVB neuron, and that *twk-40* is essential for the regulation of the rhythmic expulsion motor program (**Figure 7F**).

Discussion

Potassium currents mediated by K_{2P} channels are important modulators of neuronal activity in animal nervous systems. We show here that the previously uncharacterized K_{2P} channel TWK-40 provides a prominent leak potassium current in the DVB motor neuron that regulates the rhythmic expulsion behaviour of *C. elegans*. We present genetic evidence that loss of *twk-40* suppresses the *exp* defects of *nlp-40* and *aex-2* mutants. Additional evidence from expression pattern analysis, behavioral rescue experiments, and real-time neuronal and muscular Ca^{2+} imaging, reveal the substantial contribution of TWK-40 to rhythmic expulsion activity. Further proof collected by heterologous expression of TWK-40 (Meng *et al.*, 2022) and *in situ* whole-cell patch clamp recording, demonstrates that *twk-40* constitutes a K_{2P}-like potassium-selective current. Our results thus identify a K_{2P} channel TWK-40, through its influence on DVB neuron resting

335   membrane potential, plays a critical role in coordinating the expulsion rhythm and

locomotion, shedding light on the intricate interplay between potassium channels and

neural circuits in controlling complex motor programs in vivo. These findings suggest a

broader significance of TWK-40/K$_{2P}$ in orchestrating multiple motor rhythms, providing

insights into the molecular mechanisms underlying coordinated motor behaviors.

340   Circadian rhythms with 24-hour-long cycles like sleeping are well studied and are

controlled by the expression fluctuation of clock genes, like *Per1*, *Per2*,and *Bmal1 etc*.

(Lowrey and Takahashi, 2004). Ultradian rhythms (or episodic ultradian events) with

diverse shorter periods ranging from milliseconds to hours, are ubiquitous and

indispensable in many neuromuscular systems, such as heartbeat, respiration and

345   peristalsis (Goh et al., 2019). These ultradian rhythms are generally considered to be

generated by motor central pattern generators (CPG) (Marder and Bucher, 2001). The

ultradian DMP rhythm in *C. elegans* however is a remarkable exception, as it is initiated

by calcium oscillations in intestinal cells (Teramoto and Iwasaki, 2006). The DVB and

AVL neurons, acting as a signaling bridge through NLP-40 and GABA pathways,

350   transfer non-neuronal calcium oscillation from the intestine to enteric muscles to execute

the expulsion action.

Genetic analysis initially highlighted critical ion channel genes for DVB activity,

such as *unc-2* and *egl-19* that encode the α1 subunits of P/Q- and L- type voltage-gated

Ca^{2+} channel, respectively (Wang and Sieburth, 2013), and *egl-36*, a Shaw-type (Kv3)

355   voltage-dependent potassium channel subunit (Johnstone et al., 1997). However, the *in*

vivo cellular mechanism of how these channels contribute to the behavior is still unknown.

The recent descriptions of synchronized giant action potentials between AVL and DVB,

and of unusual compound action potentials in AVL, provide insights into how the

rhythmic expulsion behaviour is generated at the cellular level. In particular, the negative

360   potassium spikes in AVL are mediated by a repolarization-activated potassium channel

EXP-2 (Jiang *et al.*, 2022). Yet, no ion channel had been associated so far with the

rhythmic activity of DVB. Thus, this study identifies TWK-40 as a novel important

electrical component for the regulation of the defecation motor program.

Among 47 K$_{2P}$ genes in the *C. elegans* genome (Bargmann, 1998), only few have

365   been substantially investigated. Interestingly, nematode K$_{2P}$ channels have been mostly

studied using gain-of-function mutations identified in forward genetic screens. For

instance, while *twk-18(gf)* mutants cause sluggish, uncoordinated movement (Kunkel et

al., 2000), *sup-9(gf)* mutants are hyperactive with a characteristic rubberband phenotype

(de la Cruz et al., 2014). Gain-of-function mutants of *unc-58* show significant deficits in

370   locomotion, egg laying, development and aging (Kasap et al., 2018; Salkoff et al., 2005).

Except for *twk-18*, we still lack a comprehensive understanding of the electrical

properties of these channels *in vitro* and *in vivo*. The recent discovery of a universal

activating mutation that promotes the gating of vertebrate and invertebrate K$_{2P}$ channels

has opened the way to more comprehensive manipulation of ion channel activity (Ben

375   Soussia *et al.*, 2019). We used this strategy here to engineer a point mutation of TWK-40

L159N that achieved a substantial gain-of-function effect. First, we show in a related

study that the currents conducted by TWK-40(L159N) channels exhibit ~5 fold increase

compared to TWK-40(WT) when heterologously expressed in HEK293 cells (Meng *et al.*,

2022). Second, the resting membrane potential of DVB neurons was drastically

380   hyperpolarized due to the increased outward K$^+$ current. In this *twk-40(bln336)* gain-of-

function mutant, the neuronal Ca^{2+} oscillation frequency and amplitude were reduced, resulting in the significantly diminished expulsion frequency. Thus, our study not only identifies TWK-40 as a regulator of the expulsion rhythm and DVB RMP, but also provides insight into the electrical properties of this K_{2P} channel.

385   We also describe the impact of loss of *twk-40* on the expulsion behavior and DVB activity. In contrast to *gf* mutants, the resting membrane potential of DVB neuron was depolarized in *twk-40(lf)* mutant, and high Ca^{2+} oscillation activity was observed. Consequently, the expulsion number exhibited a moderate but significant increase in *twk-40(lf)* mutant. The fact that only few K_{2P} mutants with loss of function phenotypes have

390   been identified in *C. elegans* may be due to functional redundancy between K_{2P} genes. *twk-7* is a notable exception as it exhibits hyperactive locomotion (Gottschling et al., 2017; Luersen et al., 2016). Interestingly, we have found that *twk-40(lf)* mutants also exhibit increased forward and backward body bends, as well as more frequent and prolonged reversals (Meng *et al.*, 2022). A recent genetic study reported that loss of

395   function *twk-40* could reverse the reduced body curvature of *C. elegans NALCN(lf)* mutants (Zhou et al., 2022). Thus, TWK-40 appears to be required for the regulation of multiple motor rhythms.

The slight outward-rectification of the *twk-40* dependent K^+ currents in DVB neurons differs from the voltage-independent currents observed in recombinant TWK-40 recorded in HEK293 cells (Meng *et al.*, 2022). This could be due to endogenous

400   regulatory subunits expressed in DVB neurons that modify the channel kinetics of TWK-40. Alternatively, TWK-40 could form heteromers with other K_{2P} subunits *in vivo*, resulting in heterodimers with different activation kinetics (Czirjak and Enyedi, 2002).

K_{2P} channels can be modulated by a variety of biophysical parameters, such as pH,

405   temperature, and mechanic forces (Enyedi and Czirjak, 2010). Certain K_{2P} channels,

including TASK-1/3, TREK-1, and TRESK, are activated by volatile general anesthetics

at clinically relevant concentrations (Liu et al., 2004; Patel et al., 1999; Sirois et al., 2000),

indicating a major class of drug target from these K_{2P} channels. As a possible ortholog of

human TASK-1/3, identification of the critical stimulator of TWK-40 in the future could

410   promote the understanding of how the external environment and internal state affect

motor rhythm.

Materials and Methods

C. elegans strains and transgenic lines

The complete lists of constructs, transgenic lines and strains generated or acquired for

415   this study are provided in the Supplementary Table 1 and Table 2. All *C. elegans* strains

were cultured on standard Nematode Growth Medium (NGM) plates seeded with OP50,

and maintained at 22 °C. Multiple extrachromosomal transgenic lines were generated for

all rescue experiments and visually examined for consistency. One representative

transgenic line was quantified and results shown in most experiments. Unless stated

420   otherwise noted, the wild-type animal refers to the Bristol N2 strain. Only hermaphrodite

worms were used for the experiments.

Constructs and molecular biology

All expression plasmids in this study were constructed by Three-Fragment Multisite

Gateway (Invitrogen, Thermo Fisher Scientific, USA) (Magnani et al., 2006). Three-

425   Fragment Multisite gateway system consists of three entry clones. Three entry clones

comprising three PCR products (promoter, target gene, sl2-GFP or unc-54 3'UTR, in

name of slot 1, slot 2 and slot 3, respectively) were recombined into the pDEST-R4-R3 or pDEST-R4-R3-unc-54 3'UTR destination vectors by using standard LR recombination reactions to generate the expression clones.

430   All promoters used in this study were generated by PCR against a mixed-stage N2 *C. elegans* genomic DNA. Promoters include the 1.4 kb P*twk-40*, 1.2 kb P*unc-47*, 1.6 kb P*flp-10*, 1.8 kb P*unc-25s*, 2.4 kb P*exp-1* and 1.4 kb P*ges-1* genomic sequence upstream of the respective ATG start codon and were used to substitute for the *rab-3* fragment in standard BP reaction-generated entry clone A with the In-Fusion method, using

435   ClonExpress® Ⅱ One Step Cloning Kit (Vazyme, China).

To generate the entry clones slot 2 and slot 3, we used standard BP recombination reactions. An entry clone B contributing sequences of slot 2 in the expression plasmid contains sequence of a target gene, i.e., *twk-40*, GFP, or RFP. The "B" entry clones were constructed by the use of *pDONR™ 221* donor vector (Invitrogen) through BP reactions.

440   All the DNA fragments used to construct entry clone B involved in this project were amplified by PCR with the primers containing attB1 and attB2 recombination sites. An entry clone "C" contributing sequences of slot 3 in the expression plasmids contains a sequence of *unc-54* 3' UTR, *sl2d*-GFP or *sl2d*-wCherry. The "C" entry clones were constructed by use of *pDONR-P2R-P3* donor vector through standard BP reactions. The

445   corresponding PCR products with attB2R and attB3 sites were amplified by PCR.

For calcium imaging, constructs with genetic calcium sensor GCaMP6s were used for neuronal activity measurement. The GCaMP6s sequence (Chen et al., 2013) was codon-optimized for expression in *C. elegans*. The synthesized gene also contains three *C. elegans* introns and contain restriction enzyme sites to facilitate subsequent cloning. In

450   all constructs, GCaMP6s was fused in frame with a *C. elegans* codon-optimized wCherry at the C-terminus.

Genetic mutants and transgenetic arrays

Loss-of-function allele *hp834* was generated by targeted CRISPR/Cas9 genome editing with a 7 bp deletion of the sequence (GTTCGAG, base 127-133) was introduced in the

455   2nd exon of T28A8.1a. *twk-40(bln336)* was generated by CRISPR/Cas9 technology by mutating 1047-1048 base of T28A8.1a from CT to AA. Other genetic mutants used for constructing transgenic lines and compound mutants were obtained from the *Caenorhabditis Genetics Center* (*CGC*). Transgenic animals that carry non-integrated, extra-chromosomal arrays (*gaaEx*) were generated by co-injecting an injection marker

460   with one to multiple DNA construct at 5-30 ng/μl. All animals were backcrossed at least four times against N2 prior to analyses.

Behavioral assay for the defecation motor program

Defecation behavior was measured on NGM plates under standard conditions as previously described (Thomas, 1990). All behavioral tests were performed at 20-22°C.

465   Worms were synchronized and well-fed young hermaphrodites adults were transferred to fresh NGM plates with OP50 for behavior assay during freely-moving state. The defecation phenotypes were scored under a stereo fluorescence microscope (Zeiss V16, Germany). Each animal was quantified in 10 minutes after the first pBoc step pass using the Etho program software (Liu and Thomas, 1994). We simplified the observation on the

470   pBoc and the Exp steps, due to the ambiguous aBoc step (Iwasaki and Thomas, 1997). Exp/cycle was calculated as the ratio of the number of Exp/pBoc during a 10 min recording (Wang *et al.*, 2013).

Fluorescence Microscopy

Images of fluorescently tagged proteins were captured in live L4 worms. L4 stage

transgenic animals expressing fluorescence markers were picked a day before imaging.

Worms were immobilized by 2.5 mM levamisole (Sigma-Aldrich, USA) in M9 buffer.

Fluorescence signals were captured from live worms using a Plan-Apochromatic 60X

objective on a confocal microscope (FV3000, Olympus, Japan) in the same conditions.

Calcium imaging

The two integrated strains *hpIs468* (P*unc-47*::GCaMP6s::wCherry) and *hpIs582* (P*exp-1*::GCaMP6s::wCherry) were used for calcium imaging of DVB neuron and enteric

muscles, respectively. Under standard OP50 feeding, two strains exhibit normal

defecation interval and Exp step. To simplify the experiment, young hermaphrodite adult

animals were glued dorsally as described previously (Gao et al., 2015), and imaged with a

60x water objective (Nikon, Japan, numerical aperture = 1.0) (Li *et al.*, 2023). In

particular, animals are loosely immobilized with WormGlu (Histoacryl Blue, Braun,

Germany) to a sylgard-coated cover glass covered with bath-solution (Sylgard 184,

Dowcorning, USA) but leave the tail free in the recording solution. To keep the worms'

regular pumping and defecation activities, a small amount of OP50 (~100 µl, OD600 = 1)

was mixed into the recording solution, which consists of (in mM): NaCl 150; KCl 5;

$CaCl_2$ 5; $MgCl_2$ 1; glucose 10; sucrose 5; HEPES 15, pH7.3 with NaOH, ~330 mOsm.

Fluorescence images were acquired with excitation wavelength LED at 470 nm

(LED, BioLED Light Source Control Module 9-24V DC) and a digital sCMOS camera

(Hamamatsu ORCA-Flash 4.0 V2) at 100 ms per frame for 3 minute. Data was collected

from HCImage (Hamamatsu) and analyzed by Image Pro Plus 6.0 (Media Cybernetics,

Inc., Rockville, MD, USA) and Image J (National Institutes of Health). The GCaMP6s

fluorescence intensity of the region of interest (ROI) was defined as F, the background

intensity near the ROI was defined as F_0. The true neuronal calcium fluorescence signal

was obtained by subtracting the background signal from the ROI. $\Delta F/F_0 = (F-F_0)/F_0$ was

500   plotted over time as a fluorescence variation curve.

In situ electrophysiology

Dissection and recording were carried out using protocols and solutions described in the

previous studies (Gao and Zhen, 2011; Richmond and Jorgensen, 1999). Briefly, 1- or 2-

day-old hermaphrodite adults were glued (Histoacryl Blue, Braun, Germany) to a

505   sylgard-coated cover glass covered with bath solution (Sylgard 184, Dowcorning, USA)

under stereoscopic microscope M50 (Leica, Germany). After clearing the viscera by

suction through a glass pipette, the tail cuticle flap was turned and gently glued down

using WORMGLU (GluStitch Inc., Canada) to expose the DVB soma from *gaaEx0112*

[P*unc-47*::GFP::3'UTR; P*myo-2*::GFP]. The integrity of the ventral nerve cord and DVB

510   neuron were visually examined via DIC microscopy (Eclipse FN1, Nikon, Japan), and

DVB soma were patched using 15-20 MΩ-resistant borosilicate pipettes (1B100F-4,

World Precision Instruments, USA). Pipettes were pulled by micropipette puller P-1000

(Sutter, USA), and fire-polished by microforge MF-830 (Narishige, Japan). Membrane

potentials and currents were recorded in the whole-cell configuration by EPC9 amplifier

515   (HEKA, Germany), and analyzed using the Igor Pro (WaveMetrics, USA) and Clampfit

10 software (Axon Instruments, Molecular Devices, USA). Membrane potential was

recorded at 0 pA, I-V curve was collected at holding potential from –60 mV to +80 mV at

20 mV increment. Data were digitized at 10-20 kHz and filtered at 2.6 kHz. The pipette

solution contains (in mM): K-gluconate 115; KCl 25; $CaCl_2$ 0.1; $MgCl_2$ 5; BAPTA 1;

520 HEPES 10; Na_2ATP 5; Na_2GTP 0.5; cAMP 0.5; cGMP 0.5, pH7.2 with KOH, ~320

mOsm. cAMP and cGMP were included to maintain the activity and longevity of the

preparation. The bath solution consists of (in mM): NaCl 150; KCl 5; $CaCl_2$ 5; $MgCl_2$ 1;

glucose 10; sucrose 5; HEPES 15, pH7.3 with NaOH, ~330 mOsm. Chemicals were

obtained from Sigma unless stated otherwise. Experiments were performed at room

525 temperatures (20-22 °C).

Statistical analysis

The two-tailed Student's *t*-test were used to compare two groups data sets. When multiple

groups (more than two groups) of data were compared, ordinary one-way or two-way

ANOVA analysis was performed for statistical analysis. The Shapiro-Wilk test was used

530 to check for normality and either Levene's test to check for equal variance. $P < 0.05$ was

considered to be statistically significant; *, **, *** and **** denote $P < 0.05$, $P < 0.01$, P

< 0.001, $P < 0.0001$, respectively. Graphing and subsequent analysis were performed

using Igor Pro (WaveMetrics), Clampfit (Molecular Devices), Image J (National

Institutes of Health), Python, Matlab (MathWorks, USA), GraphPad Prism 8 (GraphPad

535 Software Inc., USA) and Excel (Microsoft, USA). For behavior analysis, calcium

imaging, electrophysiology and fluorescence imaging, each recording trace was obtained

from a different animal. Unless specified otherwise, data were presented as the mean ±

SEM.

Author Contributions

540 S.G. conceived experiments and wrote the manuscript. Z.Y., B.Y., Y.L., and Y.X.

performed experiments and analyzed data. L.C., J.C. J.M., W.H., Y.T., S.E.M., and C.Z.

contributed to the experiments. T.B. and M.Z. provided reagents, discussed the experiments and edited the text.

Acknowledgements

545   We thank Gao lab members for comments and discussion. This research was supported by the Major International (Regional) Joint Research Project (32020103007), the National Natural Science Foundation of China (32371189, 31871069), the National Key Research and Development Program of China (2022YFA1206001), the Overseas High-level Talents Introduction Program, and the European Research Council (TB, ERC

550   Starting Grant, Kelegans). We thank *Caenorhabditis Genetics Center*, which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440), for strains.

Declaration of Interests

The authors declare no conflict of interest.

References

555   Bargmann, C.I. (1998). Neurobiology of the Caenorhabditis elegans genome. Science *282*, 2028-2033. 10.1126/science.282.5396.2028.

Beg, A.A., and Jorgensen, E.M. (2003). EXP-1 is an excitatory GABA-gated cation channel. Nat Neurosci *6*, 1145-1152. 10.1038/nn1136.

Ben Soussia, I., El Mouridi, S., Kang, D., Leclercq-Blondel, A., Khoubza, L., Tardy, P.,
560   Zariohi, N., Gendrel, M., Lesage, F., Kim, E.J., et al. (2019). Mutation of a single residue promotes gating of vertebrate and invertebrate two-pore domain potassium channels. Nat Commun *10*, 787. 10.1038/s41467-019-08710-3.

Branicky, R., and Hekimi, S. (2006). What keeps C. elegans regular: the genetics of defecation. Trends Genet *22*, 571-579. 10.1016/j.tig.2006.08.006.

565   Brose, N., Hofmann, K., Hata, Y., and Sudhof, T.C. (1995). Mammalian homologues of Caenorhabditis elegans unc-13 gene define novel family of C2-domain proteins. J Biol Chem *270*, 25273-25280. 10.1074/jbc.270.42.25273.

Buckingham, S.D., Kidd, J.F., Law, R.J., Franks, C.J., and Sattelle, D.B. (2005). Structure and function of two-pore-domain K+ channels: contributions from genetic
570   model organisms. Trends Pharmacol Sci *26*, 361-367. 10.1016/j.tips.2005.05.003.

Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V., et al. (2013). Ultrasensitive fluorescent proteins for imaging neuronal activity. Nature *499*, 295-300. 10.1038/nature12354.

Choi, U., Wang, H., Hu, M., Kim, S., and Sieburth, D. (2021). Presynaptic coupling by
575   electrical synapses coordinates a rhythmic behavior by synchronizing the activities of a neuron pair. Proceedings of the National Academy of Sciences *118*, e2022599118.

Cook, S.J., Jarrell, T.A., Brittin, C.A., Wang, Y., Bloniarz, A.E., Yakovlev, M.A., Nguyen, K.C.Q., Tang, L.T., Bayer, E.A., Duerr, J.S., et al. (2019). Whole-animal connectomes of both Caenorhabditis elegans sexes. Nature *571*, 63-71. 10.1038/s41586-
580   019-1352-7.

Czirjak, G., and Enyedi, P. (2002). Formation of functional heterodimers between the TASK-1 and TASK-3 two-pore domain potassium channel subunits. J Biol Chem *277*, 5426-5432. 10.1074/jbc.M107138200.

Dal Santo, P., Logan, M.A., Chisholm, A.D., and Jorgensen, E.M. (1999). The inositol
585   trisphosphate receptor regulates a 50-second behavioral rhythm in C. elegans. Cell *98*, 757-767. 10.1016/s0092-8674(00)81510-x.

de la Cruz, I.P., Ma, L., and Horvitz, H.R. (2014). The Caenorhabditis elegans iodotyrosine deiodinase ortholog SUP-18 functions through a conserved channel SC-box

to regulate the muscle two-pore domain potassium channel SUP-9. PLoS Genet *10*,
e1004175. 10.1371/journal.pgen.1004175.

Enyedi, P., and Czirjak, G. (2010). Molecular background of leak K+ currents: two-pore
domain potassium channels. Physiol Rev *90*, 559-605. 10.1152/physrev.00029.2009.

Gao, S., Xie, L., Kawano, T., Po, M.D., Guan, S., Zhen, M., Pirri, J.K., and Alkema, M.J.
(2015). The NCA sodium leak channel is required for persistent motor circuit activity that
sustains locomotion. Nat Commun *6*, 6323. 10.1038/ncomms7323.

Gao, S., and Zhen, M. (2011). Action potentials drive body wall muscle contractions in
Caenorhabditis elegans. Proc Natl Acad Sci U S A *108*, 2557-2562.
10.1073/pnas.1012346108.

Goh, G.H., Maloney, S.K., Mark, P.J., and Blache, D. (2019). Episodic Ultradian Events-
Ultradian Rhythms. Biology (Basel) *8*. 10.3390/biology8010015.

Goldstein, S.A., Bayliss, D.A., Kim, D., Lesage, F., Plant, L.D., and Rajan, S. (2005).
International Union of Pharmacology. LV. Nomenclature and molecular relationships of
two-P potassium channels. Pharmacol Rev *57*, 527-540. 10.1124/pr.57.4.12.

Gottschling, D.C., Doring, F., and Luersen, K. (2017). Locomotion Behavior Is Affected
by the GalphaS Pathway and the Two-Pore-Domain K(+) Channel TWK-7 Interacting in
GABAergic Motor Neurons in Caenorhabditis elegans. Genetics *206*, 283-297.
10.1534/genetics.116.195669.

Graham, J.M., Jr., Zadeh, N., Kelley, M., Tan, E.S., Liew, W., Tan, V., Deardorff, M.A.,
Wilson, G.N., Sagi-Dain, L., and Shalev, S.A. (2016). KCNK9 imprinting syndrome-
further delineation of a possible treatable disorder. Am J Med Genet A *170*, 2632-2637.
10.1002/ajmg.a.37740.

Iwasaki, K., and Thomas, J.H. (1997). Genetics in rhythm. Trends in genetics *13*, 111-
115.

Jiang, J., Su, Y., Zhang, R., Li, H., Tao, L., and Liu, Q. (2022). C. elegans enteric motor
neurons fire synchronized action potentials underlying the defecation motor program.
Nature communications *13*, 1-15.

Jin, Y., Jorgensen, E., Hartwieg, E., and Horvitz, H.R. (1999). The Caenorhabditis
elegans gene *unc-25* encodes glutamic acid decarboxylase and is required for synaptic
transmission but not synaptic development. Journal of Neuroscience *19*, 539-548.

Johnstone, D.B., Wei, A., Butler, A., Salkoff, L., and Thomas, J.H. (1997). Behavioral
defects in C. elegans egl-36 mutants result from potassium channels shifted in voltage-
dependence of activation. Neuron *19*, 151-164. 10.1016/s0896-6273(00)80355-4.

Kasap, M., Weeks, K., Aamodt, E.J., and Dwye, D.S. (2018). Gain-of-function mutations
in a two-pore domain K + channel(unc-58) cause developmental, motor and feeding

625 defects in C. elegans modified by temperature and a channel inhibitor,loratadine. Curr Neurobiol *9*, 84-91.

Kunkel, M.T., Johnstone, D.B., Thomas, J.H., and Salkoff, L. (2000). Mutants of a temperature-sensitive two-P domain potassium channel. J Neurosci *20*, 7517-7524.

Lalevee, N., Monier, B., Senatore, S., Perrin, L., and Semeriva, M. (2006). Control of
630 cardiac rhythm by ORK1, a Drosophila two-pore domain potassium channel. Curr Biol *16*, 1502-1508. 10.1016/j.cub.2006.05.064.

LeBoeuf, B., and Garcia, L.R. (2017). Caenorhabditis elegans Male Copulation Circuitry Incorporates Sex-Shared Defecation Components To Promote Intromission and Sperm Transfer. G3 (Bethesda) *7*, 647-662. 10.1534/g3.116.036756.

635 Lee, J., Song, H.O., Jee, C., Vanoaica, L., and Ahn, J. (2005). Calcineurin regulates enteric muscle contraction through EXP-1, excitatory GABA-gated channel, in C. elegans. J Mol Biol *352*, 313-318. 10.1016/j.jmb.2005.07.032.

Lesage, F., and Lazdunski, M. (2000). Molecular and functional properties of two-pore-domain potassium channels. Am J Physiol Renal Physiol *279*, F793-801.
640 10.1152/ajprenal.2000.279.5.F793.

Li, Y., Chitturi, J., Yu, B., Zhang, Y., Wu, J., Ti, P., Hung, W., Zhen, M., and Gao, S. (2023). UBR-1 ubiquitin ligase regulates the balance between GABAergic and glutamatergic signaling. EMBO Rep *24*, e57014. 10.15252/embr.202357014.

Liu, C., Au, J.D., Zou, H.L., Cotten, J.F., and Yost, C.S. (2004). Potent activation of the
645 human tandem pore domain K channel TRESK with clinical concentrations of volatile anesthetics. Anesth Analg *99*, 1715-1722. 10.1213/01.ANE.0000136849.07384.44.

Liu, D.W., and Thomas, J.H. (1994). Regulation of a periodic motor program in C. elegans. The Journal of neuroscience : the official journal of the Society for Neuroscience *14*, 1953-1962. 10.1523/jneurosci.14-04-01953.1994.

650 Lowrey, P.L., and Takahashi, J.S. (2004). Mammalian circadian biology: elucidating genome-wide levels of temporal organization. Annu Rev Genomics Hum Genet *5*, 407-441. 10.1146/annurev.genom.5.061903.175925.

Luersen, K., Gottschling, D.C., and Doring, F. (2016). Complex Locomotion Behavior Changes Are Induced in Caenorhabditis elegans by the Lack of the Regulatory Leak K+
655 Channel TWK-7. Genetics *204*, 683-701. 10.1534/genetics.116.188896.

Magnani, E., Bartling, L., and Hake, S. (2006). From Gateway to MultiSite Gateway in one recombination event. BMC Mol Biol *7*, 46. 10.1186/1471-2199-7-46.

Mahoney, T.R., Luo, S., Round, E.K., Brauner, M., Gottschalk, A., Thomas, J.H., and Nonet, M.L. (2008). Intestinal signaling to GABAergic neurons regulates a rhythmic

660   behavior in Caenorhabditis elegans. Proc Natl Acad Sci U S A *105*, 16350-16355. 10.1073/pnas.0803617105.

Marder, E., and Bucher, D. (2001). Central pattern generators and the control of rhythmic movements. Curr Biol *11*, R986-996. 10.1016/s0960-9822(01)00581-4.

Marder, E., and Calabrese, R.L. (1996). Principles of rhythmic motor pattern generation.
665   Physiol Rev *76*, 687-717. 10.1152/physrev.1996.76.3.687.

McIntire, S.L., Jorgensen, E., and Horvitz, H.R. (1993a). Genes required for GABA function in Caenorhabditis elegans. Nature *364*, 334-337. 10.1038/364334a0.

McIntire, S.L., Jorgensen, E., Kaplan, J., and Horvitz, H.R. (1993b). The GABAergic nervous system of Caenorhabditis elegans. Nature *364*, 337-341. 10.1038/364337a0.

670   Meng, J., Ahamed, T., Yu, B., Huang, W., Mouridi, S., Leclercq-Blondel, A., Gendrel, M., Wang, Z., Chen, L., and Quan, W., et al. (2022). A descending interneuron with depolarized resting membrane potential controls C. elegans motor states. bioRxiv.

Nehrke, K., Denton, J., and Mowrey, W. (2008). Intestinal Ca2+ wave dynamics in freely moving C. elegans coordinate execution of a rhythmic motor program. Am J Physiol Cell
675   Physiol *294*, C333-344. 10.1152/ajpcell.00303.2007.

Patel, A.J., Honore, E., Lesage, F., Fink, M., Romey, G., and Lazdunski, M. (1999). Inhalational anesthetics activate two-pore-domain background K+ channels. Nat Neurosci *2*, 422-426. 10.1038/8084.

Richmond, J.E., and Jorgensen, E.M. (1999). One GABA and two acetylcholine receptors
680   function at the C. elegans neuromuscular junction. Nat Neurosci *2*, 791-797. 10.1038/12160.

Salkoff, L., Butler, A., Fawcett, G., Kunkel, M., McArdle, C., Paz-y-Mino, G., Nonet, M., Walton, N., Wang, Z.W., Yuan, A., and Wei, A. (2001). Evolution tunes the excitability of individual neurons. Neuroscience *103*, 853-859. 10.1016/s0306-
685   4522(01)00079-3.

Salkoff, L., Wei, A.D., Baban, B., Butler, A., Fawcett, G., Ferreira, G., and Santi, C.M. (2005). Potassium channels in C. elegans. WormBook, 1-15. 10.1895/wormbook.1.42.1.

Sirois, J.E., Lei, Q., Talley, E.M., Lynch, C., 3rd, and Bayliss, D.A. (2000). The TASK-1 two-pore domain K+ channel is a molecular substrate for neuronal effects of inhalation
690   anesthetics. J Neurosci *20*, 6347-6354.

Teramoto, T., and Iwasaki, K. (2006). Intestinal calcium waves coordinate a behavioral motor program in C. elegans. Cell Calcium *40*, 319-327. 10.1016/j.ceca.2006.04.009.

Thomas, J.H. (1990). Genetic analysis of defecation in Caenorhabditis elegans. Genetics *124*, 855-872.

695   Wang, H., Girskis, K., Janssen, T., Chan, J.P., Dasgupta, K., Knowles, J.A., Schoofs, L., and Sieburth, D. (2013). Neuropeptide secreted from a pacemaker activates neurons to control a rhythmic behavior. Current biology : CB *23*, 746-754. 10.1016/j.cub.2013.03.049.

Wang, H., and Sieburth, D. (2013). PKA controls calcium influx into motor neurons
700   during a rhythmic behavior. PLoS genetics *9*, e1003831. 10.1371/journal.pgen.1003831.

Wang, Z.W., Kunkel, M.T., Wei, A., Butler, A., and Salkoff, L. (1999). Genomic organization of nematode 4TM K+ channels. Ann N Y Acad Sci *868*, 286-303. 10.1111/j.1749-6632.1999.tb11294.x.

White, J.G., Southgate, E., Thomson, J.N., and Brenner, S. (1986). The structure of the
705   nervous system of the nematode Caenorhabditis elegans. Philosophical Transactions of the Royal Society B: Biological Sciences *314*, 1-340.

Witvliet, D., Mulcahy, B., Mitchell, J.K., Meirovitch, Y., Berger, D.R., Wu, Y., Liu, Y., Koh, W.X., Parvathala, R., Holmyard, D., et al. (2021). Connectomes across development reveal principles of brain maturation. Nature *596*, 257-261. 10.1038/s41586-021-03778-
710   8.

Xie, L., Gao, S., Alcaire, S.M., Aoyagi, K., Wang, Y., Griffin, J.K., Stagljar, I., Nagamatsu, S., and Zhen, M. (2013). NLF-1 delivers a sodium leak channel to regulate neuronal excitability and modulate rhythmic locomotion. Neuron *77*, 1069-1082. 10.1016/j.neuron.2013.01.018.

715   Zhou, C., Zhou, Q., He, X., He, Y., Wang, X., Zhu, X., Zhang, Y., and Ma, L. (2022). Differential modulation of C. elegans motor behavior by NALCN and two-pore domain potassium channels. PLoS Genet *18*, e1010126. 10.1371/journal.pgen.1010126.

Figure 1. Loss of function of twk-40 suppresses expulsion defects in nlp-40 and aex-2 mutants

(**A**) Schematic representation of the <u>d</u>efecation <u>m</u>otor <u>p</u>rogram (DMP), which consists of

725 three distinct sets of muscle contractions: the posterior body muscle contraction (pBoc), the anterior body muscle contraction (aBoc), and the expulsion muscle contraction (exp or EMC). The regular defecation cycle period is approximately 45 sec. (**B**) Schematic representation of the expulsion motor circuit, including the intestine, DVB neuron and enteric muscles (EM). ER, endoplasmic reticulum; ITR-1, inositol 1,4,5-trisphosphate

730 receptor; DCV: dense core vesicle. (**C, E, G**) Ethograms of defecation behavior in *nlp-40*

and *nlp-40; twk-40* double mutants, *aex-2* and *aex-2; twk-40* double mutants, and *exp-1* and *exp-1; twk-40* double mutants, respectively. Each dot and character represent 1 s. Letters 'p' and 'x' represent pBoc and exp, respectively. (**D, F, H**) Quantifications of the expulsion frequency per defecation cycle. The expulsion deficiency was partially recovered by loss of *twk-40* in *nlp-40* and *aex-2* mutants, but not in *exp-1* mutants. All data are expressed as means ± SEM. Student's *t*-test was used. Statistical significance is indicated as follows: ns, not significant, *$P < 0.05$, ***$P < 0.001$ in comparison with that as denoted.

740 *Figure 2. twk-40 regulates expulsion rhythm from DVB neuron*

(**A**) Diagram of the TWK-40 K$^+$ channel and the mutation loci. P1 and P2, pore domains;

1-4 transmembrane domains; *fs*, predicted frameshift mutation; *gf*, gain-of-function

mutation. (**B**) *twk-40* is expressed in DVB based on co-localization of P*twk-40*::TWK-

40::sl2dGFP and P*unc-47*::RFP. GFP was observed in the ventral excitatory motor

745 neurons, a few head and tail neurons, and the intestine. Scale bar, 20 μm. *Right*, zoomed

view of *twk-40* expression in DVB and DVC. Scale bar, 5 μm. (**C**) Ethograms of

defecation behavior in wild type, *twk-40(bln336, gf)* and *twk-40(hp834, lf)* animals. Each

dot and character represent 1 s. Letters 'p' and 'x' represent pBoc and exp, respectively.

(**D**) Quantifications of the expulsion frequency per defecation cycle. Compared to wild-

750   type animals, the expulsion deficiency was significantly reduced in *twk-40(bln336, gf)*,

but not in *twk-40(hp834, lf)*. (**E**) Neuronal expression of wild-type TWK-40 (TWK-

40(WT)) in DVB neurons rescues the expulsion deficiency of *twk-40(bln336, gf)* mutants.

(**F, G**) Neuronal expression of gain-of-function TWK-40(L159N) in DVB in wild type

(F) or *twk-40(hp834)* (G) recapitulates the expulsion deficiency of *twk-40(gf)* mutant. The

755   number of tested animals is indicated for each strain. All data are expressed as means ±

SEM. One-way ANOVA test was used, in which: ns, not significant, $**P < 0.01$, $****P$

< 0.0001 relative to wild type or as indicated.

Figure 3. twk-40 inhibits DVB Ca²⁺ oscillation

(**A**) Schematic representation of *C. elegans* DVB neuron and enteric muscle (EM) in a

lateral view. The genetically-encoded Ca^{2+} indicator GCaMP6s was expressed in DVB to

measure its neuronal activity. (**B**) Representative spontaneous Ca^{2+} oscillation imaged by

GCaMP in DVB neuron. *Upper*, from left to right, Ca^{2+} signal of DVB in (1) the

quiescent state, (2) at peak Ca^{2+} signal and (3) upon return to the basal state. Scale bar, 5

μm. *Bottom*, representative recording of Ca^{2+} oscillations in DVB over time. Arrow heads

indicate the corresponding time points in upper panels. (**C**) Simultaneous recording of

DVB Ca^{2+} activity and expulsion action that observed by anal bacterial fluorescence. (**D**) Cross-correlation between DVB Ca^{2+} and Exp. Faint lines indicate the results from individual Ca^{2+} transient, and the black line indicates mean value. Black dashed line

770   denotes tight correlation between DVB Ca^{2+} and Exp action with a ~ 0.8 s delay. (**E**) Representative Ca^{2+} activity (*left*) and color maps (*right*) of DVB neurons in wild type, *twk-40(bln336, gf)* and *twk-40(hp834, lf)* mutants, respectively. (**F, G**) Significant reduction of Ca^{2+} transient frequency and amplitude in *twk-40(bln336, gf)* mutant, n=20. Reduction of frequency but not amplitude in *twk-40(hp834, lf)* mutants. All data are

775   expressed as means ± SEM. One-way ANOVA test was used, in which: ns, not significant, $*P < 0.05$, $**P < 0.01$, $****P < 0.0001$ in comparison with that as denoted.

Figure 4. twk-40 regulates Ca²⁺ oscillation in DVB neuron

(**A**) Representative DVB Ca²⁺ traces (*left*) and color maps (*right*) for tissue-specific

expression of TWK-40(WT) in *twk-40(bln336, gf)* mutants. *Upper*, endogenous promoter

P*twk-40*; *middle*, P*unc-47* promoter including DVB expression; *bottom*, P*unc-25*

promoter lacking DVB expression. (**B, C**) Quantification of the frequency and amplitude of Ca^{2+} transients, n=20. (**D**) Representative DVB Ca^{2+} traces (*left*) and color maps (*right*) for tissue-specific expression of TWK-40(WT) in *twk-40(hp834, lf)* mutant. (**E, F**)

785   Quantification of the frequency and amplitude of Ca^{2+} transients, n=20 animals. All data are expressed as means ± SEM. One-way ANOVA test was used, in which: ns, not significant, *$P < 0.05$, **$P < 0.01$, ****$P < 0.0001$ in comparison with that as denoted.

790   ***Figure 5. twk-40 mutants regulate Ca²⁺ dynamics of enteric muscles***

(**A**) *i*. Schematic representation of *C. elegans* DVB neuron and enteric muscle (EM) in a lateral view. The genetically-encoded Ca^{2+} indicator GCaMP6s was expressed in EM. *ii*. Representative Ca^{2+} oscillation imaged by GCaMP in EM *in-vivo*. *Upper*, from left to right, Ca^{2+} signal of EM in (1) the quiescent state, (2) at peak Ca^{2+} signal and (3) upon return to the basal state. Scale bar, 5 μm. *Bottom*, representative recording of Ca^{2+} oscillations in EM over time. Arrow heads indicate the corresponding time points in upper panels. Color map of the rhythmic EM Ca^{2+} activities in wild-type animals. (**B**) Representative Ca^{2+} activity (*left*) and color maps (*right*) of enteric muscles in *twk-40(bln336, gf)* and *twk-40(hp834, lf)* mutants, respectively. (**C**) Quantification of EM Ca^{2+} transients frequency and amplitude , n=20. Ca^{2+} transient frequency was significantly reduced in *twk-40(bln336, gf)* mutant, but was increased in *twk-40(hp834, lf)* mutant. EM Ca^{2+} amplitude was not altered in either mutants. All data are expressed as means ± SEM. One-way ANOVA -test was used, in which: ns, not significant, **$P < 0.05$, ****$P < 0.0001$ in comparison with that as denoted.

Figure 6. DVB regulates enteric muscles Ca²⁺ dynamics in twk-40 mutants

(**A, C**) Representative EM Ca^{2+} traces (*left*) and color maps (*right*) in *twk-40(bln336, gf)*

(A) and *twk-40(hp834, lf)* (C) for tissue-specific expression of TWK-40(WT). 1,

endogenous promoter P*twk-40*; 2, P*unc-47* promoter including DVB expression; 3, P*unc-*

810   *25* promoter lacking DVB expression; 4, P*exp-1* promoter for enteric muscle expression.

(**B, D**) Rescue of EM Ca^{2+} transient frequency by expression of TWK-40(WT) in DVB,

but not enteric muscles or other neurons, n=20. (**E**) Representative EM Ca^{2+} traces (*left*)

and color maps (*right*) in *twk-40(hp834, lf)* following tissue-specific expression of TWK-

40(L159N). (**F**) Reduction of EM Ca^{2+} transient frequency following TWK-40(L159N)

815   expression in DVB, but not in enteric muscles or other neurons, n=20. All data are

expressed as means ± SEM. One-way ANOVA test was used, in which: ns, not

significant, $*P < 0.05$, $**P < 0.01$ in comparison with that as denoted.

Figure 7. TWK-40 hyperpolarizes the DVB neuron.

(**A**) *Left*, schematic representation of *C. elegans* DVB neuron for whole-cell patch clamp

recording. *Right*, GFP-labeled DVB soma and axon. (**B**) Representative outward K^+

currents recorded from DVB neurons in different genotypes. (**C**) Quantification of the

current-voltage curves from different genotypes reveals that the outward currents were

reduced in *twk-40(hp834, lf)* mutant, but increased in *twk-40(bln336, gf)* mutant. (**D, E**)

Representative traces and quantification of the resting membrane potential (RMP)

recorded from DVB in different genotypes. (**F**) The working Model: K_{2P} channel TWK-

40 regulates the rhythmic expulsion behavior by setting the resting membrane potential of DVB neuron. One-way ANOVA test was used to test the significant difference of average currents at +80 mV, in which: ns, not significant, *$P < 0.05$, **$P < 0.01$ in comparison with that as denoted.

Supplementary Figure 1. TWK-40 conservation, domain structure and gene mutations.
(**A**) TWK-40 amino acid sequence alignment with other species. Identical amino acids are highlighted in yellow. Black boxes denote the transmembrane domains, pink dashed boxes denote the pore domains. Gene mutations are labeled on top of the sequence. (**B**) TWK-40 E58K suppresses the expulsion frequency per defecation cycle defects of *nlp-40*

and *aex-2* mutants, but not *exp-1* mutants. (**C**) Expression of TWK-40(WT) in the intestine does not rescue the expulsion defect of *twk-40(bln336, gf)*. All data are expressed as means ± SEM. Student's *t*-test was used, in which: ns, not significant, *$P <$ 0.05, in comparison with that as denoted.

Supplementary Figure 2. DVB/EM Ca^{2+} activities are dependent upon nlp-40.

(**A**) Quantification of the expulsion frequency in *nlp-40(tm4085)* mutant. (B, D) Color

maps summarize the DVB Ca^{2+} and EM Ca^{2+} activities in wild type and *nlp-40* animals,

845   respectively. (C, E) Quantification of the frequency and peak amplitude of these Ca^{2+}

activities. $*P < 0.05$; $****P < 0.0001$; two-tailed unpaired Student's t-test. n = 10-30

animals. Error bars, SEM.

Supplementary Figure 3. twk-40* mutation does not *alter Ca^{2+} dynamics in DVB and

850   ***enteric muscles.***

(**A, D**) Representative single Ca^{2+} transients from DVB (A) and enteric muscle (D) in

wild type, *twk-40(bln336, gf)* and *twk-40(hp834, lf)* animals, respectively. (**B, C, E, F**)

The rise time (10-90%) and decay time (10-90%) of the Ca^{2+} transients exhibited no

difference between these genotypes. Each group, n=20 animals. All data are expressed as

855   means ± SEM. One-way ANOVA test was used, in which: ns, not significant, in

comparison with that as denoted.

Supplementary Figure 4. twk-40 regulates DVB Ca^{2+} oscillations in unc-13(lf) background.

(**A**) Representative DVB recordings (*left*) and color maps of rhythmic Ca^{2+} oscillations (*right*) in *unc-13(s69)*, *unc-13(s69); twk-40(bln336, gf),* and *unc-13(s69); twk-40(hp834, lf)* mutants, respectively. (**B, C**) Quantification of the frequency and amplitude of DVB Ca^{2+} transients in different mutants. The frequency and amplitude were significantly reduced in *unc-13(s69); twk-40(bln336, gf)* mutants compared to *unc-13(s69)*. The Ca^{2+} frequency was slightly increased in *twk-40(hp834, lf)* mutant, but the amplitude was not altered. Each group, n=40. All data are expressed as means ± SEM. One-way ANOVA test was used, in which: ns, not significant, $*P < 0.05$, $**P < 0.01$, $****P < 0.0001$ in comparison with that as denoted.

870 *Supplementary Figure 5. Ectopic expression of TWK-40(L159N) inhibits DVB*

(**A, D**) Representative Ca^{2+} traces (*left*) and color maps (*right*) for DVB neurons in wild

type (A) and *twk-40(hp834, lf)* (D) following expression of TWK-40(L159N) in different

neuron groups. *Upper*, endogenous *twk-40* promoter; *middle*, *unc-47* promoter including

DVB expression; *bottom*, short *unc-25* promoter (P*unc-25s*) lacking DVB expression. (**B,**

875 **C, E, F**) Quantification of the frequency and amplitude of the oscillated DVB Ca^{2+}

transients in different genetic backgrounds, both of which were reduced when TWK-

40(L159N) was expressed in the DVB, but not in other neurons, n=20. All data are

expressed as means ± SEM. One-way ANOVA test was used, in which: ns, not

significant, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $****P < 0.0001$ in comparison with

880   that as denoted.