1 **The admixture histories of Cabo Verde**
2

3 **Authors:**
4 Romain Laurent[1], Zachary A. Szpiech[2,3], Sergio S. da Costa[1], Valentin Thouzeau[4,5], Cesar A. Fortes-Lima[6],
5 Françoise Dessarps-Freichey[1], Laure Lémée[7], José Utgé[1], Noah A. Rosenberg[8], Marlyse Baptista[9,10], and
6 Paul Verdu[1,*].
7
Authors' affiliations :
1. UMR7206 Eco-anthropologie, CNRS-MNHN-Université Paris Cité, Paris, France
2. Department of Biology, Pennsylvania State University, University Park, PA 16802, USA
3. Institute for Computational and Data Sciences, Pennsylvania State University, University Park, PA 16802,
USA
4. UMR 7534 Centre de Recherche en Mathématiques de la Décision, CNRS-Université Paris-Dauphine-
PSL University, Paris, France
5. Département d'Etudes Cognitives, Laboratoire de Sciences Cognitives et Psycholinguistique, ENS-PSL
University-EHESS-CNRS, Paris, France
6. Department of Organismal Biology, Sub-department of Human Evolution, Evolutionary Biology Centre,
Uppsala University, Uppsala, Sweden
7. Plate-forme Technologique Biomics–Centre de Ressources et Recherches Technologiques (C2RT), Insti-
tut Pasteur, Paris, France.
8. Department of Biology, Stanford University, Stanford, CA, USA
9. Department of Linguistics, University of Michigan, Ann Arbor, MI, USA
10. Department of Afroamerican and African Studies, University of Michigan, Ann Arbor, MI, USA
24
25 ***Corresponding author:**
26 Paul Verdu, UMR7206 Eco-anthropologie, CNRS-MNHN-Université Paris Cité, Musée de l'Homme, 17,
27 place du Trocadéro, 75016 Paris, France. Email : paul.verdu@mnhn.fr. Tel : +33 144057317

28
Authors' contributions:
PV, MB, NAR designed sampling strategy
PV, MB, SSC, VT, CFL conducted sampling field-work
FD, PV, LL, JU conducted molecular-genetics experiments
RL, PV conducted genetic data quality control and merging
SSC conducted linguistic transcriptions
MB, SSC conducted linguistic data categorization
RL, PV conducted linguistic data parsing
RL, PV, ZAS, MB, SSC, VT conducted genetic and linguistic statistical descriptions
RL, PV conducted genetic inferences
PV, RL, MB, ZAS wrote the paper with the help of all authors
PV, MB, and NAR supervised the research
41
42 **Conflict of interest statement**
43 The authors declare no conflict of interests for this work.
44
Ethics statement
Research sampling protocols followed the Declaration of Helsinki guidelines and the French laws of scien-
tific research deontology (Loi n° 2016-483 du 20 avril 2016). Research and ethics authorizations were pro-

vided by the Ministério da Saúde de Cabo Verde (228/DGS/11), Stanford University IRB (Protocol ID
n°23194–IRB n°349), University of Michigan IRB (n°HUM00079335), and the French CNRS ethics com-
mittees and CNIL (Declaration n°1972648). All volunteer participants provided written and video-recorded
informed consent.
52
Data availability
The novel genetic data presented here can be accessed via the European Genome-phenome Archive (EGA)
database accession number ==#XX (pending)== upon request to PV. Linguistic data presented here can be ob-
tained directly upon request to the corresponding author (PV). Both datasets can be shared provided that
future envisioned studies comply with the informed consents provided by the participants, and in agreement
with institutional ethics committee's recommendations applying to this data.
59

60 **Software availability**
61 *MetHis* is an open source C software available with user manual on GitHub at https://github.com/romain-
62 laurent/MetHis (Fortes-Lima et al. 2021).
63

Acknowledgements
The authors would like to warmly thank all Cabo Verdean participants to this study as well as the UniCV for
facilitating the project. This project was funded in part by the French "Agence Nationale pour la Recherche"
grant ANR METHIS 15-CE32-0009-1, and by a grant from the France-Stanford Center for Interdisciplinary
Studies. MB was supported in part by the Linguistics Department at the University of Michigan (MI, USA).
ZAS was supported in part by startup funds from the Department of Biology at the Pennsylvania State Uni-
versity (PA, USA). CAFL was supported in part by the Marcus Borgströms Foundation for Genetic Research
and the Bertil Lundman Foundation for Anthropological Studies (Sweden). We thank the platform
"Paléogénomique et génétique moléculaire" (P2GM) of the French Muséum National d'Histoire Naturelle at
the Musée de l'Homme for support handling biological samples and generating genetic data. We thank the
BIOMICS platform from the Pasteur institute for performing genotyping analyses. Finally, the authors
warmly thank Frank Alvarez-Pereyre, Erkan O. Buzbas, Marta Ciccarella, Pierre Darlu, Ethan M. Jewett,
Evelyne Heyer, Marie-France Mifune, Jorge M. Rocha, and Bruno Toupance, for useful comments and dis-
cussions about this work.
78
79

80 **Keywords**
81 Admixture; anthropology; population genetics; historical inference; machine-learning; approximate Bayesian
82 computations; genetic diversity; linguistic diversity; slave-trade
83
84

85   **ABSTRACT**
86
From the 15th to the 19th century, the Trans-Atlantic Slave-Trade influenced the genetic and cultural diversi-
ty of numerous populations. We explore genomic and linguistic data from the nine islands of Cabo Verde,
the earliest European colony of the era in Africa, a major Slave-Trade platform between the 16th and 19th
centuries, and a previously uninhabited location ideal for investigating early admixture events between Eu-
ropeans and Africans. We find that genetic admixture in Cabo Verde occurred primarily between Iberian and
Senegambian populations, although forced and voluntary migrations to the archipelago involved numerous
other populations. Inter-individual genetic and linguistic variation recapitulate geographic distribution of
individuals' birth-places across Cabo Verdean islands, suggesting that Kriolu language variants have devel-
oped together with genetic divergences. Furthermore, we find that admixture occurred both early in each
island, long before the 18th-century massive TAST deportations triggered by the plantation economy, and
after this era. Our results illustrate how shifting socio-cultural relationships between enslaved and non-
enslaved communities shaped enslaved-African descendants' descent on both sides of the Atlantic.
99
100

INTRODUCTION

Between the 15th and 19th centuries, European colonization and the Trans-Atlantic Slave-Trade (TAST) put into contact groups of individuals previously isolated genetically and culturally. These forced or voluntary migrations profoundly influenced the descent of numerous European, African, and American populations, creating new cultures, languages, and genetic patterns [1,2].

Population geneticists have extensively described genetic admixture patterns in enslaved-African descendants in the Americas, and mapped their genomes for regions of recently shared ancestry with continental Africa and Europe [3,4]. This allowed for reconstructing their detailed possible origins, as this knowledge is often intractable with genealogical records alone [1]. Furthermore, genetic admixture-mapping methods have been used to identify genetic variation underlying phenotypic variation [5,6], and to identify post-admixture natural selection signatures [7], thus revealing how admixture shaped human populations' recent evolution. Maximum-likelihood approaches based on linkage-disequilibrium (LD) patterns of admixed individuals[8,9] have repeatedly highlighted the diversity of admixture processes experienced by populations historically related to the TAST. In particular, they identified different European, African, and American populations respectively at the source of genetic admixture patterns, sometimes consistent with the preferred commercial routes of each European empire [8,10]. Furthermore, they identified variable timing of admixture events during and after the TAST, sometimes consistent with major socio-historical events such as the expansion of the plantation economic system or the abolition of slavery [11,12]. From a cultural perspective, linguists have shown that novel contact-languages, such as creole languages [13,14], emerged from recurring interactions between socio-economically dominant Europeans with Africans and Americans. Furthermore, they identified the languages of origin of numerous linguistic traits in several creole languages [15–17], and emphasized the complex histories of contacts that shaped language diversity on both sides of the Atlantic.

Numerous questions remain unsolved and novel interrogations have emerged concerning the history of admixture during and after the TAST. First, while the genetic history of enslaved-African descendants in the Americas has been extensively studied, the influence of the TAST on genetic admixture in Africa remains under-investigated. Nevertheless, studying these questions in Africa would provide invaluable information about the influence of the onset and early stages of the TAST and the subsequent expansion of European empires on genetic admixture patterns on both sides of the ocean. Second, while admixture-LD inference methods have repeatedly brought novel insights into the admixture processes experienced by enslaved-African descendant populations, they could only explore historical models with one or two pulses of admixture, a methodological limitation extensively discussed by the authors [8,9]. Nevertheless, complex admixture histories may be expected as a result of the recurring flows of enslaved-Africans forcibly displaced between and within continents, changes of social relationships among enslaved and non-enslaved communities, and variable assimilation of new migrants in pre-existing communities, during and after the TAST [18,19]. Finally, while the comparison of genetic and linguistic diversities has been the focus of numerous evolutionary anthropology studies at large geographical scales, it has rarely been endeavored for creole-speaking populations in the historical context of the TAST [20–22].

Here, we propose to reconstruct the detailed genetic and linguistic admixture histories of Cabo Verde, as this archipelago represents an ideal case to address these three understudied aspects of TAST history. First, Cabo Verde is the first European peopling colony in Sub-Saharan Africa, located 500 kms West of Senegal in Western Africa (**Figure 1**), and settled in the 1460s by Portuguese immigrants and enslaved-Africans forcibly removed from the continent. After Columbus, and in particular after the 17th century expansion of plantation economy in the Americas, Cabo Verde served as a major slave-trade platform between continents [23]. Second, Cabo Verde forms an archipelago of nine inhabited islands that were settled over the course of three centuries due to the changing political, commercial, and forced or voluntary migratory contexts [24–26].

Therefore, studying the admixture history of Cabo Verde will provide unique insights into the onset of the
TAST before Columbus, and into the history of slavery thereafter. This setting further promises to illustrate,
at a micro-geographical scale island-per-island, the fundamental socio-historical and serial founding
migrations mechanisms having influenced genetic patterns in admixed populations throughout the TAST.
Finally, Cabo Verdean Kriolu is the first creole language of the TAST, born from contacts between the
Portuguese language and a variety of African languages [15,17,27,28]. The archipelago thus represents a unique
opportunity to investigate, jointly, genetic and linguistic admixture histories and their interactions since the
mid-15th century.

Previous genetic studies exploring sex-specific or genome-wide markers from several islands of the
archipelago [29–32], attested to the dual, sex-biased, origins of the Cabo Verdean gene-pool, resulting mainly
from admixture between African females and European males. Furthermore, these studies described variable
admixture patterns across islands. Interestingly, adaptive-introgression signals for malaria resistance in
Santiago island were recently identified as a result of migrations and genetic admixture during the TAST [33].
Finally, joint analyses of genetic and linguistic diversities from Santiago showed that genetic and linguistic
admixture histories possibly occurred in parallel [21].

Based on these previous studies, we propose to first determine which continental African and European
populations in fact contributed to the genetic landscape of each island today. Indeed, which enslaved-African
populations only briefly transited through the archipelago, and which remained for longer periods is largely
debated by historians [23–25]; and, while Portuguese influence is clear, further details about which European
migrations genetically influenced Cabo Verde remain to be assessed [34]. These aspects are often crucial for
understanding the genetic history of enslaved-African descendant populations on either side of the Atlantic
[1,12,35,36]. Second, we propose to further evaluate the possible parallels between genetic and linguistic
admixture histories in each island to better understand how contacts shaped cultural variation during the
TAST [21]. Finally, we reconstruct the detailed history of admixture dynamics in each island since the 15th
century. Altogether, this will highlight the socio-historical mechanisms that shaped the genetic and linguistic
diversity of this population, the first to be born from the TAST.
172
173

RESULTS

We investigate genetic and linguistic variation in 261 Kriolu speakers from the nine Cabo Verdean islands (Brava, Fogo, Santiago, Maio, Sal, Boa Vista, São Nicolau, São Vicente, Santo Antão, **Figure F1**). With novel genome-wide genotyping autosomal data (**Figure SF1**), we first describe genetic differentiation patterns in Cabo Verde and other enslaved-African descendants in the Americas from previous datasets, in particular with respect to continental Africa and Europe. Second, we deploy local-ancestry inferences and determine the best proxy source-populations for admixture patterns in each Cabo Verde island. Then, we describe Runs of Homozygosity and genetic isolation-by-distance patterns at reduced geographical scale within Cabo Verde. We also investigate Kriolu utterance diversity with respect to geography and socio-cultural co-variables and, then, investigate jointly genetic and linguistic admixture patterns throughout the archipelago. Finally, we infer the detailed genetic admixture history of each island using the machine-learning *MetHis*-Approximate Bayesian Computation (ABC) approach [37].

1. Cabo Verde and other admixed populations related to the TAST in the worldwide genetic context

We explored genetic diversity patterns captured along the first three axes of the Multi-Dimensional Scaling projection of individual pairwise Allele Sharing Dissimilarities (ASD [38]), computed from different individual subsets (**Table ST1**). Detailed ASD-MDS decompositions are provided in **Supplementary Text** and **Figures SF2-6**. Note that we considered seven geographical regions in Africa shown in **Figure F1**.

Figure F2 show that the second ASD-MDS axis distinguishes West Western African Senegambian populations from East Western Africans, while Central Western Africans are at intermediate distances between these two clusters. Moreover, the third MDS axis separates Northern and Southern European populations; the British-GBR and USA-CEU individuals clustering at intermediate distances between the Finnish-FIN, and a cluster represented by Iberian-IBS and Tuscan-TSI Western Mediterranean individuals. Consistently with previous results [3,21], on the first MDS axis, Cabo Verdean individuals cluster almost exclusively along a trajectory going from the Southern European cluster to Senegambia (**Figure F2A-C**), with little traces of affinity with other African populations. Instead, the USA African-American ASW (**Figure F2D-F**) and Barbadian-ACB (**Figure F2G-I**) cluster along a trajectory going from the GBR and CEU cluster to Central and East Western Africa; and Puerto Ricans-PUR cluster along a trajectory going from the Southern European cluster to the Central Western African cluster (**Figure F2J-L**).

2. Genetic structure in Cabo Verde and other admixed populations related to the TAST

Based on these results, we further investigated patterns of individual genetic structure among Cabo Verdean-born individuals, ASW, and ACB populations with respect to European and Western, Central, South-Western, and Southern African populations (**Table ST1**), using ADMIXTURE [39]. Extended descriptions of the results are presented in **Supplementary Text**.

At K=2, the orange genetic-cluster in **Figure F3A** is maximized in African individuals while the blue alternative cluster is maximized in Europeans. Cabo Verdean, ASW and ACB individuals exhibit intermediate genotype-membership proportions between the two clusters, consistently with patterns expected for European-African admixed individuals. ACB individuals show, on average, the highest membership to the orange "African" cluster (88.23%, SD=7.33%), followed by the ASW (78.00%, SD=10.88%), and Cabo Verdeans (59.01%, SD=11.97%). Furthermore, membership proportions to this cluster are highly variable across Cabo Verdean islands, with highest average memberships for Santiago-born individuals (71.45%, SD=10.39%) and Maio (70.39%, SD=5.26%), and lowest for Fogo (48.10%, SD=6.89%) and Brava (50.61%, SD=5.80%). Furthermore, we found that inter-individual memberships' variation within Cabo Verde islands captured as F_{ST}/F_{ST}^{max} values [40], are significantly different across pairs of islands for 32 out of 36 comparisons (Wilcoxon rank-sum test $p<3.96\times10^{-8}$), with variability across islands ranging from a lowest value of 0.010 in Santo Antao to a highest value of 0.0519 in Santiago (**Figure SF7**).

At *K*=5, the new red cluster is maximized in the YRI, Igbo and ESN populations, distinct from Western
and Southern African orange and gray clusters, respectively. Note that the former orange cluster is almost
completely replaced with red membership in the ACB and ASW populations. Conversely, Cabo Verdean
individuals exhibit small to virtually no membership to this East Western African red cluster, but their Afri-
can component, instead, resembled genetic patterns found among West Western Africans.
Ultimately, at *K*=10, the light-green cluster is maximized in Cabo Verdean individuals born on Maio,
Boa Vista, Sal, and São Nicolau, distinct form the dark green cluster maximized in Santo Antão and São
Vicente, and hence producing three distinct ADMIXTURE patterns among Cabo Verdean islands. Further-
more, an alternative mode at *K*=10 shows (**Figure SF8**), that Cabo Verdean individuals resemble more IBS
and TSI patterns for their European-like membership than ASW and ACB individuals who, instead, resemble
more CEU and GBR patterns, consistently with ASD-MDS results and previous studies [10,21].
232

3. Local-ancestry in Cabo Verde and other admixed populations related to the TAST
ASD-MDS and ADMIXTURE descriptive analyses do not formally test admixture and putative source popu-
lations of origins, they rather disentangle genetic-structure resemblances among individuals [39,41–44]. Therefore,
we conducted admixture-LD haplotypic local-ancestry inferences with the SHAPEIT2-
CHROMOPAINTER-SOURCEFIND pipeline [45–47], to identify the possible European and African popula-
tions at the sources of admixture patterns.
Figure F3B shows striking differences concerning both the European and the African source popula-
tions involved in the admixture history of ACB, ASW, and birth islands of Cabo Verdeans. We find that all
Cabo Verdean islands of birth share almost all their European haplotypic ancestry with the Iberian-IBS popu-
lation rather than other European populations. Santiago-born individuals present the smallest (27%) average
haplotypic ancestry shared with IBS, and Fogo-born the highest (51%). Conversely, the ASW and ACB both
share haplotypic ancestries only with the USA-CEU of North-Western European origin (20% and 12% re-
spectively).
Furthermore, we find that all Cabo Verdeans almost exclusively share African haplotypic ancestries
with two Senegambian populations (Mandinka and Wolof) and very reduced to no shared ancestries with
other regions of Africa. More specifically, we find that the Mandinka from Senegal are virtually the sole
African population to share haplotypic ancestry with Cabo Verdeans born in Brava, Fogo, Santiago, Maio,
and Boa Vista, and the majority of the African shared ancestry for Sal, São Nicolau, São Vicente, and Santo
Antão. In these four latter islands, we find shared haplotypic ancestry with the Wolof population ranging
from 5%-4% for individuals born on Sal (considering six or four possible sources, respectively), up to 22%-
16% for Santo Antão. Finally, we find limited (1%-6%) shared haplotypic ancestry with East Western (Igbo,
YRI, or Ga-Adangbe) or South-West Central (Kimbundu, Kongo, or Ovimbundu) African populations in all
Cabo Verdean islands, except Fogo and Brava, and the specific populations identified and their relative pro-
portion of shared haplotypic ancestries varies across analyses. Conversely, we find that the ASW and ACB
populations share African haplotypic ancestries in majority with East Western African populations (YRI, Ga-
Adangbe, and Igbo), and substantial shared ancestries with Senegambian populations (5-10%), the MSL
from Sierra Leone in Central Western Africa (5-12%), and South-West Central African populations (3-18%),
albeit variable depending on the number of putative sources considered.
261

4. Genetic and linguistic diversity and admixture within Cabo Verde
Runs of Homozygosity (ROH) patterns within Cabo Verde
Runs of homozygosity (ROH) are Identical-By-Descent haplotypes resulting from recent parental relatedness
and present in individuals as long stretches of homozygous genotypes. Their length and abundance can
reflect demographic events, such as population bottlenecks and founder events, natural selection, and cultural
preferences for endogamy [48–50].

We find higher levels of long ROH ($\geq$1cM) in Cabo Verdeans compared to most other analyzed
populations, including ASW and ACB (**Figures F4A, SF9**). We find the highest levels of long-ROH in
individuals born on Maio, Boa Vista, Sal, São Nicolau, São Vincente, and Santo Antão, with a mean
individual length of long-ROHs around 3cM (**Figure F4B**), and the lowest levels of long-ROH in Santiago
and Brava. Among long ROH (**Figure SF10C**), we find little to no correlation with total non-ROH levels for
African local-ancestry segments (Pearson ρ=-0.06689, p=0.3179), European (ρ=0.1551, p=0.01989), or
Asian (ρ=0.06239, p=0.3516).

In admixed populations, we expected that some of the long ROH spanned local ancestry breakpoint
switches (see **Materials and Methods**), indicating that the most recent common ancestor existed after the
initial admixture event having generated local-ancestry patterns. Furthermore, we expected that these
"spanning" ROH should be among the longest ROH observed if admixture occurred only in the past few
generations. We find that (**Figure F4C**), almost uniformly across Cabo Verde, the longest ROH identified
indeed spanned at least one ancestry breakpoint, excluding the very few Asian ancestry regions identified.
Furthermore, correcting ancestry-specific long-ROH sizes (**Figure SF11**) for individual's total ancestry
fraction of that ancestry, we find that individuals born in Brava and Fogo have, on average, an apparent
overrepresentation of European ancestry in long-ROH (**Figure F4D**), and that individuals from Sal and São
Vincente have, conversely, an apparent overrepresentation of African ancestry. Finally, we find that
individuals from Santiago, Maio, Boa Vista, São Nicolau, and Santo Antão have relatively similar long-ROH
levels in either African or European segments. This latter pattern may be consistent with these populations
being founded by admixed individuals, while the former patterns could indicate, in addition to such founding
effect, more recent or recurring contributions from either source, respectively (Mooney et al. 2018).

ASD and utterance diversity within Cabo Verde
Together with the above results, previous studies suggested substantial genetic and linguistic isolation across
certain Cabo Verdean groups of islands [15,17,21,27,28,32]. We aim at further characterizing possible patterns of
isolation-by-distance among and within islands, both genetically and linguistically. We explored both pair-
wise ASD and inter-individual variation in Kriolu utterance frequencies (**Material and Methods**), in the
same set of 225 Cabo Verde-born individuals, using MDS and Mantel testing (**Figure F5**, **Table T2**, and
Table ST14).

The first ASD-MDS axis mainly differentiates Brava and Fogo-born individuals compared to Santiago
(**Figure F5A-B**). The second axis differentiates mainly individuals from Santiago, Fogo, and Brava from all
other islands, while the third axis differentiates Boa Vista, São Nicolau, Sal, and Maio from all other islands.
Furthermore, we find a significant positive correlation between ASD and actual individual birth-places
across Cabo Verde (**Table T2**; Spearman ρ=0.2916, two-sided Mantel p<2x10^{-4}). This correlation increases
within-islands excluding inter-island distances (Spearman ρ=0.3460, two-sided Mantel p<2x10^{-4}), thus illus-
trating the strong signal of genetic isolation-by-distance [51] within Cabo Verde at very reduced geographical
scales.

Furthermore, the first utterance-MDS axis of pairwise inter-individual Euclidean distances between ut-
terance frequencies mainly differentiates Santiago and Santo Antão/São Vincente-born individuals' speech-
varieties; all other Cabo Verdeans clustering intermediately (**Figure F5C-D**). The third axis further separates
speech-varieties recorded in individuals from Fogo, Maio, and Brava. Analogously to genetic differentiation
patterns, we find a positive correlation between differences in utterance frequencies and actual birth-places'
distances (Spearman ρ=0.2794, two-sided Mantel p<2x10^{-4}), as well as paternal and maternal birth-places
respectively (**Table T2**). However, contrarily to ASD, we find that utterance-frequencies differences stem
from inter-birth-islands' distances, rather than shorter distances within islands only. Consistently with previ-
ous results from Santiago only (Verdu et al. 2017), these results altogether show that speech-varieties are
significantly transmitted from one generation to the next throughout Cabo Verde, anchored in individuals'

birth-places, albeit not accounting for the majority of observed linguistic variation in our dataset. Indeed, we find that age-differences also substantially correlate with utterance-frequency differences even correcting for individual birth-places (Spearman ρ=0.2294, two-sided partial-Mantel p<2x10^{-4}). Finally, while we might intuitively expect that academic education influences idiolects, we find instead that differences in education-duration do not correlate with Kriolu utterance-frequencies differences, whether correcting for residence or birth-places distances, or not (**Table T2**). This shows the modest influence of academic education on Kriolu variation. Altogether, our results highlight strong genetic and linguistic isolation-by-distance patterns at reduced geographic distances within Cabo Verde.

Genetic and linguistic admixture within Cabo Verde

Interestingly, we find that absolute differences in inter-individual genetic admixture levels from Africa, estimated with ADMIXTURE or ASD-MDS, significantly correlate with actual birth-places distances across islands (Spearman ρ=0.1865, two-sided Mantel p<2x10^{-4} and ρ=0.1813, p<2x10^{-4}, respectively), but not within-islands only (ρ=0.0342 p=0.3094 and ρ=0.0282 p=0.3385, respectively), indicating that two individuals born on far-away islands are more likely to differ in African-admixture levels.

We explored inter-individual variation in Kriolu utterance frequencies specifically for uttered items of clearly African and dual European-African origins (Categories A and B; **Material and Methods**) providing an estimate of individual African linguistic-admixture scores (Verdu et al. 2017). We find that African linguistic-admixture score differences significantly correlate with actual birth-places distances throughout Cabo Verde (Spearman ρ=0.1297, two-sided Mantel p<2x10^{-4}), and even marginally significantly correlate with birth-places distances at short distances within birth-islands (Spearman ρ=0.1209, two-sided Mantel p=0.0419).

Finally, we find a significant positive correlation (Spearman ρ=0.2070, p=0.0018) between genetic and linguistic admixture in Cabo Verde (**Figure 5E**), indicating that individuals who frequently use African-related utterances in their manner of speaking Kriolu are more likely to exhibit higher levels of African genetic-admixture. This correlation remains, respectively, marginally significant and significant when considering utterances of strictly African-origin (Category A) or utterances with a dual European-African etymology (Category B) separately (Spearman ρ=0.1631, p=0.0143, and ρ=0.1829, p=0.0059, respectively).

Altogether, our genetic and linguistic diversity results generalize our previous results obtained in Santiago only [21], and further suggest that genetic and linguistic admixture histories occurred in parallel in Cabo Verde. Together with the above genetic and linguistic isolation-by-distance signals and ROH results, this strongly suggests that different founding events and admixture histories for each Cabo Verde islands are at the root of genetic and linguistic diversity in the archipelago.

5. Genetic admixture histories in Cabo Verde inferred with *MetHis*-ABC

MetHis–ABC prior checking

We considered four competing genetic-admixture scenarios (**Figure F6**), tested separately for individuals born on each Cabo Verdean island with *MetHis*–ABC machine-learning scenario-choice and posterior parameter inferences [37,52,53]. ABC inferences are based on 42 summary statistics (**Table ST3**) calculated from 60,000 independent autosomal SNPs in individuals form each Cabo Verdean island, and in the African Mandinka and European Iberian-IBS proxy source populations.

We find that the summary-statistics calculated from the observed datasets fall well within the space of summary-statistics obtained from 10,000 simulated-datasets under each of the four competing scenarios (**Figures SF13-22**), considering non-significant (α>5%) goodness-of-fit, visual inspection of summary-statistics PCA-projections, and each summary-statistics' distribution, for each Cabo Verdean birth-island separately. Prior-checks thus demonstrate that *MetHis* simulations are appropriate for further ABC scenario-choice and posterior parameter inferences.

MetHis–Random Forest-ABC scenario-choices

Overall (**Figure F7B**), *MetHis*-RF-ABC scenario-choices indicate that multiple pulses of admixture from the European and African source populations (after the founding admixture pulse, two independent admixture pulses from both Africa and Europe: "Afr2Pulses-Eur2Pulses" scenarios, **Figure F4**), best explain the genetic history of individuals born on six out of nine Cabo Verdean islands. Furthermore, we find that more complex scenarios involving a period of recurring admixture from either source best explain the history of the remaining three islands. Conversely, periods of recurring admixture processes from both Africa and Europe ("AfrRecurring-EurRecurring") are least favored throughout Cabo Verde.

RF votes favor the Afr2Pulses-Eur2Pulses scenario with high posterior probabilities of identifying correctly this scenario for individuals born, respectively, in Santiago (RF-ABC posterior probability of choosing the Afr2Pulses-Eur2Pulses=0.6154), São Nicolau (post-prob.=0.5114), and Santo Antão (post-prob.=0.5388). While RF votes results also favor Afr2Pulses-Eur2Pulses scenarios for individuals born in Fogo, Brava and Maio, they do so less clearly with higher votes for alternative scenarios and weaker posterior probabilities (post- prob.=0.4632, 0.4152, and 0.4189, respectively). For Boa Vista and Sal, the Afr2Pulses-EurRecurring scenario is favored, while the AfrRecurring-Eur2Pulses scenario is favored for São Vicente. For São Vicente, the RF-ABC posterior probability is high (0.5151), while weaker for Boa Vista and Sal (0.3830 and 0.4910, respectively).

RF-ABC cross-validation prior-errors for each of the 40,000 simulations used, in-turn, as pseudo-observed data indicate a reasonably good, albeit not perfect, discriminatory power of the RF (**Figure SF23A**). We find a majority of correct scenario assignation for most scenarios and most islands and asymmetrical scenarios are the least confused with one-another (AfrRecurring-Eur2Pulses vs Afr2Pulses-EurRecurring, or Afr2Pulses-Eur2Pulses vs AfrRecurring-EurRecurring). These results show that scenario-nestedness induce, as expected a priori, increased assignation-errors in large parts of the parameter-space where scenarios are, by definition, biologically equivalent [37,54]. Finally (**Figure SF23B**), the mean, variance, skewness, kurtosis, minimum, and maximum of individual admixture proportions' distributions are the most informative summary-statistics for RF-ABC scenario-choice in every island, consistently with previous studies [37,55].

MetHis–Neural Network-ABC posterior parameter estimations

For each Cabo Verdean individuals from each island of birth separately, we performed NN-ABC joint posterior parameter estimation based on 100,000 *MetHis* simulations under the winning scenario. Posterior parameter estimation errors were based on NN-ABC cross-validations for the 1,000 simulations closest to the observed data considered, in turn, as pseudo-observed [37]. For each island separately, detailed posterior parameters' distributions, Credibility Intervals (CI), and cross-validation errors are provided in **Figure SF24-26** and **Tables ST5-13**. We synthesized our results considering median point-estimates and 50%-CI for each scenario parameter in the admixture history of each island in **Figure F7C-D**. We detailed our results and discussion for admixture history inferences for each island separately in **Supplementary Text,** in the light of historical data about the peopling of Cabo Verde (**Table ST2**).

Figure F7C shows that the reproductive population size of Cabo Verde islands remained very low until a strong increase in the last three generations for all islands but Santiago and Brava. In Santiago the population expansion was more linear since the founding of Cabo Verde in the 1460's, while the reproductive size of Brava remained low until today.

In brief summary, **Figure F7D** shows that European and African admixture events throughout the archipelago occurred first during the early peopling history of each island, before the mid-17th century massive expansion of the TAST due to the expansion of plantation economy [1]. We find that other admixture events from Europe or Africa, or both, likely occurred much later during, or immediately after, the 19th century abolition of the TAST and of slavery in European colonial empires. Altogether our *MetHis*-ABC results support

limited historical admixture having occurred in Cabo Verde during the most intense periods of the TAST
between the mid-17th and early 19th centuries.
Furthermore, note that we find admixture events often earlier than, or concomitant with, the first peren-
nial peopling of an island. For the islands of Santiago, Fogo, Santo Antão and, to a lesser extent, São Nicolau,
initial historical admixture events occurred synchronically to the first perennial settlement of the island. For
the islands of Brava, Maio, Boa Vista, and São Vicente, early admixture events occurred long before their
first perennial peopling, thus showing that their founding was already largely composed of admixed individ-
uals, likely originated from other Cabo Verde islands. Finally, our *MetHis*-NN-ABC posterior parameter
inferences cannot infer all scenario-parameters accurately, as some parameters hardly depart from their re-
spective priors (**Figure SF24-26**), and the admixture history of the island of Sal remains overall poorly in-
ferred.
420

DISCUSSION

Which African and European populations contributed genetically to Cabo Verde?

The genetic heritage of continental Africa to Cabo Verde

Numerous enslaved-African populations from Western, Central, and South-Western Central Africa have been forcibly deported during the TAST to both Cabo Verde and the Americas, as shown by historical demographic records [1,23]. There is still extensive debate about whether enslaved-Africans remained or more briefly transited in Cabo Verde during the most intense period of the TAST, in the 18th and 19th centuries, when the archipelago served as a slave-trade platform between continents [23,56,57]; a question also of major interest throughout the Americas [18,35].

In this context, our genetic results favor scenarios where mostly certain Senegambian populations (Mandinka and Wolof in our study) contribute to the genetic makeup of Cabo Verde (**Figures F2** and **F3**). Other Western, Central, and South-Western African populations historically deported during the TAST seem to have had only very limited contributions to the genomic diversity of Cabo Verde, and only in certain islands. This could be due to Cabo Verde being only a temporarily waypoint for these latter enslaved-African populations between Africa, the Americas, and Europe, but would also be consistent with additional socio-historical processes (see below). Interestingly, and further echoing these genetic results, the Cabo Verdean Kriolu language carries specific signatures mainly from the Mande language-family, and Wolof and Temne languages from Western Africa, and largely more limited signatures of Kikongo and Kimbundu Bantu languages from Central and South-Western Africa [15,28].

These results contrast with the admixture patterns identified in other enslaved-African descendant populations in the Americas in our dataset (African American and Barbadian, **Figures F2 and F3**), and in previous studies [3,4,10,36,58]. Indeed, the origins of African ancestries in numerous populations throughout the Caribbean and the Americas traced to varied continental African regions, from Western to South-Western Africa, thus qualitatively consistent with the known diversity of slave-trade routes used between continents and within the Americas after the Middle Passage.

The genetic heritage of continental Europe to Cabo Verde

After the initial settlement of Cabo Verde by Portuguese migrants, temporary changes of European dominion in certain islands, newly developing commercial opportunities, and intense piracy during the 16th and 17th centuries have triggered different European populations to migrate to the archipelago [24,25,34]. Nevertheless, these latter historical events do not seem to have left a major signature in the genetic landscape of Cabo Verde (**Figures F2 and F3**). Instead, we find that Cabo Verdean-born individuals in our dataset share virtually all their European genetic ancestry with Iberian populations, with extremely limited evidence of contributions from other European regions, consistent with previous studies [3,21]. Interestingly, the reduced diversity of European populations' contributions to the genomic landscape of Cabo Verde is also identified in other enslaved-African descendant populations in our study, as well as in previous studies in Caribbean and American populations [11,12,36]. Our results thus show that European admixture in enslaved-African descendant populations on both sides of the Atlantic mainly stem from the gene-pool of the European empires historically and socio-economically dominant locally, rather than subsequent European migrations [2].

Genetic and linguistic isolation-by-distance and difficult demographic expansion in Cabo Verde

A scenario of island peopling via series of founding events followed by slow-growing population sizes and relative isolation would consistently explain the increasing differentiation of island-specific patterns with increasing values of *K* found with ADMIXTURE, ASD-MDS and isolation-by-distance results (**Figures F2-F5**, **Table T2**), as well as *MetHis*-ABC demographic inferences (**Figure F7**).

Indeed, *MetHis*-ABC results (**Figure F7C**) show the long period of small reproductive sizes until very recent increases in most Cabo Verdean islands; with the notable exception of Santiago, the political and commercial center of Cabo Verde throughout the colonial history of the archipelago. This result was expected as the dry Sahelian climate of Cabo Verde with scarcely accessible water resources, recurring famines and epidemic outbreaks, and the Portuguese crown maintaining a strong control over population movements within Cabo Verde, rendered difficult the perennial peopling of most islands (see **Table ST2** and references therein, **Supplementary Text**).

Furthermore, such demographic scenario is also consistent with long-ROH patterns reflecting isolation on each Cabo Verdean islands, whereas elevated shorter ancestry-specific ROH patterns likely stemmed from admixture (**Figure F4**), similarly to our results in ASW and ACB populations and as previously identified [50]. Interestingly, we also found substantial signals of isolation-by-distance among Kriolu idiolects (i.e. individual manners of uttering Cabo Verdean Kriolu), also anchored in individual birth-places (**Figure F5**, **Table T2**), thus showing striking parallels between the history of biological and cultural isolation in Cabo Verde at a micro-geographical scale.

Importantly, we considered only random matting processes in our inferences and interpretations, albeit our ROH results together with recent work [59], may suggest that non-random matting processes significantly influenced Cabo Verdean genetic patterns. Therefore, future studies will need to evaluate how possible deviations from random-mating in Cabo Verde, such as assortative mating [60], may have influenced our results and interpretations. Note that this is a conceptually particularly challenging task in a small census-size population with strong marital stratification where mate-choices have been, by definition, limited during most of the TAST [61]. Nevertheless, such complex processes may also underlie the correlations found here between linguistic and genetic admixture patterns, and inter-island differences in both genetic and linguistic admixture levels (**Figure F5**); and would explain that genetic and linguistic histories of admixture apparently occurred in parallel in Cabo Verde.

Histories of genetic admixture in Cabo Verde
Early admixture in Cabo Verde, limited admixture during the plantation economy era
While we expected recurring African admixture processes due to the known history of regular forced migrations from Africa during most of Cabo Verde history [23], our *MetHis*–ABC scenario-choices indicate that, qualitatively, these demographic migrations did not necessarily translate into recurring gene-flow processes to shape genetic patterns in most of Cabo Verde islands (**Figure F7B**). Indeed, African admixture processes in all islands, except São Vicente, seem to have occurred during much more specific periods of Cabo Verdean history. Our *MetHis*–ABC posterior parameter estimations further highlighted often largely differing admixture histories across Cabo Verde islands (**Figure F7** and **Supplementary Text**).

We find that admixture from continental Europe and Africa occurred first early during the TAST history, concomitantly with the successive settlement of each Cabo Verdean island between the 15th and the early 17th centuries (**Figure F7D**). Furthermore, we find that the most intense period of enslaved-African deportations during the TAST via Cabo Verde, between the mid-17th and early-19th centuries during the expansion of the plantation economic system in the Americas and Africa [1,23–25], seem to have left a limited signature in the admixture patterns of most Cabo Verdean islands today. Interestingly, previous studies also highlighted that African admixture in enslaved-African descendants in North America and the Caribbean may have been limited during the height of the plantation economy era [3,12]. Therefore, our results illustrate the apparent discrepancy between intense demographic forced migrations during the TAST and genetic admixture signatures on both sides of the Atlantic.

Historians reported, in Cabo Verde and other European colonies in the Americas, that relationships between enslaved and non-enslaved communities largely changed with the expansion of plantation economy at the end of the 17th century. These changes are often referred to as the shift from Societies-with-Slaves to

Slave-Societies [1,19,62]. Slave-Societies legally enforced the socio-marital and economic segregation between
communities, and coercively controlled relationships between novel enslaved-African migrants and pre-
existing enslaved-African descendant communities more systematically and violently than Societies-with-
Slaves [19-p.15-46,95-108,23-p.281-319]. The high prevalence of segregation during the height of plantation economy
could have limited genetic admixture between enslaved-African descendants and non-enslaved communities
of European origin, as well as admixture between new migrants, forced or voluntary, and pre-existing Cabo
Verdeans, notwithstanding the known history of dramatic sexual abuses during and before this era. This
could consistently explain our observations of a relative lack of diverse African or European origins in Cabo
Verdean genomes despite the known geographical diversity of populations deported and emigrated to the
archipelago throughout the TAST. Furthermore, with legally enforced segregation, we might expect more
marital pairings than before to occur among individuals with common origins; i.e. between two individuals
with the same continental African or European origin. Such ancestry-specific marriages triggered by socio-
cultural segregation would be consistent with our ROH patterns (**Figure F4**).
In this context, the diversified African ancestries here found in the Americas (**Figure F2** and **F3**), con-
sistently with previous studies also showing relatively limited admixture events during the height of planta-
tion economy in the Americas [3,4,12], would suggest that the gene-pool of enslaved-Africans communities
admixing with Europeans in the Americas during the 16th and early 17th centuries already often involved, at a
local scale, multiple African source populations, thus reflecting the multiple slave-trade routes between con-
tinents. Conversely in Cabo Verde, the early onset of the TAST during the 15th century likely privileged
commercial routes with nearby Senegambia [23-p.31-54,281-319]), thus favoring almost exclusively admixture
events with individuals from this region. Altogether, our results in Cabo Verde contrasting with other en-
slaved-African descendant populations in the Americas, highlight the importance of early admixture process-
es and socio-cultural constraints changes on intermarriages throughout the TAST history, which likely dura-
bly influenced genetic diversities in descendant populations locally, on both sides of the Atlantic.
538
Admixture in Cabo Verde after the abolition of the TAST and of Slavery
Finally, we find that recent European and African admixture in Cabo Verde occurred mainly during the
complex historical transition after the abolition of the TAST in European colonial empires in the 1800's and
the subsequent abolition of slavery between the 1830's and the 1870's (**Figure F7D**). Historians showed that
these major historical shifts strongly disrupted pre-existing segregation systems between enslaved and non-
enslaved communities [1-p.271-290,23-p.335-362,63]. In addition, illegal slave-trade has flourished during this era,
bringing numerous African deportees to Cabo Verde outside of the official routes [23-p.363-384]. Altogether, our
results indicate increased signals of European and African admixture events in almost every island of Cabo
Verde during this period, and were thus consistent with a change of the social constraints regarding admix-
ture having prevailed over the preceding 200 years of the TAST. These results were largely consistent with
previous studies elsewhere in the Caribbean [11,64], and Central and South America [4,36]; showing, at a local
scale, the major influence of this recent and global socio-historical change in inter-community relationships
in European colonial empires on either sides of the ocean.
552

Perspectives

Altogether, our results highlight both the unity and diversity of the genetic peopling and admixture histories
of Cabo Verde islands, the first colony peopled by Europeans in Sub-Saharan Africa, resulting from the so-
ciocultural and historical complexity of the Trans-Atlantic Slave Trade and European colonial expansions
since the 15th century. Our results obtained at a micro-geographical scale reveal the fundamental importance
of the early TAST history, before the expansion of plantation economy, in durably shaping the genetic and
cultural diversity of enslaved-African descendant populations in both Africa and the Americas.

Importantly, we considered only the genome-wide autosomal admixture history of Cabo Verde in this
study, and therefore did not explore possible sex-biased admixture processes. However, previous studies
demonstrated the strong sex-biased admixture processes involved in Cabo Verde using sex-specific genetic
markers [30,31], similarly as in most enslaved-African descendant populations in the Americas [2]. Furthermore,
previous theoretical work considering sex-specific mechanistic admixture models showed that sex-biased
admixture processes may possibly bias historical inferences based only on autosomal data [65]. It will thus be
imperative, in future studies, to explore how this sex-biased admixture history may have influenced the ABC
inferences here conducted; for instance, via sex-specific developments of *MetHis*-ABC.
Our novel results together with their methodological limitations massively beg for future work further
complexifying the genetic admixture models here considered, as well as incorporating other summary-
statistics such as admixture-LD and sex-specific statistics. This will allow to further dissect the admixture
processes having given birth to enslaved-African descendant populations, on both sides of the Atlantic.
572
573

MATERIALS AND METHODS

1. Cabo Verde genetic and linguistic datasets

We conducted joint sampling of anthropological, genetic, and linguistic data in Cabo Verde with the only inclusion criteria that volunteer-participants would be healthy adults with Cabo Verdean citizenship and self-reporting speaking Kriolu [21]. Between 2010 and 2018, six interdisciplinary sampling-trips were conducted to interview 261 participants from more than thirty interview-locations throughout the archipelago (**Figure F1**).

1.a. Familial anthropology and geography data

The 261 Cabo Verdean individuals were each interviewed to record a classical familial anthropology questionnaire informing self-reported life-history mobility. In particular, we recorded primary residence location, birth location, parental and grand-parental birth and residence locations, and history of islands visited in Cabo Verde and foreign experiences [21]. Furthermore, we also recorded age, sex, marital status, and cumulated years of schooling and higher education (for 211 individuals only), and languages known and their contexts of use.

GPS coordinates for each reported location was acquired on site during field-work, supplemented by paper maps and Google Earth[TM] for foreign locations and islands that we did not visit (Sal and Maio). While participants' birth-locations were often precise, increasing levels of uncertainty arose for the reported parental and grand-parental locations. We arbitrarily assigned the GPS coordinates of the main population center of an island when only the island of birth or residence could be assessed with some certainty by participants. All other uncertain locations where recorded as missing data. Figure maps were designed with the software *QuantumGIS* v3.10 "Bucureşti" and using Natural Earth free vector and raster map data (https://www.naturalearthdata.com).

1.b. Genome-wide genotyping data

The 261 participants each provided 2mL saliva samples collected with DNAGenotek[TM] OG-250 kits, and complete DNA was extracted following manufacturer's recommendation. DNA samples were genotyped using an Illumina HumanOmni2.5Million-BeadChip genotyping array following manufacturer's instructions. We followed the quality-control procedures for genotypic data curation using Illumina's GenomeStudio® Genotyping Module v2.0 described in [21]. Genotype-calling and population-level quality-controls, and merging procedures are detailed in **Figure SF1**.

In summary, we extracted a preliminary dataset of 259 Cabo Verdean Kriolu-speaking individuals, including relatives, genotyped at 2,118,835 polymorphic di-allelic autosomal SNPs genome-wide. We then merged this dataset with 2,504 worldwide samples from the 1000 Genomes Project Phase 3 [66]; with 1,307 continental African samples from the African Genome Variation Project [67] (EGA accession number EGAD00001000959); and with 1,235 African samples [7] (EGA accession number EGAS00001002078). We retained only autosomal SNPs common to all data sets, and excluded one individual for each pair of individuals related at the 2nd degree (at least one grand-parent in common) as inferred with KING [68], following previous procedure [21].

After merging all datasets (**Figure SF1**), we considered a final working-dataset of 5,157 worldwide unrelated samples, including 233 unrelated Cabo Verdean Kriolu-speaking individuals of which 225 were Cabo-Verde-born, genotyped at 455,705 autosomal bi-allelic SNPs (**Figure F1**; **Table ST1**).

1.c. Individual utterances of Kriolu

We collected linguistic data for each of the 261 Cabo Verdean individuals using anthropological linguistics questionnaires, and semi-directed interviews. Each participant was shown a brief (~6 min) speech-less movie, *"The Pear Story"* [69], after which they were asked to narrate the story as they wanted in "the Kriolu they

speak every day at home". Discourses were fully recorded without interruption, whether individuals' discourses were related to the movie or not. Then each discourse was separately fully transcribed using the orthographic convention of Cabo Verdean Kriolu official alphabet "Alfabeto Unificado para a Escrita da Língua Cabo-verdiana (ALUPEC)" (Decreto-Lei nº 67/98, 31 de Dezembro 1998, I Série nº 48, Sup. B. O. da República de Cabo Verde).

Building on Verdu *et al.* (2017) approach, we were interested in inter-individual variation of "ways of speaking Kriolu" rather than in a prescriptivist approach to the Kriolu language. Thus, we considered each utterance as defined in [70-p.107]: "a particular instance of actually-occurring language as it is pronounced, grammatically structured, and semantically interpreted in its context". Transcripts were parsed together and revealed 4831 (L=4831) unique uttered items among the 225 discourses from the genetically-unrelated Cabo Verde-born individuals, for 92,432 uttered items in total. To obtain these counts, we considered phonetic, morphological, and syntactic variation of the same lexical root items that were uttered/pronounced differently, and we excluded from the utterance-counts onomatopoeia, interjections, and names. Note that we were here interested in the diversity of realizations in the Kriolu lexicon, including within the same individuals. In other words, we are interested in both between-speaker and within-speaker variation. Also note that a very limited number of English words were pronounced by particular individuals (10 utterances each occurring only once), and were kept in utterance-counts.

1.d. Individual Kriolu utterance frequencies

The list of unique uttered items was then used to compute individual's specific vectors of uttered items' relative frequencies as, for each genetically unrelated Cabo Verde-born individual i (in [1, 225]) and each unique uttered item l (in [1, L=4831]), $f_{i,l} = n_{i,l} / \sum_{j=1}^{L} n_{i,j}$, where $n_{i,l}$ is the absolute number of times individual i uttered the unique item l over her/his entire discourse, f_i being the vector $(f_{i,1}, f_{i,2}, \ldots, f_{i,L})$. We compared vectors of individuals' utterance-frequencies by computing the inter-individual pairwise Euclidean distance matrix as, for all pairs of individuals i and j, $d(f_i, f_j) = \sqrt{\sum_{l=1}^{L} (f_{i,l} - f_{j,l})^2}$ [21].

1.e. African origin of Kriolu utterances

We categorized each of the 4831 unique uttered items separately into five linguistic categories [21]. Category A included only unique utterances directly tracing to a known African language and comprised 88 unique items occurring 3803 times out of the 92,432 utterances. Category B included only items with a dual African and European etymology, attesting to the intense linguistic contacts at the origins of Cabo Verdean Kriolu, and comprised 254 unique items occurring 6960 times. Category C included 4432 items (occurring 73,799 times) with strictly Portuguese origin and not carrying identifiable traces of significant African linguistic origin or influence. Category D included 26 items occurring 6762 times with possible traces of African languages' phonetic or morphologic influences, albeit without certainty. Finally, Category U included the 10 English utterances occurring 10 times and the 21 unique utterances occurring 1089 times which could not be traced to African or European languages.

Following [21], we defined an "African-utterances score" based, conservatively, on the utterance frequencies obtained for items in Categories A and/or B as, for individual i and the set of utterances in each category denoted *Cat* (in [A; B; A&B]), $Z_{i,Cat} = \sum_{l=1}^{L_{Cat}} f_{i,l}$, with L_{Cat} the number of uttered items in the corresponding category, and $f_{i,l}$ defined as previously.

2. Population genetics descriptions

2.a. Allele Sharing Dissimilarities, Multidimensional Scaling, and ASD-MDS admixture estimates

We calculated pairwise Allele Sharing Dissimilarities [38] using the *asd* software (v1.1.0a;
https://github.com/szpiech/asd), among the 5,157 individuals in our worldwide dataset (**Figure F1**; **Table**
ST1), using 455,705 autosomal SNPs, considering, for a given pair of individuals, only the SNPs with no
missing data. We then projected this dissimilarity matrix in three dimensions with metric Multidimensional
Scaling using the *cmdscale* function in R [71]. We conducted successive MDS analyses on different individual
subsets of the original ASD matrix, by excluding, in turn, groups of individuals and recomputing each MDS
separately. Lists of populations included in each analysis can be found in **Table ST1**. 3D MDS animated
plots in *gif* format (**Figures SF2-6**) were obtained with the *plot3d* and *movie3d* functions from the R packag-
es *rgl* and *magick*.

Recently admixed individuals are intuitively expected to be at intermediate distances between the clus-
ters formed by their putative proxy source populations on ASD-MDS two-dimensional plots. A putative es-
timate of individual admixture rates can then be obtained by projecting admixed individuals orthogonally on
the line joining the respective centroids of each proxy-source populations and, then, calculating the distance
between the projected points and either centroid, scaled by the distance between the two centroids (**Figure**
SF12; [21,37]). We estimated such putative individual admixture rates in Cabo Verdean, ASW, and ACB indi-
viduals, considering sets of individuals for the putative proxy-source populations identified visually and re-
sulting from our ASD-MDS decomposition (**Supplementary Text**).

2.b. ADMIXTURE-CLUMPP-DISTRUCT analyses

Based on ASD-MDS explorations, we focused on the genetic structure of individuals born in Cabo Verde
compared to that of other admixed populations related to TAST migrations. Therefore, we conducted AD-
MIXTURE analyses [39] using 1,100 individuals from 22 populations: four from Europe, 14 from Africa, the
USA-CEU, the African-American ASW, the Barbadian-ACB populations, and the North Chinese-CHB pop-
ulation as an outgroup (**Table ST1**). To limit clustering biases due to uneven sampling, we randomly
resampled 50 individuals, once, for each one of these 22 populations. Furthermore, we also included all 44
Angolan individuals from four populations in the analyses, as no other samples from the region were availa-
ble in our dataset. In addition to the 1,100 individuals hence obtained, we included the 225 Cabo Verde-born
individuals.

Following constructor recommendations, we pruned the initial 455,705 SNPs set for low LD using *plink*
[72] function *--indep-pairwise* for a 50 SNPs-window moving in increments of 10 SNPs and a r^2 threshold of
0.1. We thus conducted all subsequent ADMIXTURE analyses considering 1,369 individuals genotyped at
102,543 independent autosomal SNPs.

We performed 30 independent runs of ADMIXTURE for values of K ranging from 2 to 10. For each
value of K separately, we identified groups of runs providing highly similar results (ADMIXTURE "modes"),
with Symmetric Similarity Coefficient strictly above 99.9% for all pairs of runs within a mode, using
CLUMPP [73]. We plotted each modal result comprising two ADMIXTURE runs or more, for each value of K
separately, using DISTRUCT [74]. Finally, we evaluated within-population variance of individual membership
proportions as F_{ST}/F_{ST}^{max} values using FSTruct [40] with 1000 Bootstrap replicates per population, for the AD-
MIXTURE mode result at $K=2$ (**Figure SF7**).

2.c. Local-ancestry inferences

To identify all populations sharing a likely common ancestry with the Cabo Verdean, ASW or ACB individ-
uals in our dataset using local-ancestry haplotypic inferences, we considered the same sample-set as for the
ADMIXTURE analysis (**Table ST1**), including all 455,705 SNPs from the merged dataset.

Phasing with ShapeIT2

We first phased individual genotypes using SHAPEIT2 [46] for the 22 autosomal chromosomes separately
using the joint Hap Map Phase 3 Build GRCh38 genetic recombination map [75]. We considered default pa-
rameters using phasing windows of 2Mb and 100 states per SNP. We considered by default 7 burn-in itera-
tions, 8 pruning iterations, 20 main iterations, and missing SNPs were imputed as monomorphic. Finally, we
considered a "Ne" parameter of 30,000, and all individuals were considered unrelated.
717
Chromosome painting with ChromoPainter2
We determined the possible origins of each Cabo Verdean, ASW, and ACB individual phased haplotypes
among European, African, USA-CEU, and Chinese-CHB populations using CHROMOPAINTER v2 [45] with
the same recombination map used for phasing. Following authors' recommendations, we conducted a first
set of 10 replicated analyses on a random subset of 10% of the individuals for chromosomes 2, 5, 12, and 19,
which provided a posteriori Ne=233.933 and θ=0.0004755376, on average by chromosome weighted by
chromosome sizes, to be used in the subsequent analysis. Then, we conducted a full CHROMOPAINTER
analysis using these parameters to paint all individuals in our dataset, in turn set as Donor and Recipient.
Finally, we combined painted chromosomes for each individual in the Cabo Verdean, ASW, and ACB popu-
lation, separately.
728
Estimating possible source populations for the Cabo Verde gene-pool using SOURCEFIND
We used CHROMOPAINTER results aggregated for each Cabo Verdean, ACB, and ASW individual sepa-
rately and conducted two SOURCEFIND [47] analyses using all other populations in the dataset as a possible
source, separately for four or six possible source populations ("surrogates"). We considered 400,000 MCMC
iteration steps (including 100,000 burn-in) and kept only one MCMC step every 10,000 steps for final likeli-
hood estimation. Each individual genome was divided in 100 slots with possibly different ancestry, with an
expected number of surrogates equal to 0.5 times the number of surrogates, for each SOURCEFIND analysis.
We aggregated results obtained for all individuals in the ACB, ASW, and each Cabo Verdean birth-island,
separately. We present the highest likelihood results across 20 separate iterations of the SOURCEFIND
analysis in **Figure F3B.** The second-best results were highly consistent and thus not shown.
739
740 **3. Genetic and linguistic diversities, and admixture within Cabo Verde**
741
3.a. Runs of Homozygosity (ROH)
Considering the same sample and SNP set as in the above local-ancestry analyses (**Table ST1**), we called
ROH with GARLIC [76]. For each population separately, we ran GARLIC using the weighted logarithm of the
odds (wLOD) score-computation scheme, with a genotyping-error rate of 1e^{-3} (a likely overestimate), and
using the same recombination map as for phasing, window sizes ranging from 30 to 90 SNPs in increment of
10 SNPs, 100 resampling to estimate allele frequencies, and all other GARLIC parameters set to default
values. We only considered results obtained with a window size of 40 SNPs, which was the largest windows
size associated with a bimodal wLOD score distribution and a wLOD score cutoff between the two modes,
for all populations.
For each populations and Cabo Verdean islands separately, we considered three classes of ROH that
correspond to the approximate time to the most recent common ancestor of the IBD haplotypes, which can
be estimated from the equation g=100/2L, where L is the ROH length in cM and g is the number of
generations to the most recent common ancestor of the haplotypes [48]. Short ROH are less than 0.25cM
reflecting homozygosity of haplotypes from more than 200 generations ago; medium ROH are between
0.25cM and 1cM reflecting demographic events having occurred between approximately 200 and 50
generations ago; and long ROH are longer than 1cM reflecting haplotypes with a recent common ancestor
less than 50 generations ago. In **Figure F4A-B** and **Figure SF9**, we plot the distribution of the summed

length of ROH per individual per size-classes.

3.b. Intersecting ROH and local ancestry painting

Using the same phasing results as described above, we conducted 10 EM iterations of the RFMIX2 [77] algo-
rithm to assign, for each Cabo Verdean individual and each SNP on each chromosome, separately, it's puta-
tive source population of origin among the 24 African, European, Chinese-CHB, and USA-CEU populations.
We collapsed the local ancestry assignments for each SNP in each Cabo Verdean individuals hence obtained
into three continental regions, representing broadly African, broadly European, and broadly Asian ancestries
respectively. We considered an approach similar to previous work [50], and intersected local ancestry calls with
ROH calls (**Figure 4C-D**).

RFMIX2 only makes local ancestry calls at loci that are present in the dataset. Therefore, a gap of un-
classified ancestry exists where inferred ancestry switches between two successive genotyped loci as called
by RFMIX2. These gaps necessarily each contain an odd number of ancestry switch points (≥ 1) absent from
our marker set. Therefore, when computing the total ancestry content of an ROH that overlaps one of these
ancestry breakpoints, we assign half of the length of this gap to each ancestry classification, effectively ex-
tending each local ancestry segment to meet at the midpoint.

We then plotted the length distribution of long ROH for each island and we breakout the distributions
by ancestral haplotype background: those with only African ancestry, those with only European ancestry,
and those that span at least one ancestry breakpoint (**Figure F4C**). We excluded long ROH in Asian ancestry
segments from this and the following analyses, as we found overall very limited such ancestry in the samples
(**Figure SF10**).

Finally, we explored how total African/European ancestry in long ROH varies between islands. For
each individual we summed the total amount of each ancestry in long-ROH and plot the distributions across
islands (**Figure SF11**). High levels of a given ancestry in long ROH could stem from an overall high level of
that ancestry in that individual. Therefore, for each individual, we computed their global ancestry proportions
by summing up the total length of the genome inferred as a given ancestry and dividing by the length of the
genome. We then plotted (**Figure F4D**), the difference of an individual's long-ROH ancestry proportion and
their global ancestry proportion, for African and European ancestries separately. Values above zero indicated
that a given ancestry is overrepresented in long-ROH relative to genome-wide proportions of that ancestry.

3.c. Genetic and linguistic diversity within Cabo Verde

We explored genetic pairwise levels of differentiation calculated with ASD as previously, considering the
1,899,878 non-monomorphic SNPs within Cabo Verde obtained after QC Stage 3 (**Figure SF1**). We project-
ed the matrix for the 225 unrelated Cabo Verde-born individuals using metric MDS as above. Note that prun-
ing this data sets for very low LD using *plink* function *--indep-pairwise* for a 50 SNPs window moving in
increments of 10 SNPs and a r^2 threshold of 0.025 resulted in 85,425 SNPs for which the ASD matrix was
highly correlated with the one using all SNPs (Spearman ρ=0.8745, *p*<2.2x10^{-16}). In parallel, we described
the diversity of Kriolu idiolects ("individuals' manners of speaking Kriolu") among the 225 genetically unre-
lated Cabo Verde-born individuals by projecting, with metric MDS, the matrix of pairwise Euclidean dis-
tances between vectors of individuals' utterance-frequencies (**Material and Methods 1.d**), considering the
4831 unique uttered items.

Then, we conducted series of Mantel and partial Mantel correlation tests (Legendre and Legendre 1998),
using the *partial.mantel.test* function in the R package *ncf*, with Spearman correlation and 10,000 Mantel
permutations, to explore possible isolation-by-distance [51] patterns as well as correlations with other co-
variables of interest. We conducted correlation tests between either the pairwise Euclidean distances between
vectors of individuals' utterance-frequencies or genetic ASD separately, and individual pairwise matrices of *i)*
absolute age differences, *ii)* absolute differences in academic education duration, *iii)* geographic distances

between residence-locations (logarithmic scale), *iv)* between birth-locations, *v)* between mothers' birth-locations, *vi)* between fathers' birth-locations (**Table T2**, **Table ST14**). All pairwise geographic distances were calculated with the Haversine great circle distance formulation [78], taking 6,371 km for the radius of the Earth, using the *rdist.earth* function in the R package *fields*. Before computing logarithmic distances, we added 1km to all pairwise distances.

3.d. Genetic and linguistic admixture within Cabo Verde

We further explored isolation-by-distance patterns within Cabo Verde specifically for African genetic and linguistic individual admixture levels. We considered African genetic admixture levels estimated from AD-MIXTURE at K=2 or ASD-MDS approaches (**Material and Methods 2.a** and **Supplementary Text**), and individual African linguistic admixture as "African-utterances scores" $Z_{i,Cat}$ as defined in **Material and Methods 1.e.** for utterance lists contained in Categories A and/or B [21]. For genetic or linguistic admixture levels separately, we computed the pairwise matrix of individual absolute admixture levels differences, and conducted Mantel testing with the different geographical distance matrices as above. Finally, we compared African genetic and linguistic admixture scores using Spearman correlations, throughout Cabo Verde and within all birth-islands, separately.

4. Inferring genetic admixture histories in Cabo Verde with *MetHis*-ABC

We aimed at reconstructing the detailed genetic admixture history of each Cabo Verde island separately. To do so, we first design *MetHis* v1.0 [37] forward-in-time simulations of four competing complex admixture models. We then couple *MetHis* simulation and summary-statistics calculation tools with ABC Random-Forest scenario-choice implemented in the R package *abcrf* [53], followed by Neural-Network posterior parameter estimation with the R package *abc* [52], under the winning model for each island separately.

4.a. Four competing models of complex historical admixture for Cabo Verde Islands

Following our descriptive analyses results, we considered models with only one "European" and one "African" source population and each Cabo Verde island, separately, as the "Admixed" recipient population. This corresponds to the "two-source" version of the [55] general model also explored with *MetHis* in [37].

Foundation of the admixed population and duration of the admixture history

In all four scenarios (**Figure F6**; **Table T1**), we considered, forward-in-time, that the admixed population (Population H) is founded at generation 0, 21 generations before present, which roughly corresponds to the 15th century for an average generation time of 25 years and sampled individuals born before the 1990s and on average between the 1960's and the 1980's in our dataset. This is reasonable as historical records showed that Cabo Verde was likely un-inhabited before its initial colonial settlement established in the 1460s on Santiago (Albuquerque and Madeira Santos 1991). At generation 0, the admixed population is founded with a proportion $s_{Eur,0}$ of admixed individuals' parents originating from the European source and a proportion $s_{Afr,0}$ from the African source, with $s_{Eur,0} + s_{Afr,0}$=1. Note that simulations considering parameter-values of $s_{Afr,0}$, or $s_{Eur,0}$, close to 0 correspond to scenarios where the "admixed" population is initially founded by only one of the sources, and that genetic admixture *per se* may only occur at the following admixture event.

After the founding of Population H, we considered two different admixture scenarios for either source populations' contribution to the gene-pool of population H.

Two admixture pulses after foundation

For a given source population hereafter designated "Source" ("European" or "African" in our case), after founding at generation 0, we considered scenarios with two additional pulses of admixture ("Source"-

2Pulses models). The two pulses occur, respectively, at generation $t_{Source,p1}$ and $t_{Source,p2}$ in [1,20], with $t_{Source,p1}$
$\neq t_{Source,p2}$; and with intensity $s_{Source,tSource,p1}$ and $s_{Source,tSource,p2}$ in [0,1], respectively.

Note that simulations considering $t_{Source,p1=tSource,p2}$ + 1, and simulations with either $s_{Source,tSource,p1}$ or
$s_{Source,tSource,p2}$ close to 0, may strongly resemble those expected under scenarios with only one pulse of admix-
ture after the founding of Population H.

A period of recurring admixture after foundation
For a given Source population, after founding at generation 0, we considered scenarios with a possible period
of recurring admixture, where, during this period, admixture intensity followed a monotonically decreasing
trend ("Source"-Recurring models). The period of admixture occurs between times $t_{Source,t1}$ and $t_{Source,t2}$ in
[1,20] with $t_{Source,t2} \geq t_{Source,t1}$ + 1. The beginning of the admixture period at $t_{Source,t1}$ is associated with admix-
ture intensity $s_{Source,tSource,t1}$ in [0,1]. The end of the admixture period at $t_{Source,t2}$ is associated with intensity
$s_{Source,tSource,t2}$ in [0,1] such that $s_{Source,tSource,t1} \geq s_{Source,tSource,t2}$. In between, the admixture intensity values at each
generation of the admixture period are the discrete numerical solutions of a decreasing rectangular hyperbola
function of parameter u_{Source} in [0,1/2] [37].

Intuitively, a u parameter value close to 0 corresponds to a sharp pulse of admixture occurring at the be-
ginning of the admixture period of intensity $s_{Source,tSource,t1}$, followed at the next generation by constant recur-
ring admixture of intensity $s_{Source,tSource,t2}$ until the end of the admixture period. Alternatively, a u parameter
value close to ½ corresponds to a linearly decreasing admixture at each generation of the admixture period
from $s_{Source,tSource,t1}$ to $s_{Source,tSource,t2}$.

Note that in the limit when $s_{Source,tSource,t1} \sim s_{Source,tSource,t2}$, Recurring scenarios correspond to constant re-
curring admixture of that intensity. Furthermore, simulations with u and $s_{Source,tSource,t2}$ parameter values both
close to 0 correspond to scenarios with a single pulse of admixture from a given source after the founding
pulse, occurring at time $t_{Source,t1}$ and with intensity $s_{Source,tSource,t1}$.

We combined the 2Pulses and Recurring scenarios from either the African and European source populations
in order to produce four competing scenarios of admixture for the genetic history of Cabo Verde (**Figure F6**),
with the only constraint that at each generation g between 1 and 20, $s_{Afr,g} + s_{Eur,g}=1–h_g$, where h_g is the contri-
bution of the admixed population H to itself at the following generation in [0,1].

Reproductive population size in the admixed population
We considered, for the admixed population H, a reproductive population size of N_0 diploid individuals at
generation 0, with N_0 in [10,1000], and N_{20} in [100,100000] at generation 20, such that $N_0 < N_{20}$. In between
these two values, we considered for N_g values the discrete numerical solution of an increasing rectangular
hyperbola function of parameter u_N in [0,1/2] [37].

4.b. *MetHis*-ABC Random Forest scenario-choice and Neural Network posterior parameter estima-
tions
To reconstruct highly complex admixture histories in Cabo Verde using genetic data, we conducted machine-
learning Approximate Bayesian Computation inferences. They rely on simulations conducted with model-
parameter values drawn randomly within prior distributions set explicitly by the user, and based on sets of
summary-statistics calculated on each simulation and formally compared to the same statistics computed on
the observed data [79,80]. To do so, we conducted simulations using *MetHis* [37] forward-in-time simulator, fol-
lowed by Random-Forest ABC scenario-choice [53], and Neural Network ABC posterior parameter inferences
[52], for each Cabo Verde island separately. We followed the *MetHis*-ABC approach proposed in [37] for sum-
mary-statistics calculation, prior-checking, out-of-bag cross-validation, machine-learning ABC predictions
and inferences parameterization, and posterior parameter cross-validation error calculations.

For concision here, all the details of these procedures can be found in corresponding **Supplementary**
Text, **Tables ST3-13**, and **Figures SF13-26**. Briefly, we performed 10,000 simulations of 60,000 independ-
ent SNPs under each of the four competing scenarios described above, drawing parameter values in prior
distributions detailed in **Table T1**. We then computed 42 summary-statistics on each simulated dataset sepa-
rately, described in **Supplementary Text** and **Table ST3**, by drawing sample-sets matching the observed
sample-sizes of each Cabo Verde birth-island separately. Note that we considered summary-statistics specifi-
cally aiming at describing the distribution of individual admixture fractions in each island, known theoreti-
cally and empirically to be highly informative about complex admixture history parameters [37,55]. Based on
these vectors of summary-statistics, we conducted RF-ABC scenario-choice to identify, for each birth-island
separately, the most likely admixture history scenario. Under the winning scenario in each birth-island sepa-
rately, we performed an additional 90,000 simulations for further Neural Network ABC posterior parameter
estimations and cross-validation parameter error calculations, thus each based on 100,000 total simulations.
911
912

 FIGURE AND TABLE LEGENDS

Figure F1:

Sampling and birth place locations of 225 unrelated Cabo Verdean-born individuals, merged with 77

worldwide populations.

Linguistic and familial anthropology interviews, and genetic sampling for Cabo Verde participants were

conducted during six separate interdisciplinary fieldworks between 2010 and 2018. Further details about

populations are provided in **Table ST1**.

Figure F2:

Multidimensional Scaling projections of pairwise Allele Sharing Dissimilarities in Cabo Verdeans and

continental African and European populations.

A-C) 3-Dimensional MDS projection of ASD computed among 233 unrelated Cabo Verdeans and other

continental African and European populations using 445,705 autosomal SNPs. Cabo Verdean patterns in

panels A-C are compared to **D-F** the USA African-Americans ASW, **G-I** the Barbadians-ACB, or **J-L** the

Puerto Ricans-PUR in the same African and European contexts. We evaluate the Spearman correlation be-

tween 3D MDS projections and the original ASD matrix to evaluate the precision of the dimensionality re-

duction. We find significant ($p<2.2 \times 10^{-16}$) Spearman $\rho=0.9635$ for the Cabo Verde analysis (A-C); 0.9383

for the ASW (D-F); 0.9306 for the ACB (G-I); and 0.9437 for the PUR (J--L). Each individual is represented

by a single point. See **Table ST1** for the populations used in these analyses, and **Figure F1** for sample loca-

tions and symbols.

Figure F3:

Individual genetic structure and haplotypic local ancestry inference among Cabo Verdean, Barbadian-

ACB and African-American ASW populations.

A) Unsupervised ADMIXTURE analyses using randomly resampled individual sets for populations original-

ly containing more than 50 individuals (**Table ST1**). 225 unrelated Cabo Verdean-born individuals are com-

prised in the analysis and grouped by birth-island. Number of runs highly resembling one-another using

CLUMPP are indicated below each K-value. All other modes are presented in **Figure SF8**. **B)**

SOURCEFIND results for each eleven target admixed populations (ASW, ACB, each nine Cabo Verde

birth-island), considering respectively 4 or 6 possible source populations (surrogates) among the 24 possible

European, African, and Asian populations considered in the ADMIXTURE analyses. The cumulated average

African admixture levels in each admixed population was highly consistent between SOURCEFIND esti-

mates and ADMIXTURE results at $K=2$ (Spearman $\rho=0.99772$, $p<8 \times 10^{-12}$, and 0.98861, $p<2 \times 10^{-8}$, for 6 or 4

surrogates respectively). Furthermore, individual admixture levels estimated using an ASD-MDS based ap-

proach were highly consistent with estimates based on ADMIXTURE at $K=2$ ($\rho=0.99725$; $p<2.2 \times 10^{-16}$ for

Cabo Verde; $\rho=0.99568$; $p<2.2 \times 10^{-16}$ for ASW; $\rho=0.99366$; $p<2.2 \times 10^{-16}$ for ACB, **Supplementary Text**).

Figure F4:

Distributions of long ROH in Cabo Verde.

A) The distribution of total long-ROH per person for each Cabo Verdean birth-island and other populations.

B) The distribution of individual long-ROH lengths (log10-scale). **C)** The distribution of ancestry-specific

and ancestry-spanning individual long-ROH lengths for each Cabo Verdean birth-island. **D)** The distribution

of differences between individuals' long-ROH ancestry proportion and their global ancestry proportion, for
African and European ancestries separately and for each Cabo Verdean birth-island.
961
962
Figure F5:
Utterance and genetic diversity and admixture within Cabo Verde.
A-B) 3D MDS projection of Allele Sharing Dissimilarities computed among 225 unrelated Cabo-Verde-born
individuals using 1,899,878 autosomal SNPs. Three-dimensional Euclidean distances between pairs of indi-
viduals in this MDS significantly correlated with ASD (Spearman ρ=0.6863; p<2.2x10^{-16}). **C-D)** 3D MDS
projection of individual pairwise Euclidean distances between uttered linguistic items frequencies based on
the 4831 unique uttered items obtained from semi-spontaneous discourses. Three-dimensional Euclidean
distances between pairs of individuals in this MDS significantly correlated with the utterance-frequencies
distances (Spearman ρ=0.8647; p<2.2x10^{-16}). **E)** Spearman correlation between individual African utterance
scores and individual genetic African admixture rates obtained with ADMIXTURE at K=2. **F)** Birth-
locations of 225 individuals in Cabo Verde. Symbols for individuals' birth-island in panels **A-D** are shown in
panel **F**. Panel **A-D** were procrusted according to individual actual birth-places' geographical locations in
panel **F** (Wang et al. 2010).
976
977
Figure F6:
Four competing scenarios for the genetic admixture history of each Cabo Verde island.
For all scenarios, the duration of the admixture process is set to 20 generations after the initial founding ad-
mixture event occurring at generation 0, which corresponds roughly to the initial peopling of Cabo Verde in
the 1460's, considering 25 years per generation and sampled individuals born on average between the 1960s
and 1980's.
After the initial founding pulse of admixture, *Scenario 1 Afr2Pulses-Eur2Pulses*: the admixed population
receives two separate introgression pulses from the African and European sources, respectively; *Scenario 2*
Afr2Pulses-EurRecurring: the admixed population receives two separate introgression pulses from the Afri-
can source, and a period of monotonically constant or decreasing recurring introgression from the European
source; *Scenario 3 AfrRecurring-Eur2Pulses*: the admixed population receives a period of monotonically
constant or decreasing recurring introgression from the African source, and two separate introgression pulses
from the European source; *Scenario 4 AfrRecurring-EurRecurring*: the admixed population receives a period
of monotonically constant or decreasing recurring introgression from the African source, and, separately, a
period of monotonically constant or decreasing recurring introgression from the European source. For all
scenarios, we consider the same demographic models with a possible increase in reproductive population
size N_g between the founding event and present. See **Materials and Methods** for detailed descriptions of
model-parameters.
996
997
Figure F7:
Genetic admixture histories of Cabo Verde islands.
Elements of the peopling-history of Cabo Verde islands are synthesized in **Table ST2**, stemming from his-
torical work. Islands are ordered from top to bottom in the chronological order for the first historical census
perennially above 100 individuals within an island, indicated with the purple dashed-lines. First historical
record of the settlement of an island as reported in political and administrative historical documents, tax-
records, and/or parishes establishments, are indicated with the red dashed lines.

A) Within-island birth-places of 225 Cabo-Verde-born individuals. **B)** *MetHis*-Random Forest-ABC scenar-
io-choice vote results for each island separately. Posterior probabilities of correctly identifying a scenario if
correct are indicated for the winning scenario. **C)** *MetHis*-Neural Network-ABC posterior parameter distribu-
tions with 50% Credibility Intervals for the reproductive population size history of each birth-island sepa-
rately. **D)** Synthesis of *MetHis*-NN-ABC posterior parameter median point-estimates and associated 50%CI,
for the admixture history of each island under the winning scenario identified with RF-ABC. European ad-
mixture history appears in blue, African admixture history in orange. Horizontal bars indicate 50% CI for the
corresponding admixture time parameters, vertical arrows correspond to median admixture intensity esti-
mates.
For **C)** and **D)**, posterior parameter distributions showing limited departure from their respective priors and
large Credibility Intervals are grayed, as they were largely unidentifiable in our ABC procedures. Detailed
parameter posterior distributions, 95%CI, and cross-validation errors are provided in **Figure SF24-26** and
Tables ST5-13. Detailed results description for each island are provided in **Supplementary Text**.
All ABC inference rely on forward-in-time *MetHis* simulations over 21 generations of history. Individuals in
our sample set were born on average between the 1960's and the 1980's. We translated *MetHis* generations
in Gregorian calendar dates by considering an average generation time of 25 generations. We considered the
onset of the massive increase of TAST African-enslaved individuals' deportations concomitant to the expan-
sion of plantation economy in the 1630's [1,2]. We considered the abolition of the TAST starting around the
1810's, as the British empire abolition of the TAST in 1807 marked the onset of TAST abolition in all other
European empires over the following 30 years [1]. We considered the abolition of slavery in the 1885's, as
slavery was abolished in Portugal and it's colonial empire over the course of more than 20 years between
1856 and 1878, and further took several years to be fully enacted in all regions of the empire [23]. The inde-
pendence of Cabo Verde occurred in 1975.
1028
1029
1030

1031 **Table T1:**

1032 **Prior distributions for the parameters of four competing scenarios for the admixture history of Cabo**

1033 **Verde islands.**

1034 Parameters are presented in **Figure F6** and described in **Materials and Methods**.

1035

1036

Description	Scenario	Model Parameter	Prior	Conditions
African admixture-pulse times	1, 2	$t_{Afr,p1}$	Uniform [1..20] in generations, a range corresponding to between ~1485 and ~1960 in years CE	$t_{Afr,p1} \neq t_{Afr,p2}$
		$t_{Afr,p2}$		
European admixture-pulse times	1, 3	$t_{Eur,p1}$	Uniform [1..20] in generations, a range corresponding to between ~1485 and ~1960 in years CE	$t_{Eur,p1} \neq t_{Eur,p2}$
		$t_{Eur,p2}$		
African admixture period start and end times	2, 4	$t_{Afr,t1}$	Uniform [1..20] in generations, a range corresponding to between ~1485 and ~1960 in years CE	$t_{Afr,t1} \neq t_{Afr,t2}$
		$t_{Afr,t2}$		
European admixture period start and end times	3, 4	$t_{Eur,t1}$	Uniform [1..20] in generations, a range corresponding to between ~1485 and ~1960 in years CE	$t_{Eur,t1} \neq t_{Eur,t2}$
		$t_{Eur,t2}$		
African admixture-pulse intensities	1, 2	$s_{Afr,tAfr,p1}$	Uniform [0,1]	
		$s_{Afr,tAfr,p2}$		$s_{Afr,g} + s_{Eur,g} = 1 - h_g$, with h_g in [0,1]
European admixture-pulse intensities	1, 3	$s_{Eur,tEur,p1}$	Uniform [0,1]	
		$s_{Eur,tEur,p2}$		
African admixture period intensity parameters	2, 4	$s_{Afr,tAfr,t1}$	Uniform [0,1]	$s_{Afr,tAfr,t1} \geq s_{Afr,tAfr,t2}$
		$s_{Afr,tAfr,t2}$	Uniform [0,1]	$s_{Afr,g} + s_{Eur,g} = 1 - h_g$, with h_g in [0,1]
		u_{Afr}	Uniform [0 , 0.5]	
European admixture period intensity parameters	3, 4	$s_{Eur,tEur,t1}$	Uniform [0,1]	$s_{Eur,tEur,t1} \geq s_{Eur,tEur,t2}$
		$s_{Eur,tEur,t2}$	Uniform [0,1]	$s_{Afr,g} + s_{Eur,g} = 1 - h_g$, with h_g in [0,1]
		u_{Eur}	Uniform [0 , 0.5]	
Founding reproductive population size	1, 2, 3, 4	N_0	Uniform [10 , 1000]	
Current reproductive population size	1, 2, 3, 4	N_{20}	Uniform [100 , 100000]	$N_0 < N_{20}$
Steepness of the reproductive population size increase	1, 2, 3, 4	u_N	Uniform [0 , 0.5]	

1037
1038
1039

Table T2:

Mantel and partial-Mantel correlations between utterance frequency differences and covariables, and

between genetic ASD and the same covariables, in 225 genetically unrelated Cabo Verde-born Kriolu-

speaking individuals.

Spearman correlation ρ are indicated in bold when significant after Bonferroni correction for multiple-testing

($\alpha<1\%$), and in italics when not significant after such correction. Spearman correlations and Mantel-tests

among covariables are provided in **Table ST14**.

Mantel variable	Partial-Mantel control	n	Geographic scale	Utterance-Frequency Euclidean Distance - 4831 uttered items		Genetic ASD - 1,899,978 SNPs	
				Spearman rho	10000 Mantel two-sided permutation p	Spearman rho	10000 Mantel two-sided permutation p
abs(Age difference)	--	225	within and between islands	**0.2215**	**< 2.10-4**	**0.1303**	**< 2.10-4**
abs(Age difference)	log(Birth-loc. dist.)	225	within and between islands	**0.2294**	**< 2.10-4**	**0.1348**	**< 2.10-4**
log(Birth-loc. dist.)	--	225	within and between islands	**0.2794**	**< 2.10-4**	**0.2916**	**< 2.10-4**
log(Birth-loc. dist.)	abs(Age difference)	225	within and between islands	**0.2855**	**< 2.10-4**	**0.2935**	**< 2.10-4**
abs(Education duration difference)	--	186	within and between islands	*0.0962*	*0.0024*	*0.0168*	*0.2730*
abs(Education duration difference)	log(Birth-loc. dist.)	186	within and between islands	*0.0834*	*0.0071*	*-0.0023*	*0.4900*
abs(Education duration difference)	--	185	within and between islands	*0.1001*	*0.0014*	*0.0159*	*0.2825*
abs(Education duration difference)	log(Residence dist.)	185	within and between islands	*0.0824*	*0.0068*	*-0.0041*	*0.4651*
log(Residence dist.)	--	224	within and between islands	**0.2145**	**< 2.10-4**	**0.1658**	**< 2.10-4**
log(Residence dist.)	log(Birth-loc. dist.)	224	within and between islands	*0.0306*	*0.0682*	**-0.0488**	**0.0005**
log(Birth-loc. dist.)	--	224	within and between islands	**0.2800**	**< 2.10-4**	**0.2889**	**< 2.10-4**
log(Birth-loc. dist.)	log(Residence dist.)	224	within and between islands	**0.1863**	**< 2.10-4**	**0.2445**	**< 2.10-4**
log(Father Birth-loc. dist.)	--	222	within and between islands	**0.1704**	**< 2.10-4**	**0.2424**	**< 2.10-4**
log(Father Birth-loc. dist.)	log(Birth-loc. dist.)	222	within and between islands	*0.0066*	*0.3915*	*0.0846*	*0.0014*
log(Mother Birth-loc. dist.)	--	224	within and between islands	**0.2634**	**< 2.10-4**	**0.2619**	**< 2.10-4**
log(Mother Birth-loc. dist.)	log(Birth-loc. dist.)	224	within and between islands	*0.0853*	*0.0071*	*0.0748*	*0.0057*
abs(Age difference)	--	225	within islands only	**0.2727**	**< 2.10-4**	**0.2124**	**0.0006**
abs(Age difference)	log(Birth-loc. dist.)	225	within islands only	**0.2546**	**< 2.10-4**	*0.1648*	*0.0041*
log(Birth-loc. dist.)	--	225	within islands only	*0.1412*	*0.0401*	**0.3460**	**< 2.10-4**
log(Birth-loc. dist.)	abs(Age difference)	225	within islands only	*0.0990*	*0.1030*	**0.3212**	**< 2.10-4**
abs(Education duration difference)	--	186	within islands only	*0.1287*	*0.0440*	*-0.0370*	*0.3077*
abs(Education duration difference)	log(Birth-loc. dist.)	186	within islands only	*0.1239*	*0.0496*	*-0.0537*	*0.2330*
abs(Education duration difference)	--	185	within islands only	*0.1421*	*0.0292*	*-0.0382*	*0.3037*
abs(Education duration difference)	log(Residence dist.)	185	within islands only	*0.1202*	*0.0546*	*-0.0491*	*0.2566*
log(Residence dist.)	--	224	within islands only	*0.0982*	*0.0911*	*-0.0667*	*0.1907*
log(Residence dist.)	log(Birth-loc. dist.)	224	within islands only	*0.1063*	*0.0704*	*-0.0549*	*0.2319*
log(Birth-loc. dist.)	--	224	within islands only	*0.1537*	*0.0282*	**0.3465**	**< 2.10-4**
log(Birth-loc. dist.)	log(Residence dist.)	224	within islands only	*0.1589*	*0.0230*	**0.3446**	**< 2.10-4**
log(Father Birth-loc. dist.)	--	222	within islands only	*0.0160*	*0.4123*	**0.2660**	**0.0006**
log(Father Birth-loc. dist.)	log(Birth-loc. dist.)	222	within islands only	*-0.0111*	*0.4546*	*0.2187*	*0.0045*
log(Mother Birth-loc. dist.)	--	224	within islands only	*0.1283*	*0.0423*	*0.2240*	*0.0034*
log(Mother Birth-loc. dist.)	log(Birth-loc. dist.)	224	within islands only	*0.1000*	*0.0925*	*0.1563*	*0.0303*

1047

REFERENCES

1. Eltis, D. & Richardson, D. *Atlas of the transatlantic slave trade*. (2015).
2. Fortes-Lima, C. & Verdu, P. Anthropological genetics perspectives on the transatlantic slave trade. *Human Molecular Genetics* **30**, R79–R87 (2021).
3. Micheletti, S. J. *et al.* Genetic Consequences of the Transatlantic Slave Trade in the Americas. *The American Journal of Human Genetics* **107**, 265–277 (2020).
4. Ongaro, L. *et al.* The Genomic Impact of European Colonization of the Americas. *Current Biology* **29**, 3974-3986.e4 (2019).
5. Winkler, C. A., Nelson, G. W. & Smith, M. W. Admixture Mapping Comes of Age. *Annu. Rev. Genom. Hum. Genet.* **11**, 65–89 (2010).
6. Wojcik, G. L. *et al.* Genetic analyses of diverse populations improves discovery for complex traits. *Nature* **570**, 514–518 (2019).
7. Patin, E. *et al.* Dispersals and genetic adaptation of Bantu-speaking populations in Africa and North America. *Science* **356**, 543–546 (2017).
8. Gravel, S. Population Genetics Models of Local Ancestry. *Genetics* **191**, 607–619 (2012).
9. Hellenthal, G. *et al.* A Genetic Atlas of Human Admixture History. *Science* **343**, 747–751 (2014).
10. Mathias, R. A. *et al.* A continuum of admixture in the Western Hemisphere revealed by the African Diaspora genome. *Nat Commun* **7**, 12522 (2016).
11. Moreno-Estrada, A. *et al.* Reconstructing the Population Genetic History of the Caribbean. *PLoS Genet* **9**, e1003925 (2013).
12. Baharian, S. *et al.* The Great Migration and African-American Genomic Diversity. *PLoS Genet* **12**, e1006059 (2016).
13. Holm, J. A. *An introduction to pidgin and creoles*. (Cambridge University Press, 2000).
14. Escure, G. & Schwegler, A. *Creoles, contact, and language change: linguistics and social implications*. (John Benjamins Pub, 2004).
15. Quint, N. *Le cap-verdien: origines et devenir d'une langue métisse*. (L'Harmattan, 2000).
16. Essegbey, J., Migge, B. & Winford, D. Cross-linguistic influence in language creation: Assessing the role of the Gbe languages in the formation of the Creoles of Suriname. *Lingua* **129**, 1–8 (2013).
17. Baptista, M. Continuum and variation in Creoles: Out of many voices, one language. *JPCL* **30**, 225–264 (2015).
18. Eltis, D. *Coerced and free migration: global perspectives*. (Stanford University Press, 2002).
19. Berlin, I. *Many Thousands Gone: the First Two Centuries of Slavery in North America*. (Harvard University Press, 2009).
20. Ansari-Pour, N. *et al.* Palenque de San Basilio in Colombia: genetic data support an oral history of a paternal ancestry in Congo. *Proc. R. Soc. B.* **283**, 20152980 (2016).
21. Verdu, P., Jewett, E. M., Pemberton, T. J., Rosenberg, N. A. & Baptista, M. Parallel Trajectories of Genetic and Linguistic Admixture in a Genetically Admixed Creole Population. *Current Biology* **27**, 2529-2535.e3 (2017).

22. Hagemeijer, T. & Rocha, J. Creole languages and genes: the case of São Tomé and Príncipe.
Faits Lang. **49**, 167–182 (2019).

23. Carreira, A. *Cabo Verde: formação e extinção de uma sociedade escravocrata (1460-1878).*
(IPC, 2000).

24. Albuquerque, L. de, Madeira Santos, M. E. *História geral de Cabo Verde T1.* vol. 1 (Centro de
Estudos de História e Cartografia Antiga, Instituto de Investigação Científica Tropical ;
Instituto Nacional da Cultura de Cabo Verde, 1991).

25. Albuquerque, L. de, Madeira Santos, M. E. *História geral de Cabo Verde T2.* vol. 2 (Centro de
Estudos de História e Cartografia Antiga, Instituto de Investigação Científica Tropical ;
Instituto Nacional da Cultura de Cabo Verde, 1995).

26. Albuquerque, L. de, Madeira Santos, M. E. *História concisa de Cabo Verde (Resumo da*
História geral de Cabo Verde T1-3). (Centro de Estudos de História e Cartografia Antiga,
Instituto de Investigação Científica Tropical ; Instituto Nacional da Cultura de Cabo Verde,
2007).

27. Baptista, M. *The syntax of Cape Verdean Creole: the Sotavento varieties.* (John Benjamins Pub,
2002).

28. Lang, J. *Les langues des autres dans la créolisation: théorie et exemplification par le créole*
d'empreinte wolof à l'île Santiago du Cap Vert. (Narr, 2009).

29. Brehm, A., Pereira, L., Bandelt, H.-J., Prata, M. J. & Amorim, A. Mitochondrial portrait of the
Cabo Verde archipelago: the Senegambian outpost of Atlantic slave trade. *Ann. Hum. Genet.* **66**,
49–60 (2002).

30. Gonçalves, R. *et al.* Y-chromosome lineages in Cabo Verde Islands witness the diverse
geographic origin of its first male settlers. *Human Genetics* **113**, 467–472 (2003).

31. Beleza, S. *et al.* The admixture structure and genetic variation of the archipelago of Cape Verde
and its implications for admixture mapping studies. *PLoS ONE* **7**, e51103 (2012).

32. Beleza, S. *et al.* Genetic architecture of skin and eye color in an African-European admixed
population. *PLoS Genet* **9**, e1003372 (2013).

33. Hamid, I., Korunes, K. L., Beleza, S. & Goldberg, A. Rapid adaptation to malaria facilitated by
admixture in the human population of Cabo Verde. *eLife* **10**, e63177 (2021).

34. Soares, M. J. The British Presence on the Cape Verdean Archipelago (Sixteenth to Eighteenth
centuries). *African Economic History* **39**, 129–146 (2011).

35. Berlin, I. *The making of African America: the four great migrations.* (Penguin Books, 2010).

36. Gouveia, M. H. *et al.* Origins, Admixture Dynamics, and Homogenization of the African Gene
Pool in the Americas. *Molecular Biology and Evolution* **37**, 1647–1656 (2020).

37. Fortes Lima, C. A., Laurent, R., Thouzeau, V., Toupance, B. & Verdu, P. Complex genetic
admixture histories reconstructed with Approximate Bayesian Computation. *Mol Ecol Resour*
21, 1098–1117 (2021).

38. Bowcock, A. M. *et al.* High resolution of human evolutionary trees with polymorphic
microsatellites. *Nature* **368**, 455–457 (1994).

39. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in
unrelated individuals. *Genome Research* **19**, 1655–1664 (2009).

40. Morrison, M. L., Alcala, N. & Rosenberg, N. A. *FSTruct: an FST-based tool for measuring*
ancestry variation in inference of population structure.
http://biorxiv.org/lookup/doi/10.1101/2021.09.24.461741 (2021)
doi:10.1101/2021.09.24.461741.

41. Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of population structure using multilocus
genotype data. *Genetics* **155**, 945–959 (2000).

42. Rosenberg, N. A. Genetic Structure of Human Populations. *Science* **298**, 2381–2385 (2002).

43. Falush, D., Stephens, M. & Pritchard, J. K. Inference of population structure using multilocus
genotype data: linked loci and correlated allele frequencies. *Genetics* **164**, 1567–1587 (2003).

44. Lawson, D. J., van Dorp, L. & Falush, D. A tutorial on how not to over-interpret STRUCTURE
and ADMIXTURE bar plots. *Nat Commun* **9**, 3258 (2018).

45. Lawson, D. J., Hellenthal, G., Myers, S. & Falush, D. Inference of Population Structure using
Dense Haplotype Data. *PLoS Genet* **8**, e1002453 (2012).

46. Delaneau, O., Marchini, J. & Zagury, J.-F. A linear complexity phasing method for thousands
of genomes. *Nat Methods* **9**, 179–181 (2012).

47. Chacón-Duque, J.-C. *et al.* Latin Americans show wide-spread Converso ancestry and imprint
of local Native ancestry on physical appearance. *Nat Commun* **9**, 5388 (2018).

48. Thompson, E. A. Identity by Descent: Variation in Meiosis, Across Genomes, and in
Populations. *Genetics* **194**, 301–326 (2013).

49. Mooney, J. A. *et al.* Understanding the Hidden Complexity of Latin American Population
Isolates. *The American Journal of Human Genetics* **103**, 707–726 (2018).

50. Szpiech, Z. A. *et al.* Ancestry-Dependent Enrichment of Deleterious Homozygotes in Runs of
Homozygosity. *The American Journal of Human Genetics* **105**, 747–762 (2019).

51. Rousset, F. Genetic Differentiation and Estimation of Gene Flow from F -Statistics Under
Isolation by Distance. *Genetics* **145**, 1219–1228 (1997).

52. Csilléry, K., François, O. & Blum, M. G. B. abc: an R package for approximate Bayesian
computation (ABC): *R package: abc*. *Methods in Ecology and Evolution* **3**, 475–479 (2012).

53. Pudlo, P. *et al.* Reliable ABC model choice via random forests. *Bioinformatics* **32**, 859–866
(2016).

54. Robert, C. P., Mengersen, K. & Chen, C. Model choice versus model criticism. *Proceedings of*
the National Academy of Sciences **107**, E5–E5 (2010).

55. Verdu, P. & Rosenberg, N. A. A General Mechanistic Model for Admixture Histories of Hybrid
Populations. *Genetics* **189**, 1413–1426 (2011).

56. Patterson, K. D. Epidemics, Famines, and Population in the Cape Verde Islands, 1580-1900.
The International Journal of African Historical Studies **21**, 291 (1988).

57. Brooks, G. E. Cabo Verde: Gulag of the South Atlantic: Racism, Fishing Prohibitions, and
Famines. *Hist. Afr.* **33**, 101–135 (2006).

58. Martin, A. R. *et al.* Human Demographic History Impacts Genetic Risk Prediction across
Diverse Populations. *The American Journal of Human Genetics* **100**, 635–649 (2017).

59. Korunes, K. L. *et al. Sex-biased admixture and assortative mating shape genetic variation and*
influence demographic inference in admixed Cabo Verdeans.

http://biorxiv.org/lookup/doi/10.1101/2020.12.14.422766 (2020)

doi:10.1101/2020.12.14.422766.

60. Zaitlen, N. *et al.* The Effects of Migration and Assortative Mating on Admixture Linkage

Disequilibrium. *Genetics* **205**, 375–383 (2017).

61. Versluys, T. M. M., Mas-Sandoval, A., Flintham, E. O. & Savolainen, V. Why do we pick

similar mates, or do we? *Biol. Lett.* **17**, 20210463 (2021).

62. Chaudenson, R. & Mufwene, S. S. *Creolization of language and culture*. (Routledge, 2001).

63. Cooper, F., Holt, T. C. & Scott, R. J. *Beyond slavery: explorations of race, labor, and*

citizenship in postemancipation societies. (University of North Carolina Press, 2000).

64. Fortes-Lima, C. *et al.* Exploring Cuba's population structure and demographic history using

genome-wide data. *Sci Rep* **8**, 11422 (2018).

65. Goldberg, A., Verdu, P. & Rosenberg, N. A. Autosomal Admixture Levels Are Informative

About Sex Bias in Admixed Populations. *Genetics* **198**, 1209–1229 (2014).

66. The 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature*

526, 68–74 (2015).

67. Gurdasani, D. *et al.* The African Genome Variation Project shapes medical genetics in Africa.

Nature **517**, 327–332 (2015).

68. Manichaikul, A. *et al.* Robust relationship inference in genome-wide association studies.

Bioinformatics **26**, 2867–2873 (2010).

69. Chafe, W. L. *The Pear stories: cognitive, cultural, and linguistic aspects of narrative*

production. (Ablex Pub. Corp, 1980).

70. Croft, W. Linguistic Selection: An Utterance-based Evolutionary Theory of Language Change.

Nordic Journal of Linguistic **19**, 99–139 (1996).

71. R Development Core Team. *R: A language and environment for statistical computing*. (R

Foundation for Statistical Computing, 2020).

72. Purcell, S. *et al.* PLINK: A Tool Set for Whole-Genome Association and Population-Based

Linkage Analyses. *The American Journal of Human Genetics* **81**, 559–575 (2007).

73. Jakobsson, M. & Rosenberg, N. A. CLUMPP: a cluster matching and permutation program for

dealing with label switching and multimodality in analysis of population structure.

Bioinformatics **23**, 1801–1806 (2007).

74. Rosenberg, N. A. DISTRUCT: a program for the graphical display of population structure.

Molecular Ecology Notes **4**, 137–138 (2003).

75. The International HapMap 3 Consortium. Integrating common and rare genetic variation in

diverse human populations. *Nature* **467**, 52–58 (2010).

76. Szpiech, Z. A., Blant, A. & Pemberton, T. J. GARLIC: Genomic Autozygosity Regions

Likelihood-based Inference and Classification. *Bioinformatics* **33**, 2059–2062 (2017).

77. Maples, B. K., Gravel, S., Kenny, E. E. & Bustamante, C. D. RFMix: A Discriminative

Modeling Approach for Rapid and Robust Local-Ancestry Inference. *The American Journal of*

Human Genetics **93**, 278–288 (2013).

78. Snyder, J. P. *Map projections--a working manual*. (U.S. Government Printing Office 1987).

79. Tavaré, S., Balding, D. J., Griffiths, R. C. & Donnelly, P. Inferring coalescence times from

DNA sequence data. *Genetics* **145**, 505–518 (1997).

80. Beaumont, M. A., Zhang, W. & Balding, D. J. Approximate Bayesian computation in

population genetics. *Genetics* **162**, 2025–2035 (2002).

1215

1216

 SUPPLEMENTARY FIGURES AND TABLES LEGENDS

 Figure SF1:

 Quality-Control and datasets merging procedures.

Quality controls at the genotyping call level (Stage 1) have been conducted with Illumina's GenomeStudio® software Genotyping Module. Cabo Verde original DNA samples have been collected during six separate field-trips between 2010 and 2018, genotyped in four batches using four different versions of the Illumina Omni2.5Million Beadchip genotyping array. The resulting dataset was merged with 2,504 worldwide samples from the 1000 Genomes Project Phase 3 (1000 Genomes Project Consortium, 2015); with 1,307 continental African samples from the African Genome Variation Project (EGA accession number EGAD00001000959, Gurdasani et al. 2015); and with 1,235 African samples from Patin et al. 2017 (EGA accession number EGAS00001002078). We retained only autosomal SNPs common to all datasets, and excluded one individual for each pair of individuals related at the 2^{nd} degree (at least one grand-parent in common) as inferred with KING (Manichaikul et al. 2010), following previous procedure (Verdu et al. 2017).

Figure SF2:

Multidimensional Scaling three-dimensional projection of Allele Sharing Dissimilarity pairwise dissimilarities, for all worldwide populations in our dataset.

All populations and individuals from our working-dataset are included in this analysis: see **Table ST1** for the population list used and **Figure F1** for sample locations and symbols.

Each individual is represented by a single point.

A) Axis 1 and 2; B) Axis 1 and 3. 3D animated plot is provided in *.gif* format.

We evaluate the Spearman correlation between 3D MDS projections and the original ASD pairwise matrix to evaluate the precision of the dimensionality reduction along the first three axes of the MDS. We find significant ($p<2.2 \times 10^{-16}$) Spearman ρ of 0.9796.

Figure SF3:

Multidimensional Scaling three-dimensional projection of Allele Sharing Dissimilarity pairwise dissimilarities, for African, European and other admixed populations related to the TAST.

We removed all Asian and South American samples compared to the sample set employed in **Figure SF2**, and recomputed the MDS based on this reduced sample set. See **Table ST1** for the population list used in these analyses. Sample map used in these analyses is extracted from **Figure F1** and provided in the bottom right corner. Each individual is represented by a single point.

A) Axis 1 and 2; B) Axis 2 and 3; C) Axis 1 and 3. 3D animated plot is provided in *.gif* format.

We evaluate the Spearman correlation between 3D MDS projections and the original ASD pairwise matrix to evaluate the precision of the dimensionality reduction along the first three axes of the MDS. We find significant ($p<2.2 \times 10^{-16}$) Spearman ρ of 0.9392.

Figure SF4:

Multidimensional Scaling three-dimensional projection of Allele Sharing Dissimilarity pairwise dissimilarities, for a subset of African, European and other admixed populations related to the TAST.

We removed all East African and Central African hunter-gatherer samples (Baka, Bezan, Ba.Bongo, Ba.Koya, Ba.Twa, Bi.Aka, Mbuti) compared to the sample set employed in **Figure SF3**, and recomputed the

MDS based on this reduced sample set. See **Table ST1** for the population list used in these analyses. Sample
map used in these analyses is extracted from **Figure F1** and provided in the bottom right corner. Each indi-
vidual is represented by a single point.
A) Axis 1 and 2; B) Axis 2 and 3; C) Axis 1 and 3. 3D animated plot is provided in *.gif* format.
We evaluate the Spearman correlation between 3D MDS projections and the original ASD pairwise matrix to
evaluate the precision of the dimensionality reduction along the first three axes of the MDS. We find signifi-
cant ($p < 2.2 \times 10^{-16}$) Spearman ρ of 0.9450.
1271
1272

Figure SF5:
Multidimensional Scaling three-dimensional projection of Allele Sharing Dissimilarity pairwise dis-
similarities, for a subset of African, European and and other admixed populations related to the TAST.
We removed all West and South-West Central African samples, as well as South African samples, compared
to the sample set employed in **Figure SF4**, and recomputed the MDS based on this reduced sample set. See
Table ST1 for the population list used in these analyses. Sample map used in these analyses is extracted
from **Figure F1** and provided in the bottom right corner. Each individual is represented by a single point.
A) Axis 1 and 2; B) Axis 2 and 3; C) Axis 1 and 3. 3D animated plot is provided in *.gif* format.
We evaluate the Spearman correlation between 3D MDS projections and the original ASD pairwise matrix to
evaluate the precision of the dimensionality reduction along the first three axes of the MDS. We find signifi-
cant ($p < 2.2 \times 10^{-16}$) Spearman ρ of 0.9636.
1284
1285

Figures SF6A-D:
Four separate animated Multidimensional Scaling three-dimensional projections of Allele Sharing
Dissimilarity pairwise dissimilarities, for a subset of African, European and other admixed popula-
tions related to the TAST.
Each four animated plot is provided in *.gif* format. Each individual is represented by a single point. See **Ta-**
ble ST1 for the population list used in these analyses, they correspond to each separate sample set provided
in two-dimensional plots in **Figure F2**. Spearman correlations are provided in **Figure F2** caption.
1293
1294

Figure SF7:
Population F_{st}/F_{st}^{max} values for the ASW, ACB, and each Cabo Verdean birth-island separately con-
sidering the ADMIXTURE mode result at $K=2$.
F_{st}/F_{st}^{max} values were computed using FSTruct (Morrisson et al. 2021) with 1000 bootstrap replicates per
population. All population pairwise distributions of bootstrap F_{st}/F_{st}^{max} values were significantly different
from one another after Bonferroni correction (Wilcoxon two-sided rank sum test $p < 3.96 \times 10^{-8}$), except the
following pairwise comparisons: ACB-Santiago, Brava-Maio, Fogo-Sal, and Maio-Santo Antão.
1302
1303

Figure SF8:
Alternative ADMIXTURE mode results for the individual genetic structure among Cabo Verdean,
Barbadian-ACB and African-American ASW populations related to the TAST.
ADMIXTURE (Alexander et al. 2009) analyses using resampled individual sets for the population sets origi-
nally containing more than 50 individuals (**Table ST1**). 225 unrelated Cabo Verdean-born individuals are
comprised in the analysis and grouped by island of birth (**Table ST1**). All analyses comprised 102,543 inde-
pendent autosomal SNPs, for values of K between 2 and 10. 30 independent ADMIXTURE runs were per-

formed for each value of *K*, and groups of runs (>2) providing similar results (all pairwise SSC > 99.9%)
were averaged in a single "mode" result using CLUMPP (Jakobsson and Rosenberg 2009), and plotted with
DISTRUCT (Rosenberg et al. 2004). Number of runs in the presented modes are indicated below the value
of *K*. All other mode results are presented in **Figure F3**.
1315
1316
1317   **Figure SF9:**
1318   **The distribution of (A) short, (B) medium, and (C) all ROH per individual, for each Cabo Verdean**
1319   **birth-island and other worldwide populations.**
1320
1321
1322   **Figure SF10:**
1323   **Individual total ancestry contained in (A) short, (B) medium, and (C) long ROH versus total ancestry**
1324   **not in that class ROH, for each Cabo Verdean birth-island and other worldwide populations.**
1325
1326
1327   **Figure SF11:**
1328   **The distribution of total ancestry in long ROH per individual for each Cabo Verdean birth-island.**
1329
1330
1331   **Figure SF12:**
1332   **Schematic representation of the 2-Dimensional ASD-MDS estimates of individual admixture propor-**
1333   **tions and angles, also used as summary statistics for ABC inference and implemented in *MetHis* sum-**
1334   **mary-statistics calculation tools.**
1335
1336
Figure SF13:
ABC Prior checking for *MetHis* simulations for each island separately.
10,000 simulations are conducted under each four competing scenarios of historical admixture considered in
Random-Forest ABC scenario-choice (see **Figure F6**). **A)** Goodness-of-fit histograms obtained considering
the 42 summary-statistics calculated from each simulation and 1000 permutations. Observed statistic is indi-
cated by a red vertical line**. B)** First two axes of a Principal Component Analysis performed on the 42 sum-
mary-statistics obtained for 40,000 simulations per island (10,000 simulations for each four competing-
scenario), each color corresponding to a given scenario. The red white-circled dot corresponds to the vector
of summary-statistics from the observed dataset. **C)** Axes 1 and 3 of the same PCA projection as in panels **B**).
1346
1347
1348   **Figure SF14:**
1349   **Density distributions (black line) of each 42 summary-statistics obtained from 40,000 simulations,**
1350   **10,000 under each four competing scenarios, for Santiago birth-island, compared to the observed sta-**
1351   **tistic obtained in this island (red vertical line).**
1352
1353
1354   **Figure SF15:**
1355   **Density distributions (black line) of each 42 summary-statistics obtained from 40,000 simulations,**
1356   **10,000 under each four competing scenarios, for Fogo birth-island, compared to the observed statistic**
1357   **obtained in this island (red vertical line).**

1358

1359

1360 **Figure SF16:**

1361 **Density distributions (black line) of each 42 summary-statistics obtained from 40,000 simulations,**

1362 **10,000 under each four competing scenarios, for Santo Antão birth-island, compared to the observed**

1363 **statistic obtained in this island (red vertical line).**

1364

1365

1366 **Figure SF17:**

1367 **Density distributions (black line) of each 42 summary-statistics obtained from 40,000 simulations,**

1368 **10,000 under each four competing scenarios, for São Nicolau birth-island, compared to the observed**

1369 **statistic obtained in this island (red vertical line).**

1370

1371

1372 **Figure SF18:**

1373 **Density distributions (black line) of each 42 summary-statistics obtained from 40,000 simulations,**

1374 **10,000 under each four competing scenarios, for Brava birth-island, compared to the observed statistic**

1375 **obtained in this island (red vertical line).**

1376

1377

1378 **Figure SF19:**

1379 **Density distributions (black line) of each 42 summary-statistics obtained from 40,000 simulations,**

1380 **10,000 under each four competing scenarios, for Maio birth-island, compared to the observed statistic**

1381 **obtained in this island (red vertical line).**

1382

1383

1384 **Figure SF20:**

1385 **Density distributions (black line) of each 42 summary-statistics obtained from 40,000 simulations,**

1386 **10,000 under each four competing scenarios, for Boa Vista birth-island, compared to the observed**

1387 **statistic obtained in this island (red vertical line).**

1388

1389

1390 **Figure SF21:**

1391 **Density distributions (black line) of each 42 summary-statistics obtained from 40,000 simulations,**

1392 **10,000 under each four competing scenarios, for São Vicente birth-island, compared to the observed**

1393 **statistic obtained in this island (red vertical line).**

1394

1395

1396 **Figure SF22:**

1397 **Density distributions (black line) for each 42 summary-statistics obtained from 40,000 simulations,**

1398 **10,000 under each four competing scenarios, for Sal birth-island, compared to the observed statistic**

1399 **obtained in this island (red vertical line).**

1400

1401

Figure SF23:

Island per island Random-Forest ABC cross validation results and variable importance.

We considered four competing scenarios as described in **Material and Methods** and in **Figure F6**, with
associated prior parameter distributions. Cross-validation results are obtained by conducting RF scenario-
choices using in-turn each 40,000 simulations (10,000 per competing scenario) as pseudo-observed data and
the remaining 39,999 simulations in the reference table. Note that no real data is used in this analysis. We
considered 1,000 trees in the random-forest for each analysis. Cross-validation results and associated sum-
mary-statistics' importance for the RF decisions are obtained from the same function *abcrf* in the R package
abcrf (Pudlo et al. 2016).
1411
1412
Figure SF24:
Reproductive-size posterior parameter distributions and associated priors obtained with Neural Net-
work ABC inference for each island separately.
We considered, for each island separately, 100,000 simulations computed under the winning scenario ob-
tained with RF-ABC scenario-choice procedures and provided on the left of the figure. Neural-Network tol-
erance levels and number of neurons in the hidden layer, for each birth-island separately, are chosen based
on posterior parameter cross-validation error minimization procedures conducted on 1,000 random simula-
tions used in-turn, as pseudo observed data (see **Supplementary Text**, and **Table ST4**). For each island,
posterior parameter distributions correspond to the solid lines and corresponding priors correspond to the
doted black lines. Density distributions are based on the logit transformation of parameter values (see **Sup-**
plementary Text) using an Epanechnikov kernel between the corresponding prior bounds. See **Results** and
Supplementary Text for description and discussion of the results. Results are synthesized in **Figure F7**.
1425
1426
Figure SF25:
African admixture histories posterior parameter distributions and associated priors obtained with
Neural Network ABC inference for each island separately.
We considered, for each island separately, 100,000 simulations computed under the winning scenario ob-
tained with RF-ABC scenario-choice procedures and provided on the left of the figure. Neural Network tol-
erance levels and number of neurons in the hidden layer, for each island separately, are chosen based on pos-
terior parameter cross-validation error minimization procedures conducted on 1,000 random simulations
used in-turn, as pseudo observed data (see **Supplementary Text**, and **Table ST4**). For each island, posterior
parameter distributions correspond to the solid lines and corresponding priors correspond to the doted black
lines. Density distributions are based on the logit transformation of parameter values (see **Supplementary**
Text) using an Epanechnikov kernel between the corresponding prior bounds. See **Results** and **Supplemen-**
tary Text for description and discussion of the results. Results are synthesized in **Figure F7**.
1439
1440
Figure SF26:
European admixture histories posterior parameter distributions and associated priors obtained with
Neural Network ABC inference for each island separately.
We considered, for each island separately, 100,000 simulations computed under the winning scenario ob-
tained with RF-ABC scenario-choice procedures and provided on the left of the figure. Neural Network tol-
erance levels and number of neurons in the hidden layer, for each island separately, are chosen based on pos-
terior parameter cross-validation error minimization procedures conducted on 1,000 random simulations
used in-turn, as pseudo observed data (see **Supplementary Text**, and **Table ST4**). For each island, posterior
parameter distributions correspond to the solid lines and corresponding priors correspond to the doted black
lines. Density distributions are based on the logit transformation of parameter values (see **Supplementary**

Text) using an Epanechnikov kernel between the corresponding prior bounds. See **Results** and **Supplemen-**
tary Text for description and discussion of the results. Results are synthesized in **Figure F7**.

Table ST1:
Population table corresponding to Figure F1 map and sample inclusion in all analyses, respectively.

Table ST2:
Historical landmark chronology for the peopling history of Cabo Verde as provided by previous his-
torical work, respectively for each island.
See detailed historical references in the table.

Table ST3:
Summary-statistics used for *MetHis*-machine-learning ABC inferences.

Table ST4:
Average parameter posterior random cross-validation error across all model parameters as a function
of the number of neurons in the hidden layer and the rejection tolerance level (number of simulations
retained for training the Neural-Network) under the winning scenario for each island respectively.
For each island separately, we considered, 1,000 random simulations in-turn as pseudo-observed data to es-
timate posterior parameter distributions and 100,000 total simulations in the reference table. For each cross-
validation procedure, we considered between 5 and 11 neurons in the hidden layer ("NN-HL") for the win-
ning scenarios with 12 original parameters, and between 5 and 12 NN-HL for the winning scenarios with 13
original parameters. We considered three different tolerance levels of 0.01, 0.03, and 0.1 ("Tol.") corre-
sponding, respectively, to 1,000, 3,000, and 10,000 simulations, in turn closest to each one of the 1,000
cross-validation pseudo-observed simulation retained for training the NN. The median values of posterior
parameter distributions were used as point estimates for the calculation of the error of each parameter
(Csilléry et al. 2012).
For each birth-island and corresponding winning scenario separately, we considered, for further posterior
parameter estimation using the observed "real" data, only the pair of tolerance level and number of hidden
neurons that minimized the average error on posterior parameter estimations across all parameters (indicated
in bold in the table).

Table ST5-Santiago:
NN-ABC posterior parameter estimations for Santiago island, cross-validation posterior parameter
errors, and cross-validation 95% Credibility Interval accuracies.
Cross-validation errors and 95% CI accuracies are based on 1,000 NN-ABC posterior parameter inferences
using, in turn, each one of the 1,000 simulations closest to the observed data used as pseudo-observed data,
and the remaining 99,999 simulations as the reference table.
Error calculations are based on the mode and, separately, the median point estimate for each 1,000 pseudo-
observed simulations posterior parameter estimation, compared to the known parameter used for simulation.
We considered 6 neurons in the hidden layer and a tolerance level of 0.01 (1000 simulations) for all posterior

parameter estimation in this analysis as identified in **Table ST4**. Plotted distributions for posterior parameter
estimations can be found in **Figure SF24-26**.
1499
1500
Table ST6-Fogo:
NN-ABC posterior parameter estimations for Fogo island, cross-validation posterior parameter errors,
and cross-validation 95% Credibility Interval accuracies.
Cross-validation errors and 95% CI accuracies are based on 1,000 NN-ABC posterior parameter inferences
using, in turn, each one of the 1,000 simulations closest to the observed data used as pseudo-observed data,
and the remaining 99,999 simulations as the reference table.
Error calculations are based on the mode and, separately, the median point estimate for each 1,000 pseudo-
observed simulations posterior parameter estimation, compared to the known parameter used for simulation.
We considered 10 neurons in the hidden layer and a tolerance level of 0.01 (1000 simulations) for all poste-
rior parameter estimation in this analysis as identified in **Table ST4**. Plotted distributions for posterior pa-
rameter estimations can be found in **Figure SF24-26**.
1512
1513
Table ST7-Santo Antão:
NN-ABC posterior parameter estimations for Santo Antão island, cross-validation posterior parame-
ter errors, and cross-validation 95% Credibility Interval accuracies.
Cross-validation errors and 95% CI accuracies are based on 1,000 NN-ABC posterior parameter inferences
using, in turn, each one of the 1,000 simulations closest to the observed data used as pseudo-observed data,
and the remaining 99,999 simulations as the reference table.
Error calculations are based on the mode and, separately, the median point estimate for each 1,000 pseudo-
observed simulations posterior parameter estimation, compared to the known parameter used for simulation.
We considered 11 neurons in the hidden layer and a tolerance level of 0.01 (1000 simulations) for all poste-
rior parameter estimation in this analysis as identified in **Table ST4**. Plotted distributions for posterior pa-
rameter estimations can be found in **Figure SF24-26**.
1525
1526
Table ST8-São Nicolau:
NN-ABC posterior parameter estimations for São Nicolau island, cross-validation posterior parameter
errors, and cross-validation 95% Credibility Interval accuracies.
Cross-validation errors and 95% CI accuracies are based on 1,000 NN-ABC posterior parameter inferences
using, in turn, each one of the 1,000 simulations closest to the observed data used as pseudo-observed data,
and the remaining 99,999 simulations as the reference table.
Error calculations are based on the mode and, separately, the median point estimate for each 1,000 pseudo-
observed simulations posterior parameter estimation, compared to the known parameter used for simulation.
We considered 11 neurons in the hidden layer and a tolerance level of 0.01 (1000 simulations) for all poste-
rior parameter estimation in this analysis as identified in **Table ST4**. Plotted distributions for posterior pa-
rameter estimations can be found in **Figure SF24-26**.
1538
1539
Table ST9-Brava:
NN-ABC posterior parameter estimations for Brava island, cross-validation posterior parameter er-
rors, and cross-validation 95% Credibility Interval accuracies.

Cross-validation errors and 95% CI accuracies are based on 1,000 NN-ABC posterior parameter inferences
using, in turn, each one of the 1,000 simulations closest to the observed data used as pseudo-observed data,
and the remaining 99,999 simulations as the reference table.
Error calculations are based on the mode and, separately, the median point estimate for each 1,000 pseudo-
observed simulations posterior parameter estimation, compared to the known parameter used for simulation.
We considered 10 neurons in the hidden layer and a tolerance level of 0.01 (1000 simulations) for all poste-
rior parameter estimation in this analysis as identified in **Table ST4**. Plotted distributions for posterior pa-
rameter estimations can be found in **Figure SF24-26**.
1551
1552

Table ST10-Maio:
NN-ABC posterior parameter estimations for Maio island, cross-validation posterior parameter errors,
and cross-validation 95% Credibility Interval accuracies.
Cross-validation errors and 95% CI accuracies are based on 1,000 NN-ABC posterior parameter inferences
using, in turn, each one of the 1,000 simulations closest to the observed data used as pseudo-observed data,
and the remaining 99,999 simulations as the reference table.
Error calculations are based on the mode and, separately, the median point estimate for each 1,000 pseudo-
observed simulations posterior parameter estimation, compared to the known parameter used for simulation.
We considered 10 neurons in the hidden layer and a tolerance level of 0.01 (1000 simulations) for all poste-
rior parameter estimation in this analysis as identified in **Table ST4**. Plotted distributions for posterior pa-
rameter estimations can be found in **Figure SF24-26**.
1564
1565

Table ST11-Boa Vista:
NN-ABC posterior parameter estimations for Boa Vista island, cross-validation posterior parameter
errors, and cross-validation 95% Credibility Interval accuracies.
Cross-validation errors and 95% CI accuracies are based on 1,000 NN-ABC posterior parameter inferences
using, in turn, each one of the 1,000 simulations closest to the observed data used as pseudo-observed data,
and the remaining 99,999 simulations as the reference table.
Error calculations are based on the mode and, separately, the median point estimate for each 1,000 pseudo-
observed simulations posterior parameter estimation, compared to the known parameter used for simulation.
We considered 8 neurons in the hidden layer and a tolerance level of 0.01 (1000 simulations) for all posterior
parameter estimation in this analysis as identified in **Table ST4**. Plotted distributions for posterior parameter
estimations can be found in **Figure SF24-26**.
1577
1578

Table ST12-São Vicente:
NN-ABC posterior parameter estimations for São Vicente island, cross-validation posterior parameter
errors, and cross-validation 95% Credibility Interval accuracies.
Cross-validation errors and 95% CI accuracies are based on 1,000 NN-ABC posterior parameter inferences
using, in turn, each one of the 1,000 simulations closest to the observed data used as pseudo-observed data,
and the remaining 99,999 simulations as the reference table.
Error calculations are based on the mode and, separately, the median point estimate for each 1,000 pseudo-
observed simulations posterior parameter estimation, compared to the known parameter used for simulation.
We considered 11 neurons in the hidden layer and a tolerance level of 0.01 (1000 simulations) for all poste-
rior parameter estimation in this analysis as identified in **Table ST4**. Plotted distributions for posterior pa-
rameter estimations can be found in **Figure SF24-26**.

1590

1591

Table ST13-Sal:

NN-ABC posterior parameter estimations for Sal island, cross-validation posterior parameter errors,
and cross-validation 95% Credibility Interval accuracies.

Cross-validation errors and 95% CI accuracies are based on 1,000 NN-ABC posterior parameter inferences
using, in turn, each one of the 1,000 simulations closest to the observed data used as pseudo-observed data,
and the remaining 99,999 simulations as the reference table.

Error calculations are based on the mode and, separately, the median point estimate for each 1,000 pseudo-
observed simulations posterior parameter estimation, compared to the known parameter used for simulation.

We considered 11 neurons in the hidden layer and a tolerance level of 0.01 (1000 simulations) for all poste-
rior parameter estimation in this analysis as identified in **Table ST4**. Plotted distributions for posterior pa-
rameter estimations can be found in **Figure SF24-26**.

1603

1604

1605 **Table ST14:**

1606 **Mantel correlations among individual birth-places, residence-places, maternal and paternal birth**
1607 **places, age, and academic education duration.**

1608 We considered only the 225 genetically unrelated Cabo Verde-born individuals in these analyses.