Coarsening dynamics can explain meiotic crossover patterning in both the presence and absence of the synaptonemal complex

John A Fozard[1], Chris Morgan[2,*], Martin Howard[1,*]

[1] Computational and Systems Biology and [2] Cell and Developmental Biology, John Innes Centre, Norwich Research Park, Norwich NR4 7UH, UK

* Co-corresponding authors: chris.morgan@jic.ac.uk, martin.howard@jic.ac.uk

Abstract

The shuffling of genetic material facilitated by meiotic crossovers is a critical driver of genetic variation. Therefore, the number and positions of crossover events must be carefully controlled. In *Arabidopsis,* an obligate crossover and repression of nearby crossovers on each chromosome pair are abolished in mutants that lack the synaptonemal complex (SC), a conserved protein scaffold. We use mathematical modelling and quantitative super-resolution microscopy to explore and mechanistically explain meiotic crossover pattering in *Arabidopsis* lines with full, incomplete or abolished synapsis. For *zyp1* mutants, which lack an SC, we develop a coarsening model in which crossover precursors globally compete for a limited pool of the pro-crossover factor HEI10, with dynamic HEI10 exchange mediated through the nucleoplasm. We demonstrate that this model is capable of quantitatively reproducing and predicting *zyp1* experimental crossover patterning and HEI10 foci intensity data. Additionally, we find that a model combining both SC- and nucleoplasm-mediated coarsening can explain crossover patterning in wild-type *Arabidopsis* and in *pch2* mutants, which display partial synapsis. Together, our results reveal that regulation of crossover patterning in wild-type *Arabidopsis* and SC defective mutants likely act through the same underlying coarsening mechanism, differing only in the spatial compartments through which the pro-crossover factor diffuses.

Introduction

During meiotic prophase I, pairs of threadlike homologous chromosomes are tethered together through the formation of a protein scaffold called the synaptonemal complex (SC) (Page and Hawley, 2004). Prior to SC formation, programmed double strand breaks (DSBs) form along the length of the chromosomal DNA (Gray and Cohen, 2016; Hunter, 2015). DSBs are then repaired via the formation of recombination-intermediate (RI) DNA joint-molecules, as either non-crossovers (NCOs) or crossovers (COs) (Gray and Cohen, 2016; Hunter, 2015). In most species, the number of RIs repaired as NCOs vastly outnumbers those repaired as COs (Gray and Cohen, 2016; Hunter, 2015). Importantly, nearby pairs of COs on the same chromosome are less likely to occur than more distant pairs, but each pair of chromosomes always receives at least one CO. These phenomena are known as 'crossover interference' and 'crossover assurance', respectively (Jones and Franklin, 2006; Otto and Payseur, 2019; Sturtevant, 1915; von Diezmann and Rog, 2021; Zickler and Kleckner, 2016).

Recently, we introduced a mechanistic coarsening model that quantitatively explains CO interference, CO assurance and other CO patterning features in wild-type *Arabidopsis thaliana* (Morgan et al., 2021). In our SC-mediated coarsening model, the pro-CO protein HEI10 (Chelysheva et al., 2012; Ziolkowski et al., 2017) can diffuse along synapsed homolog bivalents and can reversibly bind into immobile clusters at RI sites. Conservation of the total amount of HEI10 on individual bivalents, along with an unbinding escape rate from RIs that decreases as more HEI10 locally accumulates, causes the system to coarsen. By the end of the pachytene substage of prophase I, significant amounts of HEI10 are retained at only a small number of predominantly distantly-spaced RIs, each with high levels of HEI10, which are then designated to become COs (Morgan et al., 2021). A similar model was subsequently proposed to underlie CO patterning in *C. elegans* (Zhang et al., 2021), suggesting that this molecular coarsening paradigm represents a conserved feature of meiotic CO control.

The coarsening model fundamentally relies on the retention of HEI10 molecules on synapsed bivalent chromosomes, and redistribution of HEI10 molecules by diffusion along the bivalents. Work in *C. elegans* points to the SC acting as a conduit, or channel, through which HEI10 molecules can diffuse, promoting coarsening along individual bivalents. Consistently, in *C. elegans*, the SC is known to have liquid-crystal like properties that can spatially compartmentalise pro-CO proteins (Rog et al., 2017; Zhang et al., 2018). Furthermore, innovative live-cell imaging experiments have recently directly demonstrated that ZHP-3, a HEI10 ortholog, can diffuse along the SC (Stauffer et al., 2019; Zhang et al., 2021). In *Arabidopsis*, it was also recently shown that CO interference and assurance are abolished in plants lacking the SC transverse filament protein, ZYP1 (Capilla-Pérez et al., 2021; France et al., 2021). These observations suggest that the SC plays a direct role in mediating crossover patterning dynamics.

In this work, we therefore develop a new coarsening model for crossover interference in the absence of an SC, and use it to explore the regulation of crossover number and position in *Arabidopsis zyp1* mutants with abolished synapsis. In this new model, due to the lack of an SC, HEI10 molecules are no longer restricted to diffuse along individual bivalents. Instead, HEI10 is assumed to diffuse through a communal nucleoplasm, generating competition for HEI10 between RIs located on different bivalents. Similarly, in *C. elegans*, it has previously been suggested that the SC functions to prevent exchange of recombination proteins between RIs via the nucleoplasm (Rog et al., 2017; Zhang et al., 2021, 2018). Other studies have also recently proposed that nucleoplasm-mediated coarsening can explain CO patterning in *Arabidopsis* mutants with abolished synapsis, although the dynamics of this process have not been explicitly modelled (Durand et al., 2022; Lloyd, 2022). Using

our model, we provide a mechanistic explanation for why *zyp1* mutants without synapsis lose both
CO interference and assurance. Intriguingly, we also reveal and explain an additional layer of
crossover control, where *zyp1* mutants lose positional effects that are determined by CO
interference along individual chromosome pairs, but still exhibit constraints on total crossover
number per-cell. Importantly, we also find that a model combining nucleoplasm-mediated and SC-
mediated coarsening is still fully capable of explaining CO patterning in wild-type *Arabidopsis,* as well
as in *pch2* mutants, which display incomplete synapsis (Lambing et al., 2015). Overall, our results
demonstrate a critical role for the SC in controlling and constraining the dynamic coarsening of
HEI10 molecules.

Results

A nucleoplasm-mediated coarsening model for crossover patterning without the SC

The SC likely acts as a conduit for HEI10 diffusion along synapsed pachytene bivalents (Morgan et al.,
2021; Rog et al., 2017; Zhang et al., 2021, 2018). Without an SC, we postulated that HEI10 diffusion
could instead occur through the nucleoplasm, with little or no movement along chromosomes, as
has also previously been suggested from work in *C. elegans* (Zhang et al., 2021). We therefore
investigated this hypothesis by developing a new coarsening model for crossover interference
without an SC. This 'nucleoplasmic coarsening' model incorporates the processes shown in Figure
1A,B on the 5 pairs of homologous chromosomes in each cell, whose lengths are drawn from the
experimentally measured length distributions, and with randomly positioned recombination
intermediates (RIs) (Methods).

In this model, HEI10 is able to move from the nucleoplasmic pool to the HEI10 clusters at each of the
RIs. It can also escape from the RI clusters back into the nucleoplasmic pool, at a rate which depends
on the amount of HEI10 within that RI cluster. Similar to previous work (Morgan et al., 2021), this
rate is chosen to have the form shown in Figure 1C,D, decreasing as the amount of HEI10 within the
RI compartment increases. Again, as in our earlier model (Morgan et al., 2021), if the total amount of
HEI10 is sufficiently high then a uniform steady state becomes unstable and the system progressively
coarsens, with RIs with more HEI10 growing at the expense of those with less. Eventually, the
majority of the HEI10 will accumulate into a single focus per cell, but the limited duration of
pachytene means that this process does not complete, and instead there are a number of RIs with
significant levels of HEI10 at the end of the simulation. Full details of the simulated model can be
found in Figure 1A-G and Methods, including minor differences in how the model was simulated (in
the initial distribution of HEI10, dynamics at small HEI10 foci and criteria for HEI10 focus/CO calling)
as compared to our earlier work (Morgan et al., 2021).

Nucleoplasmic coarsening model explains CO number distribution in *zyp1* mutants

To determine whether the nucleoplasmic coarsening model was capable of explaining CO patterning
in *Arabidopsis* mutants without synapsis, we first manually adjusted model parameter values to fit
simulation outputs to existing experimental data on CO frequency from *Arabidopsis zyp1* null
mutants, which lack an SC. These data include total CO number and number of homologs without a
CO (Figure 2A,B).

By counting the number of MLH1 foci in late-prophase I cells in 7 different Col-0 *zyp1* null mutant
lines, it was previously found that there was a ~50% increase in the predominant class I COs in these
mutants compared with wild-type (data from (Capilla-Pérez et al., 2021), shown in Figure 2A). Also,
by analysing DAPI stained metaphase I cells, it was found that ~11% of metaphase I cells contained a

pair of univalent chromosomes, which form when homologs fail to form a single CO, indicating an
absence of CO assurance (data from (Capilla-Pérez et al., 2021), shown in Figure 2B). Importantly, we
found that the nucleoplasmic coarsening model was capable of recapitulating the increase in CO
number and the increased univalent frequency observed in *zyp1* mutants (Figure 2A, B). The number
of COs per cell was slightly less dispersed, with a lower sample variance, within our model than in
the experimental data. This reduction is likely because the experimental data is pooled from multiple
different mutant lines, each of which has a slightly different mean number of COs per cell, thereby
generating a broader distribution (Capilla-Pérez et al., 2021). The expected frequency of univalents is
also slightly higher in our simulation outputs than in the experimental data, but this is again
expected as chiasma number in metaphase I cells (that were used for the experimental analysis) are
influenced by both class I and class II CO numbers, whereas the nucleoplasmic coarsening model
only simulates CO patterning via the (dominant) class I pathway. Assuming ~2 additional class II COs
per cell (Mercier et al., 2005) are distributed randomly among the five chromosome pairs, in a
simple estimation we would expect the univalent frequency in our model output to be reduced by a
factor of $(0.8)^2=0.64$ (from 19% of cells to 12% of cells containing one or more univalent). However,
univalents are more likely to be associated with the shorter chromosomes, as they have fewer RIs.
This was recently experimentally demonstrated, with chromosome 4 (the shortest *Arabidopsis*
chromosome) exhibiting the highest frequency of aneuploidy in *zyp1* mutants (Durand et al., 2022).
This effect slightly reduces how much the addition of class II crossovers decreases the frequency of
cells with one or more univalent. In our simulations we thus find a value of ~13%, similar to but
slightly higher than previous observations (~11%) (Capilla-Pérez et al., 2021).

Using the nucleoplasmic coarsening model, we were also able to simulate the distribution of
crossovers on each of the five *Arabidopsis* chromosomes (Figure 2C). Consistent with previous
genetic analyses, we found that the average number of crossovers per chromosome pair is positively
correlated with their physical size (Durand et al., 2022). We also found that the distribution of the
number of crossovers on each chromosome is close to a Poisson distribution, with nearly equal
mean and variance. Processes that occur at a constant probability per unit space or time, when
integrated over a fixed interval, have a Poisson distribution (Haldane, 1931). Hence, as expected, CO-
interference along each chromosome is not present in this model. These findings are consistent with
a previous genetic analysis, which found that the distribution of CO numbers on individual
chromosomes did not deviate from a Poisson distribution (Capilla-Pérez et al., 2021).

Paradoxically, while the distribution of CO numbers on individual chromosomes in our simulations
closely followed a Poisson distribution, the total number of COs per cell did not. Applying a test for
underdispersion (Cameron and Trivedi, 1990), the distribution of simulated CO numbers was found
to be significantly under-dispersed relative to a Poisson distribution (z=-276.85, p<0.001, see also
Figure 2A). Compellingly, we found that a sub-Poissonian distribution of total CO numbers per cell
was also found in the experimental data from (Capilla-Pérez et al., 2021) (Figure 2A; test for
underdispersion z=-6.5757, p<0.001), although this experimental under-dispersion was not
previously highlighted. In the model, the total number of crossovers in each cell is regulated through
global competition for a limited pool of HEI10. This, combined with regulation of cell-to-cell total
HEI10 amounts (Methods) leads to under-dispersion, despite the fact that there is little or no control
on the number of crossovers on each chromosome pair. This under-dispersion persists in the model
even with altered initial conditions (eliminating non-uniform initial HEI10 loading, Methods, Figure 2
– figure supplement 1).

We also found that the coarsening dynamics within the nucleoplasmic coarsening model were
capable of producing the experimentally observed final CO numbers within the limited ~10h

duration of the pachytene substage of prophase I in *Arabidopsis* (Prusicki et al., 2019) (Figure 2D).
Overall, we find that the nucleoplasmic coarsening model fits well with prior data on CO numbers in
SC mutants, giving strong support to the hypothesis of nucleoplasmic HEI10 exchange.

Nucleoplasmic coarsening model explains CO number distribution in *zyp1* HEI10 overexpressing mutants

To further test the ability of the nucleoplasmic coarsening model to explain CO patterning in
mutants without synapsis, we also sought to fit the model to recent experimental data examining
the combined effect of *zyp1* loss of function and HEI10 overexpression (Durand et al., 2022). Again,
by counting the number of MLH1 foci in late-prophase cells it was found that combining the *zyp1*
mutation with HEI10 overexpression leads to an average of 45.0 (σ^2=64) class I COs per cell in male
Col-0 meiocytes (Durand et al., 2022). This represents an increase in class I COs to approximately 3.5
times the number in wild-type. Unlike in the sole *zyp1* mutant, there was no evidence of a sub-
Poissonian distribution of COs per cell within the *zyp1* + HEI10 overexpression lines (test for
underdispersion; z=1.0843, p=0.86).

Increasing the total HEI10 within our nucleoplasmic coarsening model to 3.5 times its original
amount (applied to both the mean and standard-deviation of the distribution of total cellular HEI10)
gave approximately the same increase in CO number (to mean CO number 47.3), but did not
generate the observed increase in variance ($\sigma^2 = 14$), which was significantly smaller than the
experimental observations (Levene's test for equality of variances F(1,10034)=68.602, p<0.001). To
obtain increased variance in CO number, we increased the standard-deviation in the total cellular
HEI10 levels, increasing by approximately an order of magnitude (Methods). With this additional
noise, which we attribute to presence of the HEI10 transgene, we were able to recapitulate this
experimentally observed increase in CO number and variance, with simulation outputs giving an
average of 46.9 COs per cell ($\sigma^2 = 63$), this variance not being significantly different
(F(1,10034)=0.0982, p=0.754).

Super-resolution imaging of HEI10 in SC-defective *zyp1* mutant cells

As described above, previous studies have used HEI10 and MLH1 focus counts per cell, alongside
genetic analyses, to examine the effects of SC absence on CO frequency in *Arabidopsis.* However, the
material positioning of recombination sites (in units of microns) along prophase I chromosomes has
not yet been cytologically investigated in *zyp1* mutants. This is particularly important because, as
argued in (Zickler and Kleckner, 2016), the optimal metric for measuring CO interference is microns
of axis length. Additionally, as shown previously, correlations between HEI10 focal intensity and foci
number or position can be detected that provide strong experimental support for the coarsening
model (Morgan et al., 2021).

We therefore used super-resolution microscopy and a bespoke image-analysis pipeline to quantify
the position and intensity of late-HEI10 foci (that are known to mark the positions of class I COs
(Chelysheva et al., 2012)), along late prophase I chromosomes in wild-type and *zyp1a-2/zyp1b-1* SC
null mutant lines (previously characterised in (France et al., 2021)) (Figure 3A,B). Late-prophase I
cells from wild-type and *zyp1a-2/zyp1b-1* plants were stained for SMC3 (Lam et al., 2005) and HEI10
(Chelysheva et al., 2012), which label the meiotic axis and putative CO sites, respectively, and
imaged using 3D-SIM microscopy. In total, we imaged 10 cells from 4 wild-type plants and 14 cells
from 4 *zyp1a-2/zyp1b-1* plants, which provided sufficient data to quantitively compare with model
simulations. By tracing along the linear SMC3 signals, using the SNT plugin to FIJI (Arshadi et al.,
2021), we were able to segment the 10 axial elements in each cell (Figure 3A,B, top right panels). In

accordance with previous studies, we found that axial elements of homologous chromosomes were
tightly juxtaposed along their entire lengths in wild-type cells, whilst in the *zyp1a-2/zyp1b-1* line the
axes were only loosely paired along their lengths (Figure 3A,B, bottom rows), with the majority of
HEI10 foci being clearly located between the two homologous axes at an equivalent position along
their length (Figure 3A,B, right hand boxes) (Capilla-Pérez et al., 2021; France et al., 2021). Once the
paths of segmented axes had been extracted from the images, we were then able to use an
automated image analysis pipeline to assign HEI10 foci to specific positions along individual axes
based on their local proximity to those regions (Materials and Methods). Additionally, 3D-SIM
analysis of early pachytene cells in both the wild-type and *zyp1a-2/zyp1b-1* lines revealed a greater
number of less bright HEI10 foci than were observed in late-prophase I cells, which was consistent
with expectations (Figure 3 – figure supplement 1).

Nucleoplasmic coarsening model successfully predicts HEI10 foci patterning and intensities in *zyp1* mutants

Using the experimental cytology and image analysis described above, we were able to construct
cytological late-HEI10 foci maps from both wild-type and *zyp1a-2/zyp1b-1* SC null mutant plants
(Figure 4A). In the wild-type we found an average of 9.0 late-HEI10 foci per cell, while in the *zyp1a-*
2/zyp1b-1 line we found an average of 18.7 late-HEI10 foci per cell, which was broadly consistent
with previous cytological analyses of CO numbers in similar lines (Capilla-Pérez et al., 2021; France et
al., 2021).

When analysing the relative distribution of all late-HEI10 foci along chromosome pairs, we found
that in the *zyp1a-2/zyp1b-1* line the distribution of late-HEI10 foci was shifted towards more distal
positions, nearer the chromosome ends, when compared with the wild-type, which was also
consistent with previous genetic analysis of *zyp1* null mutants (Capilla-Pérez et al., 2021) (Figure 4A).
Importantly, nucleoplasmic coarsening model fits were capable of recapitulating this distribution,
with a preference for forming COs in more distal regions (Figure 4A). Such increases are a direct
consequence of the additional initial HEI10 loading applied to RIs near chromosome ends (Figure 1F),
which was a parameter that was included to improve the fit in the original coarsening model
(Morgan et al., 2021). Without the inclusion of this parameter, we would expect a flat distribution
with no preference for forming COs in more distal regions. Note that the extra loading here (up to
25% on the most distal 60% of each chromosome pair) is lower and more spread than that used in
the original model (up to 100% on the most distal 20%) (Figure 1 – figure supplement 1). The
increased end-loading of HEI10 was included in the original model because the dynamics of SC-
mediated coarsening naturally decrease the likelihood of RIs close to chromosome ends forming
COs. As there is an absence of SC-mediated-coarsening in the nucleoplasmic coarsening model, each
RI is equally capable of forming a CO regardless of its position. Thus, even though there is less
pronounced HEI10 end-loading in our nucleoplasmic coarsening model compared with our original
model (25% vs. 100%), this lower end-bias still causes a relative increase in distally positioned COs
under nucleoplasmic coarsening conditions compared with SC-mediated coarsening conditions. In
the original coarsening model, we hypothesised this end bias results from preferential synapsis near
the chromosome ends. In the absence of an SC there is obviously no synapsis, which explains why
the amount of end loading required within our nucleoplasmic-coarsening simulations is lower.
However, the preferential pairing of telomeres (to form the meiotic bouquet) (Varas et al., 2015),
which presumably still happens in the absence of an SC, may explain why a smaller end-bias for
HEI10 loading persists.

We also examined the spacing between adjacent late-HEI10 foci on the same chromosome pair. In
the wild-type, we found a scarcity of closely spaced crossovers, with a greater frequency of distantly
spaced COs, reflecting the action of CO interference (Figure 4B). This was consistent with our
previous analysis of wild-type CO patterning (albeit noisier due to the smaller sample size in this
study) (Morgan et al., 2021). In the *zyp1a-2/zyp1b-1* line, however, we found a very high frequency
of closely spaced crossovers, with a diminishing number of distantly spaced COs (Figure 4B). This
behaviour reflects an absence of CO interference, and provides strong cytological support for
previous conclusions drawn from genetic experiments in *zyp1* null mutants (Capilla-Pérez et al.,
2021). In this instance, the nucleoplasmic coarsening model was fully capable of predicting this
pattern of closely spaced COs, without additional parameterisation, validating the model and
reflecting an absence of CO interference (Figure 4B). Note, we use the term 'predicting' here and
below to define instances where the model simulations were capable of explaining data that was not
used to explicitly fit the model.

In addition to the late-HEI10 focus positioning data, we were also able to extract fluorescence
intensity measurements from each HEI10 focus in our imaging data. In wild-type *Arabidopsis,* we
previously showed that the normalized intensity of individual late HEI10 foci is negatively correlated
with the number of foci per bivalent, which is consistent with an SC-mediated-coarsening process
(Morgan et al., 2021). Here, we again identified such a correlation in wild-type plants (Figure 4C).
However, we did not find such a negative correlation in *zyp1a-2/zyp1b-1* mutants (Figure 4C). This
would be expected from a nucleoplasmic coarsening model as, in this scenario, HEI10 is equally
capable of diffusing through the nucleoplasm from any one RI to any other RI, regardless of whether
the other RI is located on the same chromosome pair or a different chromosome pair (see Figure
1A,B). Indeed, simulations of the nucleoplasmic coarsening model again predicted these findings
without the need for additional parameterisation and produced comparable results to the *zyp1a-*
2/zyp1b-1 line (Figure 4C).

Combining synaptonemal complex- and nucleoplasm-mediated coarsening models

Although it appears likely that the SC acts as a conduit to promote HEI10 diffusion (Morgan et al.,
2021; Rog et al., 2017; Zhang et al., 2021, 2018), it remains unclear whether, in wild-type cells, the
SC is capable of irreversibly compartmentalising the diffusion of HEI10, such that there is little or no
recycling of SC-bound HEI10 back into the nucleoplasm. To address this question, we sought to
combine our original SC-mediated coarsening model, which assumes there is no recycling of SC-
bound HEI10 molecules back into the nucleoplasm (Morgan et al., 2021), with the nucleoplasmic
coarsening model (Methods, Figure 4 – figure supplement 1). We then tested whether this combined
model was still capable of robustly reproducing the experimental CO patterning results we
previously obtained from wild-type and HEI10 overexpressing *Arabidopsis* (Morgan et al., 2021)
(Figure 4 – figure supplement 2).

We used a combined version of the coarsening model, where the rates of absorption and escape
between the RI and the SC were set to be 90% of those in the original SC-mediated coarsening
model (Morgan et al., 2021), and the exchange rates between the RI and nucleoplasm were set to be
10% of those in the above nucleoplasmic coarsening model. We found that this combined model
was still fully capable of explaining the CO number and CO spacing distributions previously observed
in wild-type *Arabidopsis* (Figure 4 – figure supplement 2). Importantly, with this combined
coarsening ratio, we still observed only a very small proportion (<0.2%) of bivalents with zero COs
forming in our simulations and, hence, CO assurance is retained. This analysis demonstrates that a
scenario where the SC preferentially promotes retention and diffusion of HEI10 along the SC, but

without completely blocking HEI10 exchange between the SC and nucleoplasm, is still fully
compatible with the coarsening model for CO patterning. Additionally, by increasing total HEI10
concentration (both along the SC and at RIs) within our combined coarsening model simulations by
4.5-fold, we were able the reproduce CO number and CO spacing results previously obtained from
HEI10 over-expressor lines (Figure 4 – figure supplement 2).

The combined coarsening model explains CO patterning in mutants with partial synapsis

As well as investigating coarsening in wild-type plants, HEI10 over-expressors and in mutants that
completely lack an SC, we also sought to determine if the combined coarsening model could explain
CO patterning in mutants that exhibit partial, but incomplete, synapsis. In *Arabidopsis*, closely-
spaced crossovers have been observed in a number of mutants that exhibit partial synapsis, such as
axr1, *pss1* and *asy1* (Duroc et al., 2014; Jahns et al., 2014; Lambing et al., 2020; Pochon et al., 2022).
Intuitively, this common observation could be explained by the coarsening model. In the presence of
partial synapsis, the HEI10 protein would still preferentially load and diffuse along the SC, similar to
the wild-type situation. However, as the total length of SC available for HEI10 loading would be less
than wild-type, but the total nuclear concentration of HEI10 protein would presumably be the same
as wild-type, we anticipate that a higher concentration of HEI10 would be loaded per micron of SC
length. This situation is then similar to that found in a HEI10 over-expressor line, which has a greater
frequency of closely-spaced COs than wild-type, as previously explained by the coarsening model
(Morgan et al., 2021).

To test this theory, we investigated the patterning of late-HEI10 foci in an *Arabidopsis pch2* mutant.
PCH2 is a conserved AAA+ ATPase, with many known meiotic functions (Bhalla, 2022; Lambing et al.,
2015; Yang et al., 2022, 2020). In *Arabidopsis,* loss of PCH2 has been shown to compromise SC
polymerisation, generating a greater number of closely spaced COs, with no known effect on meiotic
DSB number (Lambing et al., 2015). Once again, we used 3D-SIM microscopy to assess and
quantitively evaluate the positions of HEI10 foci in late-pachytene *pch2-1* mutants (Figure 5A). In
total, we imaged 39 cells from 3 *pch2-1* plants. We found that late-HEI10 foci in *pch2-1* mutants
were exclusively associated with the short SC segments that form. On average, *pch2-1* cells
contained 14 (std. dev. ± 3) SC segments with an average length of 5.1 (std. dev. ± 4.4) μm. We
therefore mapped the position of HEI10 foci along these short SC segments using our bespoke image
analysis pipeline and quantified the number and position of late-HEI10 per SC segment within each
nucleus (Figure 5B-F, left hand plots). We detected 11.1 (σ^2 = 3.2) late-HEI10 foci per cell (Figure 5B),
with an average of 0.79 (σ^2 = 0.48) late-HEI10 foci per SC segment (Figure 5C). We also found that all
late-HEI10 foci were reasonably evenly distributed along SC segments (Figure 5D), and closely
spaced late-HEI10 foci were frequently detected (Figure 5E). Also, the longer SC segments tended to
have more late-HEI10 foci (Figure 5F), and the relative intensity of single late-HEI10 foci (i.e., foci on
SC segments that possess only a single focus) was positively correlated with SC segment length
(Figure 5G).

To determine if the combined coarsening model was capable of recapitulating the patterning effects
observed in *pch2-1* mutants, we ran simulations using the same number and length distribution of
SC segments per cell as in *pch2-1* mutants (Methods). This arrangement differed from wild-type
simulations where only five SC segments (corresponding to the five fully synapsed bivalents) were
simulated for each cell. Thus, each SC segment was effectively treated as a discrete bivalent within
our simulations (Figure 5 – figure supplement 1). As no HEI10 foci were detected outside SC
segments, no RIs were placed there, and these parts of the chromosomes were not included in our
simulations.

In *pch2-1* simulations, the total length of the SC per cell is shorter (roughly one-quarter in length)
than in wild-type, but the HEI10 amount is equivalent to that in wild-type simulations. This resulted
in an overall higher (roughly four times) concentration of HEI10 per μm of SC and at RIs. This is akin
to the situation in our HEI10 over-expressor simulations with SC-mediated coarsening (Morgan et al.,
2021). However, as with HEI10 over-expressor simulations, this ~4-fold increase in HEI10
concentration per μm of SC led to a smaller relative increase in the number of COs per unit length
because SC-mediated coarsening dynamics proceed more rapidly when the RIs with high levels of
HEI10 are closer together, as diffusion between them is easier. More rapid dynamics lead to
relatively fewer COs, as coarsening proceeds further in the same time period, with HEI10
accumulating in a smaller number of bright foci. We therefore reduced the rates of HEI10 exchange
between the RI and SC in order to slow the coarsening process, with an uptake rate of 20% and an
escape rate of 15% of the rates in the original SC-mediated coarsening model. Changes to SC
structure in the synaptic segments that occur in *pch2* mutants, versus wild-type SC, could account
for these modified exchange rates. These changes were combined with nucleoplasmic recycling at a
rate of 10% of that in the nucleoplasmic coarsening model. These simulations were then able to
qualitatively reproduce the patterning effects observed in our experimental data (Figure 5B-G, right
hand plots), although we note that agreement between experimental data and model outputs was
slightly less good for *pch2-1* than for the *zyp1* mutant. Specifically, using the *pch2-1* parameter
values we could generate an average of 11.0 (σ^2 = 5.2) COs per cell and 0.8 (σ^2 = 0.6) COs per SC
segment, which is similar to the mean and variance values in our imaging data (Figure 5B-C). We also
found that about 40% of simulated segments had no COs, which is similar to the proportion in our
experimental data. Additionally, we recapitulated the relatively flat distribution of COs within SC
segments (albeit with a slight peak in more central positions in simulation outputs that is not present
in the experimental data, Figure 5D), as well as reproducing the spacing between adjacent COs
(Figure 5E), and the positive correlations between both CO number (Figure 5F) and single late-HEI10
focus relative intensity (Figure 5G) and SC length.

Discussion

It was recently shown that the synaptonemal complex (SC), a highly conserved and prominent
feature of meiotic prophase I, is required to promote CO interference and CO assurance in
Arabidopsis thaliana (Capilla-Pérez et al., 2021; France et al., 2021). However, the mechanistic
details of precisely how the SC plays a role in mediating interference remained obscure. Here, using
a combination of modelling and super-resolution microscopy, we have shown that CO patterning in
the absence of an SC is consistent with a coarsening model for CO interference. Previously, we
introduced a mechanistic, mathematical coarsening model that could explain CO patterning in wild-
type *Arabidopsis*, as well as in HEI10 over and under-expressor lines (Morgan et al., 2021). Here, we
make just one major change to the model for SC mutants: that HEI10 can now exchange through a
nucleoplasmic pool rather than being restricted to individual chromosomes, as in the wild-type. With
this biologically plausible modification, the coarsening model is now also capable of explaining why
zyp1 mutants lack CO assurance and CO interference in *Arabidopsis*. Additionally, the model
successfully predicts previously unexplored observations in *zyp1* SC mutants, including the pattern of
closely-spaced COs and the absence of an anticorrelation between HEI10 focal intensity and focus
number per chromosome pair. A similar function of the SC, in preventing the diffusion of
recombination proteins via the nucleoplasm, has been proposed in *C. elegans* (Rog et al., 2017;
Zhang et al., 2021, 2018). However, unlike in *C. elegans* where successful CO formation depends
upon SC formation (Colaiácovo et al., 2003), in *Arabidopsis* SC formation and CO formation can be
uncoupled (Capilla-Pérez et al., 2021; France et al., 2021), making *Arabidopsis* an excellent system

for demonstrating the effects of nucleoplasmic coarsening. Overall, our work highlights the critical
role of the SC in controlling the spatial compartment through which HEI10 diffuses.

Recently, a study in *Arabidopsis* also successfully utilised the mathematical coarsening model
(originally derived in Morgan et al., 2021) to explain genetic CO patterning data from wild-type and
HEI10 overexpressing *Arabidopsis* (Durand et al., 2022). The authors also hypothesised that
nucleoplasmic coarsening could explain CO patterning in *zyp1* mutants, and in *zyp1* mutant and
HEI10 overexpressing lines that form extremely high numbers of COs. However, unlike in this work,
the dynamics of nucleoplasmic coarsening were not explicitly modelled (Durand et al., 2022).

Interestingly, while *Arabidopsis zyp1* mutants lose CO interference, they still display a non-random
(sub-Poissonian) distribution of COs per cell. In wild-type cells, CO interference operating along
individual chromosomes regulates total CO numbers along each chromosome. In turn, the combined
effect of interference operating along all chromosome pairs within a cell causes global regulation of
the total number of COs within a cell, with the variance in total CO number per cell being close to
the sum of the per chromosome variances. In some species, COs have been shown to co-vary,
meaning a high CO number on one chromosome generally corresponds with high CO numbers on
other chromosomes within the same cell. This leads to a broader distribution of CO numbers per cell
than might be expected (Wang et al., 2019). However, recent results suggest that CO co-variation is
not present in *Arabidopsis* (Durand et al., 2022). *zyp1* mutants lose CO interference but retain per-
cell constraints on total CO numbers, as evidenced by the Poissonian and sub-Poissonian
distributions of total CO numbers on individual chromosome pairs and in individual cells,
respectively. In other words, in the *zyp1* mutant, COs designated on different RIs are spatially
independent from one another but their numbers are not independent. As noted in (Crismani et al.,
2021), an explanation for why CO numbers are relatively low (but still higher that wild-type) in SC
mutants could be due to the influence of a limiting trans-acting factor, such as HEI10. Similarly, in
our modified coarsening model, the limited nucleoplasmic pool of HEI10 explains why the
distribution of total CO numbers per cell remains sub-Poissonian. Importantly, caution should be
exercised when interpreting sub-Poissonian or Poissonian distributions of total CO number per cell
as signs of functional or non-functional CO interference, respectively. As we show here, a sub-
Poissonian distribution is perfectly compatible with non-functional CO interference.

Importantly, as well as explaining CO patterning in mutants without synapsis, we have shown that a
version of the coarsening model that incorporates both nucleoplasmic and SC-mediated coarsening
is still fully capable of explaining CO patterning in wild-type and HEI10 overexpressing *Arabidopsis.*
This builds on a previous version of the coarsening model, where HEI10 diffusion was restricted
exclusively to the SC (Morgan et al., 2021). We anticipate that this new, combined coarsening model
likely better reflects the biological reality of the coarsening process, with the SC promoting and
enhancing the diffusion of HEI10 along individual bivalents but with some exchange of HEI10
molecules between the SC and the nucleoplasm. This leakage is incorporated here as exchange
between RIs and the nucleoplasm, but it is straightforward to also include direct exchange between
the SC and the nucleoplasm in the framework of this model.

The complex interplay between SC-mediated and nucleoplasmic coarsening can also provide a
potential mechanistic explanation for other crossover phenomena, such as the interchromosomal
effect (Miller, 2020; Termolino et al., 2019).The interchromosomal effect describes the observation
that when one pair of homologs experiences a reduction in crossover frequency (due to a structural
variation, such as an inversion) this is accompanied by a corresponding increase in crossover
frequency on the other, structurally normal, homolog pairs (Miller, 2020; Termolino et al., 2019). In
this scenario, RIs may be inhibited from occurring normally at the sites of structural variations,

leading to a lower level of HEI10 in those regions. In turn, this could lead to a redistribution of the
limited pool of HEI10 to the other chromosomes, thus increasing the likelihood of those
chromosomes receiving extra COs.

Additionally, we have shown that HEI10-coarsening can explain aspects of CO patterning in *pch2*
mutants that exhibit partial, but incomplete, synapsis (Lambing et al., 2015). In this scenario,
available HEI10 molecules could still load onto the SC and undergo coarsening, similar to the
situation in the wild-type. However, if a wild-type level of HEI10 protein is loaded onto shorter
stretches of SC, this results in an increased concentration of HEI10 per micron of SC and, akin to the
situation in an otherwise wild-type HEI10 over-expressor (Morgan et al., 2021), an increase in
closely-spaced COs within those stretches. An abundance of closely spaced COs have also been
reported in other *Arabidopsis* mutants that exhibit partial synapsis, such as *axr1*, *pss1* and *asy1*
(Duroc et al., 2014; Jahns et al., 2014; Lambing et al., 2020; Pochon et al., 2022), suggesting that the
coarsening model could explain CO patterning in a variety of synaptic mutants.

Together, the results we present here further reinforce the ability of the coarsening paradigm to
explain numerous aspects of CO control in both wild-type and mutant *Arabidopsis* lines. These
aspects include: CO placement, CO frequency, CO assurance, CO interference, CO homeostasis,
heterochiasmy, meiotic duration, interchromosomal effects and diverse HEI10 focal intensity
measurements. Given the conserved nature of HEI10 (Agarwal and Roeder, 2000; Chelysheva et al.,
2012; Jantsch et al., 2004; Lake et al., 2015; Qiao et al., 2014; Reynolds et al., 2013), and the
discovery of similar coarsening mechanisms in *C. elegans* (Zhang et al., 2021), it is likely that similar
mechanisms regulate these same crossover patterning phenomena in a wide variety of sexually
reproducing eukaryotes.

Acknowledgments

We gratefully acknowledge James Higgins for providing the *zyp1a-2/zyp1b-1* mutant seeds, Chris
Franklin for supplying the SMC3, HEI10 and ZYP1 antibodies, Eva Wegel for microscopy support and
members of the M. Howard and X. Feng labs for fruitful discussions on this topic. This work was
supported by a BBSRC Discovery Fellowship (BB/V005774/1) awarded to C.M. and by BBSRC Institute
Strategic Programme GEN (BB/P013511/1) to M.H.

Materials and methods

Key Resources Table

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
gene (*Arabidopsis*)	ZYP1A	TAIR	AT1G22260	
gene (*Arabidopsis*)	ZYP1B	TAIR	AT1G22275	

gene (*Arabidopsis*)	PCH2	TAIR	AT4G24710	
strain, strain background (*Arabidopsis*)	*zyp1a-2/zyp1b-1*	France *et al.,* 2021		Supplied by Dr James Higgins, University of Leicester
strain, strain background (*Arabidopsis*)	*pch2-1*	Syngenta Arabidopsis Insertion Library (SAIL)	SAIL_1187_C06	
antibody	anti-HEI10 (rabbit polyclonal)	Lambing et al., 2015		Supplied by Prof Chris Franklin, University of Birmingham (1:500 dilution)
antibody	anti-SMC3 (rat polyclonal)	Ferdous et al., 2012		Supplied by Prof Chris Franklin, University of Birmingham (1:500 dilution)
antibody	anti-ZYP1 (guinea-pig polyclonal)	France et al., 2021		Supplied by Prof Chris Franklin, University of Birmingham (1:500 dilution)
antibody	alexa fluor 555 goat anti-rabbit (goat polyclonal)	Thermo Fisher	RRID AB_2535849	(1:200 dilution)
antibody	alexa fluor plus 488 goat anti-rat (goat polyclonal)	Thermo Fisher	RRID AB_2896330	(1:200 dilution)
antibody	alexa fluor 647 goat anti-guinea-pig (goat polyclonal)	Thermo Fisher	RRID AB_2735091	(1:200 dilution)
sequence-based reagent	PCH2_1_FV	Lambing et al., 2015	PCR primers	CAGTGCAAATAGCCGTCGCTGAG
sequence-based reagent	PCH2_1_RV	Lambing et al., 2015	PCR primers	CTCACATGGTCCTTCTTCAATGAGC
sequence-based reagent	Sail LB2	Lambing et al., 2015	PCR primers	GCTTCCTATTATATCTTCCCAAATTACCAATACA
sequence-based reagent	zyp1_ns_1	France et al., 2021	PCR primers	CTCGCATTTGCTGGTTTAAAGAGTC
sequence-based reagent	zyp1b_sp_1	France et al., 2021	PCR primers	TGCGTATATTGCTAGGTTTATATTG

sequence-based reagent	salk_lb2	France et al., 2021	PCR primers	GTGCTTTACGGCACCTCGAC
sequence-based reagent	zyp1a_sp_1	France et al., 2021	PCR primers	GAATAGTTAGCAGATTCATATTT CAC
peptide, recombinant protein	HindIII-HF	NEB	R3104S	
chemical compound, drug	cytohelicase	Sigma-Aldrich	C8274	
chemical compound, drug	polyvinylpyrrolidone	Sigma-Aldrich	PVP40	
software, algorithm	FIJI	Schindelin et al. 2012	2.1.0/1.53f51	
software, algorithm	Zen Black	Zeiss	14.0.12.201	
software, algorithm	Python	Python	RRID:SCR_008394	https://www.python.org/
software, algorithm	R	R Project for Statistical Computing	RRID:SCR_001905	http://www.r-project.org/
software, algorithm	Julia	Julia	RRID:SCR_021666	https://julialang.org/

Plant Materials

463 *Arabidopsis thaliana* lines used in this study were wild-type Col-0, *pch2-1* (SAIL_1187_C06) and the
464 *zyp1a-2/zyp1b-1* null-mutant (previously characterised in (France et al., 2021)). The *zyp1a-2/zyp1b-1*
465 line contains a CRISPR/Cas9 derived 14 base-pair deletion in exon 3 of *ZYP1A* and a T-DNA insertion
466 in exon 3 of *ZYP1B.* Plants were grown in controlled environment rooms with 16 h of light (125 mMol
467 cool white) at 20 °C and 8 h of darkness at 16 °C.

Immunocytology

Immunocytology of spread *Arabidopsis thaliana* pachytene cells was performed as described
previously (Morgan et al., 2021; Morgan and Wegel, 2020). In brief, staged anthers were dissected
from *Arabidopsis* floral buds and macerated in a 10 μl drop of digestion medium (0.4% cytohelicase,
1.5% sucrose, 1% polyvinylpyrolidone in sterile water) on a No. 1.5H coverslip (Marienfeld).
Coverslips were then incubated in a moist chamber at 37 °C for 4 minutes before adding 10 μl of 2%
lipsol solution followed by 20 μl of 4% paraformaldehyde (pH 8). Coverslips were dried for 3 hours in
the fumehood and then blocked in blocking buffer (0.3% BSA in 1 x PBS) for 15 minutes. 50 μl of
primary antibody, diluted in blocking buffer, was added to the coverslips and they were then

incubated overnight in a moist chamber at 4 °C. 50 µl of secondary antibody, diluted in blocking
buffer, was added to the coverslips and they were then incubated in a moist chamber at 37 °C for 2
hours. Coverslips were then incubated in 10 µl DAPI (10 µg/ml) for 5 min before adding 7 µl of
Vectashield and mounting them on a glass slide. Coverslips were washed for 3 x 5 minutes in 1 X PBS
0.1% Triton before and after each incubation step. The following primary antibodies were used at
1:500 dilutions: anti-SMC3 (rat) (Ferdous et al., 2012), anti-HEI10 (rabbit) (Lambing et al., 2015) and
anti-ZYP1 (guinea-pig) (France et al., 2021). The following secondary antibodies were used at 1:200
dilutions: anti-rat Alexa Fluor Plus 488 (ThermoFisher A48262), anti-rabbit Alexa Fluor 555
(ThermoFisher A21428) and anti-guinea-pig Alexa Fluor 647 (ThermoFisher A21450). Immunostained
cells were imaged using 3D structured illumination microscopy (3D-SIM) on a Zeiss Elyra PS1
microscope equipped with an EM-CCD camera, a Plan-Apochromat ×63, NA 1.40 oil objective and
405, 488, 561 and 642 nm solid-state laser diodes. Cells were imaged with three stripe angles and
five phases and z-stacks were captured at an interval size of 0.0909 µm. An immersion oil with a
refractive index of 1.515 was used that was optimised for the green/red (SMC3/HEI10) channels of
our system. For optimal image quality and to minimise the introduction of reconstruction artefacts,
microscope laser power and camera gain values were manually adjusted for each cell. Bleaching and
contrast of raw images was assessed using the SIMcheck plugin to FIJI (Ball et al., 2015).

Image analysis

For wild-type and *zyp1a-2/zyp1b-1* nuclei, individual axes were traced and segmented in 3-
dimensions from 3D-SIM z-stacks using the SNT plugin to FIJI (Arshadi et al., 2021). Traces were
made using the linear SMC3 signal, which localises along the meiotic axis, and 3D-skeletons of each
axis-traced path were generated using SNT. For *zyp1a-2/zyp1b-1* nuclei, each axial element was
traced end-to-end to generate 10 complete skeleton traces for each nucleus. For wild-type nuclei,
due to the proximity of the two parallel SMC3 signals, a single trace was used to segment the paths
of each pair of synapsed axial elements, generating 5 complete skeleton traces for each nucleus. Foci
were detected within images using the FociPicker3D Fiji plugin (Du et al., 2011). Uniform background
intensity thresholds and minimum pixel size values were manually optimised within a range of
6,000-20,000 and 5-20, respectively, with otherwise default parameters. Further analysis was
performed using custom Python scripts. For *zyp1a-2/zyp1b-1* nuclei, foci were assigned to the
chromosome pair for which the average of the two distances from the closest points on each of the
axes was smallest, provided this was less than a threshold distance (0.5 µm). If this was not the case,
then the foci was assigned to the closest chromosome pair, provided that it was nearer to one of the
two axes than a smaller threshold (0.3 µm). Foci satisfying neither of these two criterion were
considered as off target signal and ignored. For wild-type nuclei, each focus was assigned to the
closest trace, provided that it was nearer than a threshold (0.5 µm). Again, foci not satisfying this
criterion were considered as off target signal and ignored. Relative positions along chromosome
pairs were calculated by dividing the distance along the traced axis (measured in terms of voxels) by
the total length of the trace. Foci intensities were calculated using the mean intensity within a
sphere of radius 0.1985 µm (5 voxels in the x and y planes), centred at the foci peak, and the median
intensity of the image (used as an estimate of the background fluorescence) was subtracted. All
intensity values used were normalized by dividing the foci intensity by the mean of the intensities of
all foci within the same cell (only considering foci associated with a chromosome pair). Simple linear
regression was performed using the Python statsmodels package.
520
For *pch2-1* nuclei, segments of synapsis were traced and segmented in 3-dimensions from 3D-SIM z-
stacks using the SNT plugin to FIJI (Arshadi et al., 2021). Traces were made using the linear ZYP1
signal, which localises along the SC, and 3D-skeletons of each SC-traced path were generated using

SNT. Each SC segment was traced end-to-end to generate between 8 and 21 complete skeleton
traces for each nucleus. Individual traces varied from 0.4 to 28.4 µm in length. Again, foci were
detected within images using the FociPicker3D Fiji plugin (Du et al., 2011). Uniform background
intensity thresholds and minimum pixel size values were manually optimised within a range of
10,000-15,000 and 10-20, respectively, with otherwise default parameters. Further analysis was
performed using custom Python scripts. All subsequent analysis proceeded as for wild-type nuclei.

Mathematical models

Nucleoplasmic coarsening model

In each cell, $Q = 5$ chromosome pairs are simulated. The q-th chromosome pair is taken to have a
length L_q, sampled from the (per-chromosome) normal distribution $\mathcal{N}_T(\mu_q, \sigma_q^2, -3,3)$ truncated at 3
standard deviations from the mean (to ensure positive chromosome lengths). We use the same
values as in our earlier work (Morgan et al., 2021), fitted to experimentally measured wild-type
chromosome length distributions: $\mu_1 = 37.5, \mu_2 = 40.4, \mu_3 = 45.7, \mu_4 = 53.8, \mu_5 = 59.8$, and $\sigma_1 =$
$4.93, \sigma_2 = 5.06, \sigma_3 = 5.86, \sigma_4 = 7.18, \sigma_5 = 7.13$, with all these parameters having units of μm. We
chose to retain these original values as they were broadly consistent with the experimental
chromosome lengths measured in this study, but were calculated from a greater sample size.

Recombination intermediates (RIs) are placed randomly along each of the chromosome pairs, with a
probability density ρ per µm. To implement this, for each simulated chromosome pair with length L,
a random number is generated from a Poisson process with mean ρL, and then this number of RIs
are placed uniformly at random along that chromosome pair. We note that this procedure differs
slightly from the method used in our original SC coarsening model, where floor(ρL) crossovers were
placed on a chromosome with length L. Simulation outputs with this original rule (not shown) differ
only slightly.

The pro-crossover factor HEI10 is present in a nucleoplasmic pool, with amount P, and at
compartments at each of the RIs, with amounts $C_{q,n}, 1 \leq q \leq Q = 5, 1 \leq n \leq N_q$ for n-th RI on the
q-th chromosome pair. HEI10 is able to move between the nucleoplasmic pool to each of the RI
compartments at rate α, and to escape from the RI compartments back into the pool at a rate
$\beta(C_{q,n})$, which depends on the amount of HEI10 within each individual RI compartment. Figure 1D
shows the functional form of this rate in more detail, from which we see that it is a strictly
decreasing function of the RI HEI10 amount $C_{q,n}$. As in our SC interference model, we take

$$\beta(C_{q,n}) = \frac{\beta_C}{1 + (C_{q,n}/K_C)^\gamma}.$$

555 The system of ordinary differential equations governing HEI10 amounts is

$$\frac{dC_{q,n}}{dt} = (\alpha P - \beta(C_{q,n})C_{q,n})H_s(C_{q,n} - K_H), \qquad 1 \leq q \leq Q, 1 \leq n \leq N_q,$$

$$\frac{dP}{dt} = \sum_{q=1}^{Q} \sum_{n=1}^{N_q} H_s(C_{q,n} - K_H)(\beta(C_{q,n})C_{q,n} - \alpha P).$$

Compared to our earlier model (Morgan et al., 2021), we also incorporated a further modulation to
the HEI10 exchange rates between the nucleoplasm and RIs, through the multiplicative smoothing
function H_s. For HEI10 levels in an RI well below a threshold K_H, the smoothing function becomes
close to zero, while for levels well above, the function saturates to one, with a smooth crossover in

between. This feature ensures that RIs with HE10 levels well below the threshold become
disconnected, with their HEI10 effectively disappearing from the rest of the system. This reflects the
experimental observation that smaller HEI10 foci are not detected in late-pachytene nuclei (Morgan
et al., 2021), with RIs that lack sufficient HEI10 presumably being channelled into a non-crossover or
class II repair pathway. We implement this through the smooth cutoff function H_s :

567
$$H_s(s) = \frac{1}{2}\left(1 + \tanh(K_s s)\right)$$

which multiplies the HEI10 exchange rates between the RIs and nucleoplasm. Overall, this system
conserves the total amount of HEI10 within each cell (the nucleoplasmic pool plus all RI
compartments), rather than within each SC, which was the case for our earlier SC-mediated
coarsening model (Morgan et al., 2021).

Unlike in our previous SC-mediated coarsening model, where the total HEI10 amount in each cell
depended on the total length of SC and the total number of RIs (which are relatively constant from
cell to cell), we adopted an alternative approach for controlling cellular HEI10 amount in our
simulations. This change was necessary due to the absence of an SC in the nucleoplasmic-coarsening
model, and is likely more realistic, with cellular HEI10 protein amount being controlled
independently of SC formation, as we now describe.

The initial total amount of HEI10 within each cell, F, is sampled from a truncated normal distribution

579
$$F = \mathcal{N}_T(\mu_F, \sigma_F^2, -3,3),$$

where the mean μ_F and standard deviation σ_F are taken to be the mean and standard deviation of
the total amounts of HEI10 within each cell in the original SC-mediated coarsening model (Morgan et
al., 2021). This initial cellular level of HEI10 is divided between the RIs and the nucleoplasmic pool, at
constant proportions R_{RI} and R_P, respectively, with $R_{RI} + R_P = 1$.

The total amount of RI HEI10 is then randomly divided between each RI, where, for each RI, a
random weighting $\hat{C}_{q,n}$ is generated from

586
$$\hat{C}_{q,n} = f\left(\frac{x_{q,n}}{L_q}\right) X_{q,n}, \qquad X_{q,n} = \mathcal{N}_T(1, \sigma_C^2, -3,3), \qquad 1 \le q \le Q, \, 1 \le n \le N_q,$$

587 where $\mathcal{N}_T$ again denotes a truncated normal distribution and $x_{q,n}$ is the position of the n-th RI on the
588 q-th chromosome pair. The piecewise function $f(x/L)$, included to account for the increased
589 frequency of crossovers towards the end of each chromosome, is defined as

590
$$f(\xi) = \begin{cases} 1 + (f_e - 1)(1 - \xi/x_e) & 0 \le \xi \le x_e, \\ 1 & x_e \le \xi \le 1 - x_e, \\ 1 + (f_e - 1)(\xi - 1 + x_e)/x_e & 1 - x_e \le \xi \le 1. \end{cases}$$

591 The total initial amount of HEI10 at all RIs, $R_{RI}F$, is then loaded at the RIs in proportion to these
592 weights, namely

594
$$C_{q,n}(t = 0) = R_{RI}F \frac{\hat{C}_{q,n}}{\sum_{q'=1}^{Q} \sum_{n'=1}^{N_q} \hat{C}_{q',n'}}.$$

The remaining HEI10 is placed in the nucleoplasmic pool, with initial amount $P(t = 0) = R_P F$, which
has the same average as the average amount of HEI10 loaded onto all the SCs in the original SC-
mediated coarsening model (Morgan et al., 2021).

This system is simulated for a duration $T = 10$hr, a duration of pachytene equivalent to that used in
our original model (Prusicki et al., 2019). Numerical simulations are performed using the Rodas5
solver (di Marzo, 1992) of the DifferentialEquations.jl Julia package (Rackauckas and Nie, 2017), with
a non-negativity constraint on HEI10 amounts. The initial conditions for the simulation are random,
but the evolution of the system is deterministic. Multiple simulations (10000) with different random
initial conditions are performed to determine the distribution of crossover numbers, positions and
intensities. At the end of each simulation, RIs are designated as crossover sites according to a
modified rule (see "Modified CO criterion" below).

All parameters were chosen to be the same as those as in the original SC coarsening model (Morgan
et al., 2021), except for the RI HEI10 absorption rate α, and the (maximum) escape rate β_C, which
was multiplied by a factor of 0.06 (Figure 1E). These changes reflect that HEI10 is now being
exchanged with the nucleoplasmic pool, rather than locally on the synaptonemal complex.

For simulations of the *zyp1* HEI10 over-expressor line, a 3.5-fold change in HEI10 levels was used, but
a larger fold change was chosen for the standard deviation σ_F of total cellular HEI10 amounts, in
order to generate greater cell-cell variability, with an increase from 75 a.u. to 860 a.u. We also
verified that increasing the noise to this degree in the original model of (Morgan et al., 2021) (as
implemented below) for wild-type and HEI10 overexpressing lines, did not make a substantial
difference to model predictions for the number of COs per SC, or the positioning of COs. We note
that the above 3.5-fold change in HEI10 levels resulted in roughly the same relative change in CO
number, unlike for SC-mediated coarsening where 4.5 times the level of HEI10 only resulted in
approximate doubling of the number of COs (Morgan et al., 2021). This reduced change reflects the
reduced sensitivity of the SC-mediated coarsening to changes in HEI10 levels, compared to
nucleoplasmic-mediated coarsening (for these choices of parameter values).

SC-mediated coarsening model

This model is as described in our earlier work, but with the introduction of the smooth cutoff
function preventing recycling at RIs with low levels of HEI10, with slightly different initial conditions,
and with a new rule for determining which RIs are crossovers at the end of the simulation. Again,
$Q = 5$ chromosomes are simulated in each cell, these having SCs with lengths L_q generated
randomly as in the nucleoplasmic coarsening model. HEI10 is present on each of the SCs with
concentration $c_q(x, t)$ per unit length, and is able to diffuse along them with (one-dimensional)
diffusion coefficient D. RI numbers and positions are generated as in the nucleoplasmic model.
Again, we note that this is slightly different to the rule used in (Morgan et al., 2021), but that this
change has limited effect on model outputs. The variables $C_{q,n}$ represent the amount of HEI10
associated with the n-th RI on the q-th SC. However, RIs are not able to exchange HEI10 with the
nucleoplasm, but instead exchange HEI10 with the relevant SC, with uptake rate $\hat{\alpha}$ and escape rate
$\hat{\beta}(C_{q,n})$, this escape rate taking the same functional form as in the nucleoplasmic coarsening model
but with the new parameter $\hat{\beta}_C$. This gives the system of equations

635
$$\frac{dC_{q,n}}{dt} = \left(\hat{\alpha} c_q(x_{q,n}) - \hat{\beta}(C_{q,n}) C_{q,n} \right) H_s(C_{q,n} - K_H), \qquad 1 \leq q \leq Q, \, 1 \leq n \leq N_q,$$

636
$$\frac{\partial c_q}{\partial t} = D \frac{\partial^2 c_q}{\partial x^2} + \sum_{n=1}^{N_q} \left(\hat{\beta}(C_{q,n}) C_{q,n} - \hat{\alpha} c_q (x_{q,n}) \right) H_s(C_{q,n} - K_H) \delta(x - x_{q,n}),$$

637
$$1 \le q \le Q, \, 0 \le x \le L_q,$$

638
$$\frac{\partial c_q}{\partial x} = 0, \text{ at } x = 0 \text{ and } x = L_q,$$

639
$$c_q = c_0 \text{ at } t = 0.$$

640 Initial conditions for total cellular HEI10 levels are as in the nucleoplasmic model. Initial RI HEI10
641 amounts are as in the nucleoplasmic model, but with different parameters (see Figure 4 –
642 supplementary figure 1) in the non-uniform loading function $f(\xi)$. However, instead of being placed
643 in the nucleoplasmic pool, the remaining HEI10 is placed uniformly on the SCs, giving

644
$$c_q(t = 0) = c_0 = \frac{R_{SC} F}{\sum_{q'=1}^{Q} L_{q'}},$$

where $R_{SC} = 1 - R_{RI}$. Note that these initial conditions differ from those used in (Morgan et al.,
2021), but when the new rule for determining crossovers is used, they give very similar results in
terms of crossover number and location. This model was used to produce the simulation output
shown in Figure 4 – figure supplement 2 (WT, OX), using the parameters shown in Figure 4 - figure
supplement 1D (WT column), except for the over-expressor OX case, where the parameters μ_F and
σ_F controlling initial cellular HEI10 levels were increased to 4.5x their WT values.

Combined coarsening model

The combined model includes all the above rules in the nucleoplasmic and SC-mediated coarsening
models, with HEI10 being present in RI, SC and nucleoplasm compartments. The governing equations
are a combination of those in the two other models; explicitly,

655
$$\frac{dC_{q,n}}{dt} = \left(\alpha P - \beta(C_{q,n}) C_{q,n} + \hat{\alpha} c_q (x_{q,n}) - \hat{\beta}(C_{q,n}) C_{q,n} \right) H_s(C_{q,n} - K_H),$$
656
$$1 \le q \le Q, \, 1 \le n \le N_q,$$

657
$$\frac{\partial c_q}{\partial t} = D \frac{\partial^2 c_q}{\partial x^2} + \sum_{n=1}^{N_q} \left(\hat{\beta}(C_{q,n}) C_{q,n} - \hat{\alpha} c_q (x_{q,n}) \right) H_s(C_{q,n} - K_H) \delta(x - x_{q,n}),$$

658
$$1 \le q \le Q, \, 0 \le x \le L_q,$$

659
$$\frac{dP}{dt} = \sum_{q=1}^{Q} \sum_{n=1}^{N_q} (\beta(C_{q,n}) C_{q,n} - \alpha P) H_s(C_{q,n} - K_H).$$

Initial and boundary conditions for wild-type or HEI10 over-expressor plants are as in the SC-
mediated coarsening model above. No HEI10 is placed initially in the nucleoplasm. Each SC is
discretized into m=2000 compartments, and the spatial derivatives approximated by finite
differences. This model was used to generate the simulation output shown in Figure 4 – figure
supplement 2 (WT+nuc, OX+nuc), using the parameters shown in Figure 4 - figure supplement 1D
(WT+nuc column), except for the over-expressor OX+nuc case, where the parameters μ_F and σ_F
controlling initial cellular HEI10 levels were increased to 4.5x their WT values.

For the *pch2-1* mutant, we have a variable number of SC segments within the cell, which are here
treated as distinct SCs. The number of SC segments is drawn randomly from a Poisson distribution
with mean 13.92 (close to that measured experimentally), while the length of each SC segment is
drawn from a Gamma distribution with parameters $L_\alpha = 1.693$, $L_\theta = 2.984$ μm, so having mean
$L_\alpha L_\theta \approx 5.1$μm and variance $L_\alpha L_\theta^2 \approx 15.1$μm^2. This distribution has probability distribution function

677
$$f_\Gamma(x; L_\alpha, L_\theta) = \frac{x^{L_\alpha - 1} e^{-x/L_\theta}}{\Gamma(L_\alpha) L_\theta^{L_\alpha}}, \quad x > 0,$$

where $\Gamma(x)$ is the Gamma function, and was fitted to experimental measurements of SC segment
length. This statistical model for SC segment lengths introduced substantially greater variation in
total cellular SC length than in the other simulations, which motivated our new rule for the
assignment of initial HEI10 amounts. Initial and boundary conditions are the same here as for the SC-
mediated coarsening model above (although there are a larger number of SCs here).

Owing to this large number of SCs, all of which are coupled with each other through the
nucleoplasm, for efficient simulation each SC segment is spatially discretized into $m = 200$ portions,
fewer than the number ($m = 2000$) used for each SC in the SC-mediated coarsening model above.

The smooth cutoff function H_s is included so that RIs with low levels of HEI10 are unable to interact
with the nucleoplasmic pool. This is necessary, as otherwise these RIs provide an additional route for
the transfer of HEI10 between two RIs on the same SC: HEI10 can diffuse along the SC from a RI
losing HEI10 to a nearby RI with low levels of HEI10, escape into the nucleoplasm, be reabsorbed at
another RI with low levels of HE10, and then diffuse along the SC to an RI gaining HEI10. This route
causes the dynamics to progress too quickly.

The parameterization of this combined model used for the *pch2-1* mutant has substantially smaller
values for the rates of exchange of HEI10 between RIs and SCs ($\hat{\alpha}$ and $\hat{\beta}_C$) than in the original SC-
mediated coarsening model (Figure 5 – Figure supplement 1C). These were chosen to better fit the
experimental data, which shows a less pronounced spatial pattern than the wild-type data.
Nucleoplasmic recycling was also set at a rate of 10% of that in the nucleoplasmic coarsening model.
This model was used to generate the simulation output shown in Figure 5.

SC-mediated coarsening model analysis

In the SC-mediated coarsening model, as the timescale for diffusion along the SC ($\sim L^2/D$, which is
at most $(60$μm$)^2/1.1$μm^{2}s$^{-1} \approx 3300$s) is small compared with the duration of pachytene (36000s),
the HEI10 concentration adopts a piecewise linear profile along each SC, with discontinuities in its
slope at each RI. When there are just two RIs with HEI10 amounts C_1 and C_2, separated by a distance
$\Delta x = x_2 - x_1$, we have that

700
$$\frac{dC_1}{dt} = \hat{\alpha} c(x_1) - \hat{\beta}(C_1) C_1 = -J, \quad \frac{dC_2}{dt} = \hat{\alpha} c(x_2) - \hat{\beta}(C_2) C_2 = J,$$

699 where the net flux of HEI10 from C_1 to C_2, J, is given by

701
$$J = D \frac{c(x_1) - c(x_2)}{\Delta x}.$$

702 On eliminating $c(x_1)$ and $c(x_2)$, the concentrations of HEI10 on the SC at the two RIs, we find that

703
$$\frac{d}{dt}(C_2 - C_1) = \frac{\hat{\beta}(C_1) C_1 - \hat{\beta}(C_2) C_2}{1 + \frac{\hat{\alpha} \Delta x}{2D}}.$$

We can see from this equation that closely spaced pairs of RIs, with small Δx, coarsen more quickly
than those further apart. This effect generates spatial patterns in CO positioning, even in the
absence of non-uniform initial HEI10 loading. However, this difference in coarsening rate is only
significant if $\frac{\hat{\alpha}\Delta x}{2D}$ is not small compared with 1. In the limit where $\frac{\hat{\alpha}\Delta x}{2D}$ is small, the spatial patterns in
CO positioning vanish, other than those spatial biases from non-uniform initial HEI10 loading.

With $K_C = 1$ and for $C_1, C_2 \gg 1$ and $\frac{\hat{\alpha}\Delta x}{2D} \gg 1$, which is the case for our original SC-mediated
coarsening parameters, this can be approximated further as

$$711 \qquad \frac{d}{dt}(C_2 - C_1) = \frac{2\,D\,\hat{\beta}_C}{\hat{\alpha}\Delta x}\left(C_1^{1-\gamma} - C_2^{1-\gamma}\right).$$

From this equation, we find that coarsening proceeds faster for larger diffusion coefficients, larger
escape rates from RIs to the SC, and slower for larger uptake rates from the SC to RIs, more widely
spaced RIs, and when HEI10 levels are greater (as $\gamma > 1$). When more than two RIs are participating
in coarsening, a more complex system of equations is obtained, but again containing the model
parameters in the same combination $\frac{D\,\hat{\beta}_C}{\hat{\alpha}}$. As a consequence of this, it is difficult from simulation
outputs to independently identify the parameters $\hat{\alpha}$, $\hat{\beta}_C$ and D, with many parameter sets being
potentially compatible with our original findings.

Modified crossover criterion

In our original simulations (Morgan et al., 2021) of SC-mediated coarsening in wild-type Arabidopsis
thaliana, we used a per-SC criterion to decide which RIs at the end of the simulation should be
designated as crossovers. This criterion was chosen to be a threshold of 40% of the maximum HEI10
amount at those RIs on the same SC. However, this criterion needed to be changed to study
nucleoplasmic-mediated coarsening, and a simple absolute threshold would require substantial
reparameterization of the HEI10 model.

Instead, we designed a new per-cell criterion. RIs were designated as crossovers if they contained an
amount of HEI10 more than 60% of the mean amount of HEI10 at an RI in that cell, where this mean
excluded RIs with less HEI10 than an absolute threshold of 1 a.u. A per-cell threshold based on the
mean amount rather than the maximum amount was used because of the substantial variation in
maximum HEI10 level found in simulations of the *pch-1* mutant. All simulation outputs in this
manuscript use this adjusted threshold. We found that this made only a limited change to the results
of simulating SC-mediated coarsening in WT and HEI10 overexpressing cells.

Data Availability

Imaging data associated with this study are available at doi.org/10.6084/m9.figshare.19650249.v1,
doi.org/10.6084/m9.figshare.19665810.v1 and doi.org/10.6084/m9.figshare.21989732. Custom
Python scripts for data analysis are available at https://github.com/jfozard/hei10_zyp1.

References

Agarwal S, Roeder GS. 2000. Zip3 Provides a Link between Recombination Enzymes and
Synaptonemal Complex Proteins. *Cell* **102**:245–255. doi:https://doi.org/10.1016/S0092-
8674(00)00029-5

Arshadi C, Günther U, Eddison M, Harrington KIS, Ferreira TA. 2021. SNT: a unifying toolbox for
quantification of neuronal anatomy. *Nat Methods* **18**:374–377. doi:10.1038/s41592-021-
01105-7

Ball G, Demmerle J, Kaufmann R, Davis I, Dobbie IM, Schermelleh L. 2015. SIMcheck: a Toolbox for
Successful Super-resolution Structured Illumination Microscopy. *Sci Rep* **5**:15915.
doi:10.1038/srep15915

Bhalla N. 2022. PCH-2 and meiotic HORMADs: A module for evolutionary innovation in
meiosis?Current Topics in Developmental Biology. Academic Press.
doi:https://doi.org/10.1016/bs.ctdb.2022.07.001

Cameron AC, Trivedi PK. 1990. Regression-based tests for overdispersion in the Poisson model. *J
Econom* **46**:347–364. doi:https://doi.org/10.1016/0304-4076(90)90014-K

Capilla-Pérez L, Durand S, Hurel A, Lian Q, Chambon A, Taochy C, Solier V, Grelon M, Mercier R.
2021. The synaptonemal complex imposes crossover interference and heterochiasmy in
Arabidopsis. *Proceedings of the National Academy of Sciences* **118**:e2023613118.
doi:10.1073/pnas.2023613118

Chelysheva L, Vezon D, Chambon A, Gendrot G, Pereira L, Lemhemdi A, Vrielynck N, le Guin S,
Novatchkova M, Grelon M. 2012. The Arabidopsis HEI10 is a new ZMM protein related to Zip3.
PLoS Genet. doi:10.1371/journal.pgen.1002799

Colaiácovo MP, MacQueen AJ, Martinez-Perez E, McDonald K, Adamo A, la Volpe A, Villeneuve AM.
2003. Synaptonemal Complex Assembly in C. elegans Is Dispensable for Loading Strand-
Exchange Proteins but Critical for Proper Completion of Recombination. *Dev Cell* **5**:463–474.
doi:10.1016/S1534-5807(03)00232-6

di Marzo GA. 1992. Méthodes de Rosenbrock d'ordre 5(4) adaptées aux problemes différentiels-
algébriques.

Du G, Drexler GA, Friedland W, Greubel C, Hable V, Krücken R, Kugler A, Tonelli L, Friedl AA, Dollinger
G. 2011. Spatial dynamics of DNA damage response protein foci along the ion trajectory
of high-LET particles. *Radiat Res* **176**:706–715. doi:10.1667/rr2592.1

Durand S, Lian Q, Jing J, Ernst M, Grelon M, Zwicker D, Mercier R. 2022. Joint control of meiotic
crossover patterning by the synaptonemal complex and HEI10 dosage. *Nat Commun* **13**:5999.
doi:10.1038/s41467-022-33472-w

Duroc Y, Lemhemdi A, Larchevêque C, Hurel A, Cuacos M, Cromer L, Horlow C, Armstrong SJ,
Chelysheva L, Mercier R. 2014. The Kinesin AtPSS1 Promotes Synapsis and is Required for
Proper Crossover Distribution in Meiosis. *PLoS Genet* **10**:e1004674-.

Ferdous M, Higgins JD, Osman K, Lambing C, Roitinger E, Mechtler K, Armstrong SJ, Perry R, Pradillo
M, Cuñado N, Franklin FCH. 2012. Inter-Homolog Crossing-Over and Synapsis in Arabidopsis
Meiosis Are Dependent on the Chromosome Axis Protein AtASY3. *PLoS Genet* **8**:e1002507-.

France MG, Enderle J, Röhrig S, Puchta H, Franklin FCH, Higgins JD. 2021. ZYP1 is required for
obligate cross-over formation and cross-over interference in Arabidopsis. *Proceedings of the
National Academy of Sciences* **118**. doi:10.1073/pnas.2021671118

Gray S, Cohen PE. 2016. Control of Meiotic Crossovers: From Double-Strand Break Formation to
Designation. *Annu Rev Genet*. doi:10.1146/annurev-genet-120215-035111

Haldane JBS. 1931. The Cytological Basis of Genetical Interference. *Cytologia (Tokyo)* **3**:54–65.
doi:10.1508/cytologia.3.54

Hunter N. 2015. Meiotic Recombination: The Essence of Heredity. *Cold Spring Harb Perspect Biol*
7:a016618. doi:10.1101/cshperspect.a016618

Jahns MT, Vezon D, Chambon A, Pereira L, Falque M, Martin OC, Chelysheva L, Grelon M. 2014.
Crossover Localisation Is Regulated by the Neddylation Posttranslational Regulatory Pathway.
PLoS Biol **12**:e1001930-.

Jantsch V, Pasierbek P, Mueller MM, Schweizer D, Jantsch M, Loidl J. 2004. Targeted gene knockout
reveals a role in meiotic recombination for ZHP-3, a Zip3-related protein in Caenorhabditis
elegans. *Mol Cell Biol* **24**:7998–8006. doi:10.1128/MCB.24.18.7998-8006.2004

Jones GH, Franklin FCH. 2006. Meiotic Crossing-over: Obligation and Interference. *Cell* **126**:246–248.
doi:https://doi.org/10.1016/j.cell.2006.07.010

Lake CM, Nielsen RJ, Guo F, Unruh JR, Slaughter BD, Hawley RS. 2015. Vilya, a component of the
recombination nodule, is required for meiotic double-strand break formation in Drosophila.
Elife **4**:e08287. doi:10.7554/eLife.08287

Lam WS, Yang X, Makaroff CA. 2005. Characterization of Arabidopsis thaliana SMC1 and SMC3:
evidence that AtSMC3 may function beyond chromosome cohesion. *J Cell Sci* **118**:3037–3048.
doi:10.1242/jcs.02443

Lambing C, Kuo PC, Tock AJ, Topp SD, Henderson IR. 2020. ASY1 acts as a dosage-dependent
antagonist of telomere-led recombination and mediates crossover interference in Arabidopsis.
Proceedings of the National Academy of Sciences **117**:13647–13658.
doi:10.1073/pnas.1921055117

Lambing C, Osman K, Nuntasoontorn K, West A, Higgins JD, Copenhaver GP, Yang J, Armstrong SJ,
Mechtler K, Roitinger E, Franklin FCH. 2015. Arabidopsis PCH2 Mediates Meiotic Chromosome
Remodeling and Maturation of Crossovers. *PLoS Genet* **11**:e1005372-.

Lloyd A. 2022. Crossover patterning in plants. *Plant Reprod*. doi:10.1007/s00497-022-00445-4

Mercier R, Jolivet S, Vezon D, Huppe E, Chelysheva L, Giovanni M, Nogué F, Doutriaux M-P, Horlow C,
Grelon M, Mézard C. 2005. Two Meiotic Crossover Classes Cohabit in Arabidopsis: One Is
Dependent on MER3,whereas the Other One Is Not. *Current Biology* **15**:692–701.
doi:https://doi.org/10.1016/j.cub.2005.02.056

Miller DE. 2020. The Interchromosomal Effect: Different Meanings for Different Organisms. *Genetics*
216:621–631. doi:10.1534/genetics.120.303656

Morgan C, Fozard JA, Hartley M, Henderson IR, Bomblies K, Howard M. 2021. Diffusion-mediated
HEI10 coarsening can explain meiotic crossover positioning in Arabidopsis. *Nat Commun*
12:4674. doi:10.1038/s41467-021-24827-w

Morgan C, Wegel E. 2020. Cytological Characterization of Arabidopsis arenosa Polyploids by SIM. In:
Pradillo M. HS, editor. Plant Meiosis. Methods in Molecular Biology, Vol 2061. Humana, New
York, NY. doi:https://doi.org/10.1007/978-1-4939-9818-0_4

Otto SP, Payseur BA. 2019. Crossover Interference: Shedding Light on the Evolution of
Recombination. *Annu Rev Genet*. doi:10.1146/annurev-genet-040119-093957

Page SL, Hawley RS. 2004. THE GENETICS AND MOLECULAR BIOLOGY OF THE SYNAPTONEMAL
COMPLEX. *Annu Rev Cell Dev Biol* **20**:525–558. doi:10.1146/annurev.cellbio.19.111301.155141

Pochon G, Henry IM, Yang C, Lory N, Jiménez NF-, Böwer F, Hu B, Carstens L, Tsai HT, Pradillo M,
Comai L, Schnittger A. 2022. The Arabidopsis Hop1 homolog ASY1 mediates cross-over
assurance and interference. *bioRxiv* 2022.03.17.484635. doi:10.1101/2022.03.17.484635

Prusicki MA, Keizer EM, van Rosmalen RP, Komaki S, Seifert F, Müller K, Wijnker E, Fleck C, Schnittger
A. 2019. Live cell imaging of meiosis in Arabidopsis thaliana. *Elife* **8**:e42834.
doi:10.7554/eLife.42834

Qiao H, Prasada Rao HBD, Yang Y, Fong JH, Cloutier JM, Deacon DC, Nagel KE, Swartz RK, Strong E,
Holloway JK, Cohen PE, Schimenti J, Ward J, Hunter N. 2014. Antagonistic roles of ubiquitin
ligase HEI10 and SUMO ligase RNF212 regulate meiotic recombination. *Nat Genet* **46**:194–199.
doi:10.1038/ng.2858

Rackauckas C, Nie Q. 2017. DifferentialEquations.jl - A performant and Feature-Rich Ecosystem for
Solving Differential Equations in Julia. *J Open Res Softw* **5**:15.
doi:http://doi.org/10.5334/jors.151

Reynolds A, Qiao H, Yang Y, Chen JK, Jackson N, Biswas K, Holloway JK, Baudat F, de Massy B, Wang J,
Höög C, Cohen PE, Hunter N. 2013. RNF212 is a dosage-sensitive regulator of crossing-over
during mammalian meiosis. *Nat Genet* **45**:269–278. doi:10.1038/ng.2541

Rog O, Köhler S, Dernburg AF. 2017. The synaptonemal complex has liquid crystalline properties and
spatially regulates meiotic recombination factors. *Elife* **6**:e21455. doi:10.7554/eLife.21455

Stauffer WT, Zhang L, Dernburg A. 2019. Diffusion through a liquid crystalline compartment
regulates meiotic recombinationProc.SPIE. doi:10.1117/12.2513378

Sturtevant AH. 1915. The behavior of the chromosomes as studied through linkage. *Z Indukt
Abstamm Vererbungsl*. doi:10.1007/BF01792906

Termolino P, Falque M, Aiese Cigliano R, Cremona G, Paparo R, Ederveen A, Martin OC, Consiglio FM,
Conicella C. 2019. Recombination suppression in heterozygotes for a pericentric inversion
induces the interchromosomal effect on crossovers in Arabidopsis. *The Plant Journal* **100**:1163–
1175. doi:https://doi.org/10.1111/tpj.14505

Varas J, Graumann K, Osman K, Pradillo M, Evans DE, Santos JL, Armstrong SJ. 2015. Absence of
SUN1 and SUN2 proteins in Arabidopsis thaliana leads to a delay in meiotic progression and
defects in synapsis and recombination. *The Plant Journal* **81**:329–346.
doi:https://doi.org/10.1111/tpj.12730

von Diezmann L, Rog O. 2021. Let's get physical – mechanisms of crossover interference. *J Cell Sci*
134. doi:10.1242/jcs.255745

Wang S, Veller C, Sun F, Ruiz-Herrera A, Shang Y, Liu H, Zickler D, Chen Z, Kleckner N, Zhang L. 2019.
Per-Nucleus Crossover Covariation and Implications for Evolution. *Cell* **177**:326-338.e16.
doi:https://doi.org/10.1016/j.cell.2019.02.021

Wayne C, Chloe G, Andrew L. 2021. Crossover interference: Just ZYP it. *Proceedings of the National
Academy of Sciences* **118**:e2103433118. doi:10.1073/pnas.2103433118

Yang C, Hu B, Portheine SM, Chuenban P, Schnittger A. 2020. State changes of the HORMA protein
ASY1 are mediated by an interplay between its closure motif and PCH2. *Nucleic Acids Res*
48:11521–11535. doi:10.1093/nar/gkaa527

Yang C, Sofroni K, Hamamura Y, Hu B, Elbasi HT, Balboni M, Chu L, Stang D, Heese M, Schnittger A.
2022. ZYP1-mediated recruitment of PCH2 to the synaptonemal complex remodels the
chromosome axis leading to crossover restriction. *Nucleic Acids Res* **50**:12924–12937.
doi:10.1093/nar/gkac1160

Zhang L, Köhler S, Rillo-Bohn R, Dernburg AF. 2018. A compartmentalized signaling network
mediates crossover control in meiosis. *Elife* **7**:e30789. doi:10.7554/eLife.30789

Zhang L, Stauffer W, Zwicker D, Dernburg AF. 2021. Crossover patterning through kinase-regulated
condensation and coarsening of recombination nodules. *bioRxiv* 2021.08.26.457865.
doi:10.1101/2021.08.26.457865

Zickler D, Kleckner N. 2016. A few of our favorite things: Pairing, the bouquet, crossover interference
and evolution of meiosis. *Semin Cell Dev Biol*. doi:10.1016/j.semcdb.2016.02.024

Ziolkowski PA, Underwood CJ, Lambing C, Martinez-Garcia M, Lawrence EJ, Ziolkowska L, Griffin C,
Choi K, Franklin FCH, Martienssen RA, Henderson IR. 2017. Natural variation and dosage of the
HEI10 meiotic E3 ligase control Arabidopsis crossover recombination. *Genes Dev*.
doi:10.1101/gad.295501.116

881
882

Figure 1. Mathematical model for HEI10 dynamics in a _zyp1_ SC mutant. (**A**) Each cell contains Q=5 chromosome pairs (purple line pairs). RIs (yellow stars) are placed randomly along the chromosome pairs immersed in the nucleoplasm (P). (**B**) HEI10 (red) is able to move from the nucleoplasmic pool into the RI compartments (rate α), and escape (rate β) from the RI compartments back into the pool. $C_{q,n}$ is the HEI10 molecule number at the n-th RI on the q-th chromosome pair, which has length L_q. (**C**) Differential equations governing nucleoplasmic pool (P) and RI HEI10 (C) amounts, together with functional form of escape rate function $\beta(C)$, smoothing function $H(s)$ and initial conditions for RI and nucleoplasmic HEI10 amounts. $\mathcal{N}_{\mathcal{T}}$ is a Normal distribution, truncated at three standard deviations away from its mean. (**D**) Graph showing RI escape rate function $\beta(C)$. (**E**) Default simulation parameter values. (**F**) Form of end-bias function f. (**G**) Description of terms appearing in model equations.

Figure 2. Analysis of CO number in *zyp1* mutant. (A-B) Experimental data from *zyp1* null mutant plants (top) and nucleoplasmic coarsening model simulations (bottom). Results from simulating the model for 10000 cells are shown. (**A**) Distribution of total-CO number per cell. Experimental data is from pooled MLH1 foci counts from (Capilla- Pérez *et al.,* 2021). Sample mean (μ), estimated variance (S^2), and sample size (N) inset. Green dots (joined by a line) show a Poisson distribution with the same mean. (**B**) Distribution of univalent number per cell. Experimental data is from univalent counts in metaphase I chromosomes from (Capilla- Pérez *et al.,* 2021). Again data from 10000 simulated cells (**C**) Simulation outputs showing the distribution of total-CO numbers on the five individual chromosomes pairs. Green dots and lines again show Poisson distributions with same means as simulated data, with sample mean (μ) and estimated variance (S^2) inset. Simulation output from 10000 cells. (**D**) Traces of HEI10 focus intensity against time, for all RIs within a single simulated cell. Coloured lines indicate HEI10 amounts at each RI, in arbitrary units (a.u.). All simulation parameters are listed in Figure 1E.

Figure 3. 3D-SIM imaging of late-prophase I cells. Maximum intensity projections of 3D-image stacks
from wild-type (**A**) and *zyp1a-2/zyp1b-1* mutants (**B**), labelled for SMC3 (green) and HEI10 (red). 3D-
models of segmented axial elements (with each chromosome axis labelled in a different colour),
generated using the SNT plugin to FIJI, are also shown. Each nucleus contains 5 pairs of axes (bottom
row, A, B), with individual axes in each pair having equivalent lengths and pairing tightly (A) or
roughly (B) in 3D space. Zoomed in regions (right hand panels) from merged images (yellow dashed
boxes) show the localisation of late-HEI10 foci between closely-paired axes. Scale bars = 5 μm.

Figure 4. Analysis of late-HEI10 focus patterning and intensity data in wild-type and *zyp1* mutant.
(A-C) Experimental data from wild-type plants (left), *zyp1a-2/zyp1b-1* plants (middle), and from
nucleoplasmic coarsening model simulation outputs (right). Model outputs from simulating 10000
cells are shown. (**A**) Late-HEI10 focus positions along chromosome pairs, relative to the length of the
chromosome pair. Experimental data are replicated and made symmetric about chromosome
midpoints. (N=161322 simulated foci). (**B**) Distribution of spacing between adjacent late-HEI10 foci,
in µm. (N=113278 simulated spaces between foci). (**C**) Violin plots of late-HEI10 foci intensities
against the number of late-HEI10 foci on that chromosome pair. Focus intensities are relative to the
mean intensity of all (on-chromosome) HEI10 foci within the same cell. Red lines show simple linear
regression best fits, treating the number of foci on the chromosome as a continuous independent
variable. For wild-type plants, slope = -0.1944, R^2=0.188, F(1,88)=20.36, p<0.001. For *zyp1* mutant
plants, slope=0.0028, R^2<0.001, F(1,260)=0.04538, p=0.831. For simulated data, slope=0.0004,
R^2<0.001, F(1,161320)=0.7838, p=0.376. Simulation parameters are again as listed in Figure 1E, with
simulated data from 10000 cells (50000 chromosome pairs).

Figure 5. Late-HEI10 focus patterning and intensity data in *pch2-1*. (**A**) Maximum intensity
projections of 3D-image stacks from *pch2-1* mutants labelled for HEI10, ZYP1 and SMC3. A 3D-model
of segmented SC segments (with each segment labelled in a different colour), generated using the
SNT plugin to FIJI, is also shown. (**B-G**) Experimental data and combined coarsening model
simulations showing; (**B**) late-HEI10 focus number per cell (1000 simulated cells), green dots (joined
by a line) show a Poisson distribution with the same mean, (**C**) late-HEI10 focus number per SC
segment (13870 simulated segments), (**B-C**) sample mean (μ) and estimated variance (S^2) inset, (**D**)
positioning of all late-HEI10 foci along SC segments relative to total segment length (11009
simulated foci), (**E**) spacing between neighbouring late-HEI10 foci on the same SC segment (2590
spaces between simulated foci), (**F**) SC segment length versus number of late-HEI10 foci per SC
segment (same number of observations as in C) and (**G**) single late-HEI10 focus intensity (relative to

the mean intensity of all HEI10 foci within the same cell) versus SC segment length, with a random
sample of simulation output to match experimental dataset size.

Figure 1 – figure supplement 1. Form of end-bias function f from nucleoplasmic coarsening model

(Figure 1F) and from the SC-mediated coarsening model (Morgan et al., 2021). Note that the extra

HEI10 loading in the nucleoplasmic coarsening model (up to 25% on the most distal 60% of each

chromosome pair) is lower and more spread than that used in the SC-mediated coarsening model

(up to 100% on the most distal 20%). Here x/L is relative position along chromosome pair, with x_e =

0.3 (left) and x_e = 0.1 (right), specifying starting position of end-bias, where f_e = 1.25 (left) and f_e = 2

(right).

Figure 2 – figure supplement 1. *zyp1* total-CO number per cell simulation without nonuniform
initial loading. *zyp1* nucleoplasmic coarsening model simulation showing the distribution of total-CO
number per cell, where the model has uniform initial loading (no end-bias). Results from simulating
the model for 10000 cells are shown. Sample mean (μ) and estimated variance (S^2) inset. Green dots
(joined by a line) show a Poisson distribution with the same mean.

Figure 3 – figure supplement 1. 3D-SIM imaging of early pachytene cells. Maximum intensity projections of 3D-image stacks from wild-type (**A**) and *zyp1a-2/zyp1b-1* mutants (**B**), labelled for HEI10 (red), SMC3 (green) and DAPI (blue). Scale bars = 5 µm.

Figure 4 – figure supplement 1. A combined SC- and nucleoplasm-mediated coarsening model. (**A**) Schematic representation of the combined coarsening model. (**B**) HEI10 (red) is able to move from the nucleoplasmic pool into the RI compartments (rate α), and escape (rate β) from the RI compartments back into the nucleoplasm. HEI10 can also move from SC compartments into the RI compartments (rate $\hat{\alpha}$) and escape back onto the SC (rate $\hat{\beta}$). HEI10 on the SC has concentration

$c_q(x, t)$, and is able to diffuse along the SC with one-dimensional diffusion coefficient D. $C_{q,n}$ is the

HEI10 molecule number at the *n*-th RI on the *q*-th chromosome pair, which has length L_q. (**C**)

Differential equations and initial/boundary conditions governing simulations. (**D**) Default simulation

parameter values for various scenarios: WT, original SC-mediated coarsening model with wild-type

parameters (as implemented in Methods), WT+nuc, combined SC- and nucleoplasm-mediated

coarsening model with wild-type parameters. (**E**) Description of terms appearing in model equations.

Figure 4 – figure supplement 2. The combined SC- and nucleoplasm-mediated coarsening model can explain crossover patterning in wild-type and HEI10 over-expressor lines. Columns show simulated results from the original SC-mediated coarsening model with wild-type parameters (WT, as implemented in Methods), the combined SC- and nucleoplasm-mediated coarsening model with wild-type parameters (WT+nuc), the original SC-mediated coarsening model with HEI10 over-expressor parameters (OX, as implemented in Methods), and the combined SC- and nucleoplasm-mediated coarsening model with HEI10 over-expressor parameters (OX+nuc). Results from simulating the model for 1000 cells are shown. Above simulations of WT and OX cases agree well with previously published experimental data (Morgan et al., 2021). Number of late-HEI10 foci per cell, green dots (joined by a line) show a Poisson distribution with the same mean, **(A),** number of late-HEI10 foci per SC (5000 simulated SC), **(B)**, sample mean (μ) and estimated variance (S^2) inset

(**A-B**), spacing between successive late-HEI10 foci **(C)**, relative positions of late-HEI10 foci along SC
(D), relative positions of single late-HEI10 foci (WT, WT+nuc) and double late-HEI10 foci (OX,
OX+nuc) **(E)**, and relative positions of double late-HEI10 foci (WT, WT+nuc) and triple late-HEI10 foci
(OX, OX+nuc) **(F)**. Parameters are as listed in Figure 4 – figure supplement 1D, with parameters μ_F
and σ_F controlling initial cellular HEI10 levels increased to 4.5x the WT values in the HEI10 over-
expressor simulations.

A

B

PATCH

$x = 0$

$c_q(x, t)$

$x = L_q$

$\hat{\beta}(C_{q,n})$ $\hat{\alpha}$ $\hat{\beta}(C_{q,n+1})$ $\hat{\alpha}$

$C_{q,n}$ $C_{q,n+1}$

α $\beta(C_{q,n})$ α $\beta(C_{q,n+1})$

P P

C

	Value	Unit	Description
α	0.014	s^{-1}	RI HEI10 absorption rate from nucleoplasm
β_C	0.003	s^{-1}	RI HEI10 escape rate into nucleoplasm
$\hat{\alpha}$	0.42	$\mu m\, s^{-1}$	RI HEI10 absorption rate from SC
$\hat{\beta}_C$	0.075	s^{-1}	RI HEI10 escape rate onto SC
f_e	1.0		Telomere weighting for RI loading
μ_Q	13.92		Mean number of SC patches
L_α	1.693		Alpha parameter of patch length gamma distribution
L_θ	2.984	μm	Theta parameter of patch length gamma distribution

1001

1002 **Figure 5 – figure supplement 1. Model for *pch2* simulations. (A-B)** Schematic of combined
1003 nucleoplasmic- and SC-mediated coarsening model used for *pch2* simulations, where dynamics occur
1004 on patches of SC, rather than whole chromosomes. (**C**) Parameters for *pch2* simulations that are
1005 different from those in Figure 4 – figure supplement 1D. See Methods for more details.