1

Beyond bold versus shy: Zebrafish exploratory behavior falls into several behavioral

clusters and is influenced by strain and sex

Neha Rajput, Kush Parikh, and Justin W. Kenney

Department of Biological Sciences
Wayne State University
Detroit, MI 48202

Abstract

Consistent individual differences in exploratory behavior have been found across a range of

taxa, including zebrafish, and are thought to contribute to evolutionary fitness. Animals that

explore more of a novel environment, and visit areas of high predation risk, are considered bold,

whereas animals with the opposite pattern of behavior are considered shy. Here, we examined

whether this bimodal characterization of bold versus shy adequately captures the breadth of

exploratory behavior exhibited by zebrafish or if their behavior falls into several distinct

behavioral subtypes instead. To identify the presence of behavioral subtypes, we used

unsupervised machine learning applied to exploratory behaviors extracted from three-

dimensional swim traces from over 400 adult zebrafish across four strains (AB, TL, TU, and

WIK) and both sexes and found that their exploratory behavior stratified into four distinct

clusters. These included previously described bold and shy behavior as well as two new

behavioral types: wall-huggers and active explorers. Consistent with prior descriptions of

individual differences, these behavioral subtypes were stable across days and influenced by the

strain and sex of the animals. Taken together, our work suggests that zebrafish behavior

exhibits greater complexity than is typically assumed.

Introduction

Interest in the biology of behavioral differences can be traced back to at least Roman

antiquity where Galen applied the humoral theory to explain variation in human temperament

(as translated in Grant, 2000). Today, we have a greater understanding of human personalities,

defined as behavioral tendencies that are consistent across time and context, but its biological

basis remains elusive. One avenue for progress is the modeling of human personality through

the study of individual difference in animal behavior. Often dismissed as noise around an

average, a growing body of work has found that variations in animal behavior are often

consistent across time and context (Dall et al., 2012; Sih et al., 2004). Such consistent

differences have been described for behaviors important for evolutionary fitness and in a wide

range of taxa, suggesting that they may provide grist for adaptation to an ever-changing

environment.

One of the most widely studied axes of behavioral variation in animals is bold versus

shy. Bold animals tend to explore or investigate novel environments or objects more readily than

shy animals that tend to flee or retreat in response to novelty (Réale et al., 2007; Toms et al.,

2010; Wilson et al., 1994). From a fitness perspective, boldness may be adaptive when food

resources are scarce and predation risk is low, whereas shyness may be more effective when

the opposite conditions prevail (Réale et al., 2007). Variation along this axis has been described

in animals ranging from bears (Myers and Young, 2018), lizards (López et al., 2005), birds

(Carere et al., 2005), and fish (Toms et al., 2010). Studies examining the bold-shy axis typically

begin with the assumption that animals fall into one of these two categories. However, whether

this bimodal distribution of bold versus shy fully captures animal behavior has not been tested

directly. It is likely that there is more complexity to animal behavioral types which may have

been overlooked due to use of small sample sizes or assessment of only one or two behaviors

(Forkosh et al., 2019).

Zebrafish have proven to be an excellent model organism to understand behavior and its

biological basis. With 70% of fish genes having an obvious human ortholog (Howe et al., 2013),

and a central nervous system that has the same general organization and utilizes many of the

same neurotransmitters as mammals (Kenney et al., 2021; Panula et al., 2010; Wulliman et al.,

1996), findings using zebrafish are widely applicable. Our understanding of zebrafish behavior

has expanded rapidly over the past decade (Gerlai, 2020; Kalueff et al., 2013; Kenney, 2020),

including several studies that have examined the bold-shy axis. Boldness is often probed by

exposing fish to a novel tank and examining locomotion, exploratory activity and avoidance

behaviors, like geotaxis (i.e. bottom dwelling) or thigmotaxis (i.e. proximity to tank walls)

(Mustafa et al., 2019; Oswald et al., 2012; Thörnqvist et al., 2019; Toms et al., 2010). Animals

that are more active or spend more time in parts of the tank that would increase risk of predation

(i.e. the top and/or center of the tank) are considered bolder. In zebrafish, these behaviors have

been found to be consistent over time (Baker et al., 2018; Tran and Gerlai, 2013) and can

predict other behaviors like social dominance (Dahlbom et al., 2011), aggression (Martins and

Bhat, 2014), and stress reactivity (Oswald et al., 2012), all hallmarks of personality. However,

fully utilizing zebrafish exploratory behavior to understand the biological basis of individual

differences requires that we first determine if the bold versus shy distinction adequately

captures the breadth of behavioral variability exhibited during exploration.

To determine the presence of multiple behavioral clusters during exploration of a novel

tank we captured three-dimensional swim behavior from over four hundred fish. Because

zebrafish behavior is known to be influenced by strain and sex (Volgin et al., 2019), we used

animals from four inbred strains (AB, TU, WIK, and TL) and both sexes to ensure that we

captured the full range of behavioral variability. We then used an unsupervised machine

learning approach where we found that exploratory behavior stratified into four clusters that

included the traditional bold versus shy distinction, as well as two additional behavioral types.

We also found that individual cluster membership remained consistent across days, and that the

proportion of fish in each cluster was influenced by strain and sex.

Results

Three-dimensional behavioral tracking

To capture three-dimensional zebrafish swim behavior during exploration of a novel

tank, we used Intel RealSense™ cameras mounted above five-sided tanks with frosted walls

(Figure 1A). These cameras simultaneously captured color and depth streams, resulting in

three-dimensional videos (Figure 1B). Fish posture at each frame was tracked in the color

stream using DeepLabCut (Figure 1C; Mathis et al., 2018). These points were overlayed onto

the depth stream to create three-dimensional swim traces (Figure 1D) from which we extracted

positional information (distance from bottom and center), distance travelled, and percent tank

explored (Figure 1E).

Influence of sex and strain on exploratory behaviors

We first determined if sex and genetics influences individual zebrafish exploratory

behaviors by assessing swim traces of fish from four strains (AB, TU, TL, and WIK), and both

sexes, on two consecutive days. We extracted four exploratory behaviors from each swim trace:

distance from bottom, distance from center, total distance travelled, and percentage of tank

explored (Figure 1E). We found that the distribution of several of the parameters deviated from

normality (Figure S1), so we performed non-parametric 4 × 2 (strain × sex) permutation

ANOVAs. For distance from bottom (Figure 2A), we found a main effect of strain ($P = 0.0001$),

but no effect of sex ($P = 0.40$) nor an interaction ($P = 0.44$). FDR (false discovery rate) corrected

permutation t-tests found that TL fish swam closer to the bottom of the tank than all other strains

(P's ≤ 0.0008). For distance from center (Figure 2B) there was a main effect of sex ($P = 0.042$)

and a trend towards an interaction ($P = 0.060$) where female fish of every strain, except AB,

spent more time closer to the center of the tank. We also found a main effect of strain ($P =$

0.0001) with FDR corrected permutation t-tests indicating that TL fish swam closer to the center

of the tank than all other strains (P's = 0.0004), and AB spent more time on the periphery (AB

compared to TU: P = 0.0006, WIK: P = 0.017). For distance travelled (Figure 2C), we found a

main effect of sex (P = 0.0001) where male fish swam further than female fish. There was also a

main effect of strain (P = 0.014), but no interaction (P = 0.59). FDR corrected permutation t-tests

found that TL fish swam less than AB fish (P = 0.043) and a trend towards a difference between

TLs and WIKs (P = 0.061). Finally, for percent of the tank explored (Figure 2D), we also found a

main effect of sex (P = 0.024) in which female fish explored less of the tank than their male

counterparts. There was also a main effect of strain (P = 0.0001), but no interaction (P = 0.34).

Post-hoc tests revealed that TL fish explored the tank less than all other strains (P's = 0.0004),

and that AB fish explored less than TU (P = 0.035) and WIK (P = 0.0026) fish. Taken together,

we find that there are several sex differences (center distance, distance travelled, and percent

explored) and that TL fish differ the most from other strains across all measures with no clear

strain by sex interactions.

Influence of sex and strain on habituation to the novel tank

Some exploratory behaviors have been found to habituate over a single six-minute

exposure to a novel tank (Wong et al., 2010), so we examined whether any of the parameters

we measured changed over time and were influenced by sex in the various strains (Figure S2).

For each measure, we used non-parametric 2 × 6 (sex × time interval) mixed permutation

ANOVAs and adjusted for multiple tests using an FDR correction. For distance from bottom

(Figure S2A), we found no main effects of sex (AB: P = 0.51 , TL: P = 0.46 , TU: P = 0.74, WIK:

P = 0.86), but found effects of interval in all strains (AB: P = 0.0006, TL: P = 0.0093, TU: P =

0.0006, WIK: P = 0.0084) where all fish, except WIKs, increased their distance from the bottom

over time. There were no interactions except for WIKs (AB: P = 0.086, TU: P = 0.74, TL: P =

0.46, WIK: P = 0.018), where female fish appeared to spend the first minute further from the

tank bottom than their male counterparts. For distance from center (Figure S2B), a main effect

of interval was found for AB, TU, and WIKs (P's = 0.0004, TL: P = 0.52), finding that these

strains increased their distance from center over time. There were no interactions between sex

and time interval (AB: P = 0.91, TL: P = 0.28, TU: P = 0.91, WIK: P = 0.91), but we found trends

towards an effect of sex in TU and WIKs (AB: P = 0.28, TL: P = 0.41, TU: P = 0.054, WIK: P =

0.064) where female fish spent their time closer to the center of the tank than male fish,

consistent with what was observed in the overall data (Figure 2B). Finally, for distance travelled

(Figure S2C), there was an increase in locomotor activity over time in AB, TU, and WIK fish,

with a trend in TLs (AB: P = 0.0004, TL: P = 0.093, TU: P = 0.0012, WIK: P = 0.0004).

Consistent with the overall data, there were also main effects of sex in all strains except TLs,

where there was a trend (AB: P = 0.0014, TL: P = 0.093, TU: P = 0.0004, WIK: P = 0.0048).

Only the TU fish had an interaction between time interval and sex where TU female fish

increased their distance travelled over time, but male fish did not (AB: P = 0.085, TL: P = 0.39,

TU: P = 0.0022, WIK: P = 0.068).

Correlations between behavioral parameters

To determine the extent to which exploratory behavioral parameters are associated with

each other, and if these associations are affected by strain and sex, we computed correlations

between measures (Figures 2 and S3). We used Spearman's ρ to identify monotonic

relationships because of the presence of several non-normally disturbed parameters (Figure

S1). Not surprisingly, we found that distance travelled was positively correlated with percent

explored across most strains and sexes (Figures 2J & S3F). We also found positive correlations

between bottom distance and percent explored (Figures 2G & S3C), consistent with the idea

that a higher bottom distance is associated with an increased willingness to explore. However,

bottom distance did not correlate consistently with distance travelled (Figures 2F & S3B),

suggesting that, despite positive correlations between distance travelled and percent explored,

they are capturing different aspects of exploration. Bottom distance and center distance were

also not consistently correlated with each other (Figures 2E & S3A); in three strains there was a

weak negative correlation (female TLs and TUs, and male WIKs) where fish that swam nearer

to the top also swam closer to the center, and in one strain (male ABs) the opposite relationship

was observed. Center distance was mostly negatively correlated with percent explored (Figures

2I & S3E), but not universally so (male TLs being the exception), consistent with the idea that

fish that spend more time closer to the center of the tank also explore more of the tank.

Identifying behavioral clusters

Given the presence of non-normal behavioral distributions (Figure S1) and the influence

of sex and strain on the relationship between different exploratory parameters (Figure 2E-J), we

hypothesized the presence of multiple behavioral clusters in our data set. To test this, we built a

k-nearest neighbor network and applied the Louvain community detection algorithm to identify

clusters (Blondel et al., 2008). We used all four behavioral parameters (bottom distance, center

distance, distance travelled, and percent explored) in calculating nearest neighbor distances

because none of the behavioral measures showed consistently high correlations across all

strains and sexes (Figure 2E-J). To determine 'k' for building the network, we explored a range

of values and chose a value (k = 118) that optimized internal clustering metrics and was robust

to small deviations in k (Figure S4). We found that our behavioral data stratified into four distinct

behavioral clusters (Figure 3A).

An examination of the behaviors associated with each of the clusters reveals a range of

behavioral profiles that run the gamut from shy to bold (Figure 3B). Fish in the shyest cluster

were lowest in both bottom distance and percent exploration (Figure 3B; cluster 1) whereas fish

in the boldest cluster spent the most time near the top and center of the tank while exploring

more of the tank than average (Figure 3B; cluster 4). Surprisingly, however, these bold fish were

the lowest in distance travelled, suggesting that they were particularly efficient in their tank

exploration. Two 'mixed' clusters were also identified: cluster 2 where fish were near average on

all measures except for center distance where they spent most of their time near the periphery,

a group we call 'wall-huggers' (Figure 3B), and cluster 3 where fish were above average in

distance travelled and percent explored that we call 'active explorers' (Figure 3B).

Behavioral clusters across sex and strain

To determine if fish from a given strain and sex were over- or underrepresented in

behavioral clusters, we computed p-values using permutation resampling and FDR correction

for multiple comparisons (Figure 3C). We found that for the shy cluster (cluster 1), TL fish,

irrespective of sex, were overrepresented (female: $P = 0.0011$, male: $P = 0.048$), whereas the

wall-huggers (cluster 2) had overrepresentation of female AB fish ($P = 0.0053$). Consistent with

our finding that males, on average, swam more than females, more male fish were in the active

explorers group (cluster 3), although overrepresentation was only significant in TU and WIK fish

(males: AB: $P = 0.086$, TL: $P = 0.23$, TU: $P = 0.0016$, WIK: $P = 0.0073$). Finally, in the boldest

cluster (cluster 4), females outnumbered males in all strains but there was only significant

overrepresentation in TU fish and trends in ABs and WIKs (female: AB: $P = 0.082$, TL: $P = 0.16$,

TU: $P = 0.018$, WIK: $P = 0.064$).

Habituation over two days

All 426 fish used for generating clusters were exposed to the novel tank on two

consecutive days, allowing us to determine if their behavior remained consistent across

repeated exposures. First, we analyzed individual exploratory behaviors with permutation paired

t-tests with FDR corrections (Figure S5). We found that AB and TL fish increased their bottom

distance during the second exposure, with a trend in female WIK fish (Figure S5A; female AB: P

= 0.0080, male AB: P = 0.012, female TL: P = 0.0016, male TL: P = 0.0040, female TU: P =

0.54, male TU: P = 0.54, female WIK: P = 0.055, male WIK: P = 0.62). Unexpectedly,

thigmotaxis (center distance) increased in several groups: AB fish and female WIKs with trends

towards an increase in TU females, but a decrease in TL males (Figure S5B; female AB: P =

0.0032, male AB: P = 0.043, female TL: P = 0.38, male TL: P = 0.073, female TU: P = 0.073,

male TU: P = 0.82, female WIK: P = 0.043, male WIK: P = 0.87). Distance travelled did not

change in any fish (Figure S5C; P's ≥ 0.12), and percent explored had trends towards a

decrease in AB males and increase in TL females (Figure S5D; female AB: P = 0.13, male AB:

P = 0.071, female TL: P = 0.071, male TL: P = 0.55, female TU: P = 0.82, male TU: P = 0.39,

female WIK: P = 0.55, male WIK: P = 0.31). Taken together, the changes in behavior during the

second exposure were mixed, with some changes indicating an increase in behaviors

associated with boldness, like bottom distance, and others an increase in putative shy

behaviors, like thigmotaxis.

Behavioral cluster consistency over two days

To determine if the behavioral clusters we identified remained consistent across days,

we used exploratory data from the second day of exposure to the novel tank to assign fish to

clusters uncovered from the first day (Figure 4A). We found that 52.8% of fish fell into the same

behavioral cluster on days 1 and 2 (P = 0.0001; permutation average = 25.1% overlap). Overall,

the number of fish falling into the shyest group (cluster 1) decreased, and the wall-huggers

(cluster 2) remaining the most consistent. At the level of sex and strain (Figure 4B), we again

used FDR corrected permutation resampling, finding that, as on day 1, AB females were

overrepresented in cluster 2 (P = 0.0011), but were now joined by their male counterparts (P =

0.031). TL females also remained overrepresented in the shyest cluster (P = 0.0011), but a

significant portion of male and female TL fish were now in the boldest cluster (female TL: P =

0.013, male TL: P = 0.0011). TU females were no longer overrepresented in the boldest group

(P = 0.62), although TU males were now underrepresented in this group (P = 0.025) and

maintained their overrepresentation as active explorers, cluster 3 (P = 0.0011). Finally, neither

female nor male WIKs were overrepresented in any cluster on day 2.

Habituation over five days

Some exploratory behaviors have been found to habituate over several days of

exposure to a novel tank (Wong et al., 2010). In a new cohort of AB and TU fish, exploratory

behavior was measured over five consecutive days of exposure to the novel tank (Figure S6).

We used 5 × 2 (day × sex) mixed permutation ANOVAs to assess significance. For bottom

distance in AB fish (Figure S6A), there was no effect of day (P = 0.20), sex (P = 0.14) or an

interaction (P = 0.64), whereas TU fish increased their bottom distance over the five days (P =

0.025), with no effect of sex (P = 0.18) nor an interaction (P = 0.87). AB fish increased their

center distance (Figure S6B) over time (P = 0.0001) with no effect of sex (P = 0.12) or an

interaction (P = 0.14), and TU fish had a similar trend (day: P = 0.094, sex: P = 0.27, interaction:

P = 0.93). As we saw before, male AB fish swam further than female fish (P = 0.010) with no

effect of day (P = 0.62). There was a trend towards and interaction between day and sex (P =

0.058) where AB males appeared to swim less over time, and females more (Figure S6C). In

TU fish, there was a trend towards males swimming more than females (P = 0.084), and a clear

effect of day (P = 0.0007) and no interaction (P = 0.68) as both sexes increased their distance

travelled over time. Finally, for percent explored (Figure S6D), In AB fish there was no effect of

sex (P = 0.33) or day (P = 0.77), but there was a trend towards an interaction (P 0.068) where

female fish appeared to slightly decrease their percent explored over time. In TU fish, animals

increased their exploration over days (P = 0.0089), with a trend towards males exploring more

than females (P = 0.099), and no interaction of day and sex (P = 0.31).

Behavioral cluster consistency over five days

Next, we asked whether the behavioral clusters of individual animals over the five

consecutive days remained consistent (Figure 5). We found that, for both AB and TU fish, the

exploratory behavior of approximately 50% of animals fell into the same cluster on at least 4 out

of 5 days (Figure 5A). To determine if the consistency across days was greater than chance,

each animal was assigned an overlap score, a sum of the number of pair-wise overlaps across

the five days. This score ranged from 1 (only one pair of days overlapped) to 10 (all pair-wise

overlaps). The average scores for AB and TU fish were 5.6 ± 2.6 and 5.9 ± 3.1 (mean ±

standard deviation), respectively, both of which were significantly higher than the overlap scores

from resampling (permutation mean ± standard deviation: 3.4 ± 0.2 and 2.7 ± 0.2, respectively;

P's = 0.0002). Although most animals had scores higher than the permutation average, a small

subset of both AB and TU animals were "consistently inconsistent" with their exploratory

behavior falling into all four clusters across the five days. We also found that the number of

animals falling into the shyest cluster (cluster 1) decreased over time with little change in most

other clusters except an increase in the number of AB fish that became wall-huggers (cluster 2;

Figure S7).

Finally, we examined how much cluster overlap there was between each of the five days

of the experiment (Figure 5B). In both AB and TU fish, we found that the clusters from day 1

overlapped above chance with most clusters on subsequent days, with the overlap in TU fish

averaging about 10% higher than AB fish. In AB fish, cluster overlap after day 2 increased

markedly (range 62-70%). In TU fish, an increase in overlapping clusters did not occur until day

3, where it then remained high (63-80%).

Discussion

By applying an unbiased approach to the clustering of three-dimensional exploratory behavior from over 400 zebrafish, we found that zebrafish exploratory behavior stratifies into four distinct behavioral clusters. These profiles included previously described bold and shy behaviors as well as two novel behavioral types: wall-huggers and active explorers. Notably, these individual differences in fish behavior were consistent over time, one of the key hallmarks of personality. Although there were few strain-sex interactions on individual behaviors, the distribution of clusters varied considerably across strain and sex suggesting biological modulation of these behavioral clusters.

Studies that have examined individual differences in exploratory behavior typically assume a bimodal distribution of bold versus shy. Here, enabled by our large data set, we tested this assumption by taking an unbiased approach to identifying behavioral clusters, finding the presence of four distinct behavioral subtypes (shy, wall-huggers, active explorers, and bold). Cluster 4 was the boldest cluster where fish behavior corresponded to traditional descriptions of boldness: above average time in the top and center of the tank and greater exploration. The behavior of fish in cluster 1 was designated as shy because these fish spent most of their time near the bottom of the tank. Interestingly, in this group, behaviors like center distance and percent explored exhibited a high degree of variability, suggesting the potential presence of additional subgroups that may require an even larger data set to uncover.

Of the novel behavioral clusters we uncovered, we designated cluster 2 as 'wall-huggers' because these fish spent most of their time towards the periphery of the tank and were near average on all other parameters. We do not refer to this group as 'shy' because, although thigmotaxis has been interpreted as a predator avoidance or anxiety-like behavior in zebrafish (Kalueff et al., 2013), our findings do not clearly support this interpretation. For example, we did not observe consistent negative correlations between center and bottom distance (Figure 2E),

as would be expected if these behaviors reflected the same underlying construct. We also find

that, unlike bottom dwelling, thigmotaxis increased over time within and between sessions of the

novel tank test (Figures S2B and S6B), the opposite of the habituation we would expect to see if

thigmotaxis was an anxiety-like or predator avoidance behavior. Our findings are consistent with

other studies that examined thigmotaxis over time and/or alongside other avoidance behaviors

in adult fish (Blaser et al., 2010; Champagne et al., 2010; Rosa et al., 2018; Shams et al., 2015;

Wong et al., 2012). It may be that the interpretation of thigmotaxis in zebrafish has been unduly

influenced by findings in rodents where its importance for predator avoidance is clearer

(Champagne et al., 2010; Simon et al., 1994). Thus, we propose that thigmotaxis in adult

zebrafish should not be interpreted in the context of predator avoidance or anxiety until its

relevance for fish can be clarified.

We designated behavioral cluster 3 as 'active explorers' because these fish were clearly

above average in both distance travelled and percent explored, but near average on center and

bottom distance. This group of fish may correspond to what has been referred to as low

stationary, or proactive, zebrafish in other work (Baker et al., 2018; Wong et al., 2012). In

comparison to the boldest cluster, active explorers are similar in percent tank explored, but

distinguished by a notable elevation in distance travelled. Thus, during exploration, these

animals are less efficient in their exploration as they revisit many parts of the tank. This

suggests that these animals may have poor working memory or possibly higher resting

metabolic rates that have been found to be predictive of individual differences in activity (Biro

and Stamps, 2010; Yuan et al., 2018).

Strain and sex had a significant influence on behavioral cluster identity. During the initial

exposure to the tank, female fish more often fell into the boldest cluster, although only TU

females were statistically overrepresented. Male fish were more likely to be in the active

explorer group, particularly TU and WIK male fish. This likely reflects the fact that, on average,

male fish tended to swim further, and explore more of the tank, than female fish. This increase

in locomotor activity in male fish was expected as it has been reported elsewhere (Ariyomo and

Watt, 2015; Clayman et al., 2017; Philpott et al., 2012), although not universally so (Ampatzis

and Dermon, 2016; Fontana et al., 2019; Rambo et al., 2017; Tran and Gerlai, 2013). We also

found interactions between sex and strain on cluster identity, such as underrepresentation of AB

females, and overrepresentation of TL females, in the shyest cluster with the situation reversed

in the wall-huggers (cluster 2) group; notably, this interaction persisted into the second day of

testing, suggesting it is particularly robust.

The behavioral clusters we identified demonstrated a high degree of consistency across

days, supporting the idea that they are akin to personality types (Gosling, 2001). In our initial

experiment, where fish were tested in the novel tank on two consecutive days, overlap was

above 50%, twice as high as would be expected by chance (~25%). Upon repeated exposures

to the tank over five consecutive days, we found that cluster overlap steadily increased to above

65%, and over half of the fish were in the same cluster on 4 of the 5 days. This increase in

overlap over time is likely due, in part, to habituation (Wong et al., 2010) given that we found

that there were main effects of day on several individual measures, particularly in TU fish

(Figure S6). Nonetheless, this is the first report, to our knowledge, of consistency in zebrafish

behavior across days using unbiased multidimensional behavioral clustering that approximates

key elements of personality. However, it remains to be seen if these clusters can predict

behavior in other contexts.

The present work also yielded a comprehensive account of how genetic background

influences individual exploratory behaviors of zebrafish, adding to an already rich literature. The

primary strain difference on individual behaviors we observed was that TL fish differed from

most other strains, demonstrating higher bottom dwelling, less thigmotaxis, and exploring less of

the tank. Although strain differences in zebrafish behavior during the novel tank test have been

reported (Egan et al., 2009; Maximino et al., 2013; Mustafa et al., 2019), only a few studies

have compared widely used inbred strains. Vignet and colleagues (2013) found that AB fish

exhibited more bottom dwelling than TU fish, and Audira et al (2020) found no differences

between AB, TL, and WIK fish in bottom dwelling or distance travelled, but that WIK fish

exhibited greater thigmotaxis. Others have found AB fish to exhibit greater bottom dwelling than

WIKs (Sackerman et al., 2010), or a decrease in bottom dwelling in WIK fish over 60 minutes,

but not TU fish, with no difference in locomotor activity (Pannia et al., 2014). Differences

between these studies, and the present work, may be attributable to unreported sex ratios,

housing conditions, nuances of the behavioral task, or genetic drift such that inbred strains from

different labs or suppliers may differ subtly, as has been observed in rodents (Taft et al., 2006).

We attempted to address some of these challenges in our study. For example, housing

conditions are known to influence zebrafish behavior in the novel tank test (Parker et al., 2012;

Reolon et al., 2018). We housed all animals in mixed sex pairs for one week prior to behavioral

testing. This also allowed us to avoid any effects of stress due to social isolation while non-

invasively maintaining zebrafish identity over time. To minimize genetic drift, and maximize the

likelihood our findings will translate to other labs, all fish were within 2 generations of breeders

obtained from the Zebrafish International Resource Center where they maintain a much larger,

and genetically diverse, population of animals. Nonetheless, given that inbred zebrafish lines

are not isogenic (Nasiadka and Clark, 2012), there is no obvious way to ensure genetic

similarity of lines across labs.

Taken together, we have found that the exploratory behavior of zebrafish goes beyond

bold versus shy, stratifying into four distinct categories. This finding was enabled by recent

advances in animal tracking and 3D video capture that allowed us to scale up our behavioral

assessment using inexpensive open-source tools. As would be expected of behaviors capturing

personality, the clusters we identified are consistent over time and influenced by strain and sex.

Future work will be needed to determine if these clusters are predictive of behaviors in other

contexts. Nonetheless, our findings suggests that zebrafish behavior is more complex than is

typically assumed. Combined with recent advances in zebrafish molecular genetics (Rafferty

and Quinn, 2018) and tools for mapping the brain activity of adult zebrafish (Kenney et al.,

2021), our findings lay the groundwork for understanding the biological basis for individual

differences in vertebrate behavior.

452   **Acknowledgments**

453   We thank Barbara D. Fontana for excellent comments on a prior version of this manuscript. This

454   work was funded by the National Institutes of Health (R35GM142566) to JWK.

Methods

Subjects

Subjects were male and female AB, TU, WIK, or TL zebrafish 16-32 weeks of age. All

fish used in experiments were bred and raised at Wayne State University and within two

generations of animals obtained from the Zebrafish International Resource Center at the

University of Oregon. Animals were kept on high density racks under standard conditions

(temperature 26.5 ± 0.5 C; water conductivity 500 ± 10 μS, and a pH of 7.5 ± 0.2) with a 14:10

light/dark cycle (lights on at 8:00AM). Fish were fed twice a day with a dry feed in the morning

and brine shrimp (*Artemia salina*; Brine Shrimp Direct, Ogden, UT) in the afternoon. Behavioral

testing took place between 11:00 and 14:00. Sex of fish was confirmed via dissection after

behavioral procedures were complete. Those animals that were assigned the wrong sex were

removed from analysis. All procedures were approved by the Wayne State University

Institutional Animal Care and Use Committee.

Behavioral apparatus

Five-sided novel tanks (15 x 15 x 15 cm) were made from frosted acrylic

(ShopPopDisplays, Woodland Park, NJ) and open from above. Tanks were placed in an

enclosure made of white plasticore to diffuse light and prevent fish from seeing experimenters.

Intel RealSense™ cameras (D435) were mounted 20 cm above the tank to capture three-

dimensional videos (Kuroda, 2018). Video capture was done using custom written Python

scripts and the pyrealsense2 package.

Behavioral procedures

One week prior to behavioral testing, fish were placed as male/female pairs into 2 L

tanks. The tanks were divided in half with a transparent divider with two fish in each section and

a total of four fish in each tank. This allowed us to maintain the identity of fish over days without

reverting to isolation while also creating a consistent social environment across all animals. Fish

were taken off housing racks and moved to the procedural space at least one hour prior to

behavioral testing. Following testing, fish sat for one hour before being returned to the housing

racks. The novel tanks were filled with 2.5 L of fish facility water and individual fish were placed

in the tanks for six minutes while video was recorded for offline analysis. Water in the novel tank

was replaced between animals.

Animal Tracking

Fish were tracked in the color videos using DeepLabCut (Mathis et al., 2018). We

tracked five points (head, trunk, and three points on the tail; Figure 1C). Using ResNet 101, we

initially trained the network on 160 frames equally divided across fish of all four strains and both

sexes. We refined and improved our initial training by correcting outliers and including an

additional 160 frames.

To obtain the z-coordinate of fish at each frame, we overlaid the tracked points at each

color frame with the depth stream from the cameras. One z-coordinate for the fish was identified

for each frame based on a 4-pixel search radius around the tracked points starting with the

trunk. If no point was identified from the depth stream, it was interpolated. Z-coordinates were

corrected for the diffraction of water by measuring 100 points of varying distance from the

camera in the presence and absence of water and using the slope of a least-squares fit line.

Exploratory Behavioral Parameters

To calculate bottom distance, we calculated the equation for a plane along the bottom of

the tank using least squares fit of four points. We then calculated the shortest distance between

a point (the fish) and the plane. To measure center distance, we calculated how far fish were

from a line made from points at the center top and bottom of the tank. Percent of the tank

explored was calculated by dividing the tank into 1,000 evenly spaced voxels and calculating

the number of unique voxels visited. For distance travelled, traces were first smoothed using a

Savitzky-Golay filter (Press and Teukolsky, 1990) and then Euclidean distance was calculated

between points of successive frames.

Behavioral clusters

We identified behavioral clusters using a Louvain community detection algorithm

(Blondel et al., 2008) applied to a k-nearest neighbor network on our initial dataset of 426 fish

during their first day of exposure to the novel tank. This was done by first standardizing

individual behavioral parameters (bottom distance, center distance, distance travelled, and

percent explored), and calculating a similarity score between each fish:

$$Similarity\ score = \frac{1}{1 + D}$$

Where D is the Euclidean distance between each fish in four-dimensional behavioral space. To

determine the best k for building the network, we initially used a range of k's, applied the

Louvain community finding algorithm to weighted, non-directed networks, and calculated

internal clustering metrics (Calinski-Harabsz index (Caliński and Harabasz, 1974), Silhouette

index (Rousseeuw, 1987), and Davies-Bouldin index (Davies and Bouldin, 1979)). We chose

k=118, which was in the middle of a regime that optimized internal clustering and was robust to

small changes in k (Figure S4).

To identify clusters for the second day of novel tank exposure (Figure 4) and for fish

exposed on five consecutive days (Figure 5), we first standardized the new data using the

parameters from the 426 fish exposed to the novel tank described above. Then, for each new

data point, we assigned clusters based on the proportion of connections to its 51 nearest

neighbors in the initial network; 51 is half the size of the smallest cluster identified.

Coding and statistical analysis

Statistical analysis was performed using R (R Core Team and R Development Core

Team, 2016) and visualized using ggplot2 (Wickham, 2015) and RColorBrewer (Neuwirth and

Brewer, 2014). Normality was assessed using the Shapiro-Wilks test. Because a considerable

portion of the data was not normally distributed, we used permutation ANOVAs using the

permuco package (Frossard and Renaud, 2021), and the RVAideMemoire (Hervé and Hervé,

2020) for permutation t-tests. Multiple comparisons were corrected using a false discover rate

(FDR) correction (Benjamini and Hochberg, 1995). We used packages cccd (Marchette et al.,

2015) and igraph (Csardi and Nepusz, 2006) to build and analyze the k-nearest neighbor

network, and the ClusterCrit (Desgraupes, 2013) to assess internal clustering metrics. The

chord diagram was made using circlize (Gu et al., 2014).

For all permutation tests, we resampled data 10,000 times without replacement and

calculated the P-value as the proportion of experimental observations that were more extreme

than the permutation observations. When multiple tests were performed, P-values were

corrected using the FDR. To determine if behavioral clusters were affected by strain and sex

(Figure 3C and 4B), we resampled cluster assignment and calculated how many animals from

each strain and sex fell into each cluster. For comparison of behavior across days (Figure 4B)

we resampled cluster assignments from day 2. To determine if behavior was consistent across

five days (Figure 5A), first we calculated an overlap score for each fish that was the sum of the

number of days that had overlapping clusters (ranging from 1 to 10), then we took the average

of these overlap scores. We then resampled the cluster assignments of days 2 to 5 and

calculated the permutation overlap scores for each sample for comparison to the experimental

observations. Five-day percent overlap calculations and permutation tests were as described for

the two-day data except that resampling of cluster assignments was done individually for days 2

to 5.

Figures

Figure 1. Overview of three-dimensional behavioral tracking. **A**) Individual fish were placed into

a novel tank while video was recorded from above using Intel RealSense™ cameras. **B**) Videos

included both a color (top) and a depth (bottom) stream where fish can be seen (yellow arrows).

C) Animals were tracked in the color videos using DeepLabCut to identify five points along the

length of the fish. **D**) Tracking was overlaid with the depth stream to generate a three-

dimensional trace for each animal. **E**) Four exploratory parameters were extracted from each

trace: bottom distance (top left), center distance (top right), distance travelled (bottom left), and

percent of the tank explored (bottom right).

Figure 2. Influence of sex and strain on individual exploratory behaviors. The effect of sex and

strain on **A**) bottom distance, **B**) center distance, **C**) distance travelled, and **D**) percent of the

tank explored. Boxplots indicate median (center line), interquartile range (box ends), and 1.5

times the interquartile range (whiskers). Spearman's rank correlation coefficient (ρ) with 95%

confidence intervals across strain and sex for **E**) bottom distance versus center distance, **F**)

bottom distance versus distance travelled, **G**) bottom distance versus percent explored, **H**)

center distance versus distance travelled, **I**) center distance versus percent explored, and **J**)

distance travelled versus percent explored. * - $P < 0.05$, † - $P < 0.10$ compared to all other

groups or those indicated, n = 50-58.

Figure 3. Clustering of zebrafish exploratory behavior during exposure to a novel tank. **A**) Two-

dimensional representation of the four-dimensional behavioral space using a uniform manifold

approximation (McInnes et al., 2020). Clusters (outer circles) are derived from Louvain

community finding applied to a k-nearest neighbor network using 426 fish. Each point is an

individual fish where the shape represents the sex (circle: female, triangle: male), inner color the

strain, and outer color the behavioral cluster. **B**) Individual behaviors associated with each

cluster as box plots indicating median (center line), interquartile range (box ends), and 1.5 times

the interquartile range (whiskers). **C**) Percentage of fish that fall into each cluster across strain

and sex. Striped bars: $P < 0.05$ using randomized permutation tests and FDR corrections, n =

50-58.

633   *Figure 4*. Clusters across two consecutive exposures to the novel tank. **A**) Chord diagram

634   capturing how cluster membership changes from day 1 to day 2 of novel tank exposure. **B**)

635   Percentage of fish that fall into each cluster across strain and sex on the second day of

636   exposure to the novel tank. Striped bars: P < 0.05 using randomized permutation tests and FDR

637   corrections, n = 50-58.

Figure 5. Behavioral consistency in the novel tank across five consecutive days. **A**) Cluster

membership of AB (n = 46) or TU (n = 50) fish that were exposed to the novel tank on five

consecutive days. **B**) Percent of overlapping clusters in AB and TU fish across consecutive

exposure to the novel tank across five days. * - P < 0.05 using randomized permutation test and

FDR corrections.

References

Ampatzis K, Dermon CR. 2016. Sexual dimorphisms in swimming behavior, cerebral metabolic activity
and adrenoceptors in adult zebrafish (Danio rerio). *Behavioural Brain Research* **312**:385–393.
doi:10.1016/j.bbr.2016.06.047
Ariyomo TO, Watt PJ. 2015. Effect of hunger level and time of day on boldness and aggression in the
zebrafish Danio rerio. *Journal of Fish Biology* **86**:1852–1859. doi:10.1111/jfb.12674
Audira G, Siregar P, Strungaru S-A, Huang J-C, Hsiao C-D. 2020. Which Zebrafish Strains Are More
Suitable to Perform Behavioral Studies? A Comprehensive Comparison by Phenomic Approach.
Biology **9**:200. doi:10.3390/biology9080200
Baker MR, Goodman AC, Santo JB, Wong RY. 2018. Repeatability and reliability of exploratory behavior
in proactive and reactive zebrafish, Danio rerio. *Sci Rep* **8**:12114. doi:10.1038/s41598-018-
30630-3
Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach
to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**:289–
300. doi:10.1111/j.2517-6161.1995.tb02031.x
Biro PA, Stamps JA. 2010. Do consistent individual differences in metabolic rate promote consistent
individual differences in behavior? *Trends in Ecology & Evolution* **25**:653–659.
doi:10.1016/j.tree.2010.08.003
Blaser RE, Chadwick L, McGinnis GC. 2010. Behavioral measures of anxiety in zebrafish (Danio rerio).
Behavioural Brain Research **208**:56–62. doi:10.1016/j.bbr.2009.11.009
Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E. 2008. Fast unfolding of communities in large
networks. *Journal of statistical mechanics: theory and experiment* **2008**:P10008.
Caliński T, Harabasz J. 1974. A dendrite method for cluster analysis. *Communications in Statistics-theory*
and Methods **3**:1–27.
Carere C, Drent PJ, Privitera L, Koolhaas JM, Groothuis TGG. 2005. Personalities in great tits, Parus
major: stability and consistency. *Animal Behaviour* **70**:795–805.
doi:10.1016/j.anbehav.2005.01.003
Champagne DL, Hoefnagels CCM, de Kloet RE, Richardson MK. 2010. Translating rodent behavioral
repertoire to zebrafish (Danio rerio): Relevance for stress research. *Behavioural Brain Research*
214:332–342. doi:10.1016/j.bbr.2010.06.001
Clayman CL, Malloy EJ, Kearns DN, Connaughton VP. 2017. Differential behavioral effects of ethanol pre-
exposure in male and female zebrafish (Danio rerio). *Behavioural Brain Research* **335**:174–184.
doi:10.1016/j.bbr.2017.08.007
Csardi G, Nepusz T. 2006. The igraph software package for complex network research. *InterJournal,*
Complex Systems **1695**:1–9.
Dahlbom SJ, Lagman D, Lundstedt-Enkel K, Sundström LF, Winberg S. 2011. Boldness Predicts Social
Status in Zebrafish (Danio rerio). *PLOS ONE* **6**:e23565. doi:10.1371/journal.pone.0023565
Dall SRX, Bell AM, Bolnick DI, Ratnieks FLW. 2012. An evolutionary ecology of individual differences.
Ecology Letters **15**:1189–1198. doi:10.1111/j.1461-0248.2012.01846.x
Davies DL, Bouldin DW. 1979. A cluster separation measure. *IEEE transactions on pattern analysis and*
machine intelligence 224–227.
Desgraupes B. 2013. Clustering indices. *University of Paris Ouest-Lab Modal'X* **1**:34.
Egan RJ, Bergner CL, Hart PC, Cachat JM, Canavello PR, Elegante MF, Elkhayat SI, Bartels BK, Tien AK,
Tien DH, Mohnot S, Beeson E, Glasgow E, Amri H, Zukowska Z, Kalueff AV. 2009. Understanding
behavioral and physiological phenotypes of stress and anxiety in zebrafish. *Behavioural Brain*
Research **205**:38–44. doi:10.1016/j.bbr.2009.06.022

Fontana BD, Cleal M, Parker MO. 2019. Female adult zebrafish (*Danio rerio*) show higher levels of
anxiety-like behavior than males, but do not differ in learning and memory capacity. *European*
Journal of Neuroscience ejn.14588. doi:10.1111/ejn.14588
Forkosh O, Karamihalev S, Roeh S, Alon U, Anpilov S, Touma C, Nussbaumer M, Flachskamm C, Kaplick
PM, Shemesh Y, Chen A. 2019. Identity domains capture individual differences from across the
behavioral repertoire. *Nat Neurosci* **22**:2023–2028. doi:10.1038/s41593-019-0516-y
Frossard J, Renaud O. 2021. Permutation tests for regression, ANOVA, and comparison of signals: the
permuco package. *Journal of Statistical Software* **99**:1–32.
Gerlai R. 2020. Evolutionary conservation, translational relevance and cognitive function: The future of
zebrafish in behavioral neuroscience. *Neuroscience & Biobehavioral Reviews* **116**:426–435.
doi:10.1016/j.neubiorev.2020.07.009
Gosling SD. 2001. From mice to men: What can we learn about personality from animal research?
Psychological Bulletin **127**:45–86. doi:10.1037/0033-2909.127.1.45
Grant M. 2000. Galen on Food and Diet. Florence, UNITED STATES: Taylor & Francis Group.
Gu Z, Gu L, Eils R, Schlesner M, Brors B. 2014. Circlize implements and enhances circular visualization in
R. *Bioinformatics* **30**:2811–2812.
Hervé M, Hervé MM. 2020. Package 'RVAideMemoire.' *See https://CRANR-projectorg/package=*
RVAideMemoire.
Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, Collins JE, Humphray S, McLaren K,
Matthews L, McLaren S, Sealy I, Caccamo M, Churcher C, Scott C, Barrett JC, Koch R, Rauch GJ,
White S, Chow W, Kilian B, Quintais LT, Guerra-Assunção JA, Zhou Y, Gu Y, Yen J, Vogel JH, Eyre
T, Redmond S, Banerjee R, Chi J, Fu B, Langley E, Maguire SF, Laird GK, Lloyd D, Kenyon E,
Donaldson S, Sehra H, Almeida-King J, Loveland J, Trevanion S, Jones M, Quail M, Willey D, Hunt
A, Burton J, Sims S, McLay K, Plumb B, Davis J, Clee C, Oliver K, Clark R, Riddle C, Eliott D,
Threadgold G, Harden G, Ware D, Mortimer B, Kerry G, Heath P, Phillimore B, Tracey A, Corby N,
Dunn M, Johnson C, Wood J, Clark S, Pelan S, Griffiths G, Smith M, Glithero R, Howden P, Barker
N, Stevens C, Harley J, Holt K, Panagiotidis G, Lovell J, Beasley H, Henderson C, Gordon D, Auger
K, Wright D, Collins J, Raisen C, Dyer L, Leung K, Robertson L, Ambridge K, Leongamornlert D,
McGuire S, Gilderthorp R, Griffiths C, Manthravadi D, Nichol S, Barker G, Whitehead S, Kay M,
Brown J, Murnane C, Gray E, Humphries M, Sycamore N, Barker D, Saunders D, Wallis J, Babbage
A, Hammond S, Mashreghi-Mohammadi M, Barr L, Martin S, Wray P, Ellington A, Matthews N,
Ellwood M, Woodmansey R, Clark G, Cooper J, Tromans A, Grafham D, Skuce C, Pandian R,
Andrews R, Harrison E, Kimberley A, Garnett J, Fosker N, Hall R, Garner P, Kelly D, Bird C, Palmer
S, Gehring I, Berger A, Dooley CM, Ersan-Ürün Z, Eser C, Geiger H, Geisler M, Karotki L, Kirn A,
Konantz J, Konantz M, Oberländer M, Rudolph-Geiger S, Teucke M, Osoegawa K, Zhu B, Rapp A,
Widaa S, Langford C, Yang F, Carter NP, Harrow J, Ning Z, Herrero J, Searle SMJ, Enright A,
Geisler R, Plasterk RHA, Lee C, Westerfield M, De Jong PJ, Zon LI, Postlethwait JH, Nüsslein-
Volhard C, Hubbard TJP, Crollius HR, Rogers J, Stemple DL. 2013. The zebrafish reference
genome sequence and its relationship to the human genome. *Nature* **496**:498–503.
doi:10.1038/nature12111
Kalueff AV, Gebhardt M, Stewart AM, Cachat JM, Brimmer M, Chawla JS, Craddock C, Kyzar EJ, Roth A,
Landsman S, Gaikwad S, Robinson K, Baatrup E, Tierney K, Shamchuk A, Norton W, Miller N,
Nicolson T, Braubach O, Gilman CP, Pittman J, Rosemberg DB, Gerlai R, Echevarria D, Lamb E,
Neuhauss SCFF, Weng W, Bally-Cuif L, Schneider H, Schneider, and the Zebrafish Neuros H. 2013.
Towards a comprehensive catalog of zebrafish behavior 1.0 and beyond. *Zebrafish* **10**:70–86.
doi:10.1089/zeb.2012.0861
Kenney JW. 2020. Associative and nonassociative learning in adult zebrafishBehavioral and Neural
Genetics of Zebrafish. Elsevier. pp. 187–204. doi:10.1016/b978-0-12-817528-6.00012-7

Kenney JW, Steadman PE, Young O, Shi MT, Polanco M, Dubaishi S, Covert K, Mueller T, Frankland PW.
2021. A 3D adult zebrafish brain atlas (AZBA) for the digital age. *eLife* **10**:e69988.
doi:10.7554/eLife.69988
Kuroda T. 2018. A system for the real-time tracking of operant behavior as an application of 3D camera.
Journal of the Experimental Analysis of Behavior **110**:522–544. doi:10.1002/jeab.471
López P, Hawlena D, Polo V, Amo L, Martín J. 2005. Sources of individual shy–bold variations in
antipredator behaviour of male Iberian rock lizards. *Animal Behaviour* **69**:1–9.
doi:10.1016/j.anbehav.2004.05.010
Marchette DJ, Marchette MDJ, Matrix S. 2015. Package 'cccd.'
Martins EP, Bhat A. 2014. Population-level personalities in zebrafish: aggression-boldness across but not
within populations. *Behavioral Ecology* **25**:368–373. doi:10.1093/beheco/aru007
Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M. 2018. DeepLabCut:
markerless pose estimation of user-defined body parts with deep learning. *Nature Neuroscience*
21:1281–1289. doi:10.1038/s41593-018-0209-y
Maximino C, Puty B, Matos Oliveira KR, Herculano AM. 2013. Behavioral and neurochemical changes in
the zebrafish leopard strain. *Genes, Brain and Behavior* **12**:576–582. doi:10.1111/gbb.12047
McInnes L, Healy J, Melville J. 2020. UMAP: Uniform Manifold Approximation and Projection for
Dimension Reduction. *arXiv:180203426 [cs, stat]*.
Mustafa A, Roman E, Winberg S. 2019. Boldness in Male and Female Zebrafish (Danio rerio) Is
Dependent on Strain and Test. *Frontiers in Behavioral Neuroscience* **13**:248.
doi:10.3389/fnbeh.2019.00248
Myers PJ, Young JK. 2018. Consistent individual behavior: evidence of personality in black bears. *J Ethol*
36:117–124. doi:10.1007/s10164-018-0541-4
Nasiadka A, Clark MD. 2012. Zebrafish Breeding in the Laboratory Environment. *ILAR Journal* **53**:161–
168. doi:10.1093/ilar.53.2.161
Neuwirth E, Brewer RC. 2014. ColorBrewer palettes. *R package version* **1**.
Oswald ME, Drew RE, Racine M, Murdoch GK, Robison BD. 2012. Is Behavioral Variation along the Bold-
Shy Continuum Associated with Variation in the Stress Axis in Zebrafish? *Physiological and*
Biochemical Zoology **85**:718–728. doi:10.1086/668203
Pannia E, Tran S, Rampersad M, Gerlai R. 2014. Acute ethanol exposure induces behavioural differences
in two zebrafish (Danio rerio) strains: A time course analysis. *Behavioural Brain Research*
259:174–185. doi:10.1016/j.bbr.2013.11.006
Panula P, Chen YC, Priyadarshini M, Kudo H, Semenova S, Sundvik M, Sallinen V. 2010. The comparative
neuroanatomy and neurochemistry of zebrafish CNS systems of relevance to human
neuropsychiatric diseases. *Neurobiology of Disease* **40**:46–57. doi:10.1016/j.nbd.2010.05.010
Parker MO, Millington ME, Combe FJ, Brennan CH. 2012. Housing Conditions Differentially Affect
Physiological and Behavioural Stress Responses of Zebrafish, as well as the Response to
Anxiolytics. *PLOS ONE* **7**:e34992. doi:10.1371/journal.pone.0034992
Philpott C, Donack CJ, Cousin MA, Pierret C. 2012. Reducing the noise in behavioral assays: sex and age
in adult zebrafish locomotion. *Zebrafish* **9**:191–194.
Press WH, Teukolsky SA. 1990. Savitzky-Golay smoothing filters. *Computers in Physics* **4**:669–672.
R Core Team, R Development Core Team R. 2016. R: A Language and Environment for Statistical
Computing. *R Foundation for Statistical Computing* **1**:409. doi:10.1007/978-3-540-74686-7
Rafferty SA, Quinn TA. 2018. A beginner's guide to understanding and implementing the genetic
modification of zebrafish. *Progress in Biophysics and Molecular Biology*, The Use of Zebrafish for
Cardiac Research **138**:3–19. doi:10.1016/j.pbiomolbio.2018.07.005
Rambo CL, Mocelin R, Marcon M, Villanova D, Koakoski G, de Abreu MS, Oliveira TA, Barcellos LJG, Piato
AL, Bonan CD. 2017. Gender differences in aggression and cortisol levels in zebrafish subjected

to unpredictable chronic stress. *Physiology & Behavior* **171**:50–54.
doi:10.1016/j.physbeh.2016.12.032
Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ. 2007. Integrating animal temperament within
ecology and evolution. *Biological Reviews* **82**:291–318. doi:10.1111/j.1469-185X.2007.00010.x
Reolon GK, de Melo GM, da Rosa JG dos S, Barcellos LJG, Bonan CD. 2018. Sex and the housing: Effects
on behavior, cortisol levels and weight in zebrafish. *Behavioural Brain Research* **336**:85–92.
doi:10.1016/j.bbr.2017.08.006
Rosa LV, Ardais AP, Costa FV, Fontana BD, Quadros VA, Porciúncula LO, Rosemberg DB. 2018. Different
effects of caffeine on behavioral neurophenotypes of two zebrafish populations. *Pharmacology*
Biochemistry and Behavior **165**:1–8. doi:10.1016/j.pbb.2017.12.002
Rousseeuw PJ. 1987. Silhouettes: A graphical aid to the interpretation and validation of cluster analysis.
Journal of Computational and Applied Mathematics **20**:53–65. doi:10.1016/0377-
0427(87)90125-7
Sackerman J, Donegan JJ, Cunningham CS, Nguyen NN, Lawless K, Long A, Benno RH, Gould GG. 2010.
Zebrafish Behavior in Novel Environments: Effects of Acute Exposure to Anxiolytic Compounds
and Choice of Danio rerio Line. *International journal of comparative psychology* **23**:43–61.
Shams S, Chatterjee D, Gerlai R. 2015. Chronic social isolation affects thigmotaxis and whole-brain
serotonin levels in adult zebrafish. *Behavioural Brain Research* **292**:283–287.
doi:10.1016/j.bbr.2015.05.061
Sih A, Bell A, Johnson JC. 2004. Behavioral syndromes: an ecological and evolutionary overview. *Trends*
in Ecology & Evolution **19**:372–378. doi:10.1016/j.tree.2004.04.009
Simon P, Dupuis R, Costentin J. 1994. Thigmotaxis as an index of anxiety in mice. Influence of
dopaminergic transmissions. *Behavioural Brain Research* **61**:59–64. doi:10.1016/0166-
4328(94)90008-6
Taft RA, Davisson M, Wiles MV. 2006. Know thy mouse. *Trends in Genetics* **22**:649–653.
doi:10.1016/j.tig.2006.09.010
Thörnqvist P-O, McCarrick S, Ericsson M, Roman E, Winberg S. 2019. Bold zebrafish (Danio rerio) express
higher levels of delta opioid and dopamine D2 receptors in the brain compared to shy fish.
Behavioural Brain Research **359**:927–934. doi:10.1016/j.bbr.2018.06.017
Toms CN, Echevarria DJ, Jouandot DJ. 2010. A Methodological Review of Personality-Related Studies in
Fish: Focus on the Shy-Bold Axis of Behavior. *International Journal of Comparative Psychology*
23.
Tran S, Gerlai R. 2013. Individual differences in activity levels in zebrafish (Danio rerio). *Behavioural*
Brain Research **257**:224–229. doi:10.1016/j.bbr.2013.09.040
Vignet C, Bégout M-L, Péan S, Lyphout L, Leguay D, Cousin X. 2013. Systematic Screening of Behavioral
Responses in Two Zebrafish Strains. *Zebrafish* **10**:365–375. doi:10.1089/zeb.2013.0871
Volgin AD, Yakovlev OA, Demin KA, de Abreu MS, Alekseeva PA, Friend AJ, Lakstygal AM, Amstislavskaya
TG, Bao W, Song C, Kalueff AV. 2019. Zebrafish models for personalized psychiatry: Insights from
individual, strain and sex differences, and modeling gene x environment interactions. *Journal of*
Neuroscience Research **97**:402–413. doi:10.1002/jnr.24337
Wickham H. 2015. Elegant Graphics for Data Analysis, Media. Springer. doi:10.1007/978-0-387-98141-3
Wilson DS, Clark AB, Coleman K, Dearstyne T. 1994. Shyness and boldness in humans and other animals.
Trends in Ecology & Evolution **9**:442–446. doi:10.1016/0169-5347(94)90134-1
Wong K, Elegante M, Bartels B, Elkhayat S, Tien D, Roy S, Goodspeed J, Suciu C, Tan J, Grimes C, Chung A,
Rosenberg M, Gaikwad S, Denmark A, Jackson A, Kadri F, Chung KM, Stewart A, Gilder T, Beeson
E, Zapolsky I, Wu N, Cachat J, Kalueff AV. 2010. Analyzing habituation responses to novelty in
zebrafish (Danio rerio). *Behavioural Brain Research* **208**:450–457. doi:10.1016/j.bbr.2009.12.023

Wong RY, Perrin F, Oxendine SE, Kezios ZD, Sawyer S, Zhou L, Dereje S, Godwin J. 2012. Comparing
841   behavioral responses across multiple assays of stress and anxiety in zebrafish (Danio rerio).
842   *Behaviour* **149**:1205–1240.
843   Wulliman MF, Rupp B, Reichert H, Wullimann MF, Rupp B, Reichert H. 1996. Neuroanatomy of the
844   Zebrafish Brain: A Topological Atlas, Neuroanatomy of the Zebrafish Brain. Birkhäuser Verlag,
845   Basel. doi:10.1007/978-3-0348-8979-7
846   Yuan M, Chen Y, Huang Y, Lu W. 2018. Behavioral and Metabolic Phenotype Indicate Personality in
847   Zebrafish (Danio rerio). *Frontiers in Physiology* **9**:653. doi:10.3389/fphys.2018.00653

864   **Supplemental Information**

866   *Figure S1*. Distribution of values for individual behavioral parameters. Distributions across strain

867   and sex for **A**) bottom distance, **B**) center distance, **C**) distance travelled, and **D**) percent tank

868   explored. * - P < 0.05, † - P < 0.10 Shapiro-Wilks test for normality, n = 50-58.

Figure S2. Individual exploratory behaviors across time during a single exposure to the novel

tank. The effect of strain (color) and sex (shape, circle: female, triangle: male) across time for **A**)

bottom distance, **B**) center distance, and **C**) distance travelled. * - P < 0.05, † - P < 0.10 for time

interval or sex as indicated, n = 50-58. Data presented as mean ± SEM.

Figure S3. Relationship between pair-wise individual behavioral parameters. Scatterplots across

strain and sex of **A**) bottom distance versus center distance, **B**) bottom distance versus distance

travelled, **C**) bottom distance versus percent explored, **D**) center distance versus distance

887   travelled, **E**) center distance versus percent explored, and **F**) distance travelled versus percent

888   explored. n = 50-58.

Figure S4. Internal clustering metrics for knn Louvain clustering. The parameter k was varied for

generating knn's, and the **A**) Calinski-Harbasz, **B**) silhouette, and **C**) Davies-Bouldin indices

were calculated at each k. Area highlighted in red indicates region where clustering metrics

were near optimal and robust to small changes in k.

902   *Figure S5*. Changes in individual exploratory behaviors during two consecutive exposures to the

903   novel tank. The effect of strain and sex on changes over two days in **A**) bottom distance, **B**)

904   center distance, **C**) distance travelled, and **D**) percent explored. Boxplots indicate median

905   (center line), interquartile range (box ends), and 1.5 times the interquartile range (whiskers). * -

906   P < 0.05, † - P < 0.10 compared to zero, n = 50-58.

Figure S6. Individual exploratory behaviors during five consecutive days of exposure to the novel tank. AB (top) and TU (bottom) fish were exposed to the novel tank on five consecutive days and **A**) bottom distance, **B**) center distance, **C**) distance travelled, and **D**) percent explored were measured. * - P < 0.05, † - P < 0.10 for day or sex and indicated, n = 21-26. Data presented as mean ± SEM.

929   *Figure S7*. Percent cluster membership across five consecutive days of exposure to the novel

930   tank for AB (left) and TU (right) fish.