1

Beyond bold versus shy: Zebrafish exploratory behavior falls into several behavioral clusters and is influenced by strain and sex

Neha Rajput, Kush Parikh, and Justin W. Kenney

Department of Biological Sciences
Wayne State University
Detroit, MI 48202

Abstract

Consistent individual differences in exploratory behavior have been found across a range of

taxa, including zebrafish, and are thought to contribute to evolutionary fitness. Animals that

explore more of a novel environment, and visit areas of high predation risk, are considered bold,

whereas animals with the opposite pattern of behavior are considered shy. Here, we examined

whether this bimodal characterization of bold versus shy adequately captures the breadth of

exploratory behavior exhibited by zebrafish or if, instead, behavior falls into multiple distinct

behavioral subtypes. To identify the presence of behavioral subtypes, we applied unsupervised

machine learning to behaviors extracted from three-dimensional swim traces from over 400

adult zebrafish across four strains (AB, TL, TU, and WIK) and both sexes. We found that

behavior stratified into four distinct clusters. These included previously described bold and shy

behavior as well as two new behavioral types: wall-huggers and active explorers. Consistent

with prior work that found individual differences to be stable across time and influenced by

biological factors like genetics and sex, the behavioral subtypes we identified were stable for up

to 10 weeks and were influenced by the strain and sex of the animals. Taken together, our work

suggests that individual differences in zebrafish exploratory behavior goes beyond bold versus

shy and exhibits greater complexity than is typically assumed.

Introduction

Interest in the biology of behavioral differences can be traced back to at least Roman

antiquity where Galen applied the humoral theory to explain variation in human temperament

(as translated in Grant, 2000). Today, we have a greater understanding of human personalities,

defined as behavioral tendencies that are consistent across time and context, but its biological

basis remains elusive. One avenue for progress is the modeling of human personality through

the study of individual difference in animal behavior. Often dismissed as noise around an

average, a growing body of work has found that variations in animal behavior are often

consistent across time and context (Dall et al., 2012; Sih et al., 2004). Such consistent

differences have been described for behaviors important for evolutionary fitness, and in a wide

range of taxa, suggesting that they may provide grist for adaptation to an ever-changing

environment.

One of the most widely studied axes of behavioral variation in animals is the bold-shy

axis. Bold animals tend to explore or investigate novel environments or objects more readily

than shy animals that tend to flee or retreat in response to novelty (Réale et al., 2007; Toms et

al., 2010; Wilson et al., 1994). From a fitness perspective, boldness may be adaptive when food

resources are scarce and predation risk is low, whereas shyness may be more effective when

the opposite conditions prevail (Réale et al., 2007). Variation along this axis has been described

in animals ranging from bears (Myers and Young, 2018), lizards (López et al., 2005), birds

(Carere et al., 2005), and fish (Toms et al., 2010). Studies examining the bold-shy axis typically

begin with the assumption that animals fall into one of these two categories. However, whether

this bimodal distribution of bold versus shy fully captures animal behavior has not been tested

directly. It is likely that there is more complexity to animal behavioral types which may have

been overlooked due to use of small sample sizes or assessment of only one or two specific

behaviors (Forkosh et al., 2019).

Zebrafish have proven to be an excellent model organism to understand behavior and its

biological basis. With 70% of fish genes having an obvious human ortholog (Howe et al., 2013),

and a central nervous system that has the same general organization and utilizes many of the

same neurotransmitters as mammals (Kenney et al., 2021; Panula et al., 2010; Wulliman et al.,

1996), findings using zebrafish are widely applicable. Our understanding of zebrafish behavior

has expanded rapidly over the past decade (Gerlai, 2020; Kalueff et al., 2013; Kenney, 2020),

including several studies that have examined the bold-shy axis. Boldness is often probed by

exposing fish to a novel tank and examining locomotion, exploratory activity and avoidance

behaviors, like geotaxis (i.e. bottom dwelling) or thigmotaxis (i.e. proximity to tank walls)

(Mustafa et al., 2019; Oswald et al., 2012; Thörnqvist et al., 2019; Toms et al., 2010). Animals

that are more active or spend more time in parts of the tank that would increase risk of predation

(i.e. the top and/or center of the tank) are considered bolder. In zebrafish, these behaviors have

been found to be consistent over time (Baker et al., 2018; Tran and Gerlai, 2013) and can

predict other behaviors like social dominance (Dahlbom et al., 2011), aggression (Martins and

Bhat, 2014), and stress reactivity (Oswald et al., 2012), all hallmarks of personality. However,

fully utilizing zebrafish exploratory behavior to understand the biological basis of individual

differences requires that we first determine if the bold versus shy distinction adequately

captures the breadth of behavioral variability exhibited during exploration.

To determine the presence of multiple behavioral clusters during exploration of a novel

tank we captured three-dimensional swim behavior from over four hundred fish. Because

zebrafish behavior is known to be influenced by strain and sex (Volgin et al., 2019), we used

animals from four inbred strains (AB, TU, WIK, and TL) and both sexes to ensure that we

captured the full range of behavioral variability. Using an unsupervised machine learning

approach to identify behavioral clusters we found that exploratory behavior stratified into four

clusters. These clusters included traditional descriptions of bold and shy, as well as two

additional behavioral types we call wall-huggers and active explorers. Consistent with these

behavioral subtypes being akin to personality types, we found that individual cluster

membership remained consistent across days and weeks, and that the proportion of fish in each

cluster was influenced by strain and sex.

115   **Results**

116   *Three-dimensional behavioral tracking*

117   To capture three-dimensional zebrafish swim behavior during exploration of a novel

tank, we used Intel RealSense™ cameras mounted above five-sided tanks with frosted walls

(Figure 1A). These cameras simultaneously captured color and depth streams, resulting in

three-dimensional videos (Figure 1B). Fish posture at each frame was tracked in the color

stream using DeepLabCut (Figure 1C; Mathis et al., 2018). These points were overlayed onto

the depth stream to create three-dimensional swim traces (Figure 1D) from which we extracted

positional information (distance from bottom and center), distance travelled, and percent tank

explored (Figure 1E).

Influence of sex and strain on exploratory behaviors

127   We first determined if sex and genetics influences individual zebrafish exploratory

behaviors by assessing swim traces of fish from four strains (AB, TU, TL, and WIK), and both

sexes, on two consecutive days. We extracted four exploratory behaviors from each swim trace:

distance from bottom, distance from center, total distance travelled, and percentage of tank

explored (Figure 1E). We found that the distribution of several of the parameters deviated from

normality (Figure S1), so we performed non-parametric 4 × 2 (strain × sex) permutation

ANOVAs. For distance from bottom (Figure 2A), we found a main effect of strain ($P = 0.0001$),

but no effect of sex ($P = 0.40$) nor an interaction ($P = 0.44$). FDR (false discovery rate) corrected

permutation t-tests found that TL fish swam closer to the bottom of the tank than all other strains

(P's ≤ 0.0008). For distance from center (Figure 2B) there was a main effect of sex ($P = 0.042$)

and a trend towards an interaction ($P = 0.060$) where female fish of every strain, except AB,

spent more time closer to the center of the tank. We also found a main effect of strain ($P =$

0.0001) with FDR corrected permutation t-tests indicating that TL fish swam closer to the center

of the tank than all other strains (P's = 0.0004), and AB spent more time on the periphery (AB

compared to TU: P = 0.0006, WIK: P = 0.017). For distance travelled (Figure 2C), we found a

main effect of sex (P = 0.0001) where male fish swam further than female fish. There was also a

main effect of strain (P = 0.014), but no interaction (P = 0.59). FDR corrected permutation t-tests

found that TL fish swam less than AB fish (P = 0.043) and a trend towards a difference between

TLs and WIKs (P = 0.061). Finally, for percent of the tank explored (Figure 2D), we also found a

main effect of sex (P = 0.024) in which female fish explored less of the tank than their male

counterparts. There was also a main effect of strain (P = 0.0001), but no interaction (P = 0.34).

Post-hoc tests revealed that TL fish explored the tank less than all other strains (P's = 0.0004),

and that AB fish explored less than TU (P = 0.035) and WIK (P = 0.0026) fish. Taken together,

we find that there are several sex differences (center distance, distance travelled, and percent

explored) and that TL fish differ the most from other strains across all measures with no clear

strain by sex interactions.

Influence of sex and strain on habituation to the novel tank

Some exploratory behaviors have been found to habituate over a single six-minute

exposure to a novel tank (Wong et al., 2010), so we examined whether any of the parameters

we measured changed over time and were influenced by sex in the various strains (Figure S2).

For each measure, we used non-parametric 2 × 6 (sex × time interval) mixed permutation

ANOVAs and adjusted for multiple tests using an FDR correction. For distance from bottom

(Figure S2A), we found no main effects of sex (AB: P = 0.51 , TL: P = 0.46 , TU: P = 0.74, WIK:

P = 0.86), but found effects of interval in all strains (AB: P = 0.0006, TL: P = 0.0093, TU: P =

0.0006, WIK: P = 0.0084) where all fish, except WIKs, increased their distance from the bottom

over time. There were no interactions except for WIKs (AB: P = 0.086, TU: P = 0.74, TL: P =

0.46, WIK: P = 0.018), where female fish appeared to decrease their bottom distance over time

whereas male fish showed little change across the trial. For distance from center (Figure S2B),

a main effect of interval was found for AB, TU, and WIKs (P's = 0.0004, TL: P = 0.52), finding

that these strains increased their distance from center over time. There were no interactions

between sex and time interval (AB: P = 0.91, TL: P = 0.28, TU: P = 0.91, WIK: P = 0.91), but we

found trends towards an effect of sex in TU and WIKs (AB: P = 0.28, TL: P = 0.41, TU: P =

0.054, WIK: P = 0.064) where female fish spent their time closer to the center of the tank than

male fish, consistent with what was observed in the overall data (Figure 2B). Finally, for

distance travelled (Figure S2C), there was an increase in locomotor activity over time in AB, TU,

and WIK fish, with a trend in TLs (AB: P = 0.0004, TL: P = 0.093, TU: P = 0.0012, WIK: P =

0.0004). Consistent with the overall data, there were also main effects of sex in all strains

except TLs, where there was a trend (AB: P = 0.0014, TL: P = 0.093, TU: P = 0.0004, WIK: P =

0.0048). Only the TU fish had an interaction between time interval and sex where TU female

fish increased their distance travelled over time, but male fish did not (AB: P = 0.085, TL: P =

0.39, TU: P = 0.0022, WIK: P = 0.068).

Correlations between behavioral parameters

To determine the extent to which individual behavioral parameters captured distinct

elements of exploratory behavior, and if there was any influence of sex and genetics on these

relationships, we computed correlations between individual measures (Figure 2E-J and S3). We

used Spearman's ρ to identify monotonic relationships because of the presence of several non-

normally disturbed parameters (Figure S1). As expected, we found that distance travelled was

consistently positively correlated with percent explored across all strains and sexes (Figures 2J

& S3F). We also found consistent positive correlations between bottom distance and percent

explored (Figures 2G & S3C), which is in line with the idea that a higher bottom distance is

associated with an increased willingness to explore. However, bottom distance did not correlate

consistently with distance travelled (Figures 2F & S3B), suggesting that, despite positive

correlations between distance travelled and percent explored, these two exploratory measures

are capturing different aspects of exploration. To our surprise, bottom distance and center

distance did not consistently correlate with each other (Figures 2E & S3A): in three strain/sexes

there was a clear negative correlation (female TLs and TUs, and male WIKs) where fish that

swam nearer to the top also swam closer to the center, but in one group (male ABs) the

opposite relationship was observed with no clear relationships in the remaining groups. Center

distance was mostly negatively correlated with percent explored (Figures 2I & S3E), but not

universally so (TLs being the exception), largely consistent with the idea that fish that spend

more time closer to the center of the tank also explore more of the tank.

Identifying behavioral clusters

Given the presence of non-normal behavioral distributions (Figure S1) and variability in

the relationship between different exploratory parameters (Figure 2E-J), we hypothesized the

presence of multiple behavioral clusters in our data set. To test this, we built a k-nearest

neighbor network and applied the Louvain community detection algorithm to identify clusters

(Blondel et al., 2008). We used all four behavioral parameters (bottom distance, center distance,

distance travelled, and percent explored) in calculating nearest neighbor distances because

none of the behavioral measures showed consistently high correlations across all strains and

sexes (Figure 2E-J). To determine 'k' for building the network, we explored a range of values

and chose a value (k = 118) that optimized internal clustering metrics and was robust to small

deviations in k (Figure S4) resulting in the identification of four distinct behavioral clusters

(Figure 3A).

An examination of the behaviors associated with each of the clusters reveals a range of

behavioral profiles that run the gamut from shy to bold (Figure 3B). Fish in the shyest cluster

were lowest in both bottom distance and percent exploration (Figure 3B; cluster 1) whereas fish

in the boldest cluster spent the most time near the top and center of the tank while exploring

more of the tank than average (Figure 3B; cluster 4). Bold fish were also amongst the lowest in

distance travelled, suggesting that they were particularly efficient in their tank exploration. Two

'mixed' clusters were also identified: cluster 2 where fish were near average on all measures

except for center distance where they spent most of their time near the periphery, a group we

call 'wall-huggers', and cluster 3 where fish were above average in distance travelled and

percent explored that we call 'active explorers'.

Behavioral clusters across sex and strain

To determine if fish from a given strain and sex were over- or underrepresented in

behavioral clusters, we computed p-values using permutation resampling and FDR corrections

for multiple comparisons (Figure 3C). We found that, for the shy cluster (cluster 1), TL fish,

irrespective of sex, were overrepresented (female: P = 0.0011, male: P = 0.048), whereas the

wall-huggers (cluster 2) had overrepresentation of female AB fish (P = 0.0053). Consistent with

our finding that males, on average, swam more than females, more male fish were in the active

explorers group (cluster 3), although overrepresentation was only significant in TU and WIK fish

(males: AB: P = 0.086, TL: P = 0.23, TU: P = 0.0016, WIK: P = 0.0073). Finally, in the boldest

cluster (cluster 4), females outnumbered males in all strains but there was only significant

overrepresentation in TU fish with trends in ABs and WIKs (female: AB: P = 0.082, TL: P = 0.16,

TU: P = 0.018, WIK: P = 0.064).

Habituation over two days

All 426 fish used for generating clusters were exposed to the novel tank on two

consecutive days, allowing us to determine if their behavior remained consistent across

repeated exposures. First, we analyzed individual exploratory behaviors with permutation paired

t-tests and FDR corrections (Figure S5). We found that AB and TL fish increased their bottom

distance during the second exposure, with a trend in female WIK fish (Figure S5A; female AB: P

= 0.0080, male AB: P = 0.012, female TL: P = 0.0016, male TL: P = 0.0040, female TU: P =

0.54, male TU: P = 0.54, female WIK: P = 0.055, male WIK: P = 0.62). Thigmotaxis (center

distance) also increased in several groups: AB fish and female WIKs with trends towards an

increase in TU females, but a decrease in TL males (Figure S5B; female AB: P = 0.0032, male

AB: P = 0.043, female TL: P = 0.38, male TL: P = 0.073, female TU: P = 0.073, male TU: P =

0.82, female WIK: P = 0.043, male WIK: P = 0.87). Distance travelled did not change in any fish

(Figure S5C; P's ≥ 0.12), and percent explored had trends towards a decrease in AB males and

increase in TL females (Figure S5D; female AB: P = 0.13, male AB: P = 0.071, female TL: P =

0.071, male TL: P = 0.55, female TU: P = 0.82, male TU: P = 0.39, female WIK: P = 0.55, male

WIK: P = 0.31). Taken together, the changes in behavior during the second exposure were

mixed, with some changes indicating an increase in behaviors associated with boldness, like

bottom distance, and others an increase in putative shy behaviors, like thigmotaxis (however,

please see our discussion on thigmotaxis below).

Behavioral cluster consistency over two days

To determine if the behavioral clusters we identified remained consistent across days,

we used exploratory data from the second day of exposure to the novel tank to assign fish to

clusters uncovered on the first day (Figure 4A). We found that 52.8% of fish fell into the same

behavioral cluster on days 1 and 2 (P = 0.0001; permutation average = 25.1% overlap). Overall,

the number of fish falling into the shyest group (cluster 1) decreased, and the wall-huggers

(cluster 2) remaining the most consistent. At the level of sex and strain (Figure 4B), we again

used FDR corrected permutation resampling, finding that, as on day 1, AB females were

overrepresented in the wall-huggers group (cluster 2; P = 0.0011), but were now joined by their

male counterparts (P = 0.031). TL females also remained overrepresented in the shyest cluster

(P = 0.0011), but a significant portion of male and female TL fish were now in the boldest cluster

(female TL: P = 0.013, male TL: P = 0.0011). TU females were no longer overrepresented in the

boldest group (P = 0.62), although TU males were now underrepresented in this group (P =

0.025) and maintained their overrepresentation as active explorers, cluster 3 (P = 0.0011).

Finally, neither female nor male WIKs were overrepresented in any cluster on day 2.

Habituation over multiple days and weeks

Some exploratory behaviors have been found to habituate over several days of

exposure to a novel tank (Wong et al., 2010). In a new cohort of AB and TU fish, exploratory

behavior was measured over five consecutive days of exposure to the novel tank (Figure S6).

We used 5 × 2 (day × sex) mixed permutation ANOVAs to assess significance. For bottom

distance in AB fish (Figure S6A), there was no effect of day (P = 0.20), sex (P = 0.14) or an

interaction (P = 0.64), whereas TU fish increased their bottom distance over the five days (P =

0.025), with no effect of sex (P = 0.18) nor an interaction (P = 0.87). AB fish increased their

center distance (Figure S6B) over time (P = 0.0001) with no effect of sex (P = 0.12) or an

interaction (P = 0.14), and TU fish had a similar trend (day: P = 0.094, sex: P = 0.27, interaction:

P = 0.93). As we saw before, male AB fish swam further than female fish (P = 0.010) with no

effect of day (P = 0.62). There was a trend towards and interaction between day and sex (P =

0.058) where AB males swam less over time, and females more (Figure S6C). In TU fish, there

was a trend towards males swimming more than females (P = 0.084), and a clear effect of day

(P = 0.0007) and no interaction (P = 0.68) as both sexes increased their distance travelled over

time. Finally, for percent explored (Figure S6D), In AB fish there was no effect of sex (P = 0.33)

or day (P = 0.77), but there was a trend towards an interaction (P 0.068) where female fish

appeared to slightly decrease their percent explored over time. In TU fish, animals increased

their exploration over days (P = 0.0089), with a trend towards males exploring more than

females (P = 0.099), and no interaction of day and sex (P = 0.31).

In a separate cohort of TU fish, we examined individual exploratory behaviors over 10

weeks of biweekly (every other week) exposures to the novel tank (Figure S7). We used 6 × 2

(week × sex) mixed permutation ANOVAs to assess significance. For bottom distance (Figure

S7A), we found no effect of week (P = 0.96) or sex (P = 0.23) but there was an interaction (P =

0.0023) where female fish increased, and males decreased, their distance from bottom across

weeks. For center distance (Figure S7B), there was a trend towards an effect of week (P =

0.076), and no effect of sex (P = 0.49) nor an interaction (P = 0.88). For both distance travelled

(Figure S7C) and percent explored (Figure S7D), there were main effects of sex (P = 0.0001

and P = 0.015, respectively) where, as we saw before, male fish swam further, and explored

more of the tank, than female fish. There were also main effects of week (P = 0.0001 and P =

0.0002, respectively) and no interactions (P = 0.80 and P = 0.25, respectively), where both

female and male fish increased their exploratory behaviors during repeated exposures.

Behavioral cluster consistency over days and weeks

Next, we asked whether the behavioral clusters of individual animals over five

consecutive days or across 10 weeks remained consistent (Figure 5). Across the exposures we

found that exploratory behavior of approximately 50% of animals fell into the same cluster on at

least 4 out of 5 days (Figure 5A) or 5 of 6 biweekly exposures (Figure 5B). To determine if the

consistency across time was greater than chance, each animal was assigned an overlap score:

the sum of pair-wise overlaps across consecutive exposures to the tank. For the daily data this

score ranged from 1 (only one pair of days overlapped) to 10 (all pair-wise overlaps), for the

biweekly data it ranged from 2 to 15. During 5 days of exposure the average scores for AB and

TU fish were 5.6 ± 2.6 and 5.9 ± 3.1 (mean ± standard deviation), respectively, both of which

were significantly higher than the overlap scores from resampling (permutation mean ± standard

deviation: 3.4 ± 0.2 and 2.7 ± 0.2, respectively; P's = 0.0002). For the biweekly data, the

average overlap score was 9.0 ± 4.2, which was significantly higher than chance (4.8 ± 0.2, P =

0.0002). Although most animals had scores higher than the permutation average, a small

subset of animals were "consistently inconsistent" with their exploratory behavior falling into all

four clusters across days or weeks. In both the daily and biweekly data, we found that the

number of animals falling into the shyest cluster (cluster 1) decreased over time (Figure S8).

There were few prominent changes in other clusters except for an increase in wall-huggers

(cluster 2) in AB fish over five consecutive days (Figure S8A) and an increase in active

explorers (cluster 3) during biweekly exposures (Figure S8B).

Finally, we examined how much cluster overlap there was between each of the five

days, or weeks, of the experiment (Figure 5C and D). During the daily exposures, we found that

the clusters from day 1 overlapped above chance with most clusters on subsequent days in

both AB and TU fish, with the overlap in TU fish averaging about 10% higher than AB fish

(Figure 5C). In AB fish, cluster overlap after day 2 increased markedly (range 62-70%). In TU

fish, an increase in overlapping clusters did not occur until day 3, where it also remained high

(63-80%). We saw a similar trend during biweekly exposures (Figure 5D) where cluster identity

on the first day overlapped the least with subsequent weeks (33-44%), but from the second to

the tenth week, overlap was considerably higher (60-80%).

Discussion

By applying an unbiased approach to the clustering of three-dimensional exploratory

behavior from over 400 zebrafish, we found that behavior stratifies into four distinct behavioral

clusters. These profiles included previously described bold and shy behaviors as well as two

novel behavioral types we call wall-huggers and active explorers. Notably, these individual

differences in fish behavior were consistent over days and weeks, one of the key hallmarks of

personality. Although there were few strain-sex interactions on individual behaviors, the

distribution of clusters varied considerably across strain and sex suggesting biological

modulation of these behavioral clusters.

Studies that have examined individual differences in exploratory behavior typically

assume a bimodal distribution of bold versus shy. Here, enabled by our large dataset of three-

dimensional behavior, we tested this assumption. Using an unbiased unsupervised machine

learning approach, we found four distinct behavioral subtypes (shy, wall-huggers, active

explorers, and bold). Cluster 4 was the boldest cluster where fish behavior corresponded to

traditional descriptions of boldness: above average time in the top and center of the tank and

greater exploration. The behavior of fish in cluster 1 was designated as shy because these fish

spent most of their time near the bottom of the tank. Interestingly, in this shy group, behaviors

like center distance and percent explored exhibited a high degree of variability, suggesting the

potential presence of additional subgroups that would require an even larger data set to

uncover.

Of the novel behavioral clusters we uncovered, we designated cluster 2 'wall-huggers'

because these fish spent most of their time towards the periphery of the tank and were near

average on all other parameters. We do not refer to this group as 'shy' because, although

thigmotaxis has been interpreted as a predator avoidance or anxiety-like behavior in zebrafish

(Kalueff et al., 2013), our findings do not clearly support this interpretation. For example, we did

not observe consistent negative correlations between center and bottom distance (Figure 2E),

as would be expected if these behaviors reflected the same underlying construct (i.e. predator

avoidance). We also find that, unlike bottom dwelling, thigmotaxis increased over time within

and between sessions of the novel tank test (Figures S2B and S6B), the opposite of the

habituation we would expect to see if thigmotaxis was an anxiety-like or predator avoidance

behavior. Our findings are consistent with other studies that examined thigmotaxis over time

and/or alongside other avoidance behaviors in adult fish (Blaser et al., 2010; Champagne et al.,

2010; Rosa et al., 2018; Shams et al., 2015; Wong et al., 2012). It may be that the interpretation

of thigmotaxis in zebrafish has been unduly influenced by findings in rodents where its

importance for predator avoidance is clearer (Champagne et al., 2010; Simon et al., 1994).

Thus, we propose that thigmotaxis in adult zebrafish should not be interpreted in the context of

predator avoidance or anxiety until its relevance for fish can be clarified.

We designated behavioral cluster 3 as 'active explorers' because these fish were clearly

above average in both distance travelled and percent explored, but near average on center and

bottom distance. This group of fish may correspond to what has been referred to as low

stationary, or proactive, zebrafish in other work (Baker et al., 2018; Wong et al., 2012). In

comparison to the boldest cluster, active explorers are distinguished by a notable elevation in

distance travelled with similar levels of percent exploration. Thus, during exploration, these

animals are less efficient in their exploration as they revisit many parts of the tank. This

suggests that these animals may have poor working memory or higher resting metabolic rates,

the latter having been found to be predictive of individual differences in activity (Biro and

Stamps, 2010; Yuan et al., 2018).

Strain and sex had a significant influence on behavioral cluster identity. During the initial

exposure to the tank, female fish more often fell into the boldest cluster, although only TU

females were statistically overrepresented. Male fish were more likely to be in the active

explorer group, particularly TU and WIK male fish. This likely reflects the fact that, on average,

male fish tended to swim further, and explore more of the tank, than female fish. This increase

in locomotor activity in male fish was expected as it has been reported elsewhere (Ariyomo and

Watt, 2015; Clayman et al., 2017; Philpott et al., 2012), although not universally so (Ampatzis

and Dermon, 2016; Fontana et al., 2019; Rambo et al., 2017; Tran and Gerlai, 2013). We also

found interactions between sex and strain on cluster identity, such as underrepresentation of AB

females, and overrepresentation of TL females, in the shyest cluster with the situation reversed

in the wall-huggers (cluster 2) group; notably, this interaction persisted into the second day of

testing, suggesting it is particularly robust.

413   The behavioral clusters we identified demonstrated a high degree of consistency across

days and weeks, supporting the idea that they are akin to personality types (Gosling, 2001). In

our initial experiment, where fish were tested in the novel tank on two consecutive days, overlap

was above 50%, twice as high as would be expected by chance (~25%). Upon repeated

exposures to the tank over five consecutive days we found that cluster overlap steadily

increased to above 65%, and over half of the fish were in the same cluster on 4 of the 5 days.

We found similar effects during biweekly (every other week) exposures over 10 weeks. This

increase in overlap over days is likely due, in part, to habituation (Wong et al., 2010) given that

we found that there were main effects of day on several individual measures, particularly in TU

fish (Figures S6 and S7). Nonetheless, this is the first report, to our knowledge, of consistency

in zebrafish behavior across days and weeks using unbiased multidimensional behavioral

clustering that approximates key elements of personality.

425   The present work also yielded a comprehensive account of how genetic background

influences individual exploratory behaviors of zebrafish, adding to an already rich, if

inconsistent, literature. The primary strain difference on individual behaviors we observed was

that TL fish differed from most other strains, demonstrating higher bottom dwelling, less

thigmotaxis, and exploring less of the tank. Although strain differences in zebrafish behavior

during the novel tank test have been reported (Egan et al., 2009; Maximino et al., 2013; Mustafa

et al., 2019), only a few studies have compared widely used inbred strains. Vignet and

colleagues (2013) found that AB fish exhibited more bottom dwelling than TU fish, and Audira et

al (2020) found no differences between AB, TL, and WIK fish in bottom dwelling or distance

travelled, but that WIK fish exhibited greater thigmotaxis. Others have found AB fish to exhibit

greater bottom dwelling than WIKs (Sackerman et al., 2010), or a decrease in bottom dwelling in

WIK fish over 60 minutes, but not TU fish, with no difference in locomotor activity (Pannia et al.,

2014). Differences between these studies, and the present work, may be attributable to

unreported sex ratios, housing conditions, nuances of the behavioral task, or genetic drift such

that inbred strains from different labs or suppliers may differ subtly, as has been observed in

rodents (Taft et al., 2006). We attempted to address some of these challenges in our study. For

example, housing conditions are known to influence zebrafish behavior in the novel tank test

(Parker et al., 2012; Reolon et al., 2018). We housed all animals in mixed sex pairs for one

week prior to behavioral testing. This allowed us to maintain zebrafish identify over time while

avoiding any effects of stress due to social isolation or tagging. To minimize genetic drift, and

maximize the likelihood our findings will translate to other labs, all fish were within 2 generations

of breeders obtained from the Zebrafish International Resource Center where they maintain a

much larger, and genetically diverse, population of animals. Nonetheless, given that inbred

zebrafish lines are not isogenic (Nasiadka and Clark, 2012), there is no obvious way to ensure

genetic similarity of lines across labs.

Taken together, we have found that the exploratory behavior of zebrafish goes beyond

bold versus shy, stratifying instead into four distinct categories. This finding was enabled by

recent advances in animal tracking and 3D video capture that allowed us to scale up our

behavioral assessment using inexpensive open-source tools. As would be expected of

behaviors capturing personality, the clusters we identified are consistent over days and weeks

and influenced by strain and sex. Future work will be needed to determine if these clusters are

predictive of behaviors in other contexts. Nonetheless, our findings suggests that zebrafish

behavior is more complex than is typically assumed and that this complexity needs to be taken

into consideration when using zebrafish to study the biological basis for individual differences in

behavior.

475   **Acknowledgments**

476   We thank Barbara D. Fontana for excellent comments on a prior version of this manuscript. This

477   work was funded by the National Institutes of Health (R35GM142566) to JWK.

Methods

Subjects

Subjects were female and male AB, TU, WIK, or TL zebrafish 16-32 weeks of age. All

fish used in experiments were bred and raised at Wayne State University and within two

generations of animals obtained from the Zebrafish International Resource Center at the

University of Oregon. Animals were kept on high density racks under standard conditions

(temperature 26.5 ± 0.5 C; water conductivity 500 ± 10 μS, and a pH of 7.5 ± 0.2) with a 14:10

light/dark cycle (lights on at 8:00AM). Fish were fed twice a day with a dry feed in the morning

and brine shrimp (*Artemia salina*; Brine Shrimp Direct, Ogden, UT) in the afternoon. Behavioral

testing took place between 11:00 and 14:00. Sex of fish was confirmed via dissection after

behavioral procedures were complete. Those animals that were assigned the wrong sex were

removed from analysis. All procedures were approved by the Wayne State University

Institutional Animal Care and Use Committee.

Behavioral apparatus

Five-sided novel tanks (15 x 15 x 15 cm) were made from frosted acrylic

(ShopPopDisplays, Woodland Park, NJ) and open from above. Tanks were placed in an

enclosure made of white plasticore to diffuse light and prevent fish from seeing experimenters.

Intel RealSense^{TM} cameras (D435) were mounted 20 cm above the tank to capture three-

dimensional videos (Kuroda, 2018). Video capture was done using custom written Python

scripts and the pyrealsense2 package.

Behavioral procedures

One week prior to behavioral testing, fish were placed as male/female pairs into 2 L

tanks. The tanks were divided in half with a transparent divider with two fish in each section and

a total of four fish in each tank. This allowed us to maintain the identity of fish over days without

isolation while also creating a consistent social environment across all animals. On days when

behavior was assessed, fish were taken off housing racks and moved to the procedural space

at least one hour prior to behavioral testing. Following testing, fish sat for one hour before being

returned to the housing racks. The novel tanks were filled with 2.5 L of fish facility water and

individual fish were placed in the tanks for six minutes while video was recorded for offline

analysis. Novel tanks were rinsed between animals and water was replaced.

Animal Tracking

Fish were tracked in the color videos using DeepLabCut (Mathis et al., 2018). We

tracked five points (head, trunk, and three points on the tail; Figure 1C). Using ResNet 101, we

initially trained the network on 160 frames equally divided across fish of all four strains and both

sexes. We refined and improved our initial training by correcting outliers and including an

additional 160 frames.

To obtain the z-coordinate of fish at each frame, we overlaid the tracked points at each

color frame with the depth stream from the cameras. One z-coordinate for the fish was identified

for each frame based on a 4-pixel search radius around the tracked points starting with the

trunk. If no point was identified from the depth stream, it was interpolated. Z-coordinates were

corrected for the diffraction of water by measuring 100 points of varying distance from the

camera in the presence and absence of water and using the slope of a least-squares fit line.

Exploratory Behavioral Parameters

To measure bottom distance, we calculated the equation for a plane along the bottom of

the tank using least squares fit of four points. We then calculated the shortest distance between

a point (the fish) and the plane. To measure center distance, we calculated how far fish were

from a line made from points at the center top and bottom of the tank. Percent of the tank

explored was calculated by dividing the tank into 1,000 evenly spaced voxels and calculating

the number of unique voxels visited. For distance travelled, traces were first smoothed using a

Savitzky-Golay filter with a length of 7 frames and an order of 3 (Press and Teukolsky, 1990)

and then Euclidean distance was calculated between points of successive frames.

Behavioral clusters

We identified behavioral clusters using a Louvain community detection algorithm

(Blondel et al., 2008) applied to a k-nearest neighbor network on our initial dataset of 426 fish

during their first day of exposure to the novel tank. This was done by first standardizing

individual behavioral parameters (bottom distance, center distance, distance travelled, and

percent explored), and calculating a similarity score between each fish:

$$Similarity\ score = \frac{1}{1 + D}$$

Where D is the Euclidean distance between each fish in four-dimensional behavioral space. To

determine the best k for building the network, we initially used a range of k's, applied the

Louvain community finding algorithm to weighted, non-directed networks, and calculated

internal clustering metrics (Calinski-Harabsz index (Caliński and Harabasz, 1974), Silhouette

index (Rousseeuw, 1987), and Davies-Bouldin index (Davies and Bouldin, 1979)). We chose

k=118, which was in the middle of a regime that optimized internal clustering and was robust to

small changes in k (Figure S4).

To identify clusters in new data (e.g. the second day of novel tank exposure (Figure 4)

fish exposed on consecutive days or biweekly (Figure 5)), we first standardized the new data

using the parameters from the 426 fish exposed to the novel tank described above. Then, for

each new data point, we assigned clusters based on the proportion of connections to its 51

nearest neighbors in the initial network where 51 is half the size of the smallest cluster

identified.

Coding and statistical analysis

Statistical analysis was performed using R version 4.1.2 (R Core Team, 2016) and

visualized using ggplot2 (Wickham, 2015) and RColorBrewer (Neuwirth and Brewer, 2014).

Normality was assessed using the Shapiro-Wilks test. Because a considerable portion of the

data was not normally distributed, we used permutation ANOVAs using the permuco package

(Frossard and Renaud, 2021), and the RVAideMemoire (Hervé and Hervé, 2020) for

permutation t-tests. Multiple comparisons were corrected using a false discover rate (FDR)

correction (Benjamini and Hochberg, 1995). We used packages cccd (Marchette et al., 2015)

and igraph (Csardi and Nepusz, 2006) to build and analyze the k-nearest neighbor network, and

the ClusterCrit (Desgraupes, 2013) to assess internal clustering metrics. The chord diagram

was made using circlize (Gu et al., 2014).

For all permutation tests, we resampled data 10,000 times without replacement and

calculated the P-value as the proportion of experimental observations that were more extreme

than the permutation observations. When multiple tests were performed, P-values were

corrected using the FDR as indicated. To determine if behavioral clusters were affected by

strain and sex (Figure 3C and 4B), we resampled cluster assignment and calculated how many

animals from each strain and sex fell into each cluster. For comparison of behavioral clusters

across two days (Figure 4B) we resampled cluster assignments from day 2. To determine if

behavior was consistent across daily (five day) or biweekly (every other week over ten weeks)

(Figure 5A-B), we first calculated an overlap score for each fish that was the sum of the number

of days or weeks that had overlapping clusters (ranging from 1 to 10 for five-day data, 2 to 15

for biweekly data). We then took the average of these overlap scores and compared them to

scores obtained from permutation resampling (without replacement) of cluster assignments.

Five-day and biweekly percent overlap calculations and permutation tests were as described for

the two-day data except that resampling of cluster assignments was done individually for all

days/weeks except the fist day/week.

Figures

Figure 1. Overview of three-dimensional behavioral tracking. **A**) Individual fish were placed into

a novel tank while video was recorded from above using Intel RealSense™ cameras. **B**) Videos

included both a color (top) and a depth (bottom) stream where fish can be seen (yellow arrows).

C) Animals were tracked in the color videos using DeepLabCut to identify five points along the

length of the fish. **D**) Tracking was overlaid with the depth stream to generate a three-

dimensional trace for each animal. **E**) Four exploratory parameters were extracted from each

trace: bottom distance (top left), center distance (top right), distance travelled (bottom left), and

percent of the tank explored (bottom right).

Figure 2. Influence of sex and strain on individual exploratory behaviors and their relationships.

The effect of sex and strain on **A**) bottom distance, **B**) center distance, **C**) distance travelled,

and **D**) percent of the tank explored. Boxplots indicate median (center line), interquartile range

(box ends), and hinge ± 1.5 times the interquartile range (whiskers). Spearman's rank

correlation coefficient (ρ) with 95% confidence intervals across strain and sex for **E**) bottom

distance versus center distance, **F**) bottom distance versus distance travelled, **G**) bottom

distance versus percent explored, **H**) center distance versus distance travelled, **I**) center

distance versus percent explored, and **J**) distance travelled versus percent explored. * - P <

0.05, † - P < 0.10 compared to all other groups or those indicated, n = 50-58.

Figure 3. Clustering of zebrafish exploratory behavior during exposure to a novel tank. **A**) Two-

dimensional representation of the four-dimensional behavioral space using a uniform manifold

approximation (McInnes et al., 2020). Clusters (outer circles) are derived from Louvain

community finding applied to a k-nearest neighbor network using 426 fish. Each point is an

individual fish where the shape represents the sex (circle: female, triangle: male), inner color the

strain, and outer color the behavioral cluster. **B**) Individual behaviors associated with each

cluster as box plots indicating median (center line), interquartile range (box ends), and hinge ±

1.5 times the interquartile range (whiskers). **C**) Percentage of fish that fall into each cluster

across strain and sex. Striped bars: P < 0.05 using randomized permutation tests and FDR

corrections, n = 50-58.

657   *Figure 4*. Clusters across two consecutive exposures to the novel tank. **A**) Chord diagram

658   indicating how cluster membership changes from day 1 to day 2 of novel tank exposure. **B**)

659   Percentage of fish that fall into each cluster across strain and sex on the second day of

660   exposure to the novel tank. Striped bars: P < 0.05 using randomized permutation tests and FDR

661   corrections, n = 50-58.

Figure 5. Behavioral consistency in the novel tank across days and weeks. **A**) Cluster

membership of AB (n = 46) or TU (n = 50) fish that were exposed to the novel tank on five

consecutive days. **B**) Cluster membership of TU (n = 45) fish that were exposed to the novel

tank every other week for 10 weeks. **C**) Percent of overlapping clusters in AB and TU fish

across consecutive exposures to the novel tank across five days. **D**) Percent overlapping

clusters in TU fish across biweekly (every other week) exposures to the novel tank. * - P < 0.05

using randomized permutation test and FDR corrections.

References

Ampatzis K, Dermon CR. 2016. Sexual dimorphisms in swimming behavior, cerebral metabolic activity and adrenoceptors in adult zebrafish (Danio rerio). *Behavioural Brain Research* **312**:385–393. doi:10.1016/j.bbr.2016.06.047

Ariyomo TO, Watt PJ. 2015. Effect of hunger level and time of day on boldness and aggression in the zebrafish Danio rerio. *Journal of Fish Biology* **86**:1852–1859. doi:10.1111/jfb.12674

Audira G, Siregar P, Strungaru S-A, Huang J-C, Hsiao C-D. 2020. Which Zebrafish Strains Are More Suitable to Perform Behavioral Studies? A Comprehensive Comparison by Phenomic Approach. *Biology* **9**:200. doi:10.3390/biology9080200

Baker MR, Goodman AC, Santo JB, Wong RY. 2018. Repeatability and reliability of exploratory behavior in proactive and reactive zebrafish, Danio rerio. *Sci Rep* **8**:12114. doi:10.1038/s41598-018-30630-3

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**:289–300. doi:10.1111/j.2517-6161.1995.tb02031.x

Biro PA, Stamps JA. 2010. Do consistent individual differences in metabolic rate promote consistent individual differences in behavior? *Trends in Ecology & Evolution* **25**:653–659. doi:10.1016/j.tree.2010.08.003

Blaser RE, Chadwick L, McGinnis GC. 2010. Behavioral measures of anxiety in zebrafish (Danio rerio). *Behavioural Brain Research* **208**:56–62. doi:10.1016/j.bbr.2009.11.009

Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E. 2008. Fast unfolding of communities in large networks. *Journal of statistical mechanics: theory and experiment* **2008**:P10008.

Caliński T, Harabasz J. 1974. A dendrite method for cluster analysis. *Communications in Statistics-theory and Methods* **3**:1–27.

Carere C, Drent PJ, Privitera L, Koolhaas JM, Groothuis TGG. 2005. Personalities in great tits, Parus major: stability and consistency. *Animal Behaviour* **70**:795–805. doi:10.1016/j.anbehav.2005.01.003

Champagne DL, Hoefnagels CCM, de Kloet RE, Richardson MK. 2010. Translating rodent behavioral repertoire to zebrafish (Danio rerio): Relevance for stress research. *Behavioural Brain Research* **214**:332–342. doi:10.1016/j.bbr.2010.06.001

Clayman CL, Malloy EJ, Kearns DN, Connaughton VP. 2017. Differential behavioral effects of ethanol pre-exposure in male and female zebrafish (Danio rerio). *Behavioural Brain Research* **335**:174–184. doi:10.1016/j.bbr.2017.08.007

Csardi G, Nepusz T. 2006. The igraph software package for complex network research. *InterJournal, Complex Systems* **1695**:1–9.

Dahlbom SJ, Lagman D, Lundstedt-Enkel K, Sundström LF, Winberg S. 2011. Boldness Predicts Social Status in Zebrafish (Danio rerio). *PLOS ONE* **6**:e23565. doi:10.1371/journal.pone.0023565

Dall SRX, Bell AM, Bolnick DI, Ratnieks FLW. 2012. An evolutionary ecology of individual differences. *Ecology Letters* **15**:1189–1198. doi:10.1111/j.1461-0248.2012.01846.x

Davies DL, Bouldin DW. 1979. A cluster separation measure. *IEEE transactions on pattern analysis and machine intelligence* 224–227.

Desgraupes B. 2013. Clustering indices. *University of Paris Ouest-Lab Modal'X* **1**:34.

Egan RJ, Bergner CL, Hart PC, Cachat JM, Canavello PR, Elegante MF, Elkhayat SI, Bartels BK, Tien AK, Tien DH, Mohnot S, Beeson E, Glasgow E, Amri H, Zukowska Z, Kalueff AV. 2009. Understanding behavioral and physiological phenotypes of stress and anxiety in zebrafish. *Behavioural Brain Research* **205**:38–44. doi:10.1016/j.bbr.2009.06.022

720   Fontana BD, Cleal M, Parker MO. 2019. Female adult zebrafish (*Danio rerio*) show higher levels of
721   anxiety-like behavior than males, but do not differ in learning and memory capacity. *European*
722   *Journal of Neuroscience* ejn.14588. doi:10.1111/ejn.14588
723   Forkosh O, Karamihalev S, Roeh S, Alon U, Anpilov S, Touma C, Nussbaumer M, Flachskamm C, Kaplick
724   PM, Shemesh Y, Chen A. 2019. Identity domains capture individual differences from across the
725   behavioral repertoire. *Nat Neurosci* **22**:2023–2028. doi:10.1038/s41593-019-0516-y
726   Frossard J, Renaud O. 2021. Permutation tests for regression, ANOVA, and comparison of signals: the
727   permuco package. *Journal of Statistical Software* **99**:1–32.
728   Gerlai R. 2020. Evolutionary conservation, translational relevance and cognitive function: The future of
729   zebrafish in behavioral neuroscience. *Neuroscience & Biobehavioral Reviews* **116**:426–435.
730   doi:10.1016/j.neubiorev.2020.07.009
731   Gosling SD. 2001. From mice to men: What can we learn about personality from animal research?
732   *Psychological Bulletin* **127**:45–86. doi:10.1037/0033-2909.127.1.45
733   Grant M. 2000. Galen on Food and Diet. Florence, UNITED STATES: Taylor & Francis Group.
734   Gu Z, Gu L, Eils R, Schlesner M, Brors B. 2014. Circlize implements and enhances circular visualization in
735   R. *Bioinformatics* **30**:2811–2812.
736   Hervé M, Hervé MM. 2020. Package 'RVAideMemoire.' *See https://CRANR-projectorg/package=*
737   *RVAideMemoire*.
738   Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, Collins JE, Humphray S, McLaren K,
739   Matthews L, McLaren S, Sealy I, Caccamo M, Churcher C, Scott C, Barrett JC, Koch R, Rauch GJ,
740   White S, Chow W, Kilian B, Quintais LT, Guerra-Assunção JA, Zhou Y, Gu Y, Yen J, Vogel JH, Eyre
741   T, Redmond S, Banerjee R, Chi J, Fu B, Langley E, Maguire SF, Laird GK, Lloyd D, Kenyon E,
742   Donaldson S, Sehra H, Almeida-King J, Loveland J, Trevanion S, Jones M, Quail M, Willey D, Hunt
743   A, Burton J, Sims S, McLay K, Plumb B, Davis J, Clee C, Oliver K, Clark R, Riddle C, Eliott D,
744   Threadgold G, Harden G, Ware D, Mortimer B, Kerry G, Heath P, Phillimore B, Tracey A, Corby N,
745   Dunn M, Johnson C, Wood J, Clark S, Pelan S, Griffiths G, Smith M, Glithero R, Howden P, Barker
746   N, Stevens C, Harley J, Holt K, Panagiotidis G, Lovell J, Beasley H, Henderson C, Gordon D, Auger
747   K, Wright D, Collins J, Raisen C, Dyer L, Leung K, Robertson L, Ambridge K, Leongamornlert D,
748   McGuire S, Gilderthorp R, Griffiths C, Manthravadi D, Nichol S, Barker G, Whitehead S, Kay M,
749   Brown J, Murnane C, Gray E, Humphries M, Sycamore N, Barker D, Saunders D, Wallis J, Babbage
750   A, Hammond S, Mashreghi-Mohammadi M, Barr L, Martin S, Wray P, Ellington A, Matthews N,
751   Ellwood M, Woodmansey R, Clark G, Cooper J, Tromans A, Grafham D, Skuce C, Pandian R,
752   Andrews R, Harrison E, Kimberley A, Garnett J, Fosker N, Hall R, Garner P, Kelly D, Bird C, Palmer
753   S, Gehring I, Berger A, Dooley CM, Ersan-Ürün Z, Eser C, Geiger H, Geisler M, Karotki L, Kirn A,
754   Konantz J, Konantz M, Oberländer M, Rudolph-Geiger S, Teucke M, Osoegawa K, Zhu B, Rapp A,
755   Widaa S, Langford C, Yang F, Carter NP, Harrow J, Ning Z, Herrero J, Searle SMJ, Enright A,
756   Geisler R, Plasterk RHA, Lee C, Westerfield M, De Jong PJ, Zon LI, Postlethwait JH, Nüsslein-
757   Volhard C, Hubbard TJP, Crollius HR, Rogers J, Stemple DL. 2013. The zebrafish reference
758   genome sequence and its relationship to the human genome. *Nature* **496**:498–503.
759   doi:10.1038/nature12111
760   Kalueff AV, Gebhardt M, Stewart AM, Cachat JM, Brimmer M, Chawla JS, Craddock C, Kyzar EJ, Roth A,
761   Landsman S, Gaikwad S, Robinson K, Baatrup E, Tierney K, Shamchuk A, Norton W, Miller N,
762   Nicolson T, Braubach O, Gilman CP, Pittman J, Rosemberg DB, Gerlai R, Echevarria D, Lamb E,
763   Neuhauss SCFF, Weng W, Bally-Cuif L, Schneider H, Schneider, and the Zebrafish Neuros H. 2013.
764   Towards a comprehensive catalog of zebrafish behavior 1.0 and beyond. *Zebrafish* **10**:70–86.
765   doi:10.1089/zeb.2012.0861
766   Kenney JW. 2020. Associative and nonassociative learning in adult zebrafishBehavioral and Neural
767   Genetics of Zebrafish. Elsevier. pp. 187–204. doi:10.1016/b978-0-12-817528-6.00012-7

768   Kenney JW, Steadman PE, Young O, Shi MT, Polanco M, Dubaishi S, Covert K, Mueller T, Frankland PW.
769   2021. A 3D adult zebrafish brain atlas (AZBA) for the digital age. *eLife* **10**:e69988.
770   doi:10.7554/eLife.69988
771   Kuroda T. 2018. A system for the real-time tracking of operant behavior as an application of 3D camera.
772   *Journal of the Experimental Analysis of Behavior* **110**:522–544. doi:10.1002/jeab.471
773   López P, Hawlena D, Polo V, Amo L, Martín J. 2005. Sources of individual shy–bold variations in
774   antipredator behaviour of male Iberian rock lizards. *Animal Behaviour* **69**:1–9.
775   doi:10.1016/j.anbehav.2004.05.010
776   Marchette DJ, Marchette MDJ, Matrix S. 2015. Package 'cccd.'
777   Martins EP, Bhat A. 2014. Population-level personalities in zebrafish: aggression-boldness across but not
778   within populations. *Behavioral Ecology* **25**:368–373. doi:10.1093/beheco/aru007
779   Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M. 2018. DeepLabCut:
780   markerless pose estimation of user-defined body parts with deep learning. *Nature Neuroscience*
781   **21**:1281–1289. doi:10.1038/s41593-018-0209-y
782   Maximino C, Puty B, Matos Oliveira KR, Herculano AM. 2013. Behavioral and neurochemical changes in
783   the zebrafish leopard strain. *Genes, Brain and Behavior* **12**:576–582. doi:10.1111/gbb.12047
784   McInnes L, Healy J, Melville J. 2020. UMAP: Uniform Manifold Approximation and Projection for
785   Dimension Reduction. *arXiv:180203426 [cs, stat]*.
786   Mustafa A, Roman E, Winberg S. 2019. Boldness in Male and Female Zebrafish (Danio rerio) Is
787   Dependent on Strain and Test. *Frontiers in Behavioral Neuroscience* **13**:248.
788   doi:10.3389/fnbeh.2019.00248
789   Myers PJ, Young JK. 2018. Consistent individual behavior: evidence of personality in black bears. *J Ethol*
790   **36**:117–124. doi:10.1007/s10164-018-0541-4
791   Nasiadka A, Clark MD. 2012. Zebrafish Breeding in the Laboratory Environment. *ILAR Journal* **53**:161–
792   168. doi:10.1093/ilar.53.2.161
793   Neuwirth E, Brewer RC. 2014. ColorBrewer palettes. *R package version* **1**.
794   Oswald ME, Drew RE, Racine M, Murdoch GK, Robison BD. 2012. Is Behavioral Variation along the Bold-
795   Shy Continuum Associated with Variation in the Stress Axis in Zebrafish? *Physiological and*
796   *Biochemical Zoology* **85**:718–728. doi:10.1086/668203
797   Pannia E, Tran S, Rampersad M, Gerlai R. 2014. Acute ethanol exposure induces behavioural differences
798   in two zebrafish (Danio rerio) strains: A time course analysis. *Behavioural Brain Research*
799   **259**:174–185. doi:10.1016/j.bbr.2013.11.006
800   Panula P, Chen YC, Priyadarshini M, Kudo H, Semenova S, Sundvik M, Sallinen V. 2010. The comparative
801   neuroanatomy and neurochemistry of zebrafish CNS systems of relevance to human
802   neuropsychiatric diseases. *Neurobiology of Disease* **40**:46–57. doi:10.1016/j.nbd.2010.05.010
803   Parker MO, Millington ME, Combe FJ, Brennan CH. 2012. Housing Conditions Differentially Affect
804   Physiological and Behavioural Stress Responses of Zebrafish, as well as the Response to
805   Anxiolytics. *PLOS ONE* **7**:e34992. doi:10.1371/journal.pone.0034992
806   Philpott C, Donack CJ, Cousin MA, Pierret C. 2012. Reducing the noise in behavioral assays: sex and age
807   in adult zebrafish locomotion. *Zebrafish* **9**:191–194.
808   Press WH, Teukolsky SA. 1990. Savitzky-Golay smoothing filters. *Computers in Physics* **4**:669–672.
809   R Core Team. 2016. R: A Language and Environment for Statistical Computing. *R Foundation for*
810   *Statistical Computing* **1**:409. doi:10.1007/978-3-540-74686-7
811   Rambo CL, Mocelin R, Marcon M, Villanova D, Koakoski G, de Abreu MS, Oliveira TA, Barcellos LJG, Piato
812   AL, Bonan CD. 2017. Gender differences in aggression and cortisol levels in zebrafish subjected
813   to unpredictable chronic stress. *Physiology & Behavior* **171**:50–54.
814   doi:10.1016/j.physbeh.2016.12.032

815   Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ. 2007. Integrating animal temperament within
816   ecology and evolution. *Biological Reviews* **82**:291–318. doi:10.1111/j.1469-185X.2007.00010.x
817   Reolon GK, de Melo GM, da Rosa JG dos S, Barcellos LJG, Bonan CD. 2018. Sex and the housing: Effects
818   on behavior, cortisol levels and weight in zebrafish. *Behavioural Brain Research* **336**:85–92.
819   doi:10.1016/j.bbr.2017.08.006
820   Rosa LV, Ardais AP, Costa FV, Fontana BD, Quadros VA, Porciúncula LO, Rosemberg DB. 2018. Different
821   effects of caffeine on behavioral neurophenotypes of two zebrafish populations. *Pharmacology*
822   *Biochemistry and Behavior* **165**:1–8. doi:10.1016/j.pbb.2017.12.002
823   Rousseeuw PJ. 1987. Silhouettes: A graphical aid to the interpretation and validation of cluster analysis.
824   *Journal of Computational and Applied Mathematics* **20**:53–65. doi:10.1016/0377-
825   0427(87)90125-7
826   Sackerman J, Donegan JJ, Cunningham CS, Nguyen NN, Lawless K, Long A, Benno RH, Gould GG. 2010.
827   Zebrafish Behavior in Novel Environments: Effects of Acute Exposure to Anxiolytic Compounds
828   and Choice of Danio rerio Line. *International journal of comparative psychology* **23**:43–61.
829   Shams S, Chatterjee D, Gerlai R. 2015. Chronic social isolation affects thigmotaxis and whole-brain
830   serotonin levels in adult zebrafish. *Behavioural Brain Research* **292**:283–287.
831   doi:10.1016/j.bbr.2015.05.061
832   Sih A, Bell A, Johnson JC. 2004. Behavioral syndromes: an ecological and evolutionary overview. *Trends*
833   *in Ecology & Evolution* **19**:372–378. doi:10.1016/j.tree.2004.04.009
834   Simon P, Dupuis R, Costentin J. 1994. Thigmotaxis as an index of anxiety in mice. Influence of
835   dopaminergic transmissions. *Behavioural Brain Research* **61**:59–64. doi:10.1016/0166-
836   4328(94)90008-6
837   Taft RA, Davisson M, Wiles MV. 2006. Know thy mouse. *Trends in Genetics* **22**:649–653.
838   doi:10.1016/j.tig.2006.09.010
839   Thörnqvist P-O, McCarrick S, Ericsson M, Roman E, Winberg S. 2019. Bold zebrafish (Danio rerio) express
840   higher levels of delta opioid and dopamine D2 receptors in the brain compared to shy fish.
841   *Behavioural Brain Research* **359**:927–934. doi:10.1016/j.bbr.2018.06.017
842   Toms CN, Echevarria DJ, Jouandot DJ. 2010. A Methodological Review of Personality-Related Studies in
843   Fish: Focus on the Shy-Bold Axis of Behavior. *International Journal of Comparative Psychology*
844   **23**.
845   Tran S, Gerlai R. 2013. Individual differences in activity levels in zebrafish (Danio rerio). *Behavioural*
846   *Brain Research* **257**:224–229. doi:10.1016/j.bbr.2013.09.040
847   Vignet C, Bégout M-L, Péan S, Lyphout L, Leguay D, Cousin X. 2013. Systematic Screening of Behavioral
848   Responses in Two Zebrafish Strains. *Zebrafish* **10**:365–375. doi:10.1089/zeb.2013.0871
849   Volgin AD, Yakovlev OA, Demin KA, de Abreu MS, Alekseeva PA, Friend AJ, Lakstygal AM, Amstislavskaya
850   TG, Bao W, Song C, Kalueff AV. 2019. Zebrafish models for personalized psychiatry: Insights from
851   individual, strain and sex differences, and modeling gene x environment interactions. *Journal of*
852   *Neuroscience Research* **97**:402–413. doi:10.1002/jnr.24337
853   Wickham H. 2015. Elegant Graphics for Data Analysis, Media. Springer. doi:10.1007/978-0-387-98141-3
854   Wilson DS, Clark AB, Coleman K, Dearstyne T. 1994. Shyness and boldness in humans and other animals.
855   *Trends in Ecology & Evolution* **9**:442–446. doi:10.1016/0169-5347(94)90134-1
856   Wong K, Elegante M, Bartels B, Elkhayat S, Tien D, Roy S, Goodspeed J, Suciu C, Tan J, Grimes C, Chung A,
857   Rosenberg M, Gaikwad S, Denmark A, Jackson A, Kadri F, Chung KM, Stewart A, Gilder T, Beeson
858   E, Zapolsky I, Wu N, Cachat J, Kalueff AV. 2010. Analyzing habituation responses to novelty in
859   zebrafish (Danio rerio). *Behavioural Brain Research* **208**:450–457. doi:10.1016/j.bbr.2009.12.023
860   Wong RY, Perrin F, Oxendine SE, Kezios ZD, Sawyer S, Zhou L, Dereje S, Godwin J. 2012. Comparing
861   behavioral responses across multiple assays of stress and anxiety in zebrafish (Danio rerio).
862   *Behaviour* **149**:1205–1240.

863   Wulliman MF, Rupp B, Reichert H, Wullimann MF, Rupp B, Reichert H. 1996. Neuroanatomy of the
864   Zebrafish Brain: A Topological Atlas, Neuroanatomy of the Zebrafish Brain. Birkhäuser Verlag,
865   Basel. doi:10.1007/978-3-0348-8979-7
866   Yuan M, Chen Y, Huang Y, Lu W. 2018. Behavioral and Metabolic Phenotype Indicate Personality in
867   Zebrafish (Danio rerio). *Frontiers in Physiology* **9**:653. doi:10.3389/fphys.2018.00653

886   **Supplemental Information**

888   *Figure S1*. Distribution of values for individual behavioral parameters across stain and sex.

889   Distributions for **A**) bottom distance, **B**) center distance, **C**) distance travelled, and **D**) percent

890   tank explored. * - P < 0.05, † - P < 0.10 Shapiro-Wilks test for normality, n = 50-58.

Figure S2. Individual exploratory behaviors across six minutes during a single exposure to the novel tank. The effect of strain (color) and sex (shape, circle: female, triangle: male) across time for **A**) bottom distance, **B**) center distance, and **C**) distance travelled. * - P < 0.05, † - P < 0.10 for time interval (minute) or sex as indicated, n = 50-58. Data presented as mean ± SEM.

Figure S3. Relationship between pair-wise individual behavioral parameters. Scatterplots across strain and sex of **A**) bottom distance versus center distance, **B**) bottom distance versus distance travelled, **C**) bottom distance versus percent explored, **D**) center distance versus distance

909   travelled, **E**) center distance versus percent explored, and **F**) distance travelled versus percent

910   explored. n = 50-58.

917   *Figure S4*. Internal clustering metrics for knn Louvain clustering. The parameter k was varied for

918   generating knn's and Louvain clustering followed by measuring the **A**) Calinski-Harbasz, **B**)

919   silhouette, and **C**) Davies-Bouldin indices at each k. Area highlighted in red indicates region

920   where clustering metrics were near optimal and robust to small changes in k.

Figure S5. Changes in individual exploratory behaviors during two consecutive exposures to the novel tank. The effect of strain and sex on changes (day 2 minus day 1) over two days in **A**) bottom distance, **B**) center distance, **C**) distance travelled, and **D**) percent explored. Boxplots indicate median (center line), interquartile range (box ends), and hinge ± 1.5 times the interquartile range (whiskers). * - P < 0.05, † - P < 0.10 compared to zero, n = 50-58.

Figure S6. Individual exploratory behaviors during five consecutive days of exposure to the novel tank. AB (top) and TU (bottom) fish were exposed to the novel tank on five consecutive days and **A**) bottom distance, **B**) center distance, **C**) distance travelled, and **D**) percent explored were measured. * - $P < 0.05$, † - $P < 0.10$ for day or sex and indicated, n = 21-26. Data presented as mean ± SEM.

950   *Figure S7.* Individual exploratory behaviors over ten weeks. TU fish were exposed to the novel

951   tank every other week and we measured **A**) bottom distance, **B**) center distance, **C**) distance

952   travelled, and **D**) percent explored. * - P < 0.05, † - P < 0.10 for week or sex as indicated, n =

953   22-23. Data presented as mean ± SEM.

Figure S8. Percent of fish falling into each cluster during repeated exposures to the novel tank.

A) Membership across five consecutive days for AB (left) and TU (right) fish. **B**) Cluster

membership across biweekly (every other week) exposures of TU fish.