High levels of cyclic diguanylate interfere with beneficial bacterial colonization

Ruth Y. Isenberg[1,2], David G. Christensen[3], Karen L. Visick[3], and Mark J. Mandel[1,2,*]

[1] Department of Medical Microbiology and Immunology, University of Wisconsin-Madison, Madison, WI USA

[2] Microbiology Doctoral Training Program, University of Wisconsin-Madison, Madison, WI USA

[3] Department of Microbiology and Immunology, Loyola University Stritch School of Medicine, Chicago, IL USA

Short title: c-di-GMP during *V. fischeri* colonization

Keywords: *Vibrio fischeri*, microbiome, *Euprymna scolopes*, squid, c-di-GMP, biofilm, flagellar motility

* Correspondence to:

Mark J. Mandel
University of Wisconsin-Madison
Department of Medical Microbiology and Immunology
1550 Linden Drive
Madison, WI 53706
Phone: (608) 261-1170
Fax: (608) 262-8418
Email: mmandel@wisc.edu

ABSTRACT

During colonization of the Hawaiian bobtail squid (*Euprymna scolopes*), *Vibrio fischeri* bacteria

undergo a lifestyle transition from a planktonic motile state in the environment to a biofilm state

in host mucus. Cyclic diguanylate (c-di-GMP) is a cytoplasmic signaling molecule that is

important for regulating motility-biofilm transitions in many bacterial species. *V. fischeri* encodes

50 proteins predicted to synthesize and/or degrade c-di-GMP, but a role for c-di-GMP regulation

during host colonization has not been investigated. We examined strains exhibiting either low or

high levels of c-di-GMP during squid colonization and found that while a low c-di-GMP strain

had no colonization defect, a high c-di-GMP strain was severely impaired. Expression of a

heterologous c-di-GMP phosphodiesterase restored colonization, demonstrating that the effect

is due to high c-di-GMP levels. In the constitutive high c-di-GMP state, colonizing *V. fischeri*

exhibited reduced motility, altered biofilm aggregate morphology, and a regulatory interaction

where transcription of one polysaccharide locus is inhibited by the presence of the other

polysaccharide. Our results highlight the importance of proper c-di-GMP regulation during

beneficial animal colonization, illustrate multiple pathways regulated by c-di-GMP in the host,

and uncover an interplay of multiple exopolysaccharide systems in host-associated aggregates.

IMPORTANCE

There is substantial interest in studying cyclic diguanylate (c-di-GMP) in pathogenic and

environmental bacteria, which has led to an accepted paradigm in which high c-di-GMP levels

promote biofilm formation and reduce motility. However, considerably less focus has been

placed on understanding how this compound contributes to beneficial colonization. Using the

Vibrio fischeri-Hawaiian bobtail squid study system, we took advantage of recent genetic

advances in the bacterium to modulate c-di-GMP levels and measure colonization and track c-

di-GMP phenotypes in a symbiotic interaction. Studies in the animal host revealed a c-di-GMP-

dependent genetic interaction between two distinct biofilm polysaccharides, Syp and cellulose,

that was not evident in culture-based studies: elevated c-di-GMP altered the composition and

abundance of the *in vivo* biofilm by decreasing *syp* transcription due to increased cellulose

synthesis. This study reveals important parallels between pathogenic and beneficial colonization

and additionally identifies c-di-GMP-dependent regulation that occurs specifically in the squid

host.

INTRODUCTION

In both pathogenic and beneficial associations, bacteria transition from a motile, planktonic state

in the environment to a surface-associated biofilm state within the host. Cyclic diguanylate (c-di-

GMP) is an intracellular signaling molecule that regulates this lifestyle transition for many

bacterial species, including *Vibrio cholerae*, *Pseudomonas aeruginosa*, *Caulobacter crescentus*,

and *Agrobacterium tumefaciens* (1, 2). Generally, higher intracellular c-di-GMP concentrations

lead to increased biofilm formation and decreased motility, while lower intracellular c-di-GMP

concentrations lead to the opposite phenotypes. C-di-GMP levels are regulated by diguanylate

cyclase (DGC) proteins, which synthesize c-di-GMP, and phosphodiesterase (PDE) proteins,

which degrade c-di-GMP. DGCs contain a GGDEF domain and PDEs contain either an EAL or

HD-GYP domain. While the effects of c-di-GMP on motility, biofilm formation, and infectivity of

pathogens such as *V. cholerae* have been extensively studied (3), the impact of c-di-GMP

during the establishment of beneficial bacterial symbioses is not well understood. The focus of

this study is to examine how c-di-GMP impacts beneficial colonization in a system where we

can manipulate intracellular levels of the compound and dissect the colonization process in

detail.

Vibrio fischeri is the exclusive, beneficial light organ symbiont of the Hawaiian bobtail squid

(*Euprymna scolopes*). *V. fischeri* undergoes the motility-to-biofilm lifestyle transition during the

initiation of host colonization (4–7). While *V. fischeri* is solitary and expresses polar flagella in

seawater, upon encountering the juvenile bobtail squid light organ the bacteria require the

symbiosis polysaccharide (Syp) to aggregate in the host mucus prior to commencing further

colonization (5, 8–11). The bacterial aggregates form on the outer face of the light organ

independent of flagellar motility and are influenced by physical currents generated by cilia on

the light organ surface (4, 12). Host-produced nitric oxide stimulates dispersal from the biofilm

aggregates (13), and the bacteria swim through the pore into the light organ ducts using

flagellar motility and chemotaxis toward squid-derived chitin oligosaccharides (8, 14–17). The

symbionts pass through an antechamber and bottleneck before reaching the crypts of the light

organ, where they reach high density and produce light to camouflage the shadow of the squid

host (18–20). The initiation of this process—including biofilm formation, chemotaxis, and

motility—occurs only when the squid is newly hatched, as colonizing bacteria seed the host for

its lifetime. Therefore, the mechanisms that regulate these processes are fundamental to

selection for the correct symbionts and the establishment of the association.

As with the pathogens noted above, *V. fischeri* (strain ES114) encodes a large number of

proteins predicted to regulate c-di-GMP levels: 50 total that encompass 28 DGCs, 14 PDEs, 5

DGC/PDEs, and 3 proteins that are predicted to be degenerate (21). Roles for DGCs MifA and

MifB, and PDEs BinA and PdeV in *V. fischeri* motility and/or biofilm formation have been

described (22–24). However, those studies did not examine individual mutants during squid

colonization, and a large-scale transposon-insertion sequencing study did not identify individual

mutations in genes related to c-di-GMP that were predicted to have significant phenotypes in

the squid host (25). Given the predicted complexity of the c-di-GMP network in *V. fischeri*, we

used new technology (26) to generate strains lacking multiple DGCs or PDEs to mimic either

low or high c-di-GMP levels, and we applied these strains to determine the effects of c-di-GMP

extremes on squid colonization. This approach allowed us to directly study the role of c-di-GMP

in vivo during beneficial colonization, to determine bacterial behaviors that are influenced by the

compound during animal colonization, and to reveal signaling interactions that are not evident

during growth in culture.

RESULTS

Deletion of multiple DGCs or PDEs effectively manipulates c-di-GMP levels in the

symbiont *V. fischeri*

To determine the role that c-di-GMP plays during the beneficial colonization of *E. scolopes* by *V.*

fischeri, we constructed strains that were altered in their basal levels of c-di-GMP. While it is not

yet feasible to delete all 50 c-di-GMP-related genes in a single strain, we applied recently

developed technology that facilitates rapid construction of multiple gene deletions (26) to

generate a strain that lacked seven DGCs (Δ7DGC; lacking *VF_1200*, *mifA*, *VF_1245*,

VF_A0342-A0343, *VF_1639*, and *VF_A0216*) and another that lacked six PDEs (Δ6PDE;

lacking *pdeV*, *VF_2480*, *binA*, *VF_0087*, *VF_1603*, and *VF_A0506*). These two strains were

derived using a recently generated collection of 50 single mutants (27). The specific gene

disruptions were chosen based both on technical reasons (*e.g.,* location of a gene of interest in

the chromosome or the specific antibiotic resistance cassette present) and on preliminary

phenotypic assessments. For the latter, we focused on mutations that, in single mutant

experiments, impacted known c-di-GMP-dependent phenotypes, primarily the control over

cellulose production (via Congo red binding) (24). Ultimately, our goal was to generate strains

with decreased and increased levels of c-di-GMP. To determine if we achieved this goal, we

examined the levels of c-di-GMP in these strains using the pFY4535 c-di-GMP reporter (28) and

found that the Δ7DGC strain had c-di-GMP levels over 9-fold lower than the parent strain, while

the Δ6PDE strain exhibited c-di-GMP levels about 1.5-fold higher (**Fig. 1A**). These differences

were observed in rich medium as well as in filter-sterilized Instant Ocean (FSIO) under

conditions that mimic squid inoculation (**Fig. 1A**), suggesting to us that these strains would be

effective for studying the role of c-di-GMP during squid colonization. We examined phenotypes

known to be regulated by c-di-GMP, including cellulose biofilm formation and flagellar motility.

The Δ7DGC strain displayed elevated motility on TBS soft agar, reduced Congo red binding,

and reduced activity of a *bcsQ'-gfp*[+] cellulose locus transcriptional reporter (**Fig. 1B-D**). The

Δ6PDE strain exhibited the opposite phenotypes (**Fig. 1B-D**). These data supported our

contention that we could use the multiple deletion approach to examine c-di-GMP regulation in

the symbiont. Together, these data and those below support that the Δ7DGC strain represents

cells in a low c-di-GMP state whereas the Δ6PDE cells are in a high c-di-GMP state, and for the

remainder of the study we will refer to these strains as the "Low cdG" and "High cdG" strains,

respectively.

Elevated c-di-GMP levels interfere with productive colonization initiation

To determine whether the lower and/or higher c-di-GMP levels impacted symbiotic colonization,

we introduced the Low cdG and High cdG strains to newly hatched *E. scolopes* squid. Each

strain was individually inoculated into a bowl with hatchling squid and allowed to initiate

colonization for 3 h. The squid were transferred to new water, and the bacterial load in each

host was measured at 18 and 48 hours-post-inoculation (hpi) (29). For WT *V. fischeri* ES114,

this process leads to approximately 10^5 CFU per squid at 18 and 48 hpi, and a similar yield was

observed for the Low cdG strain (**Fig. 2**). This was surprising to us, given that there are 50

genes that encode c-di-GMP synthesis/degradation factors, yet downregulation of c-di-GMP

levels by over 9-fold did not impact the ability to colonize the squid host (**Fig. 2**). To ask whether

the low cdG strain exhibited a more subtle colonization defect that we were not detecting in this

single strain colonization assay, we performed a competitive colonization assay in which the

Low cdG and WT strain were co-inoculated. We reasoned that if the Low cdG strain had a slight

defect in the host, then the competitive assay could reveal this effect. However, in the

competitive assay, the Low cdG strain exhibited similar fitness as the WT strain (**Fig. S1**). In

contrast, in the single-strain colonization assay, the High cdG strain had a substantial deficit in

bacterial yield in the host as early as 18 hpi and also at 48 hpi, with a median CFU below 10^3

per light organ (**Fig. 2**). In the competitive colonization, the High cdG strain displayed a

significant deficit, as WT outcompeted the High cdG strain by over 3000-fold (**Fig. S1**). Growth

of the High cdG strain in culture was similar to WT (**Fig. S2**), arguing that the colonization defect

is due to a developmental deficit in the host and not an inherent growth property of the strain.

From these data, we conclude that high levels of c-di-GMP are detrimental for the initiation of a

beneficial colonization in the squid light organ, whereas low levels of c-di-GMP do not impact

the colonization behavior during the first 48 h of symbiosis.

To ask whether the phenotypes we observed in the High cdG strain were due to regulation of c-

di-GMP levels or whether they were due to a specific gene product that is absent (i.e., signaling

specificity), we introduced a PDE from *V. cholerae*, VC1086, that is known to be active in

heterologous bacteria (30, 31), and that we predicted would lower c-di-GMP levels in *V. fischeri*.

Across the phenotypes tested, the plasmid-expressed VC1086 had only a slight effect in a WT

background (**Fig. 3**). However, in the High cdG background, expression of VC1086 reduced

cdG levels by 1.6-fold, restored swimming motility on TBS agar, and reduced Congo red binding

1.7-fold in the High cdG background (**Fig. 3A-C**). In a squid colonization assay, expression of

the *V. cholerae* PDE rescued colonization of the High cdG strain (**Fig. 3D**). This set of

experiments confirmed that our results were due to altering levels of c-di-GMP rather than the

absence of a single gene product. We conclude that lower levels of c-di-GMP are not harmful to

establishing a symbiotic association, while higher levels are detrimental to symbiotic initiation.

The High c-di-GMP strain exhibits flagellar motility *in vivo*

In culture, the High cdG strain had a substantial swimming motility defect (**Fig. 1B**). We

therefore considered the hypothesis that a key basis for the poor colonization by the High cdG

strain was due to the altered flagellar motility, which is required for host colonization (8, 15, 17).

In culture, we did observe a migration ring in the presence of magnesium sulfate, which

promotes *V. fischeri* motility (32), demonstrating that the strain is not amotile (**Fig. 1B**). Cells

from the outer ring of motility from the High cdG strain (**Fig. 1B**) exhibited the same phenotype

as the parent High cdG strain upon reinoculation into a new plate (i.e., low motility) (**Fig. S3**),

arguing that the motility observed was not due to spontaneous suppressor mutants. An open

question, then, was whether this strain exhibited flagellar motility under the conditions

experienced in the light organ mucus. We therefore performed a competitive colonization assay

between the High cdG strain and the amotile High cdG Δ*flrA* isogenic derivative. The High cdG

strain outcompeted the amotile Δ*flrA* derivative, with a median advantage of approximately 10-

fold (**Fig. 4**), supporting that the High cdG strain is indeed motile in the host.

Elevated c-di-GMP levels lead to altered aggregate morphology *in vivo*

The *V. fischeri*-squid system provides an opportunity to combine genetic and imaging

approaches to study the planktonic-to-biofilm transition during colonization of a symbiotic host.

Symbiosis polysaccharide (Syp) is required for host colonization (5), while the distinct cellulose

polysaccharide is not known to impact host phenotypes, so we questioned whether c-di-GMP

would impact Syp-dependent aggregation *in vivo*. At 3 hpi, we examined *in vivo* biofilm

aggregates using *V. fischeri* that constitutively expressed a fluorescent protein (GFP or

mCherry). Compared to the Low cdG strain, the High cdG strain produced larger-sized

aggregates, while the WT aggregates were an intermediate size (**Fig. 5A,B**). In over 10% of the

light organs examined, the High cdG strain produced four or more aggregates, whereas squid

colonized by the WT strain had three aggregates at most (**Fig. 5C**).

Although the pFY4535 reporter plasmid was not designed for *in vivo* squid experiments, we

attempted to use this tool to measure c-di-GMP levels in the host for the Low cdG, WT, or High

cdG strain (28). AmCyan is produced constitutively from the plasmid, enabling us to locate the

bacterial cells in the host. Turbo RFP as a reporter for c-di-GMP levels was substantially

reduced in the Low cdG strain compared to WT, consistent with what we observed *in vitro* (**Fig.**

S4, Fig. 1). Examination of the High cdG strain, however, revealed aggregate morphology that

was not consistent with what we observed above with a constitutive GFP marking the cells in

aggregates (**Fig. 5**). Specifically, we saw diffuse small groups of cells in the host in the High

cdG/pFY4535 strain, rather than the aggregates that are produced by the High cdG/pVSV102

(GFP) strain. The High cdG/pFY4535 phenotype is the outlier, as the gene reporter assays

described in the next paragraph use a constitutive mCherry to mark the cells, and their

aggregate behavior matches that of the constitutive GFP in **Fig. 5**. Put together, our

interpretation of these data is that the high level of reporter expression in the High cdG strain *in*

vivo alters the aggregative behavior of the cells. While this effect prohibits accurate

measurements of c-di-GMP levels in the host, it is consistent with a model in which high c-di-

GMP levels in the strain are retained *in vivo*.

To test if the Syp polysaccharide was required for the pattern of multiple bacterial aggregates

we observed in the High cdG strain, we deleted the gene encoding the structural Syp protein

SypQ in the High cdG background. This strain failed to colonize the host robustly at 48 hpi (**Fig.**

S5B) and did not form detectable aggregates (**Fig. 5A,B**), demonstrating that Syp-dependent

biofilm is required for *in vivo* aggregation in the High cdG background. We asked whether

transcription of the *syp* locus was altered in the High cdG strain. Activity of a *sypA'-gfp*[+] reporter

was examined during the aggregation phase, where we found that in the host, *sypA'-gfp*[+]

reporter activity was substantially reduced in the High cdG strain compared to the WT parent

(**Fig. 6A, Fig. S6A**). High levels of c-di-GMP therefore negatively impact the expression of the

symbiosis polysaccharide locus *in vivo*. Notably, when we asked whether the High cdG strain

had altered *sypA'-gfp*[+] reporter activity *in vitro*, we found no significant effects in LBS medium or

in TBS-Ca^{2+} medium that leads to a higher basal *syp* level (**Fig. 6C**) (33). Therefore, c-di-GMP

exerts a host-specific effect on symbiotic biofilm expression.

High levels of c-di-GMP repress *syp* polysaccharide transcription via the cellulose

polysaccharide and in a host-dependent manner

In previous studies in the squid host, larger bacterial aggregates have correlated with an

upregulation of *syp* gene transcription and a higher propensity for colonization initiation and

competitive fitness (5, 34, 35). In this case, however, the larger aggregates of the High cdG

strain have reduced *sypA'-gfp*[+] reporter activity and exhibit poor colonization (**Fig. 6A, Fig. S6A,**

Fig. 2). We therefore asked if there was another component of the aggregate that may

contribute to the morphology observed but that may not enhance colonization ability in the same

way that additional Syp polysaccharide promotes colonization. A candidate for this component

is cellulose, as we demonstrated above that cellulose is upregulated in the High cdG strain (**Fig.**

1C,D). Therefore, we asked whether there may be an interaction between the cellulose and Syp

polysaccharide biosynthetic pathways. We measured *bcsQ* transcription *in vivo* using a

fluorescent transcriptional reporter (*bcsQ'-gfp*[+]), and we determined that *bcsQ'-gfp*[+] reporter

activity was elevated in bacterial aggregates in the host mucus in the High cdG strain compared

to WT (**Fig. 6B, Fig. S6B**). Using a *bcsA*::Tn*erm* transposon insertion mutant that interrupts

cellulose synthase, we examined colonization in the context of the High cdG background.

Interruption of *bcsA* did not improve colonization at 18 or 48 hpi compared to the High cdG

strain (**Fig. S5**). However, compared to the High cdG strain, a *bcsA* derivative exhibited higher

sypA'-gfp⁺ activity *in vivo* (**Fig. 6B, Fig. S6B**), yet no difference in *sypA'-gfp*⁺ activity was

observed *in vitro* with the same strains, even with *syp* induced by added calcium chloride (**Fig.**

6C) (33). These data present a novel connection between transcriptional control of

polysaccharide systems that occurs specifically in the host.

DISCUSSION

Cyclic di-GMP impacts multiple pathways during beneficial colonization. Our study was

motivated by the question of how c-di-GMP impacts a beneficial microbe-host association. To

ask this question, we focused on the *V. fischeri*-squid mutualism. This well-studied binary

association allows us to examine individual stages of colonization using strains deleted for 6-7

genes and that consequently have altered levels of c-di-GMP. We found that a strain with

elevated levels of c-di-GMP exhibited striking defects during host colonization. The High cdG

strain formed aggregates that trended larger (**Fig. 4**), had altered host-specific polysaccharide

gene expression in the aggregates (**Fig. 6A,B, Fig. S6**), exhibited reduced flagellar motility (**Fig.**

1B), and displayed a significant colonization defect as enumerated by bacterial counts in the

host as early as 18 hpi (**Fig. 2**). Bacterial counts did not recover by 48 hpi, suggesting that high

levels of c-di-GMP confer both an initiation defect and an accommodation defect on the

colonizing strains (36), as discussed further below. Colonization of the High cdG strain was

rescued by overexpression of a heterologous PDE that does not have significant homology to

proteins in *V. fischeri* (**Fig. 3D**), strongly supporting that the observed effects are due to the c-di-

GMP levels and not to individual functions of the gene products affected in the multiple deletion

strains.

Our results parallel work in several pathogenic bacteria, where elevated c-di-GMP levels are

correlated with a decrease in virulence (3, 37–40). In *Yersinia pestis*, the small number of

relevant enzymes made it possible to remove all of the DGCs or PDEs. Deletion of both genes

for the functional DGCs effectively eliminated c-di-GMP in the cell (measured by LC-MS), yet

virulence in mouse infection models was not affected (40). However, deletion of the single

functional PDE gene, *hmsP*, led to approximately 10-fold higher c-di-GMP levels and led to

reduced virulence in a subcutaneous infection model (40). In *V. cholerae*, elevated c-di-GMP

(visualized by 2D-TLC) led to 5-fold reduced competitive colonization in the infant mouse small

intestine model (39). In that study, the authors tested if the reduction was due to overexpression

of biofilm (VPS), but it was not, and they attributed the phenotype to the resulting virulence gene

expression downstream of c-di-GMP. We note that the magnitude of the effect we observe in *V.*

fischeri colonization of bobtail squid is substantially higher: >100-fold in a single-strain

colonization assay. It therefore seems likely that this aspect of mutualism is broadly shared with

pathogenic colonization, and we expect there to be strong constraints to retain c-di-GMP levels

low during the initiation of the host colonization process. There are also likely to be beneficial

symbiosis-specific adaptations within c-di-GMP signaling. For example, the zebrafish gut

symbiont *Aeromonas veronii* responds to host amino acids to inhibit a diguanylate cyclase,

thereby stimulating bacterial motility into the host (41). It remains to be seen whether there are

similar specific adaptations in the *V. fischeri*-squid symbiosis.

Despite encoding 50 genes that manipulate c-di-GMP, we were surprised to find that

substantially lowering levels of c-di-GMP (approximately 9-fold) (**Fig. 1A**) led to no significant

defect in host colonization. In the Low cdG strain, we detected slightly smaller bacterial

aggregates in the host mucus (**Fig. 5**), but no consequent diminution of bacterial counts in the

light organ at 18 hpi or 48 hpi (**Fig. 2**). The Low cdG strain similarly exhibited no competitive

colonization defect (**Fig. S1**). When examining transcriptional reporters for the symbiosis

polysaccharide gene *sypA* or the cellulose biosynthesis gene *bcsQ in vivo*, the Low cdG strain

resembled the wild-type parent (**Fig. 6A,B, Fig. S6**). We note that colonization was intact

despite this strain exhibiting strong biofilm and motility phenotypes *in vitro* (**Fig. 1**), thus

emphasizing the importance of studying these phenotypes in the relevant host system. The key

bacterial behaviors during the initial stages in the host include adhering to host mucus and then

cilia within the mucus (10); forming the Syp-dependent aggregates (5, 9–11); migrating toward

the host pores in a manner independent of flagellar motility (4, 12); employing flagellar motility

and chemotaxis to swim toward chitin oligosaccharides released by the host (8, 14–17); and

resisting host innate immune attacks as they traverse the pore into the duct, antechamber,

bottleneck, and then crypts of the light organ (42–45). These processes take 18-24 h and

encompass the "initiation" and "accommodation" phases of the symbiosis. Around 18-24 h, the

interaction shifts to the "persistence" phase, where quorum sensing stimulates bioluminescence

in the host, and a daily (diel) rhythm ensues in which bacteria consume nutrient sources

provisioned from the host squid. We observe specific initiation defects in biofilm reporter gene

expression, biofilm aggregate number and architecture, flagellar motility, and bacterial counts in

the host at 18 hpi. We additionally observe an accommodation defect in that bacterial counts do

not recover by 48 hpi, arguing that the high cdG strain continues to inhibit expansion in the

crypts following the initial defect. The accommodation phenotype is unlikely to be due to

flagellar motility, as most bacteria lose their flagella during the mature association (8).

Therefore, these effects may be due to regulation of biofilm genes in the host (7) or may be due

to other unrecognized effects in the High cdG strain.

By using strains with altered c-di-GMP levels, we have uncovered regulatory differences in

transcriptional reporters in culture versus the host. In culture, we observed a 3.2-fold increase in

bcsQ'-gfp[+] reporter activity in the WT strain relative to the Low cdG strain (**Fig. 1D**). However, in

the animal, we observed similar *bcsQ'-gfp*[+] reporter activity in the two strains (**Fig. 6B, Fig.**

S6B), suggesting that the lower *bcsQ'-gfp*[+] activity of the WT strain may occur in response to

the host environment. Similarly, the upregulation of *bcs* gene expression in the background of

elevated c-di-GMP conditions led to a substantial downregulation of *sypA'-gfp*[+] reporter activity

in the host, yet no comparable effect was observed in two distinct culture conditions (**Fig. 6**).

Given the multiple cases in which gene expression in the host is distinct from that observed in

culture, it remains an intriguing question as to how the host influences pathways linked to c-di-

GMP signaling.

If the Low cdG strain thrives in the host under conditions where levels of c-di-GMP remain low,

is c-di-GMP required for *V. fischeri* biology in the squid? Given that there are another twenty-

one DGCs and five putative bifunctional DGC/PDEs still present in the Low cdG strain, our

results do not preclude a requirement for c-di-GMP during colonization. It remains possible that

additional DGCs perform localized functions that were not addressed in the current study and/or

that DGCs play critical roles at later stages in the colonization process.

Construction of multiple-gene deletions in *V. fischeri* is an effective tool to tackle gene

families. The recent development of a method for construction of serial deletions (26) enabled

the construction of multiple-mutant strains such as the Low cdG and High cdG strains in this

study. We have further extended the method to allow for detection of barcoded deletions in

competitive colonization experiments (46). The work in this study highlights the utility of such

approaches to interrogate gene families. There are many instances in which deletions of

multiple genes could clarify the role of various biological processes. Previously, we identified a

role for chemotaxis toward chitin oligosaccharides as a key developmental stage during squid

colonization (16). Efforts from our group and others have studied 23 *V. fischeri* chemoreceptors

through a combination of single and multiple mutants (47–49), but there are a total of 43 methyl-

accepting chemoreceptors encoded in the ES114 genome, and the receptor(s) for chitin

oligosaccharides remain elusive. Generation of multiple mutant strains has been especially

valuable to dissect gene families in *V. cholerae* (50), and our work suggests that a similar

approach will be useful in *V. fischeri*.

Elevated c-di-GMP levels diminish, but do not eliminate, flagellar motility *in vivo*

The High cdG strain has a significant soft agar motility defect (**Fig. 1B**), and amotile strains are

at a significant colonization disadvantage compared to WT. Our competition of the High cdG

strain with the isogenic Δ*flrA* derivative revealed a 15-fold advantage for the former during squid

colonization (**Fig. 4**). Our interpretation of this result is that the High cdG strain retains the ability

to be motile in the host despite the elevated c-di-GMP levels. The magnitude of the competitive

advantage (15-fold) is diminished compared to that of the wild-type strain competing against a

Δ*flrA* strain (>1,000-fold) (46, 51). Therefore, it seems likely that c-di-GMP does impact motility

in the host, but not to the extent that it fully prohibits colonization.

Cyclic di-GMP has a qualitative effect on the nature of the exopolysaccharide in animal

tissue. The increased number of aggregates observed in the High cdG strain in squid is, to our

knowledge, the first time such a phenotype has been observed (**Fig. 5**). Hypermotile mutants

were noted to exhibit more diffuse aggregates on the light organ surface, though these were

often smaller than their WT counterparts (15). Multiple studies have noted larger *in vivo*

aggregates upon overexpression of biofilm activator RscS or removal of biofilm inhibitor BinK (5,

35). In both cases, it was shown that Syp was the relevant pathway induced, and the cells within

the resulting larger aggregates were functional to outcompete those from WT aggregates. The

morphology and gene expression in High cdG aggregates are significantly more complicated

and implicate an interaction between cellulose and Syp exopolysaccharides. We showed that

cellulose on its own is not required for colonization (**Fig. S5B**). However, the high c-di-GMP

condition revealed a role for the *bcs* (cellulose synthase) locus in regulating *syp* (mirroring

results obtained in specific agar conditions in (52)) (**Fig. 6A, Fig. S6A**). Therefore, while

cellulose production on its own does not impact colonization, in the High cdG background it

significantly impacts expression of *syp*. In *Pseudomonas aeruginosa*, roles for multiple

exopolysaccharide systems in biofilm formation have been explored, for example as the Psl and

Pel systems have distinct but also overlapping roles in attachment and biofilm formation (53). In

V. fischeri, our work supports data in the literature that Syp is the key system for *in vivo* biofilm

formation and that the cellulose system is dispensable for host colonization. At the same time,

this study reveals a novel transcriptional interaction between two exopolysaccharide systems.

This interplay may impact the formation of multiple-exopolysaccharide biofilms during later

stages of host colonization or under environmental conditions that have yet to be explored, and

points to a novel regulatory mechanism that connects biofilm regulation in an animal host.

In conclusion, this work identified a role for c-di-GMP regulation during the initiation of a

mutualism, identified specific developmental stages in the host that were regulated by altered c-

di-GMP levels, and revealed a transcriptional interplay between multiple exopolysaccharide

systems that is apparent only in the host. Our study therefore lays the groundwork for future

investigations into how this widespread bacterial compound influences bacterial behaviors

beyond pathogenesis to impact microbiome behavior and interactions with symbiotic hosts.

MATERIALS AND METHODS

Bacterial strains, plasmids, and media

V. fischeri and *E. coli* strains used in this study are listed in Table 1. Plasmids used in this study

are listed in Table 2. *V. fischeri* strains were grown at 25°C in Luria-Bertani salt (LBS) medium

(per liter: 25 g Difco LB broth [BD], 10 g NaCl, 50 mL 1 M Tris buffer [pH 7.5]), or TBS (per liter:

10 g tryptone [VWR], 20 g NaCl, 35 mM $MgSO_4$ [where noted], 10 mM calcium chloride [where

noted], 50 mM 1 M Tris buffer [pH 7.5]) where noted. *E. coli* strains used for cloning and

conjugation were grown at 37°C in Luria-Bertani (LB) medium (per liter: 25 g Difco LB broth

[BD]). When needed, antibiotics were added to the media at the following concentrations:

kanamycin, 100 µg/mL for *V. fischeri* and 50 µg/mL for *E. coli*; chloramphenicol, 1 or 5 µg/mL for

V. fischeri and 25 µg/mL for *E. coli*; erythromycin, 2.5 or 5 µg/mL for *V. fischeri*; gentamicin, 2.5

µg/mL for *V. fischeri* and 5 µg/mL for *E. coli*; trimethoprim, 10 µg/mL for *V. fischeri*;

spectinomycin, 160 µg/mL in LB for *V. fischeri*. When needed, thymidine was added at 0.3 mM

for *E. coli*. Growth media was solidified using 1.5% agar when needed. For Congo red agar, 40

µg/mL Congo red and 15 µg/mL Coomassie blue were added to LBS. For Xgal agar, 20 µg/mL

Xgal were added to LBS. Plasmids were introduced from *E. coli* strains into *V. fischeri* strains

using standard techniques (54, 55)**.**

DNA synthesis and sequencing

Primers used in this study are listed in Table 3 and were synthesized by Integrated DNA

Technologies (Coralville, IA). Full inserts for cloned constructs and gene deletions were

confirmed by Sanger Sequencing at Functional Biosciences via UW-Madison. Sequence data

were analyzed using Benchling. PCR to amplify constructs for cloning and sequencing were

performed using Q5 High-Fidelity DNA polymerase (NEB), OneTaq DNA polymerase (NEB), or

MilliporeSigma Novagen KOD DNA Polymerase. Diagnostic PCR was performed using GoTaq

polymerase (Promega) or OneTaq DNA polymerase.

Construction of multiple gene deletion mutants

The multiple DGC and PDE mutants were constructed as follows. First, single gene deletions

were generated using PCR-SOE and *tfoX*-induced transformation methods previously described

(26, 56); some of these mutants have been previously described (23, 52). Briefly, regions

flanking a gene of interest were amplified using primers listed in Table 3 and joined to an

antibiotic-resistance cassette (e.g., Trim^R/Erm^R/Spec^R) flanked by Flp-recombinase target (FRT)

sites that was amplified with primers 2089 and 2090. The resulting fused PCR product was

transformed into a *V. fischeri* strain (generally ES114) that contained a TfoX-overproducing

plasmid (pLostfoX or pLostfoX-Kan plasmid (25, 56). Selection for the integrants was conducted

on selective agar. The deletion/insertion was verified by PCR, generally with the outside primers

used to generate the flanking fragments. Next, strains with multiple mutations were generated

using genomic DNA from specific single mutants as follows. A *tfoX*-overexpression plasmid was

introduced into a mutant of interest and transformed with genomic DNA containing a specific

mutation, followed by selection for and verification of the mutation as described above. The

resulting strain carrying a *tfoX*-overexpression plasmid could then serve as a recipient for the

introduction of the next mutation. Because some mutants were generated using the same

antibiotic resistance cassette, as needed the Flp-encoding plasmid pKV496 was introduced to

induce the loss of the antibiotic resistance cassette and the retention of a 112 bp FRT scar in

the gene. Following strain construction, the integrity of each deletion junction was verified with

diagnostic primers listed in Table 3 to ensure that only recombination at local FRT sites

occurred. As noted in the strain list (Table 1), the final cassette often was left in place. As an

example, for Δ7PDE strain KV9601, the starting strain, KV8969 (23) (Δ*pdeV*::FRT) carrying *tfoX*

plasmid pLostfoX-Kan was sequentially transformed with genomic DNA carrying the

Δ*VF_2480*::FRT-Erm, Δ*binA*::FRT-Trim, and Δ*VF_0087*::FRT-Spec mutations. The cassettes

were removed, and the *tfoX* plasmid reintroduced. Next, the Δ*VF_1603*::FRT-Trim mutation was

introduced, and the cassette removed. Finally, the Δ*VF_A0506*::FRT-Trim mutation was

introduced. Similar methods were used to generate KV9599 (Δ7DGC) and the derivatives of

KV9601 that carry mutations in *sypQ* or *bcsA* (the latter of which was derived from strain

KV8408, which carries a Tn mutation in *bcsA* (26)) rather than a cassette as described above).

Construction of *bcsA*::Tn*erm* strain

Genomic DNA isolated from KV9767 (Δ6PDE *bcsA*::Tn*erm*) was introduced into MJM1100 via

transformation using pLostfoX-Kan (25, 56). Mutant candidates were selected using

erythromycin and screened by PCR using RYI560 and RYI561 primers.

Construction of the *bcsQ'-gfp*+ transcriptional reporter plasmid pRYI063

The *bcsQ* promoter (*bcsQ* -444 to -29; P*bcsQ*) was amplified from MJM1100 genomic DNA

using RYI576 and RYI577 primers that added flanking XmaI and XbaI restriction sites. The

amplified P*bcsQ* and pTM267 were digested with XmaI and XbaI. Digested pTM267 was treated

with antarctic phosphatase. Digested P*bcsQ* was ligated into digested pTM267 using T4 DNA

ligase (NEB). The ligation reaction was transformed into chemically competent DH5α λpir.

Transformant candidates were selected using chloramphenicol and confirmed by Sanger

sequencing using the RYI526 primer.

Construction of the *VC1086* expression plasmid pRYI064

pVSV105 was amplified using RYI578 and RYI579 primers. *VC1086*, along with 101 bp

upstream and 99 bp downstream, was amplified from *Vibrio cholerae* C6706 str2 genomic DNA

using RYI580 and RYI581 primers. *VC1086* was assembled with pVSV105 by Gibson assembly

using NEBuilder HiFi DNA Assembly Master Mix (NEB). The Gibson reaction was transformed

into chemically competent DH5α λpir and transformant candidates were selected using

chloramphenicol. Plasmid was screened by PCR using M13 For. (-20) and M13 Rev. (-48)

primers. Plasmid was confirmed by Sanger sequencing using M13 For. (-20), M13 Rev. (-48),

RYI584, RYI587, RYI585, and RYI586 primers.

C-di-GMP quantification

Strains were streaked on LBS agar and incubated overnight at 25°C. Liquid LBS was inoculated

from single colonies and grown at 25°C overnight on a rotator. 8 µL of liquid culture were

spotted onto LBS agar and incubated at 25°C for 24 h. Spots were resuspended in 1 mL 70%

Instant Ocean (IO). OD_{600}, TurboRFP (555 nm excitation/585 nm emission), and AmCyan (453

nm excitation/486 nm emission) for each resuspended spot were measured in triplicate using

BioTek Synergy Neo2 plate reader. To calculate c-di-GMP levels, TurboRFP values (reports c-

di-GMP levels) were normalized to AmCyan values (constitutively expressed).To compare c-di-

GMP levels of strains in LBS vs. IO, 1 mL of overnight LBS liquid cultures were pelleted at 8000

x g for 10 min, resuspended in 600 µL filter-sterilized IO (FSIO), and both the FSIO and LBS

samples were incubated at room temperature for 3 hr; LBS and FSIO strains were pelleted at

8000 x g for 10 min, and resuspended in 600 µL FSIO. OD_{600}, TurboRFP, and AmCyan for each

sample were measured in triplicate as described above.

Motility assay

Strains were streaked on TBS and TBS-Mg^{2+} agar and incubated overnight at 25°C. Single

colonies were inoculated into TBS and TBS-Mg^{2+} soft agar (0.3% agar), respectively, and

incubated at 25°C for 5 h. Images of plates were taken using a Nikon D810 digital camera and

diameter of migration was measured using ImageJ software.

Isolation of cells from the outer edge of motility ring

Strain was streaked on TBS-Mg^{2+} agar and incubated overnight at 25°C. A single colony was

inoculated into TBS-Mg^{2+} soft agar (0.3% agar) and incubated at 25°C for 18 h. Cells from the

outer edge of the resulting motility ring were streak purified on LBS. Isolates were then

reinoculated into new TBS-Mg^{2+} motility agar as described above.

Congo red assay

Strains were streaked on LBS agar and incubated overnight at 25°C. Liquid LBS was inoculated

with single colonies and grown at 25°C overnight on a shaker. 4 µL spots of liquid culture were

spotted on LBS Congo red agar and incubated 24 h at 25°C. Spots were transferred onto white

printer paper (26) and images were scanned as TIFF files. Congo red binding was quantified

using ImageJ software.

syp and *bcs* **transcriptional reporter analysis** *in vitro*

Strains were streaked on LBS agar and incubated overnight at 25°C. Liquid media was

inoculated with single colonies and grown at 25°C for 24 hr on a shaker. 4 µL of overnight

cultures were spotted onto TBS or TBS-Ca^{2+} agar plates and grown at 25°C for 24 hr (52).

Spots were imaged using the Zeiss Axio Zoom.v16 large-field 561 fluorescent stereo

microscope and fluorescence levels were measured using the Zen Blue Software polygon tool.

To calculate gene promoter activity, GFP values (reports promoter activity) were normalized to

mCherry values (constitutively expressed).

Growth curves

Strains were streaked on LBS agar and incubated overnight at 25°C. Liquid LBS was inoculated

with single colonies of each strain in triplicate and grown at 25°C overnight on a rotator.

Cultures were diluted 1:1000 in LBS and grown at 25°C on a rotator to OD$_{600}$ ~0.5. Cultures

were diluted 1:100 in LBS (n = 6 technical replicates per culture) in a 96-well microtiter plate.

OD$_{600}$ was measured every 15 min for 20 hr at 25°C using BioTek Synergy Neo2 plate reader.

Doubling time was calculated in Prism using a nonlinear regression analysis of each strain

during exponential phase.

Squid single strain colonization assay

E. scolopes hatchlings were colonized with approximately 10^3-10^5 CFU/mL of bacteria

according to standard procedure (29). At 18 hpi and 48 hpi, luminescence of hatchlings was

measured using the Promega GloMax 20/20 luminometer and euthanized by storage at -80°C.

CFU counts per light organ were determined by plating homogenized euthanized hatchlings and

counting colonies.

Squid competitive colonization assay

Strains were grown overnight in liquid LBS at 25°C on a rotator. Strains were diluted 1:80 in

liquid LBS and grown to an OD_{600} of approximately 0.2. The strains were mixed in a 1:1 ratio

with a competing strain carrying a constitutive *lacZ* on the pVSV103 plasmid and the mixed

cultures were used to inoculate *E. scolopes* hatchlings with approximately 10^3-10^4 CFU/mL. At 3

hpi, hatchlings were washed and transferred to 40 mL filter-sterilized Instant Ocean (FSIO).

Water was changed at 24 hpi and hatchlings were euthanized at 48 hpi by storage at -80°C.

Euthanized hatchlings were homogenized and plated on LBS-Xgal agar. Competitive index of

strains was measured by calculating the blue/white colony ratios as previously described (25,

29).

Squid aggregation assays

Strains were grown overnight in liquid LBS at 25°C on a rotator. *E. scolopes* hatchlings were

inoculated with approximately 10^5-10^8 CFU/mL of bacteria. At 3 hpi, squid were anesthetized in

2% EtOH in FSIO and immediately dissected and imaged as described below, or were fixed in

4% paraformaldehyde in 1x mPBS (50 mM phosphate buffer, 0.45 M NaCl, pH 7.4) for

approximately 48 h. Fixed hatchlings were washed in 1x mPBS x4, dissected, and imaged using

the Zeiss Axio Zoom.v16 large-field 561 fluorescent stereo microscope. Aggregate area was

selected and fluorescence levels were measured using the Zen Blue Software polygon tool.

Note that the analyses were conducted with all data as well as solely with the inocula in the 10^6-

10^7 range and the same results were observed, so we included all of the data in the manuscript.

Data analysis

Congo red binding was quantified using ImageJ by subtracting the WT gray value from the

mutant gray value and multiplying the value by -1. Fluorescence of strains in liquid culture were

measured using a BioTek Synergy Neo2 plate reader. Fluorescence of strains spotted on agar

plates and bacterial aggregates in the squid were measured using Zen Blue Software.

GraphPad Prism was used to generate graphs and conduct statistical analyses. Graphs were

further refined in Adobe Illustrator.

ACKNOWLEDGMENTS

We thank Fitnat Yildiz for providing the pFY4535 reporter plasmid, Chris Waters for input on this

project, and Ali Razvi and Steven Eichinger for help with strain construction. This work was

funded by NIGMS grant R35 GM119627 to M.J.M., NIGMS grant R35 GM130355 to K.L.V, and

R.Y.I. was supported by NIGMS training grant T32 GM007215.

594
595

596   **Table 1: Strains used in this study**

Strain	Genotype	Source or Reference(s)
V. fischeri		
KV4674 = ES114	Natural isolate, squid light-organ (Visick Lab Stock)	57, 58
KV8069	KV4674 / Δ*sypQ*::FRT-Cam	33
KV8408	KV4674 / *bcsA*::Tn*erm*	26
KV8920	KV4674 / Δ*VF_1639*::FRT-Spec	59
KV8932	KV4674 / Δ*VF_1200*::FRT-Cam	60
KV8969	KV4674 / Δ*pdeV*::FRT	23
KV9599 = Δ7DGC (Low cdG)	Δ*VF_1200*::FRT Δ*mifA*::FRT Δ*VF_1245*::FRT Δ(*VF_A0342-VF_A0343*)::FRT Δ*VF_1639*::FRT Δ*mifB*::FRT-Spec	This study
KV9601 = Δ6PDE (High cdG)	Δ*pdeV*::FRT Δ*VF_2480*::FRT Δ*binA*::FRT Δ*VF_0087*::FRT Δ*VF_1603*::FRT Δ*VF_A0506*::FRT-Trim	This study
KV9767	KV9601 *bcsA*::Tn*erm*	This study
KV9769	KV9601 Δ*sypQ*::FRT-Cam	This study
MJM1100 = ES114	Natural isolate, squid light-organ (Mandel Lab Stock)	57, 58
MJM1438	MJM1100 / pM1422	35
MJM4009	MJM1100 / pFY4535	This study
MJM4135	KV9599 / pFY4535	This study
MJM4137	KV9601 / pFY4535	This study
MJM4138	KV9560 / pFY4535	This study
MJM4139	KV9569 / pFY4535	This study
MJM4140	KV9599 / pVSV102	This study
MJM4142	KV9601 / pVSV102	This study
MJM4198	KV9599 / pM1422	This study
MJM4200	KV9601 / pM1422	This study

Strain	Genotype	Source or Reference(s)
MJM4308	KV9767 / pFY4535	This study
MJM4310	KV9769 / pFY4535	This study
MJM4312	KV9767 / pVSV102	This study
MJM4314	KV9769 / pVSV102	This study
MJM4322	KV9767 / pM1422	This study
MJM4484	MJM1100 *bcsA*::Tn*erm*	This study
MJM4592	KV9601 / pVSV103	This study
MJM4593	MJM1100 / pRYI063	This study
MJM4594	KV9599 / pRYI063	This study
MJM4595	KV9601 / pRYI063	This study
MJM4596	KV9767 / pRYI063	This study
MJM4597	MJM4484 / pRYI064	This study
MJM4603	KV9601 Δ*flrA*::*erm*-bar	This study
MJM4630	MJM1100 / pRYI064	This study
MJM4631	KV9601 / pRYI064	This study
MJM4632	MJM1100 / pFY4535; pRYI064	This study
MJM4634	KV9601 / pVSV105	This study
MJM4635	MJM1100 / pFY4535; pVSV105	This study
MJM4636	KV9601 / pFY4535; pVSV105	This study
E. coli		
KV6937	DH5α / pLostfox-Kan	25
KV8052	π3813 / pKV496	26
MJM542	DH5α λpir / pVSV102	47
MJM552	DH5α λpir / pVSV103	47
MJM579	DH5α λpir / pVSV105	47
MJM1422	DH5α λpir / pM1422	35

Strain	Genotype	Source or Reference(s)
CAB1516	DH5α λpir / pVSV105	47
MJM3999	NEB5α / pFY4535	28
MJM4580	DH5α λpir / pRYI063	This study
MJM4625	DH5α λpir / pRYI064	This study

598
599

Table 2: Plasmids used in this study

Plasmid	Description	Source or Reference(s)
pEVS104	Conjugal helper plasmid (Kan^R)	54
pFY4535	c-di-GMP reporter plasmid (Gent^R)	28
pJJC4	Vector carrying *tfoX* and *litR* genes (Cam^R)	61
pKV494	Vector carrying FRT-Erm^R	26
pKV495	Vector carrying FRT-Cam^R	26
pKV496	pLostfoX-Kan backbone containing the FLP recombinase (Kan^R)	26
pKV521	Vector carrying FRT-Spec^R	26
pLostfoX-Kan	TfoX induction vector for chitin-pathway transformation (Kan^R)	25
pM1422	pTM267 *sypA'-gfp$^+$* (Cam^R)	35
pMCL2	Vector carrying FRT-Trim^R	26
pRYI063	pTM267 *bcsQ'-gfp$^+$* (Cam^R)	This study
pRYI064	pVSV105 carrying *VC1086* (Cam^R)	This study
pVSV102	Constitutive GFP (Kan^R)	62
pVSV103	Constitutive LacZ (Kan^R)	62
pVSV105	Complementation vector (Cam^R)	47

602
603

Table 3: Primer List

Primer Name	Sequence (5'-3')	Notes

2089	CCATACTTAGTGCGGCCGCCTA	Forward primer to amplify antibiotic resistance cassette
2090	CCATGGCCTTCTAGGCCTATCC	Reverse cassette to amplify antibiotic resistance cassettes
2537	AGTGCCTCATATAAGGGTTAAG	Outside forward PCR primer for the Δ(*VF_A0342-A0343*) construct
2538	taggcggccgcactaagtatggTGTTACGTCAACAACCTTTTGG	Inside reverse PCR primer for the Δ(*VF_A0342-A0343*) construct
2543	ggataggcctagaaggccatggATGGCGGATACTGCATTGTATC	Inside forward PCR primer for the Δ(*VF_A0342-A0343*) construct
2544	CCAAATCGACAGAGATCCCC	Outside reverse PCR primer for the Δ(*VF_A0342-A0343*) construct
2593	GCCAATGCAGATATGACAAGG	Outside forward PCR primer for the Δ*VF_1245* construct
2558	taggcggccgcactaagtatggTAAGCGCTGAAGACGAGATTG	Inside reverse PCR primer for the Δ*VF_1245* construct
2559	ggataggcctagaaggccatggATTGAACAAGCAGATAAGGCGC	Inside forward PCR primer for the Δ*VF_1245* construct
2560	GCATTCATTAGTGATGTGCGTG	Outside reverse PCR primer for the Δ*VF_1245* construct
2905	ggataggcctagaaggccatggCACTTCGTGTTAAAGAATTTATAC	Confirmation of Δ*VF_1639*
2564	CTAACCATTCATGCAAGAACC	Confirmation of Δ*VF_1639*
2577	TAAAGCAGCAGCTGAGCAAG	Confirmation of Δ*VF_1200*
2580	GTTTGGTGAGTACCAATCGC	Confirmation of Δ*VF_1200*
2585	CAAAGCATCAAAAACTCACTTGC	Outside forward PCR primer for the Δ*VF_0087* construct
2586	taggcggccgcactaagtatggAATGCTTTGAACACTAAATGAGAG	Inside reverse PCR primer for the Δ*VF_0087* construct
2587	ggataggcctagaaggccatggGGTGAGCCTAAATCGTTACAC	Inside forward PCR primer for the Δ*VF_0087* construct
2588	AACGAACGCCACAAACTCAC	Outside reverse PCR primer for the Δ*VF_0087* construct
944	CGCATACGTTTCTACCGTTTC	Confirmation of ΔVF_0087
945	CCAAGCTGAACGAAGTGGAC	Confirmation of ΔVF_0087
2598	CACTGATGGTTTAGAGCTTGG	Outside forward PCR primer for the Δ*mifA* construct
2599	taggcggccgcactaagtatggCAGCTCTGTTTGATATAGTGTCC	Inside reverse PCR primer for the Δ*mifA* construct
2600	ggataggcctagaaggccatggAGTGAAATGGATC	Inside forward PCR primer for the Δ*mifA*

	AAATCGCATG	construct
2601	TTGATGCGAGATTGAAATACGC	Outside reverse PCR primer for the Δ*mifA* construct
2618	AGAGCAGCTCGTGAGTTATC	Confirmation of Δ*pdeV*
2621	CTTCCATAGTAAGAACCTCTGC	Confirmation of Δ*pdeV*
2648	GCATGTGTACCAAGTAAGCTTG	Outside forward PCR primer for the Δ*VF_A0506* construct
2649	taggcggccgcactaagtatggCATTGTCCATTCATAAGATGGGC	Inside reverse PCR primer for the Δ*VF_A0506* construct
2650	ggataggcctagaaggccatggCGAGGTCAACAATTTGACCC	Inside forward PCR primer for the Δ*VF_A0506* construct
2651	GGCATTGATGAGTCTTAAGCTG	Outside reverse PCR primer for the Δ*VF_A0506* construct
2674	CTAAAGTGCACAAACCATGGC	Outside forward PCR primer for the Δ*VF_1603* construct
2675	taggcggccgcactaagtatggGTCTAAGATTGGTTGACGAGC	Inside reverse PCR primer for the Δ*VF_1603* construct
2676	ggataggcctagaaggccatggTCTGCCAAATGGGTAGATGAG	Inside forward PCR primer for the Δ*VF_1603* construct
2677	TTATTGCAGGTGCTGCACTG	Outside reverse PCR primer for the Δ*VF_1603* construct
2678	TTGGCCACGGTGATCGAAAG	Outside forward PCR primer for the Δ*VF_2480* construct
2679	taggcggccgcactaagtatggTATCGGTTGTCTGGCAACG	Inside reverse PCR primer for the Δ*VF_2480* construct
2680	ggataggcctagaaggccatggACCTACCAAGAAGCGGTTAAG	Inside forward PCR primer for the Δ*VF_2480* construct
2681	TTGCTGCTAATCGTGACAGC	Outside reverse PCR primer for the Δ*VF_2480* construct
1152	CAGTAGAGTCACTACCGTTG	Outside forward PCR primer for the Δ*binA* construct
2714	taggcggccgcactaagtatggTTGTTCAGGAATGTCGATGGC	Inside reverse PCR primer for the Δ*binA* construct
2715	ggataggcctagaaggccatggCACTTTGTGTAATGGGCACTC	Inside forward PCR primer for the Δ*binA* construct, inside forward
2716	GCCTCTGGATTTGGATAGTG	Outside reverse PCR primer for the Δ*binA* construct
850	gggcccAGAATAAACTGCTACTATCT	Confirmation of Δ*binA*
445	AAAACAGTAATCAGGGTGGAACG	Confirmation of Δ*binA*
2753	ATAGATGGTAAAGCGTCATACC	Outside forward PCR primer for the Δ*mifB*

		construct
2754	taggcggccgcactaagtatggACGTAAGCTCACATATGGGC	Inside reverse PCR primer for the Δ*mifB* construct
2755	ggataggcctagaaggccatggTTTACCGTTGCTGATGAACATATG	Inside forward PCR primer for the Δ*mifB* construct
2756	CAAACCATTGAACGCCATGATC	Outside reverse PCR primer for the Δ*mifB* construct
M13 For. (-20)	GTAAAACGACGGCCAGT	For amplification of gene inserts in pVSV105
M13 Rev. (-48)	AGCGGATAACAATTTCACACAGG	For amplification of gene inserts in pVSV105
RYI526	GTGCCCATTAACATCACCATCTAATTCA	Primer to confirm replacement of Kan^R cassette with promoter of choice
RYI560	GATATCAATGAGGTAATGCTTTTATTTTTTGCGAA	Forward primer to confirm transposon insertion in *bcsA*
RYI561	GCATAATTCTTACTTGGTTGCATGGTTTG	Reverse primer to confirm transposon insertion in *bcsA*
RYI576	gctaCCCGGGggtttgtggcatgacgatcact	Forward primer to amplify P*bcsQ* with XmaI restriction site
RYI577	cacgTCTAGAactaactttattctccaatgtattaactacacattttgg	Reverse primer to amplify P*bcsQ* with XbaI restriction site
RYI578	CAGGTCGACTCTAGAGGATCCCC	Forward primer to linearize pVSV105 for Gibson assembly
RYI579	CAGGCATGCAAGCTTGAGTATTCTAT	Reverse primer to linearize pVSV105 for Gibson assembly
RYI580	tactcaagcttgcatgcctgGAAGAACACCATGGCAAAATAAAGCTC	Forward primer to amplify *VC1086* with 101 bp upstream for Gibson assembly with pVSV105
RYI581	gatcctctagagtcgacctgCGGTTAAGCGCTTTCAAAATATCGTTTTC	Reverse primer to amplify *VC1086* with 99 bp downstream for Gibson assembly with pVSV105
RYI584	GGTGATAGAGATTAATCGCCGCTATCC	Sequencing primer for *VC1086*
RYI585	GATAGCTTTCAGAACCCAGCACCT	Sequencing primer for *VC1086*
RYI586	GTTACTGAGCGTCGCGGAGG	Sequencing primer for *VC1086*
RYI587	TTATGGAGATAGTTTGTTATGGAGATAGTTCAGGA	Sequencing primer for *VC1086*

606
607
608
609

REFERENCES

REFERENCES

1. Hengge R, Gründling A, Jenal U, Ryan R, Yildiz F. 2016. Bacterial signal transduction by
cyclic Di-GMP and other nucleotide second messengers. J Bacteriol 198:15–26.

2. Heindl JE, Wang Y, Heckel BC, Mohari B, Feirer N, Fuqua C. 2014. Mechanisms and
regulation of surface interactions and biofilm formation in *Agrobacterium*. Front Plant Sci
5:176.

3. Römling U, Galperin MY, Gomelsky M. 2013. Cyclic di-GMP: the first 25 years of a
universal bacterial second messenger. Microbiol Mol Biol Rev 77:1–52.

4. Nyholm SV, Stabb EV, Ruby EG, McFall-Ngai MJ. 2000. Establishment of an animal-
bacterial association: recruiting symbiotic vibrios from the environment. Proc Natl Acad Sci
U S A 97:10231–10235.

5. Yip ES, Geszvain K, DeLoney-Marino CR, Visick KL. 2006. The symbiosis regulator RscS
controls the *syp* gene locus, biofilm formation and symbiotic aggregation by *Vibrio fischeri*.
Mol Microbiol 62:1586–1600.

6. Visick KL. 2009. An intricate network of regulators controls biofilm formation and
colonization by *Vibrio fischeri*. Mol Microbiol 74:782–789.

7. Ludvik DA, Bultman KM, Mandel MJ. 2021. Hybrid histidine kinase BinK represses *Vibrio*
fischeri biofilm signaling at multiple developmental stages. J Bacteriol 203:e0015521.

8. Ruby EG, Asato LM. 1993. Growth and flagellation of *Vibrio fischeri* during initiation of the
sepiolid squid light organ symbiosis. Arch Microbiol 159:160–167.

9. Yip ES, Grublesky BT, Hussa EA, Visick KL. 2005. A novel, conserved cluster of genes
promotes symbiotic colonization and σ^{54}-dependent biofilm formation by *Vibrio fischeri*. Mol

Microbiol 57:1485–1498.

10. Koehler S, Gaedeke R, Thompson C, Bongrand C, Visick KL, Ruby E, McFall-Ngai M.

2019. The model squid-vibrio symbiosis provides a window into the impact of strain- and

species-level differences during the initial stages of symbiont engagement. Environ

Microbiol 21:3269–3283.

11. Rotman ER, Bultman KM, Brooks JF 2nd, Gyllborg MC, Burgos HL, Wollenberg MS,

Mandel MJ. 2019. Natural strain variation reveals diverse biofilm regulation in squid-

colonizing *Vibrio fischeri*. J Bacteriol 201:e00033–19.

12. Nawroth JC, Guo H, Koch E, Heath-Heckman EAC, Hermanson JC, Ruby EG, Dabiri JO,

Kanso E, McFall-Ngai M. 2017. Motile cilia create fluid-mechanical microhabitats for the

active recruitment of the host microbiome. Proc Natl Acad Sci U S A 114:9510–9516.

13. Thompson CM, Tischler AH, Tarnowski DA, Mandel MJ, Visick KL. 2019. Nitric oxide

inhibits biofilm formation by *Vibrio fischeri* via the nitric oxide sensor HnoX. Mol Microbiol

111:187–203.

14. Graf J, Dunlap PV, Ruby EG. 1994. Effect of transposon-induced motility mutations on

colonization of the host light organ by *Vibrio fischeri*. J Bacteriol 176:6986–6991.

15. Millikan DS, Ruby EG. 2002. Alterations in *Vibrio fischeri* motility correlate with a delay in

symbiosis initiation and are associated with additional symbiotic colonization defects. Appl

Environ Microbiol 68:2519–2528.

16. Mandel MJ, Schaefer AL, Brennan CA, Heath-Heckman EAC, Deloney-Marino CR, McFall-

Ngai MJ, Ruby EG. 2012. Squid-derived chitin oligosaccharides are a chemotactic signal

during colonization by *Vibrio fischeri*. Appl Environ Microbiol 78:4620–4626.

17. Brennan CA, Mandel MJ, Gyllborg MC, Thomasgard KA, Ruby EG. 2013. Genetic

determinants of swimming motility in the squid light-organ symbiont *Vibrio fischeri*.

Microbiologyopen 2:576–594.

18. Sycuro LK, Ruby EG, McFall-Ngai M. 2006. Confocal microscopy of the light organ crypts

in juvenile *Euprymna scolopes* reveals their morphological complexity and dynamic function

in symbiosis. J Morphol 267:555–568.

19. McFall-Ngai MJ, Ruby EG. 1991. Symbiont recognition and subsequent morphogenesis as

early events in an animal-bacterial mutualism. Science 254:1491–1494.

20. Jones BW, Nishiguchi MK. 2004. Counterillumination in the Hawaiian bobtail squid,

Euprymna scolopes Berry (Mollusca: Cephalopoda). Mar Biol 144:1151–1155.

21. Wolfe AJ, Visick KL. 2010. Roles of diguanylate cyclases and phosphodiesterases in

motility and biofilm formation in *Vibrio fischeri*, p. 186–200. *In* Wolfe, AJ, Visick, KL (eds.),

The Second Messenger Cyclic Di-GMP. American Society for Microbiology Press,

Washington, DC.

22. O'Shea TM, Klein AH, Geszvain K, Wolfe AJ, Visick KL. 2006. Diguanylate cyclases control

magnesium-dependent motility of *Vibrio fischeri*. J Bacteriol 188:8196–8205.

23. Christensen DG, Marsden AE, Hodge-Hanson K, Essock-Burns T, Visick KL. 2020. LapG

mediates biofilm dispersal in *Vibrio fischeri* by controlling maintenance of the VCBS-

containing adhesin LapV. Mol Microbiol 114:742–761.

24. Bassis CM, Visick KL. 2010. The cyclic-di-GMP phosphodiesterase BinA negatively

regulates cellulose-containing biofilms in *Vibrio fischeri*. J Bacteriol 192:1269–1278.

25. Brooks JF 2nd, Gyllborg MC, Cronin DC, Quillin SJ, Mallama CA, Foxall R, Whistler C,

Goodman AL, Mandel MJ. 2014. Global discovery of colonization determinants in the squid

symbiont *Vibrio fischeri*. Proc Natl Acad Sci U S A 111:17284–17289.

26. Visick KL, Hodge-Hanson KM, Tischler AH, Bennett AK, Mastrodomenico V. 2018. Tools

for rapid genetic engineering of *Vibrio fischeri*. Appl Environ Microbiol 84:e00850–18.

27. Shrestha P, Razvi A, Fung BL, Eichinger SJ, Visick KL. Mutational analysis of *Vibrio*

fischeri c-di-GMP-modulating genes reveals complex regulation of motility. J Bacteriol.

28. Zamorano-Sánchez D, Xian W, Lee CK, Salinas M, Thongsomboon W, Cegelski L, Wong

GCL, Yildiz FH. 2019. Functional Specialization in *Vibrio cholerae* Diguanylate Cyclases:

Distinct Modes of Motility Suppression and c-di-GMP Production. MBio 10:e00670–19.

29. Naughton LM, Mandel MJ. 2012. Colonization of *Euprymna scolopes* squid by *Vibrio*

fischeri. J Vis Exp e3758.

30. Waters CM, Lu W, Rabinowitz JD, Bassler BL. 2008. Quorum sensing controls biofilm

formation in *Vibrio cholerae* through modulation of cyclic di-GMP levels and repression of

vpsT. J Bacteriol 190:2527–2536.

31. Spurbeck RR, Tarrien RJ, Mobley HLT. 2012. Enzymatically active and inactive

phosphodiesterases and diguanylate cyclases are involved in regulation of Motility or

sessility in *Escherichia coli* CFT073. MBio 3:e00307–12.

32. O'Shea TM, Deloney-Marino CR, Shibata S, Aizawa S-I, Wolfe AJ, Visick KL. 2005.

Magnesium promotes flagellation of *Vibrio fischeri*. J Bacteriol 187:2058–2065.

33. Tischler AH, Lie L, Thompson CM, Visick KL. 2018. Discovery of calcium as a biofilm-

promoting signal for *Vibrio fischeri* reveals new phenotypes and underlying regulatory

complexity. J Bacteriol 200:e00016–18.

34. Morris AR, Darnell CL, Visick KL. 2011. Inactivation of a novel response regulator is

necessary for biofilm formation and host colonization by *Vibrio fischeri*. Mol Microbiol

82:114–130.

35. Brooks JF 2nd, Mandel MJ. 2016. The histidine kinase BinK Is a negative regulator of

biofilm formation and squid colonization. J Bacteriol 198:2596–2607.

36. Nyholm SV, McFall-Ngai MJ. 2021. A lasting symbiosis: how the Hawaiian bobtail squid

finds and keeps its bioluminescent bacterial partner. Nat Rev Microbiol 19:666–679.

37. Hall CL, Lee VT. 2018. Cyclic-di-GMP regulation of virulence in bacterial pathogens. Wiley

Interdiscip Rev RNA 9.

38. Petersen E, Chaudhuri P, Gourley C, Harms J, Splitter G. 2011. *Brucella melitensis* cyclic

di-GMP phosphodiesterase BpdA controls expression of flagellar genes. J Bacteriol

193:5683–5691.

39. Tamayo R, Schild S, Pratt JT, Camilli A. 2008. Role of cyclic Di-GMP during El Tor biotype

Vibrio cholerae infection: characterization of the in vivo-induced cyclic Di-GMP

phosphodiesterase CdpA. Infect Immun 76:1617–1627.

40. Bobrov AG, Kirillina O, Ryjenkov DA, Waters CM, Price PA, Fetherston JD, Mack D,

Goldman WE, Gomelsky M, Perry RD. 2011. Systematic analysis of cyclic di-GMP

signalling enzymes and their role in biofilm formation and virulence in *Yersinia pestis*. Mol

Microbiol 79:533–551.

41. Robinson CD, Sweeney EG, Ngo J, Ma E, Perkins A, Smith TJ, Fernandez NL, Waters CM,

Remington SJ, Bohannan BJM, Guillemin K. 2021. Host-emitted amino acid cues regulate

bacterial chemokinesis to enhance colonization. Cell Host Microbe 29:1221–1234.e8.

42. Chun CK, Troll JV, Koroleva I, Brown B, Manzella L, Snir E, Almabrazi H, Scheetz TE,

Bonaldo M de F, Casavant TL, Soares MB, Ruby EG, McFall-Ngai MJ. 2008. Effects of

colonization, luminescence, and autoinducer on host transcription during development of

the squid-vibrio association. Proc Natl Acad Sci U S A 105:11323–11328.

43. Troll JV, Adin DM, Wier AM, Paquette N, Silverman N, Goldman WE, Stadermann FJ,

Stabb EV, McFall-Ngai MJ. 2009. Peptidoglycan induces loss of a nuclear peptidoglycan

recognition protein during host tissue development in a beneficial animal-bacterial

symbiosis. Cell Microbiol 11:1114–1127.

44. Troll JV, Bent EH, Pacquette N, Wier AM, Goldman WE, Silverman N, McFall-Ngai MJ.

2010. Taming the symbiont for coexistence: a host PGRP neutralizes a bacterial symbiont

toxin. Environ Microbiol 12:2190–2203.

45. McFall-Ngai M, Nyholm SV, Castillo MG. 2010. The role of the immune system in the

initiation and persistence of the *Euprymna scolopes–Vibrio fischeri* symbiosis. Semin

Immunol 22:48–53.

46. Burgos HL, Burgos EF, Steinberger AJ, Suen G, Mandel MJ. 2020. Multiplexed competition

in a synthetic squid light organ microbiome using barcode-tagged gene deletions.

mSystems 5:e00846–20.

47. Brennan CA, DeLoney-Marino CR, Mandel MJ. 2013. Chemoreceptor VfcA mediates amino

acid chemotaxis in *Vibrio fischeri*. Appl Environ Microbiol 79:1889–1896.

48. Nikolakakis K, Monfils K, Moriano-Gutierrez S, Brennan CA, Ruby EG. 2015.

Characterization of the *Vibrio fischeri* Fatty Acid Chemoreceptors, VfcB and VfcB2. Appl

Environ Microbiol 82:696–704.

49. Bennett BD, Essock-Burns T, Ruby EG. 2020. HbtR, a Heterofunctional Homolog of the

Virulence Regulator TcpP, Facilitates the Transition between Symbiotic and Planktonic

Lifestyles in *Vibrio fischeri*. MBio 11:e01624–20.

50. Hayes CA, Dalia TN, Dalia AB. 2017. Systematic genetic dissection of PTS in *Vibrio*

cholerae uncovers a novel glucose transporter and a limited role for PTS during infection of

a mammalian host. Mol Microbiol 104:568–579.

51. Millikan DS, Ruby EG. 2003. FlrA, a σ^{54}-Dependent Transcriptional Activator in *Vibrio*

fischeri, Is Required for Motility and Symbiotic Light-Organ Colonization. J Bacteriol

185:3547–3557.

52. Dial CN, Speare L, Sharpe GC, Gifford SM, Septer AN, Visick KL. 2021. Para-

Aminobenzoic Acid, Calcium, and c-di-GMP Induce Formation of Cohesive, Syp-

Polysaccharide-Dependent Biofilms in *Vibrio fischeri*. MBio 12:e0203421.

53. Colvin KM, Irie Y, Tart CS, Urbano R, Whitney JC, Ryder C, Howell PL, Wozniak DJ,

Parsek MR. 2012. The Pel and Psl polysaccharides provide *Pseudomonas aeruginosa*

structural redundancy within the biofilm matrix. Environ Microbiol 14:1913–1928.

54. Stabb EV, Ruby EG. 2002. RP4-based plasmids for conjugation between *Escherichia coli*

and members of the Vibrionaceae. Methods Enzymol 358:413–426.

55. Christensen DG, Visick KL. 2020. *Vibrio fischeri*: Laboratory Cultivation, Storage, and

Common Phenotypic Assays. Curr Protoc Microbiol 57:e103.

56. Pollack-Berti A, Wollenberg MS, Ruby EG. 2010. Natural transformation of *Vibrio fischeri*

requires *tfoX* and *tfoY*. Environ Microbiol 12:2302–2311.

57. Boettcher KJ, Ruby EG. 1990. Depressed light emission by symbiotic *Vibrio fischeri* of the

sepiolid squid *Euprymna scolopes*. J Bacteriol 172:3701–3706.

58. Mandel MJ, Stabb EV, Ruby EG. 2008. Comparative genomics-based investigation of

resequencing targets in *Vibrio fischeri*: focus on point miscalls and artefactual expansions.

BMC Genomics 9:138.

59. Tischler AH, Vanek ME, Peterson N, Visick KL. 2021. Calcium-Responsive Diguanylate

Cyclase CasA Drives Cellulose-Dependent Biofilm Formation and Inhibits Motility in *Vibrio*

fischeri. MBio 12:e0257321.

60. Dial CN, Eichinger SJ, Foxall R, Corcoran CJ, Tischler AH, Bolz RM, Whistler CA, Visick

KL. 2021. Quorum Sensing and Cyclic di-GMP Exert Control Over Motility of *Vibrio fischeri*

KB2B1. Front Microbiol 12:690459.

61. Cohen JJ, Eichinger SJ, Witte DA, Cook CJ, Fidopiastis PM, Tepavčević J, Visick KL. 2021.

Control of Competence in *Vibrio fischeri*. Appl Environ Microbiol 87:e01962–20.

62. Dunn AK, Millikan DS, Adin DM, Bose JL, Stabb EV. 2006. New *rfp*- and pES213-derived

tools for analyzing symbiotic *Vibrio fischeri* reveal patterns of infection and *lux* expression in

situ. Appl Environ Microbiol 72:802–810.

 FIGURE LEGENDS

FIG 1 High c-di-GMP levels inhibit swimming motility, promote Congo red binding, and promote

cellulose gene transcription. (A) Quantification of c-di-GMP concentration for indicated *V.*

fischeri strains in LBS and filter-sterilized Instant Ocean (FSIO) using the pFY4535 c-di-GMP

reporter plasmid. Constitutive AmCyan was used to normalize GFP to cell density. For each

strain, n = 3 biological and n = 3 technical replicates per biological replicate, each point

represents the mean of technical replicates, average bars represent the mean of biological

replicates, and error bars represent standard error of the mean. (B) Quantification of migration

through soft (0.3%) agar for indicated *V. fischeri* strains. For each strain, n = 8-9 biological

replicates, average bars represent mean, and error bars represent standard error of the mean.

Motility plate images are representative. (C) Quantification of Congo red binding for indicated *V.*

fischeri strains. For each strain, n = 3 technical replicates, average bars represent mean, and

error bars represent standard error of the mean. Congo red spot images are representative. (D)

Quantification of *in vitro bcs* transcription by indicated *V. fischeri* strains using the pRYI063

bcsQ'-gfp⁺ transcriptional reporter plasmid. For each strain, n = 3 biological replicates. Points

represent each biological replicate, average bars represent the mean, and error bars represent

standard error of the mean. Constitutive mCherry was used to normalize GFP to cell density.

For panels A-D, one-way ANOVA was used for statistical analysis and asterisks represent

significance relative to the WT strain; ns = not significant, *$p < 0.05$, **$p < 0.006$, ****$p < 0.0001$.

FIG 2 Elevated c-di-GMP levels inhibit host colonization. (A) Quantification of squid colonization

levels at 18 hpi by indicated *V. fischeri* strains and aposymbiotic (Apo) control. Sample sizes

from left to right are 8, 14, 25, and 13 squid. (B) Quantification of squid colonization levels at 48

hpi by indicated *V. fischeri* strains and aposymbiotic (Apo) control. Sample sizes from left to

right are 23, 68, 75, and 65 squid. For panels A and B, box-and-whisker plots represent

minimum, 25th percentile, median, 75th percentile, and maximum. Kruskal-Wallis test was

performed for statistical analysis for squid that were introduced to bacteria; ns = not

significant,**p < 0.002, ****p < 0.0001.

FIG 3 Reduced swimming motility, increased Congo red binding, and diminished squid

colonization of the High cdG strain are dependent on high c-di-GMP levels. (A) Quantification of

c-di-GMP concentration for indicated *V. fischeri* strains using the pFY4535 c-di-GMP reporter

plasmid. For each strain, $n = 3$ biological and $n = 3$ technical replicates, each dot represents the

mean of technical replicates, average bars represent the mean of biological replicates, and error

bars represent standard error of the mean. (B) Quantification of migration through TBS soft

(0.3%) agar for indicated *V. fischeri* strains. For each strain, $n = 5$ biological replicates, average

bars represent mean, and error bars represent standard error of the mean. (C) Quantification of

Congo red binding for indicated *V. fischeri* strains. For each strain, $n = 3$ technical replicates,

average bars represent the mean, and error bars represent standard error of the mean. Congo

red spot images are representative. For panels A-C, one-way ANOVA was used for statistical

analysis; ns = not significant, *p = 0.017, ****p < 0.0001. (D) Quantification of squid colonization

levels at 48 hpi by indicated *V. fischeri* strains and aposymbiotic (Apo) control. Box-and-whisker

plots represent minimum, 25th percentile, median, 75th percentile, and maximum. Sample sizes

from left to right are 30, 26, 34, 37, and 11 squid. Kruskal-Wallis test was performed for

statistical analysis for squid that were introduced to bacteria; ns = not significant, ****p < 0.0001.

FIG 4 High c-di-GMP levels do not eliminate motility during host colonization. Quantification of

squid competitive colonization index at 48 hpi by indicated *V. fischeri* strains. Competitive index

represents $\log_{10}$((test strain/High cdG)$_{output}$ / (test strain/High cdG)$_{input}$). Box-and-whisker plots

represent minimum, 25th percentile, median, 75th percentile, and maximum. Sample sizes from

left to right are 43 and 75 squid. Kruskal-Wallis test was performed for statistical analysis; ****p

< 0.0001.

FIG 5 Elevated c-di-GMP levels lead to a greater number of *V. fischeri* aggregates in the squid

host. (A) Representative fluorescence microscopy images of aggregates made by indicated *V.*

fischeri strains carrying a constitutive GFP on the pVSV102 plasmid within the host mucus.

Arrows indicate location of aggregates. (B) Quantification of aggregate area per squid for

indicated *V. fischeri* strains carrying a constitutive GFP on the pVSV102 plasmid. Zero values

are displayed at the bottom of the plot. Box-and-whisker plots represent minimum, 25th

percentile, median, 75th percentile, and maximum. Sample sizes from left to right are 73, 131,

101, 38, and 12 squid. Kruskal-Wallis test was used for statistical analysis; ns = not significant,

*p < 0.04, **p < 0.02, ****p < 0.0001. (C) Quantification of number of aggregates formed per

squid by indicated *V. fischeri* strains. Sample sizes from left to right are 73, 131, and 101 squid.

FIG 6 c-di-GMP downregulates *syp* via BcsA in host mucus but not in culture. (A) Quantification

of *syp* transcription for indicated *V. fischeri* strains using the pM1422 *sypA'-gfp*⁺ transcriptional

reporter plasmid in aggregates within the host mucus. Sample sizes from left to right are 47, 40,

and 22 aggregates. (B) Quantification of *bcs* transcription for indicated *V. fischeri* strains using

the pRYI063 *bcsQ'-gfp*⁺ transcriptional reporter plasmid in aggregates within the host mucus.

Sample sizes from left to right are 19, 24 and 28 aggregates. For panels A and B, Box-and-

whisker plots represent minimum, 25th percentile, median, 75th percentile, and maximum.

Kruskal-Wallis test was used for statistical analysis; ns = not significant, ****p < 0.0001. (C)

Quantification of *in vitro syp* transcription in LBS and TBS-Ca^{2+} for indicated *V. fischeri* strains

carrying the pM1422 *sypA'-gfp*⁺ transcriptional reporter plasmid. For each strain, *n* = 3 biological

replicates. Points represent each replicate, average bars represent the mean of replicates, and

error bars represent the standard error of the mean. One-way ANOVA was used for statistical

analysis; ns = not significant relative to the WT strain in the same media. For panels A-C,

constitutive mCherry was used to normalize GFP to cell density.

FIG S1 Low levels of c-di-GMP do not inhibit host colonization. Quantification of squid

competitive colonization index at 48 hpi by indicated *V. fischeri* strains. Competitive index

represents $\log_{10}((\text{test strain/WT})_{\text{output}} / (\text{test strain/WT})_{\text{input}})$. Box-and-whisker plots represent

minimum, 25th percentile, median, 75th percentile, and maximum. Sample sizes from left to

right are 36, 26, and 20 squid. Kruskal-Wallis test was performed for statistical analysis; ns = sot

significant, ****$p < 0.0001$.

FIG S2 The High c-di-GMP *V. fischeri* strain does not have a growth defect. (A) Growth curves

for *V. fischeri* and indicated mutants. For each strain, $n = 3$ biological and $n = 6$ technical

replicates per biological replicate. Points represent the mean of technical replicates. Error bars

represent standard error of the mean. (B) Doubling times for *V. fischeri* and indicated mutants.

Average bars represent the mean of biological replicates. Points represent the mean of

technical replicates. Error bars represent standard error of the mean. One-way ANOVA was

used for statistical analysis; ns = not significant. Doubling time was calculated using a nonlinear

regression analysis of each strain during exponential growth phase. For panels A and B, outliers

were excluded from analysis.

FIG S3 High cdG motile cells are not suppressor mutants. Representative image of migration

through soft (0.3%) agar for *V. fischeri* and indicated strains.

FIG S4 Multiple gene deletions alter *in vivo* c-di-GMP levels. (A) Quantification of c-di-GMP

levels for indicated *V. fischeri* strains using the pFY4535 c-di-GMP reporter plasmid in

aggregates within the host mucus. Samples sizes from left to right are 9 and 5 aggregates. A

Mann-Whitney test was used for statistical analysis; ***p ≤ 0.01. (B) Representative fluorescent

microscopy images of squid light organs containing indicated *V. fischeri* strains carrying the

pFY4535 c-di-GMP reporter plasmid. Arrows indicate location of aggregates.

FIG S5 Absence of the cellulose synthase BcsA does not rescue the high c-di-GMP strain

colonization defect. (A) Quantification of squid colonization levels at 18 hpi by indicated *V.*

fischeri strains and aposymbiotic (Apo) control. Sample sizes from left to right are 14, 25, 17,

and 13 squid. (B) Quantification of squid colonization levels at 48 hpi by indicated *V. fischeri*

strains. Sample sizes from left to right are 68, 16, 75, 56, 17, and 65 squid. For panels A and B,

data for WT, High cdG, and Apo groups are the same as in Fig 3. Box-and-whisker plots

represent minimum, 25th percentile, median, 75th percentile, and maximum. Kruskal-Wallis test

was performed for statistical analysis for squid that were introduced to bacteria; ns = not

significant, **p = 0.002, ****p < 0.0001.

FIG S6 High c-di-GMP-dependent Bcs expression inhibits *syp* expression in bacterial

aggregates within the host mucus. (A) Representative fluorescent microscopy images of squid

light organs containing indicated *V. fischeri* strains carrying the pM1422 *sypA*'-*gfp*[+]

transcriptional reporter plasmid. (B) Representative fluorescent microscopy images of squid light

organs containing indicated *V. fischeri* strains carrying the pRYI063 *bcsQ*'-*gfp*[+] transcriptional

reporter plasmid.

A 18 hpi

B 48 hpi

A

Low cdG | WT | High cdG

High cdG *bcsA*::Tn*erm* | High cdG Δ*sypQ* | Apo

200 μM

B

Aggregate area per squid (μm²)

Low cdG | WT | High cdG | High cdG *bcsA*::Tn*erm* | High cdG Δ*sypQ*

C

Percent of squid

Low cdG | WT | High cdG

aggregates per squid