When blue turns to grey - Paleogenomic insights into the evolutionary history and extinction of the blue antelope (*Hippotragus leucophaeus*)

Elisabeth Hempel*[,1,2], Faysal Bibi[2], J. Tyler Faith[3,4,5], Klaus-Peter Koepfli[6,7], Achim M. Klittich[1], David A. Duchêne[8], James S. Brink[9,10,†], Daniela C. Kalthoff[11], Love Dalén[11,12], Michael Hofreiter[#1] and Michael V. Westbury[#8]

[1] Universität Potsdam, Evolutionary Adaptive Genomics, Karl-Liebknecht-Str. 24-25, 14476 Potsdam, Germany
[2] Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Invalidenstraße 43, 10115 Berlin, Germany
[3] Natural History Museum of Utah, University of Utah, 301 Wakara Way, Salt Lake City, UT 84108, USA
[4] Department of Anthropology, University of Utah, 260 South Central Campus Drive, Salt Lake City, UT 84112, USA
[5] Origins Centre, University of the Witwatersrand, Johannesburg, Republic of South Africa
[6] Smithsonian-Mason School of Conservation, George Mason University, Front Royal, VA 22630, USA
[7] Smithsonian Conservation Biology Institute, Center for Species Survival, National Zoological Park, Front Royal, VA 22630, USA
[8] The GLOBE Institute, University of Copenhagen, Øster Voldgade 5-7, Copenhagen, Denmark
[9] National Museum Bloemfontein, Florisbad Quaternary Research Station and Department, P.O. Box 266, Bloemfontein, 9031, Republic of South Africa
[10] Centre for Environmental Management, University of the Free State, PO Box 339, Bloemfontein 9300, Republic of South Africa
[11] Swedish Museum of Natural History, Box 50007, 10405 Stockholm, Sweden
[12] Centre for Palaeogenetics, Svante Arrhenius väg 20c, 10691 Stockholm, Sweden

† deceased
[#] shared senior authorship

*Corresponding author: Elisabeth Hempel; Hempel.Elisabeth@posteo.org

Abstract

The blue antelope (*Hippotragus leucophaeus*) is the only large African mammal to have become extinct in historical times, yet no nuclear genomic information is available for this species. A recent study showed that many alleged blue antelope museum specimens are either roan (*H. equinus*) or sable (*H. niger*) antelopes, further reducing the possibilities for obtaining genomic information for this extinct species. While the blue antelope has a rich fossil record from South Africa, climatic conditions in the region are unfavourable to the preservation of ancient DNA. Nevertheless, we recovered two blue antelope draft genomes, one at 3.4x coverage from a historical specimen (~19th century) and one at 2.1x coverage from a fossil specimen dating to 9,800–9,300 cal BP, making it currently the oldest paleogenome from Africa. Phylogenomics show that blue and sable antelope are sister species, confirming previous mitogenomic results, and demonstrate ancient gene flow from roan into blue antelope. We show that blue antelope genomic diversity was much lower than in roan and sable antelopes, indicative of a low population size since at least the early Holocene. This supports observations from the fossil record documenting major decreases in the abundance of blue antelope after the Pleistocene-Holocene transition. Finally, the persistence of this species throughout the Holocene despite low population size suggests that colonial-era human impact was likely a major factor in the blue antelope's extinction.

key words: ancient DNA, diversity, extinction, Holocene, paleogenome, South Africa

Introduction

Earth is currently undergoing a major biodiversity crisis that is a direct result of human activities (Ceballos et al. 2015). The consequences for many species include considerable reductions in population size and associated losses of genetic diversity that can lead to reduced fitness and adaptability. Today, through the continuing development of next generation sequencing technologies and advancement of ancient DNA (aDNA) techniques, it is feasible to retrieve

genomic information from extinct and non-model organism species (Westbury et al. 2020; Barlow et al. 2021; Sánchez-Barreiro et al. 2021), thereby facilitating inferences about threats facing many present-day species. One especially powerful use of these genomes is to reconstruct the evolutionary history of species in relation to environmental change and human impacts. In this regard, Africa is hugely understudied, likely due to its environmental conditions, with high temperatures being detrimental to DNA preservation (Smith et al. 2001; Bollongino et al. 2008; Hofreiter et al. 2015). To date, most aDNA studies on African samples have been restricted to DNA enrichment approaches and focused on human remains (Skoglund et al. 2017; Vicente and Schlebusch 2020; Lipson et al. 2022), with very few investigating other fossil fauna (e.g. Mathieson et al. 2020).

One species that likely fell victim to human impacts during the beginning of the current biodiversity crisis is the blue antelope, *Hippotragus leucophaeus* (Pallas, 1766). The blue antelope belongs to the bovid tribe Hippotragini, which comprises the extant genera *Hippotragus*, *Oryx* and *Addax*. It went extinct ~1800 AD (Lichtenstein 1811, 1814) and therefore represents the first – and so far only – historic extinction of a large African mammal species (Harper 1945). It had a distinct white patch in front of the eyes (Pallas 1767), and its pelt was perceived as bluish while alive (Lichtenstein 1814; von Schreber and Goldfuß 1836) – perhaps similar to the blue wildebeest (*Connochaetes taurinus*) or the nilgai (*Boselaphus tragocamelus)* (Mohr 1967) – turning greyish post-mortem (Lichtenstein 1814; von Schreber and Goldfuß 1836) (fig. 1A). Like its extant relatives, the blue antelope was a grazer (Mohr 1967; Klein 1974, 1987). Historically, the blue antelope was endemic to a very small area (~4,300 km^2) between Swellendam, Caledon, and Bredasdorp in South Africa (Skead 1980; Kerley et al. 2009). However, its Quaternary fossil record demonstrates a broader prehistoric range (fig. 1C), with fossil occurrences throughout the Cape Floristic Region and extending into the highlands of Lesotho (Klein 1974; Plug 1997; Faith and Thompson 2013; Avery 2019). Blue antelopes are particularly ubiquitous and abundant in Pleistocene fossil

assemblages from South Africa's Western Cape Province, contracting in range and abundance at the onset of the Holocene (Faith 2011).

The blue antelope's extinction is hypothesised to have been the result of several anthropogenic causal factors, including landscape transformation (Faith and Thompson 2013), overhunting by European colonists (FitzSimons 1920; Harper 1945), competition with and habitat deterioration by livestock (Klein 1974, 1987), and disruption of migratory pathways in prehistoric and colonial times (Faith and Thompson 2013). Additionally, the potentially detrimental role of stochastic processes on a small population size has also been considered (Kerley et al. 2009). Unfortunately, due to its early demise, our sole sources of knowledge of the blue antelope's ecology and evolutionary history has been limited to information from the fossil record (Klein 1974, 1987; Faith 2011; Faith and Thompson 2013) and historical mitochondrial data (Robinson et al. 1996; Themudo and Campos 2018; Hempel et al. 2021a). The relationships among the three extant *Hippotragus* species, the roan (*H. equinus*), the sable (*H. niger*) and the blue antelope, have so far only been investigated using mitochondrial sequences, leaving the phylogeny unresolved at the nuclear genome level. Considering that mitochondrial DNA data can sometimes present an incomplete picture of the evolutionary history of a species and can be confounded by past gene flow events (e.g. Larsen et al. 2010; Edwards et al. 2011; Westbury et al. 2020), nuclear data is necessary to conclusively resolve phylogenetic relationships.

Here, we present the nuclear genome of a ~200 year old blue antelope specimen as well as a paleogenome from an early Holocene fossil specimen (~9,300–9,800 years BP), which, to our knowledge, currently constitutes the oldest paleogenome from Africa. Using these genomes, we investigate the phylogenomic relationships of the blue antelope and show that the blue antelope was more closely related to the sable than to the roan antelope, but with detectable past gene flow between the roan and the blue antelope. Furthermore, we demonstrate that the blue antelope had very low diversity compared to its congeners, likely since at least the beginning of the Holocene.

FɪG. 1. Blue antelope specimens and distribution (**A**) Historical mounted skin of a young male blue antelope (*Hippotragus leucophaeus*) from the Swedish Museum of Natural History, Stockholm, Sweden (NRM 590107); arrow indicates area sampled for aDNA extraction (**B**) Fossil lower deciduous 4th premolar (NBC RB4 D3) of a blue antelope from Nelson Bay Cave (curated at Iziko Museums of South Africa, Cape Town, Republic of South Africa); scale is one centimetre; arrows indicate areas sampled for aDNA extraction (**C**) Historical distribution (light purple at the southern tip) and Holocene and Pleistocene fossil sites (Skead 1980; Kerley et al. 2009; modified from Avery 2019) (base map: https://www.naturalearthdata.com, prepared in QGIS v2.18 (https://qgis.org)). Photo credits: NBC RB4 D3: J.T. Faith, courtesy: Archaeology Unit, Iziko Museums of South Africa; NRM 590107: Hempel et al. 2021a.

Results

Using shotgun sequencing, we obtained both a nuclear and a mitochondrial genome from a fossil specimen from Nelson Bay Cave (NBC RB4 D3) (Klein 1972, 1983) dating to 9,300–9,800 cal BP (Loftus et al. 2016) at 2.14x and 234.67x coverage, respectively. In addition, we generated a nuclear genome of the ~200 year old historical specimen NRM 590107 from the Swedish Museum of Natural History, Stockholm, Sweden, with a coverage of 3.44x. Details on read numbers and mapping statistics can be found in supplementary tables S2 & S3. MapDamage v2.2.0 (Jónsson et al. 2013) analysis of DNA sequences obtained from the NBC RB4 D3 specimen showed typical patterns of damage (caused by cytosine deamination) for an aDNA sample of its age (supplementary figs. S1A & S1B). The same was true for NRM 590107, but in addition, this sample also showed an elevated level of guanine to thymine transversions (supplementary figs. S2A & S2B).

Phylogenetic relationships

A nuclear genome neighbor-joining phylogeny supported the monophyly of all three *Hippotragus* species, demonstrating them to be distinct lineages and showing that the blue antelope is more closely related to the sable than to the roan antelope (fig. 2A). The mitochondrial genome maximum-likelihood phylogeny was congruent with the nuclear neighbor-joining phylogeny (figs. 2A & 2B). The mitochondrial phylogeny using transversions only (binary format), yielded an identical topology with respect to the grouping of the three species (supplementary fig. S3).

Fig. 2. Phylogenetic analyses (**A**) Neighbor-joining tree of the three *Hippotragus* species based on nuclear genomes, showing the sister relationship of blue and sable antelopes. The phylogeny was constructed using ANGSD v0.923 (Korneliussen et al. 2014) and FastME v2.1.6.1 (Lefort et al. 2015). (**B**) Maximum-likelihood phylogeny of mitochondrial genomes of the three *Hippotragus* species. The phylogeny is congruent with the nuclear phylogeny. The maximum-likelihood phylogeny was generated using RAxML v8.2.12 with a GTR+G substitution model (Stamatakis 2014). The control region was excluded from the alignment (15,440 bp alignment length). Node labels show bootstrap support percentages (1000 replicates). In both analyses, the scimitar-horned oryx (*Oryx dammah*, SB20612, Humble et al. 2020) was used as the outgroup (not shown). Roan and sable antelope raw data are from Gonçalves et al. (2021) and Koepfli et al. (2019). Photo credits: see fig. 1.

Gene flow

Both blue antelope individuals showed significantly higher levels of shared derived alleles, suggestive of gene flow, with the roan antelope than with the sable antelope (fig. 3A, supplementary table S7). Sliding window phylogenetic tree analyses also suggested gene flow between blue and roan antelope (supplementary table S8 & S9). Moreover, by comparing multiple individuals per species, the sliding window analysis showed that gene flow between roan and blue antelope occurred after the split of blue and sable antelope but before the split of the most recent common ancestors of all roan and also the two blue antelope individuals analysed in our study.

FIG. 3. *D statistics* and geographical distribution of the three *Hippotragus* species (**A**) D values and Z-scores of *D* statistics for the fossil (NBC RB4 D3) and historical (NRM 590107) blue antelope specimens showing significant gene flow between blue and roan antelope; analysis performed with Dstat (transversions only version, https://github.com/jacahill/Admixture). (**B**) Tested topology for *D* statistics according to the neighbor-joining tree (fig. 2A) with blue antelope in position 2. The green double-arrow indicates gene flow between blue and roan antelope with unknown directionality. (**C**) Historical distribution and Holocene and Pleistocene fossil sites of blue, sable and roan antelopes; the distributions of the roan and sable antelope further north are not shown (base map: https://www.naturalearthdata.com, prepared in QGIS v2.18 (https://qgis.org) (du Plessis 1969; Kerley et al. 2009; modified from Avery 2019). NBC RB4 D3: J.T. Faith, courtesy: Archaeology Unit, Iziko Museums of South Africa; NRM 590107: Hempel et al. 2021a; roan antelope: Charles J. Sharp, wikimedia commons, CC-BY 4.0; sable antelope: Paulmaz, wikimedia commons, CC-BY 3.0; scimitar-horned oryx: E. Hempel.

To determine the directionality of the gene flow, we utilised the branch lengths between roan and sable antelope in window trees where the roan and blue antelope form sister lineages. It was expected that in case of gene flow from roan into blue antelope the branch lengths

between roan and sable antelope would be in agreement with the species tree topology, whereas gene flow from blue into roan antelope would lead to a higher similarity between roan and sable antelope and thereby to a shorter branch length compared to the species tree topology (fig. 4A). The sliding window analysis with branch length calculations showed that the branch lengths of roan and sable antelope were similar for the inferred species tree topology and the alternative topology placing blue and roan as sister lineages. Assuming that the most frequently found topology is the species tree topology, the next most frequent topology presumably results from both incomplete lineage sorting and/or gene flow. A bimodal distribution would have been expected if there had been any occurrence of gene flow from the blue into the roan antelope. One mode would reflect the relatively ancient species divergence between the roan and the sable, while the other mode would reflect the more recent divergence between the introgressed blue antelope loci and the sable. Since the distributions of branch lengths between roan and sable were unimodal and approximately equal between windows resulting in the species tree topology and the second most frequent topology, the direction of gene flow was most likely from the roan into the blue antelope (fig. 4B).

Fɪɢ. 4. Gene flow direction analysis (**A**) Expected outcome of changes in genetic distances for gene flow (green arrow) from roan into blue antelope and blue into roan antelope. Dashed red lines illustrate the expected distances between roan and sable antelope. (**B**) Genetic distances between roan and sable antelope as a proportion of the tree length performed with WindowTrees v1.0 (https://github.com/achimklittich/WindowTrees). Photo credits: see fig. 3.

Nuclear and mitochondrial diversity

Calling heterozygous positions to estimate the nuclear genomic diversity of the two low coverage genomes was not a reliable option. Therefore, we estimated diversity through an alternative measure. By uniquely evaluating each substitution type, we confirmed elevated cytosine to thymine and guanine to thymine levels found via mapDamage (supplementary figs. S1 & S2). After removing substitutions that could have been caused by DNA damage, our results show that pairwise differences are related to mean heterozygosity in roan and sable antelope. Therefore, we concluded that pairwise differences could be used to reliably estimate species-wide nuclear diversity even when using low coverage genomes (fig. 5, supplementary tables S10-12, supplementary table S16).

Pairwise difference estimates showed that the blue antelope had much lower nuclear diversity than the roan and sable antelopes (fig. 5). The roan antelope showed the highest nuclear diversity among the three *Hippotragus* species, with the sable antelope showing values that were lower than the roan antelopes' but still much higher than those of the blue antelope. Comparing subsampled and not subsampled input data for the pairwise difference estimates showed that coverage can have an influence, but not to the extent that it could result in the substantially reduced nuclear diversity of the blue antelope observed relative to the other species (supplementary fig. S4, supplementary tables S13-15).

FIG. 5. Nuclear diversity analysis. Species-wide nuclear diversity of the three *Hippotragus* species computed from pairwise comparisons (data normalised by dividing by the number of sites used in the analysis - 2,243,953 sites; genomes for heterozygosity subsampled to 4.26x, the lowest sequence coverage in the comparison data set). Each dot represents a pairwise comparison. Lines represent means. Heterozygosity was not estimated for the blue antelope due to low coverage and ancient DNA damage. Photo credits: see fig. 3.

To estimate the mitochondrial genome diversity of the blue antelope, we generated two haplotype networks, one using complete and the other using partial mitochondrial genomes. The removal of all ambiguities/missing data resulted in an alignment length of 16,492 bp for the complete mitochondrial alignment and 6,300 bp for the partial mitochondrial alignment. We found 15 segregating sites and a nucleotide diversity of 0.00061 in the alignment with complete mitochondrial genomes (fig. 6A), and 10 segregating sites and a very similar nucleotide diversity of 0.00067 for the alignment with partial mitochondrial genomes (fig. 6B).

Fig. 6. Blue antelope mitochondrial haplotype networks (**A**) Median-joining network of three complete blue antelope mitochondrial genomes (alignment length 16,492 bp) generated in POPART v1.7 (Bandelt et al. 1999; Leigh and Bryant 2015). Ambiguities/missing data were excluded. Numbers at lines represent mutational steps. Circle sizes correspond to frequency of haplotypes. City names are locations of the museum specimens, GenBank accession numbers are in parentheses. (**B**) Median-joining network of three complete and two partial mitochondrial genomes (alignment length 6,300 bp). The white circle represents an unsampled haplotype. Photo credits: see fig. 1.

Discussion

Using ancient DNA (aDNA) techniques, we sequenced the first nuclear genomes of the blue antelope to study its evolutionary history and genetic diversity and provide insights into the only historical extinction of a large African mammal species to date. The retrieval of aDNA from African fossils is a challenging task because of the prevailing environmental conditions, namely high temperatures, which facilitate DNA degradation (Smith et al. 2001; Bollongino et al. 2008; Hofreiter et al. 2015). For this reason, few studies have succeeded in recovering aDNA from Africa, and the few exceptions were mainly focused on humans and used genomic enrichment approaches (Vicente and Schlebusch 2020; Lipson et al. 2022). Nonetheless, despite such challenges, we were able to successfully sequence a sample that, with an age of 9,300–9,800 years BP, currently represents the oldest paleogenome from Africa. Our success demonstrates that palaeogenomes can be recovered successfully from African sites, setting the stage for future genomic studies from this area. It is worth noting that our fossil

specimen originates from cave deposits in South Africa's southern Cape region, which should have higher potential for aDNA recovery than, for example, fossils from fluvial sediments in tropical Africa (Hofreiter et al. 2015).

Our finding that the blue antelope is more closely related to the sable than to the roan antelope on a nuclear level confirms results from studies based on mitochondrial genome sequences (Robinson et al. 1996; Themudo and Campos 2018; Hempel et al. 2021a). It was not certain if the mitochondrial phylogeny alone could reliably settle the question of the relationships among the three *Hippotragus* species, since it could have been biased by introgression or incomplete lineage sorting as seen in other studies (e.g. Barlow et al. 2018; Rakotoarivelo et al. 2019; Westbury et al. 2020). Moreover, we show that introgression did occur between *Hippotragus* species, implying that there was a realistic possibility for the mitochondrial topology to have been biased by introgression. However, our phylogenomic result confirms the relationships among the recent species of *Hippotragus* on the nuclear level and puts to rest the long-going discussion on whether the blue antelope was a distinct species or not (Smith 1849; Kohl 1886; Mohr 1967).

Both blue antelope specimens showed significant gene flow with the roan antelope relative to that between the roan and the sable antelope. This may look surprising at first, since the extant and historical populations of the roan and the sable antelope overlap considerably. However, the ranges of the roan and the blue antelope overlapped in the southern Cape region during the late Pleistocene and Holocene (fig. 3C, Klein 1972; Faith 2012; Avery 2019).

Interspecific gene flow is known to be more likely when population size is low since the chances of finding a conspecific mating partner are reduced (Hubbs principle) (Hubbs 1955; Jansson et al. 2007; Crossman et al. 2016; Vaz Pinto et al. 2016). Although we were unable to directly date the gene flow event(s), we could show that gene flow occurred after the split of blue and sable antelope but before the last common ancestors of the roan antelope individuals included in this study. Although we did not detect evidence for more recent interspecific gene flow, it may have been biologically possible since roan and blue antelope

have the same evolutionary distance as roan and sable antelope, which are likely still able to produce viable offspring (Robinson and Harley 1995; Vaz Pinto et al. 2016).

The genomic evidence for low diversity and therefore probably low population sizes from the early Holocene confirms observations derived from the fossil record. Fossil evidence from southern Africa suggests that the blue antelope was both widespread and abundant during the last glacial period (~115 to 11.7 ka) (Klein 1987; Faith 2011). It became increasingly rare in the fossil record following the Pleistocene-Holocene transition (Faith 2011), with the long stratified sequence at Nelson Bay Cave – from where our fossil specimen originates – demonstrating that it remained a rare component of the large mammal community throughout the Holocene (Klein 1983). The onset of the population decline of the blue antelope near the Pleistocene-Holocene transition broadly coincides with the extinction of several other large-bodied grazers in southern Africa, including the Cape zebra (*Equus capensis*), long-horn buffalo (*Syncerus antiquus*), and giant wildebeest (*Megalotragus priscus*), among others. These losses have been linked at least in part to declines in the availability and year-round productivity of grassy forage (Klein 1980; Brink and Lee-Thorp 1992; Faith 2014). This is particularly evident in the Cape Floristic Region, where grassy habitats diminished in conjunction with a rise in sea levels and drowning of the Paleo-Agulhas Plain that would have contributed to dramatic habitat loss, disruption of possible migratory routes, and fragmentation of species ranges (Dingle and Rogers 1972; Klein 1983; van Andel 1989; Fisher et al. 2010; Compton 2011; Faith and Behrensmeyer 2013; Faith and Thompson 2013; Copeland et al. 2016; Venter et al. 2020). Rare Holocene occurrences of the blue antelope in the southwestern Cape (Klein and Cruz-Uribe 2016), the southern Cape (Klein 1983), and in the interior (Opperman 1987; Plug 1997), likely represent disjunct populations due to lack of dispersal corridors around the Cape Fold Mountains when sea levels were high (Compton 2011; Faith and Behrensmeyer 2013; Faith and Thompson 2013). It seems that by the onset of historical times, the blue antelope had become restricted to the southern Cape (Klein 1983, but see Loubser et al. 1990), an area that provides limited foraging opportunities for large-bodied

grazers (Klein 1983; Kerley et al. 2009; Faith and Behrensmeyer 2013; Cawthra et al. 2015; Cawthra et al. 2020). Thus, the environmental changes that marked the transition from the Pleistocene to the Holocene are likely to have resulted in a geographically isolated and small population (Skead 1980; Kerley et al. 2009), which conforms with our finding of very low nuclear diversity compared to the two extant *Hippotragus* species that were and are more broadly distributed across southern Africa (figs. 3C & 5).

Low nuclear diversity, and likely low population size have characterised the blue antelope at least since the early Holocene, raising the question of whether the blue antelope was doomed to extinction, or whether it may have been able to survive in the absence of increased human impacts in historical times. As Kerley et al. (2009) pointed out, due to increased susceptibility to drift there might have been an accumulation of mildly deleterious mutations in the small remaining blue antelope population, which might have put the species on a slow path to extinction.

Our finding of low diversity since the early Holocene, suggests that low diversity was the result of climatically induced habitat fragmentation at the Pleistocene-Holocene transition long before livestock was introduced or European colonists started hunting and land transformation. Generally, low diversity is associated with increased extinction risk (DeWoody et al. 2021; Kardos et al. 2021). However, the ability of the blue antelope to survive for at least 10,000 years despite continuously low diversity suggests that low diversity may not have impacted the survivability of the species, similar to other species in southern Africa (Westbury et al. 2018; Sánchez-Barreiro et al. 2021). Cycles of habitat restrictions through changes in sea level in the past may have led to it repeatedly having had limited population size and genetic diversity. Continuously low population sizes may have enabled more efficient purging of highly deleterious mutations by exposing them more often in a homozygous state, meaning that low population size may be an adaptive advantage rather than detriment (Xue et al. 2015; Dussex et al. 2021; Liu et al. 2021). However, hunting with firearms and land transformation

in its restricted habitat during the colonial era (FitzSimons 1920; Harper 1945; Faith and Thompson 2013) may have proved too much for the blue antelope given its low population size, ultimately culminating in its extinction at the hand of man.

In conclusion, our study shows that using genomic data from fossil specimens from Africa opens up new research possibilities to understand evolutionary dynamics. For the blue antelope, we found low population size throughout the Holocene, confirming observations derived from the fossil record. In the future, it would be interesting to see if genomic data from Pleistocene specimens would continue to reflect the patterns seen in the fossil record. Our results suggest that humans in the colonial era caused the extinction of a species that was likely already vulnerable due to habitat loss and range fragmentation at least since the onset of the Holocene.

Materials and Methods

Samples

We obtained tooth root or bone fossil samples from 25 blue antelope (*Hippotragus leucophaeus*) specimens from Archaeology Unit, Iziko Museums of South Africa, Cape Town, Republic of South Africa, ranging between 5,000–71,000 years BP (supplementary table S1). Two samples were undated. In addition, we used the single-stranded library of a historical sample from the Swedish Museum of Natural History in Stockholm, Sweden, from a previous study (Hempel et al. 2021a). Sample BNK1 O25/2 5.2 was radiocarbon-dated at the [14]Chrono Centre, Queens University Belfast, Ireland using CALIB REV 7.0.0 (Stuiver and Reimer 1993, supplementary table S1).

Laboratory procedures

DNA preparation

We extracted DNA from all fossil samples following Dabney et al. (2013), using columns of Roche's High Pure Viral Nucleic Acid Kit for purification. Subsequently, we built single-stranded libraries from these extracts according to Gansauge et al. (2017), employing an additional initial 15 min. incubation step with 0.5 µl USER Enzyme for Uracil removal (modified from Meyer et al. (2012). We used a maximum of 13 ng DNA as input for library construction. To determine the optimal number of amplification cycles for the subsequent dual-indexing PCR, we performed a qPCR (Thermo Scientific PikoReal Real-Time PCR System). Extraction and library blanks were run alongside all samples to check for the presence of contamination. All pre-PCR lab work was conducted in dedicated ancient DNA (aDNA) facilities at the University of Potsdam, Germany. The single-stranded library of the historical sample NRM 590107 was prepared in the same way as described above (Hempel et al. 2021a). All libraries from fossil specimens were test-sequenced on an Illumina NextSeq500 using custom primers (Gansauge and Meyer 2013; Paijmans et al. 2017) at the University of Potsdam to determine their endogenous DNA content. Only one sample from Nelson Bay Cave (NBC RB4 D3, a lower deciduous premolar (dP_4), fig. 1B) yielded sufficient endogenous DNA content for deeper sequencing. Therefore, we sequenced the library of NBC RB4 D3 using custom primers in one run on an Illumina NextSeq500 at the University of Potsdam, generating 75 bp single-end reads, and in another run on a NovaSeq6000 at the SciLifeLab, Sweden, generating 100 bp paired-end reads. We further sequenced the library of NRM 590107 using custom primers in three runs on an Illumina NextSeq500 at the University of Potsdam, generating 75 bp single-end reads, and on one run on a NovaSeq6000 at the SciLifeLab, generating 100 bp paired-end reads. In addition, for NRM 590107, we used 75 bp single-end read data generated from one run in Hempel et al. (2021a) (SRR number, supplementary table S2).

Bioinformatic procedures and analyses

Nuclear genome

Data preprocessing and read mapping

We processed single- and paired-end reads separately prior to duplicate removal but combined all single-end runs of NRM 590107 before processing. We used Cutadapt v.2.8 (Martin 2011) to trim Illumina adapter sequences (1 bp overlap) and to remove reads shorter than 30 bp. We merged paired-end reads with FLASH v.1.2.11 (Magoč and Salzberg 2011) using a maximum overlap of 100 bp and discarded all unmerged reads. Subsequently, we mapped the resulting reads to the nuclear genome of the scimitar-horned oryx (*Oryx dammah*) (https://www.dnazoo.org/assemblies/Oryx_dammah, Humble et al. 2020) using the BWA aln algorithm v.0.7.17 (Li and Durbin 2009) and default settings. For quality filtering, we used SAMtools view v.1.10 (Li et al. 2009) to remove reads with a mapping quality of <30, discarding unmapped reads, and then sorted bam files with SAMtools sort. We then merged all reads from a single individual with SAMtools merge and performed duplicate removal with Picard MarkDuplicates v2.22.4 (Picard Toolkit 2020 - http://broadinstitute.github.io/picard). We ran mapDamage v2.2.0 (Jónsson et al. 2013) and rescaled the bam file using the rescale option (--rescale) to decrease the quality of misincorporations that are likely caused by aDNA damage.

For comparison we used published raw sequencing read data of three roan (*Hippotragus equinus*) (Gonçalves et al. 2021), eight sable antelopes (*Hippotragus niger*) (Koepfli et al. 2019) from contemporary samples, and the scimitar-horned oryx (*Oryx dammah*) (Humble et al. 2020) (supplementary table S4). The reads for these samples were treated in the same way as described above with the exception that merged and unmerged reads were both mapped and that the maximum overlap parameter in FLASH v.1.2.11 was adjusted according to sequencing cycle length. For one roan antelope sample (10954) and the scimitar-horned oryx sample (SB20612), both from 10X Genomics libraries, 22 bp were trimmed from R1 after

adapter trimming with Cutadapt v.2.8 using FASTA/Q Trimmer from the FASTX-Toolkit v0.0.14 (http://hannonlab.cshl.edu/fastx_toolkit).

Since sex chromosomes and mitochondrial genomes have different inheritance patterns relative to autosomes and could bias downstream analyses, we removed 53 scaffolds that were identified to likely represent X and Y chromosomes and the mitochondrial genome in the scimitar-horned oryx reference genome (Humble et al. 2020; Hempel et al. 2021b).

As input fasta files for *D* statistics (Green et al. 2010; Durand et al. 2011) and the nuclear diversity estimates, we generated pseudohaploid consensus sequences for all specimens using Consensify (Barlow et al. 2020, https://github.com/jlapaijmans/Consensify). We generated the base count input file using ANGSD v0.923 (Korneliussen et al. 2014) (-minQ 30, -minMapQ 30, -uniqueonly 1, -remove_bads 1, -baq 2, -dumpCounts 3, -C 0, -only_proper_pairs 0, -doCounts 1, -trim 0). For the fossil and historical blue antelope specimens, the rescaled bam files from mapDamage were used as input files. We ran Consensify with twice the average sequence coverage of each sample as the maxDepth parameter and using only autosomal scaffolds larger than 1 Mb. To check if sequence coverage had an effect on downstream results, we generated an additional set of subsampled files for all individuals (fossil, historical and contemporary) with SAMtools view v1.10, setting the baseline to the sample with the lowest sequence coverage (2.14x, NBC RB4 D3) before running Consensify.

For the sliding window tree analyses we generated pseudohaploid consensus sequences using random read sampling (-doFasta 1) in ANGSD and the following parameter settings: -minQ 30, -minMapQ 30, -uniqueonly 1, -only_proper_pairs 0, -remove_bads 1, -baq 2, -C 0, -explode 1, -doCounts 1, -trim 0, -basesPerLine 70. We only included autosomal scaffolds larger than 1 Mb (supplementary table S6). For the fossil and historical specimens of blue antelope, the rescaled bam files from mapDamage were used as input files.

Species-wide nuclear diversity comparisons

We performed within species pairwise comparisons to estimate the species-wide nuclear diversity for each species within the genus *Hippotragus*. We implemented an approach that we show can be used for low coverage data and take the patterns of unusual DNA damage found in blue antelope specimen NRM 590107 into account.

We used the fasta files generated with Consensify from the two blue, three roan, and eight sable antelope samples, and the single outgroup scimitar-horned oryx, and filtered those by excluding positions with missing data and positions where all individuals had the same allele (uninformative sites) using a custom perl script. We then performed pairwise difference estimates for the individuals of each species by counting the number of different alleles while excluding all potential aDNA damage (transitions) and all differences resembling the damage pattern found in NRM 590107 (guanine to thymine and cytosine to adenine and vice versa). The values were normalised by dividing them by the total number of positions after filtering. We plotted the values for each pairwise comparison together with the mean for each species using RStudio v1.4.1106 (https://www.rstudio.com) (RStudio Team 2021) and R v3.6.3 (https://www.R-project.org) (R Core Team 2020) and edited the plots in Inkscape v1.1 (https://inkscape.org). We additionally ran the analysis with the input files that were subsampled to 2.14x coverage before running Consensify to test for the effects of sequence coverage.

To examine how the number of differences between individuals compares to a more standard measure of diversity, i.e. heterozygosity, we calculated allele frequencies using genotype likelihoods for each roan and sable antelope individual using ANGSD, setting the -setMaxDepth parameter to twice the average coverage and with the following parameter settings: -setMinDepthInd 5, -minInd 1, -doCounts 1, -GL 2, -doSaf 1, -fold 1, -minQ 30, -minMapQ 30, -uniqueonly 1, -remove_bads 1, -only_proper_pairs 0, -trim 0, -C 0, -baq 2. Next,

we calculated the site frequency spectrum using realSFS in ANGSD. This analysis was performed once using the full dataset and once with the bam files subsampled with SAMtools view to the lowest sequence coverage of a specimen in the comparison data set (4.26x of sable antelope SB1954) to ensure no bias could result from uneven coverage. We plotted the values for the subsampled calculations per individual together with the mean for each species in RStudio and R and edited the plots in Inkscape.

Neighbor-joining tree

We constructed a neighbor-joining tree using the nuclear genome data from the two blue, three roan (Gonçalves et al. 2021) and eight sable antelopes (Koepfli et al. 2019), along with the scimitar-horned oryx (SB20612, Humble et al. 2020) as outgroup. The rescaled bam files from mapDamage for the fossil and historical blue antelope specimens as well as the bam files for all modern individuals were used as input files. We generated the input distance matrix in ANGSD using a consensus base call approach (-doIBS 2), removing singletons (-minMinor 2) and transitions (-rmTrans 1), and using only positions where all individuals had coverage (-minInd 14) (-doCounts 1, -makeMatrix 1, -minQ 30, -minMapQ 30, -mininddepth 2, -uniqueonly 1, -only_proper_pairs 0, -remove_bads 1, -baq 2, -trim 0, -C 0, -doMajorMinor 1, -GL 2). We used FastME v2.1.6.1 (Lefort et al. 2015) to generate a neighbor-joining tree from the distance matrix using default parameters. FigTree v1.4.3 (https://github.com/rambaut/figtree) and Inkscape were used to visualise the tree.

D statistics

For *D* statistics analysis, we used the fasta files generated with Consensify of the roan and sable antelope individuals with the highest coverage (10954, Gonçalves et al. 2021, and SB2152, Koepfli et al. 2019) together with the scimitar-horned oryx as outgroup (SB20612, Humble et al. 2020) and either the fossil or the historical blue antelope specimen. We used the topology resulting from the neighbor-joining tree (fig. 2A) placing the sable antelope in

position one, the blue antelope in position two, and the roan antelope in position three (fig. 3B). We conducted *D* statistics/abbababa test with the tool Dstat (transversion only version, (https://github.com/jacahill/Admixture) and calculated the standard error using the weighted_block_jackknife_D tool with a 1 Mb window size. From this we calculated Z-scores. The results were plotted using RStudio and R and plots were edited with Inkscape.

Sliding window tree analysis

To investigate the direction and timing of gene flow we used a sliding window tree approach. As input we used the randomly sampled fasta files (see above). We then used the tool WindowTrees v1.0 (https://github.com/achimklittich/WindowTrees) to generate non-overlapping sliding window maximum-likelihood phylogenies using RAxML v8.2.12 (Stamatakis 2014). We used binary mode to exclude transitions (--binary), a window size of 100 kb (-w) and a threshold for missing data of 50% (-N 0.5). Prior to the introduction of the WindowTrees tool, this procedure was already successfully used to determine gene flow direction in other studies (Barlow et al. 2018; Westbury et al. 2020; Paijmans et al. 2021). This was performed for all possible combinations with two roan, one sable and one blue antelope using the following individuals: roan: 10954, He108: sable: SB2152, HN216; blue antelope: NBC RB4 D3, NRM 590107. The scimitar-horned oryx (SB20612) was defined as outgroup in each case (--outgroup).

In addition, we ran one sliding window tree analysis again using WindowTrees in binary mode and a missing data threshold of 50% but with a window size of 20 kb and a gap size of 80 kb (-lw 80000) between windows. For this analysis only one individual of each species was used (roan: 10954, sable: SB2152, blue antelope: NBC RB4 D3). Again, the scimitar-horned oryx was used as outgroup (SB20612). We examined the hypothesis of gene flow in the direction of blue into roan antelope using estimates of gene trees and their phylogenetic branch lengths. Under a scenario of gene flow from blue into roan antelope, the portion of introgressed loci will show a signal of more recent common ancestry between the roan and the sable antelope

relative to the species tree topology. On the other hand, a scenario of gene flow in the direction from roan into blue antelope will lead all of the loci to have a signal of similar times of common ancestry between roan and sable antelope in accordance with the species tree topology (fig. 4A). To examine these hypotheses, we extracted the sum of phylogenetic branch lengths separating roan and sable antelope instead of roan and blue antelope as to avoid biases that may influence the branch length, e.g. possible DNA damage and higher sequencing errors, due to lower coverage, as expected from the fossil specimen. We then compared this signal between the gene trees in which roan and blue antelope are sister species (potentially including introgressed regions) against gene trees in which sable and blue antelope are sister species (showing distances between common ancestors of those species). To exclude extremely high and extremely low distances, which likely arise due to the high variance in estimates of branch lengths with negligible information available, we filtered out distances of <0.0002 and >0.9. For any occurrence of gene flow from the blue into the roan antelope, a bimodal distribution would have been expected due to the relatively shorter interspecific branch lengths that would arise. Genetic distances from gene trees were extracted using the R package *ape* v5.5 (Paradis and Schliep 2019, supplementary file S7). We plotted the comparison of the alternative topology against the species topology using RStudio and R and edited the plot in Inkscape.

Mitochondrial genome

Read mapping

We treated the reads for the mitochondrial genome of NBC RB4 D3 in the same way as described for the nuclear genome but used the mitochondrial genome of the blue antelope (MW222233, Hempel et al. 2021a) as reference and removed duplicates with MarkDupsByStartEnd (https://github.com/dariober/Java-cafe/tree/master/MarkDupsByStartEnd). We generated a consensus sequence in Geneious R10 v10.2.3 (Kearse et al. 2012, https://www.geneious.com) using an 85% majority rule

threshold for base calling, a minimum coverage of 3x and the "trim to reference" option. To improve the coverage at the end of the sequence, we mapped and filtered the data again as described above, but used a reference with 200 bp shifted from the end to the start of the reference, considering the circular nature of the mitochondrial genome. We called a consensus sequence as described before and subsequently moved the 200 bp back to the end of the sequence. Then we called a consensus sequence from both sequences using a 50% majority rule threshold for base calling (option "50%—Strict: Bases matching at least 50% of the sequences").

We generated mitochondrial genomes from the raw sequencing data of all roan and sable antelope specimens and the outgroup scimitar-horned oryx specimen for which assembled mitogenomes were not available (roan: He95, He108, 10954; sable antelope: HN216, HN250; scimitar-horned oryx: 20612) (supplementary table S5). We treated the data in the same way as described for the nuclear genomes while using conspecific mitochondrial genomes as references (roan and sable antelope and scimitar-horned oryx: JN632647, JN632648, JN632677; Hassanin et al. 2012) and employing a second mapping step as described for NBC RB4 D3 but varying the length of the reference part that was moved for the second mapping (see above) to be twice the sequencing cycle length of each sample.

Haplotype network with mitochondrial genomes

To determine the extent of differences between our fossil and the previously published historical specimens of the blue antelope, we aligned the mitochondrial genome of NBC RB4 D3 to the other two available complete mitochondrial blue antelope genomes (MW222233, MW222234, Hempel et al. 2021a, supplementary table S17) using the MAFFT algorithm v7.450 (Katoh et al. 2002; Katoh and Standley 2013) with default settings as implemented in Geneious. We removed all ambiguities/missing data (supplementary files S5 & S6). Subsequently, we constructed a median-joining network with POPART v1.7 (Bandelt et al. 1999; Leigh and Bryant 2015) and calculated the number of segregating sites

and nucleotide diversity. In addition, we aligned the mitochondrial genome of NBC RB4 D3 with the two available complete and the two partial mitochondrial genomes of the blue antelope (MW228401, MW228402, supplementary table S17) using again the MAFFT algorithm with default settings as implemented in Geneious and repeated all steps described for the alignment with the complete mitochondrial genomes to generate a median-joining network and determine the number of segregating sites and the nucleotide diversity using POPART.

Phylogenetic tree with mitochondrial genomes

We built an alignment with the consensus sequences of the two blue, three roan and eight sable antelopes using the MAFFT algorithm and default parameters as implemented in Geneious. We used the scimitar-horned oryx as outgroup (SB20612). We removed the control regions from all sequences prior to aligning them due to their limited alignability for different species (supplementary files S3 & S4). We generated two maximum-likelihood phylogenies with 1000 bootstrap replicates each using RAxML v8.2.12, once with the GTR+G substitution model and the other with the BINGAMMA model (Stamatakis 2014). For the latter, the alignment was first transformed into binary format to only score transversions. We visualised the results in FigTree and edited them in Inkscape.

Supplementary Material

Supplementary data are available at Molecular Biology and Evolution online including alignments for the haplotype networks and maximum-likelihood phylogenies and the R script for the sliding window tree genetic distance analysis.

Data availability

The BioProject number of this project in GenBank is PRJNAxxxxxx. The complete mitochondrial genome of the fossil blue antelope specimen NBC RB4 D3 is available at GenBank under the accession number xxxxxxx. The untrimmed raw data were uploaded for the fossil specimen NBC RB4 D3 to the sequence read archive under SRRxxxx–SRRxxxx and for the historical specimen NRM 590107 under SRRxxxx–SRRxxxx. For NRM 590107 run SRRxxxx from BioProject PRJNAxxxxxx was used as well. Accession and BioProject numbers and SRRs will be provided upon acceptance.

Acknowledgments

This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) (project number 315696891, HO 3492/3-1, BI 1879/2-1 to M.H. and F.B.) and by the Swedish Research Council (2017-04647) and FORMAS (2018-01640 both to L.D.). We would like to acknowledge the support of Iziko Museums of South Africa, Heritage Western Cape, Cape Nature, Eastern Cape Provincial Heritage Resources Authority and South African Heritage Resources Agency. We would like to thank Wendy Black, Wilhelmina Seconna and Mark De Benedictis for access to and support while at the Archaeology Unit, Iziko Museums of South Africa. We thank Margarida Gonçalves and Raquel Godinho for providing the raw data for the roan antelope comparison sample 10954 and Gaik Tamazian and Pavel Dobrynin for providing the raw data for the sable antelope comparison samples HN216 and HN250. We would like to thank Stefanie Hartmann for providing the custom perl script for data filtering and Michaela Preick for lab work in screening samples. The authors also acknowledge support from Science for Life Laboratory, the Knut and Alice Wallenberg Foundation, the National Genomics Infrastructure funded by the Swedish Research Council, and Uppsala Multidisciplinary Center for Advanced Computational Science for assistance with massively parallel sequencing and access to the UPPMAX computational infrastructure.

Author Contributions

The study was designed by E.H., F.B., J.T.F., J.S.B., M.H. and M.V.W. Funding was acquired by F.B. and M.H. J.T.F. identified the specimens. E.H. sampled the specimens and performed lab work. E.H., D.A.D. and M.V.W. performed DNA analyses with input from M.H. A.M.K. wrote the WindowTrees program. E.H. and M.V.W. wrote the manuscript with input from F.B., J.T.F., K.-P.K., D.A.D., D.C.K. and M.H. Resources were supplied by L.D. and M.H. Final editing and manuscript preparation was coordinated by E.H. All contributing authors read and agreed to the final manuscript.

References

van Andel TH. 1989. Late Pleistocene sea levels and the human exploitation of the shore and shelf of southern South Africa. *J Field Archaeol* 16:133–155.

Avery DM. 2019. A fossil history of southern African land mammals. Cambridge, New York, Port Melbourne, New Dehli: Cambridge University Press.

Bandelt HJ, Forster P, Röhl A. 1999. Median-joining networks for inferring intraspecific phylogenies. *Mol Biol Evol* 16:37–48.

Barlow A, Cahill JA, Hartmann S, Theunert C, Xenikoudakis G, Fortes GG, Paijmans JLA, Rabeder G, Frischauf C, Grandal-d'Anglade A, et al. 2018. Partial genomic survival of cave bears in living brown bears. *Nat Ecol Evol* 2:1563–1570.

Barlow A, Hartmann S, Gonzalez J, Hofreiter M, Paijmans JLA. 2020. Consensify: A method for generating pseudohaploid genome sequences from palaeogenomic datasets with reduced error rates. *Genes* 11:50.

Barlow A, Paijmans JLA, Alberti F, Gasparyan B, Bar-Oz G, Pinhasi R, Foronova I, Puzachenko AY, Pacher M, Dalén L, et al. 2021. Middle Pleistocene genome calibrates a revised evolutionary history of extinct cave bears. *Curr Biol* 31:1771-1779.e7.

Bollongino R, Tresset A, Vigne J-D. 2008. Environment and excavation: Pre-lab impacts on ancient DNA analyses. *C R Palevol* 7:91–98.

Brink JS, Lee-Thorp JA. 1992. The feeding niche of an extinct springbok, *Antidorcas bondi* (Antelopini, Bovidae), and its paleoenvironmental meaning. *S Afr J Sci* 88:227–229.

Cawthra HC, Compton JS, Fisher EC, MacHutchon MR, Marean CW. 2015. Submerged shorelines and landscape features offshore of Mossel Bay, South Africa. *Geol Soc Spec Publ* 411:219–233.

Cawthra HC, Cowling RM, Andò S, Marean CW. 2020. Geological and soil maps of the Palaeo-Agulhas Plain for the Last Glacial Maximum. *Quat Sci Rev* 235:105858.

Ceballos G, Ehrlich PR, Barnosky AD, García A, Pringle RM, Palmer TM. 2015. Accelerated modern human–induced species losses: Entering the sixth mass extinction. *Sci Adv* 1:e1400253.

Compton JS. 2011. Pleistocene sea-level fluctuations and human evolution on the southern coastal plain of South Africa. *Quat Sci Rev* 30:506–527.

Copeland SR, Cawthra HC, Fisher EC, Lee-Thorp JA, Cowling RM, le Roux PJ, Hodgkins J, Marean CW. 2016. Strontium isotope investigation of ungulate movement patterns on the Pleistocene Paleo-Agulhas Plain of the Greater Cape Floristic Region, South Africa. *Quat Sci Rev* 141:65–84.

Crossman CA, Taylor EB, Barrett-Lennard LG. 2016. Hybridization in the Cetacea: Widespread occurrence and associated morphological, behavioral, and ecological factors. *Ecol Evol* 6:1293–1303.

Dabney J, Knapp M, Glocke I, Gansauge M-T, Weihmann A, Nickel B, Valdiosera C, García N, Pääbo S, Arsuaga J-L, et al. 2013. Complete mitochondrial genome sequence of a middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proc Natl Acad Sci USA* 110:15758–15763.

DeWoody JA, Harder AM, Mathur S, Willoughby JR. 2021. The long-standing significance of genetic diversity in conservation. *Mol Ecol* 30:4147–4154.

Dingle RV, Rogers J. 1972. Pleistocene palaeogeography of the Agulhas Bank. *Trans R Soc S Afr* 40:155–165.

Durand EY, Patterson N, Reich D, Slatkin M. 2011. Testing for ancient admixture between closely related populations. *Mol Biol Evol* 28:2239–2252.

Dussex N, van der Valk T, Morales HE, Wheat CW, Díez-del-Molino D, von Seth J, Foster Y, Kutschera VE, Guschanski K, Rhie A, et al. 2021. Population genomics of the critically endangered kākāpō. *Cell Genomics* 1:100002.

Edwards CJ, Suchard MA, Lemey P, Welch JJ, Barnes I, Fulton TL, Barnett R, O'Connell TC, Coxon P, Monaghan N, et al. 2011. Ancient hybridization and an Irish origin for the modern polar bear matriline. *Curr Biol* 21:1251–1258.

Faith JT. 2011. Late Quaternary megafaunal extinctions in southern Africa's Cape Floral Region.

Faith JT. 2012. Conservation implications of fossil roan antelope (*Hippotragus equinus*) in southern Africa's Cape Floristic Region. In: Louys J, editor. Paleontology in Ecology and Conservation. Berlin, Heidelberg: Springer. p. 239–251.

Faith JT. 2014. Late Pleistocene and Holocene mammal extinctions on continental Africa. *Earth-Sci Rev* 128:105–121.

Faith JT, Behrensmeyer AK. 2013. Climate change and faunal turnover: Testing the mechanics of the turnover-pulse hypothesis with South African fossil data. *Paleobiology* 39:609–627.

Faith JT, Thompson JC. 2013. Fossil evidence for seasonal calving and migration of extinct blue antelope (*Hippotragus leucophaeus*) in southern Africa. *J Biogeogr* 40:2108–2118.

Fisher EC, Bar-Matthews M, Jerardino A, Marean CW. 2010. Middle and late Pleistocene paleoscape modeling along the southern coast of South Africa. *Quat Sci Rev* 29:1382–1398.

FitzSimons FW. 1920. The bluebuck or blaauwbok (*Hippotragus leucophaeus*). In: The natural history of South Africa. Vol. 3. London: Longmans, Green and Co. p. 97–105.

Gansauge M-T, Gerber T, Glocke I, Korlević P, Lippik L, Nagel S, Riehl LM, Schmidt A, Meyer M. 2017. Single-stranded DNA library preparation from highly degraded DNA using T4 DNA ligase. *Nucleic Acids Res* 45:e79.

Gansauge M-T, Meyer M. 2013. Single-stranded DNA library preparation for the sequencing of ancient or damaged DNA. *Nat Protoc* 8:737–748.

Gonçalves M, Siegismund HR, Jansen van Vuuren B, Koepfli K-P, Ferrand N, Godinho R. 2021. *De novo* whole-genome assembly and resequencing resources for the roan (*Hippotragus equinus*), an iconic African antelope. *G3* 11:jkab002.

Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, Patterson N, Li H, Zhai W, Fritz MHY, et al. 2010. A draft sequence of the neandertal genome. *Science* 328:710–722.

Harper F. 1945. Blaauwbok; Blue-buck. In: Extinct and vanishing mammals of the old world. Baltimore: The Lord Baltimore Press. p. 698–700.

Hassanin A, Delsuc F, Ropiquet A, Hammer C, Jansen van Vuuren B, Matthee C, Ruiz-Garcia M, Catzeflis F, Areskoug V, Nguyen TT, et al. 2012. Pattern and timing of diversification of Cetartiodactyla (Mammalia, Laurasiatheria), as revealed by a comprehensive analysis of mitochondrial genomes. *C R Biol* 335:32–50.

Hempel E, Bibi F, Faith JT, Brink JS, Kalthoff DC, Kamminga P, Paijmans JLA, Westbury MV, Hofreiter M, Zachos FE. 2021a. Identifying the true number of specimens of the extinct blue antelope (*Hippotragus leucophaeus*). *Sci Rep* 11:2100.

Hempel E, Westbury MV, Grau JH, Trinks A, Paijmans JLA, Kliver S, Barlow A, Mayer F, Müller J, Chen L, et al. 2021b. Diversity and paleodemography of the Addax (*Addax nasomaculatus*), a Saharan antelope on the verge of extinction. *Genes* 12:1236.

Hofreiter M, Paijmans JLA, Goodchild H, Speller CF, Barlow A, Fortes GG, Thomas JA, Ludwig A, Collins MJ. 2015. The future of ancient DNA: Technical advances and conceptual shifts: Prospects & overviews. *BioEssays* 37:284–293.

Hubbs CL. 1955. Hybridization between fish species in nature. *Syst Zool* 4:1–20.

Humble E, Dobrynin P, Senn H, Chuven J, Scott AF, Mohr DW, Dudchenko O, Omer AD, Colaric Z, Aiden EL, et al. 2020. Chromosomal-level genome assembly of the scimitar-horned oryx: Insights into diversity and demography of a species extinct in the wild. *Mol Ecol Resour* 20:1668–1681.

Jansson G, Thulin C-G, Pehrson Å. 2007. Factors related to the occurrence of hybrids between brown hares *Lepus europaeus* and mountain hares *L. timidus* in Sweden. *Ecography* 30:709–715.

Jónsson H, Ginolhac A, Schubert M, Johnson PLF, Orlando L. 2013. mapDamage2.0: Fast approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics* 29:1682–1684.

Kardos M, Armstrong EE, Fitzpatrick SW, Hauser S, Hedrick PW, Miller JM, Tallmon DA, Funk WC. 2021. The crucial role of genome-wide genetic variation in conservation. *Proc Natl Acad Sci USA* 118:e2104642118.

Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: A novel method for rapid multiple sequence alignment based on fast fourier transform. *Nucleic Acids Res* 30:3059–3066.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Mol Biol Evol* 30:772–780.

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, et al. 2012. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649.

Kerley GIH, Sims-Castley R, Boshoff AF, Cowling RM. 2009. Extinction of the blue antelope *Hippotragus leucophaeus*: Modeling predicts non-viable global population size as the primary driver. *Biodivers Conserv* 18:3235–3242.

Klein RG. 1972. The late Quaternary mammalian fauna of Nelson Bay Cave (Cape Province, South Africa): Its implications for megafaunal extinctions and environmental and cultural change. *Quat res* 2:135–142.

Klein RG. 1974. On the taxonomic status, distribution and ecology of the blue antelope, *Hippotragus leucophaeus* (Pallas, 1766). *Ann S Afr Mus* 65:99–143.

Klein RG. 1980. Environmental and ecological implications of large mammals from upper Pleistocene and Holocene sites in southern Africa. *Ann S Afr Mus* 81:223–283.

Klein RG. 1983. Palaeoenvironmental implications of Quaternary large mammals in the fynbos region. In: Deacon HJ, Hendrey QB, Lambrechts JNN, editors. Fynbos paleoecology: A preliminary synthesis. South African National Scientific Programmes Rep. Pretoria, Republic of South Africa: CSIR. p. 116–138.

Klein RG. 1987. The extinct blue antelope. *Sagittarius* 2:20–23.

Klein RG, Cruz-Uribe K. 2016. Large mammal and tortoise bones from Elands Bay Cave (South Africa): Implications for later Stone Age environment and ecology. *South Afr Humanit* 29:259–282.

Koepfli K-P, Tamazian G, Wildt D, Dobrynin P, Kim C, Frandsen PB, Godinho R, Yurchenko AA, Komissarov A, Krasheninnikova K, et al. 2019. Whole genome sequencing and re-sequencing of the sable antelope (*Hippotragus niger*): A resource for monitoring diversity in *ex situ* and *in situ* populations. *G3* 9:1785–1793.

Kohl FF. 1886. Ueber neue und seltene Antilopen des k. k. naturhistorischen Hofmuseums. *Ann K K Naturh Hofmus Wien* 1:75–86.

Korneliussen TS, Albrechtsen A, Nielsen R. 2014. ANGSD: Analysis of next generation sequencing data. *BMC Bioinform* 15:356.

Larsen PA, Marchán-Rivadeneira MR, Baker RJ. 2010. Natural hybridization generates mammalian lineage with species characteristics. *Proc Natl Acad Sci USA* 107:11447–11452.

Lefort V, Desper R, Gascuel O. 2015. FastME 2.0: A comprehensive, accurate, and fast distance-based phylogeny inference program. *Mol Biol Evol* 32:2798–2800.

Leigh JW, Bryant D. 2015. POPART: Full-feature software for haplotype network construction. *Methods Ecol Evol* 6:1110–1116.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics* 25:2078–2079.

Lichtenstein H. 1811. Reisen im südlichen Africa in den Jahren 1803, 1804, 1805 und 1806. Berlin: C. Salfeld.

Lichtenstein H. 1814. Die Gattung Antilope. *Ges Naturf Freunde Berlin Mag neuesten Entdeck gesammten Naturk* 6:147–160, 163–182.

Lipson M, Sawchuk EA, Thompson JC, Oppenheimer J, Tryon CA, Ranhorn KL, de Luna KM, Sirak KA, Olalde I, Ambrose SH, et al. 2022. Ancient DNA and deep population structure in sub-Saharan African foragers. *Nature* 603:290–296.

Liu S, Westbury MV, Dussex N, Mitchell KJ, Sinding M-HS, Heintzman PD, Duchêne DA, Kapp JD, von Seth J, Heiniger H, et al. 2021. Ancient and modern genomes unravel the evolutionary history of the rhinoceros family. *Cell* 184:4874-4885.e16.

Loftus E, Sealy J, Lee-Thorp J. 2016. New radiocarbon dates and Bayesian models for Nelson Bay Cave and Byneskranskop 1: Implications for the South African later Stone Age sequence. *Radiocarbon* 58:365–381.

Loubser J, Brink J, Laurens G. 1990. Paintings of the extinct blue antelope, *Hippotragus leucophaeus*, in the eastern Orange Free State. *S Afr Archaeol Bull* 45:106–111.

Magoč T, Salzberg SL. 2011. FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27:2957–2963.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* 17:10–12.

Mathieson I, Abascal F, Vinner L, Skoglund P, Pomilla C, Mitchell P, Arthur C, Gurdasani D, Willerslev E, Sandhu MS, et al. 2020. An ancient baboon genome demonstrates long-term population continuity in southern Africa. *Genome Biol Evol* 12:407–412.

Meyer M, Kircher M, Gansauge M-T, Li H, Racimo F, Mallick S, Schraiber J G, Jay F, Prüfer K, de Filippo C, et al. 2012. A high-coverage genome sequence from an archaic Denisovan individual. *Science* 338:222–226.

Mohr E. 1967. Der Blaubock *Hippotragus leucophaeus* (Pallas, 1766) - Eine Dokumentation. (Herre W, Röhrs M, editors.). Hamburg, Berlin: Paul Parey.

Opperman H. 1987. The later Stone Age of the Drakensberg Range and its foothills. Oxford: B.A.R.

Paijmans JLA, Baleka S, Henneberger K, Taron UH, Trinks A, Westbury MV, Barlow A. 2017. Sequencing single-stranded libraries on the Illumina NextSeq 500 platform. Unpublished data. http://arxiv.org/abs/1711.11004. Last accessed 12 April 2022.

Paijmans JLA, Barlow A, Becker MS, Cahill JA, Fickel J, Förster DWG, Gries K, Hartmann S, Havmøller RW, Henneberger K, et al. 2021. African and Asian leopards are highly differentiated at the genomic level. *Curr Biol* 31:1872-1882.e5.

Pallas PS. 1767. Spicilegia zoologica quibus novae imprimis et obscurae animalium species. Berlin: Gottlieb August Lange.

Paradis E, Schliep K. 2019. ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* 35:526–528.

du Plessis SF. 1969. The past and present geographical distribution of the Perissodactyla and Artiodactyla in southern Africa.

Plug I. 1997. Late Pleistocene and Holocene hunter–gatherers in the eastern highlands of South Africa and Lesotho: A faunal interpretation. *J Archaeol Sci* 24:715–727.

R Core Team. 2020. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing.

Rakotoarivelo AR, O'Donoghue P, Bruford MW, Moodley Y. 2019. An ancient hybridization event reconciles mito-nuclear discordance among spiral-horned antelopes. *J Mammal* 100:1144–1155.

Robinson TJ, Bastos AD, Halanych KM, Herzig B. 1996. Mitochondrial DNA sequence relationships of the extinct blue antelope *Hippotragus leucophaeus*. *Naturwissenschaften* 83:178–182.

Robinson TJ, Harley EH. 1995. Absence of geographic chromosomal variation in the roan and sable antelope and the cytogenetics of a naturally occurring hybrid. *Cytogenet Cell Genet* 71:363–369.

RStudio Team. 2021. RStudio: Integrated development environment for R. Boston, MA: RStudio, PBC.

Sánchez-Barreiro F, Gopalakrishnan S, Ramos-Madrigal J, Westbury MV, de Manuel M, Margaryan A, Ciucani MM, Vieira FG, Patramanis Y, Kalthoff DC, et al. 2021. Historical population declines prompted significant genomic erosion in the northern and southern white rhinoceros (*Ceratotherium simum*). *Mol Ecol* 30:6355–6369.

von Schreber JCD, Goldfuß A. 1836. Die Säugthiere in Abbildungen nach der Natur mit Beschreibungen. Erlangen.

Skead CJ. 1980. Blue antelope (extinct blaauwbok), *Hippotragus leucophaeus* (Pallas, 1766). In: Historical mammal incidence in the Cape Province. Vol. 1. 2nd ed. Cape Town: Department of Nature and Environmental Conservation of the Provincial Administration of the Cape of Good Hope. p. 526–537.

Skoglund P, Thompson JC, Prendergast ME, Mittnik A, Sirak K, Hajdinjak M, Salie T, Rohland N, Mallick S, Peltzer A, et al. 2017. Reconstructing prehistoric African population structure. *Cell* 171:59-71.e21.

Smith A. 1849. Illustrations of the zoology of South Africa. London: Smith, Elder and Co.

Smith CI, Chamberlain AT, Riley MS, Cooper A, Stringer CB, Collins MJ. 2001. Not just old but old and cold? *Nature* 410:771–772.

Stamatakis A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30:1312–1313.

Stuiver M, Reimer PJ. 1993. Extended ^{14}C data base and revised CALIB 3.0 ^{14}C age calibration program. *Radiocarbon* 35:215–230.

Themudo GE, Campos PF. 2018. Phylogenetic position of the extinct blue antelope, *Hippotragus leucophaeus* (Pallas, 1766) (Bovidae: Hippotraginae), based on complete mitochondrial genomes. *Zool J Linn Soc* 182:225–235.

Vaz Pinto P, Beja P, Ferrand N, Godinho R. 2016. Hybridization following population collapse in a critically endangered antelope. *Sci Rep* 6:18788.

Venter JA, Brooke CF, Marean CW, Fritz H, Helm CW. 2020. Large mammals of the Palaeo-Agulhas Plain showed resilience to extreme climate change but vulnerability to modern human impacts. *Quat Sci Rev* 235:106050.

Vicente M, Schlebusch CM. 2020. African population history: An ancient DNA perspective. *Curr Opin Genet Dev* 62:8–15.

Westbury MV, Hartmann S, Barlow A, Preick M, Ridush B, Nagel D, Rathgeber T, Ziegler R, Baryshnikov G, Sheng G, et al. 2020. Hyena paleogenomes reveal a complex evolutionary history of cross-continental gene flow between spotted and cave hyena. *Sci Adv* 6:eaay0456.

Westbury MV, Hartmann S, Barlow A, Wiesel I, Leo V, Welch R, Parker DM, Sicks F, Ludwig A, Dalén L, et al. 2018. Extended and continuous decline in effective population size results in low genomic diversity in the world's rarest hyena species, the brown hyena. *Mol Biol Evol* 35:1225–1237.

Xue Y, Prado-Martinez J, Sudmant PH, Narasimhan V, Ayub Q, Szpak M, Frandsen P, Chen Y, Yngvadottir B, Cooper DN, et al. 2015. Mountain gorilla genomes reveal the impact of long-term population decline and inbreeding. *Science* 348:242–245.