1  A systematic examination of learning in the invasive ant
2  *Linepithema humile* reveals very rapid development of short and
3  long-term memories
4

5  Thomas Wagner[1],*, Henrique Galante[1], Tomer J. Czaczkes[1]

6  [1] Animal Comparative Economics laboratory, University of Regensburg, Germany

7  * Corresponding author, email: t.wagner.science@gmail.com

## Abstract

The Argentine ant (*Linepithema humile*) is one of the most damaging and widespread invasive

ant species worldwide. However, control attempts often fail due to insufficient bait uptake,

or bait abandonment. Increasing preference for, and consumption of, is thus an important

requirement for successful control. Associative learning and within-nest information transfer

might be a potential tool for achieving this goal. Here, we conducted a detailed and systematic

investigation of olfactory learning and side learning in Argentine ants. The ants showed very

strong and rapid side learning, choosing the correct arm in a Y-maze 65% of time after just one

visit, and 84% correct after two. Odour learning was even more rapid, with just one visit to a

flavoured food source, reached by a scented runway, leading to 85% choices for the

corresponding scent on a Y-maze. Further experiments demonstrated that having two cues

(runway odour and food flavour) does not improve learning significantly over just one cue.

This rapid learning is long-lasting, with one exposure to a runway odour associated with a

reward resulting in a strong preference (73%) for this odour even after 48 hours. Food flavour

information is transferred efficiently between nestmates in the nest, driving preference: naïve

ants housed with ants fed on flavoured food show a strong preference (77%) for that odour

after 24 hours. Our results demonstrate the impressive learning abilities of *Linepithema*

*humile*, which coupled with efficient intranidal information transfer and strong use of

pheromonal recruitment may help explain their ability to discover and then dominate

resources. However, these strengths could potentially be used against them, by exploiting

learning and information transfer to increase toxic bait uptake during control efforts. Steering

ant preference by leveraging learning might be an underappreciated tool in invasive alien

species control.

**Keywords**: Associative learning, memory, Argentine ants, invasive species, odour

## Introduction

With increasing trade, and the concomitant breakdown of biogeographic barriers, invasive species are spreading globally (Mack and Lonsdale 2001; Mooney and Cleland 2001; Perrings et al. 2005). Invasive species are often economically damaging and ecologically devastating (Marbuah et al. 2014; Escobar et al. 2018), with invertebrates being amongst the most damaging invasive groups (Ricciardi 2015). Ants are especially effective at invading habitats outside their native range (Tsutsui and Suarez 2003), where they displace native species through competition and predation (Ness and Bronstein 2004; Abbott 2006; Matsui et al. 2009).

The Argentine ant (*Linepithema humile* (Mayr, 1868)) is one of the most damaging ant species worldwide, and the most widespread invasive ant in Europe (Human and Gordon 1999; Trigos-Peral et al. 2021). The presence of *L. humile* in invaded areas causes a massive decrease in invertebrate diversity and even affects vertebrates (Wetterer et al. 2001; Suarez et al. 2005; Alvarez-Blanco et al. 2021). In addition, *L. humile* can act as an important agriculture pest by enhancing Hemipteran populations, which then increase the likelihood of fungal and viral infections (Wetterer et al. 2009).

The economic damage through invasive species costs the EU an estimated €12.5 billion a year (European Commission, 2011). Especially combatting invasive ants has become a top priority for conservation programs (Hoffmann et al. 2016). Unfortunatly, in addition to being ecologically and economically damaging, invasive ants are also difficult to control. Two-thirds of *L. humile* eradication attempts fail (Hoffmann et al. 2016). Insect control methods usually rely on the use of insecticide spraying, which is known to be ineffective against ants. The reason behind this ineffectiveness is primarily the strongly sheltered reproductive part of a colony, the queens (Williams et al. 2001). The queens and the brood are not only physically protected, but may also be protected by tuned interaction networks within the colony, which regulate the exposure of vulnerable individuals, at least to pathogens (Stroeymeyt et al. 2018). The current gold standard approach for eradication of ants is the use of baits with a slow-acting poison, which allows workers to bring the food-poison mixture back to the nest to distribute it to the brood and the queen (Hoffmann et al. 2016). Even so, the success rate of such eradication attempts is still low (Souza et al. 2008; Hoffmann 2011). A big issue is the availability of high-quality natural food, which is often preferred by ants and acts as a competitor for the poisoned bait, leading to low bait consumption rates (Rust et al. 2003;

Silverman and Brightwell 2008). Driving bait preference and increasing consumption is thus a

critical step towards successful ant control.

One approach to increasing bait attractiveness, beyond a change in bait formulation,

is to steer ant behaviour. Studies in leaf-cutting ants showed that alarm pheromones can be

used to increase bait consumption and foraging activity (Hughes et al. 2001, 2002). A study in

Argentine ants showed that synthetic (Z)-9-hexadecenal, the main component of the trail

pheromones in this species, increased bait (liquid sucrose) consumption (Greenberg and Klotz

2000). Pheromones can not only be used to lure ants to a bait, but also to disrupt their trail-

following behaviour (Tatsuki et al. 2005; Tanaka et al. 2009). An open field experiment

demonstrated that the combination of a highly-concentrated synthetic pheromone and

insecticidal baits may provide effective control of Argentine ant populations (Sunamura et al.

2011).

However, preference and consumption can also be manipulated by exposing

individuals to tailored information. Insights from behavioural economics and cognition

research are regularly used to drive preference and consumption behaviour in humans. A

widespread example is the exploitation of the endowment effect, where the perceived value

of a good is increased by the subject already owning it. For example, someone who would be

willing to pay €5 for a mug may, if given the mug for free, demand €8 to sell it (Kahneman et

al. 1991). Companies and advertisers exploit this by providing a service at a discount for a

limited period, to cause endowment, before raising the price. This demonstrates that insights

from behavioural economics and cognition research can be used to effectively drive

preferences and consumption behaviour. Driving preference using such behavioural economic

effects has been shown also in animals. Primates and birds are known to be sensitive to

irrational behaviours like the endowment effect (Shafir et al. 2002; Bateson 2002;

Lakshminaryanan et al. 2008). A related concept to an endowment effect was shown in *Lasius*

*niger* ants, where foragers were trained to expect one food flavour, and then provided another

food source of equal value but with a different flavour. These ants undervalued the novel

flavoured food, as demonstrated by reduced food and recruitment (Oberhauser and Czaczkes

2018). However, we are still far from a sophisticated understanding of insect value perception.

Possibly the easiest way of changing the perceived value of a cue is via associative

learning. Associative learning is perhaps the most fundamental form of learning, and strongly

impacts perceived value. Classical association experiments involve the pairing of an

unconditioned stimulus (e.g. food), which innately produces a response, with a neutral

stimulus (e.g. odour), which usually triggers no response. After training, exposure to the

conditioned stimulus can trigger the unconditional response which the unconditional stimulus

usually elicits, or allows the subject to predict that the unconditional stimulus is imminent

(Fanselow 1998; Clark 2004).

Ants have been shown to be good associative learners. Ants can form impressive route

memories which allow them to associate a direction, panorama, or a route with a food reward,

nest (Aron et al. 1988; Harrison et al. 1989; Graham and Collett 2006; Grüter et al. 2011;

Knaden and Graham 2016) or even with a negative outcome (i.e. getting trapped) (Wystrach

et al. 2020). Many ant species have been shown to associate an odour with a food reward

(Roces 1990; Helmy and Jander 2003; Dupuy et al. 2006; Czaczkes et al. 2014; Oberhauser et

al. 2019; Czaczkes and Kumar 2020). Food-odour associations can be rapidly formed and may

last for days (Josens et al. 2009; Arenas and Roces 2018; Piqueret et al. 2019).

However, very little is known about the associative learning ability of invasive ants. In

one study it was shown that Argentine ants (*L. humile*) can use visual and spatial cues to find

a food source, but the experimental setting was a binary choice test between visual cues and

a pheromone trail (Aron et al. 1988, 1993). A recent study in *L. humile* showed that they are

also able to associate an odour spot in a circular arena with a sucrose solution (Rossi et al.

2020). Preexposure to trail pheromone increased food acceptance rates, but had no effect on

associative learning. Rossi et al. (2020) is a very important paper for us, as it is the only

currently available investigation of associative learning in free-running Argentine ants. In that

study, ants were allowed to visit the sucrose reward three times. It is thus unclear how many

exposures to the reward would be required in order for the individual to form a short-term

association. Furthermore, no long-term memory tests were conducted, nor has information

transfer of food-associated odours between nestmates been investigated, as previously

shown in bees, wasps, and other ants (Farina et al. 2005; Provecho and Josens 2009; Schueller

et al. 2010).

Here, we conduct a comprehensive in-depth investigation of learning in the

ecologically important ant *L. humile*. We study how rapidly *L. humile* form short- and long-

term memories, which types of cues best support this, and whether food-related cues are

transferred between nestmates intranidally. Such information is critical if we hope to develop

cognition-based control strategies.

# Materials and Methods

## Colony maintenance

*Linepithema humile* ants were collected in 2021 from Girona, Spain and Proença-a-Nova, Portugal, and were all part of the same European supercolony. Colony fragments (henceforth colonies), consisting of one or more queens and 300-1000 workers, were kept in plastic foraging boxes (32.5 x 22.2 x 11.4 cm) with a plaster of Paris on the bottom. The walls were coated in fluon to prevent escape. Each box contained several 15mL red plastic tubes partly filled with water, plugged with cotton, for use asas nests. The ants were maintained on a 12:12 light:dark cycle at room temperature (21-25 °C) and provided with water *ad libitum*. Colonies were fed for three days with *ad libitum* 0.5M sucrose solution and *Drosophila melanogaster*, and deprived of food for four days prior to testing. For all experiments using odours and/or flavours, donor and recipient colony pairs were used. Such colonies were collected in the same location at the same time, however the donor colonies had never experienced any of the chemicals used, whilst the recipient colonies received individuals that had experienced these.

## Solutions and odours

1M sucrose solutions, (Südzucker AG, Mannheim, Germany), were used as a reward during training for all experiments. Where a negative reinforcement was also presented, 0.6mM quinine (Merck KGaA, Darmstadt, Germany) solutions were used. Paper overlays, used during odour experiments, were stored for at least 1 week prior to the experiments in airtight plastic boxes (19.4 x 13.8 x 6.6 cm) containing a glass plate with 500µl of either strawberry or apple food flavouring (Seeger, Springe, Germany). For experiments were flavoured food was used, 1µL of the respective flavouring was added per mL of 1M sucrose solution. Pilot studies showed *L. humile* workers having a slight preference for strawberry over apple flavour (58% of the ants prefered strawberry, N = 158, see supplement 3).

## Experiment 1 – Short-term spatial memory

Can *L. humile* ants learn to associate a location, one arm of a Y-maze, with the presence of a reward? Here we investigate this by offering a reward, sucrose, on one of the Y-maze arms and a punishment, quinine, on the opposing arm. 6-8 ants per day were tested, testing in total 46 ants from 7 colonies

Y-maze setup and training methods follow Czaczkes (2018). A colony was connected via a

drawbridge to a Y-maze (arms 10cm long, 1cm wide, tapering to 2mm at the bifurcation, see

fig. 1a) covered in unscented disposable paper overlays. A c. 20μl drop of sucrose solution was

placed at the end of one arm of the maze, and a drop of quinine solution on the other.

The first 1-3 ants to initially chose the arm leading to the punishment was marked with acrylic

paint after it subsequently found the sucrose reward. This protocol ensures that the focal ants

do not have an innate preference for the rewarded side. From this point on, only one by one

of the marked ants were selectively allowed to move onto the setup. Upon satiation, ants ran

back over the bridge to the nest and unloaded their collected food to their nestmates. While

unloading, the Y-maze paper overlays were replaced with fresh overlays, to remove any

pheromone trails or cuticular hydrocarbons left by the ant. After unloading, the ants were

allowed back onto the Y-maze via the bridge. We recorded the ant's initial decision (defined

as the antennae crossing a line 2cm from the bifurcation) and final decision (crossing a line

8cm from the bifurcation, 2cm from the arm end). We then allowed the ant to carry out 3

more such visits (5 visits to the sucrose, 4 trained decisions in total). Half the ants were

rewarded on the left, the other half rewarded on the right.

| Rewarded cues | Training | Test |
|---|---|---|
| (a) side-learning experiment | | |
| (b) odour & flavour experiment basic setup | | |

Δ = 1M sucrose    ● = apple    ▢ = unscented paper
◇ = 0.60mM quinine    ● = strawberry    ▢ = water

Figure 1: Basic setup used for all experiments: a) Ants were trained to associate a side of the

Y-maze with a 1M sucrose reward, while the other side contained 0.60mM quinine or water

(exp 1). In all other experiments (b), ants were trained to associate an odour (apple or

strawberry) or/and a flavour (apple or strawberry) with a 1M sucrose reward on a linear

runway and were then tested on a Y-maze. In experiments where quinine was used as

negative reinforcer (2,4,6,8) the flavour and odour of the quinine was the converse of that of

the reward. In the experiments (3,5,8,9) the quinine punishment training was removed. In

the test trials the arms of the Y-maze contained a water drop. The ants' decision was noted

when they crossed the initial decision line (2cm after arm start) and a final decision line (8cm

after arm start, 2 cm before end).

| Experiment | Runway odour | Food flavour | Punishment visit (quinine) | N | Set | Short or long-term memory | % correct choices |
|---|---|---|---|---|---|---|---|
| *2) Scented & flavoured reward vs. punishment* | X | x | X | 72 | 1 | short | **84%** |
| *3) scented & flavoured reward vs. neutral stimulus* | X | X | | 48 | 1 | Short | **85%** |
| *4) scented & unflavoured reward vs. punishment* | X | - | X | 72 | 1 | Short | **72%** |
| *5) unscented & flavoured reward vs. neutral stimulus* | | X | | 48 | 1 | Short | 65.5% |
| *6) unscented & flavoured reward vs. punishment* | | X | X | 44 | 2 | Short | **86%** |
| *7) scented & unflavoured reward vs. punishment* | X | - | X | 48 | 2 | short | **78.5%** |
| *8) scented & flavoured reward vs. neutral stimulus* | X | X | | 70 | 1 | Long (8, 24, 48 hours) | 8h = **85%** 24h = **81%** 48h = **80%** <br><br> Nestmates: 24h = **77%,** 48h = 67% |
| *9) scented & unflavoured reward vs. neutral stimulus vs. neutral stimulus* | x | - | X | 56 | 1 | Long (8, 24, 48 hours) | 8h = **67%,** 24h = **78%** 48h = **73%** <br><br> Nestmates: 24h = 40% 48h = **30%** |
| *1) Side learning: Training on a Y-maze, reward on one arm, punishment on the other* | | | X | | 1 | short | Visit # 2 = **65%,** 3 = **84%** 4 = **84%** 5 = **88%** |

Table 1) Experimental overview. Experiments 2-9 involved training on a linear runway and testing on a Y-maze with scented arms. Experiment 1 involved training over multiple visits on a Y-maze. Runway odour: was the runway scented during training? Food flavour: was the food (and punishment where present) flavoured during training? Experiments in set 1 were carried out between April and August. Experiments in set 2 were carried out between November and December. In short-term learning experiments training occurred immediately after testing, c. 1-5 minutes. In long-term memory experiments testing occurred the stated number of hours after training. % correct choices are printed in **bold** if significantly different from random choice. In experiments 8 and 9 untrained nestmates, housed with the trained nestmates, were also tested.

<u>Experiment 2 – Short-term memory: scented and flavoured reward versus punishment</u>
<u>(experimental set 1)</u>

Experiments conducted in summer are considered as experimental set 1.

The aim of this experiment was to examine if ants can associate two environmental cues
(odour and a flavour) with a sucrose reward. An ant was allowed access to a 10cm-long straight
runway covered by a scented paper overlay, which had a drop of flavoured sucrose solution
at the end (apple or strawberry). After the ant offloaded the collected food in a recipient sub-
colony nest, created to prevent them from sharing odour or flavour information with future
tested ants, it was allowed to visit the runway again. The reward odours and flavours used
were alternated between different ants, where if an ant experienced strawberry as a reward
then it would experience apple as a punishment and vice-versa. The training phase was
concluded once the ant experienced both the reward and punishment in a straight runway
and was allowed back into the recipient colony. The ant was then allowed onto a Y-maze (see
above), in which one arm was covered in a reward-associated scented overlay and the other
in the punishment-scented overlay, and the ant's preference was tested as in experiment 1
(see "test" in fig. 1b & table 1). 72 ants from three donor colonies were tested, ensuring an
equal number of individuals was tested for each odour/flavour and Y-maze side combination
daily.

<u>Experiment 3 – Short-term memory: scented and flavoured reward versus neutral stimulus</u>
<u>(experimental set 1)</u>

The aim of this experiment was to test whether ants need a positive and a negative stimulus
for successful learning, or if only a positive stimulus is sufficient. Ants were trained as in
experiment 2 (runway odour and flavoured food), but no second visit to a punished runway
was performed. The test was carried out as in experiment 2 (see also fig. 1 & table. 1). 48 ants
from two donor colonies were tested, and again both odours/flavours and Y-maze sides were
alternated in every combination possible to remove any potential bias.

<u>Experiment 4 – Short-term memory: scented and unflavoured reward versus punishment</u>
<u>(experimental set 1)</u>

The aim of this experiment was to determine whether ants can associate runway odour cues
alone with a reward or punishment. Ants were trained as in Experiment 2 (see fig. 1 & table

1), except this time, the food reward and the quinine were not flavoured. 72 ants from 6

colonies were tested and both odours and Y-maze sides were alternated daily.

<u>Experiment 5 – Short-term memory: unscented and flavoured reward versus neutral stimulus</u>

<u>(experimental set 1)</u>

The aim of this experiment was to test whether flavoured food, without runway odour or

negative reinforcement, is sufficient to form an association between flavoured food and an

runway odour presented on a Y-maze arm. This is especially important for potential future

applications in pest control. Ants were trained as in Experiment 3, except that the runway

overlay was unscented (see fig. 1 & table.1 ). In the test, ants had the choice between a

scented arm and an unscented one. 48 ants from two colonies were tested.

<u>Experiment 6 – Short-term memory: unscented and flavoured reward versus punishment</u>

<u>(experimental set 2)</u>

We had the oppertunity to collect additional data in winter, which we considered as

experimental set 2.

The aim of this experiment was to test whether flavoured food, without runway odour, is

sufficient to form an association between flavoured food and an runway odour presented on

a Y-maze arm when we add a quinine visit as punishment (contrary to experiment 5). Ants

were trained as in experiment 5 except that a quinine visit was added. The test was carried

out as in experiment 2 (see fig. 1 & table 1). 44 ants from three colonies were tested.

<u>Experiment 7 – **Short-term memory: scented and unflavoured reward versus punishment**</u>

<u>(experimental set 2)</u>

The results of Experiment 6 were unexpected, considering the ones from Experiment 3.

Therefore Experiment 4 was repeated to ensure these surprising results were not caused by

seasonal effects or some other systematic change between experimental sets (see fig. 1 &

table 1). 48 ants from three colonies were tested.

<u>Experiment 8 – Long-term memory: scented and flavoured reward versus neutral stimulus</u>

<u>(experimental set 1)</u>

In this experiment, ants were trained as in experiment 3 (see fig. 1 & table 1) with the

exception that every colony was only conditioned to one reward odour (strawberry or

apple). Two colonies were conditioned to strawberry and two to apple. 50 ants were trained

and, after training, housed in a small sub-colony with 60 naïve ants from their nest. Trained

ants were tested in a Y-maze, as in experiment 3, after 6 hours, 24 hours and 48 hours (4

colonies, 14 ants per period). Each individual ant was only tested once and was then

removed. After the trained ants were tested we also tested the untrained nestmates (4

colonies, 20 ants per period, 24h and 48h), to test whether contact with the trained ants,

which had fed on flavoured food, allowed them to learn the food flavour and thus follow this

odour cue. The colony test order was alternately between a colony conditioned to apple or

strawberry. In total, we tested 70 trained and 80 untrained ants per time period (trained =

6h, 24h and 48; untrained = 24h and 48h).

<u>Experiment 9 – Long-term memory: scented and unflavoured reward versus neutral stimulus</u>

The aim of this experiment was to exclude the possibility that the ants in experiment 8 were

not remembering their association for the whole time period, but rather were refreshing

their memory by repeatedly sampling the flavoured food from other nestmates. To this end,

this experiment was identical to experiment 8 but with an unflavoured reward - only a

scented runway was used (see fig. 1 & table 1). From the previous experiments, we expected

that the trained ants would form a strong association even without food flavour, but that the

untrained ants would not be able to gain any relevant odour information from their sisters.

In total, we tested 56 trained and 80 untrained ants from 4 colonies per time period.

**Statistical analysis**

Only 3.6% – 10.9% of initial and final choices differed, depending on experiment, so we

focused our analysis on the final choices. Data were analysed using generalized linear mixed-

effect models (GLMM) (Bolker et al. 2009) in R version 4.1.0 (R Core Team 2021). GLMMs

were fitted using the lme4 package (Bates et al. 2015). As the data were binomial (correct /

incorrect), a binomial error distribution was used. Since multiple ants were tested per

colony, we included colony as random factor. Each model was validated using the DHARMa

package (Hartig 2018). Results were plotted using the gglot2 package (Wickham 2016). The

complete code and analysis output is provided in supplement 1.

## Results

The complete dataset is provided in supplement 2.

### Experiment 1 - Side-learning

This experiment tested whether Argentine ants can associate a side of the Y-maze with the reward (food), and how this memory develops over subsequent visits.

Over all visits, significantly more choices were made for the arm leading to the food (GLMM, n = 46, z-ratio = 2.513 p = 0.045, see fig. 2). In visit 2, after only one visit to the food, 65% (30/46) of ants chose the arm leading to the food (n =46, z-ratio = 2.581, p = 0.042). This rose to 84% (39/46) in visit 3 (n = 46, z-ratio = 1.860, p = 0.026), 84% in visit 4 (38/45) (n = 45, z-ratio = 1.469, p = 0.032) and 88% (39/44) in visit 5 (n = 44, z-ratio = 2.245, p = 0.009). For a pairwise comparison between visits, see online supplement 3

Figure 2: Side-learning: Proportion of ants choosing the side associated with reward (food) per visit. Ants chose the side associated with reward significantly more often than chance after all visits. Bars depict mean, whiskers 95% confidence intervals derived from the fitted GLMM. The dotted horizontal line displays chance level of 50%. *p < 0.05; **p < 0.01; ***p < 0.001.

**Odour association: Short-term memory**

The aim of experiments 2 – 7 was to understand if and how well Argentine ants can
associate odours and food flavours with food rewards, and whether punishment improves
learning (as has been shown in honeybees, Avargues-Weber et al. 2010).

Figure 3: Short-term odour associative learning: Proportion of ants reaching the end of a Y
maze arm scented with an odour associated with a sucrose reward. Ants were able to form a
strong association to both training runway odour and food flavour. Punishment of the
contrasting odour using quinine did not improve learning. Ants learned better in
experimental set 2 (conducted in the winter) – this is especially noteable when ants were
trained to flavoured food without a scented runway. Offering both odour and flavour cues
did not increase learning compared to offering only one cue. Bars depict means, whiskers
95% confidence intervals derived from the fitted GLMM. The dotted horizontal line displays
chance level of 50%. *$p < 0.05$; **$p < 0.01$; ***p < 0.001.

Experiment 2 - Runway odour and flavoured reward (experimental set 1)

This experiment tested if ants learn to associate the sucrose reward with both
environmental cues, a food flavour and the corresponding runway odour (apple or
strawberry). 84% (61/72) of the ants chose the food-associated odour arm (GLMM, z-ratio =
3.375, p < 0.001, see fig. 3). The specific odour rewarded did not significantly affect choice
accuracy (apple 80% and strawberry 88%, z-ratio = -0.972, p = 0.3312).

Experiment 3 – Runway odour and flavoured reward, no quinine visit
(experimental set 1)

This experiment tested whether a reward alone, and no punishment (quinine), results in an
equally high proportion of choices for the food-associated odour. 85% (41/48) of the ants

chose food-associated odour arm (GLMM, z-ratio = 2.938, p = 0.003, see fig. 3). Again, the

specific odour rewarded did not significantly affect choice accuracy (apple 83% and

strawberry 87%, z-ratio = -0.408, p = 0.683).

Experiment 4 - Runway odour, unflavoured reward (experimental set 1)

This experiment tested if Argentine ants can associate a runway odour (apple or strawberry)

with an unflavoured reward. 72% (52/72) of the ants chose the side of the Y-maze with a

scent associated with a reward (GLMM, z-ratio = 2.568, p = 0.010). Ants performed

identically on both rewarded odours (72% correct decisions for both odours, z-ratio = 0.00, p

= 1.0000, see fig. 3).

Experiment 5 - Flavoured reward, no runway odour, no quinine (experimental set 1)

This experiment was run to test the ant´s learning ability without negative reinforcement

and using only one cue, the flavoured food. 62.5% (30/48) of ants chose the food-associated

odour arm, which does not differ significantly from random choice (GLMM, z-ratio = 1.212, p

= 0.226, see fig. 3). Again, there was no significant difference between performance with the

two odours (62.5% correct decisions for both odours, z-ratio = 0.000, p = 1.000).

Experiment 6 - Flavoured reward, no runway odour (experimental set 2)

This experiment was similar as the 5$^{th}$ experiment except that a quinine visit was added and

with only one cue (flavour). Again, the purpose of this study was to test for the need of a

punishment visit. Unlike in summer (experimental set 1), the winter experiment

(experimental set 2) showed that ants were able to learn to associate just one cue (flavour)

with a sucrose reward. 86% (38/44) of the ants chose the side of the Y-maze with a scent

associated with reward (GLMM, z-ratio = 2.721, p = 0.0065, see fig. 3. There was no

significant difference between the two odours (apple 81% and strawberry 90%, z-ratio = -

0.863, p = 0.388)

Experiment 7 - Runway odour, unflavoured reward (experimental set 2)

This experiment, conducted in winter, was the same as the 4$^{th}$ experiment. The purpose was

to test if ants show a systematic difference in learning between the two experimental sets.

The ants again formed a strong association between just one cue (odour) and the sucrose

reward. 87.5% (42/48) of the ants chose the side of the Y-maze with a scent associated with reward a reward (GLMM, , z-ratio = 2.656, p = 0.007, see fig. 3). Again, the specific odour rewarded did not significantly affect choice accuracy (apple 79% and strawberry 95%, z-ratio = -1.582, p = 0.1138).

The roles of runway odour and food flavour in learning

In order to characterise the relative roles of runway odour and food flavour in driving learning, we compared the behaviour of ants in experiments 2 (runway odour and food flavour during training) with the results from experiment 4 (only runway odour) and experiment 5 (only food flavour). Note, however, that experiment 5 did not include a negative reinforcement, while experiments 2 and 4 did. We consider this appropriate, since negative reinforcement did not improve learning (compare experiments 2 & 3).

Combined runway odour and flavoured food led to a significantly higher proportion of correct choices than only having flavoured food (z-ratio = -2.714, p = 0.018). However, runway odour alone was not significantly weaker than the combined cues (z-ratio = -1.118, p = 0.502). Runway odour alone did not result in significantly more correct choices than food flavour alone (z-ratio = 1.803, p = 0.168).

**Long-term odour and flavour association**

The aim of experiments 8 and9 was to examine if Argentine ants also form a long-term (6h, 24h, 48h) association between runway odours and food rewards (experiment 9) and with flavoured food rewards (experiment 8).

Figure 4: Long-term odour associative learning: Proportion of ants choosing the odour associated with a sucrose reward. Trained ants were tested after 6h, 24h, and 48h. **4a:** Untrained nestmates also showed a preference for the rewarded odour when food was flavoured, which was not significantly weaker than that of trained ants. **4b:** Trained ants were able to associate a runway odour alone with an unflavoured sucrose reward after 6h, 24h and 48h. By contrast, untrained nestmates did not show any odour preference after 24 hours, and showed an aversion for the food odour after 48 hour. Bars depict means, whiskers 95% confidence intervals derived from the fitted GLMM. The dotted horizontal line displays chance level of 50%. *$p < 0.05$; **$p < 0.01$; ***p < 0.001.

### Experiment 8 - Long-term memory, runway odour and flavoured food

*Trained ants and their ability to build a long-term association*

Ants trained on combined runway odour and food flavour significantly preferred the odour-associated Y-maze arm after 6 hours, 24 hours and 48 hours (GLMM, z-ratio = 4.204, p <

0.0001, see fig. 4a), with 85% (60/70), 81% (57/70), and 80% (56/70) correct choices after 6, 24, and 48 hours respectively (6 hours: GLMM, z-ratio = 5.246 , p < 0.0001, 24 hours: GLMM, z-ratio = 4.809, p < 0.0001 , 48 hours: GLMM, z-ratio = 8.515, p < 0.0001).

*Odour preference in untrained ants housed with flavoured-food fed nestmates (trained)*

Untrained ants housed with ants trained with flavoured food showed a strong preference for that food flavour after 24 and 48 hours (z-ratio = 3.830, p < 0.0001, see fig. 4a). 77% (62/80) and 67% (54/80) of ants chose food-associated odour after 24 and 48 hours respectively (24 hours: GLMM, z-ratio = 4.619, p < 0.0001, 48 hours: GLMM, n = 80, z-ratio = 2.511, p = 0.012).

Experiment 9 - Long-term memory, only runway odour, unflavoured reward

This experiment was run to quantify long term learning without access to informative flavoured food, since this could refresh the ants' memories.

*Trained ants and their ability to build a long-term association with unflavoured food*

Ants trained with runway odour and unflavoured food nonetheless significantly preferred the odour-associated Y-maze arm after 6, 24 and 48 hours (GLMM, z-ratio = 3.990, p < 0.0001, see fig. 4b), with 67% (38/56), 78% (44/56), 73% (41/56) correct choices after 6, 24, and 48 hours respectively (6 hours: GLMM, z-ratio = 2.611 , p = 0.009, 24 hours: GLMM, z-ratio = 3.990, p < 0.0001 , 48 hours: GLMM, z-ratio = 3.332, p < 0.0001.

*Odour preference in untrained ants housed with unflavoured-food fed nestmates (trained)*

Unlike naïve ants housed with nestmates trained on flavoured food, naïve ants housed with nestmates trained on unflavoured food showed no preference for the food-associated odour (z-ratio = -1.777, p = 0.075, see fig. 4b). 40% (32/80) and 30% (24/80) of ants chose the food-odour covered Y-maze arm after 24 and 48 hours, respectively. Indeed, while the choice of naïve ants after 24 hours did not differ from chance (GLMM, z-ratio = -1.381, p = 0.268), after 48 hours ants significantly avoided the scented arm (z-ratio = -3.473, p < 0.0001).

*The effect of food flavour presence on the long-term memory of trained ants*

Here we compared how well the Argentine ants performed on the long-term memory test

(6h, 24h, 48h) when they had a flavoured reward vs. an unflavoured reward. Do they form a

significantly better memories when they had a flavoured reward?

There was no significant difference between trained ants who had an unflavoured reward

treatment (Experiment 9) and those trained on flavoured food (experiment 8) after 6 hours,

24 hours and 48 hours (6 hours: GLMM, n =56[exp.8] and n=70[exp.9] , z-ratio = 2.344 , p =

0.176 ,24 hours: GLMM, n =56[exp.8] and n=70[exp.9] , z-ratio = 0.399 , p = 0.998, 48 hours:

GLMM, n =56[exp.8] and n=70[exp.9] , z-ratio = 0.897, p = 0.947).

However, untrained ants who were treated with an unflavoured reward (9. Experiment)

performed significantly worse than ants with a flavoured reward treatment (8. Experiment)

after 24 hours and 48 hours (24 hours: GLMM, n =80[exp.8] and n=80[exp.9], z-ratio = 4.709,

p < 0.001, 48 hours: GLMM, n =80[exp.8] and n=80[exp.9], z-ratio = 4.631, p < 0.001).

## Discussion

Linepithema humile are adept learners: The side-learning results show that Argentine ants are capable of learning to associate a direction or an arm of a Y-maze with a food reward after just one visit, and accuracy improves with further visits (see fig. 2). Such rapid and reliable learning is surprising, given the fact that L. humile relies strongly on social information (pheromone trails), while other recruiting social insects which learn equally fast, such as the ants Lasius niger and Paraponera clavata ants, and the honeybee Apis melifera, prioritize memories over recruitment signals when the two conflict (Harrison et al. 1989; Aron et al. 1993; Grüter et al. 2008, 2011; Czaczkes et al. 2013; von Thienen et al. 2016). L. humile's strong individual memory may allow them to reliably recruit even to distant resources, while their strong reliance on social information, and ability to recruit from active trails (Flanagan et al. 2013), may allow them to rapidly dominate these resources once found. Indeed, the rapid discovery of new resources and massive, rapid recruitment might be an important contribution to making Argentine ants competitive against native ants (Holway 1999), but see (Cordonnier et al. 2020). Route learning and navigation is hindered by higher complexity, at least in La. niger ants, but when trail pheromones are present, error rates decrease (Czaczkes et al. 2013). The combination of social information and a strong individual side-learning memory could hence be extremely beneficial and explain why Argentine ants developed such a strong individual route learning ability in parallel to their pheromone recruitment system. However, a systematic exploration of learning in (social) insects has not been carried out, so it is also possible that many or most ants are equally good learners.

Alongside their impressive route learning, Argentine ants also show even more rapid learning of olfactory cues. Ants were able to form a strong association after only one rewarded visit (see fig. 3 & table 1). These findings are again comparable to results from Lasius niger (Czaczkes et al. 2014; Czaczkes and Kumar 2020). Somewhat surprisingly, results from both the short-term and long-term learning experiments showed that punishing the alternative odour with quinine does not improve response accuracy. While some studies report ants (Camponotus fellah) failing to form a negative association to a scent (Josens et al. 2009), studies on Las. niger ants and honeybees do (Avarguès-Weber et al. 2010; Wenig et al. 2021). Also remarkable is that L. humile is able to form an association with a food source given only one cue: a food flavour or a runway odour. Although the difference between choice accuracy

when trained on only one cue or both was not significantly, the gain of 10-16 (see fig. 3 & table

1) percentage points when both are available, and the fact that both single cues were

sufficient to form an association, suggests to us that cue effects could have an additive or sub-

additive effect on learning.

To our knowledge, the learning ability of *Linepithema humile* has never before been

described in such detail, and their very rapid learning of olfactory cues never fully appreciated.

Rossi et al. (2020) previously demonstrated that *L. humile* could associate an odour with a

food reward. Our results support their findings using a different experimental approach, and

with a higher resolution: the Rossi study allowed 3 visits to a circular open foraging arena, so

did not explore the limits of ant learning. Our study controlled the experiences of the

individual ants more tightly, by reducing the number of rewarded visits to one, and separating

rewarded and unrewarded learning events. While other studies have also demonstrated that

ants can associate an odour alone with a reward, in combination with a quinine negative trial,

these studies required extensive training, with 6-24 visits to the reward, and the odour was

blown by an air stream instead of using scented paper overlays (Dupuy et al. 2006; Guerrieri

and d'Ettorre 2010). To our knowledge, our study is also the first study to demonstrated that

*Linepithema humile* can learn to associate the flavour of the food alone with the food reward.

Alongside the strong short-term olfactory memory, our study demonstrates that

Argentine ants also possess a strong long-term memory, which lasts for at least 48 hours (see

fig. 4 & table 1). As in our short-term memory tests, an runway odour alone, without food

flavour, was sufficient to drive strong and stable learning. This is in line with studies on other

species (*Formica fusca* and *Camponotus fellah)*, which show stable memory for at least 72

hours, and a decay in learning after a week (Josens et al. 2009; Piqueret et al. 2019). However,

the study on *Camponotus fellah* again used very extensive training (16 training visits). We

found that one exposure alone, without a punishment visit, was all that was required to elicit

strong and stable learning. One similar study showed, that *Cataglyphis fortis* desert ants can

also build after only visit a strong association between a food source and an odour (Huber and

Knaden 2018). This again suggests that *L. humile* are an extraordinarily good olfactory

learners, as well as strong route learners, although differences in experimental protocol may

also have contributed to the impressive *L. humile* results.

It is also important to note that, in the long-term memory experiments, tested ants

were housed in a small sub-nest where they fed their nestmates after their training. As a

result, flavoured food was likely distributed amongst the workers via trophallaxis. This could

have acted to 'refresh' the memory of the tested ants for some time. However, learning

remained strong even if ants were trained only with runway odours, and without food odours.

These impressive learning skills are not limited to trained ants. Tellingly, inclusion of food

flavour resulted in the untrained nestmates also developing a preference for that odour,

demonstrating a very effective distribution of food-related information within the nest. Such

intranidal information spread affecting future foraging decision has been reported in both

other ants (Roces 1990; Provecho and Josens 2009; Arenas and Roces 2018), bees (Farina et

al. 2005) and wasps (Jeanne and Taylor 2009).

Finally, our results indicate that there might be seasonal variation in learning, although

this was not tested systematically. While our summer experiments showed that flavour alone

is not sufficient for ants to form a strong association with the food reward, repeating the

experiment in the winter resulted in very strong learning. Furthermore, our only runway odour

experiment conducted in winter revealed an even stronger learning than in summer (see. fig.

3 & table 1). However, it has to be mentioned that the flavour experiment in winter had a

quinine visit, which summer experiments did not. Nevertheless, as mentioned above, quinine

does not seem to improve learning within a season, which makes a seasonal effect more likely.

The apparent improvement in learning in the winter may be ecologically relevant, but may

also be a (to us) inexplicable laboratory artefact. We note that the opposite pattern was found

in *Lasius niger* in a route-learning paradigm (Oberhauser et al. 2018). Season and temperature

effects on foraging and recruitment behaviour are commonly reported in social insects (Porter

and Tschinkel 1987; Stuble et al. 2013; Stroeymeyt et al. 2014; Park and Nieh 2017)

*Linepithema humile* are adept learners and rapidly disseminate food-related

information in the nest – abilities which might help them to adapt to new environments and

dominate resources, and could be one reason for their success as invaders. However, their

strength could potentially become a weakness, if their memory can be used against them. One

possibility would be to use associative learning and intranidal information transfer to steer

foraging preference towards poisoned baits, much as honeybee preference can be steered for

pollination purposes (Farina et al. 2020). Similarly, foraging preference of leaf-cutter ant

colonies can be steered to a specific food target by exploiting their preference for foraging on

resources which smell or taste like those already being collected (Arenas and Roces 2018).

While leveraging learning to control animal behaviour is commonly used in behavioural

conservation of vertebrates, it is mostly used to reduce crop damage or to minimize negative

interactions with humans (Matsuzawa et al. 1983; Webb et al. 2015; Valenta et al. 2021).

Leveraging learning to improve the control of invasive animals may be an underappreciated

tool in the conservation toolbox.

Acknowledgements

We would like to thank Silvia Abril and Eduardo Sequeira for providing ant colonies. A special

thank you to Roxana Josens for comments on earlier versions of this manuscript. Thomas

Wagner and Henrique Galantea were supported by an ERC starter grant (Cognitive control:

948181) to Tomer J. Czaczkes. Tomer J. Czaczkes was supported by a Heisenberg fellowship

from the Deutsche Forschungsgemeinschaft (CZ 237 / 4-1).

# References

Abbott KL (2006) Spatial dynamics of supercolonies of the invasive yellow crazy ant, *Anoplolepis gracilipes*, on Christmas Island, Indian Ocean. Divers Distrib 12:101–110. https://doi.org/10.1111/j.1366-9516.2006.00193.x

Alvarez-Blanco P, Cerdá X, Hefetz A, et al (2021) Effects of the Argentine ant venom on terrestrial amphibians. Conserv Biol 35:216–226. https://doi.org/10.1111/cobi.13604

Arenas A, Roces F (2018) Appetitive and aversive learning of plants odors inside different nest compartments by foraging leaf-cutting ants. J Insect Physiol 109:85–92. https://doi.org/10.1016/j.jinsphys.2018.07.001

Aron S, Beckers R, Deneubourg JL, Pasteels JM (1993) Memory and chemical communication in the orientation of two mass-recruiting ant species. Insectes Sociaux 40:369–380. https://doi.org/10.1007/BF01253900

Aron S, Deneubourg JL, Pasteels JM (1988) Visual cues and trail-following idiosyncrasy *inleptothorax unifasciatus*: An orientation process during foraging. Insectes Sociaux 35:355–366. https://doi.org/10.1007/BF02225811

Avarguès-Weber A, de Brito Sanchez MG, Giurfa M, Dyer AG (2010) Aversive Reinforcement Improves Visual Discrimination Learning in Free-Flying Honeybees. PLoS ONE 5:e15370. https://doi.org/10.1371/journal.pone.0015370

Bateson M (2002) Context-dependent foraging choices in risk-sensitive starlings. Anim Behav 64:251–260. https://doi.org/10.1006/anbe.2002.3059

Clark RE (2004) The classical origins of Pavlov's conditioning. Integr Physiol Behav Sci 39:279–294. https://doi.org/10.1007/BF02734167

Cordonnier M, Blight O, Angulo E, Courchamp F (2020) The Native Ant *Lasius niger* Can Limit the Access to Resources of the Invasive Argentine Ant. Animals 10:2451. https://doi.org/10.3390/ani10122451

Czaczkes TJ, Grüter C, Ellis L, et al (2013) Ant foraging on complex trails: route learning and the role of trail pheromones in *Lasius niger*. J Exp Biol jeb.076570. https://doi.org/10.1242/jeb.076570

Czaczkes TJ, Kumar P (2020) Very rapid multi-odour discrimination learning in the ant *Lasius niger*. Insectes Sociaux 67:541–545. https://doi.org/10.1007/s00040-020-00787-0

Czaczkes TJ, Schlosser L, Heinze J, Witte V (2014) Ants use directionless odour cues to recall odour-associated locations. Behav Ecol Sociobiol 68:981–988. https://doi.org/10.1007/s00265-014-1710-2

Dupuy F, Sandoz J-C, Giurfa M, Josens R (2006) Individual olfactory learning in *Camponotus ants.* Anim Behav 72:1081–1091. https://doi.org/10.1016/j.anbehav.2006.03.011

Escobar LE, Mallez S, McCartney M, et al (2018) Aquatic Invasive Species in the Great Lakes Region: An Overview. Rev Fish Sci Aquac 26:121–138. https://doi.org/10.1080/23308249.2017.1363715

Fanselow MS (1998) Pavlovian Conditioning, Negative Feedback, and Blocking: Mechanisms that Regulate Association Formation. Neuron 20:625–627. https://doi.org/10.1016/S0896-6273(00)81002-8

Farina WM, Arenas A, Díaz PC, et al (2020) Learning of a Mimic Odor within Beehives Improves Pollination Service Efficiency in a Commercial Crop. Curr Biol 30:4284-4290.e5. https://doi.org/10.1016/j.cub.2020.08.018

Farina WM, Grüter C, Díaz PC (2005) Social learning of floral odours inside the honeybee hive. Proc R Soc B Biol Sci 272:1923–1928. https://doi.org/10.1098/rspb.2005.3172

Flanagan TP, Pinter-Wollman NM, Moses ME, Gordon DM (2013) Fast and Flexible: Argentine Ants Recruit from Nearby Trails. PLoS ONE 8:e70888. https://doi.org/10.1371/journal.pone.0070888

Graham P, Collett TS (2006) Bi-directional route learning in wood ants. J Exp Biol 209:3677–3684. https://doi.org/10.1242/jeb.02414

Greenberg L, Klotz JH (2000) Argentine Ant (Hymenoptera: Formicidae) Trail Pheromone Enhances Consumption of Liquid Sucrose Solution. J Econ Entomol 93:119–122. https://doi.org/10.1603/0022-0493-93.1.119

Grüter C, Balbuena MS, Farina WM (2008) Informational conflicts created by the waggle dance. Proc R Soc B Biol Sci 275:1321–1327. https://doi.org/10.1098/rspb.2008.0186

Grüter C, Czaczkes TJ, Ratnieks FLW (2011) Decision making in ant foragers (*Lasius niger*) facing conflicting private and social information. Behav Ecol Sociobiol 65:141–148. https://doi.org/10.1007/s00265-010-1020-2

Guerrieri FJ, d'Ettorre P (2010) Associative learning in ants: Conditioning of the maxilla-labium extension response in *Camponotus aethio*ps. J Insect Physiol 56:88–92. https://doi.org/10.1016/j.jinsphys.2009.09.007

Harrison JF, Fewell JH, Stiller TM, Breed MD (1989) Effects of experience on use of orientation cues in the giant tropical ant. Anim Behav 37:869–871. https://doi.org/10.1016/0003-3472(89)90076-6

Helmy O, Jander R (2003) Topochemical learning in black carpenter ants (*Camponotus pennsylvanicus*). Insectes Sociaux 50:32–37. https://doi.org/10.1007/s000400300005

Hoffmann BD (2011) Eradication of populations of an invasive ant in northern Australia: successes, failures and lessons for management. Biodivers Conserv 20:3267–3278. https://doi.org/10.1007/s10531-011-0106-0

Hoffmann BD, Luque GM, Bellard C, et al (2016) Improving invasive ant eradication as a conservation tool: A review. Biol Conserv 198:37–49. https://doi.org/10.1016/j.biocon.2016.03.036

Holway DA (1999) Competitive mechanisms underlying the displacement of native ants by the invasive Argentine ant. Ecology 80:238–251. https://doi.org/10.1890/0012-9658(1999)080[0238:CMUTDO]2.0.CO;2

Huber R, Knaden M (2018) Desert ants possess distinct memories for food and nest odors. Proc Natl Acad Sci 115:10470–10474. https://doi.org/10.1073/pnas.1809433115

Hughes WOH, Howse PE, Vilela EF, et al (2002) Field Evaluation of Potential of Alarm Pheromone Compounds to Enhance Baits for Control of Grass-Cutting Ants (Hymenoptera: Formicidae). J Econ Entomol 95:537–543. https://doi.org/10.1603/0022-0493-95.3.537

Hughes WOH, Howse PE, Vilela EF, Goulson D (2001) The response of grass-cutting ants to natural and synthetic versions of their alarm pheromone: Alarm response of grass-cutting ants. Physiol Entomol 26:165–172. https://doi.org/10.1046/j.1365-3032.2001.00230.x

Human KG, Gordon DM (1999) Behavioral interactions of the invasive Argentine ant with native ant species. Insectes Sociaux 46:159–163. https://doi.org/10.1007/s000400050127

Jeanne R, Taylor B (2009) Individual and Social Foraging in Social Wasps. CRC Press, chapter 3, pp 53–79

Josens R, Eschbach C, Giurfa M (2009) Differential conditioning and long-term olfactory memory in individual *Camponotus fellah* ants. J Exp Biol 212:1904–1911. https://doi.org/10.1242/jeb.030080

Kahneman D, Knetsch JL, Thaler RH (1991) Anomalies: The Endowment Effect, Loss Aversion, and Status Quo Bias. J Econ Perspect 5:193–206. https://doi.org/10.1257/jep.5.1.193

Knaden M, Graham P (2016) The Sensory Ecology of Ant Navigation: From Natural Environments to Neural Mechanisms. Annu Rev Entomol 61:63–76. https://doi.org/10.1146/annurev-ento-010715-023703

Lakshminaryanan V, Keith Chen M, Santos LR (2008) Endowment effect in capuchin monkeys. Philos Trans R Soc B Biol Sci 363:3837–3844. https://doi.org/10.1098/rstb.2008.0149

Mack RN, Lonsdale WM (2001) Humans as Global Plant Dispersers: Getting More Than We Bargained For. BioScience 51:95. https://doi.org/10.1641/0006-3568(2001)051[0095:HAGPDG]2.0.CO;2

Marbuah G, Gren I-M, McKie B (2014) Economics of Harmful Invasive Species: A Review. Diversity 6:500–523. https://doi.org/10.3390/d6030500

Matsui S, Kikuchi T, Akatani K, et al (2009) Harmful Effects of Invasive Yellow Crazy Ant *Anoplolepis gracilipes* on Three Land Bird Species of Minami-Daito Island. Ornithol Sci 8:81–86. https://doi.org/10.2326/048.008.0111

Matsuzawa T, Hasegawa Y, Gotoh S, Wada K (1983) One-trial long-lasting food-aversion
learning in wild Japanese monkeys (*Macaca fuscata*). Behav Neural Biol 39:155–159.
https://doi.org/10.1016/S0163-1047(83)90791-4

Mooney HA, Cleland EE (2001) The evolutionary impact of invasive species. Proc Natl Acad
Sci 98:5446–5451. https://doi.org/10.1073/pnas.091093398

Ness JH, Bronstein JL (2004) The Effects of Invasive Ants on Prospective Ant Mutualists. Biol
Invasions 6:445–461. https://doi.org/10.1023/B:BINV.0000041556.88920.dd

Oberhauser FB, Czaczkes TJ (2018) Tasting the unexpected: disconfirmation of expectations
leads to lower perceived food value in an invertebrate. Biol Lett 14:20180440.
https://doi.org/10.1098/rsbl.2018.0440

Oberhauser FB, Koch A, Czaczkes TJ (2018) Small differences in learning speed for different
food qualities can drive efficient collective foraging in ant colonies. Behav Ecol
Sociobiol 72:164. https://doi.org/10.1007/s00265-018-2583-6

Oberhauser FB, Schlemm A, Wendt S, Czaczkes TJ (2019) Private information conflict: *Lasius*
*niger* ants prefer olfactory cues to route memory. Anim Cogn 22:355–364.
https://doi.org/10.1007/s10071-019-01248-3

Park B, Nieh JC (2017) Seasonal trends in honey bee pollen foraging revealed through DNA
barcoding of bee-collected pollen. Insectes Sociaux 64:425–437.
https://doi.org/10.1007/s00040-017-0565-8

Perrings C, Dehnen-Schmutz K, Touza J, Williamson M (2005) How to manage biological
invasions under globalization. Trends Ecol Evol 20:212–215.
https://doi.org/10.1016/j.tree.2005.02.011

Piqueret B, Sandoz J-C, d'Ettorre P (2019) Ants learn fast and do not forget: associative
olfactory learning, memory and extinction in *Formica fusca*. R Soc Open Sci 6:190778.
https://doi.org/10.1098/rsos.190778

Porter SD, Tschinkel WR (1987) Foraging in *Solenopsis invicta* (Hymenoptera: Formicidae):
Effects of Weather and Season. Environ Entomol 16:802–808.
https://doi.org/10.1093/ee/16.3.802

Provecho Y, Josens R (2009) Olfactory memory established during trophallaxis affects food
search behaviour in ants. J Exp Biol 212:3221–3227.
https://doi.org/10.1242/jeb.033506

Ricciardi A (2015) Ecology of Invasive Alien Invertebrates. In: Thorp and Covich's Freshwater
Invertebrates. Elsevier, 2015, pp 83–91

Roces F (1990) Olfactory conditioning during the recruitment process in a leaf-cutting ant.
Oecologia 83:261–262. https://doi.org/10.1007/BF00317762

Rossi N, Pereyra M, Moauro MA, et al (2020) Trail pheromone modulates subjective reward
evaluation in Argentine ants. J Exp Biol 223:jeb230532.
https://doi.org/10.1242/jeb.230532

Rust MK, Reierson DA, Klotz JH (2003) Pest Management of Argentine Ants (Hymenoptera: Formicidae)1. J Entomol Sci 38:159–169. https://doi.org/10.18474/0749-8004-38.2.159

Schueller TI, Nordheim EV, Taylor BJ, Jeanne RL (2010) The cues have it; nest-based, cue-mediated recruitment to carbohydrate resources in a swarm-founding social wasp. Naturwissenschaften 97:1017–1022. https://doi.org/10.1007/s00114-010-0712-9

Shafir S, Waite T, Smith B (2002) Context-dependent violations of rational choice in honeybees (*Apis mellifera*) and gray jays (*Perisoreus canadensis*). Behav Ecol Sociobiol 51:180–187. https://doi.org/10.1007/s00265-001-0420-8

Silverman J, Brightwell RJ (2008) The Argentine Ant: Challenges in Managing an Invasive Unicolonial Pest. Annu Rev Entomol 53:231–252. https://doi.org/10.1146/annurev.ento.53.103106.093450

Souza E, Follett PA, Price DK, Stacy EA (2008) Field Suppression of the Invasive Ant *Wasmannia auropunctata* (Hymenoptera: Formicidae) in a Tropical Fruit Orchard in Hawaii. J Econ Entomol 101:7

Stroeymeyt N, Grasse AV, Crespi A, et al (2018) Social network plasticity decreases disease transmission in a eusocial insect. Science 362:941–945. https://doi.org/10.1126/science.aat4793

Stroeymeyt N, Jordan C, Mayer G, et al (2014) Seasonality in communication and collective decision-making in ants. Proc R Soc B Biol Sci 281:20133108. https://doi.org/10.1098/rspb.2013.3108

Stuble KL, Pelini SL, Diamond SE, et al (2013) Foraging by forest ants under experimental climatic warming: a test at two sites. Ecol Evol 3:482–491. https://doi.org/10.1002/ece3.473

Suarez AV, Yeh P, Case TJ (2005) Impacts of Argentine ants on avian nesting success. Insectes Sociaux 52:378–382. https://doi.org/10.1007/s00040-005-0824-y

Sunamura E, Suzuki S, Nishisue K, et al (2011) Combined use of a synthetic trail pheromone and insecticidal bait provides effective control of an invasive ant. Pest Manag Sci 67:1230–1236. https://doi.org/10.1002/ps.2172

Tanaka Y, Nishisue K, Sunamura E, et al (2009) Trail-following Disruption in the Invasive Argentine Ant with a Synthetic Trail Pheromone Component (Z)-9-hexadecenal. Sociobiology 54:15

Tatsuki S, Terayama M, Fukumoto T (2005) (54). Behavior-disrupting agent and behavior disrupting method of Argentine ant. U.S. patent 0209344 A1

Trigos-Peral G, Abril S, Angulo E (2021) Behavioral responses to numerical differences when two invasive ants meet: the case of *Lasius neglectus* and *Linepithema humile*. Biol Invasions 23:935–953. https://doi.org/10.1007/s10530-020-02412-4

Tsutsui ND, Suarez AV (2003) The Colony Structure and Population Biology of Invasive Ants.
Conserv Biol 17:48–58. https://doi.org/10.1046/j.1523-1739.2003.02018.x

Valenta K, Schmitt MH, Ayasse M, Nevo O (2021) The sensory ecology of fear: African
elephants show aversion to olfactory predator signals. Conserv Sci Pract 3:e333.
https://doi.org/10.1111/csp2.333

von Thienen W, Metzler D, Witte V (2016) How memory and motivation modulate the
responses to trail pheromones in three ant species. Behav Ecol Sociobiol 70:393–407.
https://doi.org/10.1007/s00265-016-2059-5

Webb J, Legge S, Tuft K, et al (2015) Can we mitigate cane toad impacts on northern quolls?
Charles Darwin University| Final report. 20

Wenig K, Bach R, Czaczkes TJ (2021) Hard limits to cognitive flexibility: ants can learn to
ignore but not avoid pheromone trails. J Exp Biol 224:jeb242454.
https://doi.org/10.1242/jeb.242454

Wetterer JK, Wetterer AL, Hebard E (2001) Impact of the Argentine Ant, *Linepithema humile*
on the Native Ants of Santa Cruz Island, California. 38:13

Wetterer JK, Wild AL, Suarez AV, et al (2009) Worldwide spread of the Argentine ant,
*Linepithema humile* (Hymenoptera: Formicidae). Myrmecol. News 12: 187-194

Williams DF, Collins HL, Oi DH (2001) (Hymenoptera: Formicidae): An Historical Perspective
of Treatment Programs and the Development of Chemical Baits for Control. Am
Entomol 47:14

Wystrach A, Buehlmann C, Schwarz S, et al (2020) Rapid Aversive and Memory Trace
Learning during Route Navigation in Desert Ants. Curr Biol 30:1927-1933.e2.
https://doi.org/10.1016/j.cub.2020.02.082

762