Light targeting the melanopic system suppresses melatonin, but does not alter sleepiness, vigilance, sensory processing, or sleep

Christine Blume[1,2][0000-0003-2328-9612], Maria Niedernhuber[3][0000-0002-4459-6796],

Manuel Spitschan[1,2,4,5][0000-0002-8572-9268], Helen C. Slawik[6], Martin P. Meyer[6][0000-0002-8041-5884],

Tristan Bekinschtein[3]§[0000-0001-5501-8628], Christian Cajochen[1,2]§[0000-0003-2699-7171]

§ These authors contributed equally

[1]Centre for Chronobiology, Psychiatric Hospital of the University of Basel, Basel, Switzerland

[2]Transfaculty Research Platform Molecular and Cognitive Neurosciences, University of Basel, Basel, Switzerland

[3]Consciousness and Cognition Lab, Department of Psychology, University of Cambridge, Cambridge, United Kingdom

[4]Translational Sensory and Circadian Neuroscience, Max Planck Institute for Biological Cybernetics, Tübingen, Germany

[5]TUM Department of Sport and Health Sciences (TUM SG), Technical University of Munich, Munich, Germany

[6]Psychiatric Hospital of the University of Basel, Basel, Switzerland

Corresponding Author:

Dr. Christine Blume

Centre for Chronobiology

Psychiatric Hospital of the University of Basel

Wilhelm-Klein-Str. 27

CH-4002 Basel, Switzerland

Email: christine.blume@unibas.ch

Authors' email addresses:

mn473@cam.ac.uk

tb419@cam.ac.uk

manuel.spitschan@tuebingen.mpg.de

helenchristina.slawik@upk.ch

martin.meyer@upk.ch

christian.cajochen@upk.ch

Funding

The project was supported by the following grants awarded to CB: a fellowship of the Austrian Science Fund (FWF; J-4243), a grant from the Research Fund for Junior Researchers of the University of Basel, and funds from the Freiwillige Akademische Gesellschaft (FAG), the Novartis Foundation for Biological-Medical Research, and the Psychiatric Hospital of the University of Basel (UPK).

Abstract

Study objectives: Pre-sleep exposure to short-wavelength light acutely suppresses melatonin, increases vigilance, and decrease sleepiness with the activating effects extending to sleep. The effects have mainly been attributed to melanopic effects of light, but mechanistic insights are missing. Thus, we investigated whether two metameric light conditions designed to only differ in their melanopic effects (123 vs. 59 lux melanopic EDI) also differentially affect sleep besides melatonin effects. Beyond this, we also studied whether the light conditions modulate sensory processing during wakefulness and sleep.

Methods: In a preregistered study, twenty-nine healthy participants aged 18-30 y (15 women) were exposed to two metameric light conditions (high- vs. low-melanopic, differing by a factor of 2x, 7-day washout period) for 1 hour prior to their habitual bed time. This was followed by an 8-h sleep opportunity with polysomnography. Objective sleep measurements were complemented by self-reported sleep evaluation after wake-up. Salivary melatonin levels, subjective sleepiness, and behavioural vigilance were sampled in regular intervals. Sensory processing was evaluated using an oddball paradigm participants completed during the light exposure in the evening and the following sleep episode. Specifically, we were interested in event-related potentials (ERPs) and time-frequency responses related to violations of expectations, that is, the mismatch response.

Results: Despite ≈14% melatonin suppression in the high- compared to the low-melanopic condition, light conditions did not differentially affect objectively assessed sleep, self-perceived sleep quality, sleepiness, or vigilance. A neural mismatch response was evident during all sleep stages but not differentially modulated by pre-sleep light.

Conclusions: Suppression of melatonin by light targeting the melanopic system does not necessarily translate to acutely altered levels of vigilance or sleepiness and neither to changes in sleep, sleep quality, or basic sensory processing. This may suggest that an interaction between melanopsin and cone-rod signals could be at work for such effects to occur.

Keywords: artificial light, melanopsin, melatonin, sleep, sensory processing, circadian timing system

Introduction

The effects of short-wavelength light on sleep and the circadian timing system are thought to be primarily mediated by specialised retinal ganglion cells that contain the photopigment melanopsin, wherefore they are intrinsically sensitive to light between 460 and 480 nm (intrinsically photosensitive retinal ganglion cells [ipRGCs]; e.g. Berson, Dunn, & Takao, 2002; Dacey et al., 2005; Lockley, Brainard, & Czeisler, 2003). Short-wavelength light exposure in the evening can acutely suppress melatonin secretion, increase alertness, and delay the onset of melatonin secretion. However, although the neuroendocrine sensitivity by light is characterised by considerable interindividual variability (Phillips et al., 2019). Moreover, evening light exposure has been shown to improve performance on tasks requiring sustained attention acutely (e.g. Lockley et al., 2006; Cajochen et al., 2011) and increase the available processing resources on a simple cognitive task as indicated by the amplitude of the P300 component of the human event-related potential (ERP; Okamoto & Nakagawa, 2013). Further, it can decrease sleepiness in the evening, prolong sleep onset latency and consequently increase sleepiness in the morning (cf. e.g. Zeitzer, Dijk, Kronauer, Brown, & Czeisler, 2000; Chang, Aeschbach, Duffy, & Czeisler, 2015; Santhi et al., 2012; Dijk, Cajochen, & Borbély, 1991). Moreover, evaluating sleep objectively with electroencephalography (EEG), Münch et al. (2006) demonstrated that exposure to short-wavelength light compared to longer wavelengths leads to decreased slow-wave activity (SWA) during the first sleep cycle. The authors concluded this finding to reflect that the alerting effect of the light persisted into sleep, which is well in line with findings by Chellappa et al. (2013), who reported a decrease in homeostatic sleep pressure following 'blue' light exposure in the evening. Chang et al. (2015) additionally found the use of light-emitting e-readers to decrease and delay REM sleep, but no change in total sleep time (TST), sleep efficiency, or the duration of non-REM (NREM) sleep compared to reading a printed book.

While many of the studies point to a special relevance of the short-wavelength-sensitive melanopic system in mediating the abovementioned effects, only recently so-called metameric light conditions have allowed to specifically target specific classes of retinal receptors in selected, isolated fashion while producing no responses in other classes (Spitschan & Woelders, 2018). Such studies yield strong mechanistic insights regarding the relevance of specific receptors and thus light characteristics.

Contrasting two metameric light conditions (73.5 photopic lux), that is, high- vs. low-melanopic light, Allen, Hazelhoff, Martial, Cajochen, and Lucas (2018) demonstrated that 5-h exposure to high-melanopic light during the evening (18:00-23:00) resulted in stronger suppression of melatonin and a decrease of subjective sleepiness. Similarly, Souman et al. (2018a) found that 3-h high-melanopic exposure from two hours before to one hour after habitual bed time suppressed melatonin compared with a low-melanopic (both low-/high-melanopic: 175 photopic lux) and a dim light condition, however without affecting alertness. Thus, while high-melanopic light is effective in suppressing melatonin secretion, the effects on cognition are less clear. Whether the effects of light on subsequent sleep that have been reported in other studies can largely be attributed to short-wavelength light acting on ipRGCs or rather result from an interaction with photopic or cone-mediated light characteristics is still unknown. Thus, this project aimed at investigating the effects of two metameric light conditions designed to only differ in their effects on ipRGCs but not cones on subjective and objective sleep parameters beyond melatonin suppression. We expected melatonin to be suppressed and SWA in the first sleep cycle to be decreased more effectively by high compared to low-melanopic light. Besides this, we explored the effects on sleep, sleepiness, and behavioural vigilance using self-report scales and objective measurements, respectively.

Going beyond classic, sleep parameters (i.e., SWA, sleep architecture) that describe global brain states and reflect a *macroscopic* perspective, we were also interested in whether and how light exposure alters brain processes on the *microscopic* level. More specifically, we propose this could inform whether high-melanopic artificial light exposure leads to more 'wake-like' processing during sleep that underlies the observed effects on 'global brain states' and sleep quality. To this end, we investigated whether metameric light conditions would differentially alter basic sensory processing during wakefulness and subsequent sleep. Classically, basic sensory processing has been studied using so-called oddball paradigms, where rare deviating tones are included in a sequence of frequent standard tones. These deviating stimuli are well-known to give rise to a distinct component in the human event-related potential (ERP), the so-called mismatch negativity (MMN; Sams, Alho, & Näätänen, 1983; Näätänen, Paavilainen, Tiitinen, Jiang, & Alho, 1993; for a review see Näätänen, Paavilainen, Rinne, & Alho, 2007). Although the MMN is often considered to reflect a low-level or 'pre-attentive' prediction error

(Näätänen et al., 1993), it is best explained by an active 'top-down' predictive mechanism and only to a lesser extent by passive sensory adaptation and violations thereof (Wacongne, Changeux, & Dehaene, 2012). Additionally, the MMN amplitude has been shown to be modulated by (selective) attention in some studies, although it does not require (waking) attention (Woldorff et al., 1993; Alain & Woods, 1997; Woldorff, Hillyard, Gallen, Hampson, & Bloom, 1998). Besides this, Vandewalle et al. (2006) found light (during daytime) to enhance brain responses in cortical areas that support attentional oddball effects. Thus, we here investigated whether 1-h pre-sleep exposure to two metameric light conditions would differentially alter early sensory processing as reflected in the mismatch response (i) during the light exposure and (ii) during a subsequent 8-h sleep opportunity (cf. Figure 1). Specifically, we hypothesised that the light exposure would differentially affect the active prediction system during sleep thus yielding stronger ERP responses in the high-melanopic condition.

Methods and Materials

Pre-registration

This project was pre-registered on Open Science Framework (OSF) including details about the light exposure, the analytic strategy as well as the code for the acoustic stimulation paradigm (https://www.doi.org/10.17605/OSF.IO/M23EH). As it is classified as a clinical study in Switzerland, it has also been registered on the German Clinical Trials Register (DRKS00023602). For reasons of conciseness, we will address the pre-registered analyses relating to global and omission effects as well as analyses including the data from the post-sleep recordings in (a) separate publication(s).

Participants

Following screening procedures, between April 2019 and May 2020 we had invited 40 volunteers to participate in the study. All volunteers had provided written informed consent before final study inclusion. Ten volunteers dropped out or were excluded (e.g., due to non-adherence to the study protocol) after the adaptation night or the first experimental night. The final study sample thus comprised 30 participants (15 women). Data from one male participant had to be excluded as melatonin analyses (see below) revealed that he was a high-secretor (i.e., no value below 29 pg/ml between 5 hours and 15 minutes prior to habitual bed time [HBT]), wherefore the final sample comprised 29 participants (15

women). German was the volunteers' mother tongue (or comparable level) and they scored normally on questionnaires screening for sleep disorders (Pittsburgh Sleep Quality Index [PSQI], inclusion if sum values ≤ 5; Buysse, Reynolds, Monk, Berman, & Kupfer, 1989) and psychiatric symptoms (Brief Symptom Inventory [BSI]; sum scores on nine subscales and 'global severity index' outside clinical range; Franke & Derogatis, 2000). Furthermore, they did not habitually sleep less than 6 or more than 10 hours on work days and did not have an extreme chronotype according to the Munich Chronotype Questionnaire (MCTQ; Roenneberg, Wirz-Justice, & Merrow, 2003) or the Morningness-Eveningness Questionnaire (MEQ - German version; Horne & Östberg, 1976). The average self-selected HBT was 11:05 pm (range 10:15 pm – 00:00). Participants had to be right-handed, between 18 and 30 years of age (M = 23.2±2.8 years; range 19-29 years), and did not report any vision disorders nor hearing problems. They had normal or corrected-to-normal vision, and in case they wore glasses, the glasses did not have a blue light filter. All volunteers were healthy non-smokers with a body-mass-index (BMI) above 18.5 and below 25 (i.e., normal weight) and did not report any previous brain injury or relevant medical history regarding neurological or psychiatric disorders. To ensure participants did not take drugs, volunteers had to take a urine-based multi-drug panel (nal von Minden GmbH, Germany) screening for cocaine, benzodiazepines, amphetamines, tetrahydrocannabinol (THC), morphine, and methadone on the evening of every visit to the laboratory. Participants were seen by a study physician and only included if s/he did not identify reasons for exclusion. Shift-work within the three months or travelling across >2 time zones in the month prior to study participation served as additional exclusion criteria. The study protocol was approved by the ethics commission (Ethikkommission Nordwest- und Zentralschweiz; 2018-02003). Upon completion of the ambulatory and laboratory parts, participants received a remuneration of CHF 450.

Study procedures

Data were recorded continuously between April 2019 and May 2020 with a break between mid-March and mid-May 2020 due to the Coronavirus pandemic. The experimental protocol comprised an adaptation night as well as two experimental lab visits, which were interspaced by two ambulatory phases (cf. Figure 1). During the adaptation night, volunteers slept in the laboratory during the pre-defined habitual sleep window with an 8-h sleep opportunity with full polysomnography equipment (for

details see below) to familiarise them with the laboratory setting and rule out any undiscovered sleep disorders. Following the adaptation night, they kept a pre-defined constant sleep-wake cycle for the next 7 days, which was verified by sleep logs and wrist actigraphy (ActiGraph LL.C, Pensacola, FL 32502, USA). Specifically, they had to switch off lights at a specific time (±30min to the agreed [HBT]) and get up at a specific time (±30 min habitual wake time) while maintaining an 8:16h wake-sleep cycle. In one case, the HBT was delayed by 30 minutes on the second visit as the participant had consistently gone to sleep at the end of the agreed time window for lights off. In another case, HBT was delayed due to technical failure of the screen. All participants then came to the laboratory for the first experimental visit (see below), after which they underwent the ambulatory protocol for a second time before coming to the laboratory for a second and final experimental visit.

Figure 1. *Experimental protocol for a participant with a habitual bed time (HBT) at 00:00. An adaptation night was followed by two experimental visits, which were spaced by one week and took place on the same day of the week. For 7 days before the first experimental visit and in between the two visits volunteers adhered to fixed bed and wake times to stabilise their circadian rhythms. For the experimental visits, participants arrived at the laboratory 5h 15min before their HBT, at the latest at 5:45 pm. From 6 pm onwards melatonin samples (red stars) were collected every 30 min. From just before the light exposure (1h 50 min before until 50 min before HBT) until the end of the protocol, they rated their sleepiness on the Karolinska Sleepiness Scale (blue triangles) with each melatonin sample. Additionally, there were 4 psychomotor vigilance tests (PVT, 10 min). The order of the light exposure conditions (high- vs. low-melanopic) was counter-balanced across participants and sexes. The red arrow marks the point of inclusion in the study. LE = Light exposure.*

Experimental nights took place on the same day of the week (i.e., Monday through Friday) and were usually spaced by exactly one week. In three cases, the two experimental visits were spaced by eleven, 14, and 15 days, respectively, due to unforeseen circumstances, wherefore the visits took place on a different day of the week in two cases. Only in one case, the two nights were spaced by 62 days, because the first night took place just before the laboratory was closed due to the Coronavirus pandemic and the second night took place right afterwards. Importantly, if the adaptation night and experimental visits were not exactly spaced by seven days, participants still had to adhere to a 7-day ambulatory phase to stabilise the sleep-wake cycle. We aimed at counter-balancing the order of the conditions across participants and gender subgroups. Drop-outs precluded counter-balancing among gender subgroups. Thus, the high-melanopic condition was the first for 8 men and 6 women.

Experimental visits

On the evenings of both experimental visits, which only differed regarding the light condition (i.e., mel-high [123 lux melanopic EDI] or mel-low [59 lux melanopic EDI], cf. Table 3), volunteers came to the lab 5 h 15 min before HBT or at 5:45 pm at the latest. Upon arrival, the room lighting was characterised by 3.2 cd/m^2 melanopic EDL (cf. Table 1). They were served a light dinner (i.e., a sandwich) and from then on, they were seated in a comfortable chair facing the screen used for light exposure (see below), which was switched off at this time. From 6 pm (irrespective of the individual HBT), salivary melatonin samples were taken using salivettes (Sarstedt®) in 30 min intervals until the light source was switched on, 5 min into the light exposure, and again in 30 min intervals until 15 min before HBT. Saliva samples were centrifuged and frozen at -18°C immediately for later assaying. We started mounting of the EEG 4h 20 min before HBT. From 3h 20min until 2h 20min prior to HBT volunteers were in dim light (0.3 cd/m^2 melanopic EDL; see Table 1 for details and Fig. 1) listening to podcasts while keeping their eyes open. Following dim light, they were in complete darkness for 30 min with eyes open. We ensured open eyes by regularly checking the EOG for blinks and reminded participants to keep their eyes open if necessary. Ten minutes into the dark adaptation, participants performed a 10 min auditory version of the psychomotor vigilance task (PVT; Dinges, Powell, & Computers, 1985) and a 3 min resting-state EEG with open eyes, still in darkness. Note that for conciseness, we will not report the results of the resting EEG in this manuscript. Afterwards, the dim

light was switched on again and served as a background light for the remainder of the evening. Just before the start of the light exposure, volunteers also completed the Karolinska Sleepiness Scale (KSS; Åkerstedt & Gillberg, 1990) for the first time. After this, they underwent 60 min of light exposure (i.e., either high- or low-melanopic screen light), during which they completed the auditory local-global oddball task (see below for details, overall duration approx. 41 minutes without breaks). This auditory task started approx. eight minutes into the light exposure following another resting-state EEG measurement to assess the acute effects of the light exposure. Further KSS ratings took place 5 minutes and again 30 minutes into the light exposure as well as just after it and 15 min before HBT. Note that the third KSS 30 minutes into the exposure was only introduced after the first 13 participants. Just after the light exposure, they completed another PVT as well as a resting-state EEG recording. During sleep, the auditory oddball task continued without interruptions. Importantly, participants were instructed to follow the instructions they had followed during wakefulness even during falling asleep and during the night. It has previously been shown that volunteers can follow such task instructions even during sleep (e.g., Andrillon, Poulsen, Hansen, Léger, & Kouider, 2016; Kouider, Andrillon, Barbosa, Goupil, & Bekinschtein, 2014). In the morning, just after having been woken up following an 8-h sleep opportunity, the volunteers resumed saliva sampling for later melatonin assays at 30 min intervals. Additionally, volunteers filled in the KSS just after waking up as well as the morning protocol of the sleep diary assessing sleep and awakening quality (Saletu, Wessely, Grünberger, & Schultes, 1987). Thirty minutes after wake-up, the morning block of assessments began. Here, two recordings of resting-state EEG activity and the completion of two more PVTs surrounded the oddball paradigm in the morning. Volunteers again indicated subjective sleepiness on the KSS after the completion of the oddball task. The morning recordings were completed under conditions of brighter light than the dim light in the evening (14.3 cd/m^2 melanopic EDL, cf. Table 1).

Light exposure protocol and conditions

Upon arrival at 5:45pm or 5 h 15 min before HBT, participants were seated at a table in *evening room light* (colour rendering index [CRI CIE Ra] 94; 8.66 cd/m^2). Starting 3 h 20 min before HBT, they were in *evening dim light* (CRI 93.88; 0.82 cd/m^2) until HBT, except for 30 min dark adaptation from 2 h 20 min until 1 h 50 min before HBT. For an overview of the radiance-derived α-opic responses and

equivalent daylight (D65) luminances for the background lighting (i.e., no experimental light) please see Table 1.

Table 1. Overview of the radiance-derived α-opic responses ($L_{e,\Omega}$; in mW/m^2*sr) for the background light in the evening and morning and radiance-derived α-opic equivalent daylight (D65) luminances (EDL, in cd/m^2) for the background light in the evening and morning. Photopic luminance was 8.7 cd/m^2 (evening room),0.8 cd/m^2 (evening dim) and 38.9 cd/m^2 (morning room). Radiance-derived chromaticity values (CIE 1931 xy standard observer for a 2° field) were x = 0.5 and y = 0.4 for all three background light settings. Values were calculated using the luox app (Spitschan, 2021).

	S-cone-opic		M-cone-opic		L-cone-opic		Rhodopic		**Melanopic**	
α-opic	$L_{e,\Omega}$	EDL	$L_{e,\Omega}$	EDL	$L_{e,\Omega}$	EDL	$L_{e,\Omega}$	EDL	$L_{e,\Omega}$	EDL
Evening room	1.2	1.5	9.6	6.6	14.3	8.8	5.9	4.0	4.2	3.2
Evening dim	0.1	0.2	0.9	0.6	1.4	0.8	0.6	0.4	0.4	0.3
Morning room	5.5	6.7	43.2	29.7	64.5	39.6	26.4	18.2	19.0	14.4

From 1 h 50 min until 50 min before HBT, participants were exposed to either *high-* (CRI 7.13; 59.87 photopic lux) or *low-melanopic* (CRI -112.25; 50.35 photopic lux) screen light. The two conditions were matched in the activation of the L, M and S cones as quantified using CIE S026 (Commission International de l'Eclairage (CIE), 2018), which incorporate 10° Stockman-Sharpe cone fundamentals. As a consequence, the photopic illuminance, quantified using the 2° V(λ) function, was slightly different between the two conditions. For a graphical illustration of the spectra, please see Figure 2 A. Please see Tables 2 and 3 for an overview of the radiance- and irradiance-derived α-opic responses and α-opic equivalent daylight (D65) irradiances/ illuminances for the two experimental screen light conditions, respectively. All participants underwent both experimental lighting conditions, the order was counterbalanced across participants. The light was presented on a 24-inch custom-made visual display (i.e., computer screen). This screen contains seven LEDs in total, two red LEDs with peak wavelengths (λ_p) of 630 and 660nm, two green LEDs (λ_p = 2x 520 and 550nm) and three blue LEDs (λ_p = 430, 450, and 470nm). The metameric stimuli were generated using constrained numerical optimisation implementing the method of silent substitution (Spitschan & Woelders, 2018; Spitschan, Aguirre, & Brainard, 2015) non-linear light sources using Matlab's 'fmincon' function (The Mathworks, Natick, MA).

*Figure 2. (**A**) Spectra of the two experimental screen light conditions. The mel-low condition is indicated by the solid line, the mel-high condition by the dashed line. For an overview of the spectral distributions for the two conditions reported in 1nm steps, please see the supplemental CSV file provided. (**B**) Photo illustrating the screen and the participant's position relative to it during the light exposure.*

Table 2. Overview of the radiance-derived α-opic responses ($L_{e,\Omega}$; in mW/m²*sr) for the two experimental screen light conditions and radiance-derived α-opic equivalent daylight (D65) luminances (EDL, in cd/m²). Photopic luminance was 10.4 cd/m² (mel-low) and 8.9 cd/m² (mel-high) and radiance-derived chromaticity values (CIE 1931 xy standard observer for a 2° field) were x = 0.3 and y = 0.2 for the mel-low and x = 0.3 and y = 0.2 for the mel-high condition. Values were calculated using the luox app (Spitschan, 2021).

	S-cone-opic		*M-cone-opic*		*L-cone-opic*		*Rhodopic*		***Melanopic***	
α-opic	$L_{e,\Omega}$	EDL	$L_{e,\Omega}$	EDL	$L_{e,\Omega}$	EDL	$L_{e,\Omega}$	EDL	$L_{e,\Omega}$	EDL
Mel-high screen light	22.1	27.1	14.1	9.7	17.5	10.7	15.7	10.8	15.0	11.3
Mel-low screen light	20.1	24.6	14.8	10.1	17.2	10.5	27.2	18.7	31.2	23.5
Contrast (mel-high – mel-low)/mel-low [%]	-9.3		4.7		-1.8		72.9		107.3	
Ratio mel-high/ mel-low	0.9		1.0		1.0		1.7		2.1	

Table 3. Overview of the irradiance-derived α-opic responses (Ee; in mW/m²) and equivalent daylight (D65) illuminances (in lux) for the two experimental screen light conditions (background light: evening dim light). Photopic illuminance was 59.9 lux (mel-low) and 50.4 lux (mel-high) and radiance-derived chromaticity values (CIE 1931 xy standard observer for a 2° field) were x = 0.3 and y = 0.2 for the mel-low and x = 0.3 and y = 0.2 for the mel-high condition. Values were calculated using the luox app (Spitschan, 2021).

α-opic	S-cone-opic E_e	EDI	M-cone-opic E_e	EDI	L-cone-opic E_e	EDI	Rhodopic E_e	EDI	*Melanopic* E_e	EDI
Mel-high screen light	103.5	126.7	79.9	54.9	96.9	59.5	142.7	98.4	162.8	122.7
Mel-low screen light	110.9	135.7	79.0	54.2	100.4	61.6	83.6	57.7	78.5	59.2
Contrast (mel-high – mel-low)/mel-low [%]	-6.7		1.1		-3.5		70.7		107.4	
Ratio mel-high/ mel-low	0.9		1.0		1.0		1.7		2.1	

A white background with a black circle in the centre subtending to a visual angle of 8.6° was presented on the screen. In the middle of the black circle, there was a small white circle, which served as a fixation point and subtended to a visual angle of 0.5°. Participants' eyes were at a distance of 70 cm from the screen. They were instructed to lean back in the chair in a comfortable position and fixate the centre of the screen for the duration of the light exposure. We refrained from using a chin rest to increase the ecological validity of the exposure. For an illustration of the display setup, please see Figure 2B. In the morning, participants woke up and they spent the time from wake-up until the end of the protocol in *morning room light* (CRI 94.38; 38.88 cd/m²). A supplemental CSV file contains the spectral distributions of radiance and irradiance for the *mel-high* and the *mel-low* light conditions at each wavelength between 380 and 780nm from the observer's point of view. It also contains the spectral radiance distributions for the background light, that is, the *evening room light*, the *evening dim light*, and the *morning room light*. Spectral measurements (range 380–780 nm; optical resolution of device 4.5 nm; reported wavelength sampling 1 nm) were performed with JETI spectroval 1501 (JETI Technische Instrumente GmbH, Jena, Germany).

Acoustic stimulation – Oddball paradigm

Recordings during wakefulness took place in the evening during light exposure to evaluate the acute effects of light exposure as well as in the mornings (delayed effects). We largely adopted the design by Strauss et al. (2015) have previously used, in which participants hear sequences of (in this case German) vowels (duration: 100 ms each). Note that we here focus on the early 'local' mismatch

effects, that is, the difference between *standards* and *deviants* (cf. Figure 2). Additionally, we also focus on data obtained during the evening and the night.

In each block, one vowel was the *standard* and presented more frequently than the other, *deviant* vowel. The location of articulation of vowels that were presented in the same sequence was maximally distant (for possible combinations see Fig. 2C). Stimuli were presented in trials of five vowels, where four *standard* vowels could either be followed by another *standard* (78% of the trials in "aaaaa" blocks, 11% of the trials in "aaaaB" blocks) or a *deviant* (78% in "aaaaB" and 11% in "aaaaa" blocks, cf. Fig. 2B). Each block started with a habituation phase of 20 (10 during sleep) trials that were excluded from the analyses. In type "aaaaa" blocks, participants were instructed to count the number of *deviants* whereas in the "aaaaB" blocks they counted the number of *standards* during wakefulness and sleep. The effect of interest, the 'local' mismatch effect, was the comparison between deviants and standards. It was previously been shown that participants can, to some extent, also follow such instructions during sleep (Kouider et al., 2014; Andrillon et al., 2016). One block comprised 146 (±1) trials (82 [±1] during sleep). Each vowel was separated from the following one by a short gap of 50 ms and trials were separated by an interval jittered between 850-1150 ms (in 50 ms steps, average 1000 ms). Two vowel combinations were used in the evening and two during the morning recordings. During sleep, participants heard the same vowel combinations as in the evening at first, and after approx. 40 min, when participants were expected to have reached N2 sleep, the other two vowel combinations were included. Our data confirmed that participants needed an average of 10 min to reach N2 sleep (range 0.5-32.5 min). Note that besides the 'normal' blocks, there were also 'control blocks', which however are not relevant in the context of the analyses presented here. In total, participants were presented with eight standard blocks and two control blocks during wakefulness (approx. 41 min without breaks, self-paced). During sleep, there was a total of 140 standard blocks and 52 control blocks. Blocks during sleep were separated by a 6-s inter-block-interval (IBI) to mark the transition between blocks. Thus, the stimulation during sleep summed up to 7.7 h.

*Figure 1: **Oddball Paradigm.** (**A**) Two different block types can serve as a standard. (**B**) Upper panel: In a sequence of five identical stimuli ("aaaaa") the fifth stimulus can be a standard (78%) or a deviant (11%). In "aaaaa" blocks, participants counted the deviants. Lower panel: the standard can also be an "aaaaB" block. In this case, participants counted the standards. (**C**) Possible vowel combinations. Modified from Strauss et al. (2015). Note that the investigated effect is equivalent to the 'local' mismatch effect in the context of the local-global oddball. The red rectangle marks the stimuli of interest for the analyses.*

Before the wakefulness recordings in the evening, participants were given printed instructions explaining the structure of the paradigm and it was ensured that they understood the structure and task instructions (i.e., the stimuli they had to count). For further details on stimulus generation and delivery, please see the supplemental material.

Data collection and reduction

Electrophysiological data

We recorded the EEG signal using a BrainProducts® (BrainProducts GmbH, Gilching, Germany) LiveAmp® device. Data were collected at a sampling rate of 250 Hz from 23 scalp electrodes, 2 ECG, and 2 chin EMG electrodes. Eye movements were recorded from electrodes placed according to the guidelines of the American Academy of Sleep Medicine and Iber (2007) below the outer canthus of the left and above the outer canthus of the right eye. For the polysomnography setup during the adaptation night and further details please see the supplemental material. Impedances were kept below 5kΩ and checked several times throughout the recording.

For EEG analyses we used Matlab 2019a and the Fieldtrip toolbox (Oostenveld, Fries, Maris, & Schoffelen, 2010) in the version distributed by the Salzburg Brain Dynamics lab (date downloaded: 31-05-2019). For details on EEG pre-processing, the analyses of event-related potentials (ERPs) and time-frequency responses as well as analysis of EEG slow-wave activity (SWA), please see the supplemental material.

Sleep staging

Sleep was scored semi-automatically by the Siesta Group GmbH (Vienna, Austria; Anderer et al., 2005; Anderer et al., 2010). To this end, we down-sampled data from electrodes F3, F4, C3, C4, O1, and O2 as well as the EMG and EOG data acquired according to Rechtschaffen & Kales recommendations to 128 Hz.

Sleep cycle detection

Sleep cycles were defined according to criteria adapted from Feinberg and Floyd (1979) and as implemented in the 'SleepCycles' package for R (R Core Team, 2015; Blume, 2020; Blume & Cajochen, 2020). We only considered the first three sleep cycles during every night in accordance with Chellappa et al. (2013).

Melatonin

Salivary samples were assayed for melatonin using direct double-antibody radioimmunoassays (RIA) which has been validated by gas chromatography-mass spectroscopy with an analytical least detectable dose of 0.9 pg/ml and an analytical sensitivity of 0.2 pg/ml (NovoLytiX GmbH, Pfeffingen, Switzerland; Weber, Schwander, Unger, & Meier, 1997). Samples from two volunteers were reanalysed using an enzyme-linked immunosorbent assay (ELISA, NovoLytiX GmbH, Pfeffingen, Switzerland) as two RIA measurements had led to diverging results. In one case, the profile that better resembled the ELISA results was used for further analyses, in the other case, the average of the two measurements was used. For further details on the handling of melatonin data, please see the supplemental material.

Subjective sleepiness

Subjective sleepiness was repeatedly assessed with the Karolinska Sleepiness Scale (Åkerstedt & Gillberg, 1990) throughout the evenings and mornings. Specifically, participants rated their sleepiness

on a 9-point Likert scale ranging from "extremely alert" to "very sleepy, great effort to keep awake, fighting sleep".

Behavioural vigilance

Behavioural vigilance was assessed twice in the evenings and twice in the mornings using a 6-minute auditory version of the psychomotor vigilance task (Dinges et al., 1985; Basner & Dinges, 2011). Please note that a visual task would always have involved participants looking at a screen, which required an auditory solution. Participants had to press the space key on a keyboard in response to a 65 dB tone. The interstimulus interval varied between 2 and 9 seconds. For pre-processing, we excluded non-valid trials, that is, trials with a response time (RT) <100 ms (Basner & Dinges, 2011). We then computed the median RT as well as the fastest and slowest 10% RTs for further processing.

Subjective sleep quality

Subjective sleep and awakening quality was assessed using a sleep diary by Saletu et al. (1987). More precisely, participants answered several questions in the mornings on a four-point Likert scale ("no", "somewhat", "moderate", "very much"), which were then combined into the subscales "sleep quality" and "awakening quality" by averaging. Additionally, the volunteers detailed the subjectively perceived number of awakenings, and the time it took them to fall asleep.

Visual comfort ratings

Following the light exposure, participants also rated the screen light regarding brightness, pleasantness, glare and warmth and indicated how activating they felt the light was on a 11-step Likert scale ("not at all" to "very much").

Statistical analyses

All statistical analyses were done in R version 4.0.4 except for the analyses of ERPs and time-frequency analyses, which were done in Fieldtrip®. All analyses except ERP and time-frequency analyses were two-sided with the critical alpha being set to 0.05. Analyses of sleep architecture and measures of objective sleep parameters as well as visual comfort were two-sided as there were directed a priori hypotheses for a subset of analyses only relating to sleep latency. If applicable, we adjusted *p*-values for multiple comparisons using the method by Benjamini and Hochberg (1995).

Melatonin

For the statistical evaluation of the melatonin data, we ran three separate analyses for (i) the six measurements preceding the light exposure, (ii) the four values between the onset of the light exposure and habitual bed time (HBT), and (iii) the four assessments in the morning following wake-up. Specifically, we used advanced non-parametric analyses as implemented in the 'nparLD' package (Noguchi, Gel, Brunner, & Konietschke, 2012) to assess the effect of the light condition (i.e., *mel-high* vs. *mel-low*) on melatonin secretion across time. A comparison of the time-series data between conditions was followed up by comparisons at each time point. Besides time series, we also compared the area under the curve (AUC) between the conditions. For the calculation of the AUC, we used the 'AUC' function implemented in the 'DescTools' package (Signorell & et al., 2021) with spline interpolation. We here report the ANOVA-type statistic (ATS) with degrees of freedom rounded to the next integer and relative treatment effects (RTE) as a measure of the effect size for the lighting condition effects. The RTE indicates the probability with which a randomly chosen value from the whole dataset is smaller than a randomly chosen value from a specific light condition.

Subjective sleepiness, behavioural vigilance, and visual comfort ratings

For the statistical analysis of subjectively reported sleepiness as assessed with the Karolinska Sleepiness Scale (KSS; Åkerstedt & Gillberg, 1990), behavioural vigilance assessed with the PVT (Dinges et al., 1985; Basner & Dinges, 2011), and visual comfort ratings for the two light sources, we also used the 'nparLD' package (Noguchi et al., 2012). More precisely, we evaluated condition differences and variability across time (if applicable).

For the KSS, we compared (i) the values before the light exposure, (ii) values from five minutes into the exposure until 15 minutes prior to HBT, and (iii) the values obtained in the morning. For the PVT, we compared all four assessments regarding median RT and fastest and slowest 10% RTs. Regarding visual comfort ratings, we compared ratings obtained after light exposure to each of the two lighting conditions.

Sleep architecture, objective sleep parameters, subjective sleep and awakening quality

Sleep architecture, objective sleep parameters, as well as subjective sleep and awakening quality were analysed with mixed linear models as implemented in R's 'lme4' package (Bates, Mächler, Bolker, & Walker, 2015). Specifically, we investigated the influence of the light exposure condition (i.e., *mel-*

low vs. *mel-high*; fixed effect) and the experimental visit (i.e., *visit 1* vs. *visit 2*; fixed effect). Participant ID was modelled as a random factor. Normal distribution of the data was assessed using Shapiro-Wilk normality tests and if violated, the outcome variable was transformed to ranks using R's 'rank' function prior to statistical analysis. The following variables were rank-transformed: sleep latencies, WASO, sleep efficiency, number of awakenings. For fixed effects, we report *t*-values along with degrees of freedom (*df*) rounded to the next integer as the Satterthwaite approximation results in decimals. We also report 95% confidence interval for the fixed effects, which were computed using the 'lme4' bootstrapping approach. In the case of a significant interaction, this was followed with contrasts using the 'emmeans' package (van Lenth, 2021) with Kenward-Roger degrees of freedom and a Tukey adjustment for multiple comparisons.

Sleep – EEG slow-wave activity

For the comparison of SWA during NREM sleep across the first three sleep cycles, we likewise used mixed linear models based on rank-transformed data. Again, participant ID was modelled as a random factor and the two light exposure conditions as well as the cycle number (i.e., first vs. second vs. third) were included as fixed effects with an interaction between the two.

Electrophysiological data

The mismatch effect was investigated within each light condition and compared between the *mel-low* and *mel-high* conditions for each sleep stage. For these comparisons, we used cluster-based permutation tests as implemented in the Fieldtrip toolbox (Maris & Oostenveld, 2007).

Results

Melatonin

On average, melatonin values from 5 min into the light exposure until habitual bed time were lower in the *mel-high* condition than in the *mel-low* condition ($F_{ATS}(1)=6.85$, *p*=.019, $RTE_{mel-low}$=0.537, cf. Figure 2). The RTE indicates that, with a probability of 53.7% (46.3%), a randomly chosen value from the whole dataset was smaller than a randomly chosen value from the *mel-low* (*mel-high*) condition. There was also a significant effect of the sampling time ($F_{ATS}(2)=31.18$, *p*<.001) and a trend for a sampling time × condition interaction ($F_{ATS}(3)=2.97$, *p*=.074). Follow-up analyses indicated that

melatonin values 30 minutes into ($F_{ATS}(1)$=13.88, p<.001, $RTE_{mel-low}$=0.559) as well as just after the light exposure ($F_{ATS}(1)$=8.03, p=.012, $RTE_{mel-low}$=0.554) differed significantly whereas values 5 minutes into the light exposure ($F_{ATS}(1)$=0.91, p=0.4, $RTE_{mel-low}$=0.521) and 15 min before HBT ($F_{ATS}(1)$=2.93, p=.14, $RTE_{mel-low}$=0.533) did not. Taking into account melatonin values at the two significant time points yielded an overall suppression of melatonin (i.e. difference between *mel-low* and *mel-high*) by 14.05% (SD = 34.7%).

There was no condition difference among the six values preceding the light exposure ($F_{ATS}(1)$=1.03, p=.4, $RTE_{mel-low}$=0.511) nor a time × condition interaction ($F_{ATS}(5)$=0.39, p=.8). As expected, values however varied across time ($F_{ATS}(3)$=68.18, p<.001). Likewise, there was no difference in melatonin values assessed in the morning following wake-up ($F_{ATS}(1)$=1.13, p=.4, $RTE_{mel-low}$=0.488) or time × condition interaction ($F_{ATS}(3)$=0.56, p=.66) but variability across time ($F_{ATS}(2)$=163.01, p<.001). Please see Figure 3A for a graphical illustration of the melatonin results.

In line with these results, the AUC for the melatonin values from 5 min into the light exposure until HBT also differed between the two light conditions with the AUC in the *mel-low* condition being larger than in the *mel-high* condition ($F_{ATS}(1)$=10.14, p=.004, $RTE_{mel-low}$=0.549). At the two time points (i.e., 30 minutes in to the light exposure and just after the exposure), the AUC was smaller by 14.6% in the *mel-high* relative to the *mel-low* condition. There were no differences between the AUCs for the values prior to the start of the light exposure ($F_{ATS}(1)$=0.77, p=.57, $RTE_{mel-low}$=0.518) nor the values in the morning ($F_{ATS}(1)$=0.27, p=.6, $RTE_{mel-low}$=0.49).

*Figure 2. (**A**) Time course of median salivary melatonin levels (n=29, pg/ml) for the melanopic high and melanopic low condition. Vertical bars indicate the 95% confidence intervals. The high-melanopic light was associated with a relevant suppression of melatonin secretion 30 minutes into light exposure and just after light exposure. HBT = habitual bed time. Asterisks indicate a significant difference between the light exposure conditions. (**B**) Subjective sleepiness (median and 95%*

CI) assessed with the Karolinska Sleepiness Scale. There were no significant differences in sleepiness between the two lighting conditions. Please note that the third value, 30 minutes into the light exposure was only available from 17 participants as this measurement had only been introduced later. (C) EEG slow-wave activity (SWA) between 0.5 and 4.5 Hz across the first three sleep cycles. Left panel: Boxplots of the SWA during the NREM and REM parts of each of the first three sleep cycles. Right panel: Average power for each percentile of each NREM and REM part of a sleep cycle. Vertical bars correspond to standard errors. Generally, SWA decreased across the cycles. There were no differences between the conditions. (D) Subjective sleep and awakening quality. The values are averaged values from subscales pertaining to sleep and awakening quality in the sleep diary by Saletu et al. (1987), respectively. There were no significant differences in self-perceived sleep or awakening quality. (E) Psychomotor vigilance task (PVT, 10 min auditory version). Median and 95% CI reaction times (left panel), slowest 10% (middle panel) and fastest 10% (right panel) of reaction times. There were no light condition differences and no interaction between assessment point and condition. However, there was a modulation across time for all three measures (cf. main text). In boxplots, the lower and upper hinges of the boxplot correspond to the 25% and 75% quartiles, the thick black line indicates the median. Whiskers extend to the lowest/largest value at most 1.5× the interquartile range (IQR) from the hinges. Gray circles in boxplots represent individual values of participants. Colour code: high melanopic condition: dark purple; low melanopic condition: coral.

Subjective sleepiness – Karolinska Sleepiness Scale

Subjective sleepiness during the light exposure prior to HBT did not differ between the two lighting conditions ($F_{ATS}(1)$=0.28, p=.69, $RTE_{mel-low}$=0.49, cf. Fig. 3B for an illustration of the time course of KSS measurements). The RTE indicates that, with a probability of 49% or 51%, a randomly chosen value from the whole dataset was smaller than a randomly chosen value from the *mel-low* or *mel-high* condition, respectively. While KSS values varied across time ($F_{ATS}(2)$=84.99, p<.001), there was no interaction between condition and time ($F_{ATS}(2)$=1.36, p=.36). As expected, KSS ratings before the beginning of the light exposure did not differ between the two lighting conditions ($F_{ATS}(1)$=0.12, p=.73, $RTE_{mel-low}$=0.488). Likewise, there was no condition difference in the morning ($F_{ATS}(1)$=2.11, p=.34, $RTE_{mel-low}$=0.477). While KSS values varied across time ($F_{ATS}(1)$=14.89, p<.001), there was no condition × time interaction ($F_{ATS}(2)$=1.66, p=.34).

Behavioural Vigilance – Psychomotor Vigilance Task

There were no condition differences regarding the median reaction time across the four measurement points ($F_{ATS}(1)$=0.73, p=.44, $RTE_{mel-low}$=0.492) with RTE suggesting that, with a probability of 49.2% (50.8%), a randomly chosen value from the whole dataset was smaller than a randomly chosen value from the *mel-low* (*mel-high*) condition. While RTs varied across time ($F_{ATS}(1)$=12.14, p<.001), there was no time × condition interaction ($F_{ATS}(2)$=1.03, p=.44, cf. Fig. 3E). Specifically, RT slowed from before (mdn = 182.25 ms) to after light exposure (mdn = 193 ms; $F_{ATS}(1)$=14.62, p<.001), became shorter in the morning again (mdn = 186.75 ms; $F_{ATS}(1)$=15.7, p<.001)

and slowed again until after the completion of the local-global oddball paradigm (mdn = 191.00 ms; $F_{ATS}(1){=}17.47$, $p{<}.001$).

Likewise, there was no condition difference for the 10% fastest RTs ($F_{ATS}(1){=}1.09$, $p{=}.38$, $RTE_{mel-low}{=}0.491$) and no time × condition interaction ($F_{ATS}(3){=}1.72$, $p{=}.22$), but RTs varied across time ($F_{ATS}(2){=}9.31$, $p{<}.001$, cf. Fig. 3E). RTs among the fastest 10% of the trials were again faster before the light exposure (mdn = 151.75 ms) than afterwards (mdn = 160.3 ms; $F_{ATS}(1){=}13.31$, $p{<}.001$). In the morning, volunteers' reaction times were again faster than in the evening (mdn = 155.65 ms; $F_{ATS}(1){=}29.54$, $p{<}.001$), and they slowed again in the morning from before to after the oddball paradigm recording (mdn = 158.0 ms; $F_{ATS}(1){=}11.54$, $p{=}.001$).

In line with these results, also the 10% slowest RTs did not differ between lighting conditions ($F_{ATS}(1){=}0.09$, $p{=}.8$, $RTE_{mel-low}{=}0.497$) and there was no time × condition interaction ($F_{ATS}(2){=}0.28$, $p{=}.8$). Again, the results confirmed a variability of the 10% slowest response times throughout the experimental visit ($F_{ATS}(2){=}11.79$, $p{<}.001$, cf. Fig. 3E). Specifically, we observed a slowing of RTs throughout the evening light exposure (mdn_{pre} = 227.25 ms; mdn_{post} = 242.60 ms; $F_{ATS}(1){=}8.59$, $p{=}.005$), RTs became again faster following sleep (mdn = 230.39 ms; $F_{ATS}(1){=}21.76$, $p{<}.001$) and then again slowed until the end of the experimental visit in the morning (mdn_{post} = 243.9 ms; $F_{ATS}(1){=}30.65$, $p{<}.001$).

Visual comfort

Participants perceived the *mel-high* condition as brighter than the *mel-low* condition ($F_{ATS}(1){=}4.62$, $p{=}.031$, $RTE_{mel-low}{=}0.432$, cf. Suppl. Fig. 2 for an illustration of the subjective ratings). There were no conditions differences in the perceived warmth of the light ($F_{ATS}(1){=}0.34$, $p{=}.56$, $RTE_{mel-low}{=}0.517$), its pleasantness ($F_{ATS}(1){=}0.1$, $p{=}.75$, $RTE_{mel-low}{=}0.51$), its glare ($F_{ATS}(1){=}2.53$, $p{=}.11$, $RTE_{mel-low}{=}0.451$), or regarding how activating it was perceived ($F_{ATS}(1){=}0.04$, $p{=}.85$, $RTE_{mel-low}{=}0.507$).

Subjective sleep and awakening quality

Subjective sleep quality did not differ between conditions (b = 0.11, $t(46)$ = 0.48, p = .64) nor the visit (b = -0.06, $t(46)$ = -0.26, p = .79) and neither was there an interaction between condition and visit (b = 0.14, $t(27)$ = 0.37, p = .71). Likewise, awakening quality did not differ between light exposure conditions (b = 0.5, $t(34)$ = 1.69, p = .1), nor between visits (b = 0.12, $t(34)$ = 1.49, p = .15), nor was

there an interaction between condition and visit ($b = -0.21$, $t(27) = -1.33$, $p = .19$). Supplementary Tables S1 and S2 provide a detailed overview of the results and Fig. 3D (right panel) a graphical illustration.

Objective sleep parameters & sleep architecture

Differences in latency to continuous 10 minutes of sleep did not differ between light exposure conditions ($b = -9.52$, $t(50) = -1.61$, $p = .12$; $mdn_{mel-low} = 5$ min, $mdn_{mel-high} = 7$ min; cf. Fig. 3D), but latency was decreased at the second experimental visit ($b = -12.96$, $t(50) = -2.2$, $p = .033$; $mdn_{V1} = 10$ min, $mdn_{V2} = 5.5$ min). There was no condition × visit interaction ($b = 3.8$, $t(27) = 0.4$, $p = .69$). Latency to N2 tended to be shorter in the *mel-low* condition (N2: $b = -11.27$, $t(48) = -1.93$, $p = .06$; $mdn_{mel-low} = 9$ min, $mdn_{mel-high} = 10.5$ min), and at the second visit (N2: $b = -17.03$, $t(48) = -2.92$, $p = .006$; $mdn_{V1} = 10.5$ min, $mdn_{V2} = 7.5$ min) but there was no interaction (N2: $b = 10.7$, $t(27) = 1.11$, $p = .28$). Latency to N1, N3, or REM did not differ between conditions (N1: $b = -2.75$, $t(46) = -0.43$, $p = .66$; N3: $b = 2.05$, $t(45) = 0.33$, $p = .74$; REM: $b = 1.08$, $t(53) = 0.17$, $p = .86$; cf. Fig. 3D) or experimental visits (N1: $b = -4.15$, $t(46) = -0.66$, $p = .52$; N3: $b = -6.69$, $t(45) = -1.09$, $p = .28$; REM: $b = 5.98$, $t(53) = 0.94$, $p = .34$) nor was there an interaction (N1: $b = -1.11$, $t(27) = -0.1$, $p = .92$; N3: $b = -6.08$, $t(27) = -0.59$, $p = .56$; REM: $b = -9.6$, $t(27) = -1$, $p = .32$). Supplementary Table S3 provides an overview of the results.

Percentages of N1, N2, N3, or REM did not differ between conditions (N1: $b = -1.85$, $t(33) = -0.93$, $p = .36$; N2: $b = 0.59$, $t(40) = 0.28$, $p = .78$; N3: $b = 2.65$, $t(34) = 1.14$, $p = .26$; REM: $b = -1.38$, $t(40) = -0.76$, $p = .46$) or experimental visits (N1: $b = -2.17$, $t(33) = -1.09$, $p = .28$; N2: $b = 0.40$, $t(40) = 0.19$, $p = .84$; N3: $b = 2.26$, $t(34) = 0.97$, $p = .34$; REM: $b = -0.49$, $t(40) = -0.27$, $p = .78$) nor was there an interaction (N1: $b = 2.17$, $t(27) = 0.57$, $p = .86$; N2: $b = -1.12$, $t(27) = -0.3$, $p = .76$; N3: $b = -3.83$, $t(27) = -0.88$, $p = .38$; REM: $b = 2.77$, $t(27) = 0.86$, $p = .40$). An overview of the results can be found in supplementary table S4.

Wake after sleep onset (WASO) did not differ between conditions ($b = 5.96$, $t(39) = 0.98$, $p = .34$) or visits ($b = 6.13$, $t(39) = 1.0$, $p = .32$) following rank transformation. There was no condition × visit interaction ($b = -18.99$, $t(39) = -1.73$, $p = .096$). For an overview of the results for WASO see supplementary table S5.

Neither did sleep efficiency differ between conditions (b = -7.44, t(39) = -1.23, p = .22) or visits (b = -5.34, t(39) = -0.92, p = .36). There was no condition × visit interaction (b = 20.51, t(27) = 1.88, p = .072). Suppl. table S6 provides an overview of the results for sleep efficiency.

Also the number of awakenings did not differ between conditions (b = 0.61, t(36) = 0.1, p = .92) or visits (b = -1.45, t(36) = -0.23, p = .82) and there was no interaction (b = -3.73, t(27) = -0.32, p = .76). Likewise, for the arousal index, that is, the number of arousals per total sleep time, there was no difference between conditions (b = -0.63, t(31) = -0.15, p = .88) or visits (b = -1.42, t(31) = -0.34, p = .74) and there was no condition × visit interaction (b = 0.29, t(27) = 0.036, p = .98). For an overview of the results for the number of awakenings and the arousal index see supplementary tables S7 and S8, respectively.

EEG slow-wave activity

There was no difference in SWA (0.5-4.5 Hz) between the two light exposure conditions (b = 52.45, t(1708) = 0.75, p = .46), neither was there an interaction between condition and cycle number (i.e., first, second, or third; b = -28.37, t(1708) = -0.87, p = .38). As expected, SWA decreased across the first three cycles (b = -165.4, t(1708) = -7.2, p < .001). For an overview of the results for SWA see Fig. 3C and supplementary table S9.

Event-related potentials (ERPs)

Please note that additional time-frequency analyses confirmed the ERP results detailed below. For the results of these analyses, please see the supplemental material.

Wakefulness

During wakefulness, there was a significant mismatch effect with deviants resulting in stronger early (mel-low: 52-148 ms, p = .004; mel-high: 52-148 ms, p = .004) as well as late (mel-low: 160-424 ms, p < .001; mel-high: 160-444 ms, p < .001) responses (cf. Fig. 4 A/F). There were no differences between the two light exposure conditions (all p > .28).

Figure 3: Event-related potential (ERP) effects for the mismatch effect during wakefulness (A, F), N1 (B, G), N2 (C, H), N3 (D, I), and REM (E, J) in the mel-high and mel-low conditions. The plots show the average potential across all electrodes that were part of the significant cluster. The shaded area corresponds to the standard error. Significant electrodes are indicated

Sleep

N1

During N1, there was a mismatch effect with an early (mel-low: 48-116 ms, $p = .029$ and 144-348 ms, $p < .001$; mel-high: 48-120 ms, $p = .029$ and 148-316 ms, $p < .001$) and a late (mel-low: 416-816 ms, $p < .001$; mel-high: 344-792 ms, $p < .001$) component (cf. Fig. 4B/G). There was no difference between the two light conditions (all $p > .22$).

N2

During N2, the mismatch effect persisted with early (mel-low: 0-372 ms, $p < .001$; mel-high: 0-364 ms, $p < .001$) and late components (mel-low: 308-844 ms, $p < .001$; mel-high: 324-863 ms, $p < .001$; cf. Fig. 4 C/H). There were no differences between the two light exposure conditions (all $p > .5$).

N3

The mismatch effect continued to be present with an early (mel-low: 24-320 ms, $p = .035$; mel-high: 44-308 ms, $p < .001$) and a late cluster (mel-low: 332-792 ms, $p < .001$; mel-high: 340-744 ms, $p < .001$; cf. Fig. 4 D/I). There was no difference between the two light conditions (no clusters).

REM

For the mismatch effect, there was again an early (mel-low: 0-312 ms, $p < .001$; mel-high: 48-124 ms, $p < .001$ and 140-364 ms, $p < .001$) and a late (mel-low: 324-900 ms, $p < .001$; mel-high: 332-900 ms, $p < .001$; cf. Fig. 4 E/J) component during REM. There was a difference between the light exposure conditions ($p = .038$) with the mel-low condition being associated with a larger amplitude between 0 and 92 ms above frontocentral electrodes.

Discussion

One-hour pre-sleep exposure to two metameric lighting conditions that specifically targeted intrinsically photosensitive retinal ganglion cells (ipRGCs; melanopic ratio 2.1, photopic illuminance ≈60 lux) differentially affected melatonin secretion. As expected, high-melanopic light acutely suppressed melatonin more effectively than low-melanopic light resulting in a relative decrease of about 14%. Going beyond earlier studies using metameric light, we also assessed the effects of pre-sleep light exposure on self-reported sleep quality as well as objective PSG-derived sleep variables. Sleep did not differ between the two light conditions despite their neuroendocrine effectiveness. In line with this, the two light conditions differentially modulated neither subjective sleepiness nor behavioural vigilance. We also found that basic sensory processing of a neural mismatch response was retained during all sleep stages but likewise not differentially affected by the light conditions. Our findings underline that melatonin suppression does not automatically translate into alterations of sleepiness or sleep, as well as the notion that other light characteristics besides melanopic effects may be involved in modulating the effects of evening light beyond the neuroendocrine response (Revell, Arendt, Fogg, & Skene, 2006; Brown, Thapan, Arendt, Revell, & Skene, 2021).

The absence of any acute or delayed (i.e., subsequent to light exposure) differential effects besides melatonin suppression is somewhat astonishing as it has repeatedly been reported that pre-sleep light exposure is associated with reduced subjective sleepiness and increased behavioural vigilance, longer sleep onset latencies, and the suppression of melatonin secretion (Chang et al., 2015; Cajochen, Kräuchi, Danilenko, & Wirz-Justice, 1998; for a review see Souman, Tinga, te Pas, van Ee, & Vlaskamp, 2018b; Gooley et al., 2011; Chellappa et al., 2011). In particular, the suppression of melatonin and possible downstream effects such as delays in sleep onset (Santhi et al., 2012) or alterations in sleep architecture and homeostatic sleep pressure (Münch et al., 2006), have largely been attributed to the melanopic rather than the cone-mediated system. This notion has received support from findings that melatonin suppression is particularly strong when short wavelength proportions are high (Cajochen et al., 2005; Chellappa et al., 2011; Revell et al., 2006) and that 'blue'-blocking glasses can mitigate neuroendocrine and alerting responses (van der Lely et al., 2015). A recent meta-analysis even confirmed that the relationship between melanopic illuminance and the delay in sleep onset follows a

dose-response relationship (Giménez et al., 2022). Further mechanistic evidence comes from studies using metameric conditions designed to affect the melanopic system only, as in the present study (Allen et al., 2018; Souman et al., 2018a). However, comparing conditions with high vs. low power between 450-500 lux at constant photopic illuminance (175 lux), Souman et al. (2018a) likewise found no differential effects on vigilance levels despite melatonin suppression reaching almost 50% (55 vs. 187 melanopic lux in the *mel-low* and the *mel-high* conditions, respectively). In line with this, Lok, van Koningsveld, Gordijn, Beersma, and Hut (2019) showed that light and melatonin affected behavioural vigilance independently, at least during daytime. In their study, melatonin ingestion in the afternoon increased sleepiness, but bright light exposure did not alter sleepiness nor behavioural vigilance. Future studies will have to verify whether this independence also holds true for the relationship between endogenous melatonin and light exposure in the evening. Taking this notion even a step further, whether vigilance, subjective sleepiness, melatonin secretion and eventually sleep may be sensitive to different light characteristics, is still an open question. Another reason for the absence of effects on subjective sleepiness ratings or vigilance as assessed with the PVT could be that the contrast in melanopic EDI was not strong enough or it requires longer light exposures. In particular, in the study using metameric light conditions by Allen et al. (2018), participants underwent 5 hours of light exposure while the contrast between low- and high-melanopic conditions was 24.7 vs. 77.7 melanopic lux, and light's alerting and melatonin supressing effects only appeared in the last third of the light exposure. Likewise, also the effects on slow-wave activity may be sensitive to exposure duration (Münch et al., 2006; Chellappa et al., 2013: both 2 hours, conditions were not metameric though), which we here found to be unmodulated by prior light exposure. In sum, one-hour exposure as in the present study may have been too short for differential effects beyond melatonin suppression to occur.

Besides this, cone and rod signals may well be relevant in addition to melanopic effects. As ipRGCs receive cone and rod inputs, it is plausible that the effects seen here are due to an interaction with melanopsin signals. In line with this, a recent study reported that behavioural vigilance (i.e., hit rate) was improved by both blue and red light compared to dimmer white light in day- and night shift workers (Figueiro & Pedler, 2020). The exact nature of this potential interaction between different light characteristics remains to be clarified in future studies. Whether and how metameric light affects sleep

beyond melatonin suppression has, to the best of our knowledge, not previously been investigated. While it has been known for a long time that melatonin is neither sufficient nor necessary for sleep (Cajochen, Kräuchi, & Wirz-Justice, 2003), our finding again underlines that sleep and melatonin suppression are not necessarily linked.

Regarding basic sensory processing as reflected by the mismatch response, we did not find evidence for differential effects of the low- vs. high-melanopic light exposure conditions. This is despite earlier research suggesting that particularly the melanopic photoreceptor system may mediate effects of short-wavelength light on cognition (Vandewalle et al., 2007). However, considering the absence of any differential effects on sleepiness, vigilance, or sleep apart from melatonin suppression, the absence of such effects may not be particularly surprising. Moreover, although bright light enhances responses in areas supporting attentional oddball effects (Vandewalle et al., 2006) and can affect later components such as the P300 (Okamoto & Nakagawa, 2013), especially early responses such as a mismatch effect may be relatively insensitive to rather subtle variations in the available processing resources resulting from light exposure. Future studies should evaluate whether later processing stages are sensitive to the effects of light and whether larger contrasts or variations of the effects on other retinal receptors are involved. Besides the absence of lighting-related differential effects, we found the mismatch response to be retained during all sleep stages, although the shape of the ERP considerably changed when participants fell asleep. This is in contrast to previous research by Strauss et al. (2015), who used the same auditory oddball paradigm during an afternoon nap, and who had concluded that sleep disrupts the mismatch response leaving only (passive) sensory adaptation mechanisms intact. Generally, with a high number of trials across a whole night and thus an excellent signal-to-noise ratio (SNR), we observed a stronger K-complex (KC)-like response elicited by deviants compared to standards, most prominently during N2 and deep N3 sleep. During N1 and REM, the effect was less strong albeit still present, which may be due to a decreased SNR in these sleep stages due to the presence of eye movements and larger trial-to-trial variability of the EEG signal. Interestingly, the observed pattern in ERPs as well as time-frequency analyses is very similar to what has been reported to be evoked by salient or subjectively relevant stimuli such as one's own name or an unfamiliar voice (Blume et al., 2016; Blume, del Giudice,

Wislowska, Heib, & Schabus, 2018). This suggests that deviants continued to be salient during all sleep stages.

Several limitations have to be considered that at the same time stake out the scope for future research. First, the sample only comprised young healthy sleepers. With increasing age or the presence of vulnerability factors, the sensitivity to light may well change. Thus, the results presented here should not be generalised beyond the investigated sample. Moreover, as participants only reported to the lab in the early evening, we cannot exclude that the light history during the day had an effect. This could be relevant as it has previously been shown that bright light during the day may decrease the susceptibility to light in the evening (Rångtell et al., 2016; Smith, Schoen, & Czeisler, 2004; Hébert, Martin, Lee, & Eastman, 2002). In our sample though, the laboratory conditions took place seven days apart in most participants, wherefore seasonal differences are unlikely. Furthermore, the time they reported having spent under the open sky before coming to the lab did not differ between the two conditions (mel-high: 102.4 min $\pm$ 101.9; mel-low: 99.1 min $\pm$ 72.6; $t(28) = 0.62$, $p = .8$). Likewise, there was no difference in the perceived brightness on a scale from 1 (very cloudy day) to 10 (bright summer's day) during this time (mel-high: 5.5 $\pm$ 2.7; mel-low: 5.8 min $\pm$ 2.6; $t(28) = -0.41$, $p = .68$). Thus, it seems unlikely that the individual prior light history affected our results.

To conclude, using two metameric light conditions that exclusively differ in their effects on ipRGCs, we find that, early sensory processing was not differentially modulated, neither during wakefulness nor sleep. Beyond this, our findings support the notion that suppression of melatonin in strictly controlled light settings does not automatically translate to altered levels of behavioural vigilance or experienced sleepiness and neither to changes in sleep or sleep quality. Thus, neuroendocrine and sleep-related mechanisms should be considered as relatively independent and investigated as such.

Conflicts of interest

CC and MS declare the following interests related to lighting. MS is currently an unpaid member of CIE Technical Committee TC 1-98 ("A Roadmap Toward Basing CIE Colorimetry on Cone Fundamentals"). MS was an unpaid advisor to the Division Reportership DR 6-45 of Division 3 ("Publication and maintenance of the CIE S026 Toolbox") and a member of the CIE Joint Technical Committee 9 on the definition of CIE S 026:2018. Since 2020, MS is an elected member of the Daylight Academy, an unpaid member of the Board of Advisors of the Center for Environmental Therapeutics. MS is named inventor on a patent application entitled on optimising non-linear multi-primary LED system filed by Oxford University Innovation Ltd. (US Patent Application no. 17/428,073, European Patent Application No 20705492.5). CC has had the following commercial interests related to lighting: honoraria, travel, accommodation and/or meals for invited keynote lectures, conference presentations or teaching from Toshiba Materials, Velux, Firalux, Lighting Europe, Electrosuisse, Novartis, Roche, Elite, Servier, and WIR Bank. CC is a member of the Daylight Academy.

Ethical approval

Ethical approval was provided by the cantonal ethics commission (Ethikommission Nordwest- und Zentralschweiz; 2018-02003). The study was conducted in accordance with Swiss law and the Declaration of Helsinki.

Author Contributions

CB conceptualised the project, supported by MN, TB, MS, and CC. CB acquired the data, conducted the analyses, and wrote the first draft of the manuscript. All authors provided critical review of the manuscript and approved the submitted version.

Acknowledgements

We thank all participants, who volunteered to participate in the project. We especially also thank the interns and master students Anaité del Río, Patricia Egli, Corinna M. Hofer, Jessica Jacobs, Melina Koller, Daniela Lindegger, Marlene Schmidt, and Natascha Stoffel without whom the study could not have been conducted. We also thank our colleague Isabel Schöllhorn for her valuable expertise and help with measuring the light conditions.

References

Åkerstedt, T., & Gillberg, M. (1990). Subjective and objective sleepiness in the active individual. *International Journal of Neuroscience, 52*(1-2), 29-37.

Alain, C., & Woods, D. L. (1997). Attention modulates auditory pattern memory as indexed by event-related brain potentials. *Psychophysiology, 34*(5), 534-546. doi:10.1111/j.1469-8986.1997.tb01740.x

Allen, A. E., Hazelhoff, E. M., Martial, F. P., Cajochen, C., & Lucas, R. J. (2018). Exploiting metamerism to regulate the impact of a visual display on alertness and melatonin suppression independent of visual appearance. *Sleep, 41*(8). doi:10.1093/sleep/zsy100

American Academy of Sleep Medicine, & Iber, C. (2007). *The AASM manual for the scoring of sleep and associated events: rules, terminology and technical specifications*: American Academy of Sleep Medicine.

Anderer, P., Gruber, G., Parapatics, S., Woertz, M., Miazhynskaia, T., Klösch, G., Saletu, B., Zeitlhofer, J., Barbanoj, M. J., & Danker-Hopfe, H. (2005). An E-health solution for automatic sleep classification according to Rechtschaffen and Kales: validation study of the Somnolyzer 24× 7 utilizing the Siesta database. *Neuropsychobiology, 51*(3), 115-133.

Anderer, P., Moreau, A., Woertz, M., Ross, M., Gruber, G., Parapatics, S., Loretz, E., Heller, E., Schmidt, A., Boeck, M., Moser, D., Kloesch, G., Saletu, B., Saletu-Zyhlarz, G. M., Danker-Hopfe, H., Zeitlhofer, J., & Dorffner, G. (2010). Computer-Assisted Sleep Classification according to the Standard of the American Academy of Sleep Medicine: Validation Study of the AASM Version of the Somnolyzer 24 × 7. *Neuropsychobiology, 62*(4), 250-264. Retrieved from https://www.karger.com/DOI/10.1159/000320864

Andrillon, T., Poulsen, A. T., Hansen, L. K., Léger, D., & Kouider, S. (2016). Neural Markers of Responsiveness to the Environment in Human Sleep. *The Journal of Neuroscience, 36*(24), 6583-6596. doi:10.1523/jneurosci.0902-16.2016

Basner, M., & Dinges, D. F. (2011). Maximizing sensitivity of the psychomotor vigilance test (PVT) to sleep loss. *Sleep, 34*(5), 581-591.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *2015, 67*(1), 48. doi:http://dx.doi.org/10.18637/jss.v067.i01

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B (Methodological), 57*(1), 289-300. doi:10.2307/2346101

Berson, D. M., Dunn, F. A., & Takao, M. (2002). Phototransduction by retinal ganglion cells that set the circadian clock. *Science, 295*(5557), 1070-1073.

Blume, C. (2020). SleepCycles: Sleep Cycle Detection (Version 1.0.1). Retrieved from https://CRAN.R-project.org/package=SleepCycles

Blume, C., & Cajochen, C. (2020). 'SleepCycles' package for R - A free software tool for the detection of sleep cycles from sleep staging. *Open Science Framework*. doi:https://doi.org/10.31219/osf.io/r2q8v

Blume, C., del Giudice, R., Lechinger, J., Wislowska, M., Heib, D. P., Hoedlmoser, K., & Schabus, M. (2016). Preferential processing of emotionally and self-relevant stimuli persists in unconscious N2 sleep. *Brain and Language*. doi:http://dx.doi.org/10.1016/j.bandl.2016.02.004

Blume, C., del Giudice, R., Wislowska, M., Heib, D. P., & Schabus, M. (2018). Standing sentinel during human sleep: Continued evaluation of environmental stimuli in the absence of consciousness. *Neuroimage, 178*, 638-648.

Brown, T. M., Thapan, K., Arendt, J., Revell, V. L., & Skene, D. J. (2021). S-cone contribution to the acute melatonin suppression response in humans. *Journal of pineal research, 71*(1), e12719. doi:https://doi.org/10.1111/jpi.12719

Buysse, D. J., Reynolds, C. F., Monk, T. H., Berman, S. R., & Kupfer, D. J. (1989). The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry Research, 28*(2), 193-213. doi:https://doi.org/10.1016/0165-1781(89)90047-4

Cajochen, C., Frey, S., Anders, D., Späti, J., Bues, M., Pross, A., Mager, R., Wirz-Justice, A., & Stefani, O. (2011). Evening exposure to a light-emitting diodes (LED)-backlit computer screen affects circadian physiology and cognitive performance. *Journal of Applied Physiology, 110*(5), 1432-1438.

Cajochen, C., Kräuchi, K., Danilenko, K. V., & Wirz-Justice, A. (1998). Evening administration of melatonin and bright light: Interactions on the EEG during sleep and wakefulness. *Journal of Sleep Research, 7*(3), 145-157. doi:https://doi.org/10.1046/j.1365-2869.1998.00106.x

Cajochen, C., Kräuchi, K., & Wirz-Justice, A. (2003). Role of melatonin in the regulation of human circadian rhythms and sleep. *Journal of Neuroendocrinology, 15*(4), 432-437.

Cajochen, C., Munch, M., Kobialka, S., Krauchi, K., Steiner, R., Oelhafen, P., Orgul, S., & Wirz-Justice, A. (2005). High sensitivity of human melatonin, alertness, thermoregulation, and heart rate to short wavelength light. *The Journal of Clinical Endocrinology & Metabolism, 90*(3), 1311-1316.

Chang, A.-M., Aeschbach, D., Duffy, J. F., & Czeisler, C. A. (2015). Evening use of light-emitting eReaders negatively affects sleep, circadian timing, and next-morning alertness. *Proceedings of the National Academy of Sciences, 112*(4), 1232-1237.

Chellappa, S. L., Steiner, R., Blattner, P., Oelhafen, P., Götz, T., & Cajochen, C. (2011). Non-visual effects of light on melatonin, alertness and cognitive performance: can blue-enriched light keep us alert? *PLoS ONE, 6*(1), e16429.

Chellappa, S. L., Steiner, R., Oelhafen, P., Lang, D., Götz, T., Krebs, J., & Cajochen, C. (2013). Acute exposure to evening blue-enriched light impacts on human sleep. *Journal of Sleep Research, 22*(5), 573-580.

Commission International de l'Eclairage (CIE). (2018). CIE S 026/E:2018: CIE System for Metrology of Optical Radiation for ipRGC-Influenced Responses to Light. In. Vienna: CIE Central Bureau.

Dacey, D. M., Liao, H.-W., Peterson, B. B., Robinson, F. R., Smith, V. C., Pokorny, J., Yau, K.-W., & Gamlin, P. D. (2005). Melanopsin-expressing ganglion cells in primate retina signal colour and irradiance and project to the LGN. *Nature, 433*(7027), 749-754. Retrieved from http://dx.doi.org/10.1038/nature03387

Dijk, D.-J., Cajochen, C., & Borbély, A. A. (1991). Effect of a single 3-hour exposure to bright light on core body temperature and sleep in humans. *Neuroscience Letters, 121*(1), 59-62. doi:https://doi.org/10.1016/0304-3940(91)90649-E

Dinges, D. F., Powell, J. W. J. B. R. M., Instruments,, & Computers. (1985). Microcomputer analyses of performance on a portable, simple visual RT task during sustained operations. *17*(6), 652-655. doi:10.3758/bf03200977

Feinberg, I., & Floyd, T. (1979). Systematic trends across the night in human sleep cycles. *Psychophysiology, 16*(3), 283-291.

Figueiro, M. G., & Pedler, D. (2020). Red light: A novel, non-pharmacological intervention to promote alertness in shift workers. *Journal of Safety Research, 74*, 169-177. doi:https://doi.org/10.1016/j.jsr.2020.06.003

Franke, G. H., & Derogatis, L. R. (2000). *BSI: Brief Symptom Inventory von LR Derogatis:(Kurzform der SCL-90-R): deutsche Version: Testmappe*: Beltz Test.

Giménez, M. C., Stefani, O., Cajochen, C., Lang, D., Deuring, G., & Schlangen, L. J. M. (2022). Predicting melatonin suppression by light in humans: Unifying photoreceptor-based equivalent daylight illuminances, spectral composition, timing and duration of light exposure. *Journal of pineal research, n/a*(n/a), e12786. doi:https://doi.org/10.1111/jpi.12786

Gooley, J. J., Chamberlain, K., Smith, K. A., Khalsa, S. B. S., Rajaratnam, S. M., Van Reen, E., Zeitzer, J. M., Czeisler, C. A., & Lockley, S. W. (2011). Exposure to room light before bedtime suppresses melatonin onset and shortens melatonin duration in humans. *The Journal of Clinical Endocrinology & Metabolism, 96*(3), E463-E472.

Hébert, M., Martin, S. K., Lee, C., & Eastman, C. I. (2002). The effects of prior light history on the suppression of melatonin by light in humans. *Journal of pineal research, 33*(4), 198-203. doi:10.1034/j.1600-079X.2002.01885.x

Horne, J. A., & Östberg, O. (1976). A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *International journal of chronobiology*.

Kouider, S., Andrillon, T., Barbosa, Leonardo S., Goupil, L., & Bekinschtein, Tristan A. (2014). Inducing Task-Relevant Responses to Speech in the Sleeping Brain. *Current Biology, 24*(18), 2208-2214. doi:http://dx.doi.org/10.1016/j.cub.2014.08.016

Lockley, S. W., Brainard, G. C., & Czeisler, C. A. (2003). High Sensitivity of the Human Circadian Melatonin Rhythm to Resetting by Short Wavelength Light. *The Journal of Clinical Endocrinology & Metabolism, 88*(9), 4502-4505. doi:10.1210/jc.2003-030570

Lockley, S. W., Evans, E. E., Scheer, F. A. J. L., Brainard, G. C., Czeisler, C. A., & Aeschbach, D. (2006). Short-Wavelength Sensitivity for the Direct Effects of Light on Alertness, Vigilance, and the Waking Electroencephalogram in Humans. *Sleep, 29*(2), 161-168. doi:10.1093/sleep/29.2.161

Lok, R., van Koningsveld, M. J., Gordijn, M. C. M., Beersma, D. G. M., & Hut, R. A. (2019). Daytime melatonin and light independently affect human alertness and body temperature. *J Pineal Res, 67*(1), e12583. doi:10.1111/jpi.12583

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG-and MEG-data. *Journal of Neuroscience Methods, 164*(1), 177-190.

Münch, M., Kobialka, S., Steiner, R., Oelhafen, P., Wirz-Justice, A., & Cajochen, C. (2006). Wavelength-dependent effects of evening light exposure on sleep architecture and sleep EEG power density in men. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology, 290*(5), R1421-R1428. doi:10.1152/ajpregu.00478.2005

Näätänen, R., Paavilainen, P., Rinne, T., & Alho, K. (2007). The mismatch negativity (MMN) in basic research of central auditory processing: a review. *Clin Neurophysiol, 118*(12), 2544-2590. doi:10.1016/j.clinph.2007.04.026

Näätänen, R., Paavilainen, P., Tiitinen, H., Jiang, D., & Alho, K. (1993). Attention and mismatch negativity. *Psychophysiology, 30*(5), 436-450. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/8416070

Noguchi, K., Gel, Y. R., Brunner, E., & Konietschke, F. (2012). nparLD: An R Software Package for the Nonparametric Analysis of Longitudinal Data in Factorial Experiments. *Journal of Statistical Software, 50*(12), 1-23. Retrieved from http://www.jstatsoft.org/v50/i12/

Okamoto, Y., & Nakagawa, S. (2013, 3-7 July 2013). *Effects of daytime exposures to short- and middle-wavelength lights on cortical activity during a cognitive task.* Paper presented at the 2013 35th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC).

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2010). FieldTrip: open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience, 2011*.

Phillips, A. J., Vidafar, P., Burns, A. C., McGlashan, E. M., Anderson, C., Rajaratnam, S. M., Lockley, S. W., & Cain, S. W. (2019). High sensitivity and interindividual variability in the response of the human circadian system to evening light. *Proceedings of the National Academy of Sciences, 116*(24), 12019-12024.

R Core Team. (2015). R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from https://www.R-project.org

Rångtell, F. H., Ekstrand, E., Rapp, L., Lagermalm, A., Liethof, L., Búcaro, M. O., Lingfors, D., Broman, J.-E., Schiöth, H. B., & Benedict, C. (2016). Two hours of evening reading on a self-luminous tablet vs. reading a physical book does not alter sleep after daytime bright light exposure. *Sleep medicine, 23*, 111-118. doi:https://doi.org/10.1016/j.sleep.2016.06.016

Revell, V. L., Arendt, J., Fogg, L. F., & Skene, D. J. (2006). Alerting effects of light are sensitive to very short wavelengths. *Neuroscience Letters, 399*(1), 96-100. doi:https://doi.org/10.1016/j.neulet.2006.01.032

Roenneberg, T., Wirz-Justice, A., & Merrow, M. (2003). Life between Clocks: Daily Temporal Patterns of Human Chronotypes. *Journal of Biological Rhythms, 18*(1), 80-90. doi:10.1177/0748730402239679

Saletu, B., Wessely, P., Grünberger, J., & Schultes, M. (1987). Erste klinische Erfahrungen mit einem neuen schlafanstoßenden Benzodiazepin, Cinolazepam, mittels eines Selbstbeurteilungsbogens für Schlaf-und Aufwachqualität (SSA). *Neuropsychiatrie, 1*(4), 169-176.

Sams, M., Alho, K., & Näätänen, R. (1983). Sequential effects on the ERP in discriminating two stimuli. *Biol Psychol, 17*(1), 41-58. doi:https://doi.org/10.1016/0301-0511(83)90065-0

Santhi, N., Thorne, H. C., van der Veen, D. R., Johnsen, S., Mills, S. L., Hommes, V., Schlangen, L. J., Archer, S. N., & Dijk, D. J. (2012). The spectral composition of evening light and individual

differences in the suppression of melatonin and delay of sleep in humans. *Journal of pineal research, 53*(1), 47-59.

Signorell, A., & et al. (2021). DescTools: Tools for descriptive statistics (Version R package version 0.99.40). Retrieved from https://cran.r-project.org/package=DescTools

Smith, K. A., Schoen, M. W., & Czeisler, C. A. (2004). Adaptation of Human Pineal Melatonin Suppression by Recent Photic History. *The Journal of Clinical Endocrinology & Metabolism, 89*(7), 3610-3614. doi:10.1210/jc.2003-032100

Souman, J. L., Borra, T., de Goijer, I., Schlangen, L. J. M., Vlaskamp, B. N. S., & Lucassen, M. P. (2018a). Spectral Tuning of White Light Allows for Strong Reduction in Melatonin Suppression without Changing Illumination Level or Color Temperature. *Journal of Biological Rhythms, 33*(4), 420-431. doi:10.1177/0748730418784041

Souman, J. L., Tinga, A. M., te Pas, S. F., van Ee, R., & Vlaskamp, B. N. S. (2018b). Acute alerting effects of light: A systematic literature review. *Behavioural brain research, 337*, 228-239. doi:https://doi.org/10.1016/j.bbr.2017.09.016

Spitschan, M. (2021). luox: Platform for calculating quantities related to light and lighting. Retrieved from https://luox.app/

Spitschan, M., Aguirre, G. K., & Brainard, D. H. (2015). Selective stimulation of penumbral cones reveals perception in the shadow of retinal blood vessels. *PLoS ONE, 10*(4), e0124328. doi:10.1371/journal.pone.0124328

Spitschan, M., & Woelders, T. (2018). The Method of Silent Substitution for Examining Melanopsin Contributions to Pupil Control. *Frontiers in Neurology, 9*. doi:10.3389/fneur.2018.00941

Strauss, M., Sitt, J. D., King, J.-R., Elbaz, M., Azizi, L., Buiatti, M., Naccache, L., van Wassenhove, V., & Dehaene, S. (2015). Disruption of hierarchical predictive coding during sleep. *Proceedings of the National Academy of Sciences, 112*(11), E1353-E1362.

van der Lely, S., Frey, S., Garbazza, C., Wirz-Justice, A., Jenni, O. G., Steiner, R., Wolf, S., Cajochen, C., Bromundt, V., & Schmidt, C. (2015). Blue Blocker Glasses as a Countermeasure for Alerting Effects of Evening Light-Emitting Diode Screen Exposure in Male Teenagers. *Journal of Adolescent Health, 56*(1), 113-119. doi:https://doi.org/10.1016/j.jadohealth.2014.08.002

van Lenth, R. (2021). emmeans: Estimated Marginal Means, aka Least-Squares Mean (Version 1.5.4). Retrieved from https://CRAN.R-project.org/package=emmeans

Vandewalle, G., Balteau, E., Phillips, C., Degueldre, C., Moreau, V., Sterpenich, V., Albouy, G., Darsaud, A., Desseilles, M., Dang-Vu, T. T., Peigneux, P., Luxen, A., Dijk, D.-J., & Maquet, P. (2006). Daytime Light Exposure Dynamically Enhances Brain Responses. *Current Biology, 16*(16), 1616-1621. doi:https://doi.org/10.1016/j.cub.2006.06.031

Vandewalle, G., Gais, S., Schabus, M., Balteau, E., Carrier, J., Darsaud, A., Sterpenich, V., Albouy, G., Dijk, D. J., & Maquet, P. (2007). Wavelength-Dependent Modulation of Brain Responses to a Working Memory Task by Daytime Light Exposure. *Cerebral Cortex, 17*(12), 2788-2795. doi:10.1093/cercor/bhm007

Wacongne, C., Changeux, J.-P., & Dehaene, S. (2012). A Neuronal Model of Predictive Coding Accounting for the Mismatch Negativity. *The Journal of Neuroscience, 32*(11), 3665-3678. doi:10.1523/jneurosci.5003-11.2012

Weber, J., Schwander, J., Unger, I., & Meier, D. (1997). A direct ultrasensitive RIA for the determination of melatonin in human saliva: comparison with serum levels. *J Sleep Res, 26*, 757.

Woldorff, M. G., Gallen, C. C., Hampson, S. A., Hillyard, S. A., Pantev, C., Sobel, D., & Bloom, F. E. (1993). Modulation of early sensory processing in human auditory cortex during auditory selective attention. *Proc Natl Acad Sci U S A, 90*(18), 8722-8726. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/8378354

Woldorff, M. G., Hillyard, S. A., Gallen, C. C., Hampson, S. R., & Bloom, F. E. (1998). Magnetoencephalographic recordings demonstrate attentional modulation of mismatch-related neural activity in human auditory cortex. *Psychophysiology, 35*(3), 283-292. doi:10.1017/s0048577298961601

Zeitzer, J. M., Dijk, D.-J., Kronauer, R. E., Brown, E. N., & Czeisler, C. A. (2000). Sensitivity of the human circadian pacemaker to nocturnal light: melatonin phase resetting and suppression. *The Journal of physiology, 526*(3), 695-702. doi:10.1111/j.1469-7793.2000.00695.x