An updated *C. elegans* body muscle transcriptome for studies in muscle

formation and function

Anna L. Schorr[1,2,3]

Email: aschorr@asu.edu

Alejandro Felix Mejia[3]

Email: Aflelixme@asu.edu

Martina Y. Miranda[4]

Email: MMiranda26@my.gcu.edu

Marco Mangone[1,2,3*]

* Corresponding author

Email: mangone@asu.edu

[1] Molecular and Cellular Biology Graduate Program, Arizona State University, Tempe,

AZ; [2] Virginia G. Piper Center for Personalized Diagnostics, The Biodesign Institute at

Arizona State University 1001 S McAllister Ave, Tempe, AZ; [3] School of Life Sciences,

Arizona State University, 751 E Lemon Mall, Tempe, AZ 85287; [4] Helios Scholars at the

Translational Genomics Research Institute, 445 N 5th St 4th Floor, Phoenix, AZ 85004

Running Title: Body muscle miRNAs in *C. elegans*

Keywords: miRNAs, 3'UTR, muscle tissue, interactome, *C. elegans*

ABSTRACT

The body muscle is an important tissue used in organisms for proper viability and

locomotion. The contractile unit of the muscle is the sarcomere, which is ultimately

responsible for the contraction reaction leading to movement. Although this tissue is

generally well studied and characterized, and many pathways have been elucidated

throughout the years, we still lack a comprehensive understanding of its transcriptome,

and how it controls muscle development and function. Here, we have updated a nuclear

FACS sorting-based approach to isolate and sequence a high-quality muscle

transcriptome from *C. elegans* mixed stage animals. We have identified 2,848 muscle-

specific protein-coding genes, including 78 transcription factors and 206 protein-coding

genes containing an RNA binding domain. We studied their interaction network,

performed a detailed promoter analysis, and identified novel muscle-specific *cis*-acting

elements. We have also identified 16 high-quality muscle-specific miRNAs, studied their

function *in vivo* using fluorochrome-based analyses, and developed the first high-quality

miRNA Interactome in a living organism, incorporating other muscle-specific datasets

produced by our lab and others.

Our study expands our understanding of how muscle tissue functions in *C. elegans* and

in turn, provide results that can in the future be applied to humans to study muscular-

related diseases.

INTRODUCTION

The body muscle is an important tissue used in organisms for proper viability and locomotion. Muscle tissue's structure, shape, and size change depending on its function and location. In humans, there are three main types of muscles: skeletal, smooth, and cardiac muscles, each responsible for a different form of contraction. Smooth and cardiac tissue contract involuntarily and are mainly located on the surface of organs. In contrast, skeletal muscles only contract voluntarily.

The contractile unit of the muscle is the sarcomere. It is composed of at least thirty different proteins, of which the most abundant are myosin and actin, and is ultimately responsible for the contraction reaction leading to movement.

The model organism *C. elegans* is ideal for studying muscle development and morphogenesis due to its well-characterized transcriptome, fully mapped cell lineage, and the availability of several methods for genetic engineering.

The physiology of its body muscle shares a high degree of similarity to the skeletal muscle of vertebrates with several mutually conserved genes (Lecroisey et al. 2007; Gieseler et al. 2017; Hrach et al. 2020; Ellwood et al. 2021b). In addition, *C. elegans* is a popular choice for studying disorders affecting humans, including muscular dystrophy (Hrach et al. 2020; Chaya et al. 2021; Ellwood et al. 2021a), since many human disease genes when introduced to worms phenocopy the symptoms (Markaki and Tavernarakis 2020; Natale et al. 2020; Chandler et al. 2021; Yue et al. 2021).

C. elegans possess only two large muscles: the pharynx and the body muscle. The pharynx is localized to the front of the animal and is responsible for food intake and physical crushing of the worm bacterial diet (Song and Avery 2013). It comprises eight

layers of non-striated muscles, surrounded by epithelial and neural tissue. The body

muscle tissue comprises 95 striated muscle cells localized throughout the animal's body

and is functionally equivalent to the vertebrate skeletal muscles (Gieseler et al. 2017).

The muscle's basic functional unit is the sarcomere. It is highly conserved between

nematodes and vertebrates and is composed of repeated units of a complex mesh of

thin (I-band) and thick (A-band) filaments. Conversely, the thin filaments are mainly

composed of actin. They anchor each sarcomere to the dense bodies and stabilize the

entire structure. Instead, thick filaments are mostly made of myosin and are attached to

the M-line, and mediate the tension generated during muscle contraction.

To properly develop and maintain functional body muscle tissue, gene

expression must be strictly regulated. Our group and others in the past finely

characterized the transcriptome of the *C. elegans* body muscle in the adult stage (Blazie

et al. 2015; Blazie et al. 2017) and during the embryonic development (Fox et al. 2007),

and identified important regulatory mechanisms used by muscle cells to maintain their

homeostasis.

In addition to the identification of important genes in this tissue, our group, and

others recently identified that tissue-specific alternative polyadenylation (APA), a poorly

characterized mechanism that produces genes with different 3'untranslated region

(3'UTR) isoforms, is an important regulatory mechanism used by cells to evade the

negative regulation effected by microRNAs (miRNAs) (Blazie et al. 2015; Blazie et al.

2017). Still, the complexity and depth of this form of regulation are currently unknown.

miRNAs are short ncRNAs that target regulatory elements in 3'UTRs, repressing gene

expression in the cytoplasm (Lee et al. 1993; Reinhart et al. 2000).

Since their discovery, several groups, including our lab, have shown that miRNAs

influence gene regulation considerably (Bartel 2018). The canonical miRNA biogenesis

pathway begins with the transcription of pri-miRNAs in the nucleus, which fold into a

hairpin structure (Bartel 2018). The microprocessor complex then cleaves the pri-

miRNAs into a shorter stem-loop structure, the pre-miRNAs, which are then exported to

the cytoplasm (Bartel 2018). Once in the cytoplasm the pre-miRNAs are cleaved into

mature miRNAs which associate with an Argonaute protein as part of the RNA-induced

silencing complex (RISC) (Bernstein et al. 2001; Kawamata and Tomari 2010; Bartel

2018). When loaded into the RISC, miRNAs bind to complementary target sequences

within the 3'UTRs of genes to be repressed (Bartel 2018). This implies that the 3'UTRs

and their isoforms are integral for miRNA-based regulation.

Several strategies have been developed to sequence tissue-specific

transcriptomes and miRNAs in *C. elegans*. These methodologies utilize either

fluorescence-activated cell sorting (FACS) (Fox et al. 2007; Haenni et al. 2012),

immunoprecipitation (Blazie et al. 2015; Blazie et al. 2017; Brosnan et al. 2021), or RNA

modification techniques (Alberti et al. 2018). Each of these approaches has unique

advantages but also present several caveats. Methodologies utilizing FACS allow for

high purity and stringency when isolating RNA from cells or nuclei. These approaches

rely on tissue-specific promoters to drive fluorochromes expression in cells or in nuclei.

After isolation, the fluorescent cells or nuclei may be readily sorted using FACS, then

either sequenced (Haenni et al. 2012) or processed using microarrays (Fox et al. 2007).

Immunoprecipitation and antibody-based approaches have also been widely

used to isolate the transcriptomes of specific tissues. This approach provides

considerably high yields of RNA compared to FACS-based methods and is generally

much more cost-efficient.

Brosnan *et al.* (2021) used an epitope labeled Argonaute protein expressed in

the specific *C. elegans* tissues to immunoprecipitate tissue-specific miRNAs populations

from the body muscle, intestine, and neurons (Brosnan et al. 2021).

Alberti *et al.* (2018) developed a methodology named "microRNAome by

methylation-dependent sequencing (Mime-seq)" to identify tissue-specific miRNA

populations from complex animals at the level of single-cell specificity (Alberti et al.

2018). The Mime-seq approach expresses a transgenic methyltransferase (Ath-HEN1)

from *Arabidopsis thaliana* in an animal system which allows the labeling of small RNAs.

Mime-seq has been utilized in *C. elegans* and *Drosophila* and improved tissue-specific

miRNA identification when compared with previous immunoprecipitation-based methods

(Alberti et al. 2018).

Our group also used an immunoprecipitation-based approach to identify miRNA

targets within the *C. elegans* intestine and body muscle (Kotagama et al. 2019), and

observed transcripts were targeted differentially between the two tissues. While

identifying the miRNA targets provided valuable information regarding tissue-specific

gene regulation, a major limitation of this study was that we were unable to identify the

specific miRNAs involved.

Here, we describe a novel nuclear FACS sorting-based approach, we named

"Nuc-Seq," to allow the isolation and sequencing of high-quality muscle transcriptomes

and miRNA populations from *C. elegans* mixed stage animals. Using this method, we

have identified 2,848 muscle-specific protein-coding genes, studied their interaction

network, performed a detailed promoter analysis, and identified novel muscle-specific

cis-acting elements. Using fluorochrome-based analyses *in viv*o, we developed a high-

quality muscle miRNA interactome, incorporating other muscle-specific datasets

produced by our lab and others.

RESULTS

Nuc-Seq: an updated approach to identify muscle-specific transcriptome

We wanted to improve the annotation of the muscle transcriptome by developing

a nuclear FACS-based strategy to isolate and sequence transcriptomes from *C. elegans*

body muscle nuclei. We named this approach 'Nuc-Seq'. Our final goal was to identify

the body muscle transcriptome and its miRNA population expressed explicitly in this

tissue.

To test the feasibility and optimize this approach, we first decided to perform

experiments using the *C. elegans* strain BN452, which ubiquitously expresses the

mCherry fluorochrome fused to the histone H2B ortholog gene *his-58* (Gomez-Saldivar

et al. 2016) (**Figure 1**). We fractionated this worm strain using mechanical stress and

separated two fractions: one cytoplasmic and one enriched with mCherry fluorescent

nuclei. (**Figure 1A-B and Materials and Methods**). We successfully tested these two

fractions for the presence of known cytoplasmic and nuclear genes using a Western blot

approach (**Figure 1C**). Importantly, these nuclei are stable over time with minimal

degradation after several hours (**Supplemental Figure S1**). We then performed the

FACS sorting step and successfully isolated a large pool of mCherry positive nuclei,

with over 34% of all the events being mCherry-positive (**Figure 1C**). We named this

updated approach Nuclear Sequencing (Nuc-Seq).

With these results in our hands, we decided to move forward and limit the

expression of the *mCherry*::*his-58* cassette in the body muscle tissue. We cloned the

his-58 gene, fused it to the mCherry fluorochrome, and forced its expression only in the

body muscle tissue using the promoter region of the *C. elegans* ortholog of the myosin

heavy chain gene (*myo-3*) (**Figure 2A upper panel**). As expected, the resultant

transgenic worm restricts the expression of the mCherry fluorochrome in the body

muscle nuclei (**Figure 2A lower panel**), with fewer fluorescent nuclei when compared

to the experiments in the BN452 strain (**Figure 2A**). We then performed the FACS

sorting step (**Figure 2B**). Although the number in sorted nuclei is significantly less than

those obtained with the positive control BN452, we could still isolate a large population

of mCherry-positive nuclei across from each of our replicates, which we then extracted

and sequenced RNA from.

The identification of the C. elegans body muscle transcriptome

We performed three sequencing reactions in duplicates (technical triplicates), two

from our body muscle-expressing nuclei (body muscle enriched), and one from nuclei

isolated from BN452 (negative controls) (total of six samples). We obtained

approximately 55M mappable reads for each sample (including our technical replicates).

For most of the samples, we could map more than 97% of the total reads

(**Supplemental Figure S2A**). The results obtained with our biological replicates

correlate well (**Supplemental Figure S1B-C**).

We have identified 2,849 protein-coding genes in the *C. elegans* body muscle tissue,

corresponding to ~14% of all *C. elegans* protein-coding genes (20,362 protein-coding

genes; WS250) (**Fig. 3 and Supplemental Table S1**). While some of our top hits have

not been characterized yet, as expected, we detected many known components of the

sarcomere, such as many of the proteins that form the thick filaments (myosin heavy

chain genes, *myo-3*, *unc-54*) and myosin light chain (*mlc-1*), and sarcomere

organization and structural factors (*unc-15*, *unc-45*), as well as those that form the thin

filaments, including several actin genes (*act-1, act-3, act-4*) and structural factors (*unc-*

60, *unc-73*) (**Main Figure 3A and Supplemental Figure S3)**. We also detect orthologs

of the tropomyosin gene *lev-11*, which in resting sarcomere blocks the binding of

myosin to actin, and other notable genes previously known to be in the *C. elegans* body

muscle tissue (**Main Figure 3A and Supplemental Figure S3)**. The genes in this tissue

are highly interconnected (**Main Figure 3B)** and produce several distinguished sub-

networks involved in body muscle formation and maintenance and protein production,

and energy storage (**Main Figure 3B)**.

Our dataset identified 78 transcription factors, including *mdl-1*, *skn-1*, *xbp-1*, *mxl-*

3, all previously known to be expressed in the body muscle tissue, and 207 non-

ribosomal proteins containing an RNA binding domain **Supplemental Table S1**. Within

this group, we identified K08D12.3, an orthologous of the human ZNF9, which is

mutated in myotonic dystrophy type 2, *mbl-1*, a MUSCLEBLIND-type of mRNA splicing

regulator required for muscle dense body organization, locomotion, and vulval

morphogenesis, and *etr-1*, an ortholog of the human CUG-binding protein CUGBP1,

required for embryonic muscle development, and has been implicated in myotonic

dystrophy **Supplemental Table S1**.

The C. elegans body muscle PROMOTERome

Next, we sought to study the core promoters of the genes identified in this study

to pinpoint potential motifs transactivated by muscle-specific transcription factors. We

have extracted 500 nucleotides upstream and 100 nucleotides downstream of the

transcription start site of the 2,849 genes we mapped in this tissue, and plotted their

nucleotide distribution (**Supplemental Figure S4**). We detected an enrichment of

thymidine nucleotides just upstream of the transcription start site in these genes

(**Supplemental Figure S4 Panel A**). Polythymine motifs (T-blocks) have been

previously reported in *C. elegans* core promoters, and their presence correlates with

gene expression levels (Grishkevich et al. 2011). We also performed a motif enrichment

analysis in these core promoters and identified several novel *cis*-acting elements

(**Supplemental Figure S4 Panel B** and **Supplemental Table S2**). We mapped 12

members of the human Krueppel-like transcription factors family (KLF), which are

important regulators of gene expression in vertebrate development and are involved in

muscle health and disease (Prosdocimo et al. 2015), Myogenin (MYOG), a muscle-

specific basic-helix-loop-helix (bHLH) transcription factor involved in muscle

development, myogenesis, and repair (Nabeshima et al. 1993), and NHLH1, a helix-

loop-helix transcription factor that is involved in growth and development of a wide

variety of tissues and species (Cogliati et al. 2002) and has been linked to muscle

growth and development (Miao et al. 2021). The complete list is shown in

Supplemental Table S2.

The C. elegans body muscle miRNAs

We then aimed to detect and study the body muscle-specific miRNA population

identified in our study. Using stringent parameters (**see Materials and Methods**), we

identified 18 miRNAs expressed in the *C. elegans* body muscle tissue (**Main Figure 5A**

and Supplemental Table S1). Several of these miRNAs have also been previously

described to be expressed in this tissue by other groups (Martinez et al. 2008), and

impairing their function either leads to defects in the muscular fiber formation or shows

an uncoordinated phenotype (Miska et al. 2007; Alberti et al. 2018) (**Supplemental**

Table S1). We validated the body muscle localization of some of these miRNAs using

previously developed strains expressing GFP under the control of specific miRNA

promoters (**Main Figure 4B**). Our most interconnected miRNA is *miR-5551*, a not

characterized *C. elegans*-specific miRNA that is predicted to target 26 muscle-

expressed protein-coding genes we detected in this tissue (**Main Figure 5A**). A GO

term analysis shows that its targets control contractile fibers, sarcomere, and myofibril

formation (**Supplemental Figure S5**). *Let-7* and the closely related *miR-241* are also

present. These miRNAs temporally regulate larval development and have been

previously identified in the pharyngeal and vulval muscles (Martinez et al. 2008;

Andachi and Kohara 2016). Among the predicted target genes, we identified several

genes in the DAF insulin pathway that control muscle health and aging (Mallick et al.

2020; Basu et al. 2021).

DISCUSSION

Here we have developed an updated FACS-based nuclear sorting method we
named Nuc-Seq, to identify tissue-specific transcriptomes from cell lysates and applied
it to determine the *C. elegans* muscle transcriptome. Nuc-Seq is robust, reproducible,
has minimal background noise, and allows the simultaneous identification of both
muscle-specific protein-coding genes and ncRNAs, such as miRNAs (**Main Figure 1**).
When applied to the *C. elegans* muscle tissue, Nuc-Seq allowed the precise
identification of 2,849 protein-coding genes, including 78 muscle transcription factors
and 207 non-ribosomal proteins containing an RNA-binding domain. Taken together,
the muscle transcriptome we identified covers ~14% of the entire *C. elegans* protein-
coding transcriptome (**Main Figure 3**). Some of these genes, such as *unc-54*, *mlc-1*,
unc-27, and others, have been previously identified in the muscle tissue and play an
important role in muscle contraction and muscle health in general. We have identified
almost all genes that form the sarcomere structure and are required for muscle
contraction (**Supplemental Figure S3**). When clustered using previously published
protein-protein interaction data (Remmelzwaal and Boxem 2019), the *C. elegans*
muscle transcriptome is highly interconnected (**Main Figure 3**), with several notable
subnetworks corresponding to mRNA processing, protein synthesis, ATP metabolism,
and muscle cell development (**Main Figure 3B**), all consistent with the function of this
tissue. We have also performed a promoter analysis, studying the promoter composition
of the genes identified in this study (**Supplemental Figure S4**). We have found a
Polythymine motif (T-blocks) directly upstream of the transcription start site and, using

prediction software (Meme Suite), we have identified many potential elements used by

muscle-specific transcription factors to control gene expression in this tissue.

Our approach also allowed us to identify 18 high-quality muscle-specific miRNAs.

Unfortunately, since our Nuc-Seq identifies nuclear pre-miRNAs, we can only identify

the miRNAs, but not which strand (5p or 3p or both) is loaded in the Argonaute protein.

Because of this limitation, one of the caveats of our miRNA prediction analysis is that

we process only the 5p strands through the miRanda algorithm, ignoring the 3p strands.

We also ignore miRNAs described in miRbase with less than 1,000 reads. Another

stringent filter we used excluded the miRNAs located within introns of other genes

(mirtron). The *C. elegans* WS250 release contains 257 miRNAs, including 118 mirtrons.

We decided to remove the mirtrons because the density of reads within their genomic

regions in genes with small introns made it too difficult to map them with high

confidence. This step automatically removed 46% of all *C. elegans* miRNAs from our

analysis and may explain why several body muscle miRNAs identified by others (Alberti

et al. 2018) are not present in our list.

In addition, we used very stringent filters during the miRNA identification step

using the miRanda algorithm, which may have also reduced the number of target

predictions.

We believe that all these stringent filters and rules, while significantly reducing

the number of miRNAs and their identified target genes, increase the quality of our hits.

We have validated the body muscle localization of several miRNAs identified in

our study (**Main Figure 4B**).

The body muscle miRNA interactome produced by clustering our 16 high-quality

miRNAs and the miRanda predictions using 3'UTRs data from our 2,849 protein-coding

genes identified in this study, showed seven subnetworks. One large subnetwork

contained eight miRNAs (**Main Figure 4)**. Most of the 88 genes in this large subnetwork

are connected to single miRNAs, but exceptions occur with *gld-1*, *daf-16* and *lin-14*

targeted by three miRNAs (*miR-241*, *let-7,* miR-392, and *miR-555*1), and *flh-1*, *ntl-9*,

rga-4, *mboa-2*, *alh-7*, *lgg-2*, and *cab-1*, each connected to two miRNAs. More

experiments need to be performed to validate these hits and understand how and why

these genes are specifically regulated in the body muscle tissue.

In conclusion, we have produced an updated *C. elegans* body muscle

transcriptome and miRNA Interactome, which will allow future studies to better

understand the function of this tissue in normal states and diseases.

MATERIALS AND METHODS

Preparation of the transgenic strains: The *C. elegans* strain bqSi189 II; mel-

28(bq5[GFP::mel-28]) III, or BN452, which ubiquitously expresses GFP and mCherry in

all nuclei, was obtained from the CGC, which is funded by NIH Office of Research

Infrastructure Programs (P40 OD010440). The genomic DNA was extracted from mixed

stage BN452 worms, and the mCherry-tagged *his-58* sequence was cloned using PCR

using BP Gateway-flanked primers specific for the Gateway entry vector pDONR221.

The *mCherry*::*his-58* sequence was successfully cloned using Gateway technology into

the entry vector, pDONR221, as evidenced by sequencing data. To assemble the

finalized clone, we performed a Gateway LR Clonase II plus reaction (cat. 12538-013;

Invitrogen) using the destination vector pCFJ150 (Frøkjær-Jensen *et al.* 2012) and entry

clones containing the body-muscle-specific promoter *myo-3*, the *mCherry*::*his-58*

sequence, and the *unc-54* 3′UTR as previously published (Blazie *et al.* 2017). The

pCFJ150 construct and pCFJ601 (50 ng/µl), which contains a Mos1 transposon, were

injected into the *C. elegans* strain EG6699 [ttTi5605 II; unc-119(ed3) III; oxEx1578]

(Frøkjær-Jensen et al. 2012), which is designed for MosI-mediated single-copy

integration (MosSCI), using standard injection techniques.

Nuclei extraction: Several medium plates of mixed stage worms were washed 4 times

at 1,500 rpm for 3 min and resuspended in 10 mL of ice-cold NPB buffer (10 mM

HEPES pH 7.6, 10 mM KCl, 1.5 mM MgCl2, 1 mM EGTA, 0.25 mM sucrose) per

250 µl volume of worm pellet. Nuclei were released by douncing 8-12 times in a chilled

Wheaton stainless-steel tissue grinder (clearance 0.0005 inches = 12.5 µm) in 5 mL

batches. The homogenate was sequentially filtered through 40 µm, 20 µm, and 10 µm

nylon filters (pluriStrainer), then centrifuged for 10 min at 2,500 *g* at 4 °C to pellet the

nuclei. The supernatant was aspirated, and the pellets containing nuclei were

resuspended in 4 volumes of ice-cold NPB buffer. To remove larger debris, the

resuspended nuclei pellets were centrifuged at 300 *g* for 1 min, then the supernatant

containing the nuclei was transferred to a new tube on ice.

Nuclei stability: The nuclei were isolated from the *myo-3::mCherry::his-58::unc-54*

strain as described above. Nuclei were imaged in triplicates using DAPI and mCherry

filters using a Leica DMi8 inverted microscope over eight hours. Images were obtained

using 1 s exposure times. The result of this experiment is shown in **Supplemental**

Figure S1.

FACS sorting: Two replicates of resuspended nuclei samples for the body muscle

tissue were FACS isolated into TRIzol solution using a BD Biosciences FACSAria III cell

sorter with a 70 µm nozzle, and a 560 nm laser with a temperature-controlled tube

holder at 4°C. The Eppendorf tubes were filled until a 1:1 TRIzol reagent, and sorted

nuclei prep was achieved after periodic gentle inversion. We set the gating parameters

with 10,000 events from our positive control BN452 and negative control N2, and

30,000 events for the *myo-3::mCherry::his-58::unc-54* strain. We sorted 239,000 and

353,000 nuclei for the sample and the replicate for the set1 and 340,000 and 353,000

sorted nuclei for the sample and the replicate of the set2.

RNA extraction and library preparation: The RNA from FACS isolated nuclei was

extracted using an RNA MiniPrep kit (ZR2070, Zymo Research) per the manufacturer's

protocol. RNA was quantified by using a bioanalyzer. The cDNA libraries for the two

replicates were generated using Nextera XT RNA sequencing. We prepared six mixed-

stages cDNA libraries from the following worm strains: BN452 (two replicates), *myo-*

3::mCherry::his-58::unc-54 (four replicated). The libraries were depleted of ribosomal

RNA contaminants using the Low Input RiboMinus Eukaryotic System v2

(ThermoFisher Scientific Catalog # A15027). The libraries were prepared from the

ribosomal RNA depleted samples using the Nextflex Rapid Directional RNAseq Kit

(PerkinElmer Catalog # NOVIA-5138-08)) as per the manufacturer's protocol and

sequenced on the Nextseq 500 instrument (Illumina, San Diego, CA), using 75bp SE

chemistry. The sequencing runs were performed at Girihlet Inc. Oakland (CA). We

obtained between 50M to 60M of mappable reads across all six datasets.

Bioinformatics analysis: The FASTQ files corresponding to the six datasets and the

controls (total of six datasets) were mapped to the *C. elegans* gene model WS250 using

the Bowtie 2 algorithm (Langmead and Salzberg 2012) with the following parameters: --

local -D 20 -R 3 -L 11 -N 1 -p 40 --gbar 1 -mp 3. The mapped reads were converted into

a bam format and sorted using SAMtools software using standard parameters (Li et al.

2009). We processed ~285M reads obtained from all our datasets combined and

obtained a median mapping success of ~93%. We used the Cufflinks software (Trapnell

et al. 2010) to estimate the expression levels of the genes obtained in each dataset as

per their BAM files. We calculated the fragment per kilobase per million bases (FPKM)

number in each experiment and performed the comparison using the Cuffdiff algorithm

(Trapnell et al. 2010). We have pooled the four experimental samples into two sets,

named set1 and set2, and the two replicates using the BN452 into a new set named

'BN452 nuclear'. We used the median FPKM value >=5 in the set1 and set2 dataset as

a threshold to define positive gene expression levels. The results are shown in

Supplemental Fig. S2 and Supplemental Table S1 using scores obtained by the

Cuffdiff algorithm (Trapnell et al. 2010) and plotted using the CummeRbund package.

miRNA target identification: The predicted miRNA targeting network was constructed

by extracting the longest 3'UTR sequence for each of the 2,848 protein-coding genes

identified in our study. We converted the 3'UTR sequences to FASTA format and

parsed the file using the miRanda algorithm (Enright et al. 2003) using a mature 5p *C.*

elegans miRNA list containing the 18 miRNAs detected with an FPKM value >=1

(**Supplemental Table S1**), and using stringent parameters (-strict -sc -1.2). We only

used miRNAs that have been previously detected in *C. elegans* with more than 1,000

reads in the miRbase database (www.mirbase.org) and that are not present in introns of

protein-coding genes. The miRanda algorithm produced high-quality 141 predicted

targets for 16 miRNAs. The networks were then built using the Cytoscape software

(Shannon et al. 2003) and uploaded to the Network Analyst online software (Xia et al.

2015) to produce the images shown in **Main Figure 4**.

miRNA validation: To validate the RNA sequencing data shown in **Main Figure 4B**,

three specific transgenic *C. elegans* strains were purchased from the CGC (VT1153,

VL347, and VT1189). These strains drive GFP expression under the control of *let-7*,

miR-230, or *miR-241* promoters respectively. Each strain was imaged using identical

settings for brightfield and GFP expression. These transgenic strains were imaged

using a Leica DMi8 Inverted microscope.

Western Blot experiments: The western blot validation experiments shown in **Main**

Figure 1C were performed using total protein for each fraction, and the input was

measured using a Bradford assay; 0.75 ng of protein was used for both the nuclear and

cytoplasmic fractions, and 3 ng of protein was used for the input. Primary monoclonal

antibodies for RFP (6G6-100, Chromotek) (1:1000) and GAPDH (ab125247, Abcam)

(1:2000) were used, followed by IRDye 800CW goat: mouse secondary antibodies (LI-

COR, 925–32210) (1:5000). The membrane was imaged using the ODYSSEY CLX

system (LI-COR Biosciences, NE).

Imaging analysis: Confocal images of isolated nuclei pre-FACS stained with DAPI for

the BN452 strain were acquired in the Biodesign Imaging Core, Division of the Arizona

State University Bioimaging Facility. Fluorescent microscopy images of isolated nuclei

pre-FACS stained with DAPI for the *myo-3*::*MH58*::*unc-54* strain were acquired using a

Leica DMi8 inverted microscope.

Network Analysis: The network shown in **Main Figure 3B** was constructed parsing the

top 2,000 hits identified in this study using the STRING algorithm (v. 11.5) (Szklarczyk

et al. 2021), run with standard parameters using only 'protein-protein interactions' as

input. The produced network possesses 1,995 nodes and 31,449 edges, with an

average node degree of 31.5 and an average local clustering coefficient of 0.356.

Promoter Analysis: We extracted 600 nt from the transcription start site for the

top 100 genes identified in this study. We then used different custom Perl scripts

to calculate the nucleotide distribution. The transcription factor predictions were

produced by parsing these promoters to the Simple Enrichment Analysis script

from the MEME suite software (Bailey et al. 2015). The results are shown in

Supplemental Figure S4 and **Supplemental Table S2**.

FIGURE LEGEND

Main Figure 1: Nuc-Seq schematics. A) Transgenic worms expressing

mCherry fluorochrome fused to histone H2B (*his-58*) are (1) homogenized, (2)

then subjected to several rounds of mechanical filtration. (3) The resultant nuclei

are then collected and (4) subjected to FACS sorting. At the completion of the

sorting step, (5) the mCherry-positive nuclei were incubated into TRIzol reagent

and the recovered RNA is sequenced. B) Top panel: The strain BN452

ubiquitously expresses the mCherry::*his-58* transgene. Bottom Panel: mCherry

positive nuclei recovered after Step 2 described in Panel A. (Red: mCherry, Blue:

DAPI). C) Western blot of nuclear and cytoplasmic fractions from BN452 worms

blotted with α-RFP and α-GAPDH antibodies. D) Left Panel: nuclear FACS profile

from N2 and BN452 nuclei. The red dots show the sorted mCherry-positive

population. Right Panel: Pie chart showing the percentage of BN452 sorted

mCherry-positive nuclei.

Main Figure 2: FACS sorting of body muscle nuclei: A: Top Panel:

Schematics of the construct used to isolate the body muscle transcriptome. The

body muscle-specific promoter *myo-3* drives the expression of the mCherry:::*his-*

58 cassette. Bottom Panel: Bright field and fluorescent images of the resultant

transgenic worm strain. B) nuclear FACS profile from N2 and

myo3p::mCherry:::his-58::unc-54 3'UTR nuclei. The red dots show the sorted

mCherry-positive population. Right Panel: Pie chart showing the percentage of

sorted mCherry-positive and body muscle-specific nuclei.

Main Figure 3: The *C. elegans* body muscle transcriptome. A) Left Panel:

Heat map showing the body muscle expression levels log(fpkm) of the 20,362

protein-coding genes in WS250. (a) BN452 positive control, (m) muscle

transcriptome. The muscle-specific transcriptome is highlighted with a dashed

yellow line. Right Panel: several examples of expression levels in genes

identified in the (a) BN452 and (m) muscle-specific nuclear fractions. The vertical

axes mark expression levels as the log(FPKM). The muscle-specific genes *unc-*

54, *mlc-1*, *unc-27,* and *lev-11* were all detected in our study, while the intestine-

specific *elt-2* and the GABA neuronal-specific *unc-47* were not present. B) The *C.*

elegans body muscle Interactome. We clustered all 2,849 protein-coding genes

identified in this study using previously published protein-protein interaction data.

Each gene is shown as a pink dot. Within this network, we identified six smaller

highly interconnected subnetworks shown in different colors. The genes in each

subnetwork were independently subjected to GO term enrichment analyses

shown with colored panels above each respective subnetwork.

Main Figure 4: The *C. elegans* body muscle Interactome. A) The 3'UTRs of

the 2,849 protein-coding genes identified in this study were screened for miRNA

targets in their 3'UTRs using 16 high-quality body-muscle miRNAs identified in

this study. We identified seven subnetworks. Each miRNA is shown in grayscale

and varies in size, depending on its number of predicted targets. Each target is

shown in a different color, depending on the number of predicted miRNA targets

in its 3'UTR, as from the miRanda algorithm. Red: one target, Light pink: two

targets, Orange: three targets. B) transgenic *C. elegans* strains expressing GFP

fluorochrome under the control of three miRNAs identified in this study. *Let-7*

(VT1153), *miR-230* (VL347), and *miR-241* (VT1189). All three strains show

strong body muscle expression in young adult worms.

Supplemental Materials:

Supplemental Figures S1-S4.

Supplemental Table S1: Comprehensive list of transcripts identified in this

study.

Supplemental Table S2: Simple Enrichment Analysis of sequence motifs

identified in promoters of transcripts identified in this study.

DATA AVAILABILITY

Raw reads were submitted to the NCBI Sequence Read Archive

(http://trace.ncbi.nlm.nih.gov/Traces/sra/) with BioProject ID: PRJNA820874 and

Submission ID: SUB11236444. The results of our analyses are available in Excel

format as **Supplemental Table S1**.

AUTHOR CONTRIBUTION

ALS and MM designed the experiments. ALS developed and performed the nuclear

FACS sorting experiments in collaboration with AFM and MYM, and ALS prepared the

sequencing reactions. MM performed the bioinformatic analysis. ALS and MM analyzed

the data, led the analysis and interpretation, assembled the figures, and wrote the

manuscript. All authors read and approved the final manuscript.

ACKNOWLEDGMENTS

This work is supported by NIH grant 1R01GM118796 to M.M. Some of the strains used

in this study were provided by the CGC, funded by the NIH Office of Research

Infrastructure Programs (P40 OD010440).

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

REFERENCES

Alberti C, Manzenreither RA, Sowemimo I, Burkard TR, Wang J, Mahofsky K, Ameres SL,
Cochella L. 2018. Cell-type specific sequencing of microRNAs from complex animal
tissues. *Nat Methods* **15**: 283-289.
Andachi Y, Kohara Y. 2016. A whole-mount in situ hybridization method for microRNA detection
in Caenorhabditis elegans. *RNA* **22**: 1099-1106.
Bailey TL, Johnson J, Grant CE, Noble WS. 2015. The MEME Suite. *Nucleic Acids Res* **43**: W39-49.
Bartel DP. 2018. Metazoan MicroRNAs. *Cell* **173**: 20-51.
Basu A, Behera S, Bhardwaj S, Dey S, Ghosh-Roy A. 2021. Regulation of UNC-40/DCC and UNC-
6/Netrin by DAF-16 promotes functional rewiring of the injured axon. *Development* **148**.
Bernstein E, Caudy AA, Hammond SM, Hannon GJ. 2001. Role for a bidentate ribonuclease in
the initiation step of RNA interference. *Nature* **409**: 363-366.
Blazie SM, Babb C, Wilky H, Rawls A, Park JG, Mangone M. 2015. Comparative RNA-Seq analysis
reveals pervasive tissue-specific alternative polyadenylation in Caenorhabditis elegans
intestine and muscles. *BMC Biol* **13**: 4.
Blazie SM, Geissel HC, Wilky H, Joshi R, Newbern J, Mangone M. 2017. Alternative
Polyadenylation Directs Tissue-Specific miRNA Targeting in Caenorhabditis elegans
Somatic Tissues. *Genetics* **206**: 757-774.
Brosnan CA, Palmer AJ, Zuryn S. 2021. Cell-type-specific profiling of loaded miRNAs from
Caenorhabditis elegans reveals spatial and temporal flexibility in Argonaute loading. *Nat*
Commun **12**: 2194.
Chandler RJ, Cogo S, Lewis PA, Kevei E. 2021. Modelling the functional genomics of Parkinson's
disease in Caenorhabditis elegans: LRRK2 and beyond. *Biosci Rep* **41**.
Chaya T, Patel S, Smith EM, Lam A, Miller EN, Clupper M, Kervin K, Tanis JE. 2021. A C. elegans
genome-wide RNAi screen for altered levamisole sensitivity identifies genes required for
muscle function. *G3 (Bethesda)* **11**.
Cogliati T, Good DJ, Haigney M, Delgado-Romero P, Eckhaus MA, Koch WJ, Kirsch IR. 2002.
Predisposition to arrhythmia and autonomic dysfunction in Nhlh1-deficient mice. *Mol*
Cell Biol **22**: 4977-4983.
Ellwood RA, Hewitt JE, Torregrossa R, Philp AM, Hardee JP, Hughes S, van de Klashorst D,
Gharahdaghi N, Anupom T, Slade L et al. 2021a. Mitochondrial hydrogen sulfide
supplementation improves health in the C. elegans Duchenne muscular dystrophy
model. *Proc Natl Acad Sci U S A* **118**.
Ellwood RA, Piasecki M, Szewczyk NJ. 2021b. Caenorhabditis elegans as a Model System for
Duchenne Muscular Dystrophy. *Int J Mol Sci* **22**.
Enright AJ, John B, Gaul U, Tuschl T, Sander C, Marks DS. 2003. MicroRNA targets in Drosophila.
Genome Biol **5**: R1.
Fox RM, Watson JD, Von Stetina SE, McDermott J, Brodigan TM, Fukushige T, Krause M, Miller
DM, 3rd. 2007. The embryonic muscle transcriptome of Caenorhabditis elegans.
Genome Biol **8**: R188.
Gieseler K, Qadota H, Benian GM. 2017. Development, structure, and maintenance of C.
elegans body wall muscle. *WormBook* **2017**: 1-59.

Gomez-Saldivar G, Fernandez A, Hirano Y, Mauro M, Lai A, Ayuso C, Haraguchi T, Hiraoka Y, Piano F, Askjaer P. 2016. Identification of Conserved MEL-28/ELYS Domains with Essential Roles in Nuclear Assembly and Chromosome Segregation. *PLoS Genet* **12**: e1006131.

Grishkevich V, Hashimshony T, Yanai I. 2011. Core promoter T-blocks correlate with gene expression levels in C. elegans. *Genome Res* **21**: 707-717.

Haenni S, Ji Z, Hoque M, Rust N, Sharpe H, Eberhard R, Browne C, Hengartner MO, Mellor J, Tian B et al. 2012. Analysis of C. elegans intestinal gene expression and polyadenylation by fluorescence-activated nuclei sorting and 3'-end-seq. *Nucleic Acids Res* **40**: 6304-6318.

Hrach HC, O'Brien S, Steber HS, Newbern J, Rawls A, Mangone M. 2020. Transcriptome changes during the initiation and progression of Duchenne muscular dystrophy in Caenorhabditis elegans. *Hum Mol Genet* **29**: 1607-1623.

Kawamata T, Tomari Y. 2010. Making RISC. *Trends Biochem Sci* **35**: 368-376.

Kotagama K, Schorr AL, Steber HS, Mangone M. 2019. ALG-1 Influences Accurate mRNA Splicing Patterns in the Caenorhabditis elegans Intestine and Body Muscle Tissues by Modulating Splicing Factor Activities. *Genetics* **212**: 931-951.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**: 357-359.

Lecroisey C, Segalat L, Gieseler K. 2007. The C. elegans dense body: anchoring and signaling structure of the muscle. *J Muscle Res Cell Motil* **28**: 79-87.

Lee RC, Feinbaum RL, Ambros V. 1993. The C. elegans heterochronic gene lin-4 encodes small RNAs with antisense complementarity to lin-14. *Cell* **75**: 843-854.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome Project Data Processing S. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078-2079.

Mallick A, Ranawade A, van den Berg W, Gupta BP. 2020. Axin-Mediated Regulation of Lifespan and Muscle Health in C. elegans Requires AMPK-FOXO Signaling. *iScience* **23**: 101843.

Markaki M, Tavernarakis N. 2020. Caenorhabditis elegans as a model system for human diseases. *Curr Opin Biotechnol* **63**: 118-125.

Martinez NJ, Ow MC, Reece-Hoyes JS, Barrasa MI, Ambros VR, Walhout AJ. 2008. Genome-scale spatiotemporal analysis of Caenorhabditis elegans microRNA promoter activity. *Genome Res* **18**: 2005-2015.

Miao W, Ma Z, Tang Z, Yu L, Liu S, Huang T, Wang P, Wu T, Song Z, Zhang H et al. 2021. Integrative ATAC-seq and RNA-seq Analysis of the Longissimus Muscle of Luchuan and Duroc Pigs. *Front Nutr* **8**: 742672.

Miska EA, Alvarez-Saavedra E, Abbott AL, Lau NC, Hellman AB, McGonagle SM, Bartel DP, Ambros VR, Horvitz HR. 2007. Most Caenorhabditis elegans microRNAs are individually not essential for development or viability. *PLoS Genet* **3**: e215.

Nabeshima Y, Hanaoka K, Hayasaka M, Esumi E, Li S, Nonaka I, Nabeshima Y. 1993. Myogenin gene disruption results in perinatal lethality because of severe muscle defect. *Nature* **364**: 532-535.

Natale C, Barzago MM, Diomede L. 2020. Caenorhabditis elegans Models to Investigate the Mechanisms Underlying Tau Toxicity in Tauopathies. *Brain Sci* **10**.

586   Prosdocimo DA, Sabeh MK, Jain MK. 2015. Kruppel-like factors in muscle health and disease.
587   *Trends Cardiovasc Med* **25**: 278-287.
588   Reinhart BJ, Slack FJ, Basson M, Pasquinelli AE, Bettinger JC, Rougvie AE, Horvitz HR, Ruvkun G.
589   2000. The 21-nucleotide let-7 RNA regulates developmental timing in Caenorhabditis
590   elegans. *Nature* **403**: 901-906.
591   Remmelzwaal S, Boxem M. 2019. Protein interactome mapping in Caenorhabditis elegans. *Curr*
592   *Opin Syst Biol* **13**: 1-9.
593   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T.
594   2003. Cytoscape: a software environment for integrated models of biomolecular
595   interaction networks. *Genome Res* **13**: 2498-2504.
596   Song BM, Avery L. 2013. The pharynx of the nematode C. elegans: A model system for the study
597   of motor control. *Worm* **2**: e21833.
598   Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, Doncheva NT, Legeay M, Fang T,
599   Bork P et al. 2021. The STRING database in 2021: customizable protein-protein
600   networks, and functional characterization of user-uploaded gene/measurement sets.
601   *Nucleic Acids Res* **49**: D605-D612.
602   Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ,
603   Pachter L. 2010. Transcript assembly and quantification by RNA-Seq reveals
604   unannotated transcripts and isoform switching during cell differentiation. *Nat*
605   *Biotechnol* **28**: 511-515.
606   Xia J, Gill EE, Hancock RE. 2015. NetworkAnalyst for statistical, visual and network-based meta-
607   analysis of gene expression data. *Nat Protoc* **10**: 823-844.
608   Yue Y, Li S, Shen P, Park Y. 2021. Caenorhabditis elegans as a model for obesity research. *Curr*
609   *Res Food Sci* **4**: 692-697.
610

A

2. Filter

4. FACS sort
into TRIzol

1. Homogenize

Waste

3. Pellet,
resuspend
nuclei

5. Extract RNA,
sequence

B

BN452

C

	Input	Cyto	Nuc

50 kDa

α-RFP

37 kDa

α-GAPDH

D

N2 Negative Control

GFP-A

mCherry+

mCherry PE-Texas Red-A

BN452 Positive Control

GFP-A

mCherry+

mCherry PE-Texas Red-A

BN452 mCherry+ Events

34.3%

65.7%

☐mCherry+ ☐Negative

A

B

A

B

Term ID	Term Name	E value
0003824	catalytic activity	3.794x10⁻¹⁶
0016071	small molecule metabolic process	6.546x10⁻¹⁴
0006091	generation of precursor metabolites and energy	6.482x10⁻¹³

Term ID	Term Name	E value
0022904	respiratory electron transport chain	6.242x10⁻⁴⁰
0042775	mitochondrial ATP synthesis	2.72x10⁻³⁹
0042773	ATP synthesis coupled electron transport	4.807x10⁻³⁹
0019646	aerobic electron transport chain	4.807x10⁻³⁹

Term ID	Term Name	E value
0006379	mRNA processing	5.104x10⁻²⁹
0016071	mRNA metabolic process	2.565x10⁻²⁸

Term ID	Term Name	E value
0051015	actin filament binding	8.32x10⁻¹⁵
003779	actin binding	2.743x10⁻¹³
0009653	anatomical structure morphogenesis	9.902x10⁻¹⁰
0030239	myofibril assembly	2.173x10⁻⁶
0055001	muscle cell development	3.684x10⁻⁶
0055002	striated muscle cell development	2.846x10⁻⁶

Term ID	Term Name	E value
0005793	endoplasmic reticulum	4.422x10⁻¹⁰
0030135	coated vesicle	1.219x10⁻⁹
0005794	Golgi apparatus	4.269x10⁻⁹

Term ID	Term Name	E value
1902600	proton transmembrane transport	2.084x10⁻⁶⁰
0098662	inorganic cation transmembrane transport	4.313x10⁻³⁹
0006754	ATP biosynthetic process	3.425x10⁻³²
0046034	ATP metabolic process	6.021x10⁻²⁶

Main Figure 4