An updated *C. elegans* nuclear body muscle transcriptome for studies in muscle

formation and function

Anna L. Schorr[1,2,3]

Email: aschorr@asu.edu

Alejandro Felix Mejia[3]

Email: Aflelixme@asu.edu

Martina Y. Miranda[4]

Email: MMiranda26@my.gcu.edu

Marco Mangone[1,2,3*]

* Corresponding author

Email: mangone@asu.edu

[1)] Molecular and Cellular Biology Graduate Program, Arizona State University, Tempe,

AZ; [2)] Virginia G. Piper Center for Personalized Diagnostics, The Biodesign Institute at

Arizona State University 1001 S McAllister Ave, Tempe, AZ; [3)] School of Life Sciences,

Arizona State University, 751 E Lemon Mall, Tempe, AZ 85287; [4)] Helios Scholars at the

Translational Genomics Research Institute, 445 N 5th St 4th Floor, Phoenix, AZ 85004

Running Title: Body muscle miRNAs in *C. elegans*

Keywords: miRNAs, 3'UTR, muscle tissue, interactome, *C. elegans*

ABSTRACT

The body muscle is an important tissue used in organisms for proper viability and locomotion. Although this tissue is generally well studied and characterized, and many pathways have been elucidated throughout the years, we still lack a comprehensive understanding of its transcriptome and how it controls muscle development and function. Here, we have updated a nuclear FACS sorting-based methodology to isolate and sequence a high-quality muscle transcriptome from *C. elegans* mixed-stage animals. We have identified 2,848 muscle-specific protein-coding genes, including 78 transcription factors and 206 protein-coding genes containing an RNA binding domain. We studied their interaction network, performed a detailed promoter analysis, and identified novel muscle-specific *cis*-acting elements. We have also identified 16 high-quality muscle-specific miRNAs, studied their function *in vivo* using fluorochrome-based analyses, and developed a high-quality *C. elegans* miRNA Interactome incorporating other muscle-specific datasets produced by our lab and others. Our study expands our understanding of how muscle tissue functions in *C. elegans* and in turn, provide results that can in the future be applied to humans to study muscular-related diseases.

INTRODUCTION

The body muscle is an important tissue used in organisms for proper viability and locomotion. Muscle tissue's structure, shape, and size change depending on its function and location. In humans, there are three main types of muscles: skeletal, smooth, and cardiac muscles, each responsible for a different form of contraction. Smooth and cardiac tissue contract involuntarily and are mainly located on the surface of organs. In contrast, skeletal muscles contract voluntarily.

The contractile unit of the muscle is the sarcomere. It is composed of at least thirty different proteins, of which the most abundant are myosin and actin, and is ultimately responsible for the contraction reaction leading to movement.

The model organism *C. elegans* is ideal for studying muscle development and morphogenesis due to its well-characterized transcriptome, fully mapped cell lineage, and the availability of several methods for genetic engineering.

The basic contractile unit of the muscle is the sarcomere, which is strongly conserved in its structure and function from *C. elegans* to mammals (WormBook). *C. elegans* sarcomeres are composed of myosin-containing thick filaments and actin-containing thin filaments, which are responsible for the contraction reaction leading to movement through the myosin heads pulling on the actin filaments (WormBook). A notable difference between the worm and vertebrates is the absence of the costamere and Z-disc, which in vertebrates anchors the actin filaments to the extracellular matrix. However, worms contain a structure called the dense body, which performs the same function [1]. In addition, over 200 proteins have been identified in *C. elegans* to be essential for the assembly, maintenance, and function of the sarcomere, many of which

have human orthologs [2]. *C. elegans* is also a popular choice for studying disorders

affecting humans, including muscular dystrophy [3-5], since many human disease

genes, when introduced in worms phenocopy the symptoms [6-9].

C. elegans possess two large muscles: the pharynx and the body muscle. The

pharynx is localized to the front of the animal and is responsible for food intake and the

physical crushing [10]. It is composed of eight layers of non-striated muscles,

surrounded by epithelial and neural tissue. The body muscle tissue comprises 95

striated muscle cells localized throughout the animal's body and is functionally

equivalent to the vertebrate skeletal muscles [11]. The contractile unit of the muscle is

the sarcomere. It is composed of at least 30 different proteins, of which the most

abundant are myosin and actin, and is ultimately responsible for the contraction reaction

leading to movement. The sarcomere is highly conserved between nematodes and

vertebrates and is composed of repeated units of a complex mesh of thin (I-band) and

thick (A-band) filaments. The thin filaments are mainly composed of actin and anchor

each sarcomere to the dense bodies and stabilize the entire structure. Conversely, the

thick filaments are mostly made of myosin and are attached to the M-line and mediate

tension generated during muscle contraction.

86   To properly develop and maintain functional body muscle tissue, gene

expression must be strictly regulated. Our group and others in the past finely

characterized the transcriptome of the *C. elegans* body muscle in the adult stage [12,

13] and during the embryonic development [14] and identified important regulatory

mechanisms used by muscle cells to maintain their homeostasis.

In addition to the identification of important genes in this tissue, our group and

others recently identified that tissue-specific alternative polyadenylation (APA), a poorly

characterized mechanism that produces genes with different 3'untranslated region

(3'UTR) isoforms, is an important regulatory mechanism used by cells to evade the

negative regulation effected by microRNAs (miRNAs) [12, 13]. Still, the complexity and

depth of this form of regulation are currently unknown. Regulation of gene expression

also depends on miRNAs, which are short ncRNAs that target regulatory elements in

3'UTRs and repress gene expression in the cytoplasm [15, 16].

Since their discovery, several groups, including our lab, have shown that miRNAs

influence gene regulation considerably [17]. The canonical miRNA biogenesis pathway

begins with the transcription of pri-miRNAs in the nucleus, which fold into a hairpin

structure [17]. The microprocessor complex cleaves the pri-miRNAs into a shorter stem-

loop structure, the pre-miRNAs, which are then exported to the cytoplasm [17]. Once in

the cytoplasm, the pre-miRNAs are cleaved into mature miRNAs, which associate with

an Argonaute protein as part of the RNA-induced silencing complex (RISC) [17-19].

When loaded into the RISC, miRNAs bind to complementary target sequences within

the 3'UTRs of genes to be repressed [17]. This implies that the 3'UTRs and their

isoforms are integral for miRNA-based regulation.

Several strategies have been developed to sequence tissue-specific

transcriptomes and miRNAs in *C. elegans*. These methodologies utilize either

fluorescence-activated cell sorting (FACS) [14, 20], immunoprecipitation [12, 13, 21], or

RNA modification techniques [22]. Each of these approaches has unique advantages

but also presents several caveats. Methodologies utilizing FACS allow for high purity

and stringency when isolating RNA from cells or nuclei. These approaches rely on

tissue-specific promoters to drive fluorochromes expression in cells or nuclei. After

isolation, the fluorescent cells or nuclei may be readily sorted using FACS, then either

sequenced [20] or processed using microarrays [14].

Immunoprecipitation and antibody-based approaches have also been widely

used to isolate the transcriptomes of specific tissues. This approach provides

considerably high yields of RNA compared to FACS-based methods and is generally

much more cost-efficient.

Brosnan *et al*. (2021) used an epitope labeled Argonaute protein expressed in

specific *C. elegans* tissues to immunoprecipitate tissue-specific miRNAs populations

from the body muscle, intestine, and neurons [21].

Alberti *et al*. (2018) developed a methodology named "microRNAome by

methylation-dependent sequencing (Mime-seq)" to identify tissue-specific miRNA

populations from complex animals at the level of single-cell specificity [22]. The Mime-

seq approach expresses a transgenic methyltransferase (Ath-HEN1) from *Arabidopsis*

thaliana in an animal system, allowing the labeling of small RNAs. Mime-seq has been

utilized in *C. elegans* and *Drosophila* and improved tissue-specific miRNA identification

compared to previous immunoprecipitation-based methods [22].

Our group also used an immunoprecipitation-based approach to identify

transcripts targeted by miRNAs within the *C. elegans* intestine, and body muscle [23]

and observed transcripts were targeted differentially between the two tissues. While

identifying the miRNA targets provided valuable information regarding tissue-specific

gene regulation, a major limitation of this study was that we could not identify the

specific miRNAs involved.

Here, we describe a novel nuclear FACS sorting-based approach named "Nuc-

Seq," to allow the isolation and sequencing of high-quality muscle transcriptomes and

miRNA populations from *C. elegans* mixed-stage animals. Using this method, we have

identified 2,848 muscle-specific protein-coding genes, studied their interaction network,

performed a detailed promoter analysis, and identified novel muscle-specific *cis*-acting

elements. Using fluorochrome-based analyses *in vivo*, we developed a high-quality

muscle miRNA interactome, which incorporates other muscle-specific datasets

produced by our lab and others.

RESULTS

Nuc-Seq: an updated approach to identify muscle-specific transcriptome

We wanted to improve the annotation of the body muscle transcriptome by

developing a nuclear FACS-based strategy to isolate and sequence transcriptomes

from *C. elegans* body muscle nuclei. We named this approach 'Nuc-Seq.' Our final goal

was to identify the body muscle transcriptome and its miRNA population expressed

explicitly in this tissue.

To test the feasibility and optimize this approach, we first performed experiments

using the *C. elegans* strain BN452, which ubiquitously expresses the mCherry

fluorochrome fused to the histone H2B ortholog gene *his-58* [24] (**Main Figure 1**). We

fractionated this worm strain using mechanical stress and separated two fractions: one

cytoplasmic and one enriched with mCherry fluorescent nuclei. (**Main Figure 1A-B and**

Materials and Methods). Using a Western blot approach, we successfully tested these

two fractions for the presence of known cytoplasmic and nuclear proteins (**Main Figure**

1C). Importantly, these nuclei are stable over time with minimal degradation after

several hours (**Supplemental Figure S1**). We then performed the FACS sorting step

and successfully isolated a large pool of mCherry-positive nuclei, with over 34% of all

the events being mCherry-positive (**Main Figure 1D**). We named this updated approach

Nuclear Sequencing (Nuc-Seq).

With these results in our hands, we decided to move forward and limit the

expression of the *mCherry*::*his-58* cassette in the body muscle tissue. We cloned the

his-58 gene*,* fused it to the mCherry fluorochrome, and then forced its expression only

in the body muscle tissue using the promoter region of the *C. elegans* ortholog of the

myosin heavy chain gene (*myo-3*) (**Main Figure 2A upper panel**). As expected, the

resultant transgenic worm restricts the expression of the mCherry fluorochrome in the

body muscle nuclei (**Main Figure 2A lower panel**), with fewer fluorescent nuclei

compared to the experiments in the BN452 strain (**Main Figure 2A**). We then performed

the FACS sorting step (**Main Figure 2B**). Although the number in sorted nuclei is

significantly less than those obtained with the positive control BN452, we could still

isolate a large population of mCherry-positive nuclei across from each of our replicates,

which we then extracted and sequenced RNA from.

The identification of the C. elegans body muscle transcriptome

We performed three sequencing reactions in duplicates (technical triplicates), two

from our body muscle-expressing nuclei (body muscle enriched), and one from nuclei

isolated from BN452 (negative controls) (a total of six samples). We obtained

approximately 55M mappable reads for each sample (including our technical replicates).

For most of the samples, we could map more than 97% of the total reads

(**Supplemental Figure S2A**). The results obtained with our biological replicates

correlate well (**Supplemental Figure S2B-C**).

We have identified 2,849 protein-coding genes in the *C. elegans* body muscle tissue,

corresponding to ~14% of all *C. elegans* protein-coding genes (20,362 protein-coding

genes; WS250) (**Main Figure 3 and Supplemental Table S1**). Using comparative data

from Kim et al. [25], we identified human orthologs for 1,872 worm genes (66%).

While some of our top hits have not been characterized yet, as expected, we

detected many known components of the sarcomere, such as many of the proteins that

form the thick filaments (myosin heavy chain genes, *myo-3*, *unc-54*) and myosin light

chain (*mlc-1*), and sarcomere organization and structural factors (*unc-15*, *unc-45*), as

well as those that form the thin filaments, including several actin genes (*act-1, act-3,*

act-4) and structural factors (*unc-60*, *unc-73*) (**Main Figure 3A and Supplemental**

Figure S3). We also detect orthologs of the tropomyosin gene *lev-11*, which in resting

sarcomere blocks myosin binding to actin, and other notable genes previously known to

be in the *C. elegans* body muscle tissue (**Main Figure 3A and Supplemental Figure**

S3).

Our dataset identified 78 transcription factors, including *mdl-1*, *skn-1*, *xbp-1*, *mxl-*

3, all previously known to be expressed in the body muscle tissue, and 207 non-

ribosomal proteins containing an RNA binding domain **Supplemental Table S1**. Within

this group, we identified K08D12.3, an ortholog of the human ZNF9, which is mutated in

myotonic dystrophy type 2, *mbl-1*, a MUSCLEBLIND-type of mRNA splicing regulator

required for muscle dense body organization, locomotion, and vulval morphogenesis,

and *etr-1*, an ortholog of the human CUG-binding protein CUGBP1, which is required for

embryonic muscle development and has been implicated in myotonic dystrophy

Supplemental Table S1.

Next, we studied the interaction network of the identified C. *elegans* muscle

genes. We used the STRING database [26] for this analysis, which contains known *C.*

elegans predicted and functional protein-protein interactions. Using our top 2,000

genes, we produced a large network containing 1,995 nodes and 31,449 edges (**Main**

Figure 3B). The genes are highly interconnected (**Main Figure 3B),** producing six large

distinguished sub-networks involved in body muscle formation, maintenance, protein

production, and energy storage (**Main Figure 3B)**. A GO term analysis shows functional

enrichments for actin filament binding, skeletal muscle myosin filament assembly,

mitochondrial functions, and ATP biosynthesis: all processes functioning in the body

muscle tissue (**Main Figure 3B)**. 71% of all genes identified in this study are protein-

coding genes, followed by ncRNAs, which are also very abundant (**Main Figure 4A;**

Top left). The largest functional category is receptors, followed by RNA-binding

proteins, cytoskeleton, and various enzymes (**Main Figure 4A; Top Right**). Based on

WormBase data, 31% of the genes identified in this study show unique muscle tissue

expression (**Main Figure 4B; Top Left**). Almost all genes are present throughout

226 development (**Main Figure 4B; Top Right**). Most genes show cytoplasmic localization,

227 and 36% of them are expressed in operons (**Main Figure 4B; Bottom Panels**).

229 ***The C. elegans body muscle PROMOTERome***

230  Next, we sought to study the core promoters of the genes identified in this study

231 to pinpoint potential motifs transactivated by muscle-specific transcription factors. We

232 have extracted 500 nucleotides upstream and 100 nucleotides downstream of the

233 transcription start site of the 2,849 genes we mapped in this tissue and plotted their

234 nucleotide distribution (**Supplemental Figure S4**). We detected an enrichment of

235 thymidine nucleotides just upstream of the transcription start site in these genes

236 (**Supplemental Figure S4 Panel A**). Repeated polythymine motifs (T-blocks) have

237 been previously reported in *C. elegans* core promoters, and their presence correlates

238 with gene expression levels [26]. We also performed a motif enrichment analysis in

239 these core promoters and identified several novel *cis*-acting elements (**Supplemental**

240 **Figure S4 Panel B** and **Supplemental Table S2**). We mapped 12 members of the

241 human Krueppel-like transcription factors family (KLF), which are important regulators of

242 gene expression in vertebrate development and are involved in muscle health and

243 disease [27], Myogenin (MYOG), a muscle-specific basic-helix-loop-helix (bHLH)

244 transcription factor involved in muscle development, myogenesis, and repair [28], and

245 NHLH1, a helix-loop-helix transcription factor that is involved in growth and

246 development of a wide variety of tissues and species [29] and has been linked to

247 muscle growth and development [30]. The complete list is shown in **Supplemental**

248 **Table S2**.

The C. elegans body muscle miRNAs

We then aimed to detect and study the body muscle-specific miRNA population

identified in our study. Using stringent parameters (**see Materials and Methods**), we

identified 16 miRNAs expressed in the *C. elegans* body muscle tissue (**Main Figure 5A**

and Supplemental Table S3). Several of these miRNAs have also been previously

described to be expressed in this tissue by other groups [31], and impairing their

function either leads to defects in the muscular fiber formation or shows an

uncoordinated phenotype [22, 32] (**Supplemental Table S1).** We validated the body

muscle localization of some of these miRNAs using two independent approaches: a

GFP-based approach using worm strain expressing this fluorochrome under the control

of specific miRNA promoters (**Main Figure 5B**) and a modified RT-qPCR approach

using muscle-specific miRNAs immunoprecipitated from a *C. elegans* strain expressing

a GFP::ALG-1 chimera in the body muscle tissue (**Supplemental Figure S5-S6**) [23,

33].

Both approaches provided strong evidence that our Nuc-Seq approach identified

bona fide muscle-specific miRNAs. Importantly, this RT qPCR approach also allowed us

to determine which one of the two miRNA strands (5p or 3p) was loaded in Argonaute.

We determined that the 5p strand was preferred for *let-7, miR-34, miR-228, miR-239a,*

miR-239b and *miR-5551*. The 3p strand was preferred for *miR-1, miR-77, miR-80, miR-*

230 and *miR-250*. *miR-392* did not have a distinct strand preference and was detected

at lower levels than other miRNAs. We also performed RT qPCR with primers for *miR-1,*

which was surprisingly below our detection threshold from our nuclear sequencing data.

(**Supplemental Figures S5-S6**). With this data, we constructed a body-muscle miRNA

Interactome. We performed MiRanda miRNA prediction using stringent parameters,

querying the 16 miRNAs with the preferred 5p or 3p strand identified in this study to the

body muscle-specific 3'UTRome data we published in the past [12, 13] network (**Main**

Figure 5A).

While most body-muscle genes are targeted by a single miRNA, 11 genes are

targeted by two miRNAs, two genes (*gld-1*, *daf-16*) are targeted by three miRNAs, and

one gene (*lin-14*) is targeted by four miRNAs (**Main Figure 5A**). Our most

interconnected miRNA is *miR-5551*, an uncharacterized *C. elegans*-specific miRNA

predicted to target 22 muscle-expressed protein-coding genes we detected in this tissue

(**Main Figure 5A**). A GO term analysis shows that its targets control contractile fibers,

sarcomere, and myofibril formation (**Supplemental Figure S3**). *Let-7* and the closely

related *miR-241* are also present. These miRNAs temporally regulate larval

development and have been previously identified in the pharyngeal and vulval muscles

[31, 34]. Among the predicted target genes, we identified several genes in the DAF

insulin pathway that control muscle health and aging [35, 36].

DISCUSSION

Here we have developed an updated FACS-based nuclear sorting method we

named Nuc-Seq, to identify tissue-specific transcriptomes from cell lysates and applied

it to determine the *C. elegans* muscle transcriptome. Nuc-Seq is robust, reproducible,

has minimal background noise, and allows the simultaneous identification of both

muscle-specific protein-coding genes and ncRNAs, such as miRNAs (**Main Figure 1**).

When applied to the *C. elegans* muscle tissue, Nuc-Seq allowed the precise

identification of 2,849 protein-coding genes, including 78 muscle transcription factors

and 207 non-ribosomal proteins containing an RNA-binding domain. Taken together,

the muscle transcriptome we identified covers ~14% of the entire *C. elegans* protein-

coding transcriptome (**Main Figure 3**). Some of these genes, such as *unc-54*, *mlc-1*,

unc-27, and others, have been previously identified in the muscle tissue and play an

important role in muscle contraction and muscle health in general. We have identified

almost all genes that form the sarcomere structure and are required for muscle

contraction (**Supplemental Figure S3**). When clustered using previously published

protein-protein interaction data [37], the *C. elegans* muscle transcriptome is highly

interconnected (**Main Figure 3B**), with several notable subnetworks corresponding to

mRNA processing, protein synthesis, ATP metabolism, and muscle cell development

(**Main Figure 3B**), all consistent with the function of this tissue.

Recently, similar methodologies have been utilized with success in fruit flies and mice.

For example, the Fly Cell Atlas published high-quality transcriptome data from 15

somatic tissues, including other tissues that cannot be directly dissected [38]. Similarly,

in mice, fluorescent nuclei have been purified from skeletal myofibers [39] and in

syncytial skeletal muscle cells throughout development [40]. Together with ours, these

studies highlight complex regulatory networks occurring in the muscle tissue.

We have also performed a promoter analysis, studying the promoter composition of the

genes identified in this study (**Supplemental Figure S4**). Our study identified the

presence of polythymine motif (T-blocks), previously reported by others, directly

upstream of the transcription start site, and using prediction software (Meme Suite [41]),

we have also identified many potential elements of muscle-specific transcription factors

used to control gene expression in this tissue.

Our approach also allowed us to identify 16 high-quality muscle-specific miRNAs.

Unfortunately, since Nuc-Seq identifies nuclear pre-miRNAs, we can only identify the

miRNAs, but not which strand (5p or 3p or both) is loaded in the Argonaute protein. To

bypass this limitation, we used a modified miRNA-CLIP approach coupled with RT

qPCR analysis to determine which miRNA strand, the 5p or the 3p, was loaded into the

Argonaute complex. We tested 11 miRNAs we identified from nuclear sequencing and

additionally tested *miR-1*, which was not identified from nuclear sequencing.

Interestingly, we found that *miR-1-3p* was highly expressed from our pulldown of ALG-1

in the body muscle. In contrast, some miRNAs that had high FPKM values, such as

miR-5551, had noticeably lower expression levels from our RT qPCR data

(**Supplemental Figure S6**). One critical difference between our two methods of miRNA

profiling was that the miRNAs were sequenced from the nucleus instead of the

cytoplasm. A possible explanation for the differences in expression is that certain

miRNAs may localize to the nucleus or not be exported and associate with Argonaute

proteins as mature miRNAs. Previous studies have also described nuclear-localized

miRNAs [42-44] and characterized their functions [45]. Future studies investigating the

miRNAs found in high abundance in our nuclear sorting but not from our cytoplasmic RT

qPCR could help explain these discrepancies and if these miRNAs have biological

functions.

To strengthen our miRNA Interactome results, we also ignore miRNAs described

in miRbase with less than 1,000 reads. Another stringent filter we used excluded the

miRNAs located within introns of other genes (mirtrons). The *C. elegans* WS250 release

contains 257 miRNAs, including 118 mirtrons. We removed the mirtrons because the

density of reads within their genomic regions in genes with small introns made it too

difficult to map them with high confidence. This step automatically removed 46% of all

C. elegans miRNAs from our analysis and may explain why several body muscle

miRNAs identified by others [22] are not present in our list. In addition, we used very

stringent filters during the miRNA identification step using the miRanda algorithm, which

may have also reduced the number of target predictions.

349   We believe that all these stringent filters and rules, while significantly reducing

350   the number of miRNAs and their identified target genes, increase the quality of our hits.

351   We have validated the body muscle localization of several miRNAs identified in

352   our study (**Main Figure 5B and Supplemental Figures S5-S6**).

353   The body muscle miRNA interactome produced by clustering our 16 high-quality

miRNAs and the miRanda predictions using 3'UTRs data from our 2,849 protein-coding

genes identified in this study showed seven subnetworks. One large subnetwork

contained 12 miRNAs (**Main Figure 5)**. 88 genes in this large subnetwork are

connected to single miRNAs, but exceptions occur with *gld-1*, and *daf-16,* each targeted

by three miRNAs (*let-7-5p*, *miR-392-5p* and *miR-4937-5p* for *gld-1,* and *let-7-5p, miR-*

241-5p and *miR-793-5p* for *daf-16*). Interestingly, the heterochronic gene *lin-14* is

targeted by four miRNAs in this tissue. More experiments must be performed to validate

these hits and understand how and why these genes are specifically regulated in the

body-muscle tissue. It this also important to note that our approach did not allow us to

identify the pharyngeal muscle transcriptome, which represents the second-largest

muscle tissue in *C. elegans* and could provide an important new angle to study the

complete muscle tissue transcriptome and miRNAome in this organism.

In conclusion, we have produced an updated *C. elegans* body muscle

transcriptome and miRNA Interactome, which will allow future studies to better

understand the function of this tissue in normal states and diseases.

MATERIALS AND METHODS

Preparation of the transgenic strains: The *C. elegans* strain bqSi189 II; mel-

28(bq5[GFP::mel-28]) III, or BN452, which ubiquitously expresses GFP and mCherry in

all nuclei, was obtained from the CGC, which is funded by NIH Office of Research

Infrastructure Programs (P40 OD010440). The genomic DNA was extracted from mixed

stage BN452 worms, and the mCherry-tagged *his-58* sequence was cloned using PCR

using BP Gateway-flanked primers specific for the Gateway entry vector pDONR221.

The *mCherry*::*his-58* sequence was successfully cloned using Gateway technology into

the entry vector, pDONR221, as evidenced by sequencing data. To assemble the

finalized clone, we performed a Gateway LR Clonase II plus reaction (cat. 12538-013;

Invitrogen) using the destination vector pCFJ150 (Frøkjær-Jensen *et al.* 2012) and entry

clones containing the body-muscle-specific promoter *myo-3*, the *mCherry*::*his-58*

sequence, and the *unc-54* 3′UTR as previously published (Blazie *et al.* 2017). The

pCFJ150 construct and pCFJ601 (50 ng/µl), which contains a Mos1 transposon, were

injected into the *C. elegans* strain EG6699 [ttTi5605 II; unc-119(ed3) III; oxEx1578]

(Frøkjær-Jensen et al. 2012), which is designed for MosI-mediated single-copy

integration (MosSCI), using standard injection techniques.

Nuclei extraction: To obtain a mixed-stage population, we synchronized a population

by bleaching gravid adult *C. elegans* as described previously [46]. We allowed worms to

reproduce several times over approximately two weeks, which resulted in a population

containing every developmental stage. Plates were inspected before nuclei isolation to

confirm all life stages were present in approximately equal ratios. Six plates

(100x15mm) were washed per replicate. Several medium plates of mixed stage worms

were washed four times at 1,500 rpm for 3 min and resuspended in 10 mL of ice-cold

NPB buffer (10 mM HEPES pH 7.6, 10 mM KCl, 1.5 mM MgCl2, 1 mM EGTA,

0.25 mM sucrose) per 250 µl volume of worm pellet. Nuclei were released by douncing

8-12 times in a chilled Wheaton stainless-steel tissue grinder (clearance 0.0005

inches = 12.5 µm) in 5 mL batches. The homogenate was sequentially filtered

through 40 µm, 20 µm, and 10 µm nylon filters (pluriStrainer), then centrifuged for 10

min at 2,500 *g* at 4 °C to pellet the nuclei. The supernatant was aspirated, and the

pellets containing nuclei were resuspended in 4 volumes of ice-cold NPB buffer. To

remove larger debris, the resuspended nuclei pellets were centrifuged at 300 *g* for 1

min, then the supernatant containing the nuclei was transferred to a new tube on ice.

Nuclei stability: The nuclei were isolated from the *myo-3::mCherry::his-58::unc-54*

strain as described above. Nuclei were imaged in triplicates using DAPI and mCherry

filters using a Leica DMi8 inverted microscope over eight hours. Images were obtained

using 1 s exposure times. The result of this experiment is shown in **Supplemental**

Figure S1.

FACS sorting*:* Two replicates of resuspended nuclei samples for the body muscle

tissue were FACS isolated into TRIzol solution using a BD Biosciences FACSAria III cell

sorter with a 70 μm nozzle, and a 560 nm laser with a temperature-controlled tube

holder at 4°C. The Eppendorf tubes were filled until a 1:1 TRIzol reagent and sorted

nuclei prep were achieved after periodic gentle inversion. We set the gating parameters

with 10,000 events from our positive control BN452 and negative control N2 and 30,000

events for the *myo-3::mCherry::his-58::unc-54* strain. We sorted 239,000 and 353,000

nuclei for the sample and the replicate for the set1 and 340,000 and 353,000 sorted

nuclei for the sample and the replicate of the set2.

RNA extraction and library preparation: The RNA from FACS isolated nuclei was

extracted using an RNA MiniPrep kit (ZR2070, Zymo Research) per the manufacturer's

protocol. RNA was quantified by using a bioanalyzer. The cDNA libraries for the two

replicates were generated using Nextera XT RNA sequencing. We prepared six mixed-

stages cDNA libraries from the following worm strains: BN452 (two replicates), *myo-*

3::mCherry::his-58::unc-54 (four replicates). The libraries were depleted of ribosomal

RNA contaminants using the Low Input RiboMinus Eukaryotic System v2

(ThermoFisher Scientific Catalog # A15027). The libraries were prepared from the

ribosomal RNA depleted samples using the Nextflex Rapid Directional RNAseq Kit

(PerkinElmer Catalog # NOVIA-5138-08) as per the manufacturer's protocol and

sequenced on the Nextseq 500 instrument (Illumina, San Diego, CA), using 75bp SE

chemistry. The sequencing runs were performed at Girihlet Inc. Oakland (CA). We

obtained between 50M to 60M of mappable reads across all six datasets.

Bioinformatics analysis: The FASTQ files corresponding to the six datasets and the

controls (total of six datasets) were mapped to the *C. elegans* gene model WS250 using

the Bowtie 2 algorithm [47] with the following parameters: --local -D 20 -R 3 -L 11 -N 1 -

p 40 --gbar 1 -mp 3. The mapped reads were converted into a bam format and sorted

using SAMtools software using standard parameters [48]. We processed ~285M reads

from all our datasets combined and obtained a median mapping success of ~93%. We

used the Cufflinks software [49] to estimate the expression levels of the genes obtained

in each dataset as per their BAM files. We calculated the fragment per kilobase per

million bases (FPKM) number in each experiment and performed the comparison using

the Cuffdiff algorithm [49]. We have pooled the four experimental samples into two sets,

named set1 and set2, and the two replicates using the BN452 into a new set named

'BN452 nuclear'. We used the median FPKM value >=5 in the set1 and set2 dataset as

a threshold to define positive gene expression levels. The results are shown in

Supplemental Fig. S2 and Supplemental Table S1 using scores obtained by the

Cuffdiff algorithm [49] and plotted using the CummeRbund package.

miRNA target identification: The predicted miRNA targeting network was constructed

by extracting the longest 3'UTR sequence for each of the 2,848 protein-coding genes

identified in our study from the body-muscle transcriptome dataset we identified in past

studies [12, 13]. We converted the 3'UTR sequences to FASTA format and parsed the

file using the miRanda algorithm [50] using stringent parameters (-strict -sc -1.2). We

used either the 5p or the 3p strand identified in our IP-coupled RT-qPCR approach

shown in **Supplemental Figures S5-S6**. All 16 miRNAs used were identified in this

study with an FPKM value >=1 (**Supplemental Table S1**). This value represents the top

5% of miRSVR scores produced by the miRanda algorithm. We also restrict the

analysis to pairing scores >150 and energy score < -7. We only used miRNAs that have

been previously detected in *C. elegans* with more than 1,000 reads in the miRbase

database (www.mirbase.org) and that are not present in introns of protein-coding

genes. The miRanda algorithm produced 118 high-quality predicted targets for 16

miRNAs. The networks were then built using the Cytoscape software [51] and uploaded

to the Network Analyst online software [52] to produce the images shown in **Main**

Figure 5.

miRNA validation*:* To validate the RNA sequencing data shown in **Main Figure 3B**,

three specific transgenic *C. elegans* strains were purchased from the CGC (VT1153,

VL347, and VT1189). These strains drive GFP expression under the control of *let-7*,

miR-230, or *miR-241* promoters, respectively. Each strain was imaged using identical

settings for brightfield and GFP expression. These transgenic strains were imaged

using a Leica DMi8 Inverted microscope. The miRNA validation in Supplemental Figure

S5-S6 was performed using miRNA immunoprecipitated from a *C. elegans* strain

expressing ALG-1 fused to the GFP fluorochrome expressed specifically in the body

muscle tissue [23]. The immunoprecipitation step was performed using anti-GFP

antibodies (Chromotek GTMA-10) as previously described [23]. Briefly, six mixed-stage

100x15mm confluent plates of transgenic *C. elegans* strains expressing the *pmyo-*

3::ALG-1::GFP construct were washed 3x at 1500 rpm for 3 min. Crosslinking and RNA

extraction was performed as previously described [23]. RNA was quantified, then 465ng

of immunoprecipitated RNA was polyuridylated (New England Biolabs, cat no. M0337S)

and reverse transcribed (Invitrogen; cat: no. 8080044) using a stem-loop poly-A reverse

primer for the first strand reaction as previously described [33]. 11 miRNA-specific

forward primers were designed and shown in Supplemental Table S4. Primers with low

melting temperatures were not used. Polyuridynated cDNA was diluted to a 1:2 or 1:10

ratio depending on the miRNAs' abundance. We used a 1:10 cDNA dilution for *let-7,*

miR-34, miR-77, miR-80, *miR-228* and *miR-1* and a 1:2 cDNA dilution for *miR-230, miR-*

239a, miR-239b, *miR-250, miR-392* and *miR-5551*. The relative abundance of each

miRNA was calculated based on an exogenous positive control.

Western Blot experiments*:* The western blot validation experiments shown in **Main**

Figure 1C were performed using total protein for each fraction, and the input was

measured using a Bradford assay; 0.75 ng of protein was used for both the nuclear and

cytoplasmic fractions, and 3 ng of protein was used for the input. Primary monoclonal

antibodies for RFP (6G6-100, Chromotek) (1:1000) and GAPDH (ab125247, Abcam)

(1:2000) were used, followed by IRDye 800CW goat: mouse secondary antibodies (LI-

COR, 925–32210) (1:5000). The membrane was imaged using the ODYSSEY CLX

system (LI-COR Biosciences, NE).

Imaging analysis: Confocal images of isolated nuclei pre-FACS stained with DAPI for

the BN452 strain were acquired in the Biodesign Imaging Core, Division of the Arizona

State University Bioimaging Facility. Fluorescent microscopy images of isolated nuclei

pre-FACS stained with DAPI for the *myo-3*::*MH58*::*unc-54* strain were acquired using a

Leica DMi8 inverted microscope.

Network Analysis: The network shown in **Main Figure 3B** was constructed by parsing

the top 2,000 hits identified in this study using the STRING algorithm (v. 11.5) [53], run

with standard parameters using only 'protein-protein interactions' as input and set the

minimum required interaction score to 'high-confidence' (0.700). The produced network

possesses 1,995 nodes and 31,449 edges, with an average node degree of 31.5 and an

average local clustering coefficient of 0.356. The network was then uploaded in

Cytoscape [54] to prepare the figure.

Promoter Analysis: We extracted 600 nt from the transcription start site for the

top 100 genes identified in this study. We then used different custom Perl scripts

to calculate the nucleotide distribution. The transcription factor predictions were

produced by parsing these promoters to the Simple Enrichment Analysis script

from the MEME suite software [41]. The results are shown in **Supplemental**

Figure S4 and **Supplemental Table S2**.

Expression Analysis

The expression analysis shown in Main Figure 4 was performed using the

SimpleMine tool in WormBase Field [46], which retrieves essential gene

information in this database. We used as input the 2,848 protein-coding genes

identified in this study. The output was then parsed with simple Perl scripts to

extract and assign gene function (GO Term output), previously published tissue

localization data, larval stage presence, cellular localization, and presence in

operons.

FIGURE LEGEND

Main Figure 1: Nuc-Seq schematics. A) Transgenic worms expressing

mCherry fluorochrome fused to histone H2B (*his-58*) are (1) homogenized and

(2) then subjected to several rounds of mechanical filtration. (3) The resultant

nuclei are then collected and (4) subjected to FACS sorting. After the sorting

step, (5) the mCherry-positive nuclei were sorted directly into TRIzol, and the

recovered RNA was sequenced. B) Top panel: The strain BN452 ubiquitously

expresses the mCherry::*his-58* transgene. Bottom Panel: mCherry positive nuclei

recovered after Step 2 described in Panel A. (Red: mCherry, Blue: DAPI). C)

Western blot of nuclear and cytoplasmic fractions from BN452 worms blotted with

α-RFP and α-GAPDH antibodies. D) Left Panel: nuclear FACS profile from N2

and BN452 nuclei. The BN452 sample is a positive control that represents all

nuclei. The red dots show the sorted mCherry-positive population. Right Panel:

Pie chart showing the percentage of BN452 sorted mCherry-positive nuclei.

Main Figure 2: FACS sorting of body muscle nuclei: A: Top Panel:

Schematics of the construct used to isolate the body muscle transcriptome. The

body muscle-specific promoter *myo-3* drives the expression of the mCherry:::*his-*

58 cassette. Bottom Panel: Bright field and fluorescent images of the resultant

transgenic worm strain. B) nuclear FACS profile from N2 and

myo3p::mCherry:::his-58::unc-54 3'UTR nuclei. The red dots show the sorted

mCherry-positive population. Right Panel: Pie chart showing the percentage of

sorted mCherry-positive and body muscle-specific nuclei.

Main Figure 3: The *C. elegans* body muscle transcriptome. A) Left Panel:

Heat map showing the body muscle expression levels log(fpkm) of the 20,362

protein-coding genes in WS250. (a) BN452 positive control, (m) muscle

transcriptome. The body muscle-specific transcriptome is highlighted with a

dashed yellow line. Right Panel: several examples of expression levels in genes

identified in the (a) BN452 (ubiquitously expresses GFP and mCherry in all

nuclei; positive control) and (m) body muscle-specific nuclear fractions. The

vertical axes mark expression levels as the log(FPKM). The body muscle-specific

genes *unc-54*, *mlc-1*, *unc-27,* and *lev-11* were all detected in our study, while the

intestine-specific *elt-2* and the GABA neuron-specific *unc-47* were not present.

(a: median log(fpkm) of the 'BN452 all nuclei' dataset; m: median fpkm values of

set1 and set2 muscle datasets). B) The *C. elegans* body muscle Interactome. We

clustered all 2,849 protein-coding genes identified in this study using previously

published protein-protein interaction data. Each gene is shown as a pink dot.

Within this network, we identified six smaller, highly interconnected subnetworks

shown in different colors. The genes in each subnetwork were independently

subjected to GO term enrichment analyses shown with colored panels above

each respective subnetwork.

Main Figure 4: Comparative Data Analysis. A) Left: Pie chart showing the

breakout of the RNA types identified in this study. Right: Out of 2,848 protein-

coding genes, 1,044 genes have been previously assigned to specific biological

functions and shown as a pie-chart. B) Top Left: Pie-chart showing the tissue

localization of the genes identified in this study. Top Right: gene expression

profile in different *C. elegans* developmental stages. Bottom Right: cellular

localization. Bottom left: the number of genes identified in this study that are in

operons. The comparative data used to prepare this analysis was extracted from

WormBase.

Main Figure 5: The *C. elegans* body muscle Interactome. A) The 3'UTRs of

the 2,849 protein-coding genes identified in this study were screened for miRNA

targets in their 3'UTRs using 16 high-quality body-muscle miRNAs identified in

this study. We identified four subnetworks. Each miRNA is shown in grayscale

and varies in size, depending on its number of predicted targets. Each target is

shown in a different color, depending on the number of predicted miRNA targets

in its 3'UTR, as from the miRanda algorithm. Red: one target, yellow: two targets,

green: three or more targets. B) transgenic *C. elegans* strains expressing GFP

fluorochrome under the control of three miRNAs identified in this study. *Let-7*

(VT1153), *miR-230* (VL347), and *miR-241* (VT1189). All three strains show

strong body muscle expression in young adult worms.

SUPPLEMENTAL MATERIALS:

Supplemental Figures S1-S6.

Supplemental Table S1: Comprehensive list of transcripts identified in this

study.

Supplemental Table S2: Simple Enrichment Analysis of sequence motifs

identified in promoters of transcripts identified in this study.

Supplemental Table S3: miRNA target prediction of the 16 miRNAs identified in

this study using the miRanda algorithm.

Supplemental Table S4: Primer list used in this study.

ETHICAL APPROVAL

Not applicable.

COMPETING INTERESTS

The authors declare that they have no competing interests.

AUTHORS' CONTRIBUTIONS

ALS and MM designed the experiments. ALS developed and performed the

nuclear FACS sorting experiments in collaboration with AFM and MYM, and ALS

prepared the sequencing reactions. MM performed the bioinformatic analysis.

ALS and MM analyzed the data, led the analysis and interpretation, assembled

the figures, and wrote the manuscript. All authors read and approved the final

manuscript.

FUNDING

604 This work is supported by NIH grant 1R01GM118796.

AVAILABILITY OF DATA AND MATERIALS

607 Raw reads were submitted to the NCBI Sequence Read Archive

608 (http://trace.ncbi.nlm.nih.gov/Traces/sra/) with BioProject ID: PRJNA820874 and

609 Submission ID: SUB11236444. The results of our analyses are available in Excel

610 format as **Supplemental Table S1**.

ACKNOWLEDGMENTS

613 Some of the strains used in this study were provided by the CGC, funded by the NIH

614 Office of Research Infrastructure Programs (P40 OD010440). We thank Emma Murari

615   and Dalton Meadows for their suggestions and advice during the preparation of the

616   manuscript. In memory of Amelia.

REFERENCES

1.   Lecroisey C, Segalat L, Gieseler K: **The C. elegans dense body: anchoring and signaling**
structure of the muscle. *J Muscle Res Cell Motil* 2007, **28**(1):79-87.
2.   Benian GM, Epstein HF: **Caenorhabditis elegans muscle: a genetic and molecular model**
for protein interactions in the heart. *Circ Res* 2011, **109**(9):1082-1095.
3.   Hrach HC, O'Brien S, Steber HS, Newbern J, Rawls A, Mangone M: **Transcriptome**
changes during the initiation and progression of Duchenne muscular dystrophy in
Caenorhabditis elegans. *Hum Mol Genet* 2020, **29**(10):1607-1623.
4.   Chaya T, Patel S, Smith EM, Lam A, Miller EN, Clupper M, Kervin K, Tanis JE: **A C. elegans**
genome-wide RNAi screen for altered levamisole sensitivity identifies genes required
for muscle function. *G3 (Bethesda)* 2021, **11**(4).
5.   Ellwood RA, Hewitt JE, Torregrossa R, Philp AM, Hardee JP, Hughes S, van de Klashorst
D, Gharahdaghi N, Anupom T, Slade L *et al*: **Mitochondrial hydrogen sulfide**
supplementation improves health in the C. elegans Duchenne muscular dystrophy
model. *Proc Natl Acad Sci U S A* 2021, **118**(9).
6.   Yue Y, Li S, Shen P, Park Y: **Caenorhabditis elegans as a model for obesity research.** *Curr*
Res Food Sci 2021, **4**:692-697.
7.   Chandler RJ, Cogo S, Lewis PA, Kevei E: **Modelling the functional genomics of**
Parkinson's disease in Caenorhabditis elegans: LRRK2 and beyond. *Biosci Rep* 2021,
41(9).
8.   Natale C, Barzago MM, Diomede L: **Caenorhabditis elegans Models to Investigate the**
Mechanisms Underlying Tau Toxicity in Tauopathies. *Brain Sci* 2020, **10**(11).
9.   Markaki M, Tavernarakis N: **Caenorhabditis elegans as a model system for human**
diseases. *Curr Opin Biotechnol* 2020, **63**:118-125.
10.   Song BM, Avery L: **The pharynx of the nematode C. elegans: A model system for the**
study of motor control. *Worm* 2013, **2**(1):e21833.
11.   Gieseler K, Qadota H, Benian GM: **Development, structure, and maintenance of C.**
elegans body wall muscle. *WormBook* 2017, **2017**:1-59.
12.   Blazie SM, Babb C, Wilky H, Rawls A, Park JG, Mangone M: **Comparative RNA-Seq**
analysis reveals pervasive tissue-specific alternative polyadenylation in
Caenorhabditis elegans intestine and muscles. *BMC Biol* 2015, **13**:4.
13.   Blazie SM, Geissel HC, Wilky H, Joshi R, Newbern J, Mangone M: **Alternative**
Polyadenylation Directs Tissue-Specific miRNA Targeting in Caenorhabditis elegans
Somatic Tissues. *Genetics* 2017, **206**(2):757-774.
14.   Fox RM, Watson JD, Von Stetina SE, McDermott J, Brodigan TM, Fukushige T, Krause M,
Miller DM, 3rd: **The embryonic muscle transcriptome of Caenorhabditis elegans.**
Genome Biol 2007, **8**(9):R188.

15.   Lee RC, Feinbaum RL, Ambros V: **The C. elegans heterochronic gene lin-4 encodes small**
RNAs with antisense complementarity to lin-14. *Cell* 1993, **75**(5):843-854.

16.   Reinhart BJ, Slack FJ, Basson M, Pasquinelli AE, Bettinger JC, Rougvie AE, Horvitz HR,
Ruvkun G: **The 21-nucleotide let-7 RNA regulates developmental timing in**
Caenorhabditis elegans. *Nature* 2000, **403**(6772):901-906.

17.   Bartel DP: **Metazoan MicroRNAs**. *Cell* 2018, **173**(1):20-51.

18.   Bernstein E, Caudy AA, Hammond SM, Hannon GJ: **Role for a bidentate ribonuclease in**
the initiation step of RNA interference. *Nature* 2001, **409**(6818):363-366.

19.   Kawamata T, Tomari Y: **Making RISC**. *Trends Biochem Sci* 2010, **35**(7):368-376.

20.   Haenni S, Ji Z, Hoque M, Rust N, Sharpe H, Eberhard R, Browne C, Hengartner MO,
Mellor J, Tian B *et al*: **Analysis of C. elegans intestinal gene expression and**
polyadenylation by fluorescence-activated nuclei sorting and 3'-end-seq. *Nucleic Acids*
Res 2012, **40**(13):6304-6318.

21.   Brosnan CA, Palmer AJ, Zuryn S: **Cell-type-specific profiling of loaded miRNAs from**
Caenorhabditis elegans reveals spatial and temporal flexibility in Argonaute loading.
Nat Commun 2021, **12**(1):2194.

22.   Alberti C, Manzenreither RA, Sowemimo I, Burkard TR, Wang J, Mahofsky K, Ameres SL,
Cochella L: **Cell-type specific sequencing of microRNAs from complex animal tissues**.
Nat Methods 2018, **15**(4):283-289.

23.   Kotagama K, Schorr AL, Steber HS, Mangone M: **ALG-1 Influences Accurate mRNA**
Splicing Patterns in the Caenorhabditis elegans Intestine and Body Muscle Tissues by
Modulating Splicing Factor Activities. *Genetics* 2019, **212**(3):931-951.

24.   Gomez-Saldivar G, Fernandez A, Hirano Y, Mauro M, Lai A, Ayuso C, Haraguchi T, Hiraoka
Y, Piano F, Askjaer P: **Identification of Conserved MEL-28/ELYS Domains with Essential**
Roles in Nuclear Assembly and Chromosome Segregation. *PLoS Genet* 2016,
12(6):e1006131.

25.   Kim W, Underwood RS, Greenwald I, Shaye DD: **OrthoList 2: A New Comparative**
Genomic Analysis of Human and Caenorhabditis elegans Genes. *Genetics* 2018,
210(2):445-461.

26.   Grishkevich V, Hashimshony T, Yanai I: **Core promoter T-blocks correlate with gene**
expression levels in C. elegans. *Genome Res* 2011, **21**(5):707-717.

27.   Prosdocimo DA, Sabeh MK, Jain MK: **Kruppel-like factors in muscle health and disease**.
Trends Cardiovasc Med 2015, **25**(4):278-287.

28.   Nabeshima Y, Hanaoka K, Hayasaka M, Esumi E, Li S, Nonaka I, Nabeshima Y: **Myogenin**
gene disruption results in perinatal lethality because of severe muscle defect. *Nature*
1993, **364**(6437):532-535.

29.   Cogliati T, Good DJ, Haigney M, Delgado-Romero P, Eckhaus MA, Koch WJ, Kirsch IR:
Predisposition to arrhythmia and autonomic dysfunction in Nhlh1-deficient mice. *Mol*
Cell Biol 2002, **22**(14):4977-4983.

30.   Miao W, Ma Z, Tang Z, Yu L, Liu S, Huang T, Wang P, Wu T, Song Z, Zhang H *et al*:
Integrative ATAC-seq and RNA-seq Analysis of the Longissimus Muscle of Luchuan and
Duroc Pigs. *Front Nutr* 2021, **8**:742672.

31.   Martinez NJ, Ow MC, Reece-Hoyes JS, Barrasa MI, Ambros VR, Walhout AJ: **Genome-**
698   **scale spatiotemporal analysis of Caenorhabditis elegans microRNA promoter activity**.
699   *Genome Res* 2008, **18**(12):2005-2015.
700   32.   Miska EA, Alvarez-Saavedra E, Abbott AL, Lau NC, Hellman AB, McGonagle SM, Bartel
701   DP, Ambros VR, Horvitz HR: **Most Caenorhabditis elegans microRNAs are individually**
702   **not essential for development or viability**. *PLoS Genet* 2007, **3**(12):e215.
703   33.   Mei Q, Li X, Meng Y, Wu Z, Guo M, Zhao Y, Fu X, Han W: **A facile and specific assay for**
704   **quantifying microRNA by an optimized RT-qPCR approach**. *PLoS One* 2012,
705   **7**(10):e46890.
706   34.   Andachi Y, Kohara Y: **A whole-mount in situ hybridization method for microRNA**
707   **detection in Caenorhabditis elegans**. *RNA* 2016, **22**(7):1099-1106.
708   35.   Mallick A, Ranawade A, van den Berg W, Gupta BP: **Axin-Mediated Regulation of**
709   **Lifespan and Muscle Health in C. elegans Requires AMPK-FOXO Signaling**. *iScience*
710   2020, **23**(12):101843.
711   36.   Basu A, Behera S, Bhardwaj S, Dey S, Ghosh-Roy A: **Regulation of UNC-40/DCC and UNC-**
712   **6/Netrin by DAF-16 promotes functional rewiring of the injured axon**. *Development*
713   2021, **148**(11).
714   37.   Remmelzwaal S, Boxem M: **Protein interactome mapping in Caenorhabditis elegans**.
715   *Curr Opin Syst Biol* 2019, **13**:1-9.
716   38.   Li H, Janssens J, De Waegeneer M, Kolluru SS, Davie K, Gardeux V, Saelens W, David FPA,
717   Brbic M, Spanier K *et al*: **Fly Cell Atlas: A single-nucleus transcriptomic atlas of the adult**
718   **fruit fly**. *Science* 2022, **375**(6584):eabk2432.
719   39.   Petrany MJ, Swoboda CO, Sun C, Chetal K, Chen X, Weirauch MT, Salomonis N, Millay
720   DP: **Single-nucleus RNA-seq identifies transcriptional heterogeneity in multinucleated**
721   **skeletal myofibers**. *Nat Commun* 2020, **11**(1):6374.
722   40.   Kim M, Franke V, Brandt B, Lowenstein ED, Schowel V, Spuler S, Akalin A, Birchmeier C:
723   **Single-nucleus transcriptomics reveals functional compartmentalization in syncytial**
724   **skeletal muscle cells**. *Nat Commun* 2020, **11**(1):6375.
725   41.   Bailey TL, Johnson J, Grant CE, Noble WS: **The MEME Suite**. *Nucleic Acids Res* 2015,
726   **43**(W1):W39-49.
727   42.   Liao JY, Ma LM, Guo YH, Zhang YC, Zhou H, Shao P, Chen YQ, Qu LH: **Deep sequencing of**
728   **human nuclear and cytoplasmic small RNAs reveals an unexpectedly complex**
729   **subcellular distribution of miRNAs and tRNA 3' trailers**. *PLoS One* 2010, **5**(5):e10563.
730   43.   Reyes-Gutierrez P, Ritland Politz JC, Pederson T: **A mRNA and cognate microRNAs**
731   **localize in the nucleolus**. *Nucleus* 2014, **5**(6):636-642.
732   44.   Politz JC, Hogan EM, Pederson T: **MicroRNAs with a nucleolar location**. *RNA* 2009,
733   **15**(9):1705-1715.
734   45.   Liu H, Lei C, He Q, Pan Z, Xiao D, Tao Y: **Nuclear functions of mammalian MicroRNAs in**
735   **gene regulation, immunity and cancer**. *Mol Cancer* 2018, **17**(1):64.
736   46.   Porta-de-la-Riva M, Fontrodona L, Villanueva A, Ceron J: **Basic Caenorhabditis elegans**
737   **methods: synchronization and observation**. *J Vis Exp* 2012(64):e4019.
738   47.   Langmead B, Salzberg SL: **Fast gapped-read alignment with Bowtie 2**. *Nat Methods*
739   2012, **9**(4):357-359.

48. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
R, Genome Project Data Processing S: **The Sequence Alignment/Map format and**
SAMtools. *Bioinformatics* 2009, **25**(16):2078-2079.
49. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold
BJ, Pachter L: **Transcript assembly and quantification by RNA-Seq reveals unannotated**
transcripts and isoform switching during cell differentiation. *Nat Biotechnol* 2010,
28(5):511-515.
50. Enright AJ, John B, Gaul U, Tuschl T, Sander C, Marks DS: **MicroRNA targets in**
Drosophila. *Genome Biol* 2003, **5**(1):R1.
51. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B,
Ideker T: **Cytoscape: a software environment for integrated models of biomolecular**
interaction networks. *Genome Res* 2003, **13**(11):2498-2504.
52. Xia J, Gill EE, Hancock RE: **NetworkAnalyst for statistical, visual and network-based**
meta-analysis of gene expression data. *Nat Protoc* 2015, **10**(6):823-844.
53. Szklarczyk D, Gable AL, Nastou KC, Lyon D, Kirsch R, Pyysalo S, Doncheva NT, Legeay M,
Fang T, Bork P *et al*: **The STRING database in 2021: customizable protein-protein**
networks, and functional characterization of user-uploaded gene/measurement sets.
Nucleic Acids Res 2021, **49**(D1):D605-D612.
54. Shen B, Zhang Y, Yu S, Yuan Y, Zhong Y, Lu J, Feng J: **MicroRNA-339, an epigenetic**
modulating target is involved in human gastric carcinogenesis through targeting
NOVA1. *FEBS Lett* 2015, **589**(20 Pt B):3205-3211.
761

A

2. Filter

4. FACS sort into TRIzol

1. Homogenize

3. Pellet, resuspend nuclei

Waste

5. Extract RNA, sequence

B

BN452

C

Input   Cyto   Nuc

50 kDa

α-RFP

37 kDa

α-GAPDH

D

N2 Negative Control

BN452 Positive Control

BN452 mCherry+ Events

GFP-A

mCherry+

mCherry PE-Texas Red-A

GFP-A

mCherry+

mCherry PE-Texas Red-A

34.3%

65.7%

☐mCherry+ ☐Negative

A

myo-3p

| mCherry | *his-58* | *unc-54 3'UTR* |

B

N2

myo-3p::mCherry::his-58

myo3::mCherry::his-58
mCherry+ Events

5.3%

94.7%

☐ mCherry+   ☐ Negative

A

B

Term ID	Term Name	E value
0003824	catalytic activity	3.794x10^-16
0016071	small molecule metabolic process	6.546x10^-14
0006091	generation of precursor metabolites and energy	6.482x10^-13

Term ID	Term Name	E value
0006379	mRNA processing	5.104x10^-29
0016071	mRNA metabolic process	2.565x10^-28

Term ID	Term Name	E value
0022904	respiratory electron transport chain	6.242x10^-40
0042775	mitochondrial ATP synthesis	2.72x10^-39
0042773	ATP synthesis coupled electron transport	4.807x10^-39
0019646	aerobic electron transport chain	4.807x10^-39

Term ID	Term Name	E value
0051015	actin filament binding	8.32x10^-15
003779	actin binding	2.743x10^-13
0009653	anatomical structure morphogenesis	9.902x10^-10
0030239	myofibril assembly	2.173x10^-6
0055001	muscle cell development	3.684x10^-6
0055002	striated muscle cell development	2.846x10^-6

Term ID	Term Name	E value
0005793	endoplasmic reticulum	4.422x10^-10
0030135	coated vesicle	1.219x10^-9
0005794	Golgi apparatus	4.269x10^-9

Term ID	Term Name	E value
1902600	proton transmembrane transport	2.084x10^-60
0098662	inorganic cation transmembrane transport	4.313x10^-39
0006754	ATP biosynthetic process	3.425x10^-32
0046034	ATP metabolic process	6.021x10^-26

A

RNA types

function

n= 3,994

n= 1,044

B

tissue

n= 2,848

developmental stage

n= 1,397

cellular localization

n= 1,165

operons

n= 2,848

Main Figure 5