Ventral attention network connectivity is linked to cortical maturation

and cognitive ability in childhood

Hao-Ming Dong[1]*, Xi-Han Zhang[1], Loïc Labache[1], Shaoshi Zhang[2], Leon Qi Rong

Ooi[2], B.T. Thomas Yeo[2,3,4], Daniel S. Margulies[5], Avram J. Holmes[6]*#, Xi-Nian

Zuo[7,8,9]*#

[1]Department of Psychology, Yale University, New Haven, Connecticut, USA

[2]Centre for Sleep and Cognition & Centre for Translational Magnetic Resonance

Research, Yong Loo Lin School of Medicine, Singapore, National University of

Singapore, Singapore, Singapore

[3]Department of Electrical and Computer Engineering, National University of

Singapore, Singapore, Singapore

[4]N.1 Institute for Health & Institute for Digital Medicine, National University of

Singapore, Singapore, Singapore

[5]Centre National de la Recherche Scientifique, Frontlab, Institut du Cerveau et de la

Moelle Epinière, Paris, France

[6]Department of Psychiatry, Brain Health Institute, Rutgers University, Piscataway,

USA

[7]State Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal

University, Beijing 100875, China

[8]National Basic Science Data Center, Beijing 100190, China

[9]Developmental Population Neuroscience Research Center, IDG/McGovern Institute

for Brain Research, Beijing Normal University, Beijing 100875, China

Correspondence: * To whom correspondence may be addressed. Emails:

donghaomingnd@gmail.com, avram.holmes@rutgers.edu, xinian.zuo@bnu.edu.cn.

# These authors have contributed equally to this work

Keywords: ventral attention network, connectivity gradient, cortical organization,

development, adolescence

31   **Abstract**

The human brain experiences functional changes through childhood and

adolescence, shifting from an organizational framework anchored within

sensorimotor and visual regions into one that is balanced through interactions

with later-maturing aspects of association cortex. Here, we link this profile of

functional reorganization to the development of ventral attention network

connectivity across independent datasets. We demonstrate that maturational

changes in cortical organization preferentially link to within-network

connectivity and heightened degree centrality in the ventral attention network,

while connectivity within network-linked vertices predicts cognitive ability. This

connectivity is closely associated with maturational refinement of cortical

organization. Children with low ventral attention network connectivity exhibit

adolescent-like topographical profiles, suggesting that attentional systems

may be relevant in understanding how brain functions are refined across

development. These data suggest a role for attention networks in supporting

age-dependent shifts in cortical organization and cognition across childhood

and adolescence.

 Introduction

The human brain undergoes a series of staged developmental cascades across

childhood and adolescence progressing from unimodal somatosensory/motor and

visual regions through the transmodal association cortex territories that support

complex cognitive functions[1-3]. Evidence for the scheduled timing of these

neurodevelopmental events has emerged across biological scales, from regional

profiles of cellular maturation[4], synapse formation and dendritic pruning[5], and

intracortical myelination[6] through macro-scale morphological features including

folding patterns[7] and associated areal expansion[8]. These processes are imbedded

within age dependent anatomical changes across lifespan[9], particularly the

prolonged development of association cortex territories. In parallel, substantial

progress has been made characterizing the organization[10,11] and spatiotemporal

maturation of large-scale functional systems across the cortex[12,13]. Here, *in vivo*

imaging work strongly supports the development of a hierarchical axis, or gradient, of

cortical organization, with association territories anchored at the opposite end of a

broad functional spectrum from primary sensory and motor regions[14]. Despite clear

evidence for age-dependent shifts in the macroscale organization of the cortex from

childhood through adolescence, the manner and extent to which specific functional

networks may contribute to the widespread process of cortical maturation remains to

be determined.

The focused study of discrete functional circuits has provided foundational

insights in core maturational processes. For example, discoveries have linked

hierarchical changes within amygdala- and ventral striatal-medial prefrontal cortex

(mPFC) circuitry to the development of emotional and social functioning in

adolescence[1,3,15,16]. Yet, the maturational refinement of these subcortical-cortical

circuits does not occur in isolation. Rather, they are embedded within a broad

restructuring of functional systems across the cortical sheet[17]. Here, areal and

network boundaries become more clearly defined throughout development[18], while

the predominance of local connectivity patterns in childhood gradually gives way to

long distance, integrative, connections in adolescence[13,19,20]. This reflects a

developmental transition from an anatomically constrained organizational motif to a

topographically distributed system[21]. In children, this complex functional architecture

is situated within the unimodal cortex, between somatosensory/motor and visual

regions. Conversely, adolescents transition into an adult-like gradient, anchored at

one end by unimodal regions supporting primary sensory/motor functions and at the

other end by the association cortex[14]. While the organizational profiles of large-scale

cortical networks are distinct across childhood and adolescence[22], the extent to

which developmental changes within select functional couplings may contribute to

the drastic reorganization in the brain hierarchy is an open question. By one view, the

developmental transition from unimodal through association cortices reflects the

coordinated and shared influence of maturational changes across multiple functional

systems spanning the entire connectome. An alternative, although not mutually

exclusive, possibility is that specific brain networks may play a preferential role in the

widespread developmental refinement of cortical connectivity.

Individual cortical parcels are functionally organized along a global gradient that

transitions from somato/motor and visual regions at one end and multimodal

association cortex at the other[23]. The hierarchical nature of these functional

relationships reflects a core feature of brain organization in both adolescents[14] and

adults[23]. Incoming sensory information undergoes a process of extensive elaboration

and attentional modulation as it cascades into deeper layers of cortical processing.

Visual system connectivity, as one example, moves along the dorsal and ventral

visual streams, uniting within aspects of the dorsal and ventral attention networks

including the anterior insula, superior parietal cortex, and operculum parietal before

eventually filtering through multimodal convergence zones, particularly within the

default network[24]. Although speculative, these data suggest a possible preferential

role for sensory orienting and attentional systems in the integrity of the information

processing hierarchies in the human brain. Intriguingly, there is mounting evidence to

suggest the staged development of a ventral attention network, encompassing

aspects of anterior insula, anterior prefrontal cortex and anterior cingulate cortex[11,25]

(see also, cingulo-opercular network[26] and salience network[27]), that follows the age-

dependent shifts in cortical organization across childhood and adolescence[14]. The

salience/ventral attentional network, together with frontoparietal network, have been

proposed to constitute a dual-network system for the 'top-down' and 'bottom-up'

processing necessary for adaptive behavioral responses[26,28,29], supporting the

functional propagation of information across primary somato/motor, visual, auditory

cortex through the default network[24]. These dissociable attentional and control

systems are interconnected in children but later segregate over the course of

adolescence to eventually form the parallel architectures that support adaptive

behavior in adulthood[18]. These data suggest that the attention system may play a

preferential role in the transformative brain changes occurring throughout childhood

and adolescence. Recent work has also revealed that the lateralization of functional

gradients may coincide with attention system lateralization[30]. As such, characterizing

the relationships linking attention network connectivity and age-dependent changes

in the macroscale brain organization would provide a tremendous opportunity to

understand how the functional architecture of cortex is shaped and sculpted across

the human lifespan. In turn, this would provide the opportunity to examine how the

hierarchical reorganization of the cortical sheet may contribute to the emergence of

cognitive and emotional abilities that mark the transition from childhood to

adolescence.

In the present study we examined the extent to which specific functional

networks may serve to underpin the age-dependent maturation of functional gradient

patterns across the cortical sheet. To directly address this open question, we first

established the cortical territories exhibiting pronounced functional changes in a

longitudinal sample of children and adolescents, revealing the preferential presence

of developmental shifts within the ventral attention network. Follow-up analyses

excluding regions exhibiting the maximal developmental change in children resulted

in the emergence of adolescent-like gradient patterns, suggesting ventral attention

territories may play a core role in the expression of adolescent-like connectivity

gradients. Moreover, across independent datasets, children with low ventral attention

connectivity exhibited a profile of cortical organization that closely resembles prior

reports in adolescents and adults. Highlighting the importance of attention network

connectivity in cognitive functioning, standardized measures of intelligence linked

with reduced attention network degree centrality in children and adolescents.

Collectively, these data suggest that ventral attention system functioning in childhood

and adolescence may underpin the developmental reorganization and maturation of

functional networks across the cortical sheet.

Results

Ventral attention network territories demonstrate high degree centrality and

pronounced shifts across development

Vertex-level functional connectivity (FC) matrices (20,484×20,484) were first

generated using the data provided by the Chinese Color Nest Project (CCNP)[21,32]. In

line with prior work[10,11,14], the top 10% connections of each vertex were retained to

enforce sparsity. Degree centrality maps for both children (6 to 12 years of age;

n=202) and adolescents (12 to 18 years of age; n=176) were generated to

characterize the broad organizational properties of the functional connectome across

development (Figure 1A,B). Here, degree centrality reflects the count of above

threshold connections for a given vertex to all other vertices (see Methods).

The observed profiles of degree centrality were highly similar between children

and adolescents (Pearson r=0.947, p_{spin}≤0.001). Significance was established using

permuted spin tests, which preserve the spatial autocorrelation structure of the

data[33]. Heightened degree centrality values in both children and adolescents were

preferentially evident in aspects of the ventral attention network, including portions of

anterior insula, medial prefrontal cortex, and anterior cingulate cortex/midline

supplementary motor area (Figure 1). Increased degree centrality was also present in

adolescents within default network territories including portions of posterior inferior

parietal lobule, posterior cingulate cortex, and precuneus. Additionally, visual system

areas including superior and transverse occipital sulcus at the boundary between

dorsal and visual network demonstrated high degree centrality values. Conversely,

primary somatosensory and motor areas as well as regions within the lateral

prefrontal cortex and temporal lobe exhibited relatively low degree centrality. Broadly,

these data reflect the presence of dense connectivity within medial and posterior

territories along the cortical sheet, while relatively low centrality was evident in lateral

prefrontal and somato/motor areas, highlighting a stable pattern of degree centrality

across childhood and adolescence.

Degree centrality broadly summarizes profiles of cortical connectivity, to examine

developmental changes in functional connectivity strength at the vertex level, we

calculated the associated Euclidean distance in functional connectivity similarity

between children and adolescents (Figure 1C). Despite the presence of broadly

consistent population-level patterns of degree centrality, analyses revealed spatially

non-uniform shifts in hub regions of functional connectivity across groups. The

maximum developmental changes were anchored within the ventral attention network

(Figure 1C-E), encompassing aspects of anterior and posterior insula as well as

cingulate cortex. One-way ANOVA revealed the presence of between-network

differences (F=790.94, df=6, p≤0.001), with increased Euclidean distance in the

ventral attention network, relative to other networks (see multiple comparisons results

in Supplemental Table 2). Prior work indicates that maturational age broadly follows

the theorized hierarchy of cortical information processing[23], with somato/motor and

visual networks maturing in childhood, while medial prefrontal aspects of default and

limbic networks, peak later during adolescence[14,34]. However, in the present

analyses, the default network exhibited relatively less developmental change in

Euclidean distance between groups, followed by visual network (Figure 1E). Although

speculative, these data suggest the presence of specific network-level similarities in

connectivity between children and adolescents that may precede broader age-

dependent shifts in the macroscale organization of cortex, highlighting the need to

consider the manner, in which individual functional networks (e.g., default and

attention) are embedded within the broader functional architecture of the brain.

A core role for the ventral attention network in the macroscale organization of cortex across childhood and adolescence

The transition from childhood to adolescence is marked by pronounced changes in the functional organization of cortex[14,22]. Broadly, this is reflected in the presence of age-dependent transitions across macroscale gradients that extend from unimodal (somato/motor and visual) regions through the cortical association areas that support complex cognition[2,3,15,16] (Figure 2A). Of note, such a transition is observed not only when restricting analyses to retain the top 10% connections of each vertex to enforce sparsity, but also revealed by when varying sets connections are included in the gradient analyses. Here, the primary transmodal gradient emerges in a higher percentile of excluded connections at each vertex (90%) in adolescents than in children (85%; see (Extended Data Fig.1). Next, we examined whether age-dependent alterations in ventral attention network connectivity might partly account for the maturation of the cortical processing hierarchy as reflected in these overlapping organizing axes, or gradients. Brain areas with maximum differences in Euclidean distance were extracted (See Methods; Figure 2B), and then removed from brain connectivity matrix while we rederived the functional gradients. Here, diffusion map embedding[10,35,36] was used to decompose participant-level connectivity matrices into a lower dimensional space. The resulting functional components, or gradients, reflect dissociable spatial patterns of cortical connectivity, ordered by the variance explained in the initial functional connectivity matrix[10,14].

As identified in our prior study[14], the primary gradient in children closely matches the second gradient in adolescents and adults. Here, dropping ventral attention areas (a simulation of lesion) generates adolescent- and adult-like gradient architectures in children. The simulated removal of ventral attention network regions led to the formation of a primary gradient in children that closely assembles the first gradient in both adolescents (r=0.68, p_{spin}≤0.001) and adults (r=0.84, p_{spin}≤0.001). The rederived second gradient in children most closely assembled the second gradient in both

adolescents (r=0.66, $p_{spin} \leq 0.001$) and adults (r=0.85, $p_{spin} \leq 0.001$). However, while the

primary gradient derived from ventral attention network simulated lesioned data in

children broadly recapitulated the primary gradient in adults[10], several

inconsistencies were observed. Notably, in the simulated lesioned data from the child

group, one end of the primary gradient of connectivity was anchored in the visual

areas, with the regions at the other end encompassed broad swaths of the

association cortex. Prior work in adults has revealed visual territories along with

somato/motor and auditory cortex serve to anchor one end of the primary cortical

gradient[10]. Additionally, although the second gradient derived from ventral attention

network simulated lesioned data in children closely resembles the second gradient in

adults, a muted default network profile can still be observed. While the dropped

clusters are not anchored at the extreme end of the primary gradient in children

(Figure 2A), it is densely connected and spatially adjacent to somato/motor

territories. Further control analysis revealed that areas within ventral attention

network primarily contributed to the reversal in gradients. Although speculative,

dropping of ventral attention vertices from the gradient analyses may decrease the

number of functional connections attributed to somato/motor network, indirectly

shifting its position along the gradient spectrum.

Most vertexes from the dropped clusters were from the ventral attention

(45.96%) and somato/motor networks (36.57%). Accordingly, the observed transition

to an adolescent- and adult-like functional architecture in children may reflect an

artifact, resulting from lesioning the data in a manner that removes of aspects of the

unimodal territories that anchor the primary gradient in children (Figure 2A). To

address this possibility and identify the primary drivers of adolescent-like profiles of

brain function in children, we subdivided the dropped clusters into 2 categories:

clusters falling within the borders of the somato/motor network and clusters outside

the somato/motor network. Gradients were then rederived in the child group to

explore the consequences associated with the individual removal of each cluster

component. The removal of somato/motor cluster preserves the developmentally

typical gradient architecture, in children the first gradient closely resembles the

second gradient in adolescents (r=0.94, $p_{spin}\leqslant$0.001) and adults (r=0.91, $p_{spin}\leqslant$

0.001), while the second gradient closely assemble the first gradient in adolescents

(r=0.95, $p_{spin}\leqslant$0.001) and adults (r=0.91, $p_{spin}\leqslant$0.001). Conversely, and consistent

with the analyses reported above (Figure 2B), dropping the cluster outside

somato/motor network led to the generation of an adolescent- and adult-like gradient

architecture in children. Here, the first gradient in children matched the first gradient

in adolescents (r=0.78, $p_{spin}\leqslant$0.001), while the second gradient in children resembled

the second gradient in adolescents (r=0.78, $p_{spin}\leqslant$0.001). Namely, dropping the

clusters outside somato/motor network in children results in a gradient organization

that resembles prior reports in adolescents.

We next examined the extent to which the observed elimination of age-

dependent shifts in the macroscale organization of the cortex is specific to the

removal of ventral attention network-dominated vertices. Here, we generated 500 null

models in the child group with clusters dropped at random locations across the

cortical sheet, but with shapes and sizes that match the ventral attention network

vertices (reflecting the maximum differences in Euclidean Distance between children

and adolescents). For each random model, the first and second gradients of child

group were extracted and correlated with the corresponding first and second

gradients in adults. Providing evidence for the role of the ventral attention network in

the formation of adult-like gradient architectures in children, for the primary gradient

the observed correlation was greater than the correlations from the null distribution

(p≤0.002) across 499/500 permutations (Figure 2C). For the second gradient the

observed correlation was greater than the correlations from the null distribution

(p≤0.001) across all 500 permutations. While the present analyses are consistent

with a core role for the ventral attention network in age-dependent changes in the

macroscale organization of the cortex, longitudinal future work should examine the

role of person level factors and possible relationships linking large-scale gradient

transitions with shifts in attention network functioning across development.

Cortical development is linked to a pattern of heightened within-ventral

attention network connectivity

Extending upon the prior Euclidean Distance analyses, connectome-level

changes in functional connectivity between children and adolescents are displayed in

a chord diagram (Figure 3) and grouped into networks according to Yeo's 7-network

solution[11]. These data reveal broad increases in within network connectivity for the

ventral attention, somato/motor, and default networks as well as a general flattening

of cross-network connectivity across development. The within-network connections

for the ventral attention, between-network connections for the dorsal attention, and

visual networks were increased, while other between-network connectivity with

somato/motor, limbic, frontoparietal and default networks are decreased from

childhood to adolescence. Individual developmental changes were examined by

linear mixed effect (LME) model. Decreased between-network connectivity for the

somato/motor network was observed ($p \leq 0.05$), while heightened between-network

connectivity was evident for the visual ($p \leq 0.05$), dorsal attention ($p \leq 0.001$), limbic

($p \leq 0.05$), and frontoparietal control ($p \leq 0.01$) networks. The degree centrality of

ventral attention network increased with age in children ($p \leq 0.05$), stabilizing in

adolescence where no other age-related associations were revealed in within-

network connections for ventral attention network.

When considering the other large-scale association networks, the present

analyses suggest a pattern where default network connectivity with other functional

systems is pruned during development, indicating a bias towards within network

connectivity and an associated differentiation from other network processes. A

developmental profile that may coincide with the emergence of the default network at

the apex of the network hierarchy in adolescence. Conversely, the frontoparietal

310 control network exhibited increased connectivity with dorsal and ventral attention

311 networks in adolescents, suggesting a potential association between cognitive

312 processes and attentional resources allocation during development.

Attention network connectivity links with cognitive ability and reveals the presence of adult-like gradient architectures in childhood

The analyses above provide evidence for a relationship between ventral attention network connectivity and the formation of adult-like gradient architectures in children. Significant connectome-level changes were also observed in heightened within-network functional connectivity and degree centrality of ventral attention network (Figure 3). However, it is not yet clear the extent to which individual variability in attention network functioning may link with the adult-like gradient architectures across development. To distinguish it from the typical developmental pattern, we refer to these early-emerging adult-like gradient architectures as an "accelerated maturation pattern". To examine this potential relationship, we divided participants into subgroups based on their individual ventral attention network connection counts (Figure 4A). Here, to avoid potential bias introduced by confounding factors like scan parameters, population and preprocessing steps, the group split was determined based on the median connection count in the CCNP oldest participants (>17 years of age; dotted line in Figure 4A) instead of independent adult population samples. Participants with fewer connections than this median value were assigned to a low ventral attention group (child n=95 adolescent n=76), all other participants were assigned to a high ventral attention group (child n=107; adolescent n=100). The gradients were then rederived for high and low attention groups across both children and adolescents. No significant associations were observed between the degree centrality of ventral attention network and demographic factors including in age (p=0.45), gender (p=0.82) and head motion (p=0.87). Further comparisons reveal matched demographic features between child the high and low ventral attention groups in age (p=0.0572), gender (p=0.7716) and head motion (p=0.5346).

In adolescents, both the primary (r=0.98, p_{spin}≤0.001) and secondary (r=0.99, p_{spin}≤0.001) gradient architectures were consistent within the low and high ventral attention network participants (Figure 4B,C). For children with high salience and

ventral attention connectivity, their functional organization follows the typical pattern

of brain maturation[14] (Figure 2A). Here the primary gradient in the high ventral

attention children matched the secondary gradient in both adolescents (r=0.89,

p_{spin}≤0.001) and adults (r=0.91, p_{spin}≤0.001), while the second gradient in high ventral

attention children matched the primary gradient in both adolescents (r=0.89,

p_{spin}≤0.001) and adults (r=0.91607, p_{spin}≤0.001). In these children, somato/motor

areas were anchored at the opposite extreme from visual regions, revealing a

unimodal dominant gradient architecture. However, in children with low ventral

attention connectivity we observed a developmentally accelerated pattern of gradient

organization that broadly matches the primary and secondary gradients previously

identified in both adolescents (Gradient 1: r=0.93, p_{spin}≤0.001; Gradient 2: r=0.92,

p_{spin}≤0.001) and adults (Gradient 1: r=0.68, p_{spin}≤0.001; Gradient 2: r=0.65,

p_{spin}≤0.001).

Permutation analyses were conducted to examine the significance of the

dissociable gradient architectures across the groups. Here, 95 and 107 child

participants were randomly assigned to two groups corresponding to the number of

participants in low and high ventral attention child group for 500 permutations, In the

adolescent group, 76 and 100 participants were randomly assigned to two groups

corresponding to the number of participants in low and high ventral attention

adolescent group for 500 times, then the gradient maps were recomputed and the

associated variances were extracted to generate a set of null models.

Children with low ventral attention connectivity exhibit a gradient organization

that broadly matches the primary and secondary gradients previously identified in

both adolescents and adults. Conversely, children with high ventral attention

connectivity exhibit functional organization that follows the typical pattern of brain

maturation. The permutation analyses demonstrate that the amount of variance

accounted for by the association cortex anchored gradient is increased in the low

(Gradient 1 variance: 0.3648) relative to high ventral attention connectivity

participants (Gradient 2 variance: 0.1052; p=0, 0/500 permutations revealed greater

difference than the real situation). Consistent with this profile the amount of variance

accounted for by the unimodal anchored gradient is decreased in the low (Gradient 2

variance: 0.1161) relative to high ventral attention connectivity participants (Gradient

1 variance: 0.3846; p=0, 0/500 revealed greater difference than the real situation).

Conversely, in the adolescence, we did not observe a significant difference

between the high and low ventral attention groups in the first association cortex

anchored gradient (0.3807 vs. 0.3595, p=0.0620), while the second gradient in high

ventral attention adolescent group (0.1052) accounted for a significantly lower

amount of variance (p=0.008, 4 in 500 permutations revealed lower difference than

the real situation) than low ventral attention adolescent group (0.1161).

The accelerated longitudinal design of the Chinese Color Nest Project[32,37], which

includes longitudinal tracking data with a visiting interval of 1.25 years, provided for

additional analyses in participants who were scanned both before (child group) and

after their 12th birthday (adolescent group). Here, we identified a set of child

participants (n=22) from the low ventral attention group who were also subsequently

scanned in their adolescence (n=26, mean age of initial scan=10.98±0.64; mean age

of second scan=12.90±0.69). In the participants classified as low ventral attention in

childhood, when they transition to adolescence, their first gradient in childhood is

highly correlated (r=0.9429, p<0.01) with the first gradient that in their adolescence. A

consistent group profile that was also evident when considering their second

gradients in both childhood and adolescence (r=0.9353, p<0.01; see Figure 4D).

Conversely, high ventral attention children (n=21), who were also scanned in

their adolescence (n=24, mean age of initial scan=11.09±0.63; mean age of second

scan=12.94±0.71), displayed a normative developmental trajectory. Here, their first

gradient in childhood was highly correlated with their second gradient in adolescence

(absolute r=0.9793, p<0.01), while their second gradient in childhood were highly

correlated with the first gradient in their adolescence (absolute r=0.9748, p<0.01; see

Figure 4D).

If the salience/ventral attention network contributes to the maturation of cortical

hierarchy, the functional integrity of attentional systems may also associate with

changes in cognitive and behavioral performance. To examine this hypothesis, we

next assessed the relationship between standardized measures of cognitive

functioning (IQ) and the degree centrality of the ventral attention network with an

LME model. LME models for IQ subdomain scores in verbal, perceptual reasoning,

working memory, processing speed, and a composite total score were constructed

separately. Age, gender, head motion and the vertex-level connectivity counts with

ventral attention network were fed into each LME model as fixed effects, repeated

measurements were set as the random effect. Results revealed significant negative

associations between ventral attention network connectivity and verbal ($p \leq 0.05$),

perceptual reasoning ($p \leq 0.01$) and the composite IQ scores ($p \leq 0.05$). Associations

with participant age and verbal score ($p \leq 8 \times 10^{-6}$), perceptual reasoning score

($p \leq 0.05$) and total score ($p \leq 0.001$), as well as associations with gender and verbal

score ($p \leq 0.05$), perceptual reasoning score ($p \leq 0.05$), and total score ($p \leq 0.05$) were

also observed. Suggesting that this pattern was not due to variability in data quality,

no significant associations between IQ scores and head motion were evident

($p > 0.05$; see LME results in Supplemental Tables 3-7 for each IQ subdomain score).

Collectively, these results demonstrate that individual variability within the

attention network in children associates with dissociable motifs of connectivity across

the functional connectome and covaries with the cognitive functioning. Children with

low ventral attention connectivity exhibit an accelerated profile of cortical maturation

that closely resembles prior reports in adolescents and adults. Although additional

longitudinal analyses are warranted, these data suggest a fundamental relationship

between individual variability within the ventral attention network and the age-

dependent changes in the macroscale properties of human brain organization and

cognitive ability.

Ventral attention network connectivity reliably associates with accelerated cortical maturation and cognitive ability across populations

The analyses above reveal a relationship between ventral attention network connectivity, the functional maturation of the cortical connectome, and cognitive functioning in a population of healthy developing children and adolescents from the Chinese Color Nest Project[32]. To examine the generalizability of the above results, we utilized 2186 participants from the ABCD[38] study, where participant data differed from the CCNP in sample population, study site, MRI scanner, acquisition parameters and longitudinal designs. Here, the ABCD data was limited to participants between 9 to 11 years old. Highlighting the robustness of the analyses reported above, we observed a profile of heightened degree centrality in ventral attention network areas that was consistent with the CCNP analyses (Figure 1A). Broadly, the ABCD degree centrality map demonstrated a lower degree centrality distribution along the cortex than the CCNP dataset (Figure 5A). Here, the inferior parietal gyrus and supramarginal gyrus, together with middle anterior cingulate cortex exhibited highly connected architecture. Hub regions of default network including the angular gyrus, middle temporal gyrus, and post cingulate cortex also revealed heightened degree centrality in the ABCD data, which might again indicate the role of default network as a significant cortical core across the transition to adolescence.

Next, we assessed the reliability of a relationship linking the maturation of functional gradient architecture of the cortical sheet with individual variability in ventral attention connectivity across populations. To do so, we split the children in the ABCD study in a manner consistent with the CCNP analyses to examine whether the ventral attention network connections are coupled with the functional gradient maturation. No significant associations were observed between the degree centrality of ventral attention network and demographic factors like age (p=0.21), gender (p=0.60), head motion (p=0.12) and family income (p=0.15). Further comparisons reveal matched demographic features between child groups in age (p=0.80), gender

(p=0.39), head motion (p=0.51) and family income (p=0.15). Consistent with our

findings in the CCNP children group, the gradient profiles of children with high ventral

attention network connectivity in the ABCD study (n=1367) displayed the

developmentally typical gradient profile (Figure 4B). In this group, the primary visual

and somato/motor areas occupied the ends of the primary gradient, matched the

second gradients in CCNP adolescents (r=0.896, p_{spin}≤0.001) and HCP adults

(r=0.93, p_{spin}≤0.001); while the second gradient revealed transmodal organization

matching the first gradients in CCNP adolescents (r=0.918, p_{spin}≤0.001) and HCP

adults (r=0.944, p_{spin}≤0.001). An accelerated developmental profile was revealed in

the ABCD children with low ventral attention network links (n=819), broadly matching

the primary and secondary gradients previously identified in CCNP adolescents

(Gradient 1: r=0.865, p_{spin}≤0.001; Gradient 2: r=0.856, p_{spin}≤0.001), and

demonstrating a hybrid organization comparing with HCP adults (correlations

between the first gradient in ABCD low attention group and gradients in HCP adults:

r=0.61, p_{spin}≤0.001, r=0.73, p_{spin}≤0.001; correlations between the second gradient in

ABCD low attention group and gradients in HCP adults: r=0.71, p_{spin}≤0.001, r=0.62,

p_{spin}=0.002). These data provide converging evidence, across independent collection

efforts, of an association between attention system connectivity and the broader

functional organization and maturational properties of cortex.

As a final step we repeated the behavior association analysis in the ABCD

project. Here, NIH toolbox scores were used to access the cognitive abilities

including crystallized (picture vocabulary and oral reading recognition) and fluid

components (pattern comparison processing speed, list sorting working memory,

picture sequence memory, flanker test and dimensional change card sort)[39]. As

above, a linear regression model was conducted to examine the relationship between

the degree centrality of the ventral attention network and cognitive functioning. Age,

sex and head motion were set as covariates in the model. Consistent with our CCNP

results, these analyses revealed the functional connections of ventral attention

network were significantly associated with the cognition total composite standard

score (p=0.0024), crystallized composite standard score (p=0.022), cognition fluid

composite standard score (p=0.0058), picture vocabulary (p=0.0048), and list sorting

working memory (p=0.0025). Other significant associations are reported in the

Supplemental Tables. Only the cognition total composite standard score (p=0.0024)

and list sorting working memory (p=0.0025) in the ABCD dataset were significant

after Bonferroni correction to address multiple comparisons across a total of 15

Linear Mixed Effects (LME) and Linear regression models, no significant association

was revealed in the CCNP dataset after the correction.

The replication analysis in the ABCD dataset highlight a potentially key role for

the ventral attention network in the maturation process of both cortical hierarchy and

cognitive ability. Notably, speaking to the robustness of the observed results, the

differences across datasets are quite substantial, including but not limited to

participant race, ethnicity, culture, and environment as well as scanner, scanning

parameters, socioeconomic factors, and education. In particular, the distinct

ethnic/racial composition of study samples can impact the generalizability of brain-

behavior associations[40,41]. Here, we applied the connection counts with ventral

attention network derived from the CCNP dataset directly on the children in the

ABCD dataset, and revealed consistent findings. Across sample collections, children

with high ventral attention network connectivity demonstrated typical developmental

patterns in functional gradients, while those with lower relative connectivity exhibited

adolescent- and adult-liked gradients. The degree centrality of the ventral attention

network is significantly associated with similar cognitive components in the CCNP

and ABCD datasets. Collectively, these analyses suggest a close and generalizable

relationship between the ventral attention network and the process of cortical

maturation, as reflected in the presence of macroscale functional gradients. These

data are in line with the hypothesis that the ventral attention network may

preferentially drive the refinement in the macroscale organization of cortex and

cognitive ability during the transition from childhood to adolescence.

512   **Discussion**

A fundamental goal of developmental systems neuroscience is to identify the

mechanisms that underlie the scheduled emergence and maturation of the large-

scale functional networks central to the information processing capabilities of the

human brain. Prior work has demonstrated age-dependent shifts in macroscale

organization of cortex[6,9], revealing the presence of a dominant profile of unimodal

cortical organization in children that is later replaced in adolescence by a distributed

transmodal architecture spanning from unimodal regions through association cortex.

In the current study, we further demonstrated that more connections were required in

children to attain a transmodal dominant feature (Extended Data Fig.1), indicating

that diffuse connections across broad swaths of association cortex gradually replace

the local connections in higher percentiles during adolescence. While this sweeping

developmental reorganization of network relationships across the cortical sheet

coincides with the transition from childhood to adolescence, reflecting an inflection

point at around age 12 to 14, the neurobiological processes underpinning these

profound functional changes have yet to be established. Here, by first identifying the

densely connected hub regions across development, we demonstrate that ventral

attention network regions[11] encompassing aspects of anterior insula, medial

prefrontal cortex, and dorsal anterior cingulate cortex/midline supplementary motor

area (also see salience[27] and cingulo-opercular[26] networks) may be linked to age-

dependent shifts in the macroscale organization of cortex across childhood and

adolescence.

Although the attention regions identified in the present analyses are recognized

as a single system at a coarse scale[42,43], this broad architecture is comprised of

multiple spatially adjacent but functionally dissociable networks[11,26,27,44]. This complex

system is theorized to underpin a cascade of functions supporting information

processing, from the control of stimulus driven attention[43] to the initialization of task

sets and maintenance of sustained attention during goal pursuit[29]. Further analysis

revealed that virtually lesioning the entire frontoparietal network also generates

transmodal organization in the first gradient (See Extended Data Fig.4). Importantly,

the salience/ventral attentional network was proposed to have a causal effect on

switching between default and frontoparietal networks during task execution[45]. The

hierarchical flow of information from sensory regions to deeper levels of cortical

processing is also reflected in the spatial continuity of associated functional parcels

along cortical surface, for example, the salience network[25,27], spanning orbital fronto-

insular and dorsal anterior cingulate areas through broad posterior areas of insula

and dorsal anterior cingulate cortex[25,29]. Broadly, information propagates along

posterior to anterior as well as dorsal to ventral axes across cortex and the

associated organization of functionally linked parcels is reflected through the

presence of continuous functional gradients[10,14]. Here, the ventral attention network is

situated at an intermediate position along this functional spectrum transiting from

primary sensory/motor networks to the default network that anchors association

cortex, which is commonly identified as the first gradient in adolescence and

adulthood, while reflecting the second gradient in childhood[14]. This transmodal

organizational profile has been inferred to reflect the hierarchy of information flow

across cortical territories[10]. Converging evidence for this functional motif has been

revealed through analyses of step-wise connectivity. Consistent with classic theories

regarding the integration of perceptual modalities into deeper layers of cortical

processing[23], functional relationships spread from primary somato/motor, visual and

auditory cortex before converging within ventral attention territories, and eventually

the default network[24]. As a whole, these data provide clear evidence situating the

ventral attention network between primary unimodal and association cortices,

highlighting a role for attention systems in the functional propagation of sensory

information to the multimodal regions that support higher-order cognitive functions.

The cingulo-opercular/ventral attention network, together with frontoparietal

network, have been proposed to constitute a parallel architecture of executive

functioning and cognitive control, supporting adaptive goal pursuit and flexible

behavioral adjustments[28]. Although the frontoparietal network, encompassing aspects

of dorsolateral prefrontal, dorsomedial prefrontal, lateral parietal, and posterior

temporal cortices, is functionally dissociable from cingulo-opercular/ventral attention

network in adulthood, studies in developmental populations have revealed that they

are functionally linked prior to adolescence[18]. Moreover, in children, the ventral

attention network possesses a broad distributed connectivity profile with reduced

segregation from salience networks[46], perhaps reflecting fluid community boundaries

in those areas. Anterior prefrontal cortex is shared by the two networks in children but

later segregated into ventral attention network in early adolescence, with dorsal

anterior cingulate cortex incorporated into frontoparietal network in adulthood[18]. This

aspect of network development is consistent with evidence for increased, but more

diffuse, patterns of task-evoked activity in children relative to adults[47], reflecting the

presence of scattered pattern of community assignments in prefrontal territories,

particularly those supporting attentional processes[22].

Suggesting a key role for the attentional systems in cognition, our present

analyses revealed reliable associations between attention network connectivity and

broad measures of intellectual functioning across populations. A recent study also

revealed abnormal functioning within limbic and insular networks and associated

cognitive deficits in adolescents who were extremely preterm at birth[48]. The

maturational course of large-scale brain network coincides with the emergence of

adult-like performance on cognitive tasks (e.g., processing speed, shifting and

response inhibition)[49,50]. A pattern of attention network development that is theorized

to reflect a transition from the over-representation of bottom-up attention to greater

top-down attentional and executive functioning capabilites[46]. Converging evidence

indicates that ventral attention network functions are intricately linked with the

frontoparietal network in both anatomy and function. Although speculative, the

corresponding segregation and integration of associated processes likely underpins

the maturation of adaptive human cognition.

598   In the present data, the observed maturational changes were not uniformly

distributed across cortical surface, suggesting a key role for the ventral attention

network during development. Intriguingly, the greatest functional changes were not

evident within the default or primary sensory territories that anchor the functional

gradients in adolescence and adulthood. Rather, ventral attention and frontoparietal

networks reflected the predominate sources of functional variation across childhood

and adolescence (Figure 1E). Of note, developmental changes were not exclusive to

attentional network regions, default network and the somato/motor network also

demonstrated age related variations across childhood and adolescence. These data

are consistent with prior reports suggesting that somato/motor and ventral attention

networks are functionally coupled in children, highlighting a distinct community

architecture in children relative to adults[22]. Even in adulthood, the ventral areas in

somato/motor network can be identified as a separate cluster from dorsal areas as

network parcellations increase in their granularity (i.e., 7-network relative to 17-

network resolutions in Yeo et al., 2011[11]; or in Gordon et al., 2017[44]).

613   Prior work has revealed a dramatic developmental change in the topography of

ventral attention network, while the spatial organization of default network

approaches an adult-like form in childhood[22]. However, it should be noted that the

developmental changes in brain's functional hierarchy are driven by comprehensive

reconfigurations across broad swaths of cortex, rather than solely reflecting

developmentally mediated shifts in the ventral attention network alone. As illustrated

in Figure 3, there are substantial alterations in both between- and within-network

connections across different age groups. This is particularly evident in the default

network, where a profile of age-linked increases in within-network connectivity

replaced between-network connections. These extensive changes along the cortical

surface indicate that network organization is not stabilized during development,

raising a critical issue that whether a given cortical area belongs to a fixed network

across puberty and into adulthood. Future studies should further establish an

accurate network affiliation for the developing population. Of note, the size and shape

of the human brain undergoes significant changes throughout the lifespan. While it's

indicated that the growth velocities of brain tissues peak before 6 years old[9], caution

is essential when interpreting MRI studies that span development populations.

Factors such as head motion and registration errors might bias the results.

Therefore, validating with large, independent datasets and alternate methods of

assessing brain functioning is crucial to assess the reproducibility, robustness, and

generalizability of the current results.

To further characterize how changes in ventral attention network connectivity

might underlie the broad functional maturation of cortex, we directly excluded the

developmentally dissociable aspects of ventral attention network from the brain

connectome and rederived the functional gradients previously characterized in

children, adolescence[14] and adults[10]. The present analyses revealed that virtually

lesioning ventral attention clusters characterized by the greatest developmental

changes generates adolescent- and adult-like gradient architectures in children, a

profile that was specific to attention network territories. Although follow-up work is

necessary, our analyses are consistent with the presence of a relative increase in

local connections between ventral attention network and adjacent somato/motor

territories in childhood, biasing a functional motif in childhood dominated by unimodal

organization. Later, diffuse connections across broad swaths of association cortex

emerge with age, until eventually the presence of interactions across distinct systems

becomes the dominant feature of cortical organization. The increased emphasis on

such transmodal integration is consistent with the developmental principle that, over

time, local connections are replaced by remote distributed interactions[13].

Individual level analysis further reveals the influence of ventral attention network

on the formation of adult-like functional organization. Children with fewer ventral

attention network connections exhibited an accelerated transmodal profile of cortical

organization, while the typical unimodal dominated architecture was presented in

children with more ventral attention connections. However, an open unanswered

question remains the manner through which ventral attention network functioning

may reshape the macroscale organization of cortex. Although speculative, the

observed results might reflect a consequence of repeated co-activation between

ventral attention network regions and associated cortical systems[18]. Accordingly, the

dense associations with somato/motor areas[22] as well as frontoparietal areas[18] in

children indicates ventral attention network is preferentially connected by both

primary and association cortex during development, which is also evident in our

degree centrality map (Figure 1A and B). While a dense profile of connections may

ensure system integrity, excessive or redundant connections are pruned across

development in order to optimize efficient information processing as neurons that are

not fully integrated within local circuits are eliminated to ensure stable network

function[51]. A profile of network segregation and integration that is hypothesized to

underpin functional specialization and computational efficiency[19,52].

Along with the stabilization of its functional organization, cortical maturation is

also marked by increased flexibility in resource allocation during task execution.

Evidence from task-based fMRI suggests that the salience/ventral attention network,

especially the right fronto-insular cortex, plays a critical role in switching between the

default mode and frontoparietal networks[45]. This finding supports our hypothesis that

the ventral attention network not only serves as a transfer hub between unimodal and

transmodal cortices but also coordinates the information flow within the transmodal

cortex. In addition to refining the unstable local connections associated with unimodal

areas, the ventral attention network may drive the predominance of transmodal

organization through a parallel mechanism, which likely involves strengthening the

transmodal interactions among higher-order networks to ensure the flexible

reallocation of resources. Future work should focus on characterizing the specific

properties of ventral attention network across development, for instance, extending to

early points in the lifespan, examining its interactions with other network under the

background of task execution or considering the roll of individual experience on

network development[53].

Accumulating evidence has revealed a dramatic restructuring of the macroscale

organization of the cortex across development, suggesting the scheduled maturation

of functional gradient patterns may be critically important for understanding how

cognitive and behavioral capabilities are refined across development. However, the

underlying developmental processes driving these functional changes remain to be

established. Here, by first localizing the significant maturational changes in functional

connectome across childhood and adolescence, we demonstrate that the ventral

attention network may play a critical role in the onset of the age-dependent shifts that

characterize the macroscale organization of cortex across development. Children

with fewer functional connections within the ventral attention network exhibit the

appearance of an accelerated maturation of gradient architecture and increased

cognitive functioning. Although the process of brain maturation emerges through

complex interactions across environmental experience and biological systems that

span genes and molecules through cells, networks, and behavior, the current

findings suggest a core role for attention network-linked territories. The multiscale

interactions linking the longitudinal development of the ventral attention network with

these maturational processes, and the associated consequences on behavior across

health and disease, remains an open question to be answered in future work.

Acknowledgements: This work was supported by the STI 2030 - the major projects

of the Brain Science and Brain-Inspired Intelligence Technology (2021ZD0200500 to

X.N.Z.), the National Institute of Mental Health (Grants R01MH120080 and

R01MH123245 to A.J.H.), the Major Fund for International Collaboration of National

Natural Science Foundation of China (81220108014 to X.N.Z.), the National Basic

Science Data Center "Interdisciplinary Brain Database for In vivo Population Imaging"

(ID-BRAIN to X.N.Z.) and the Start-up Funds for Leading Talents at Beijing Normal

University (to X.N.Z.). BTTY is supported by the NUS Yong Loo Lin School of

Medicine (NUHSRO/2020/124/TMR/LOA), the Singapore National Medical Research

Council (NMRC) LCG (OFLCG19May-0035), NMRC CTG-IIT (CTGIIT23jan-0001),

NMRC STaR (STaR20nov-0003), Singapore Ministry of Health (MOH) Centre Grant

(CG21APR1009), the Temasek Foundation (TF2223-IMH-01), and the United States

National Institutes of Health (R01MH120080 & R01MH133334). Any opinions,

findings and conclusions or recommendations expressed in this material are those of

the authors and do not reflect the views of the Singapore NRF, NMRC or MOH.

Author Contributions: H-MD, AJH and X-NZ designed the research. AJH and X-NZ

supervised the research. H-MD and AJH conducted analyses and made figures. X-

HZ, LL, S-SZ and LRO conducted validation analyses based on the ABCD dataset.

All authors contributed during writing and edited the paper.

Declaration of Interests: The authors declare they have no competing interests.

<div align="center">**Figure Legends**</div>

Figure 1. Ventral attention network areas demonstrate high population-level

degree centrality but pronounced functional changes across development.

Degree Centrality maps in children (A) and adolescents (B) reveal consistent dense

connectivity in ventral attention network areas throughout development. Scale bar

reflects the count of above threshold connections from a given vertex to all other

vertices. Larger values indicate higher degree centrality. (C) Euclidean Distance of

the functional connectome at each vertex between children and adolescents reveals

a clear switch within the ventral attention network. Larger values indicate greater

dissimilarity. (D) Regions based on the Schaefer et al., 400-parcel[31] atlas and colored

by the Yeo et al., 7-network solution[11]. (E) Bar graph reflects changes in the

Euclidean Distance of functional connectome at network level (mean network values

± standard error). The ventral attention network (6.5518±1.2002) shows the largest

developmental change while visual network (4.2584±1.1257) and default networks

(4.9813±1.1207) are most stable between children and adolescents. DorsAttn, dorsal

attention (5.1157±1.2499); Sal, salience; Som/Mot, somato/motor (5.8618±1.6067);

VentAttn, ventral attention; Control (5.4065±1.1681); and Limbic (6.2337±2.6297).

Figure 2: Ventral attention territories play a core role in the expression of

adolescent-like connectivity gradients. (A) The principal cortical gradients of

connectivity in children, adolescents (data from Dong et al., 2021[14]) and adults (data

from Margulies et al., 2016[10]). (B) Clusters with maximum developmental changes in

Euclidean Distance were extracted (Figure 1c), denoted as black on the cortical

surface. Associated vertices were then dropped from the cortical connectome in the

child group prior to rederiving the gradients. Results reveal that default and visual

networks anchor the extremes along the principal gradient (Gradient 1), mirroring the

principal gradient in both adolescents (r=0.68, p_{spin}≤0.001, two-sided spin test) and

adults (r=0.84, p_{spin}≤0.001, two-sided spin test). In children, the rederived second

gradient (Gradient 2) revealed a unimodal architecture separating somato/motor

network from visual network, which closely corresponds on the second gradient in

both adolescents (r=0.66, p_{spin}≤0.001, two-sided spin test) and adults (r=0.85,

p_{spin}≤0.001, two-sided spin test). (C) To further assess the significance of the results,

we constructed a permuted null model in the present data. Clusters with same size

and shape as (B) but shuffled locations on the cortical sheet were generated and

excluded from analyses in child group 500 times. For each shuffle, the principal

gradient was extracted and compared with the principal gradient in adults. The

permuted null model shows only 1 case revealing a higher correlation than that

observed in the real data, the x-axis indicates the absolute correlation values, the y-

axis indicates the frequency of correlations, dotted line refers to the hypothesis being

tested.

Figure 3: Developmental shifts in functional connectivity between childhood

and adolescence. Differences in network-level connectome (The degree centrality

matrix of adolescents subtracts the degree centrality matrix of children) functioning

between children and adolescents are demonstrated in a chord diagram. For each

network, the number connections within the top 10% of each vertex are first

summarized at network level, displayed as the links from right half circle with larger

radius to left half circle with smaller radius. Reflecting the asymmetrical nature of the

thresholded connectivity matrix, links from the left to right half circle indicate the

connections within each network that are included in the top 10% of connections to

other vertices. Line width highlights the number of connections, with broader lines

indicating increased connectivity. Dark colored lines indicate increased values in

adolescents, relative to children. Light color lines indicate decrease values in the

adolescent group.

Figure 4: Individual differences in ventral attention network connectivity reveal

a functional profile that resembles accelerated cortical maturation in some

children. (A) The number of functional connections linked with ventral attention

network at individual level. Male participants are marked in blue, female participants

are marked in red. Repeated imaging sessions within the same participants are

linked by lines. The x-axis represents age range from 6 to 18 years. The y-axis

reflects the number of connections that are linked to the ventral attention network

following thresholding. Participants were divided into high and low ventral attention

connectivity groups according to the median value in age 17-year-old participants

(dotted line). (B) The gradient maps in high ventral attention connectivity groups

reveal typical patterns identified in our previous work in both children and

adolescents[14]. (C) The gradient maps in low ventral attention connectivity groups

reveal an accelerated maturation process in children, with both the primary and

secondary gradients demonstrated a transmodal architectures. Functional

organization was broadly preserved across the high and low attention groups in

adolescents, with a subtle muting of the ventral attention gradient values in the low

attention participant group. The primary gradient maps in (D) high and low ventral

attention connectivity groups derived from longitudinal data. These surface maps

display a developmentally normative pattern of gradient reversals in the high ventral

attention group children, later scanned in adolescence. Conversely, low ventral

attention group children exhibit a stable adolescent-like gradient architecture in both

childhood and adolescence. See Extended Data Fig.2 & 3 for the longitudinal

analyses of Gradient 1 and Gradient 2 in both hemispheres.

Figure 5: Relationship between ventral attention network connectivity and

accelerated cortical maturation reliable across independent datasets. (A)

Degree Centrality maps in children (9~11 years old, n=2186) from the ABCD project

reveals dense connectivity in ventral attention network areas in a pattern that is

consistent with the CCNP participants (Figure 1A). Scale bar reflects the count of

above threshold connections from a given vertex to all other vertices. Larger values

indicate higher degree centrality. (B) The gradient maps in the ABCD high ventral

attention connectivity groups reveal typical patterns identified in our previous work in

children[14] as well as the CCNP high ventral attention group (Figure 4B,C children).

(C) The gradient maps in the ABCD low ventral attention connectivity groups closely

resemble the profile of accelerated maturation observed in the CCNP sample.

References

1 Casey, B. J., Heller, A. S., Gee, D. G. & Cohen, A. O. Development of the

emotional brain. *Neurosci Lett* **693**, 29-34, doi:10.1016/j.neulet.2017.11.055

(2019).

2 Casey, B. J., Getz, S. & Galvan, A. The adolescent brain. *Dev Rev* **28**, 62-77,

doi:10.1016/j.dr.2007.08.003 (2008).

3 Luna, B. *et al.* Maturation of widely distributed brain function subserves

cognitive development. *Neuroimage* **13**, 786-793,

doi:10.1006/nimg.2000.0743 (2001).

4 Kang, H. J. *et al.* Spatio-temporal transcriptome of the human brain. *Nature*

478, 483-489, doi:10.1038/nature10523 (2011).

5 Huttenlocher, P. R. & Dabholkar, A. S. Regional differences in

synaptogenesis in human cerebral cortex. *J Comp Neurol* **387**, 167-178,

doi:10.1002/(Sici)1096-9861(19971020)387:2<167::Aid-Cne1>3.0.Co;2-Z

(1997).

6 Paquola, C. *et al.* Shifts in myeloarchitecture characterise adolescent

development of cortical gradients. *Elife* **8**, doi:10.7554/eLife.50482 (2019).

7 Zilles, K., Palomero-Gallagher, N. & Amunts, K. Development of cortical

folding during evolution and ontogeny. *Trends Neurosci* **36**, 275-284,

doi:10.1016/j.tins.2013.01.006 (2013).

8 Reardon, P. K. *et al.* Normative brain size variation and brain shape diversity

in humans. *Science* **360**, 1222-1227, doi:10.1126/science.aar2578 (2018).

9 Bethlehem, R. A. I. *et al.* Brain charts for the human lifespan. *Nature*, **604**,

525-533, doi:10.1038/s41586-022-04554-y (2022).

10 Margulies, D. S. *et al.* Situating the default-mode network along a principal

gradient of macroscale cortical organization. *Proc Natl Acad Sci U S A* **113**,

12574-12579, doi:10.1073/pnas.1608282113 (2016).

852 11 Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by

853   intrinsic functional connectivity. *J Neurophysiol* **106**, 1125-1165,

854   doi:10.1152/jn.00338.2011 (2011).

855 12 Gao, W. *et al.* Temporal and spatial evolution of brain network topology

856   during the first two years of life. *PLoS One* **6**, e25278,

857   doi:10.1371/journal.pone.0025278 (2011).

858 13 Fair, D. A. *et al.* Functional Brain Networks Develop from a "Local to

859   Distributed" Organization. *Plos Comput Biol* **5**, e1000381,

860   doi:10.1371/journal.pcbi.1000381 (2009).

861 14 Dong, H. M., Margulies, D. S., Zuo, X. N. & Holmes, A. J. Shifting gradients of

862   macroscale cortical organization mark the transition from childhood to

863   adolescence. *Proc Natl Acad Sci U S A* **118**, e2024448118,

864   doi:10.1073/pnas.2024448118 (2021).

865 15 Somerville, L. H., Hare, T. & Casey, B. J. Frontostriatal maturation predicts

866   cognitive control failure to appetitive cues in adolescents. *J Cogn Neurosci*

867   **23**, 2123-2134, doi:10.1162/jocn.2010.21572 (2011).

868 16 Tottenham, N. & Sheridan, M. A. A review of adversity, the amygdala and the

869   hippocampus: a consideration of developmental timing. *Front Hum Neurosci*

870   **3**, 68, doi:10.3389/neuro.09.068.2009 (2009).

871 17 Sydnor, V. J. *et al.* Neurodevelopment of the association cortices: Patterns,

872   mechanisms, and implications for psychopathology. *Neuron* **109**, 2820-2846,

873   doi:10.1016/j.neuron.2021.06.016 (2021).

874 18 Fair, D. A. *et al.* Development of distinct control networks through segregation

875   and integration. *Proc Natl Acad Sci USA* **104**, 13507-13512,

876   doi:10.1073/pnas.0705843104 (2007).

877 19 Betzel, R. F. *et al.* Changes in structural and functional connectivity among

878   resting-state networks across the human lifespan. *Neuroimage* **102 Pt 2**, 345-

879   357, doi:10.1016/j.neuroimage.2014.07.067 (2014).

20   Betzel, R. F. *et al.* Generative models of the human connectome. *Neuroimage*

124, 1054-1064, doi:10.1016/j.neuroimage.2015.09.041 (2016).

21   Zuo, X. N. *et al.* Human Connectomics across the Life Span. *Trends Cogn Sci*

21, 32-45, doi:10.1016/j.tics.2016.10.005 (2017).

22   Tooley, U. A., Bassett, D. S. & Mackey, A. P. Functional brain network

community structure in childhood: Unfinished territories and fuzzy boundaries.

Neuroimage **247**, 118843, doi:10.1016/j.neuroimage.2021.118843 (2022).

23   Mesulam, M. M. From sensation to cognition. *Brain* **121 (Pt 6)**, 1013-1052,

doi:10.1093/brain/121.6.1013 (1998).

24   Sepulcre, J., Sabuncu, M. R., Yeo, T. B., Liu, H. & Johnson, K. A. Stepwise

connectivity of the modal cortex reveals the multimodal organization of the

human brain. *J Neurosci* **32**, 10649-10661, doi:10.1523/JNEUROSCI.0759-

12.2012 (2012).

25   Power, J. D. *et al.* Functional network organization of the human brain.

Neuron **72**, 665-678, doi:10.1016/j.neuron.2011.09.006 (2011).

26   Dosenbach, N. U. *et al.* Distinct brain networks for adaptive and stable task

control in humans. *Proc Natl Acad Sci U S A* **104**, 11073-11078,

doi:10.1073/pnas.0704320104 (2007).

27   Seeley, W. W. *et al.* Dissociable intrinsic connectivity networks for salience

processing and executive control. *J Neurosci* **27**, 2349-2356,

doi:10.1523/JNEUROSCI.5587-06.2007 (2007).

28   Dosenbach, N. U., Fair, D. A., Cohen, A. L., Schlaggar, B. L. & Petersen, S.

E. A dual-networks architecture of top-down control. *Trends Cogn Sci* **12**, 99-

105, doi:10.1016/j.tics.2008.01.001 (2008).

29   Dosenbach, N. U. *et al.* A core system for the implementation of task sets.

Neuron **50**, 799-812, doi:10.1016/j.neuron.2006.04.031 (2006).

30   Labache, L., Ge, T., Yeo, B. T. T. & Holmes, A. J. Language network
lateralization is reflected throughout the macroscale functional organization of
cortex. *Nat Communn* **14**, 3405, doi: 10.1038/s41467-023-39131-y (2023).
31   Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex
from Intrinsic Functional Connectivity MRI. *Cereb Cortex* **28**, 3095-3114,
doi:10.1093/cercor/bhx179 (2018).
32   Liu, S. *et al.* Chinese Color Nest Project : An accelerated longitudinal brain-
mind cohort. *Dev Cogn Neurosci* **52**, 101020, doi:10.1016/j.dcn.2021.101020
(2021).
33   Alexander-Bloch, A. F. *et al.* On testing for spatial correspondence between
maps of human brain structure and function. *Neuroimage* **178**, 540-551,
doi:10.1016/j.neuroimage.2018.05.070 (2018).
34   Gogtay, N. *et al.* Dynamic mapping of human cortical development during
childhood through early adulthood. *Proc Natl Acad Sci USA* **101**, 8174-8179,
doi:DOI 10.1073/pnas.0402680101 (2004).
35   Langs, G. *et al.* Identifying Shared Brain Networks in Individuals by
Decoupling Functional and Anatomical Variability. *Cereb Cortex* **26**, 4004-
4014, doi:10.1093/cercor/bhv189 (2016).
36   Coifman, R. R. & Lafon, S. Diffusion maps. *Appl Comput Harmon A* **21**, 5-30,
doi:10.1016/j.acha.2006.04.006 (2006).
37   Fan, X. R. *et al.* A longitudinal resource for population neuroscience of
school-age children and adolescents in China. *Sci Data* **10**, 545,
doi:10.1038/s41597-023-02377-8 (2023).
38   Casey, B. J. *et al.* The Adolescent Brain Cognitive Development (ABCD)
study: Imaging acquisition across 21 sites. *Dev Cogn Neurosci* **32**, 43-54,
doi:10.1016/j.dcn.2018.03.001 (2018).
39   Luciana, M. *et al.* Adolescent neurocognitive development and impacts of
substance use: Overview of the adolescent brain cognitive development

(ABCD) baseline neurocognition battery. *Dev Cogn Neurosci* **32**, 67-79,

doi:10.1016/j.dcn.2018.02.006 (2018).

40   Ricard, J. A. *et al.* Confronting racially exclusionary practices in the

acquisition and analyses of neuroimaging data. *Nat Neurosci* **26**, 4-11,

doi:10.1038/s41593-022-01218-y (2023).

41   Li, J. *et al.* Cross-ethnicity/race generalization failure of behavioral prediction

from resting-state functional connectivity. *Sci Adv* **8**, eabj1812,

doi:10.1126/sciadv.abj1812 (2022).

42   Fox, M. D. *et al.* The human brain is intrinsically organized into dynamic,

anticorrelated functional networks. *Proc Natl Acad Sci U S A* **102**, 9673-9678,

doi:10.1073/pnas.0504136102 (2005).

43   Corbetta, M. & Shulman, G. L. Control of goal-directed and stimulus-driven

attention in the brain. *Nat Rev Neurosci* **3**, 201-215, doi:10.1038/nrn755

(2002).

44   Gordon, E. M. *et al.* Precision Functional Mapping of Individual Human

Brains. *Neuron* **95**, 791-807 e797, doi:10.1016/j.neuron.2017.07.011 (2017).

45   Sridharan, D., Levitin, D. J., & Menon, V. A critical role for the right fronto-

insular cortex in switching between central-executive and default-mode

networks. *Proc Natl Acad Sci U S A* **105**, 12569-12574,

doi:10.1073/pnas.0800005105 (2008).

45   Farrant, K. & Uddin, L. Q. Asymmetric development of dorsal and ventral

attention networks in the human brain. *Dev Cogn Neurosci* **12**, 165-174,

doi:10.1016/j.dcn.2015.02.001 (2015).

46   Casey, B. J., Giedd, J. N. & Thomas, K. M. Structural and functional brain

development and its relation to cognitive development. *Biol Psychol* **54**, 241-

257, doi:10.1016/s0301-0511(00)00058-2 (2000).

47   Molloy, M. F. *et al.* Effect of Extremely Preterm Birth on Adolescent Brain

Network Organization. *Brain Connect* **13**, 394-409,

doi:10.1089/brain.2022.0077 (2023).

48   Huizinga, M., Dolan, C. V. & van der Molen, M. W. Age-related change in

executive function: developmental trends and a latent variable analysis.

Neuropsychologia **44**, 2017-2036,

doi:10.1016/j.neuropsychologia.2006.01.010 (2006).

49   Luna, B., Garver, K. E., Urban, T. A., Lazar, N. A. & Sweeney, J. A.

Maturation of cognitive processes from late childhood to adulthood. *Child Dev*

75, 1357-1372, doi:10.1111/j.1467-8624.2004.00745.x (2004).

50   Pfisterer, U. & Khodosevich, K. Neuronal survival in the brain: neuron type-

specific mechanisms. *Cell Death Dis* **8**, e2643, doi:10.1038/cddis.2017.64

(2017).

51   Bullmore, E. & Sporns, O. The economy of brain network organization. Nat

Rev Neurosci **13**, 336-349, doi:10.1038/nrn3214 (2012).

52   Gee, D. G. *et al.* Early developmental emergence of human amygdala-

prefrontal connectivity after maternal deprivation. *Proc Natl Acad Sci U S A*

110, 15638-15643, doi:10.1073/pnas.1307893110 (2013).

53   Dong, H. M. *et al.* Charting brain growth in tandem with brain templates at

school age. *Sci Bull* **65**, 1924-1934, doi:10.1016/j.scib.2020.07.027 (2020).

 Methods

Datasets

Chinese Color Nest Project (CCNP): CCNP is a five-year accelerated longitudinal

study across the human life span[21,32,54]. A total of 176 scans in adolescents and 202

scans in typically developing children were included in the analysis, details of the

dataset and exclusion criteria can be found in our previous work[14]. All MRI data was

obtained with a Siemens Trio 3.0T scanner at the Faculty of Psychology, Southwest

University in Chongqing. The reported experiments were approved by the

Institutional Review Board from Institute of Psychology, Chinese Academy of

Sciences. All participants and their parents/guardians provided written informed

consent before participating in the study.

Adolescent Brain Cognitive Development Study (ABCD): ABCD is a multi-site

longitudinal cohort following the brain and cognition development of over ten

thousand 9~10 years old children. MRI scans including T1-weighted, T2-weighted

and resting-state fMRI was obtained with 3T Siemens Prisma, General Electric 750

and Phillips scanners across 21 sites, details of the scan parameters can be found in

reference[5]. Here, we unitized the MRI baseline data from 2186 children (female

54.4%, mean age 10.01 years old) for the reproducibility analysis. The study was

approved by the Institutional Review Board from the University of California, San

Diego[55]. All participants and their parents/guardians provided written informed

consent[56].

1005

MRI Data Preprocessing

CCNP dataset: Anatomical T1 images were visually inspected to exclude individuals

with substantial head motion and structural abnormalities. Next, T1 images were fed

into the volBrain pipeline (http://volbrain.upv.es)[57] for noise removal, bias correction,

intensity normalization and brain extraction. All brain extractions underwent visual

inspection to ensure tissue integrity. After initial quality checks, T1 images were

passed into the Connectome Computation System (CCS)[58] for surface-based

analyses. CCS pipeline is designed for preprocessing multimodal MRI datasets and

integrates publicly available software including SPM[59], FSL[60], AFNI[61] and

FreeSurfer[62]. Resting-state fMRI data preprocessing included a series of steps

common to intrinsic functional connectivity analyses: (1) dropping the first 10s (4

TRs) for the equilibrium of the magnetic field; (2) estimating head motion parameters

and head motion correction; (3) slicing time correction; (4) time series de-spiking; (5)

registering functional images to high resolution T1 images using boundary-based

registration; (6) removing nuisance factors such as head motion, CSF and white

matter signals using ICA-AROMA[63]; (7) removing linear and quadratic trends of the

time series; (8) projecting volumetric time series to surface space (the *fsaverage5*

model with medial wall masked out); (9) 6mm spatial smoothing. All preprocessing

scripts are publicly available on GitHub (https://github.com/zuoxinian/CCS). Any

resting-state scan with a mean head motion above 0.5 mm was excluded from further

analysis. The demographic information of subjects included in the analyses is listed

in Supplemental Table 1.

1028

ABCD dataset: Minimally preprocessed T1 images[64] were fed into Freesurfer[62] for

surface reconstruction. Resting-state fMRI data[64] preprocessing included a series of

steps as following: (1) dropping the initial frames for the equilibrium of the magnetic

field; (2) estimating head motion parameters and voxel-wise differentiated signal

variance (DVARS) and head motion correction; (3) registering functional images to

high resolution T1 images using boundary-based registration; (4) Scrubbing the

frames with FD > 0.3 mm or DVARS > 50, along with one volume before and two

volumes after. (5) removing nuisance factors such as global signal, head motion,

CSF and white matter signals; (6) band-pass filtered ($0.009\,Hz \leq f \leq 0.08\,Hz$); (7)

projecting volumetric time series to surface space (the fsaverage5 model with medial

wall masked out); (8) 6mm spatial smoothing. Full details of data preprocessing can

be found in previous study[65]. All preprocessing scripts are publicly available

(https://github.com/ThomasYeoLab/ABCD_scripts) on GitHub. Any resting-state scan

with a max head motion above 5 mm and over half of their volumes censored was

excluded from further analysis.

1044

Degree Centrality Mapping. Functional connectivity (FC) matrices and the

corresponding Fisher-z transformed values were first generated for each resting-scan

per visit. Then the two test-retest FC fisher-z (FCz) matrices within one visit were

averaged to increase signal-to-noise ratio for generating individual FCz matrix for

each visit, which was later averaged across individuals to form group-level FCz

matrices. For the group-level FCz matrices, only the top 10% functional

connenctivities of each vertex were retained, other elements and negative FCs in the

matrix were set to 0 to enforce sparsity, yielding an asymmetrical matrix, of which the

rows are corresponding to the connectome of each vertex. Degree centrality map

was obtained by counting the non-zero elements in each column of the FCz matrix.

We then calculated the cosine distance between any two rows of the FCz matrix and

subtracted from 1 to obtain a symmetrical similarity matrix, this similarity matrix was

later used to derive the gradients.

1058

Euclidean Distance. To characterize the developmental changes between children

and adolescents, the Euclidean distance was computed for each row of the cosine

distance matrix between children and adolescents. Then most changed clusters were

extracted according to the following two criteria: the top 10% in Euclidean distance

map and the cluster size above 500 vertexes. One-way ANOVAs were performed to

test the statistical differences between networks.

1065

Gradients Analysis. The extracted clusters and their FCs were firstly dropped in the

initial FCz matrix of children group, and then cosine similarity matrix was calculated.

Diffusion map embedding[10,35] was implemented on the similarity matrix to derive

gradients (https://github.com/NeuroanatomyAndConnectivity/gradient_analysis).

Within each age bin, the functional connectivity matrix from participants with repeated

imaging scans were averaged with scans of other participants to generate a group-

level matrix, and then used to derive functional gradients. Pearson correlations were

computed between the derived gradients in children and adults. To examine the

statistical significance of the observed Pearson correlations, null distributions were

generated by randomly rotated the locations of the extracted clusters across the

cortical surface while keeping the shape and size fixed. For each permutation, the

gradients and correlations were rederived. A total of 500 permutations were

performed to generate the null model.

1079

Chord diagram. Chord diagram was utilized to demonstrate the differences in

numbers of functional connectivity between children and adolescents at network

level. For each vertex, its connectome was represented as the corresponding row in

the FCz matrix. The associated number of network-level connections was obtained

by counting the non-0 elements in each network according to the Yeo 7-network

solution, generating a matrix with dimension 20,484 (number of vertex) by 7 (number

of networks). Vertexes were grouped into networks to generate the final network to

network FC matrix (7 by 7). The rows of this network-to-network matrix are displayed

as the links from right half circus with larger radius to the left smaller half circus,

referring to the FCs with networks in its own connectome, the opposite links from left

to right circus represent the columns in the network-to-network matrix, referring to the

FCs of each network that were existed in other networks' connectome (Figure 3).

1092

High and Low Ventral Attention Network Group Definition. Ventral attention network linked connections were extracted as the corresponding column in network-to-network matrix, referring to the functional connectivity other networks linked with ventral attention network. As we found in prior work[14], gradient maps became stable in 15-year-old participants, although they still possess a hybrid organization relative to adult participants. The gradient pattern in 17-year-old participants closely resembles what is observed in adults. As such, we hypothesized that the functional connections may reach a stable status at the end of adolescence when compared with younger participants, thus we take the median value of functional connectivity number in 17-year-old age group as the reference. Any single scan with a connectivity number above the threshold was assigned to the "high" ventral attention subgroup, other scans were assigned to the "low" ventral attention group. The gradients were then rederived for each group.

Association Analysis with IQ Score in the CCNP Dataset. The association between connections number with ventral attention network and IQ scores was estimated with LME model. IQ scores were obtained by Wechsler Children Intelligence Scale IV, including scores in following subdomains: verbal, perception reasoning, working memory, processing speed ability. The LME was conducted using the following formula:

$$\text{IQ\_score} \sim 1 + \text{Age} + \text{Gender} + \text{(Ventral attention)\_DC} + \text{HeadMotion} + (1 \mid \text{Subid})$$

Here (Ventral attention)_DC refers to the connections with ventral attention network, together with age, gender, head motion and the intercept set as the fixed effect factor. Subid refers to the participant IDs, multi measurements for a single participant were coded as an identical nominal variable, set as the random effect factor. LME models were applied for the total IQ and subdomain scores separately.

1121

Association Analysis with Cognitive Score in the ABCD Dataset. Linear regression

model was applied to estimate the association between connections number with

ventral attention network and cognitive scores in the ABCD dataset. Cognitive scores

were accessed by NIH toolbox, including scores in following domains: crystallized

(picture vocabulary and oral reading recognition) and fluid components (pattern

comparison processing speed, list sorting working memory, picture sequence

memory, flanker test and dimensional change card sort)[10]. The model was conducted

using the following formula in MATLAB:

$$\text{Cognitive\_score} \sim 1 + \text{Age} + \text{Gender} + (\text{Ventral attention})\_\text{DC} + \text{HeadMotion}$$

1131

Developmental Effects on Degree Centrality of Ventral Attention Network. The

association between connections number with ventral attention network and age,

gender head motion was also estimated with LME model, which was conducted

using the following formula in MATLAB:

$$(\text{Ventral attention})\_\text{DC} \sim 1 + \text{Age} + \text{Gender} + \text{HeadMotion} + (1 \mid \text{Subid})$$

1137

1138

Data Availability: Data from the CCNP dataset used here are available at Chinese

Color Nest Project (CCNP) – Lifespan Brain-Mind Development Data Community at

Science Data Bank (https://ccnp.scidb.cn/en) including both anonymized

neuroimaging data (https://doi.org/10.57760/sciencedb.07860) and unthresholded

whole-brain connectivity matrices grouped by relevant ages (children and

adolescents) (https://doi.org/10.11922/sciencedb.00886). The raw CCNP data are

available from the website upon reasonable request. The ABCD data used in this

report came from the Annual Release 2.0 (https://doi.org/10.15154/1503209) of the

ABCD BIDS Community Collection (ABCC; NDA Collection 3165). Source data are

provided with this paper.

1149

1150   **Code Availability:** Code is available online at our GitHub page:

1151   https://github.com/HolmesLab/GradientMaturation

1152

		Methods References

1153   **Methods References**

Auchter, A. M. *et al.* A description of the ABCD organizational structure and communication framework. *Dev Cogn Neurosci* **32**, 8-15, doi:10.1016/j.dcn.2018.04.003 (2018).

Clark, D. B. *et al.* Biomedical ethics and clinical oversight in multisite observational neuroimaging studies with children and adolescents: The ABCD experience. *Dev Cogn Neurosci* **32**, 143-154, doi:10.1016/j.dcn.2017.06.005 (2018).

Manjon, J. V. & Coupe, P. volBrain: An Online MRI Brain Volumetry System. *Front Neuroinform* **10**, 30, doi:10.3389/fninf.2016.00030 (2016).

Xu, T. *et al.* A Connectome Computation System for discovery science of brain. *Sci Bull* **60**, 86-95, doi:10.1007/s11434-014-0698-3 (2015).

Friston, K. J. *et al.* Statistical parametric maps in functional imaging: a general linear approach. *Hum Brain Mapp* **2**, 189-210, doi:10.1002/HBM.460020402 (1994).

Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W. & Smith, S. M. FSL. *Neuroimage* **62**, 782-790 (2012).

Cox, R. W. AFNI: Software for analysis and visualization of functional magnetic resonance neuroimages. *Comput Biomed Res* **29**, 162-173 (1996).

Fischl, B. FreeSurfer. *Neuroimage* **62**, 774-781, doi:10.1016/j.neuroimage.2012.01.021 (2012).

Pruim, R. H. R. *et al.* ICA-AROMA: A robust ICA-based strategy for removing motion artifacts from fMRI data. *Neuroimage* **112**, 267-277, doi:10.1016/j.neuroimage.2015.02.064 (2015).

Hagler, D. J., Jr. *et al.* Image processing and analysis methods for the Adolescent Brain Cognitive Development Study. *Neuroimage* **202**, 116091, doi:10.1016/j.neuroimage.2019.116091 (2019).

1180   64   Chen, J. *et al.* Shared and unique brain network features predict cognitive,

1181   personality, and mental health scores in the ABCD study. *Nat Commun* **13**,

1182   2217, doi:10.1038/s41467-022-29766-8 (2022).

A Children

B Adolescents

Degree Centrality
1   5260

C

Euclidean Distance
2.6   10.0

D

● Default   ● Limbic   ● Som/Motor
● Control   ● Sal/VentAttn   ● Visual
● DorsAttn

E Functional Network-level
Developmental Changes

A  Children     Adolescents     Adults

-10 ▮ 8    -10 ▮ 10    -6 ▮ 6
Gradient 1 values   Gradient 1 values   Gradient 1 values

Dropping Ventral Attention Clusters
Generates Adolescent-Like Gradient Architectures in Children

B  Gradient 1    Gradient 2   C Permuted Null Model Dropping
                    Random Clusters (Spin Test)

-7 ▮ 8     -7 ▮ 9
Gradient Values    Gradient Values

Pearson Correlations of Primary
Gradients in Children and Adults

Functional Connectivity Developmental Changes
Degree Centrality$_{\text{Adolescents}}$ - Degree Centrality$_{\text{Children}}$

A

High Salience and Ventral Attention Connectivity Participants

B

Gradient 1		Gradient 2	
Children	Adolescents	Children	Adolescents

Low Salience and Ventral Attention Connectivity Participants

C

Gradient 1		Gradient 2	
Children	Adolescents	Children	Adolescents

A ABCD
Degree Centrality

1 — 4349
Degree Centrality

B ABCD
High Ventral Attention

Gradient 1   Gradient 2

ABCD
Low Ventral Attention

Gradient 1   Gradient 2