Title: Reduced ventral attention network connectivity is linked to the accelerated maturation of adult-like cortical organization in childhood

Authors: Hao-Ming Dong[1,2], Avram Holmes[1,3,4]*, Xi-Nian Zuo[2,5,6]*

[1] Department of Psychology, Yale University, New Haven, Connecticut, USA

[2] State Key Laboratory of Cognitive Neuroscience and Learning, Beijing Normal University, Beijing 100875, China

[3] Department of Psychiatry, Yale University, New Haven, Connecticut, USA

[4] Wu Tsai Institute, Yale University, New Haven, Connecticut, USA

[5] National Basic Science Data Center, Beijing 100190, China

[6] Developmental Population Neuroscience Research Center, IDG/McGovern Institute for Brain Research, Beijing Normal University, Beijing 100875, China

Correspondence: * To whom correspondence may be addressed. Email: avram.holmes@yale.edu or xinian.zuo@bnu.edu.cn.

Author Contributions: All authors designed the research. H-MD conducted analyses and made figures. All authors contributed during writing. All authors edited the paper.

Keywords: salience/ventral attention network, gradients, cortical organization, development, adolescence

Acknowledgements: This work was supported by the National Institute of Mental Health (Grants R01MH120080 and R01MH123245 to A.J.H.), the Child Brain-Mind Development Cohort Study in China Brain Initiative (SQ2021AAA010024 to X.N.Z.), the Start-up Funds for Leading Talents at Beijing Normal University (to X.N.Z.), the National Natural Science Foundation of China (Major Fund for International Collaboration: 81220108014 to X.N.Z.) and the National Basic Science Data Center

"Chinese Data-sharing Warehouse for In-vivo Imaging Brain" (NBSDC-DB-15 to X.N.Z.).

Abstract

Across the transition from childhood to adolescence the human brain experiences profound functional changes, shifting from an organizational framework anchored within somatosensory/motor and visual regions into one that is balanced through interactions with later-maturing aspects of association cortex. Here, we provide evidence linking this profile of large-scale functional reorganization to developmental shifts in attention network connectivity. We demonstrate that changes in functional connectivity between childhood and adolescence are preferential to the salience/ventral attention network. Excluding these network-linked vertices from analysis effectively removes developmental changes in cortical organization, while connectivity within the ventral attention network predicts intellectual ability. Children with low ventral attention network connectivity exhibit adolescent-like topographical profiles, suggesting attentional systems may be critically important for understanding how brain functions are refined across development. These data highlight a core role for the ventral attention network in supporting age-dependent shifts in the macroscale organization of cortex across development.

Introduction

The human brain undergoes a series of staged developmental cascades across childhood and adolescence progressing from unimodal somatosensory/motor and visual regions through the transmodal association cortex territories that support complex cognitive functions[1-3]. Evidence for the scheduled timing of these neurodevelopmental events has emerged across biological scales, from regional profiles of cellular maturation[4], synapse formation and dendritic pruning[5], and intracortical myelination[6] through macro-scale morphological features including folding patterns[7] and associated areal expansion[8]. These processes are imbedded within age dependent anatomical changes across lifespan[9], particularly the prolonged development of association cortex territories. In parallel, substantial progress has been made characterizing the organization[10,11] and spatiotemporal maturation of large-scale functional systems across cortex[12,13]. Here, *in vivo* imaging work strongly supports the development of a hierarchical axis, or gradient, of cortical organization, with association territories anchored at the opposite end of a broad functional spectrum from primary sensory and motor regions[14]. Despite clear evidence for age-dependent shifts in the macroscale organization of the cortex from childhood through adolescence, the manner and extent to which specific functional networks may contribute to the widespread process of cortical maturation remains elusive.

The focused study of discrete functional circuits has provided foundational insights in core maturational processes. For example, discoveries linking hierarchical changes within amygdala- and ventral striatal-medial prefrontal cortex (mPFC) circuitry to the development of emotional and social functioning in adolescence[2,3,15,16]. Yet, the maturational refinement these subcortical-cortical circuits does not occur in isolation. Rather, they are embedded within a broad restructuring of functional systems across the cortical sheet[17]. Here, areal and network boundaries become more clearly defined throughout development[18], while the predominance of local connectivity patterns in childhood gradually gives way to long distance, integrative, connections in adolescence[12,19,20]. This reflects a developmental transition from an

anatomically constrained organizational motif to a topographically distributed system[21]. In children, this complex functional architecture is situated within the unimodal cortex, between somatosensory/motor and visual regions. Conversely, adolescents transition into an adult-like gradient, anchored at one end by unimodal regions supporting primary sensory/motor functions and at the other end by the association cortex[14]. While the organizational profiles of large-scale cortical networks are distinct across childhood and adolescence[22], the extent to which developmental changes within select functional couplings may account for the drastic reorganization in the brain hierarchy remains an open question. By one view, the developmental transition from unimodal though association cortices reflect the coordinated and shared influence of maturational changes across multiple functional systems spanning the entire connectome. An alternative, although not mutually exclusive, possibility is that specific brain networks may play a preferential role in the widespread developmental refinement of cortical connectivity.

Individual cortical parcels are functionally organized along a global gradient that transitions from somato/motor and visual regions at one end and multimodal association cortex at the other[23]. The hierarchical nature of these functional relationships reflects a core feature of brain organization in both adolescents[14] and adults[23]. Incoming sensory information undergoes a process of extensive elaboration and attentional modulation as it cascades into deeper layers of cortical processing. Visual system connectivity, as one example, moves along the dorsal and ventral visual streams, eventually converging within aspects of the dorsal and ventral attention networks including the anterior insula, superior parietal cortex, and operculum parietal before eventually filtering through multimodal convergence zones, particularly the default network[24]. Although speculative, these data suggest a possible preferential role for sensory orienting and attentional systems in the integrity of the information processing hierarchies in the human brain. Intriguingly, there is converging evidence to suggest the staged development of a ventral attention network, encompassing aspects of anterior insula, anterior prefrontal cortex and anterior cingulate cortex[11,25] (see also, cingulo-opercular network[26] and salience

network[27]), that follows the age-dependent shifts in cortical organization across childhood and adolescence[14]. The salience/ventral attentional network, together with frontoparietal network, have been proposed to constitute a dual-network system for the 'top-down' and 'bottom-up' processing necessary for adaptive behavioral responses[26,28,29], supporting the functional propagation of information from primary somato/motor, visual, auditory cortex through the default network[24]. These dissociable attentional and control systems are interconnected in children but later segregate over the course of adolescence to eventually form the parallel architectures that support adaptive behavior in adulthood[18]. This indicates the attention system may play a preferential role in the transformative brain changes occurring throughout childhood and adolescence. As such, characterizing the relationships linking attention network connectivity and age-dependent changes in the macroscale brain organization would provide a tremendous opportunity to understand how the functional architecture of cortex is shaped and sculpted across the human lifespan.

In the present study we examined the extent to which specific functional networks may serve to underpin the age-dependent maturation of functional gradient patterns across the cortical sheet. To directly address this open question, we first established the cortical territories exhibiting pronounced functional changes in a longitudinal sample of children and adolescents, revealing the preferential presence of developmental shifts within the ventral attention network. Gradient analyses excluding regions exhibiting the maximal developmental change in children resulted in an accelerated maturation of adolescent-like gradient patterns, suggesting ventral attention territories play a core role in the expression of adolescent-like connectivity gradients. Moreover, children with low ventral attention connectivity exhibited an accelerated profile of cortical maturation that closely resembles prior reports in adolescents and adults. Highlighting the importance of attention network connectivity in cognitive functioning, standardized measures of intelligence linked with reduced attention network degree centrality in children and adolescents. Collectively, these data suggest that ventral attention system functioning in childhood and adolescence

may underpin the developmental reorganization and maturation of functional networks across the cortical sheet.

Figure 1. Salience/Ventral attention network areas demonstrate high population-level degree centrality but pronounced functional changes across development. Degree Centrality maps in children (A) and adolescents (B) reveal consistent dense connectivity in ventral attention network areas throughout development. Scale bar reflects the count of above threshold connections from a given vertex to all other vertices. Larger values indicate higher degree centrality. (C) Euclidean Distance of the functional connectome at each vertex between children and adolescents reveals a clear switch within the ventral attention network. Larger values indicate greater dissimilarity. (D) Regions based on the Schaefer et al., 400-parcel[30] atlas and colored by the Yeo et al., 7-network solution[11]. (E) Bar graph reflects changes in the Euclidean Distance of functional connectome at network level (mean network values ± standard error). The ventral attention network shows the largest developmental change while visual network and default networks are most stable between children and adolescents. DorsAttn, dorsal attention; Sal, salience; Som/Mot, somato/motor; and VentAttn, ventral attention.

Results

Ventral attention network areas demonstrate high degree centrality and pronounced shifts across development

Data for this study were provided by the Chinese Color Nest Project (CCNP)[21,31]. Vertex-level functional connectivity (FC) matrices (20,484 × 20,484) were first generated. In line with prior work[10,11,14], only the top 10% connections of each vertex were retained to enforce sparsity. Degree centrality maps for both children (6 to 12 y of age; n=202) and adolescents (12 to 18 years of age; n=176) were generated to characterize the broad organizational properties of the functional connectome across development (Figure 1a,b). Here, degree centrality reflects the count of above threshold connections for a given vertex to all other vertices (see Methods).

The observed profiles of degree centrality were highly similar between children and adolescents (Pearson r=0.947, $p_{spin} \leq 0.001$). Here, significance was established using permuted spin tests, which preserve the spatial autocorrelation structure of effects[32]. Heightened degree centrality values in both children and adolescents were preferentially evident in aspects of the ventral attention network, including portions of anterior insula, medial prefrontal cortex, and anterior cingulate cortex/midline supplementary motor area (Figure 1). Increased degree centrality was also present within default network territories including portions of posterior inferior parietal lobule, posterior cingulate cortex, and precuneus. Additionally, visual system areas including superior and transverse occipital sulcus at the boundary between dorsal and visual network demonstrated high degree centrality values. Conversely, primary somatosensory and motor areas as well as regions within the lateral prefrontal cortex and temporal lobe exhibited relatively low degree centrality. Broadly, these data reflect the presence of dense connectivity within medial and posterior territories along the cortical sheet, while relatively low centrality was evident in lateral prefrontal and somato/motor areas, highlighting a stable pattern of degree centrality across childhood and adolescence.

Degree centrality broadly summarizes profiles of cortical connectivity, to examine developmental changes in strength of functional connectivity at vertex level, we

calculated the associated Euclidean Distance in functional connectivity similarity between children and adolescents (Figure 1c). Despite the presence of consistent population-level patterns of degree centrality, analyses revealed spatially non-uniform shifts in hub regions of functional connectivity across groups. The maximum developmental changes were anchored within the ventral attention network (Figure 1c-e), encompassing aspects of anterior and posterior insula as well as cingulate cortex. One-way ANOVA revealed the presence of between-network differences (F=790.94, df=6, p≤0.001), with increased Euclidean distance in the ventral attention network, relative to other networks (see multiple comparisons results in Supplemental Table 2). Prior work indicates that maturational age broadly follows the theorized hierarchy of cortical information processing[23], with somato/motor and visual networks maturing in childhood, while medial prefrontal aspects of default and limbic networks, peak later during adolescence[14,33]. However, in the present analyses, the default network exhibited relatively less developmental change in Euclidian distance between groups, followed by visual network (Figure 1e). Although speculative, these data suggest the presence of specific network-level similarities in connectivity between children and adolescents that may precede broader age-dependent shifts in the macroscale organization of cortex, highlighting the need for developmental neuroscientists to consider the manner in which individual functional networks (e.g., default and attention) are embedded within the broader functional architecture of the brain.

To assess the importance of attention network connectivity and cognitive functioning, the relationship between standardized measures of intelligence (IQ) and the degree centrality of the ventral attention network were examined using Linear Mixed Effect (LME) model. LME models for IQ subdomain scores in verbal, perceptual reasoning, working memory, processing speed, and a composite total score were constructed separately. Age, gender, head motion and the vertex-level connectivity counts with ventral attention network were fed into each LME model as fixed effects, repeated measurements were set as the random effect. Results revealed significant negative associations between the ventral attention network and

verbal (p≤0.05), perceptual reasoning (p≤0.01) and the composite IQ scores (p≤0.05). Associations with participant age and verbal score (p≤8×10⁻⁶), perceptual reasoning score (p≤0.05) and total score (p≤0.001), as well as associations with gender and verbal score (p≤0.05), perceptual reasoning score (p≤0.05), and total score (p≤0.05) were also observed. No significant associations between IQ scores and head motion were evident (p>0.05; see LME results in Supplemental Tables 3-7 for each IQ subdomain score).

Figure 2: Ventral attention territories play a core role in the expression of adolescent-like connectivity gradients. (A) The principle cortical gradients of connectivity in children, adolescents (data from Dong et al., 2021) and adults (data from Margulies et al., 2016). (B) Clusters with maximum developmental changes in Euclidean Distance were extracted (Figure 1c), denoted as black on the cortical surface. Associated vertices were then dropped from the cortical connectome in the child group prior to rederiving the gradients. Results reveal that default and visual networks anchor the extremes along the principal gradient (Gradient 1), mirroring the principal gradient in both adolescents (r=0.68, p_{spin}≤0.001) and adults (r=0.84, p_{spin}≤0.001). In children, the rederived second gradient (Gradient 2) revealed a unimodal architecture separating somatosensory network from visual network, which closely corresponds on the second gradient in both adolescents (r=0.66, p_{spin}≤0.001) and adults (r=0.85, p_{spin}≤0.001). (C) To further assess the significance of the results, we constructed a permuted null model in the present data. Clusters with same size and shape as (B) but shuffled locations on the cortical sheet were generated and excluded from analyses in child group 500 times. For each shuffle, the principal

gradient was extracted and compared with the principal gradient in adults. The permuted null model shows only 1 case revealing a higher correlation than that observed in the real data, the x-axis indicates the absolute correlation values, the y-axis indicates the frequency of correlations, dotted line refers to the hypothesis being tested.

A core role for the ventral attention network in the macroscale organization of cortex across childhood and adolescence

The transition from childhood to adolescence is marked by pronounced changes in the functional organization of cortex[14,22]. Broadly, this is reflected in the presence of age-dependent transitions across macroscale gradients that extend from unimodal (somato/motor and visual) regions through the cortical association areas that support complex cognition[1,3,15,16](Figure 2a). Next, we examined whether age-dependent alterations in ventral attention network connectivity might partly account for the maturation of the cortical processing hierarchy as reflected in these overlapping organizing axes, or gradients. Brain areas with maximum differences in Euclidean Distance were extracted (See Methods; Figure 2b), and then removed from brain connectivity matrix while we rederived the functional gradients. Here, diffusion map embedding[10,34,35] was used to decompose participant connectivity matrices into a lower dimensional space. The resulting functional components, or gradients, reflect dissociable spatial patterns of cortical connectivity, ordered by the variance explained in the initial functional connectivity matrix[10,14].

Dropping ventral attention areas generates adolescent- and adult-like gradient architectures in children. The removal of ventral attention network regions led to the formation of a primary gradient in children that closely assembles the first gradient in both adolescents (r=0.68, p_{spin}≤0.001) and adults (r=0.84, p_{spin}≤0.001). The rederived second gradient in children most closely assembled the second gradient in both adolescents (r=0.66, p_{spin}≤0.001) and adults (r=0.85, p_{spin}≤0.001). However, while the primary gradient derived from ventral attention network lesioned data in children broadly recapitulated the primary gradient in adults[10], several inconsistencies were observed. Notably, in the lesioned data from the child group, one end of the primary gradient of connectivity was anchored in the visual areas, with the regions at the other end encompassed broad swaths of the association cortex. Prior work in adults has revealed visual territories along with somato/motor and auditory cortex serve to anchor one end of the primary cortical gradient[10]. Additionally, although the second gradient derived from ventral attention network lesioned data in children closely

resembles the second gradient in adults, a muted default network profile can still be observed. While the dropped clusters are not anchored at the extreme end of the primary gradient in children (Figure 2a), it is densely connected and spatial adjacent with somato/motor territories. Although speculative, dropping of ventral attention vertices from the gradient analyses may decrease the number of functional connections attributed to somato/motor network, indirectly shifting its position along the gradient spectrum.

The majority of vertexes from the dropped clusters were from the ventral attention (45.96%) and somato/motor networks (36.57%). Accordingly, the observed transition to an adolescent- and adult-like gradient architecture in children may reflect an artifact, resulting from lesioning the data in a manner that removes of aspects of the unimodal territories that anchor the primary gradient in children (Figure 2a). To address this possibility and identify the primary drivers of adolescent-like profiles of brain function in children we subdivided the dropped clusters into 2 categories: clusters falling within the borders of the somato/motor network and clusters outside the somato/motor network. Gradients were then rederived in the child group to explore the consequences associated with the individual removal of each cluster component. Suggesting that the removal of somato/motor cluster preserves the developmentally typical gradient architecture, in children the first gradient closely resembles the second gradient in adolescents (r=0.94, p_{spin}≤0.001) and adults (r=0.91, p_{spin}≤0.001), while the second gradient closely assemble the first gradient in adolescents (r=0.95, p_{spin}≤0.001) and adults (r=0.91, p_{spin}≤0.001). Conversely, and consistent with the analyses reported above (Figure 2b), dropping the cluster outside somato/motor network led to the generation of an adolescent- and adult-like gradient architecture in children. Here, the first gradient in children matched the first gradient in adolescents (r=0.78, p_{spin}≤0.001), while the second gradient in children resembled the second gradient in adolescents (r=0.78, p_{spin}≤0.001). Namely, dropping the clusters outside somatosensory network in children results in a gradient organization that resembles prior reports in adolescents.

We next examined the extent to which the observed elimination of age-dependent shifts in the macroscale organization of the cortex is specific to the removal of ventral attention network-dominated vertices. Here, we generated 500 null models in the child group with clusters dropped at random locations but with shapes and sizes that match the ventral attention network vertices (reflecting the maximum differences in Euclidean Distance between children and adolescents). For each random model, the first and second gradients of child group were extracted and correlated with the corresponding first and second gradients in adults. Providing evidence for the role of the ventral attention network in the formation of adult-like gradient architectures in children, for the primary gradient the observed correlation was greater than the correlations from the null distribution ($p \leq 0.002$) across 499/500 permutations (Figure 2c). For the second gradient the observed correlation was greater than the correlations from the null distribution ($p \leq 0.001$) across all 500 permutations. While the present analyses are consistent with a core role for the ventral attention network in age-dependent changes in the macroscale organization of the cortex, longitudinal future work should examine possible relationships linking large-scale gradient transitions with shifts in attention network functioning across development.

Figure 3: Developmental shifts in functional connectivity between childhood and adolescence. Differences in network-level connectome functioning between children and adolescents are demonstrated in a Circos plot. For each network, the number connections within the top 10% of each vertex are first summarized at network level, displayed as the links from right half circle with larger radius to left half circle with smaller radius. Reflecting the asymmetrical nature of the thresholded connectivity matrix, links from the left to right half circle indicate the connections within each network that are included in the top 10% of connections to other vertices. Line width highlights the number of connections, with broader lines indicating increased connectivity. Dark colored lines indicate increased values in adolescents, relative to children. Light color lines indicate decrease values in the adolescent group.

Cortical development is linked to a pattern of heightened within-ventral attention network connectivity.

Extending upon the prior Euclidian Distance analyses, connectome-level changes in functional connectivity between children and adolescents are revealed by a Circos plot in Figure 3 and grouped into networks according to Yeo's 7-network solution[11]. These data reveal broad increases in within network connectivity for the ventral attention, somato/motor, and default networks as well as a general flattening of cross-network connectivity across development. The within-network connections for the ventral attention, between-network connections for the dorsal attention, and visual networks were increased, while other between-network connectivities with somatosensory, limbic, frontoparietal and default networks are decreased from childhood to adolescence. Individual developmental changes were examined by linear mixed effect model. Decreased between-network connectivity for the somato/motor network was observed (p≤0.05), while heightened between-network connectivity was evident for the visual (p≤0.05), dorsal attention (p≤0.001), limbic (p≤0.05), and frontoparietal control (p≤0.01) networks. The degree centrality of ventral attention network increased with age in children (p≤0.05), stabilizing in adolescence where no other age-related associations were revealed in within-network connections for ventral attention network.

When considering the other large-scale association networks, the present analyses suggest a pattern where default network connectivity with other functional systems is pruned during development, indicating a bias towards within network connectivity and an associated differentiation from other network processes. A developmental profile that may coincide with the emergence of the default network at the apex of the network hierarchy in adolescence. Conversely, the frontoparietal control network exhibited increased connectivity with dorsal and ventral attention networks in adolescents, suggesting a potential association between cognitive processes and attentional resources allocation during development.

Individual differences in salience/ventral attention network connectivity reveal adult-like gradient architectures in some children

The analyses above provide evidence for a specific relationship between ventral attention network connectivity and the formation of adult-like gradient architectures in children. However, it is not yet clear the extent to which individual variability in attention network functioning may link with the accelerated maturation of gradient architectures across development. To examine this potential relationship, we divided participants into subgroups based on their individual ventral attention network connection counts (Figure 4a). Here, the group split was determined based on the median connection count in the oldest participants (>17 years of age; dotted line in Figure 4a). Participants with fewer connections than this median value were assigned to a low ventral attention group (child n=95 adolescent n=76), all other participants were assigned to a high ventral attention group (child n=107; adolescent n=100). The gradients were then rederived for high and low attention groups across both children and adolescents. No significant associations were observed between the high and low attention groups in age (p=0.45), gender (p=0.82) and head motion (p=0.87).

In adolescents, both the primary (r=0.98, p_{spin}≤0.001) and secondary (r=0.99, p_{spin}≤0.001) gradient architectures were consistent within the low and high ventral attention network participants (Figure 4b,c). For children with high salience and ventral attention connectivity, their functional organization follows the typical pattern of brain maturation (cite; Figure 2a). Here the primary gradient in the high ventral attention children matched the secondary gradient in both adolescents (r=0.89, p_{spin}≤0.001) and adults (r=0.91, p_{spin}≤0.001), while the second gradient in high ventral attention children matched the primary gradient in both adolescents (r=0.89, p_{spin}≤0.001) and adults (r=0.91607, p_{spin}≤0.001). In these children, somato/motor areas were anchored at the opposite extreme from visual regions, revealing a unimodal dominant gradient architecture. However, in children with low ventral attention connectivity we observed a developmentally accelerated pattern of gradient organization that broadly matches the primary and secondary gradients previously

identified in both adolescents (Gradient 1: r=0.93, $p_{spin} \leq 0.001$; Gradient 2: r=0.92, $p_{spin} \leq 0.001$) and adults (Gradient 1: r=0.68, $p_{spin} \leq 0.001$; Gradient 2: r=0.65, $p_{spin} \leq 0.001$).

Collectively, these results demonstrate that individual variability within the attention network in children associates with dissociable motifs of connectivity across the functional connectome. Here, children with low ventral attention connectivity exhibit an accelerated profile of cortical maturation that closely resembles prior reports in adolescents and adults. Although future work is warranted, these data suggest a fundamental relationship between individual variability within the ventral attention network and the age-dependent changes in the macroscale properties of human brain organization.

Figure 4: Individual differences in ventral attention network connectivity reveal accelerated cortical maturation in some children. (A) The number of functional connections linked with ventral attention network at individual level. Male participants are marked in blue, female participants are marked in red. Repeated imaging sessions within the same participants are linked by lines. The x-axis represents age range from 6 to 18 years. The y-axis reflects the number of connections that are linked to the ventral attention network following thresholding. Participants were divided into high and low ventral attention connectivity groups according to the median value in age 17-year-old participants (dotted line). (B) The gradient maps in high ventral attention connectivity groups reveal typical patterns identified in our previous work in both children and adolescents (cite). (C) The gradient maps in low

ventral attention connectivity groups reveal an accelerated maturation process in children, with both the primary and secondary gradients demonstrated a transmodal architectures. Functional organization was broadly preserved across the high and low attention groups in adolescents, with a subtle muting of the ventral attention gradient values in the low attention participant group.

Discussion

A fundamental goal of systems neuroscience is to identify the mechanisms that underlie the scheduled emergence and maturation of the large-scale functional networks central to the information processing capabilities of the human brain. Prior work has demonstrated age-dependent shifts in macroscale organization of cortex[14,17], revealing the presence of a dominant profile of unimodal cortical organization in children that is later replaced in adolescence by a distributed transmodal architecture spanning from unimodal regions through association cortex. While this sweeping developmental reorganization of network relationships across the cortical sheet coincides with the transition from childhood to adolescence, reflecting an inflection point at around age 12 to 14, the neurobiological processes underpinning these profound functional changes have yet to be established. Here, by first identifying the densely connected hub regions across development, we demonstrate that ventral attention network regions[11] encompassing aspects of anterior insula, medial prefrontal cortex, and dorsal anterior cingulate cortex/midline supplementary motor area (also see salience[27] and cingulo-opercular[26] networks) may drive age-dependent shifts in the macroscale organization of cortex across childhood and adolescence.

Although the attention regions identified in the present analyses are recognized as a single system at a coarse scale[36,37], this broad functional architecture is comprised of multiple spatial adjacent but functionally dissociable networks[6,11,27,38]. This complex architecture underpins a cascade of functions supporting information processing, from the control of stimulus driven attention[36] to the initialization of task sets and maintenance of sustained attention during goal pursuit[28]. The hierarchical flow of information from sensory regions to deeper levels of cortical processing is also reflected in the spatial continuity of associated functional parcels along cortical surface, for example, the salience network[25,27], spanning orbital fronto-insular and dorsal anterior cingulate areas through broad posterior areas of insula and dorsal anterior cingulate cortex[25,28]. Broadly, information propagates along posterior to

anterior as well as dorsal to ventral axes across cortex and the associated organization of functionally linked parcels is reflected through the presence of continuous functional gradients[10,14]. Here, the ventral attention network is situated at an intermediate position along this functional spectrum transiting from primary sensory/motor networks to the default network that anchors association cortex. Converging evidence for this hierarchical functional motif has been revealed through analyses of step-wise connectivity. Consistent with classic theories regarding the integration of perceptual modalities into deeper layers of cortical processing[23], functional relationships spread from primary somato/motor, visual and auditory cortex before converging within ventral attention territories, and eventually the default network[24]. As a whole, these data provide clear evidence situating the ventral attention network between primary unimodal and association cortices, highlighting a key role for attention systems in the functional propagation of sensory information to the multimodal regions that support higher-order cognitive functions.

The cingulo-opercular/ventral attention network, together with frontoparietal network, have been proposed to constitute a parallel architecture of executive functioning and cognitive control, supporting adaptive goal pursuit and flexible behavioral adjustments[29]. Although the frontoparietal network, encompassing aspects of dorsolateral prefrontal, dorsomedial prefrontal, lateral parietal, and posterior temporal cortices, is functionally dissociable from cingulo-opercular/ventral attention network in adulthood, studies in developmental populations have revealed that they are functionally linked prior to adolescence[18]. Moreover, in children, the ventral attention network possesses a broad distributed connectivity profile with reduced segregation from salience networks[39], perhaps reflecting fluid community boundaries in those areas. Anterior prefrontal cortex is shared by the two networks in children but later segregated into ventral attention network in early adolescence, with dorsal anterior cingulate cortex incorporated into frontoparietal network in adulthood[18]. This aspect of network development is consistent with evidence for increased, but more diffuse, patterns of task-evoked activity in children relative to adults[40], reflecting the

presence of scattered pattern of community assignments in prefrontal territories, particularly those supporting attentional processes[22].

Suggesting a key role for the attentional systems in cognition, our present analyses revealed associations between the reduced ventral attention network connectivity and broad measures of intellectual functioning. The maturation course of large-scale brain network coincides with the emergence of adult-like performance on cognitive tasks (e.g., processing speed, shifting and response inhibition)[41,42]. A pattern of attention network development that is theorized to reflect a transition from the over-representation of bottom-up attention to greater top-down attentional and executive functioning capacities across development[39]. Converging evidence indicates that ventral attention network functions are intricately linked with the frontoparietal network in both anatomy and function. Although speculative, the corresponding segregation and integration of associated processes likely underpins the maturation of adaptive human cognition. Here, the observed maturational changes were not uniformly distributed across cortical surface, suggesting a key role for the ventral attention network during development. Intriguingly, the greatest functional changes were not evident within the default or primary sensory territories that anchor the functional gradients in adolescence and adulthood. Rather, ventral attention and frontoparietal networks reflected the predominate sources of functional variation across childhood and adolescence (Figure 1E). Of note, developmental changes were not exclusive to attentional network regions, and the somato/sensory network also demonstrated age related variations across childhood and adolescence. These data are consistent with prior reports suggesting that somato/sensory and ventral attention networks are functionally coupled in children, reflecting a distinct community architecture in children relative to adults[22]. Even in adulthood, the ventral areas in somato/sensory network can be identified as a separate cluster from dorsal areas as network parcellations increase in their granularity (i.e., 7-network relative to 17-network resolutions in Yeo et al., 2011[11]; or in Gordon et al., 2017[38]).

To further characterize how changes in ventral attention network connectivity might underlie the broad functional maturation of cortex, we directly excluded the

developmentally dissociable aspects of ventral attention network from the brain connectome and rederived the functional gradients previously characterized in children, adolescence[14] and adults[10]. The present analyses revealed that dropping ventral attention clusters characterized by the greatest developmental changes generates adolescent- and adult-like gradient architectures in children, a profile that was specific to attention network territories. These data suggest a possible role for the ventral attention network in the staged maturation of cortical functioning. Although follow-up work is necessary, the present analyses are consistent with the presence of excessive local connections between ventral attention network and adjacent somato/motor areas in children, biasing the presence of a functional motif dominated by unimodal organization in cortical hierarchy. Later, diffuse connections across broad areas of association cortex emerge with age, until eventually the presence of interactions across distinct systems becomes the dominant feature of cortical organization. The increased emphasis on such transmodal integration is consistent with the developmental principle that local connections are replaced by remote distributed interactions with age[12].

Individual level analysis further reveals the particular influence of ventral attention network on the formation of adult-like functional organization. Children with fewer ventral attention network connections exhibited an accelerated transmodal profile of cortical organization, while the typical unimodal dominated architecture was presented in children with more ventral attention connections. However, an open unanswered question remains the manner through which ventral attention network functioning may reshape the macroscale organization of cortex. Although speculative, this might reflect a consequence of repeated co-activation between ventral attention network regions and associated cortical systems[18]. Accordingly, the dense associations with somato/motor areas[22] as well as frontoparietal areas[18] in children indicates ventral attention network is preferentially connected by both primary and association cortex during development, which is also evident in our degree centrality map (Figure 1A and B). While a dense profile of connections may ensure system integrity, excessive or redundant connections are pruned across

development in order to optimize efficient information processing as neurons that are not fully integrated within local circuits are eliminated to ensure stable network function[43]. The segregation and integration of large-scale networks underpins functional specialization as well as computational efficiency[20,44]. The present data suggest that the ventral attention system, which is both activated by external stimuli (i.e., salience network[27]) as well as the execution of cognitive control (i.e., cingulo-opercular network[28]), may play a preferential role in the broader optimization of network functions across cortex. Future work should focus on characterizing the specific properties of ventral attention network across development. For instance, there is evidence for the lateralization of functional gradients that may coincide with attention system lateralization. However, the extent to which this may reflect a global feature of brain organization attributable to functional variability within the attention system remains an open question.

Accumulating evidence has revealed a dramatic restructuring in the macroscale organization of the cortex across development and suggested the scheduled maturation of functional gradient patterns may be critically important for understanding how cognitive and behavioral capabilities are refined across development. However, the underlying developmental processes driving these functional changes remain to be established. Here, by first localizing the significant maturational changes in functional connectome across childhood and adolescence, we demonstrate that the ventral attention network plays a critical role in onset of the age-dependent shifts that characterize the macroscale organization of cortex across development. Children with fewer functional connections within the ventral attention network exhibited an accelerated maturation of gradient architecture and higher IQ scores. Although the process of brain maturation emerges from a host complex interaction of biological systems that span genes and molecules through cells, networks, and behavior, the current findings suggest a core role for attention network-linked territories. The manner though which the development of ventral attention network functioning interacts with the complex developmental relationships

that span multiple biological systems and link with associated with behaviors across health and disease, remains an open question to be answered in future work.

Methods

Dataset. Participants are from the Chinese Color Nest Project (CCNP), a five-year accelerated longitudinal study across the human life span[21,31,45]. A total of 176 adolescents and 202 typically developing children were included in the analysis, details of the dataset and exclusion criteria can be found in our previous work[14]. All MRI data was obtained with a Siemens Trio 3.0T scanner at the Faculty of Psychology, Southwest University in Chongqing. The reported experiments were approved by the Institutional Review Board from Institute of Psychology, Chinese Academy of Sciences. All participants and their parents/guardians provided written informed consent before participating in the study.

MRI data preprocessing. Anatomical T1 images were visually inspected to exclude individuals with substantial head motion and structural abnormalities. Next, T1 images were fed into the volBrain pipeline (http://volbrain.upv.es)[46] for noise removal, bias correction, intensity normalization and brain extraction. All brain extractions underwent visual inspection to ensure tissue integrity. After initial quality checks, T1 images were passed into the Connectome Computation System (CCS)[47] for surface-based analyses. CCS pipeline is designed for preprocessing multimodal MRI datasets and integrates publicly available software including SPM[48], FSL[49], AFNI[50] and FreeSurfer[51]. Resting-state fMRI data preprocessing included a series of steps common to intrinsic functional connectivity analyses: (1) dropping the first 10s (4 TRs) for the equilibrium of the magnetic field; (2) head motion correction; (3) slicing time correction; (4) time series de-spiking; (5) estimating head motion parameters; (6) registering functional images to high resolution T1 images using boundary-based registration; (7) removing nuisance factors such as head motion, CSF and white matter signals using ICA-AROMA[52]; (8) removing linear and quadratic trends of the time series; (9) projecting volumetric time series to surface space (the *fsaverage5* model with medial wall masked out); (10) 6mm spatial smoothing. All preprocessing scripts are publicly available on github (https://github.com/zuoxinian/CCS). Any resting-state scan with a mean head motion above 0.5 mm was excluded from further

analysis. The demographic information of subjects included in the analyses is listed in Supplemental Table 1.

Degree Centrality Map. Functional connectivity (FC) matrices and the corresponding Fisher-z transformed values were first generated for each resting-scan per visit. Then the two test-retest FC fisher-z matrices within one visit were averaged to increase signal-to-noise ratio for generating individual FCz matrix for each visit, which was later averaged across individuals to form group-level FCz matrices. For the group-level FCz matrices, only the top 10% connections of each vertex were retained, other elements and negative FCs in the matrix were set to 0 to enforce sparsity, yielding an asymmetrical matrix, of which the rows are corresponding to the connectome of each vertex. Degree centrality map was obtained by counting the non-zero elements in each column of the FCz matrix. We then calculated the cosine distance between any two rows of the FCz matrix and subtracted from 1 to obtain a symmetrical similarity matrix, this similarity matrix was later used to derive the gradients.

Euclidean Distance. To characterize the developmental changes between children and adolescents, the Euclidean distance was computed for each row of the cosine distance matrix between children and adolescents. Then most changed clusters were extracted according to the following two criteria: the top 10% in Euclidean distance map and the cluster size above 500 vertexes. One-way ANOVAs were performed to test the statistical differences between networks.

Gradients Analysis. The extracted clusters and their FCs were firstly dropped in the initial FCz matrix of children group, and then cosine similarity matrix was calculated. Diffusion map embedding[10,35] was implemented on the similarity matrix to derive gradients (https://github.com/NeuroanatomyAndConnectivity/gradient_analysis). Pearson correlations were computed between the derived gradients in children and adults.

To examine the statistical significance of the observed Pearson correlations, null distributions were generated by randomly rotated the locations of the extracted clusters across the cortical surface while keeping the shape and size fixed. For each permutation, the gradients and correlations were rederived. A total of 500 permutations were performed to generate the null model.

Circos Plot. Circos Plot was utilized to demonstrated the differences in numbers of functional connectivity between children and adolescents at network level. For each vertex, its connectome was represented as the corresponding row in the FCz matrix. The associated number of network-level connections was obtained by counting the non-0 elements in each network according to the Yeo 7-network solution, generating a matrix with dimension 20484 (number of vertex) by 7 (number of networks). Vertexes were grouped into networks to generate the final network to network FC matrix (7 by 7). The rows of this network-to-network matrix are displayed as the links from right half circus with larger radius to the left smaller half circus, referring to the FCs with networks in its own connectome, the opposite links from left to right circus represent the columns in the network-to-network matrix, referring to the FCs of each network that were existed in other networks' connectome (Figure 3).

High and Low ventral attention network group definition. Ventral attention network linked connections were extracted as the corresponding column in network-to-network matrix, referring to the functional connectivity other networks linked with ventral attention network. Then we take the median value of functional connectivity number in 17-year-old age group. Any single scan with a connectivity number above the threshold was assigned to the "high" ventral attention subgroup, other scans were assigned to the "low" ventral attention group. The gradients were then rederived for each group.

Association analysis with IQ score. Linear Mixed Effect (LME) model was applied to estimate the association between connections number with ventral attention network

and IQ scores. IQ scores were obtained by Wechsler Children Intelligence Scale IV, including scores in following subdomains: verbal, perception reasoning, working memory, processing speed ability. LME was conducted using the following formula:

$$\text{IQ\_score} \sim 1 + \text{Age} + \text{Gender} + (\text{Ventral attention})\_\text{DC} + \text{HeadMotion} + (1 \mid \text{Subid})$$

Here (Ventral attention)_DC refers to the connections with ventral attention network, together with age, gender, head motion and the intercept set as the fixed effect factor. Subid refers to the participant IDs, multi measurements for a single participant were coded as an identical nominal variable, set as the random effect factor. LME models were applied for the total IQ and subdomain scores separately.

Developmental effects on degree centrality of Ventral Attention network. Linear Mixed Effect (LME) model was also applied to estimate the association between connections number with ventral attention network and age, gender head motion. LME was conducted using the following formula:

$$(\text{Ventral attention})\_\text{DC} \sim 1 + \text{Age} + \text{Gender} + \text{HeadMotion} + (1 \mid \text{Subid})$$

References

Casey, B. J., Getz, S. & Galvan, A. The adolescent brain. *Dev Rev* **28**, 62-77, doi:10.1016/j.dr.2007.08.003 (2008).

Casey, B. J., Heller, A. S., Gee, D. G. & Cohen, A. O. Development of the emotional brain. *Neurosci Lett* **693**, 29-34, doi:10.1016/j.neulet.2017.11.055 (2019).

Luna, B. *et al.* Maturation of widely distributed brain function subserves cognitive development. *Neuroimage* **13**, 786-793, doi:10.1006/nimg.2000.0743 (2001).

Kang, H. J. *et al.* Spatio-temporal transcriptome of the human brain. *Nature* **478**, 483-489, doi:10.1038/nature10523 (2011).

Huttenlocher, P. R. & Dabholkar, A. S. Regional differences in synaptogenesis in human cerebral cortex. *J Comp Neurol* **387**, 167-178, doi:Doi 10.1002/(Sici)1096-9861(19971020)387:2<167::Aid-Cne1>3.0.Co;2-Z (1997).

Paquola, C. *et al.* Shifts in myeloarchitecture characterise adolescent development of cortical gradients. *Elife* **8**, doi:10.7554/eLife.50482 (2019).

Zilles, K., Palomero-Gallagher, N. & Amunts, K. Development of cortical folding during evolution and ontogeny. *Trends Neurosci* **36**, 275-284, doi:10.1016/j.tins.2013.01.006 (2013).

Reardon, P. K. *et al.* Normative brain size variation and brain shape diversity in humans. *Science* **360**, 1222-1227, doi:10.1126/science.aar2578 (2018).

Bethlehem, R. A. I. *et al.* Brain charts for the human lifespan. *Nature*, doi:10.1038/s41586-022-04554-y (2022).

Margulies, D. S. *et al.* Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proc Natl Acad Sci U S A* **113**, 12574-12579, doi:10.1073/pnas.1608282113 (2016).

Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol* **106**, 1125-1165, doi:10.1152/jn.00338.2011 (2011).

Fair, D. A. *et al.* Functional Brain Networks Develop from a "Local to Distributed" Organization. *Plos Comput Biol* **5**, doi:10.1371/journal.pcbi.1000381 (2009).

Gao, W. *et al.* Temporal and spatial evolution of brain network topology during the first two years of life. *PLoS One* **6**, e25278, doi:10.1371/journal.pone.0025278 (2011).

Dong, H. M., Margulies, D. S., Zuo, X. N. & Holmes, A. J. Shifting gradients of macroscale cortical organization mark the transition from childhood to adolescence. *Proc Natl Acad Sci U S A* **118**, doi:10.1073/pnas.2024448118 (2021).

Tottenham, N. & Sheridan, M. A. A review of adversity, the amygdala and the hippocampus: a consideration of developmental timing. *Front Hum Neurosci* **3**, 68, doi:10.3389/neuro.09.068.2009 (2009).

Somerville, L. H., Hare, T. & Casey, B. J. Frontostriatal maturation predicts cognitive control failure to appetitive cues in adolescents. *J Cogn Neurosci* **23**, 2123-2134, doi:10.1162/jocn.2010.21572 (2011).

Sydnor, V. J. *et al.* Neurodevelopment of the association cortices: Patterns, mechanisms, and implications for psychopathology. *Neuron* **109**, 2820-2846, doi:10.1016/j.neuron.2021.06.016 (2021).

Fair, D. A. *et al.* Development of distinct control networks through segregation and integration. *P Natl Acad Sci USA* **104**, 13507-13512, doi:10.1073/pnas.0705843104 (2007).

Betzel, R. F. *et al.* Changes in structural and functional connectivity among resting-state networks across the human lifespan. *Neuroimage* **102 Pt 2**, 345-357, doi:10.1016/j.neuroimage.2014.07.067 (2014).

Betzel, R. F. *et al.* Generative models of the human connectome. *Neuroimage* **124**, 1054-1064, doi:10.1016/j.neuroimage.2015.09.041 (2016).

Zuo, X. N. *et al.* Human Connectomics across the Life Span. *Trends Cogn Sci* **21**, 32-45, doi:10.1016/j.tics.2016.10.005 (2017).

Tooley, U. A., Bassett, D. S. & Mackey, A. P. Functional brain network community structure in childhood: Unfinished territories and fuzzy boundaries. *Neuroimage* **247**, 118843, doi:10.1016/j.neuroimage.2021.118843 (2022).

Mesulam, M. M. From sensation to cognition. *Brain* **121 (Pt 6)**, 1013-1052, doi:10.1093/brain/121.6.1013 (1998).

Sepulcre, J., Sabuncu, M. R., Yeo, T. B., Liu, H. & Johnson, K. A. Stepwise connectivity of the modal cortex reveals the multimodal organization of the human brain. *J Neurosci* **32**, 10649-10661, doi:10.1523/JNEUROSCI.0759-12.2012 (2012).

Power, J. D. *et al.* Functional network organization of the human brain. *Neuron* **72**, 665-678, doi:10.1016/j.neuron.2011.09.006 (2011).

Dosenbach, N. U. *et al.* Distinct brain networks for adaptive and stable task control in humans. *Proc Natl Acad Sci U S A* **104**, 11073-11078, doi:10.1073/pnas.0704320104 (2007).

Seeley, W. W. *et al.* Dissociable intrinsic connectivity networks for salience processing and executive control. *J Neurosci* **27**, 2349-2356, doi:10.1523/JNEUROSCI.5587-06.2007 (2007).

Dosenbach, N. U. *et al.* A core system for the implementation of task sets. *Neuron* **50**, 799-812, doi:10.1016/j.neuron.2006.04.031 (2006).

Dosenbach, N. U., Fair, D. A., Cohen, A. L., Schlaggar, B. L. & Petersen, S. E. A dual-networks architecture of top-down control. *Trends Cogn Sci* **12**, 99-105, doi:10.1016/j.tics.2008.01.001 (2008).

Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cereb Cortex* **28**, 3095-3114, doi:10.1093/cercor/bhx179 (2018).

Liu, S. *et al.* Chinese Color Nest Project : An accelerated longitudinal brain-mind cohort. *Dev Cogn Neurosci* **52**, 101020, doi:10.1016/j.dcn.2021.101020 (2021).

Alexander-Bloch, A. F. *et al.* On testing for spatial correspondence between maps of human brain structure and function. *Neuroimage* **178**, 540-551, doi:10.1016/j.neuroimage.2018.05.070 (2018).

Gogtay, N. *et al.* Dynamic mapping of human cortical development during childhood through early adulthood. *P Natl Acad Sci USA* **101**, 8174-8179, doi:DOI 10.1073/pnas.0402680101 (2004).

Coifman, R. R. & Lafon, S. Diffusion maps. *Appl Comput Harmon A* **21**, 5-30, doi:10.1016/j.acha.2006.04.006 (2006).

Langs, G. *et al.* Identifying Shared Brain Networks in Individuals by Decoupling Functional and Anatomical Variability. *Cereb Cortex* **26**, 4004-4014, doi:10.1093/cercor/bhv189 (2016).

Corbetta, M. & Shulman, G. L. Control of goal-directed and stimulus-driven attention in the brain. *Nat Rev Neurosci* **3**, 201-215, doi:10.1038/nrn755 (2002).

Fox, M. D. *et al.* The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proc Natl Acad Sci U S A* **102**, 9673-9678, doi:10.1073/pnas.0504136102 (2005).

Gordon, E. M. *et al.* Precision Functional Mapping of Individual Human Brains. *Neuron* **95**, 791-807 e797, doi:10.1016/j.neuron.2017.07.011 (2017).

Farrant, K. & Uddin, L. Q. Asymmetric development of dorsal and ventral attention networks in the human brain. *Dev Cogn Neurosci* **12**, 165-174, doi:10.1016/j.dcn.2015.02.001 (2015).

Casey, B. J., Giedd, J. N. & Thomas, K. M. Structural and functional brain development and its relation to cognitive development. *Biol Psychol* **54**, 241-257, doi:10.1016/s0301-0511(00)00058-2 (2000).

Luna, B., Garver, K. E., Urban, T. A., Lazar, N. A. & Sweeney, J. A. Maturation of cognitive processes from late childhood to adulthood. *Child Dev* **75**, 1357-1372, doi:10.1111/j.1467-8624.2004.00745.x (2004).

Huizinga, M., Dolan, C. V. & van der Molen, M. W. Age-related change in executive function: developmental trends and a latent variable analysis. *Neuropsychologia* **44**, 2017-2036, doi:10.1016/j.neuropsychologia.2006.01.010 (2006).

Pfisterer, U. & Khodosevich, K. Neuronal survival in the brain: neuron type-specific mechanisms. *Cell Death Dis* **8**, e2643, doi:10.1038/cddis.2017.64 (2017).

Bullmore, E. & Sporns, O. The economy of brain network organization. *Nat Rev Neurosci* **13**, 336-349, doi:10.1038/nrn3214 (2012).

Dong, H. M. *et al.* Charting brain growth in tandem with brain templates at school age. *Science Bulletin* **65**, 1924-1934, doi:10.1016/j.scib.2020.07.027 (2020).

Manjon, J. V. & Coupe, P. volBrain: An Online MRI Brain Volumetry System. *Front Neuroinform* **10**, 30, doi:10.3389/fninf.2016.00030 (2016).

Xu, T., Yang, Z., Jiang, L. L., Xing, X. X. & Zuo, X. N. A Connectome Computation System for discovery science of brain. *Science Bulletin* **60**, 86-95, doi:10.1007/s11434-014-0698-3 (2015).

Friston, K. J. *et al.* Statistical parametric maps in functional imaging: a general linear approach. *Human brain mapping* **2**, 189-210 (1994).

Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W. & Smith, S. M. Fsl. *Neuroimage* **62**, 782-790 (2012).

Cox, R. W. AFNI: Software for analysis and visualization of functional magnetic resonance neuroimages. *Comput Biomed Res* **29**, 162-173 (1996).

Fischl, B. FreeSurfer. *Neuroimage* **62**, 774-781, doi:10.1016/j.neuroimage.2012.01.021 (2012).

Pruim, R. H. R. *et al.* ICA-AROMA: A robust ICA-based strategy for removing motion artifacts from fMRI data. *Neuroimage* **112**, 267-277, doi:10.1016/j.neuroimage.2015.02.064 (2015).