Regulation of Store-Operated Ca^{2+} Entry by IP$_3$ Receptors Independent of Their Ability to Release Ca^{2+}

Pragnya Chakraborty[1,2], Bipan Kumar Deb[1,3], Colin W. Taylor[4,*] and Gaiti Hasan[1,5,*]

[1]National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bellary Road, Bangalore, 560065, India.

[2]SASTRA University, Thanjavur, Tamil Nadu, 613401, India.

[3]Present address: Department of Molecular and Cell Biology, University of California, Berkeley, CA, 94720, USA.

[4]Department of Pharmacology, University of Cambridge, Cambridge, CB2 1PD, UK.

[5]Lead Contact

[*]Correspondence: cwt1000@cam.ac.uk, gaiti@ncbs.res.in

ABSTRACT

Loss of endoplasmic reticular (ER) Ca^{2+} activates store-operated Ca^{2+} entry (SOCE)

by causing the ER localized Ca^{2+} sensor STIM to unfurl domains that activate Orai

channels in the plasma membrane at membrane contact sites (MCS). Here we

demonstrate a novel mechanism by which the inositol 1,4,5 trisphosphate receptor

(IP_3R), an ER-localized IP_3-gated Ca^{2+} channel, regulates neuronal SOCE. In human

neurons, SOCE evoked by pharmacological depletion of ER-Ca^{2+} is attenuated by

loss of IP_3Rs, and restored by expression of IP_3Rs even when they cannot release

Ca^{2+}, but only if the IP_3Rs can bind IP_3. Imaging studies demonstrate that IP_3Rs

enhance association of STIM1 with Orai1 in neuronal cells with empty stores; this

requires an IP_3-binding site, but not a pore. Convergent regulation by IP_3Rs, may tune

neuronal SOCE to respond selectively to receptors that generate IP_3.

Keywords: Ca^{2+} signaling, extended synaptotagmin, human neural progenitor cell, IP_3

receptor, membrane contact site, neuron, Orai, proximity ligation assay, STIM, store-

operated Ca^{2+} entry

INTRODUCTION

The activities of all eukaryotic cells are regulated by increases in cytosolic free Ca^{2+} concentration ($[Ca^{2+}]_c$), which are almost invariably evoked by the opening of Ca^{2+}-permeable ion channels in biological membranes. The presence of these Ca^{2+} channels within the plasma membrane (PM) and the membranes of intracellular Ca^{2+} stores, most notably the endoplasmic reticulum (ER), allows cells to use both intracellular and extracellular sources of Ca^{2+} to evoke Ca^{2+} signals. In animal cells, the most widely expressed Ca^{2+} signaling sequence links extracellular stimuli, through their specific receptors and activation of phospholipase C, to formation of inositol 1,4,5-trisphosphate (IP_3), which then stimulates Ca^{2+} release from the ER through IP_3 receptors (IP_3R) (Foskett *et al.*, 2007; Prole and Taylor, 2019). IP_3Rs occupy a central role in Ca^{2+} signaling by releasing Ca^{2+} from the ER. IP_3Rs thereby elicit cytosolic Ca^{2+} signals, and by depleting the ER of Ca^{2+} they initiate a sequence that leads to activation of store-operated Ca^{2+} entry (SOCE) across the PM (Putney, 1986; Thillaiappan *et al.*, 2019). SOCE occurs when loss of Ca^{2+} from the ER causes Ca^{2+} to dissociate from the luminal Ca^{2+}-binding sites of an integral ER protein, stromal interaction molecule 1 (STIM1). STIM1 then unfolds its cytosolic domains to expose a region that binds directly to a Ca^{2+} channel within the PM, Orai, causing it to open and Ca^{2+} to flow into the cell across the PM (Parekh and Putney, 2005; Prakriya and Lewis, 2015; Lewis, 2020). The interactions between STIM1 and Orai occur across a narrow gap between the ER and PM, a membrane contact site (MCS), where STIM1 puncta trap Orai channels. While STIM1 and Orai are undoubtedly the core components of SOCE, many additional proteins modulate their interactions (Rosado,

Jenner and Sage, 2000; Palty *et al.*, 2012; Deb, Pathak and Hasan, 2016; Srivats *et*

al., 2016) and other proteins contribute by regulating the assembly of MCS (Chang *et*

al., 2013; Giordano *et al.*, 2013; Kang *et al.*, 2019).

It is accepted that IP_3-evoked Ca^{2+} release from the ER through IP_3Rs is the usual

means by which extracellular stimuli evoke SOCE. Here, the role of the IP_3R is widely

assumed to be restricted to its ability to mediate Ca^{2+} release from the ER and thereby

activate STIM1. Evidence from *Drosophila*, where we suggested an additional role for

IP_3Rs in regulating SOCE (Agrawal et al., 2010; Chakraborty et al., 2016), motivated

the present study, wherein we examined the contribution of IP_3Rs to SOCE in

mammalian neurons. We show that in addition to their ability to activate STIM1 by

evoking ER Ca^{2+} release, IP_3Rs also facilitate interactions between active STIM1 and

Orai1. This additional role for IP_3Rs, which is regulated by IP_3 but does not require a

functional pore, reveals an unexpected link between IP_3, IP_3Rs and Ca^{2+} signaling that

is not mediated by IP_3-evoked Ca^{2+} release. We speculate that dual regulation of

SOCE by IP_3Rs may allow Ca^{2+} release evoked by IP_3 to be preferentially coupled to

SOCE.

RESULTS

Loss of IP_3R1 Attenuates SOCE in Human Neural Stem Cells and Neurons

We investigated the effects of IP_3Rs on SOCE by measuring $[Ca^{2+}]_c$ in human neural

stem cells and neurons prepared from embryonic stem cells. Human neural progenitor

cells (hNPCs) were derived from H9 embryonic stem cells using small molecules that

mimic cues provided during human brain development (Gopurappilly *et al.*, 2018). We

confirmed that hNPCs express canonical markers of neural stem cells (**Figure 1A**)

and that IP_3R1 is the predominant IP_3R subtype (GEO accession no. GSE109111)

(Gopurappilly *et al.*, 2018). An inducible lentiviral shRNA-miR construct targeting

IP_3R1 reduced IP_3R1 expression by $93 \pm 0.4\%$ relative to a non-silencing (NS)

construct (**Figures 1B and 1C**). Carbachol stimulates muscarinic acetylcholine

receptors, which are expressed at low levels in hNPCs (Gopurappilly *et al.*, 2018). In

Ca^{2+}-free medium, carbachol evoked an increase in $[Ca^{2+}]_c$ in about 10% of hNPCs,

consistent with it stimulating Ca^{2+} release from the ER through IP_3Rs. Restoration of

extracellular Ca^{2+} then evoked an increase in $[Ca^{2+}]_c$ in all cells that responded to

carbachol. Both carbachol-evoked Ca^{2+} release and SOCE were abolished in hNPCs

expressing IP_3R1-shRNA, confirming the effectiveness of the IP_3R1 knockdown

(**Figure 1- figure supplement 1A-C**).

Thapsigargin, a selective and irreversible inhibitor of the ER Ca^{2+} pump

(sarcoplasmic/endoplasmic reticulum Ca^{2+}-ATPase, SERCA), was used to deplete the

ER of Ca^{2+} and thereby activate SOCE (**Figure 1D**) (Parekh and Putney, 2005).

Restoration of extracellular Ca^{2+} to thapsigargin-treated hNPCs evoked a large

increase in $[Ca^{2+}]_c$, reflecting the activity of SOCE (**Figure 1D**). The maximal

amplitude and rate of SOCE were significantly reduced in cells lacking IP_3R1, but the

resting $[Ca^{2+}]_c$ and thapsigargin-evoked Ca^{2+} release were unaffected (**Figures 1D-1F**

and **Figure 1- figure supplement 1D and 1E**). STIM1 and Orai1 expression were

also unaltered in hNPC lacking IP_3R1 (**Figure 1- figure supplement 1G**). After

spontaneous differentiation of hNPC, cells expressed markers typical of mature

neurons, and the cells responded to depolarization with an increase in $[Ca^{2+}]_c$ (**Figure**

1- figure supplement 1F and Figure 1- figure supplement 1H-J). Thapsigargin

evoked SOCE in these differentiated neurons; and expression of IP_3R1-shRNA

significantly reduced the SOCE response without affecting depolarization-evoked Ca^{2+}

signals (**Figures 1H-1J** and **Figure 1- figure supplement 1H-1L**).

Loss of IP_3R1 Attenuates SOCE in Human Neuroblastoma Cells

IP_3Rs link physiological stimuli that evoke Ca^{2+} release from the ER to SOCE, but the

contribution of IP_3Rs is thought to be limited to their ability to deplete the ER of Ca^{2+}.

We have reported that in *Drosophila* neurons there is an additional requirement for

IP_3Rs independent of ER Ca^{2+} release (Venkiteswaran and Hasan, 2009; Agrawal *et

al.*, 2010; Chakraborty *et al.*, 2016). Our results with hNPCs and stem cell-derived

neurons suggest a similar requirement for IP_3Rs in regulating SOCE in mammalian

neurons. To explore the mechanisms underlying this additional role for IP_3Rs, we

turned to a more tractable cell line, SH-SY5Y cells. These cells are derived from a

human neuroblastoma; they exhibit many neuronal characteristics (Agholme *et al.*,

2010); they express M3 muscarinic acetylcholine receptors that evoke IP_3-mediated

Ca^{2+} release and SOCE (Grudt, Usowicz and Henderson, 1996); and they express

predominantly IP_3R1 (Wojcikiewicz, 1995; Tovey *et al.*, 2001), with detectable IP_3R3,

but no IP_3R2 (**Figure 2A**). We used inducible expression of IP_3R1-shRNA to

significantly reduce IP_3R1 expression (by $74 \pm 1.2\%$), without affecting IP_3R3 (**Figures

2A and 2B**). As expected, carbachol-evoked Ca^{2+} signals in individual SH-SY5Y cells

were heterogenous and the carbachol-evoked Ca^{2+} release was significantly reduced

by knockdown of IP_3R1 (**Figures 2C and 2D** and **Figure 2 - figure supplement 1A**

and 1B). Thapsigargin evoked SOCE in SH-SY5Y cells (Grudt, Usowicz and

Henderson, 1996), and it was significantly attenuated after knockdown of IP$_3$R1

without affecting resting $[Ca^{2+}]_c$, the Ca^{2+} release evoked by thapsigargin or

expression of STIM1 and Orai1 (**Figures 2E-2G** and **Figure 2 - figure supplement**

1C-1E).

We also used CRISPR/Cas9n and Cas9 to disrupt one or both copies of the

IP$_3$R1 gene, subsequently referred to as IKO (one copy knockout) and IKO null (both

copies knocked out) in SH-SY5Y cells. IP$_3$R1 expression was absent in the IKO null

(**Figure 2 - figure supplement 1F**) whereas expression of STIM1, STIM2 and Orai1

were unperturbed (**Figure 2 - figure supplement 1G**). Carbachol-evoked Ca^{2+}

release and thapisgargin-evoked SOCE were significantly reduced (**Figure 2 - figure**

supplement 1H-1J). Since the IKO null cells were fragile and grew slowly, we

examined SOCE in SH-SY5Y cells with disruption of one copy of the IP$_3$R1 gene. In

the IKO cells, IP$_3$R1 expression, carbachol-evoked Ca^{2+} signals and thapsigargin-

evoked SOCE were all reduced. A minimal effect on thapsigargin-evoked store Ca^{2+}

release was observed in IKO null and IKO cells (**Figure 2 - figure supplement 1J**

and 1Q).

These observations, which replicate those from hNPCs and neurons (**Figure 1**),

vindicate our use of SH-SY5Y cells to explore the mechanisms linking IP$_3$Rs to SOCE

in human neurons.

Expression of IP$_3$R1 or IP$_3$R3 in SH-SY5Y cells expressing IP$_3$R1-shRNA

restored both carbachol-evoked Ca^{2+} release and thapsigargin-evoked SOCE without

affecting resting $[Ca^{2+}]_c$ or thapsigargin-evoked Ca^{2+} release (**Figures 2H-2J** and

Figure 2 - figure supplement 2A-2D). Over-expression of STIM1 in cells expressing

NS-shRNA had no effect on SOCE (**Figure 2 - figure supplement 2E and 2F**), but it

restored thapsigargin-evoked SOCE in cells expressing IP$_3$R1-shRNA, without

affecting resting $[Ca^{2+}]_c$ or thapsigargin-evoked Ca^{2+} release (**Figures 2K-2M**). We

conclude that IP$_3$Rs are required for optimal SOCE, but they are not essential

because additional STIM1 can replace the need for IP$_3$Rs (**Figure 3A**).

It has been reported that SOCE is unaffected by loss of IP$_3$R in non-neuronal

cells (Ma *et al.*, 2001; Chakraborty *et al.*, 2016). Consistent with these observations,

the SOCE evoked in HEK cells by stores emptied fully by treatment with thapsigargin

was unaffected by expression of IP$_3$R1 shRNA (**Figure 2 - figure supplement 3A-3C**)

or by knockout of all three IP$_3$R subtypes using CRISPR/cas9 (HEK-TKO cells)

(**Figure 2 - figure supplement 3D and 3E**). The association of STIM1 with Orai1 in

wild type HEK cells and HEK TKO cells after thapsigargin-evoked store depletion also

appeared identical as tested by a proximity ligation assay (PLA, described further in

Figure 5) (**Figure 2 - figure supplement 3F**). Neuronal and non-neuronal cells may,

therefore, differ in the contribution of IP$_3$R to SOCE. We return to this point later.

Binding of IP$_3$ to IP$_3$R Without a Functional Pore Stimulates SOCE

IP$_3$Rs are large tetrameric channels that open when they bind IP$_3$ and Ca^{2+}, but they

also associate with many other proteins (Prole and Taylor, 2019), and many IP$_3$Rs

within cells appear not to release Ca^{2+} (Thillaiappan *et al.*, 2019). A point mutation

(D2550A, IP$_3$R1$^{D/A}$) within the IP$_3$R1 pore prevents it from conducting Ca^{2+} (Dellis *et*

al., 2008). As expected, expression of IP$_3$R1$^{D/A}$ in cells lacking IP$_3$R1 failed to rescue

carbachol-evoked Ca^{2+} release, but it unexpectedly restored thapsigargin-evoked

SOCE (**Figures 3B-3D** and **Figure 3 - figure supplement 1A-E**). We confirmed that

rescue of thapsigargin-evoked Ca^{2+} entry by this pore-dead IP_3R was mediated by a

conventional SOCE pathway by demonstrating that it was substantially attenuated by

siRNA-mediated knockdown of Orai1 (**Figures 3C and 3D** and **Figure 3 - figure**

supplement 1F-1H).

Activation of IP_3Rs is initiated by IP_3 binding to the N-terminal IP_3-binding core of

each IP_3R subunit (Prole and Taylor, 2019). Mutation of two conserved phosphate-

coordinating residues in the α-domain of the binding core (R568Q and K569Q of

IP_3R1, $IP_3R1^{RQ/KQ}$) almost abolishes IP_3 binding (Yoshikawa *et al.*, 1996; Iwai *et al.*,

2007), while mutation of a single residue (R568Q, IP_3R1^{RQ}) reduces the IP_3 affinity by

~10-fold (Dellis *et al.*, 2008). Expression of rat $IP_3R1^{RQ/KQ}$ rescued neither carbachol-

evoked Ca^{2+} release nor thapsigargin-evoked SOCE in cells lacking IP_3R1 (**Figures**

3E and 3F and **Figure 3 - figure supplement 1C)**. However, expression of IP_3R1^{RQ}

substantially rescued thapsigargin-evoked SOCE (**Figures 3E and 3F** and **Figure 3 -**

figure supplement 1I and 1J). Expression of an N-terminal fragment of rat IP_3R

(IP_3R1^{1-604}), to which IP_3 binds normally (Iwai *et al.*, 2007), failed to rescue

thapsigargin-evoked SOCE (**Figure 3 - figure supplement 1K and 1L)**. These results

establish that a functional IP_3-binding site within a full-length IP_3R is required for IP_3Rs

to facilitate thapsigargin-evoked SOCE. Hence in cells with empty Ca^{2+} stores, IP_3

binding, but not pore-opening, is required for regulation of SOCE by IP_3Rs. In cells

stimulated only with thapsigargin and expressing IP_3Rs with deficient IP_3 binding,

basal levels of IP_3 are probably insufficient to meet this need.

We further examined the need for IP_3 by partially depleting the ER of Ca^{2+} using

cyclopiazonic acid (CPA), a reversible inhibitor of SERCA, to allow submaximal

activation of SOCE (**Figure 3 - figure supplement 1M and 1N**). Under these

conditions, addition of carbachol in Ca^{2+}-free HBSS to cells expressing IP_3R1-shRNA

caused a small increase in $[Ca^{2+}]_c$ (**Figures 4A-4C**). In the same cells expressing

IP_3R1^{DA}, the carbachol-evoked Ca^{2+} release was indistinguishable from that observed

in cells without IP_3R^{DA} (**Figures 4B and 4C**), indicating that the small response was

entirely mediated by residual native IP_3R1 and/or IP_3R3. Hence, the experiment allows

carbachol to stimulate IP_3 production in cells expressing IP_3R1^{DA} without causing

additional Ca^{2+} release. The key result is that in cells expressing IP_3R1^{DA}, carbachol

substantially increased SOCE (**Figures 4A-4C**). We conclude that IP_3, through IP_3Rs,

regulates coupling of empty stores to SOCE. This is the first example of an IP_3R

mediating a response to IP_3 that does not require the pore of the channel.

G-protein-coupled receptors are linked to IP_3 formation through the G-protein

Gq, which stimulates phospholipase C β (PLC β). We used YM-254890 to inhibit Gq

(Kostenis, Pfeil and Annala, 2020; Patt *et al.*, 2021). As expected, addition of YM-

254890 to wild type (WT) or NS-shRNA transfected SH-SY5Y cells abolished the Ca^{2+}

signals evoked by carbachol **(Figure 4 - figure supplement 1B)**, but it also reduced

the maximal amplitude and rate of thapsigargin-evoked SOCE (**Figures 4D- 4E** and

Figure 3 - figure supplement 1O). YM-254890 had no effect on the residual

thapsigargin-evoked SOCE in SH-SY5Y cells expressing IP_3R1-shRNA **(Figure 4F**

and **Figure 3 - figure supplement 1O**). The latter result is important because it

demonstrates that the inhibition of SOCE in cells with functional IP$_3$Rs is not an off-

target effect causing a direct inhibition of SOCE.

In wild type or HEK-TKO cells, YM-254890 had no effect on thapsigargin-evoked

SOCE, but it did inhibit SOCE in HEK cells lacking IP$_3$R1 (**Figures 4G-4I** and **Figure 4**

- figure supplement 1C-1G). These results suggest that in HEK cells, which

normally express all three IP$_3$R subtypes (Mataragka & Taylor, 2018), neither loss of

IP$_3$R1 nor inhibition of Gαq is sufficient on its own to inhibit thapsigargin-evoked

SOCE, but when combined there is a synergistic loss of SOCE.

IP$_3$Rs Promote Interaction of STIM1 With Orai1 Within MCS

Our evidence that IP$_3$Rs intercept coupling between empty stores and SOCE (**Figure**

3A) prompted us to investigate the coupling of STIM1 with Orai1 across the narrow

junctions between ER and PM (Carrasco and Meyer, 2011). An *in situ* proximity

ligation assay (PLA) is well suited to analyzing this interaction because it provides a

signal when two immunolabelled proteins are within ~40 nm of each other (Derangère

et al., 2016), a distance comparable to the dimensions of the junctions wherein STIM1

and Orai1 interact (Poteser *et al.*, 2016). We confirmed the specificity of the PLA and

demonstrated that it reports increased association of STIM1 with Orai1 after treating

SH-SY5Y cells with thapsigargin (**Figure 5A** and **Figure 5 - figure supplement 1A-**

1F). In cells expressing IP$_3$R1-shRNA, thapsigargin had no effect on the STIM1-Orai1

interaction reported by PLA, but the interaction was rescued by expression of IP$_3$R1 or

IP$_3$R1^{DA}. There was no rescue with IP$_3$R1$^{RQ/KQ}$ (**Figures 5B-5E**). WT SH-SY5Y cells

that were depleted of basal IP$_3$ by treatment with the Gq inhibitor YM-254890,

showed significantly reduced STIM1-Orai1 interaction after thapsigargin-evoked

depletion of Ca^{2+} stores (**Figure 4 - figure supplement 1A**). The results with PLA

exactly mirror those from functional analyses (**Figures 1** to **4**), suggesting that IP_3

binding to IP_3R enhances SOCE by facilitating interaction of STIM1 with Orai1 (**Figure**

3A).

Extended synaptotagmins (E-Syts) are ER proteins that stabilize ER-PM

junctions including STIM1-Orai1 MCS (Maléth *et al.*, 2014; Kang *et al.*, 2019; Woo *et*

al., 2020). Over-expression of E-Syt1 in SH-SY5Y cells expressing IP3R1-shRNA

rescued thapsigargin-evoked Ca^{2+} entry without affecting resting $[Ca^{2+}]_c$ or

thapsigargin-evoked Ca^{2+} release (**Figures 6A-6C**). The rescued Ca^{2+} entry is likely to

be mediated by conventional SOCE because it was substantially attenuated by

knockdown of STIM1 (**Figures 6D-6F)**. Over-expression of E-Syt1 had no effect on

SOCE in cells with unperturbed IP_3Rs (**Figures 6G-6I**). These results suggest that

attenuated SOCE after loss of IP_3Rs can be restored by exaggerating ER-PM MCS.

DISCUSSION

After identification of STIM1 and Orai1 as core components of SOCE (Prakriya and

Lewis, 2015; Thillaiappan *et al.*, 2019), the sole role of IP_3Rs within the SOCE

pathway was assumed to be the release of ER Ca^{2+} that triggers STIM1 activation.

The assumption is consistent with evidence that thapsigargin-evoked SOCE can occur

in avian (Sugawara *et al.*, 1997; Ma *et al.*, 2002; Chakraborty *et al.*, 2016) and

mammalian cells without IP_3Rs (Prakriya and Lewis, 2001). Although SOCE in

mammalian HEK cells was unaffected by loss of IP$_3$Rs in our study (**Figure 4G and**

4I), it was modestly reduced in other studies of mammalian cells (Bartok *et al.*, 2019;

Yue *et al.*, 2020). However, additional complexity is suggested by evidence that

SOCE may be reduced in cells without IP$_3$Rs (Chakraborty *et al.*, 2016; Bartok *et al.*,

2019; Yue *et al.*, 2020), by observations implicating phospholipase C in SOCE

regulation (Rosado, Jenner and Sage, 2000; Broad *et al.*, 2001), by evidence that

SOCE responds differently to IP$_3$Rs activated by different synthetic ligands (Parekh,

Riley and Potter, 2002) and by some, albeit conflicting reports (Woodard *et al.*, 2010;

Santoso, Cebotaru and Guggino, 2011; Béliveau, Lessard and Guillemette, 2014;

Sampieri *et al.*, 2018; Ahmad *et al.*, 2022), that IP$_3$Rs may interact with STIM and/or

Orai (Woodard *et al.*, 2010; Santoso, Cebotaru and Guggino, 2011; Béliveau, Lessard

and Guillemette, 2014; Sampieri *et al.*, 2018).

We identified two roles for IP$_3$Rs in controlling endogenous SOCE in human

neurons. As widely reported, IP$_3$Rs activate STIM1 by releasing Ca^{2+} from the ER, but

they also, and independent of their ability to release Ca^{2+}, enhance interactions

between active STIM1 and Orai1 (**Figure 7**). The second role for IP$_3$Rs can be

supplanted by over-expressing other components of the SOCE complex, notably

STIM1 or ESyt1 (**Figures 2K-2M** and **Figures 6A and 6B**). It is intriguing that STIM1

(Carrasco and Meyer, 2011; Lewis, 2020), ESyt1 (Giordano *et al.*, 2013) and perhaps

IP$_3$Rs (through the IP$_3$-binding core) interact with phosphatidylinositol 4,5-

bisphosphate (PIP$_2$), which is dynamically associated with SOCE-MCS (Kang *et al.*,

2019). We suggest that the extent to which IP$_3$Rs tune SOCE in different cells is

probably determined by the strength of Gq signaling and endogenous interactions

between STIM1 and Orai1. The latter is likely to depend on the relative expression of

STIM1 and Orai1 (Woo *et al.*, 2020), the STIM isoforms expressed, expression of

proteins that stabilize STIM1-Orai1 interactions (Darbellay *et al.*, 2011; Rana *et al.*,

2015; Rosado *et al.*, 2016; Knapp *et al.*, 2022), and the size and number of the MCS

where STIM1 and Orai1 interact (Kang *et al.*, 2019). The multifarious contributors to

SOCE suggest that cells may differ in whether they express "spare capacity". In

neuronal cells, loss of IP_3 (**Figure 4D**) or of the dominant IP_3R isoform (IP$_3$R1-shRNA;

Figures 1 and 2) is sufficient to unveil the contribution of IP_3R to SOCE, whereas

HEK cells require loss of both IP_3 and IP$_3$R1 to unveil the contribution (**Figures 4H**

and 4I). The persistence of SOCE in cells devoid of IP_3Rs (**Figure 2 - figure**

supplement 3D and 3E) (Prakriya and Lewis, 2001; Ma *et al.*, 2002) probably arises

from adaptive changes within the SOCE pathway. This does not detract from our

conclusion that under physiological conditions, where receptors through IP_3 initiate

SOCE, IP_3Rs actively regulate SOCE.

The IP_3Rs that initiate Ca^{2+} signals reside in ER immediately beneath the PM and

alongside, but not within, the MCS where STIM1 accumulates after store depletion

(Thillaiappan *et al.*, 2017). In migrating cells too, IP_3Rs and STIM1 remain separated

as they redistribute to the leading edge (Okeke *et al.*, 2016). Furthermore, there is

evidence that neither STIM1 nor STIM2 co-immmunoprecipitate with IP$_3$R1 (Ahmad *et*

al., 2022). We suggest, and consistent with evidence that SOCE in cells without IP_3Rs

can be restored by over-expressing E-Syt1 (**Figures 6A-6C**), that IP_3Rs probably

facilitate SOCE by stabilizing the MCS wherein STIM1 and Orai1 interact, rather than

by directly regulating either protein. This proposal provides an analogy with similar

structural roles for IP$_3$Rs in maintaining MCS between ER and mitochondria (Bartok *et*

al., 2019) or lysosomes (Atakpa *et al.*, 2018) (**Figure 7C**). Since both contributions of

IP$_3$Rs to SOCE require IP$_3$ binding (**Figures 3E and 3F**), each is ultimately controlled

by receptors that stimulate IP$_3$ formation (**Figures 4B and 4C**). Convergent regulation

by IP$_3$Rs at two steps in the SOCE pathway may ensure that receptor-regulated PLC

activity provides the most effective stimulus for SOCE; more effective, for example,

than ryanodine receptors, which are also expressed in neurons (**Figure 7B**). By

opening IP$_3$Rs parked alongside SOCE MCS (Thillaiappan *et al.*, 2017; Ahmad *et al.*,

2022), IP$_3$ selectively releases Ca^{2+} from ER that is optimally placed to stimulate

SOCE, and by facilitating Orai1-STIM1 interactions IP$_3$ reinforces this local activation

of SOCE (**Figure 7A and 7B**).

We conclude that IP$_3$-regulated IP$_3$Rs regulate SOCE by mediating Ca^{2+} release

from the ER, thereby activating STIM1 and/or STIM2 (Ahmad *et al.*, 2022) and,

independent of their ability to release Ca^{2+}, IP$_3$Rs facilitate the interactions between

STIM and Orai that activate SOCE. Dual regulation of SOCE by IP$_3$ and IP$_3$Rs allows

robust control by cell-surface receptors and may reinforce local stimulation of Ca^{2+}

entry.

ACKNOWLEDGEMENTS

This research was supported by grants to G.H. from the Dept. of Biotechnology, Govt.

of India (BT/PR6371/COE/34/19/2013) and NCBS-TIFR core support, and to C.W.T.

from the Wellcome Trust (101844) and Biotechnology and Biological Sciences

Research Council (BB/T012986/1). P.C is supported by a DST-INSPIRE fellowship

(DST/INSPIRE Fellowship/2017/IF170360) and she received an Infosys-NCBS travel

award to visit C.W.T.'s lab at Cambridge. We are grateful to Renjitha Gopurappilly

(NCBS, TIFR) for the derivation of human neural precursor cells. We acknowledge

use of the Central Imaging and Flow Cytometry Facility (CIFF), Stem Cell Culture

Facility and Biosafety level-2 laboratory facility at NCBS, TIFR.

DISCLOSURE AND COMPETING INTEREST STATEMENT

The authors declare no competing interests.

REFERENCES

Agholme, L., Lindström, T., Kågedal, K., Marcusson, J. and Hallbeck, M. 2010. An in

vitro model for neuroscience: differentiation of SH-SY5Y cells into cells with

morphological and biochemical characteristics of mature neurons. *J. Alzheimer's*

Dis. 20:1069–1082.

Agrawal, N., Venkiteswaran, G., Sadaf, S., Padmanabhan, N., Banerjee, S. and

Hasan, G. 2010. Inositol 1, 4, 5-trisphosphate receptor and dSTIM function in

Drosophila insulin-producing neurons regulates systemic intracellular calcium

homeostasis and flight. *J. Neurosci.* 30:1301-1313.

Ahmad, M. et al. 2022. Functional communication between IP$_3$R and STIM2 at

subthreshold stimuli is a critical checkpoint for initiation of SOCE. *Proc. Natl.*

Acad. Sci. 119: e2114928118.

Atakpa, P., Thillaiappan, N.B., Mataragka, S., Prole, D.L. and Taylor, C.W. 2018. IP$_3$

receptors preferentially associate with ER-lysosome contact sites and selectively

deliver Ca^{2+} to lysosomes. *Cell Rep.* 25:3180-3193.

Auyeung, V.C., Ulitsky, I., McGeary, S.E. and Bartel, D.P. 2013. Beyond secondary

structure: primary-sequence determinants license pri-miRNA hairpins for

processing. *Cell.* 152:844-858.

Bartok, A., Weaver, D., Golenár, T., Nichtova, Z., Katona, M., Bánsághi, S., Alzayady,

K.J., Thomas, V.K., Ando, H., Mikoshiba, K. and Joseph, S.K. 2019. IP$_3$ receptor

isoforms differently regulate ER-mitochondrial contacts and local calcium

transfer. *Nat. Commun.* 10:1-14.

Béliveau, É., Lessard, V. and Guillemette, G. 2014. STIM1 positively regulates the

Ca^{2+} release activity of the inositol 1,4,5-trisphosphate receptor in bovine aortic

endothelial cells. *PLoS One.* 9: e114718.

Broad, L.M., Braun, F.J., Lievremont, J.P., Bird, G.S.J., Kurosaki, T. and Putney Jr,

J.W. 2001. Role of the phospholipase C-inositol 1, 4, 5-trisphosphate pathway in

calcium release-activated calcium current and capacitative calcium entry. *J. Biol.*

Chem. 276:15945–15952.

Carrasco, S. and Meyer, T. 2011. STIM proteins and the endoplasmic reticulum-

plasma membrane junctions. *Annu. Rev. Biochem.* 80:973–1000.

Chakraborty, S., Deb, B.K., Chorna, T., Konieczny, V., Taylor, C.W. and Hasan, G.

2016. Mutant IP$_3$ receptors attenuate store-operated Ca^{2+} entry by destabilizing

STIM–Orai interactions in *Drosophila* neurons. *J. Cell Sci.* 129:3903–3910.

Chang, C.L., Hsieh, T.S., Yang, T.T., Rothberg, K.G., Azizoglu, D.B., Volk, E., Liao,

J.C. and Liou, J. 2013. Feedback regulation of receptor-induced Ca^{2+} signaling

mediated by E-Syt1 and Nir2 at endoplasmic reticulum-plasma membrane

junctions. *Cell Rep.* 5:813–825.

Darbellay, B., Arnaudeau, S., Bader, C.R., Konig, S. and Bernheim, L. 2011. STIM1L

is a new actin-binding splice variant involved in fast repetitive Ca^{2+} release. *J.*

Cell Biol. 194:335-346.

Deb, B. K., Pathak, T. and Hasan, G. 2016. Store-independent modulation of Ca^{2+}

entry through Orai by Septin 7. *Nat. Commun.* 7:1-15.

Dellis, O., Rossi, A.M., Dedos, S.G. and Taylor, C.W. 2008. Counting functional

inositol 1, 4, 5-trisphosphate receptors into the plasma membrane. *J Biol. Chem.*

283:751-755.

Derangère, V., Bruchard, M., Végran, F. and Ghiringhelli, F. 2016. Proximity Ligation

Assay (PLA) protocol using Duolink® for T cells. *Bio-protocol.* 6:e1811.

Forostyak, O., Romanyuk, N., Verkhratsky, A., Sykova, E. and Dayanithi, G. 2013.

Plasticity of calcium signaling cascades in human embryonic stem cell-derived

neural precursors. *Stem Cells Dev.* 22:1506-1521.

Foskett, J.K., White, C., Cheung, K.H. and Mak, D.O.D. 2007. Inositol trisphosphate

receptor Ca^{2+} release channels. *Physiol. Rev.* 87:593–658.

Giordano, F., Saheki, Y., Idevall-Hagren, O., Colombo, S.F., Pirruccello, M., Milosevic,

I., Gracheva, E.O., Bagriantsev, S.N., Borgese, N. and De Camilli, P. 2013. PI(4,

5) P_2-dependent and Ca^{2+}-regulated ER-PM interactions mediated by the

extended synaptotagmins. *Cell.* 153:1494-1509.

Gopurappilly, R., Deb, B.K., Chakraborty, P. and Hasan, G. 2018. Stable STIM1

knockdown in self-renewing human neural precursors promotes premature

neural differentiation. *Front. Mol. Neurosci.* 11:178.

Grudt, T. J., Usowicz, M. M. and Henderson, G. 1996. Ca^{2+} entry following store

depletion in SH-SY5Y neuroblastoma cells. *Mol. Brain Res.* 36:93–100.

Grynkiewicz, G., Poenie, M. and Tsien, R. Y. 1985. A new generation of Ca^{2+}

indicators with greatly improved fluorescence properties. *J. Biol. Chem.*

260:3440–3450.

Iwai, M., Michikawa, T., Bosanac, I., Ikura, M. and Mikoshiba, K. 2007. Molecular

basis of the isoform-specific ligand-binding affinity of inositol 1, 4, 5-

trisphosphate receptors. *J. Biol. Chem.* 282:12755–12764.

Kang, F., Zhou, M., Huang, X., Fan, J., Wei, L., Boulanger, J., Liu, Z., Salamero, J.,

Liu, Y. and Chen, L. 2019. E-syt1 Re-arranges STIM1 clusters to stabilize ring-

shaped ER-PM contact sites and accelerate Ca^{2+} store replenishment. *Sci Rep.*

9:1-11.

Knapp, M.L., Alansary, D., Poth, V., Förderer, K., Sommer, F., Zimmer, D., Schwarz,

Y., Künzel, N., Kless, A., Machaca, K. and Helms, V., 2022. A longer isoform of

Stim1 is a negative SOCE regulator but increases cAMP-modulated NFAT

signaling. *EMBO reports*, *23:* e53135.

Knott, S.R., Maceli, A.R., Erard, N., Chang, K., Marran, K., Zhou, X., Gordon, A., El

Demerdash, O., Wagenblast, E., Kim, S. and Fellmann, C. 2014. A

computational algorithm to predict shRNA potency. *Mol. Cell.* 56:796-807.

Lewis, R. S. 2020. Store-operated calcium channels: From function to structure and

back again. *Cold Spring Harb. Perspect. Biol.* 12: a035055.

Ma, H. T., Venkatachalam, K., Li, H. S., Montell, C., Kurosaki, T., Patterson, R. L., &

Gill, D. L. 2001. Assessment of the role of the inositol 1,4,5-trisphosphate

receptor in the activation of transient receptor potential channels and store-

operated Ca^{2+} entry channels. *J Biol Chem*, 276: 18888–18896.

Ma, H. T., Venkatachalam, K., Parys, J. B., & Gill, D. L. 2002. Modification of store-

operated channel coupling and inositol trisphosphate receptor function by 2-

aminoethoxydiphenyl borate in DT40 lymphocytes. *J Biol Chem*, 277: 6915–

6922.

Maléth, J., Choi, S., Muallem, S. and Ahuja, M. 2014. Translocation between PI(4,

5)P_2-poor and PI(4, 5)P_2-rich microdomains during store depletion determines

STIM1 conformation and Orai1 gating. *Nat Commun.* 5:1–10.

Mataragka, S. and Taylor, C. W. 2018. All three IP_3 receptor subtypes generate Ca^{2+}

puffs, the universal building blocks of IP_3-evoked Ca^{2+} signals. *J Cell Sci.* 131:

jcs220848.

Miotke, L., Lau, B. T., Rumma, R. T., & Ji, H. P. 2014. High sensitivity detection and

quantitation of DNA copy number and single nucleotide variants with single color

droplet digital PCR. *Anal. Chem.*, *86:* 2618-2624.

Nunes-Hasler, P., Maschalidi, S., Lippens, C., Castelbou, C., Bouvet, S., Guido, D.,

Bermont, F., Bassoy, E.Y., Page, N., Merkler, D. and Hugues, S. 2017. STIM1

promotes migration, phagosomal maturation and antigen cross-presentation in

dendritic cells. *Nat Commun.* 8:1-15.

Okeke, E., Parker, T., Dingsdale, H., Concannon, M., Awais, M., Voronina, S., Molgó,

J., Begg, M., Metcalf, D., Knight, A.E. and Sutton, R. 2016. Epithelial–

mesenchymal transition, IP_3 receptors and ER–PM junctions: Translocation of

Ca^{2+} signalling complexes and regulation of migration. *Biochem. J.* 473:757–767.

Palty, R., Raveh, A., Kaminsky, I., Meller, R. and Reuveny, E. 2012. SARAF

inactivates the store operated calcium entry machinery to prevent excess

calcium refilling. *Cell.* 149:425-438.

Parekh, A. B. and Putney, J. W. 2005. Store-operated calcium channels. *Physiol. Rev.*

85:757–810.

Parekh, A. B., Riley, A. M. and Potter, B. V. L. 2002. Adenophostin A and ribophostin,

but not inositol 1,4,5-trisphosphate or manno-adenophostin, activate the Ca^{2+}

release-activated Ca^{2+} current, ICRAC, in weak intracellular Ca^{2+} buffer.

Biochem. J. 361:133–141.

Patt, J., Alenfelder, J., Pfeil, E.M., Voss, J.H., Merten, N., Eryilmaz, F., Heycke, N.,

Rick, U., Inoue, A., Kehraus, S. and Deupi, X. 2021. An experimental strategy to

probe Gq contribution to signal transduction in living cells. *J. Biol. Chem.*

296:100472.

Poteser, M., Leitinger, G., Pritz, E., Platzer, D., Frischauf, I., Romanin, C. and

Groschner, K. 2016. Live-cell imaging of ER-PM contact architecture by a novel

TIRFM approach reveals extension of junctions in response to store-operated

Ca^{2+}-entry. *Sci. Rep.* 6:1-13.

Prakriya, M. and Lewis, R. S. 2001. Potentiation and inhibition of Ca^{2+} release-

activated Ca^{2+} channels by 2-aminoethyldiphenyl borate (2-APB) occurs

independently of IP$_3$ receptors. *J Physiol.* 536:3–19.

Prakriya, M. and Lewis, R. S. 2015. Store-operated calcium channels pharmacology.

Physiol. Rev. 307:1383–1436.

Prole, D. L. and Taylor, C. W. 2019. Structure and function of IP$_3$ receptors. *Cold*

Spring Harb. Perspect. Biol. 11: a035063.

Putney, J. W. 1986. A model for receptor-regulated calcium entry. *Cell Calcium*. 7:1–

12.

Ran, F. A., Hsu, P. D., Wright, J., Agarwala, V., Scott, D. A., & Zhang, F. 2013.

Genome engineering using the CRISPR-Cas9 system. *Nat. Protoc.* 8: 2281-

2308.

Rana, A., Yen, M., Sadaghiani, A.M., Malmersjö, S., Park, C.Y., Dolmetsch, R.E. and

Lewis, R.S. 2015. Alternative splicing converts STIM2 from an activator to an

inhibitor of store-operated calcium channels. *J Cell Biol.* 209: 653-670.

Reinhardt, P., Glatza, M., Hemmer, K., Tsytsyura, Y., Thiel, C.S., Höing, S., Moritz, S.,

Parga, J.A., Wagner, L., Bruder, J.M. and Wu, G. 2013. Derivation and

expansion using only small molecules of human neural progenitors for

neurodegenerative disease modeling. *PLoS One.* 8: e59252.

Rosado, J.A., Diez, R., Smani, T. and Jardín, I. 2016. STIM and Orai1 variants in

store-operated calcium entry. *Front. Pharmacol.* 6:325.

Rosado, J. A., Jenner, S. and Sage, S. O. 2000. A role for the actin cytoskeleton in

the initiation and maintenance of store-mediated calcium entry in human

platelets. Evidence for conformational coupling. *J Biol. Chem.* 275: 7527–7533.

Saleem, H., Tovey, S.C., Rahman, T., Riley, A.M., Potter, B.V. and Taylor, C.W. 2013.

Stimulation of inositol 1, 4, 5-trisphosphate (IP$_3$) receptor subtypes by analogues

of IP$_3$. *PLoS One.* 8:e54877.

Sampieri, A., Santoyo, K., Asanov, A. and Vaca, L. 2018. Association of the IP$_3$R to

STIM1 provides a reduced intraluminal calcium microenvironment, resulting in

enhanced store-operated calcium entry. *Sci Rep.* 8:1–13.

Santoso, N. G., Cebotaru, L. and Guggino, W. B. 2011. Polycystin-1, 2, and STIM1

interact with IP$_3$R to modulate ER Ca^{2+} release through the PI3K/Akt pathway.

Cell Physiol Biochem. 27:715–726.

Srivats, S., Balasuriya, D., Pasche, M., Vistal, G., Edwardson, J.M., Taylor, C.W. and

Murrell-Lagnado, R.D. 2016. Sigma-1 receptors inhibit store-operated Ca^{2+} entry

by attenuating coupling of STIM1 to Orai1. *J. Cell Biol.* 213: 65–79.

Sugawara, H., Kurosaki, M., Takata, M. and Kurosaki, T. 1997. Genetic evidence for

involvement of type 1, type 2 and type 3 inositol 1, 4, 5-trisphosphate receptors

in signal transduction through the B-cell antigen receptor. *EMBO J.* 16:3078–

3088.

Thillaiappan, N.B., Chavda, A.P., Tovey, S.C., Prole, D.L. and Taylor, C.W. 2017.

Ca^{2+} signals initiate at immobile IP$_3$ receptors adjacent to ER-plasma membrane

junctions. *Nat. Commun.* 8:1-16.

Thillaiappan, N.B., Chakraborty, P., Hasan, G. and Taylor, C.W. 2019. IP$_3$ receptors

and Ca^{2+} entry. *Biochim Biophys Acta Mol Cell Res.* 1866:1092–1100.

Tovey, S.C., De Smet, P., Lipp, P., Thomas, D., Young, K.W., Missiaen, L., De Smedt,

H., Parys, J.B., Berridge, M.J., Thuring, J. and Holmes, A. 2001. Calcium puffs

are generic InsP$_3$-activated elementary calcium signals and are downregulated

by prolonged hormonal stimulation to inhibit cellular calcium responses. *J. Cell*

Sci. 114:3979-3989.

Venkiteswaran, G. and Hasan, G. 2009. Intracellular Ca^{2+} signaling and store-

operated Ca^{2+} entry are required in *Drosophila* neurons for flight. *Proc. Natl.*

Acad. Sci. U. S. A. 106:10326–10331.

Wojcikiewicz, R. J. H. 1995. Type I, II, and III inositol 1,4,5-trisphosphate receptors

are unequally susceptible to down-regulation and are expressed in markedly

different proportions in different cell types. *J. Biol. Chem.* 270:11678–11683.

Woo, J.S., Sun, Z., Srikanth, S. and Gwack, Y. 2020. The short isoform of extended

synaptotagmin-2 controls Ca^{2+} dynamics in T cells via interaction with STIM1.

Sci. Rep. 10:1-13.

Woodard, G.E., López, J.J., Jardín, I., Salido, G.M. and Rosado, J.A. 2010. TRPC3

regulates agonist-stimulated Ca^{2+} mobilization by mediating the interaction

between type I inositol 1, 4, 5-trisphosphate receptor, RACK1, and Orai1. *J. Biol.*

Chem. 285: 8045–8053.

Yoshikawa, F., Morita, M., Monkawa, T., Michikawa, T., Furuichi, T. and Mikoshiba, K.

1996. Mutational analysis of the ligand binding site of the inositol 1, 4, 5-

trisphosphate receptor. *J. Biol. Chem.* 271:18277-18284.

Yue, L., Wang, L., Du, Y., Zhang, W., Hamada, K., Matsumoto, Y., Jin, X., Zhou, Y.,

Mikoshiba, K., Gill, D.L. and Han, S. 2020. Type 3 inositol 1, 4, 5-trisphosphate

receptor is a crucial regulator of calcium dynamics mediated by endoplasmic

reticulum in HEK cells. *Cells.* 9:275.

Figure 1. Loss of IP$_3$R1 Attenuates SOCE in Human Neural Stem Cells

(**A**) Confocal images of hNPCs (passage 6) stained for DAPI and neural stem cell proteins: Pax6 and Ki67 (proliferation marker). Scale bars, 50 μm.

(**B**) WB for IP$_3$R1 of hNPCs expressing non-silencing (NS) or IP$_3$R1-shRNA.

(**C**) Summary results (mean ± s.d., n = 3) show IP$_3$R1 expression relative to actin. $^{**}P$ < 0.01, Student's t-test with unequal variances.

(**D**) Changes in [Ca^{2+}]$_c$ evoked by thapsigargin (Tg, 10 μM) in Ca^{2+}-free HBSS and then restoration of extracellular Ca^{2+} (2 mM) in hNPCs expressing NS or IP$_3$R1-shRNA. Mean ± s.e.m. from 3 independent experiments, each with 4 replicates that together included 100-254 cells. Inset shows the target of Tg.

(**E-G**) Summary results (individual cells, median (bar), 25th and 75th percentiles (box) and mean (circle)) show Ca^{2+} signals evoked by Tg or Ca^{2+} restoration (**E**), rate of Ca^{2+} entry (**F**) and resting [Ca^{2+}]$_c$ (**G**). $^{***}P$ < 0.001, Mann-Whitney U-test.

(**H**) Changes in [Ca^{2+}]$_c$ evoked by Tg (10 μM) in Ca^{2+}-free HBSS and after restoring extracellular Ca^{2+} (2 mM) in neurons (differentiated hNPCs) expressing NS or IP$_3$R1-shRNA. Mean ± s.e.m. from 3 experiments with ~200 cells.

(**I,J**) Summary results (presented as in E-G) show Ca^{2+} signals evoked by Tg or Ca^{2+} restoration (**I**) and rate of Ca^{2+} entry (**J**). $^{***}P$ < 0.001. Mann-Whitney U-test.

See also **Figure 1 - figure supplement 1.**

Source data in Figure 1 source data file

Figure 2. Loss of IP$_3$R1 Attenuates SOCE in SH-SY5Y Cells

(**A**) WB for IP$_3$R1-3 of SH-SY5Y cells expressing non-silencing (NS) or IP$_3$R1-shRNA.

(**B**) Summary results (mean ± s.d., $n = 4$) show IP$_3$R expression relative to actin

normalized to control NS cells. $^{**}P < 0.01$, Student's t-test with unequal variances.

(**C**) Ca^{2+} signals evoked by carbachol (CCh, 3 µM) in SH-SY5Y cells expressing NS or

IP$_3$R1-shRNA. Mean ± s.e.m. from 3 experiments with 70-90 cells.

(**D**) Summary results show peak changes in [Ca^{2+}]$_c$ (Δ[Ca^{2+}]$_c$) evoked by CCh. $^{***}P <$

0.001, Mann-Whitney U-test.

(**E**) Ca^{2+} signals evoked by thapsigargin (Tg, 10 µM) in Ca^{2+}-free HBSS and then after

restoration of extracellular Ca^{2+} (2 mM) in cells expressing NS or IP$_3$R1-shRNA. Mean

± s.e.m. from 3 experiments with ~50 cells.

(**F, G**) Summary results (individual cells, mean ± s.e.m., $n = 3$, ~50 cells) show peak

changes in [Ca^{2+}]$_c$ evoked by Ca^{2+} restoration (Δ[Ca^{2+}]$_c$) (**F**) and rate of Ca^{2+} entry (**G**).

$^{***}P < 0.001$, Mann-Whitney U-test.

(**H**) Ca^{2+} signals evoked by Tg and then Ca^{2+} restoration in cells expressing NS-

shRNA, or IP$_3$R1-shRNA alone or with IP$_3$R1 or IP$_3$R3. Traces show mean ± s.e.m.

(50-115 cells from 3 experiments).

(**I, J**) Summary results (mean ± s.e.m, 50-115 cells from 3 experiments) show peak

increases in [Ca^{2+}]$_c$ (Δ[Ca^{2+}]$_c$) evoked by Ca^{2+} restoration (**I**) and rates of Ca^{2+} entry

(**J**) evoked by restoring extracellular Ca^{2+}.

(**K**) Effects of thapsigargin (Tg, 10 µM) in Ca^{2+}-free HBSS and then after Ca^{2+}

restoration (2 mM) in cells expressing IP$_3$R1-shRNA alone or with IP$_3$R1 or mCh-

STIM1. Traces show mean ± s.e.m. (100-150 cells from 3 experiments).

(**L, M**) Summary results (mean ± s.e.m.) show peak increase in $[Ca^{2+}]_c$ after Ca^{2+}

restoration ($\Delta[Ca^{2+}]_c$) (**L**) and rate of Ca^{2+} entry (**M**). Different letters indicate

significant differences (panels I, J, L, M), $P < 0.001$, one-way ANOVA with pair-wise

Tukey's test.

See also **Figure 2 with 3 supplements (Figure 2 - figure supplements 1 - 3).**

Source data in Figure 2 source data file

Figure 3. Regulation of SOCE by IP$_3$R Requires IP$_3$ Binding but Not a Functional

Pore in SH-SY5Y cells

(**A**) SOCE is activated when loss of Ca^{2+} from the ER through IP$_3$Rs activates STIM1

(i). Our results suggest an additional role for IP$_3$Rs (ii).

(**B**) SH-SY5Y cells expressing IP$_3$R1-shRNA alone or with IP$_3$R1 or IP$_3$R1^{DA} were

stimulated with thapsigargin (Tg, 1 µM) in Ca^{2+}-free HBSS before restoring

extracellular Ca^{2+} (2 mM). Traces show mean ± s.e.m, for 100-150 cells from 3

experiments.

(**C**) Cells expressing IP$_3$R1-shRNA and IP$_3$R1^{DA} were treated with NS-siRNA or Orai1-

siRNA before measuring Tg-evoked Ca^{2+} entry. Traces show mean ± s.e.m. for 85-

100 cells from 3 experiments.

(**D**) Summary results (mean ± s.e.m.) show peak increases in [Ca^{2+}]$_c$ (Δ[Ca^{2+}]$_c$)

evoked by Ca^{2+} restoration.

(**E**) Tg-evoked Ca^{2+} entry in cells expressing IP$_3$R1-shRNA with IP$_3$R1, IP$_3$R1^{RQ} or

IP$_3$R1$^{RQ/KQ}$. Traces show mean ± s.e.m, for 90-150 cells from 3 experiments.

(**F**) Summary results (mean ± s.e.m.) show peak increases in [Ca^{2+}]$_c$ (Δ[Ca^{2+}]$_c$)

evoked by Ca^{2+} restoration. Different letter codes (panels D, F) indicate significantly

different values, $P < 0.001$, one-way ANOVA and pair-wise Tukey's test.

See also **Figure S3 - figure supplement 1.**

Source data in Figure 3 source data file

Figure 4.   Receptor-Regulated IP$_3$ Production Stimulates SOCE in Cells With Empty Ca^{2+} Stores and Expressing Pore-Dead IP$_3$R

(**A, B**) SH-SY5Y cells expressing IP$_3$R1-shRNA alone (**A**) or with IP$_3$R1^{DA} (**B**) were treated with a low concentration of CPA (2 µM) in Ca^{2+}-free HBSS to partially deplete the ER of Ca^{2+} and sub-maximally activate SOCE (see **Figures S5M and S5N**). Carbachol (CCh, 1 µM) was then added to stimulate IP$_3$ formation through muscarinic receptors, and extracellular Ca^{2+} (2 mM) was then restored. Traces (mean ± s.e.m of 68-130 cells from 3 experiments) show responses with and without the CCh addition.

(**C**) Summary results show the peak increases in [Ca^{2+}]$_c$ (Δ[Ca^{2+}]$_c$) after addition of CCh (CCh-induced Ca^{2+} release) and then after restoring extracellular Ca^{2+} (SOCE).

(**D-F**) SH-SY5Y cells wild type (WT) (**D**) and expressing NS-shRNA (**E**) or IP$_3$R1-shRNA (**F**) were treated with YM-254890 (YM, 1 µM, 5 min) in Ca^{2+}-free HBSS to inhibit Gαq and then with thapsigargin (Tg, 1 µM) before restoring extracellular Ca^{2+} (2 mM). Traces show mean ± s.e.m of ~120 cells from 3 experiments. .

(**G-I**) Similar analyses of HEK cells. Summary results (mean ± s.e.m, 50-100 cells from 3 experiments) are shown in (I).

Different letter codes (panels C and I) indicate significantly different values within the store Ca^{2+} release or SOCE groups, *P* < 0.001, one-way ANOVA and pair-wise Tukey's test.

See also **Figure 4 - figure supplement 1.**

Source data in Figure 4 source data file

Figure 5. IP$_3$Rs Promote Interaction of STIM1 With Orai1

(**A-E**) PLA analyses of interactions between STIM1 and Orai1 in SH-SY5Y cells

expressing NS-shRNA (**A**) or IP$_3$R1-shRNA alone (**B**) or with IP$_3$R1 (**C**), IP$_3$R1^{DA} (**D**)

or IP$_3$R1$^{RQ/KQ}$ (**E**). Confocal images are shown for control cells or after treatment with

thapsigargin (Tg, 1 µM) in Ca^{2+}-free HBSS. PLA reaction product is red, and nuclei

are stained with DAPI (blue). Scale bars, 5 µm. Summary results show the surface

area of the PLA spots for 8-10 cells from 2 independent analyses. Individual values,

median (bar) and 25th and 75th percentiles (box). $^{***}P < 0.001$, Student's t-test with

unequal variances.

See also **Figure 5 - figure supplement 1.**

Source data in Figure 5 source data file

Figure 6. Extended Synaptotagmins Rescue SOCE in Cells Lacking IP$_3$R1

(**A**) SH-SY5Y cells expressing IP$_3$R1-shRNA alone or with E-Syt1 were stimulated with Tg (1 μM) in Ca^{2+}-free HBSS before restoring extracellular Ca^{2+} (2 mM). Traces show mean ± s.e.m, for 20-80 cells from 3 experiments.

(**B**) Summary results show Δ[Ca^{2+}]$_c$ evoked by restoring Ca^{2+} (SOCE). Mean ± s.e.m, ***P < 0.001, Mann-Whitney U- test.

(**C**) Summary results (mean ± s.e.m, n = 20-80 cells) show resting [Ca^{2+}]$_c$ (left) and the peak Ca^{2+} signals (Δ[Ca^{2+}]$_c$) evoked by thapsigargin (Tg, 1 μM) in Ca^{2+}-free HBSS for SH-SY5Y cells expressing IP$_3$R1-shRNA alone or with human E-Syt1.

(**D**) Cells over-expressing E-Syt1 and treated with IP$_3$R1-shRNA in combination with either NS or STIM1 siRNA were stimulated with Tg (1 μM) in Ca^{2+}-free HBSS before restoration of extracellular Ca^{2+} (2 mM). Mean ± s.e.m. from 3 experiments with 30-40 cells.

(**E, F**) Summary results (mean ± s.e.m, n = 30-40 cells) show SOCE evoked by Tg (**E**), resting [Ca^{2+}]$_c$ and the Tg-evoked Ca^{2+} release from intracellular stores (F). ***P < 0.001, one-way ANOVA with pair-wise Tukey's test.

(**G**) Similar analyses of cells expressing NS shRNA alone or with human E-Syt1 and then treated with Tg (1 μM) in Ca^{2+}-free HBSS before restoring extracellular Ca^{2+} (2 mM). Mean ± s.e.m. from 3 experiments with 115-135 cells.

(**H, I**) Summary results (mean ± s.e.m, n = 115-135 cells) show resting [Ca^{2+}]$_c$ (**H**) and Δ[Ca^{2+}]$_c$ evoked by Tg (store release) or Ca^{2+} restoration (SOCE) (**I**). No significant difference, one-way ANOVA with pair-wise Tukey's test.

Source data in Figure 6 source data file

Figure 7. Dual Regulation of SOCE by IP$_3$Rs

(**A**) SOCE is activated when loss of Ca^{2+} from the ER, usually mediated by opening of IP$_3$Rs when they bind IP$_3$, causes STIM to unfurl cytosolic domains (2). The exposed cytosolic domains of STIM1 reach across a narrow gap between the ER and PM at a MCS to interact with PIP$_2$ and Orai1 in the PM. Binding of STIM1 to Orai1 causes pore opening, and SOCE then occurs through the open Orai1 channel. We show that IP$_3$Rs when they bind IP$_3$ also facilitate interactions between Orai1 and STIM, perhaps by stabilizing the MCS (1). Receptors that stimulate IP$_3$ formation thereby promote both activation of STIM (by emptying Ca^{2+} stores) and independently promote interaction of active STIM1 with Orai1.

(**B**) Other mechanisms, including ryanodine receptors (RyR), can also release Ca^{2+} from the ER. We suggest that convergent regulation of SOCE by IP$_3$R with bound IP$_3$ allows receptors that stimulate IP$_3$ formation to selectively control SOCE.

 Figure 1 - figure supplement 1

 Loss of IP$_3$R1 Attenuates SOCE In Neural Precursor Cells and Differentiated

 Neurons.

(**A-C**) hNPCs expressing NS (**A, C**) or IP$_3$R1-shRNA (**B**) were stimulated with

carbachol (CCh, 100 µM) in Ca^{2+}-free HBSS before restoring extracellular Ca^{2+} (10

mM). Each trace shows the Fura 2 fluorescence ratio (F$_{340}$/F$_{380}$) from a single cell (>50

cells from 3 experiments). Traces in (**C**) show only cells that responded to CCh.

(**D, E**) hNPCs expressing shRNA were stimulated with thapsigargin (Tg, 10 µM) in

Ca^{2+}-free HBSS before restoring extracellular Ca^{2+} (2 mM). Traces are from individual

cells (>180 cells from 3 experiments), with the mean response shown by a thick line.

Summary results in **Figures 1D-1G.**

(**F**) Confocal images of hNPCs and neurons spontaneously differentiated (10 days)

from hNPCs stained for DAPI and neuronal markers: Tuj1 (βIII-tubulin) and NeuN

(neuronal nuclear antigen). Scale bars, 50 µm. Only differentiated hNPCs express

neuronal markers.

(**G**) WB for STIM1 and Orai1 in lysates from hNPCs cells expressing non-silencing

(NS) or IP$_3$R1-shRNA. Summary results (mean ± s.d., n = 3) show relative expression

of STIM1 and Orai1 relative to control shRNA (NS) cells. $P > 0.01$, Student's t-test

with unequal variances.

(**H, I**) Spontaneously differentiated hNPCs (25 days) expressing NS (**H**) or IP$_3$R1-

shRNA (**I**) were depolarized by addition of KCl (75 mM). Traces show responses from

single cells (>30 cells from 3 experiments) and mean response (thick line).

(**J**) Summary results (mean ± s.e.m., 3 experiments) show peak increases in $[Ca^{2+}]_c$

($\Delta[Ca^{2+}]_c$) evoked by depolarization. No significant difference, Student's *t*-test with

unequal variances.

(**K, L**) Spontaneously differentiated hNPCs expressing NS (**K**) or IP$_3$R1-shRNA (**L**)

were stimulated with Tg (10 µM) in Ca^{2+}-free HBSS before restoring extracellular Ca^{2+}

(2 mM). Traces from individual cells (>100 cells from 3 experiments) and the mean

response (bold) are shown. Summary results in **Figure 1H-1J.**

Source data in Figure 1 - figure supplement 1 source data file.

 Figure 2 - figure supplement 1

 Reduced Expression of IP$_3$R1 Using either shRNA or CRISPR/Cas9n Attenuates

 SOCE in SH-SY5Y Cells

(**A, B**) Ca^{2+} signals evoked by carbachol (CCh, 3 µM) in Ca^{2+}-free HBSS in SH-SY5Y

cells expressing NS (**A**) or IP$_3$R1-shRNA (**B**). Traces show responses from single cells

(100-200 cells from 3 experiments).

(**C, D**) SH-SY5Y cells expressing NS (**C**) or IP$_3$R1-shRNA (**D**) were treated with

thapsigargin (Tg, 1 µM) in Ca^{2+}-free HBSS before restoration of extracellular Ca^{2+} (2

mM). Traces show responses from single cells (50-100 cells from 3 experiments) and

mean response (thick lines). Summary results in **Figures 2C-2G**.

(**E**) Western blots (WB) for STIM1 and Orai1 from lysates of SH-SY5Y cells

expressing non-silencing (NS) or IP$_3$R1-shRNA. Summary results (mean ± s.d., n = 3)

show STIM1 and Orai1 expression relative to NS-shRNA cells. $P > 0.05$, Student's t-

test with unequal variances.

(**F**) Western blot for IP$_3$R1 and tubulin from lysates from WT SH-SY5Y cells and for

cells from a cell line in which the IP$_3$R1 gene was targeted using CRISPR/Cas9 (IKO

null). The blot is typical of 3 similar analyses. Summary results (mean ± s.d., n = 3)

show IP$_3$R1 expression relative to WT cells. $P < 0.001$, Student's t-test with unequal

variances.

(**G**) Western blots (WB) for STIM1 (S1), STIM2 (S2) and Orai1 (O1) from lysates of

WT and IKO null SH-SY5Y cells. Summary results (mean ± s.d., n = 3) show STIM1,2

and Orai1 expression relative to WT cells. $P > 0.05$, Student's t-test with unequal

variances.

(**H**) Ca^{2+} signals evoked by carbachol (CCh, 1 μM) in WT and IKO null SH-SY5Y cells

in Ca^{2+}-free HBSS.

(**I**) SH-SY5Y cells (WT or IKO null) were treated with thapsigargin (Tg, 1 μM) in Ca^{2+}-

free HBSS before restoring extracellular Ca^{2+} (2 mM). Mean ± s.e.m. from 3

experiments with WT cells (black, 110 cells), IKO null cells (red, 117 cells).

(**J**) Summary results show the peak increases in $[Ca^{2+}]_c$ evoked by Tg (1 μM) in Ca^{2+}-

free HBSS ($\Delta[Ca^{2+}]_c$) and after Ca^{2+} restoration (SOCE). Different letters indicate P <

0.001, Mann-Whitney U-test.

(**K**) PCRs for IP_3R1 and GAPDH (control gene) of genomic DNA isolated from SH-

SY5Y cells. Results (mean $\pm$ s.e.m. from 3 independent analyses of a single cell line)

are shown for wild type (WT) cells and for cells in which the IP_3R1 gene was targeted

using CRISPR/Cas9n (IKO). Expression of IP_3R1 has been normalized to expression

of GAPDH. $^{**}P < 0.01$, Student's *t*-test with unequal variances.

(**L**) Droplet digital PCR amplification was used to determine gene copy number for

IP_3R1 and a control gene RPP30 for WT and IKO SH-SY5Y cells. $**P < 0.05$,

Student's *t*-test with unequal variances.

(**M**) Western blot for IP_3R1 and tubulin from lysates from WT SH-SY5Y cells and for

cells from a cell line in which the IP_3R1 gene was targeted using CRISPR/Cas9n

(IKO). The blot is typical of 3 similar analyses. Summary results (mean ± s.d., n = 3)

show IP_3R1 expression relative to WT cells. P < 0.01, Student's *t*-test with unequal

variances.

(**N**) Ca^{2+} signals evoked by carbachol (CCh, 1 μM) in WT and IKO SH-SY5Y cells in

Ca^{2+}-free HBSS. Results show (mean ± s.e.m.) for each of three independently edited

cell lines (IKO1, 2 and 3), each with at least 30 cells; and the overall mean ± s.e.m.

(red line). Results from a total of 89 WT cells and 130 IKO cells are shown.

(**O**) Summary results (mean ± s.e.m. from 3 experiments with 80-100 cells) show peak

changes in $[Ca^{2+}]_c$ ($\Delta[Ca^{2+}]_c$) evoked by CCh. ***P < 0.001, Mann-Whitney U-test.

(**P**) SH-SY5Y cells (WT or IKO) were treated with thapsigargin (Tg, 1 µM) in Ca^{2+}-free

HBSS before restoring extracellular Ca^{2+} (2 mM). Mean ± s.e.m. from 3 experiments

with WT cells (black, 85 cells), 3 independently edited cell lines (IKO 1, 2 and 3; at

least 30 cells for each) and the pooled results from all edited cells (97; red).

(**Q**) Summary results show the peak increases in $[Ca^{2+}]_c$ evoked by Tg (1 µM) in Ca^{2+}-

free HBSS ($\Delta[Ca^{2+}]_c$) and after Ca^{2+} restoration (SOCE). Different letters indicate P <

0.001, Mann-Whitney U-test.

Source data in Figure 2 - figure supplement 1 source data file.

Figure 2 - figure supplement 2

Attenuated SOCE In SH-SY5Y Cells Lacking IP$_3$R1 is Rescued by Expression of

IP$_3$R1, IP$_3$R3 or STIM1.

(**A**) Ca^{2+} signals evoked by carbachol (CCh, 3 µM) in SH-SY5Y cells expressing NS or

IP$_3$R1-shRNA, and the latter with IP$_3$R1 or IP$_3$R3. Traces show mean ± s.e.m. from >3

experiments with 50-70 cells.

(**B-D**) Summary results show peak Ca^{2+} signals (Δ[Ca^{2+}]$_c$) evoked by CCh (**B**), resting

[Ca^{2+}]$_c$ (**C**) and the peak Ca^{2+} signal evoked by Tg (10 µM) in Ca^{2+}-free HBSS

(Δ[Ca^{2+}]$_c$, a reporter of ER Ca^{2+} content) (**D**). Different letter codes indicate

significantly different values, $P < 0.001$, one-way ANOVA and pair-wise Tukey's test.

(**E**) Effects of expressing mCh-STIM1 in SH-SY5Y cells expressing NS shRNA on the

Ca^{2+} signals evoked by Tg (1 µM) in Ca^{2+}-free HBSS and then after restoration of

extracellular Ca^{2+} (2 mM). Traces show mean ± s.e.m. from 3 experiments with 30-

110 cells.

(**F**) Summary results show the peak Δ[Ca^{2+}]$_c$ evoked by restoring extracellular Ca^{2+}

after Tg. Mean ± s.e.m. $P > 0.05$, Mann-Whitney U-test.

Source data in Figure 2 - figure supplement 2 source data file.

Figure 2 - figure supplement 3

Loss of IP$_3$R1 Does Not affect SOCE in HEK Cells.

(**A**) WB for IP$_3$R1 in HEK cells expressing non-silencing (NS) or IP$_3$R1-shRNA. Summary results (mean ± s.d., n = 3) show IP$_3$R1 expression relative to tubulin. [***]$P <$ 0.001, Student's t-test with unequal variances.

(**B**) HEK cells expressing NS or IP$_3$R1-shRNA were treated with thapsigargin (Tg, 1 µM) in Ca^{2+}-free HBSS before restoration of extracellular Ca^{2+} (2 mM). Traces show mean response from >100 cells from 3 experiments.

(**C**) Summary results show the peak increases in [Ca^{2+}]$_c$ evoked by Tg (1 µM) in Ca^{2+}-free HBSS (Δ[Ca^{2+}]$_c$, a reporter of ER Ca^{2+} content) and after Ca^{2+} restoration (SOCE).

(**D**) Wild type HEK cells (HEK WT) and cells lacking all IP$_3$Rs (HEK-TKO) were treated with thapsigargin (Tg, 1 µM) in Ca^{2+}-free HBSS before restoration of extracellular Ca^{2+} (2 mM). Traces show mean ± s.e.m. from 3 experiments with 80-120 cells.

(**E**) Summary results show the peak Δ[Ca^{2+}]$_c$ evoked by restoring extracellular Ca^{2+} after Tg. Mean ± s.e.m. $P > 0.01$, Mann-Whitney U-test.

(**F**) Proximity ligation assay (PLA) analyses of interactions between STIM1 and Orai1 in wild type HEK (WT) and TKO cells. Confocal images are shown for control cells (-Tg) or after treatment with thapsigargin (+Tg, 1 µM) in Ca^{2+}-free HBSS. PLA reaction product is red, and nuclei are stained with DAPI (blue). Scale bars, 5 µm. Summary results show the surface area of the PLA spots for ~20 cells from 2 independent analyses. Individual values, median (bar) and 25th and 75th percentiles (box). ***P < 0.01, Student's t-test with unequal variances.

803   **Source data in Figure 2 - figure supplement 3 source data file.**

Figure 3 - figure supplement 1

Attenuated SOCE In SH-SY5Y Cells Lacking IP$_3$R1 is Rescued by Expression of Pore-Dead IP$_3$R1 with a Functional IP$_3$-Binding Site.

(**A**) WB with IP$_3$R1 antibody showing expression of IP$_3$R1 and IP$_3$R1^{DA} in SH-SY5Y cells expressing IP$_3$R1-shRNA.

(**B**) Summary (mean ± s.e.m, n = 3) shows IP$_3$R1 expression normalized to actin. $P <$ 0.001, one-way ANOVA with pair-wise Tukey's test. Here, and in most subsequent panels, different letters indicate groups that are statistically different, with the test defined for each panel.

(**C**) Ca^{2+} signals evoked by carbachol (CCh 3 µM) in SH-SY5Y cells expressing IP$_3$R1-shRNA alone or with IP$_3$R1, IP$_3$R1^{DA}, IP$_3$R1$^{RQ/KQ}$ or IP$_3$R1$^{1-604}$. Traces show mean ± s.e.m. from 3 experiments.

(**D**) Summary results (mean ± s.e.m) show peak changes in [Ca^{2+}]$_c$ evoked by CCh (Δ[Ca^{2+}]$_c$). $P <$ 0.001, one-way ANOVA with pair-wise Tukey's test.

(**E**) Resting [Ca^{2+}]$_c$ (left) and thapsigargin (Tg)-evoked Ca^{2+} release (1 µM, Δ[Ca^{2+}]$_c$) (right) in cells treated as indicated. $P <$ 0.001, one-way ANOVA with pair-wise Tukey's test.

(**F**) WB showing Orai1 expression in cells expressing IP$_3$R1-shRNA and IP$_3$R1^{DA}, and then treated with NS or Orai1-siRNA.

(**G**) Summary results (normalized to GAPDH expression, mean ± s.d, $n =$3). $^{**}P <$ 0.01, Student's t-test with unequal variances.

(**H**) Resting $[Ca^{2+}]_c$ (left) and Tg-evoked Ca^{2+} release (right) in cells expressing IP_3R1-

shRNA and IP_3R1^{DA}, and then treated with NS or Orai1-siRNA. $P < 0.05$, Mann-

Whitney U-test.

(**I**) WB for IP_3R1 in cells expressing IP_3R1-shRNA alone or with IP_3R1 or $IP_3R1^{RQ/KQ}$.

Results typical of 2 independent experiments.

(**J**) Resting $[Ca^{2+}]_c$ (left) and Ca^{2+} release evoked by Tg (right) in cells expressing

IP_3R1-shRNA alone or with the indicated IP_3R1. Mean $\pm$ s.e.m., $n = 3$ experiments. P

< 0.001, one-way ANOVA with pair-wise Tukey's test.

(**K**) Changes in $[Ca^{2+}]_c$ evoked by Tg (10 µM) in Ca^{2+}-free HBSS and then after

restoration of extracellular Ca^{2+} (2 mM) in cells expressing IP_3R1-shRNA alone or with

IP_3R1 or IP_3R1^{1-604}. Mean $\pm$ s.e.m. from 3 experiments with >100 cells.

(**L**) Summary results show peak increases in $[Ca^{2+}]_c$ evoked by Ca^{2+} restoration in the

indicated cells. $P < 0.001$, one-way ANOVA with pair-wise Tukey's test.

(**M**) SH-SY5Y cells expressing NS-shRNA were stimulated with the indicated

concentrations of CPA in Ca^{2+}-free HBSS to partially deplete ER Ca^{2+} stores and then

with ionomycin (Iono, 1 µM) to release all remaining Ca^{2+}. Results show mean $\pm$

s.e.m. for >30 cells from 3 experiments.

(**N**) Summary results (mean $\pm$ s.e.m.) show $\Delta[Ca^{2+}]_c$ evoked by CPA or ionomycin. $P <$

0.001, one-way ANOVA with pair-wise Tukey's test. The results confirm that the lower

concentrations of CPA partially deplete the ER of Ca^{2+}.

(**O**) Summary results (mean $\pm$ s.e.m, 80-100 cells from 3 experiments) show the peak

increases in $[Ca^{2+}]_c$ ($\Delta[Ca^{2+}]_c$) after adding Tg (store Ca^{2+} release) and after restoring

extracellular Ca^{2+} (SOCE) with or without YM-254890 treatment. Different letter codes

indicate significantly different values within the Ca^{2+} release or SOCE groups, $P <$

0.001, one-way ANOVA and pair-wise Tukey's test.

Source data in Figure 3 - figure supplement 1 source data file.

Figure 4 - figure supplement 1

Effects of Inhibiting Gq on Ca^{2+} Signals and STIM1-Orai1 interactions in SH-SY5Y and HEK Cells.

(**A**) PLA analyses of interactions between STIM1 and Orai1 in wild type (WT) SH-SY5Y cells untreated (-YM) or treated (+YM, 1µm) with Gq inhibitor YM-254890. Confocal images are shown for control cells (-Tg) or after treatment with thapsigargin (+Tg, 1 µM) in Ca^{2+}-free HBSS. PLA reaction product is red, and nuclei are stained with DAPI (blue). Scale bars, 5 µm. Summary results show the surface area of the PLA spots for ~20 cells from 2 independent analyses. Individual values, median (bar) and 25th and 75th percentiles (box). ***P < 0.01, Student's t-test with unequal variances.

(**B**) HEK cells expressing NS shRNA and treated with the Gq inhibitor, YM-254890 (YM, 1 µM, 5 min) were stimulated with carbachol (CCh, 100 µM). Traces show mean ± s.e.m. from 3 experiments with 30-40 cells. The results confirm that treatment with YM-254890 effectively uncouples CCh from IP$_3$-evoked Ca^{2+} signals.

(**C**) From results shown in **Figures 4G and 4H**, rates of thapsigargin-evoked Ca^{2+} entry were determined. Summary shows mean ± s.e.m, from 100-120 cells. *P* < 0.001, Mann-Whitney U-test.

(**D-E**) Changes in [Ca^{2+}]$_c$ evoked by thapsigargin (Tg, 1 µM) in Ca^{2+}-free HBSS and then after restoration of extracellular Ca^{2+} (2 mM) in WT (**C**) and TKO (**D**) HEK cells after treatment with YM-254890 (YM, 1 µM, 5 min). Traces show mean ± s.e.m. from 3 experiments with 100-120 cells.

873   (**F**) Summary results show peak increase in $[Ca^{2+}]_c$ ($\Delta[Ca^{2+}]_c$) evoked by addition of

874   thapsigargin and then after restoring extracellular Ca^{2+} (SOCE). Mean ± s.e.m.

875   Different letter codes (C and G) indicate significantly different values, $P < 0.001$, one-

876   way ANOVA and pair-wise Tukey's test.

877   **Source data in Figure 4 - figure supplement 1 source data file.**

Figure 5 - figure supplement 1

Validation of PLA Measurements of Orai1-STIM1 Interactions.

(**A**) PLA using antibodies to Orai1 and STIM1 to establish whether they are associated. Complementary oligonucleotides conjugated to a secondary antibody (2[nd]) targeting their respective primary antibodies (1[st]) hybridize when they are close to each other, allowing rolling circle amplification (RCA) and production of a red fluorescent product. The latter can be quantified from the surface area of the fluorescent spots or their intensity. We obtained indistinguishable results with both methods of quantification. We therefore show only results obtained from surface-area measurements.

(**B-D**) Confocal images from PLA assays of SH-SY5Y cells expressing NS-shRNA performed with no primary antibody (**B**), or with primary antibody for only STIM1 (**C**) or only Orai1 (**D**). Scale bars, 5 μm.

(**E, F**) Confocal images of SH-SY5Y cells expressing NS-shRNA with (**F**) or without thapsigargin treatment (**E**) (Tg, 1 μM, in Ca^{2+}-free HBSS). Cells are stained with DAPI and immunostained for Orai1 or STIM1. Scale bars, 5 μm.

(G-N) Similar analyses of cells expressing IP_3R1-shRNA alone (**G, H**), or with IP_3R1 (**I, J**), IP_3R1^{DA} (**K, L**) or $IP_3R1^{RQ/KQ}$ (**M, N**). Results (**E-N**) are typical of 2 experiments. The PLA results are shown in **Figure 5**.

RESOURCE AVAILABILITY

Lead Contact

All requests for resources and reagents should be directed to the lead contact, Dr.

Gaiti Hasan (gaiti@ncbs.res.in).

Materials Availability

Constructs and cell lines are available upon request. MTA required for cell lines.

Data and Code Availability

This study did not generate any computer code. The data supporting the findings of

this study are available within the manuscript. All other data supporting the findings of

this study are available in source data file of respective figures.

METHOD DETAILS

Culture of Human Neural Precursor Cells

Human neural precursor cells (hNPCS) were derived from a human embryonic stem

cell (hESC) line, H9/WA09 (RRID: CVCL_9773), using a protocol that inhibits dual

SMAD signalling and stimulates Wnt signaling (Reinhardt *et al.*, 2013) as described

previously (Gopurappilly *et al.*, 2018, 2019). hNPCs were grown as adherent

dispersed cells on growth factor-reduced Matrigel (0.5%, Corning, Cat#356230) in

hNPC maintenance medium (NMM) at 37°C in humidified air with 5% CO_2. NMM

comprised a 1:1 mixture of Dulbecco's Modified Eagle Medium with Nutrient Mixture

F-12 (DMEM/F-12, Invitrogen, Cat#10565018) and Neurobasal medium

(ThermoFisher, Cat#21103049), supplemented with GlutaMAX (0.5x, ThermoFisher,

Cat#35050061), N2 (1:200, ThermoFisher, 17502048), B27 without vitamin A (1:100,

ThermoFisher, Cat#12587010), Antibiotic-Antimycotic (ThermoFisher,

Cat#15240112), CHIR99021 (3 µM, STEMCELL Technologies, Cat#72052),

purmorphamine (0.5 mM, STEMCELL Technologies, Cat#72202) and ascorbic acid

(150 µM, Sigma, Cat#A92902). Doubling time was ~24 hr. Cells were passaged every

4-5 days by treatment with StemPro Accutase (ThermoFisher, Cat#A1110501), stored

in liquid nitrogen, and thawed as required. Cells were confirmed to be mycoplasma-

free by monthly screening (MycoAlert, Lonza, Cat#LT07-318). hNPCs between

passages 16-19 were used.

All experiments performed with hESC lines were approved by the Institutional

Committee for Stem Cell Research, registered under the National Apex Committee for

Stem Cell Research and Therapy, Indian Council of Medical Research, Ministry of

Health, New Delhi.

Stable Knockdown of IP$_3$R1

An UltramiR lentiviral inducible shRNA-mir based on the shERWOOD algorithm

(Auyeung *et al.*, 2013; Knott *et al.*, 2014) was used to inhibit IP$_3$R1 expression. The

all-in-one pZIP vector, which allows puromycin-selection and doxycycline-induced

expression of both shRNA-mir and Zs-Green for visualization, was from TransOMIC

Technologies (Huntsville, AL). Lentiviral pZIP transfer vectors encoding non-silencing

shRNA (NS, NT#3-TTGGATGGGAAGTTCACCCCG) or IP$_3$R1-targeting shRNA

(ULTRA3316782- TTTCTTGATCACTTCCACCAG) were packaged as lentiviral

particles using packaging (pCMV- dR8.2 dpvr, Addgene, plasmid #8455) and

envelope vectors (pCMV-VSV-G, Addgene, plasmid #8454) by transfection of

HEK293T cells (referred as HEK, ATCC, Cat# CRL-3216). Viral particles were

collected and processed and hNPCs (passage 9) or SH-SY5Y cells were transduced

(multiplicity of infection, MOI = 10) using Lipofectamine LTX with PLUS reagent

(ThermoFisher, Cat#15338100). Cells were maintained in media containing

doxycycline (2 µg/ml, Sigma, Cat# D3072) to induce shRNA expression, and

puromycin to select transduced cells (1 µg/ml for hNPCs; 3 µg/ml for SH-SY5Y cells;

Sigma, Cat# P9620). Cells were passaged 4-5 times after lentiviral transduction to

select for stable expression of shRNAs.

Derivation of Neurons From hNPCs

Neurons were differentiated from hNPCs stably transduced with shRNA. hNPCs were

seeded at 50-60% confluence in NMM on coverslips coated with poly-d-lysine (0.2

mg/ml, Sigma, Cat#P7280). After 1-2 days, the medium was replaced with neuronal

differentiation medium, which comprised a 1:1 mixture of DMEM/F-12 with Neurobasal

supplemented with B27 (1:100), N2 (1:200), GlutaMAX (0.5x) and Antibiotic-

Antimycotic solution. Medium was replaced on alternate days. Neurons were used

after 15-20 days.

Culture and Transfection of SH-SY5Y Cells

SH-SY5Y cells (ATCC, USA, Cat# CRL-2266) were grown on culture dishes in

DMEM/F-12 with 10% fetal bovine serum (Sigma, Cat# F4135) at 37□°C in humidified

air with 5% CO_2. Cells were passaged every 3-4 days using TrypLE Express

(ThermoFisher, Cat# 12605036) and confirmed to be free of mycoplasma. Cells

expressing shRNA were transiently transfected using TransIT-LT1 reagent (Mirus,

Cat# MIR-2300) in Opti-MEM (ThermoFisher, Cat# 31985062). Plasmids (250 ng)

and/or siRNA (200 ng) in transfection reagent (1 μg/2.5 μl) were added to cells grown

to 50% confluence on glass coverslips attached to an imaging dish. Cells were used

after 48 hrs. The siRNAs used were to human Orai1 (100 nM, Dharmacon, Cat# L-

014998-00-0005) or non-silencing (NS, Dharmacon, Cat# D-495 001810-10-05), to

human STIM1 (Santa Cruz Biotechnology, Cat# sc-76589) or NS (Santa Cruz

Biotechnology, Cat# sc-37007). The expression plasmids were IP$_3$R1 (rat type 1

IP$_3$R1 in pcDNA3.2/V5DEST vector) (Dellis *et al.*, 2008), rat IP$_3$R1^{DA} (D2550 replaced

by A in pcDNA3.2 vector) (Dellis *et al.*, 2008), rat IP$_3$R1^{RQ} (R568 replaced by Q of type

1 IP$_3$R in pCDNA3.2/V5DEST vector) (Dellis *et al.*, 2008), rat IP$_3$R1$^{RQ/KQ}$ (R568 and

K569 replaced by Q of type 1 IP$_3$R in pCDNA3.2/V5DEST vector), rat IP$_3$R1$^{1-604}$

(residues 1-604 of IP$_3$R with N-terminal GST tag in pCDNA3.2/V5DEST vector) (Dellis

et al., 2008), rat IP$_3$R3 (rat type 3 IP$_3$R in pcDNA3.2/V5DEST vector) (Saleem *et al.*,

2013), human mCherry-STIM1 (N terminal mCherry tagged human STIM1 in

pENTR1a vector) (Nunes-Hasler *et al.*, 2017) and human extended synaptotagmin 1

(E-Syt1), a kind gift from Dr S. Muallem, NIDCR, USA (Maléth *et al.*, 2014).

CRISPR/Cas9 and Cas9n Editing of SH-SY5Y Cells

To allow either CRISPR/Cas9 or Cas9n-mediated disruption of IP$_3$R1 expression, we

used a published method to clone gRNAs into the backbone vector (pSpCas9n(BB)-

2A-Puro PX462 V2.0, Addgene, Cat#62987) (Ran *et al.*, 2013). Forward and reverse

sgRNA oligonucleotides (100 μM) were annealed and ligated using T4 DNA ligase by

incubation (10 μl, 37°C, 30 min) before slow cooling to 20°C. Plasmids encoding

Cas9n were digested with *BbsI-HF* (37°C, 12 hr), gel-purified (NucleoSpin Gel and

PCR Clean-up kit from Takara) and the purified fragment was stored at -20°C. A

mixture (final volume 20 µl) of gRNA duplex (1 µl, 0.5 µM), digested px459 (for IKO

null) or pX462 vector (for IKO) (30 ng), 10×⬜T4 DNA ligase buffer (2 µl) and T4 DNA

ligase (1 µl) was incubated (20°C, 1 hr). After transformation of DH5-α competent *E.*

coli with the ligation mixture, plasmids encoding Cas9 or Cas9n and the sgRNAs were

extracted, and the coding sequences were confirmed (Ran *et al.*, 2013). The plasmid

(2 µg) was then transfected into SH-SY5Y cells (50-60% confluent) in a 6-well plate

using TransIT LT-1 reagent (Mirus Bio, Cat# MIR-2300). After 48 hr, puromycin (3

µg/ml, 72 hr) was added to kill non-transfected cells. IKO colonies were propagated

and screened for Ca^{2+} signals evoked by carbachol and for the presence of the IP_3R

gene by genomic DNA PCR and droplet digital PCR using primers close to the region

targeted by the gRNAs (Miotke *et al.*, 2014). Three independently derived IKO lines,

each with one residual IP_3R1 gene, were used for analyses of Ca^{2+} signaling (see

Figure S2N-S2Q). For one of the cell lines (IKO 2), disruption of one copy of the

IP_3R1 gene was confirmed by genomic PCR, droplet digital PCR and Western blotting

(**Figures S2K-S2M**). For the IKO null line, single cell selection was done in a 96 well

plate setup followed by screening for carbachol-evoked Ca^{2+} signals from multiple

clones. A single clone was selected (**Figure S2H**) and a western blot performed to

confirm absence of IP_3R1 expression (**Figure S2F**). All the oligonucleotide

sequences are described in **supplementary file 1.**

Ca^{2+} Imaging

Methods for single-cell Ca^{2+} imaging were described previously (Gopurappilly *et al.*,

2019)**.** Briefly, cells grown as a monolayer (~70% confluence) on homemade

coverslip-bottomed dishes were washed and loaded with Fura 2 by incubation with

Fura 2 AM (4 µM, 45 min, 37°C, ThermoFisher, Cat# F1221), washed and imaged at

room temperature in HEPES-buffered saline solution (HBSS). HBSS comprised: 20

mM HEPES, 137 mM NaCl, 5 mM KCl, 2 mM $MgCl_2$, 2 mM $CaCl_2$, 10 mM glucose, pH

7.3. $CaCl_2$ was omitted from Ca^{2+}-free HBSS. Treatments with carbachol (CCh,

Sigma, Cat# C4382), thapsigargin (Tg, ThermoFisher, Cat# 7458), cyclopiazonic acid

(CPA, Sigma Cat# C1530) or high-K^+ HBSS (HBSS supplemented with 75 mM KCl)

are described in legends.

Responses were recorded at 2-s intervals using an Olympus IX81-ZDC2 Focus

Drift-Compensating Inverted Microscope with 60× oil immersion objective (numerical

aperture, NA = 1.35) with excitation at 340 nm and 380 nm. Emitted light (505 nm)

was collected with an Andor iXON 897E EMCCD camera and AndoriQ 2.4.2 imaging

software (RRID: SCR_014461). Maximal (R_{max}) and minimal (R_{min}) fluorescence ratios

were determined by addition of ionomycin (10 µM, Sigma, Cat# 407953) in HBSS

containing 10 mM $CaCl_2$ or by addition of Ca^{2+}-free HBSS containing BAPTA (10 mM,

Sigma, Cat# 196418) and Triton X100 (0.1%). Background-corrected fluorescence

recorded from regions of interest ROI) drawn to include an entire cell was used to

determine mean fluorescence ratios (R = F_{340}/F_{380}) (ImageJ), and calibrated to $[Ca^{2+}]_c$

from (Grynkiewicz, Poenie and Tsien, 1985):

$$[Ca^{2+}]_c = K_D \cdot \frac{F_{380}^{min}}{F_{380}^{max}} \cdot \frac{(R - R_{min})}{(R_{max} - R)}$$

where, K_D = 225 nM (Forostyak $et\ al.$, 2013).

1033

Western Blots

1034

Proteins were isolated in RIPA buffer (Sigma, Cat# R0278) with protease inhibitor

cocktail (Sigma, Cat# P8340) or, for WB of Orai1, in medium containing 150 mM

NaCl, 50 mM Tris, 1% Triton-X-100, 0.1% SDS and protease inhibitor cocktail. After

30 min on ice with intermittent shaking, samples were collected by centrifugation

(11,000 ×g, 20 min) and their protein content was determined (Thermo Pierce BCA

Protein Assay kit, ThermoFisher, Cat# 23225). Proteins (~30 µg/lane) were separated

on 8% SDS-PAGE gels for IP$_3$R or 10% SDS-PAGE gels for STIM1 and Orai1, and

transferred to a Protran 0.45-µm nitrocellulose membrane (Merck, Cat# GE10600003)

using a TransBlot semi-dry transfer system (BioRad, Cat# 1703940). Membranes

were blocked by incubation (1 hr, 20°C) in TBST containing skimmed milk or bovine

serum albumin (5%, Sigma, Cat# A9418). TBST (Tris-buffered saline with Tween)

comprised: 137 mM NaCl, 20 mM Tris, 0.1% Tween-20, pH 7.5. Membranes were

incubated with primary antibody in TBST (16 hr, 4°C), washed with TBST (3□×10

min), incubated (1 hr, 20°C) in TBST containing HRP-conjugated secondary antibody

(1:3000 anti-mouse, Cell Signaling Technology Cat# 7076S; or 1:5000 anti-rabbit,

ThermoScientific Cat# 32260). After 3 washes, HRP was detected using Pierce ECL

Western Blotting Substrate (ThermoFisher, Cat# 32106) and quantified using

ImageQuant LAS 4000 (GE Healthcare) and Image J. The primary antibodies used

were to: IP$_3$R1 (1:1000, ThermoFisher, Cat# PA1-901, RRID: AB_2129984); β-actin

(1:5000, BD Biosciences, Cat# 612656, RRID: AB_2289199); STIM1 (1:1000, Cell

Signaling Technology, Cat# 5668S, RRID: AB_10828699); Orai1 (1:500, ProSci, Cat#

PM-5205, RRID: AB_10941192); IP$_3$R2 (1:1000, custom made by Pocono Rabbit

Farm and Laboratory) (Mataragka and Taylor, 2018); and IP$_3$R3 (1:500, BD

Biosciences, Cat# 610313, RRID: AB_397705).

Immunocytochemistry

After appropriate treatments, cells on a coverslip-bottomed plate were washed twice

with cold PBS, fixed in PBS with paraformaldehyde (4%, 20°C, 20 min), washed (3 × 5

min) with PBS containing Triton-X100 (0.1%, PBST) and blocked by incubation (1 hr,

20°C) in PBST containing goat serum (5%). After incubation with primary antibody in

PBST (16 hr, 4°C) and washing with PBST (3 × 5 min), cells were incubated (1 hr,

20°C) with secondary antibody in PBST containing goat serum, washed (3 × 5 min),

stained (10 min, 20°C) with DAPI (1 μg/ml in PBS; Sigma, Cat# D9542) and washed

(5 min, PBST). Cells were then covered with glycerol (60% v/v) and imaged using an

Olympus FV300 confocal laser scanning microscope with 20× or 60× oil-immersion

objectives. Fluorescence was analyzed using ImageJ. The primary antibodies used

were to: PAX6 (1:500, Abcam, Cat# ab195045, RRID: AB_2750924); Nestin (1:500,

Abcam, Cat# 92391, RRID: AB_10561437); Ki67 (1:250, Abcam, Cat# ab16667,

RRID: AB_302459); SOX1 (1:1000, Abcam, Cat# ab87775, RRID: AB_2616563); Tuj1

(βIII Tubulin 1:1000, Promega, Cat# G712, RRID: AB_430874); NeuN (1:300, Abcam,

Cat# ab177487, RRID: AB_2532109); Doublecortin (1:500, Abcam, Cat# 18723,

RRID: AB_732011); MAP2 (1:200, Abcam, Cat# ab32454, RRID: AB_776174); STIM1

(1:1000, Cell Signaling Technology, Cat# 5668S, RRID: AB_10828699); and Orai1

(1:500, ProSci, Cat# PM-5205, RRID: AB_10941192).

1078

1079

Proximity Ligation Assay

1080

The Duolink *In Situ* Red Starter Mouse/Rabbit kit was from Sigma (#Cat DUO92101)

and used according to the manufacturer's protocol with primary antibodies to Orai1

(mouse 1:500) and STIM1 (rabbit 1:1000). Cells (~30% confluent) were treated with

thapsigargin (1 µM, 5 min) in Ca^{2+}-free HBSS before fixation, permeabilization, and

incubation with primary antibodies (16 hr, 4°C) and the PLA reactants. Red

fluorescent PLA signals were imaged using an Olympus FV300 confocal laser

scanning microscope, with excitation at 561 nm, and a 60× oil-immersion objective.

Quantitative analysis of the intensity and surface area of PLA spots used the "Analyze

particle" plugin of Fiji. Results are shown for 8-10 cells from two biological replicates of

each genotype.

Statistical Analyses

1091

All experiments were performed without blinding or prior power analyses. Independent

biological replicates are reported as the number of experiments (n), with the number

of cells contributing to each experiment indicated in legends. The limited availability of

materials for PLA restricted the number of independent replicates (n) to 2 (each with

8-10 cells). Most plots show means ± s.e.m. (or s.d.). Box plots show 25^{th} and 75^{th}

percentiles, median and mean (see legends). Where parametric analyses were

justified by a Normality test, we used Student's *t*-test with unequal variances for 2-way

comparisons and ANOVA followed by pair-wise Tukey's test for multiple comparisons.

Non-parametric analyses used the Mann-Whitney U-test. Statistical significance is

shown by $^{***}P < 0.001$, $^{**}P < 0.01$, $^{*}P < 0.05$, or by letter codes wherein different letters

indicate significantly different values ($P < 0.001$, details in legends). All analyses used

Origin 8.5 software.

1104

1105 **Details of the plasmids and recombinant DNAs are given in Supplementary file**

1106 **1.**

1107

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 1 figure supplement 1

Figure 2-figure supplement 1

Figure 2-figure supplement 2

Figure 2-figure supplement 3

Figure 3-figure supplement 1

Figure 4-figure supplement 1

Figure 5-figure supplement 1