Evolution and co-evolution of the suck behaviour, a postcopulatory female

resistance trait that manipulates received ejaculate

Pragya Singh[1,2*], Jeremias N. Brand[1] and Lukas Schärer[1]

[1]University of Basel, Department of Environmental Sciences, Zoological Institute, Basel,

Switzerland

*Corresponding Author: Email: pragya.singh42019@gmail.com

ORCiD: https://orcid.org/0000-0002-7411-3206

[2]Current address: Chemical Ecology, Bielefeld University, Universitätsstr. 25, 33615

Bielefeld, Germany

Running title: Evolution of a postcopulatory female resistance trait

Author Contributions: PS made most of the mating movies, scored the reproductive

behaviour, analysed the data, and wrote the manuscript, with input from JNB and LS. JNB

provided the phylogeny and morphological data. All authors have read and approved the final

version.

Data accessibility: All data in this manuscript will be deposited online in a public data

repository.

Abstract

Sexual conflicts over the post-mating fate of received ejaculate can favour traits in one sex

that are costly to the other. Reciprocally mating hermaphrodites face unique challenges as

they mate simultaneously in both the male and female role, potentially leading to receipt of

unwanted ejaculate. Reciprocal mating can then give rise to postcopulatory female resistance

traits that allow manipulation of received ejaculate. A putative example is the suck behaviour,

observed in the flatworm genus *Macrostomum*. It involves the sperm recipient placing its

pharynx over its own female genital opening and appearing to suck, likely removing received

ejaculate after mating. The genus also contains hypodermically-inseminating species that

presumably exhibit unilateral mating and have not been observed to suck. Here, we examine

the evolution of the suck behaviour in *Macrostomum*, aiming to document the mating

behaviour in 64 species. First, we provide videographic evidence that ejaculate is indeed

removed during the suck behaviour in a reciprocally mating species, *Macrostomum hamatum*.

Next, we show evolutionary positive correlations between the presence, duration and

frequency of reciprocal mating behaviour and the suck behaviour, providing clear evidence

that the suck behaviour co-evolves with reciprocal mating behaviour. Finally, we show an

association between reproductive behaviour and reproductive morphology, suggesting that

reproductive morphology can be used for inferring the behavioural mating strategy of a

species. Together our study demonstrates sexual antagonistic coevolution leading to the

evolution of a postcopulatory behavioural trait that functions as a female counter-adaptation

allowing individuals to gain control over received ejaculate in a hermaphroditic sexual

system.

Keywords

Sexual conflict; sexual selection; simultaneous hermaphrodite; co-evolution; reproductive

traits; postcopulatory mechanisms; mating behaviour

Introduction

Sexual conflict is defined as the conflict between the two sexes over their evolutionary interests involving reproduction (Charnov, 1979; Parker, 1979; Arnqvist & Rowe, 2005). The primordial cause of sexual conflict is anisogamy, in which the male sex produces more but smaller gametes (called sperm in animals), whereas the female sex produces fewer but larger gametes (called eggs in animals) (Parker, 2011). Because of this asymmetry, eggs are often a limiting resource for reproductive success, resulting in divergent interests between the two sexes (Bateman, 1948; Lehtonen *et al.*, 2016). Furthermore, these conflicting interests can give rise to traits expressed by one sex that are costly to the other sex, resulting in antagonistic co-evolution between the sexes (Holland & Rice, 1998; Arnqvist & Rowe, 2005). Although work on sexual conflict has primarily focussed on separate-sexed organisms, sexual conflict is also pervasive in the lesser-studied hermaphroditic organisms (Charnov, 1979; Leonard, 1991; Michiels, 1998; Abbott, 2011; Schärer *et al.*, 2015).

Of particular interest is the biology of simultaneous hermaphrodites (referred to as hermaphrodites hereafter), which involves unique sexual conflicts. For example, there can be conflicts between the mating partners over the sex role exhibited in a mating, namely mating as a sperm donor, a sperm recipient, or both. Depending on the costs and benefits of mating in each role, this may lead to sex role preferences (Michiels, 1998; Schärer *et al.*, 2015). These are linked to Bateman's principle, a term coined by Charnov (1979), which reflects the notion that there is a "greater dependence of males for their fertility on frequency of insemination" (Bateman, 1948). In his seminal paper, Charnov (1979) explored the proposal that Bateman's principle also applies to simultaneous hermaphrodites. He concluded that, if true, hermaphroditic individuals may often mate more in order to give away sperm than to receive sperm, resulting in a mating conflict between the partners.

This conflict over the sex roles can be resolved via different mating strategies. One such

strategy is reciprocal mating (also called reciprocal copulation), in which the partners

simultaneously mate in both the male and female role. Each sperm donor is thus also a sperm

recipient, and while multiple mating offers more opportunities to donate sperm, it may also

lead to receipt of unwanted ejaculate from the partners. While this strategy seems like a

cooperative conflict resolution, it could shift the conflict from the precopulatory to the

postcopulatory arena (Schärer *et al.*, 2015). In the presence of sperm competition, a donor—in

order to secure a greater share of paternity—may often donate more sperm than the recipient

requires for fertilisation, thereby potentially causing direct costs, such as a risk of polyspermy

(Frank, 2000). But even if there are no direct costs posed by the received ejaculate, mating

with multiple partners—which is probably the norm for most species (Jennions & Petrie,

2007; Kokko & Mappes, 2013; Arbuthnott *et al.*, 2015)—could lead to the evolution of

cryptic female choice (Charnov, 1979; Eberhard, 1996; Hemmings & Birkhead, 2017). Thus,

receipt of excessive or unwanted ejaculate can favour the evolution of female resistance traits

that allow postcopulatory control and rejection of the received ejaculate, e.g. via sperm

digestion (Charnov, 1979).

Female resistance traits can in turn favour the evolution of male persistence traits, including

other mating strategies. Such counter-adaptations may allow the sperm donor to either

counteract or bypass the female resistance traits, thereby retain or regain access to the

recipient's eggs (Charnov, 1979; Schärer *et al.*, 2015). An example of such an alternative

mating strategy involves forced unilateral hypodermic insemination (also called hemocoelic

insemination; Charnov 1979). Here one of the partners mates in the male role and donates

sperm, while the other mates in the female role, potentially against its interests, and receives

sperm hypodermically via a traumatic male copulatory organ (Lange *et al.*, 2013; Reinhardt *et*

al., 2015). With both types of mating strategy these sexual conflicts could then lead to the

evolution of multiple male persistence and female resistance traits (spanning behaviour,

morphology and physiology) that act jointly to either gain access to eggs, or to control and

reject the received ejaculate, respectively (Arnqvist & Rowe, 2005). Therefore, we might

expect behavioural mating strategies to be involved in sexually antagonistic coevolution, and

thus to be correlated with morphological and/or physiological traits.

A putative example of a behavioural female resistance trait is the suck behaviour, originally

documented in the free-living flatworm, *Macrostomum lignano* (Schärer *et al.*, 2004). Studies

in this reciprocally-mating simultaneous hermaphrodite have shown that matings are often

followed by the suck behaviour, during which the worm bends down and places its pharynx

over its own female genital opening (which is connected to the female antrum, the sperm-

receiving organ) and then appears to suck. The suck behaviour is hypothesised to be a

postcopulatory behaviour used for removing sperm or other ejaculate components received

during mating and thus to function as a female resistance trait (Vizoso *et al.*, 2010). However,

while there have been multiple studies on this behaviour (Schärer *et al.*, 2004, 2011, 2020;

Marie-Orleach *et al.*, 2013, 2017; Patlar *et al.*, 2020), there has to date been no direct

evidence for sperm and/or ejaculate actually being removed during the suck behaviour.

Moreover, if the suck functions as a postcopulatory sexual selection process, it could affect

the strength of sperm competition and potentially impact the optimal sex allocation (i.e. the

amount of resources allocated to the male and female function) (van Velzen *et al.*, 2009;

Schärer & Pen, 2013). Indeed, studies have documented both inter- and intra-specific

variation in sex allocation in *Macrostomum* (Singh *et al.*, 2020b; Brand *et al.*, 2022a; Singh &

Schärer, 2021), with mating behaviour predicting the evolution of a species' sex allocation

(Brand *et al.*, 2022a), but not the evolution of its sex allocation plasticity (Singh & Schärer,

2021).

Interestingly, *Macrostomum* species exhibit different combinations of reproductive

morphological traits that are likely associated with the reciprocal mating and hypodermic

insemination strategies (Figure 1A,B). Indeed, a previous study demonstrated an association

between certain male and female reproductive traits and the mating strategy in 16

Macrostomum species, naming the two alternative outcomes the reciprocal and hypodermic

mating syndrome, respectively (Schärer *et al.*, 2011). A more recent study has used a refined

composite measure, called the inferred mating syndrome, derived from the observation of

additional components of the reproductive morphology, in an attempt to classify 145

Macrostomum species as showing either the reciprocal or hypodermic inferred mating

syndrome, respectively (Figure 1B) (Brand *et al.*, 2022b). The lateral bristles on the sperm in

reciprocally mating species are hypothesized to represent a male persistence trait that allows

the sperm to remain anchored in the female antrum and not be pulled out during the suck

behaviour (Vizoso *et al.*, 2010), whereas the thick female antrum wall might prevent internal

injury resulting from the male genitalia during mating. In contrast, the sharp needle-like stylet

tip of hypodermically inseminating species likely allows sperm injection through the partner's

epidermis, while the simple sperm design presumably aides its movement through the

partner's body (Schärer *et al.*, 2011; Brand *et al.*, 2022b).

Although sexual conflict has been studied in many organisms spanning different reproductive

systems, studies on female resistance traits in hermaphrodites have been fewer, particularly in

a phylogenetic context (Koene & Schulenburg, 2005; Beese *et al.*, 2006, 2009; Anthes *et al.*,

2008; Sauer & Hausdorf, 2009; Schärer *et al.*, 2011; Brand *et al.*, 2022b). In our study, we

examine the evolution of the suck behaviour, aiming to document reproductive behaviour in a

total of 64 *Macrostomum* species. As a result of this, we, for the first time, provide

videographic evidence that ejaculate is indeed removed during the suck behaviour, supporting

the previously proposed hypothesis for the function of this postcopulatory behaviour (Vizoso

et al., 2010). Using this extensive behavioural data set, we examine correlations between

different aspects of the mating and suck behaviour, and between reproductive morphology

and the behavioural mating strategies, while accounting for the phylogenetic

interrelationships. If the suck behaviour has indeed evolved as a postcopulatory strategy, we

predict positive correlations between the presence, duration, and frequency of the mating

behaviour and the suck behaviour. This could occur, e.g., if longer/frequent matings lead to

more ejaculate being transferred, which would need longer/frequent sucks to remove the

ejaculate. We might also expect a trade-off between copulation duration and frequency, if

species that spend a lot of time in copulation cannot copulate that often, e.g. due to ejaculate

limitation or mating taking up a lot of the total time (so that fewer mating could be done over

a period, i.e., an autocorrelation). Similarly, if suck functions to remove ejaculate, we may

expect a trade-off between suck duration and frequency, if shorter sucks necessitate the need

for more frequent sucks to remove the ejaculate. Finally, we also expect the reproductive

morphology to be a good proxy for inferring the behavioural mating strategy as a result of

coevolution.

Materials and Methods

Study organisms

Species in the genus *Macrostomum* are small (~0.3 to 3.0 mm body length) aquatic free-living flatworms that are highly transparent, permitting detailed observations of internal structures (for the general morphology see Figure 1A,B). The sperm and eggs are produced in the paired testes and paired ovaries, respectively, with studies documenting inter- and intra-specific variation in both testis and ovary size across the genus (Singh *et al.*, 2020b; Brand *et al.*, 2022a; Singh & Schärer, 2021). The female antrum is located anterior to the male antrum, connected to the outside, respectively, via a female genital opening (also female genital pore or vagina) and the male genital opening (also male genital pore). The stylet (male intromittent organ) resides within the male antrum and it is proximately connected via the vesicula granulorum (not shown) to the seminal vesicle, which contains sperm to be transferred during mating. In both reciprocally-mating and hypodermically-inseminating species, the female antrum serves as the egg-laying organ, while in reciprocally-mating species it additionally serves to receive the stylet during mating and as the sperm-storage organ (Vizoso *et al.*, 2010; Schärer *et al.*, 2011).

We obtained multiple specimens for a large number of *Macrostomum* species, collected from a range of locations and habitats, using a variety of extraction techniques, which we report on in more detail as part of separate studies on the phylogenetic interrelationships (Brand *et al.*, 2022c) and reproductive character evolution in this genus (Brand *et al.*, 2022b). Briefly, most specimens were sampled directly from natural field sites, while some were sampled from artificial ponds, or from aquaria containing other study organisms, and they were generally observed within a few days of collection. Other specimens were obtained from short- and long-term laboratory cultures maintained either by our group or by colleagues.

Following Brand *et al.* (2022b), 38 of a total of 64 species *Macrostomum* included in the

current study were classified as exhibiting the reciprocal inferred mating syndrome, because

they had a blunt tip of the stylet (the male intromittent organ), and of these all but one had

received sperm in the antrum. A further 6 species with a sharp stylet were also classified as

exhibiting the reciprocal inferred mating syndrome because they had complex sperm with

lateral bristles and we observed sperm in the antrum (Figure 2). In contrast, 15 species were

classified as exhibiting the hypodermic inferred mating syndrome because allosperm was

exclusively found hypodermically. An additional 4 species without observation of received

sperm were classified as exhibiting the hypodermic inferred mating syndrome because they

had a simple female antrum (no thickening of the antrum wall and no visible cellular valve), a

sperm design with reduced or absent bristles and a sharp stylet tip. Finally, one species

(*Macrostomum sp.* 101) was classified as intermediate because received sperm was observed

both in the antrum and within the tissue (Figure 2).

Observation methodology

We aimed at documenting the mating behaviour of all 64 *Macrostomum* species, by placing

the worms in mating chambers (Schärer *et al.*, 2004). A mating chamber consisted of the

worms being placed between two microscope slides in small drops (i.e. either freshwater or

water with different salinity, depending on the collection habitat), with a certain number of

spacers (separating the slides), and sealed with pure white Vaseline (note that we generally

also placed 4-6 empty drops around, to reduce evaporation). We adjusted the spacer number

and drop volume depending on the size and number of worms in a drop, respectively.

Usually, for a pair of worms of the size of *M. lignano* (~1.5 mm body length), we used 2

spacers (each spacer being ~105 μm) and a drop size of ~3 μl. Movies were recorded when

specimens were available and therefore across several sampling campaigns. Consequently, the

recording setups differed (macro lenses, cameras or lighting conditions). However from a

previous detailed study of two *Macrostomum* species we know that these minor setup

differences are unlikely to bias our observations (Singh *et al.*, 2020a). Usually, the movies

were recorded in QuickTime Format using BTV Pro (http://www.bensoftware.com/) at 1

frame s^{-1}, but for some species we also generated detailed close-up movies, where worms

were manually tracked at higher magnifications under a compound microscope and filmed at

higher frame rates (see next section). All worms were visually checked for sexual maturity

(defined as having visible gonads or eggs), either before or after filming.

Detailed observation of mating and suck behaviour in Macrostomum hamatum

While earlier work documented sperm sticking out of the female antrum after the suck

behaviour in *M. lignano* (Schärer *et al.*, 2004, 2011), direct observations of ejaculate removal

have not been reported to date. Here we could document ejaculate removal in detailed close-

up movies of *M. hamatum*, possibly since field-collected specimens of this species appeared

to be more transparent than other species. This allowed us to clearly visualise the deposition

and subsequent removal of ejaculate during the mating and suck behaviour, respectively.

Specifically, we examine the mating behaviour of *M. hamatum*, collected on 27. July 2017

directly in front of the Tvärminne Zoological Station, Finland (N 59.84452, E 23.24986), in a

detailed close-up movie (Supplementary Movie S1). Note that while we describe and

illustrate only one such instance (in an extract from a longer movie), we also observed

ejaculate removal in other detailed close-up movies of *M. hamatum* that we also deposit

(https://doi.org/10.5281/zenodo.6354683), and these observations corroborated our finding as

described here.

Scoring of mating and suck behaviour across species

We scored the mating behaviours from the mating movies by visual frame-by-frame analysis

(Supplementary Table S1). A reciprocal mating was scored when the tail plates of two worms

were in ventral contact and intertwined, such that the female antrum was accessible to the

partner's stylet and vice-versa, which would allow reciprocal transfer of ejaculate. In most

species, the copulatory posture is accompanied by the pair being tightly interlinked (like two

interlocking Gs, see Figure 3), and thus similar to the mating behaviour originally described

for *M. lignano* (Schärer *et al.*, 2004). Note that in some species the mating posture can deviate

from that observed in *M. lignano* (see Supplementary Table S2), such as, for example, in

Macrostomum sp. 57 and *Macrostomum* sp. 61 (Supplementary Movie S2A,B). The mating

duration was measured from the frame when the tail plates were in ventral contact (and

usually tightly intertwined), to the frame where the tail plates were no longer attached to each

other. We defined behaviours as matings only if the pair was in the above-described posture

for at least 3 s. The suck duration was measured starting from the frame when an individual

placed its pharynx over its female genital opening, up to the frame where the pharynx

disengaged. Note that in some cases, individuals do not lie on their side while sucking (as

generally seen in *M. lignano*), which can sometimes make it more difficult to observe the

suck behaviour. For each replicate drop, we divided the total number of matings and sucks by

the number of worms and the movie duration to obtain a standardized value. We then

averaged the frequency and duration estimates across all replicate drops for each species to

obtain the species estimates of the respective behaviours.

While we also invested significant effort into observing hypodermic insemination (see Table

S1 and Results), we only saw some rare behavioural instances in a few species that could

possibly represent cases of hypodermic insemination, such as, for example, in

Macrostomum sp. 1 and *M. gabriellae* (Supplementary Movie S3A,B). Possible reasons for

not observing hypodermic insemination could be that in many species such matings occur

very rapidly or that they mate less frequently, possibly since they try to avoid sperm receipt

(Apelt, 1969; Michiels, 1998). Given that we could not confirm the presence of hypodermic

insemination, we scored species either as having reciprocal mating being present (when it was

observed) or absent (when it was not observed) (Figure 2). Note, however, that the absence of

observations of reciprocal mating does not necessarily imply the presence of hypodermic

insemination. Instead, it could also result from a reciprocally-mating species not mating under

laboratory conditions and/or from an overall low mating frequency of a species. Similarly,

while the presence of the suck behaviour can clearly be identified in many species, the

absence of observations of the suck behaviour does not necessarily mean that a species never

shows this behaviour.

Evolution of the mating and suck behaviour across the genus Macrostomum

To perform phylogenetic comparative analyses, we used a trimmed version of a recently

published ultrametric large-scale phylogeny of the genus *Macrostomum* (i.e. the C-IQ-TREE

phylogeny of Brand *et al.*, 2022c). This phylogeny is based on an amino acid alignment of

385 genes from 98 species, supplemented with Sanger sequences from a *28S rRNA* fragment,

which allowed the addition of a further 47 species, and calculated using a maximum

likelihood approach (Brand *et al.*, 2022c), covering all the species we included in the current

study. Specifically, we determined 1) whether the presence/absence of reciprocal mating is

correlated with the presence/absence of the suck behaviour, 2) whether the presence/absence

of reciprocal mating is correlated to the presence/absence of the reciprocal inferred mating

syndrome, and 3) whether there are correlations between the frequency and the duration of the

reciprocal mating and suck behaviours among the species that show these behaviours.

Presence/absence of reciprocal mating and the suck behaviour: We used the DISCRETE

model in BayesTraits V.3.0.1 to test for correlated evolution between reciprocal mating and

the suck behaviour (both scored as present/absent), using the Reversible Jump Markov Chain

Monte Carlo (RJ MCMC) approach (Pagel, 1994; Pagel & Meade, 2006; Meade & Pagel,

2016). Specifically, we compared the marginal likelihood of a dependent model, in which the

presence of suck depends on the presence of reciprocal mating, to an independent model, in

which the suck behaviour and reciprocal mating evolve independently. Each RJ MCMC chain

was run for twelve million iterations and the first one million iterations were discarded as

burn-in, after which the chain was sampled every 1000^{th} iteration. We used a gamma

hyperprior (gamma 0 1 0 1), and placed 1000 stepping stones (with each iterating 10000

times) to obtain the marginal likelihood values for the models. We ran three separate chains

each for the dependent and independent model to check for the stability of the likelihood

values and convergence. Using the R package coda (Plummer *et al.*, 2006), we confirmed that

the chains had converged (Gelman & Rubin, 1992; Brooks & Gelman, 1998) and that the

Effective Sample Size was >200 for all parameters. In addition, we also confirmed that the

acceptance rate was between 20-40% (Pagel & Meade, 2006). We compared the alternative

models with the Log Bayes Factor (BF), using the convention that BF values > 2 are

considered as positive support for the best-fit model, while values between 5-10 and > 10 are

considered as strong and very strong support for the model, respectively (Pagel & Meade,

2006). To examine the robustness, we repeated the analysis for a reduced dataset, by

excluding six species that had in total been observed for < 21 h (~10% quantile). For the

dependent models of the full dataset, we estimated the transition rates among the different

trait states by calculating Z values. This value can be understood as the percentage of times a

transition rate was set to zero, with a high value thus indicating that the transition between

two states is unlikely. We expect a correlation between the presence of reciprocal mating and

the presence of the suck behaviour, which would corroborate that the suck behaviour indeed

is a postcopulatory behaviour that is linked to reciprocal mating, rather than possibly serving

a function that is also present in species with hypodermic insemination.

Presence/absence of reciprocal mating and the reciprocal inferred mating syndrome: We

checked for an association between reciprocal mating (scored as present/absent) and the

inferred mating syndrome (scored as reciprocal/hypodermic), using the DISCRETE model in

BayesTraits V.3.0.1 (as above). One of the species, *Macrostomum* sp. 101, had a morphology

that was scored intermediate between reciprocal and hypodermic (Brand *et al.*, 2022b), but

since the discrete method in BayesTraits only allows binary trait states, we excluded this

species from this analysis. We expect a correlation between the presence of reciprocal mating

behaviour and the reciprocal inferred mating syndrome, which could indicate that behaviour

and morphology coevolve.

Correlations between the frequency and the duration of mating behaviours: In preparation for

phylogenetic correlation analyses we estimated the phylogenetic signal for the continuous

traits (i.e. the duration and frequency of both the reciprocal mating and the suck behaviour;

log-transformed for all the analyses) using Pagel's λ (Pagel, 1999; Revell, 2012). A λ value of

1 indicates a strong phylogenetic signal, while a value around 0 indicates no/low phylogenetic

signal (Pagel, 1999). We found phylogenetic signal that was significantly different from 0 for

the suck frequency ($\lambda=0.67$, $P=0.02$), the suck duration ($\lambda=0.76$, $P=0.005$), and the reciprocal

mating frequency ($\lambda=0.50$, $P=0.05$), but only marginally so for the mating duration ($\lambda=0.46$,

$P=0.06$). For each trait, we then fitted four different models of trait evolution, i.e. Brownian

motion, Ornstein–Uhlenbeck, Early-burst, and Lambda models (Harmon *et al.*, 2008). We

found that the Lambda model had the highest sample-size corrected Akaike Information

Criterion (AICc) weights (ω_l) (Supplementary Table S3), and this model was hence chosen

for further PGLS analysis.

For the species that exhibited both reciprocal mating and the suck behaviour, we then

investigated if there was a correlation between the frequency and duration of the reciprocal

mating and suck behaviours, using phylogenetic generalized least-squares (PGLS) regression

implemented in the *caper* package version 1.0.1 (Orme *et al.*, 2014). PGLS accounts for the

non-independence of the data by incorporating the phylogenetic relationships between species

into the error structure of the model. For each analysis using the frequency and duration of the

reciprocal mating and suck behaviours, the phylogenetic signal (Pagel's λ) was estimated

using the maximum likelihood approach. We examined the residuals of each model for

normality and homogeneity (Mundry, 2014). Additionally, we scrutinized for influential cases

(species) in each PGLS model, by excluding one species at a time from the data and rerunning

the analysis, and comparing the results obtained with the results for the entire dataset

(Mundry, 2014). And finally, we evaluated the robustness of our results by repeating the

PGLS for a reduced dataset, which excluded five species in which mating or suck had only

been observed in one replicate (note that this reduced dataset is different from the reduced

dataset used in the above BayesTraits analysis).

We performed our analysis in R, version 3.6.1 (R Core Team, 2019).

Results

Sperm deposition and removal during mating and suck behaviour in Macrostomum hamatum

The general anatomy of the reproductive organs of *M. hamatum* is similar to that of many other reciprocally-mating *Macrostomum* species (Figure 1A,B). In the detailed movie of *M. hamatum*, the worms are already interlinked in the reciprocal copulatory position at the beginning of the clip (Figure 3, Supplementary Movie S1), and we consider this as $t = 0$ s (hereafter we refer to the worm on the right as Orange and the worm on the left as Grey, respectively). At this timepoint "the tail plates touch each other ventrally in opposing directions, while the anterior ventral surface of each worm touches the posterior dorsal surface of the partner", as previously described for the copulatory position in *M. lignano* (Schärer *et al.*, 2004). Interestingly, in *M. hamatum* the copulatory position resembles a square with rounded corners, as opposed to *M. lignano,* where it is more circular. This may in part be due to a strikingly different position of the tail plate, which in *M. hamatum* stands at a 90° angle from the posterior body axis and appears to poke into the anterior ventral surface of the partner, leading to a dorsal bulge in both Orange and Grey.

Moreover, *M. hamatum* has a much more prominent erection (i.e., a translucent finger-like structure on the ventral tail plate, likely formed by the eversion of the muscular male antrum), which pokes into the posterior ventral surface of the partner in the region of the female genital opening (although it is unclear if the erection actually enters the partner). The stylet of Grey—while moving inside of the relatively stationary erection—then performs poking movements that are directed towards Orange's female antrum, initially without any transfer of ejaculate. At $t = 3\text{-}5$ s, the stylet of Grey is seen repeatedly poking against the dorsal side of Orange's female antrum wall, each time leading to a visible bulge on Orange's dorsal side. In some of

these frames one can see the sharp hook-shaped distal end of the stylet that is typical for

M. hamatum. Eventually, Grey begins to deposit ejaculate (seen as a visible darkening of

Orange's female antrum lumen starting at about t = 5s). During this process the seminal

vesicle of Grey empties (as seen at the base of the erection, see also drawing in Figure 3 at 0 s

for location), while the female antrum of Orange fills up with ejaculate over the next ~21 s

(see Figure 3 from 3.7 s). Note that we here mainly focus on the sperm transfer from Grey to

Orange, but in the meantime, Orange also pokes and eventually enters the female antrum of

Grey (t = 16-20 s) and sperm is also transferred from Orange to Grey (between t = 21-27 s),

although this is more difficult to follow in the movie.

At t = 28 s, Orange pushes out its female antrum region, places its pharynx over its female

genital opening, and then sucks. The received ejaculate can be seen leaving Orange's female

antrum (i.e. the visible darkening in the female antrum lumen moves towards the pharynx

between t = 29-30s). In total, the suck behaviour lasts for 7 s. Interestingly, during the suck

the stylet of Grey remains anchored in Orange's female genital opening (probably involving

the above-mentioned hook). At t = 52 s, the mating ends after a mating duration of ~64 s

(recall that the worms were already in copula at t = 0 s). At t = 56 s, only Grey is in frame and

the received ejaculate in its female antrum is clearly visible. It continues to have a small

erection despite the mating being over. At t = 78 s, Grey pushes its female antrum region out

and some sperm is ejected from the female antrum at t = 80 s, notably before the pharynx

makes contact (Figure 4, Supplementary Movie S1). At t = 81 s, Grey puts its pharynx over

its female genital opening and then sucks for 10 s. After the suck, some sperm can still be

seen sticking out of the female antrum (similar to *M. lignano*, Schärer *et al.*, 2004), especially

at 92 s, but most of the ejaculate has been removed from the female antrum. The female

antrum remains slightly everted and the erection somewhat visible until at least 108 s.

Evolution of the mating and suck behaviour across the genus Macrostomum

We observed a total of 2796 worms across 64 *Macrostomum* species, with a mean of 44

worms and 76.7 hours of observation time per species, for a total observation time of 4908

hours. Of the 64 species, 30 species exhibited reciprocal mating behaviour, 31 species

exhibited the suck behaviour, and 25 species exhibited both the reciprocal mating and suck

behaviour (Figure 2).

Presence/absence of reciprocal mating and the suck behaviour: We found very strong support

for the dependent model over the independent model of evolution for the correlation between

the presence of reciprocal mating and the presence of the suck behaviour, with all three runs

for each model providing highly consistent values (average marginal likelihood,

independent=-89.25, dependent=-83.37, BF: 11.75; see also Supplementary Table S4a). This

showed that the presence of reciprocal mating and the presence of the suck behaviour are

strongly correlated. And the result was robust to observation time, since excluding the 6

species that were observed for < 21 h gave similar results (Supplementary Table S4a).

The transitions from the absence of both the reciprocal mating and suck behaviour to the

presence of either of these traits were found to be the most unlikely, as is evident from the

low transition rates and the high Z values (Figure 5a). Interestingly, the other transitions,

including losing reciprocal mating or the suck behaviour from the state when they are both

present, are all similarly likely. This contrast suggests that once both reciprocal mating and

suck are lost or absent in a species, it is highly unlikely to regain either.

Presence/absence of reciprocal mating and the reciprocal inferred mating syndrome: There

was a clear correlation between the presence of reciprocal mating behaviour and the

reciprocal inferred mating syndrome, as evident from the strong support for the dependent

model over the independent model of evolution, with similar values for the three independent

runs of each model (average marginal likelihood, independent=-69.09, dependent=-65.60, BF:

6.99; see also Supplementary Table S4b), suggesting that the reproductive morphology of a

species can serve as a good proxy for its mating behaviour. And as before, our result was

robust, as the reduced dataset gave us similar results (Supplementary Table S4b).

Transitions from the presence of reciprocal mating and the presence of the reciprocal inferred

mating syndrome to the absence of either were moderately likely, while the converse

transitions were very likely (Figure 5b). Similarly, transitions from the absence of reciprocal

mating and the absence of the reciprocal inferred mating syndrome to the presence of either

were either unlikely or relatively unlikely, while the converse transitions were very likely.

Together this suggests that there is a strong association between reciprocal mating behaviour

and morphological traits characterizing the reciprocal inferred mating syndrome (and between

absence of reciprocal mating behaviour and the hypodermic inferred mating syndrome), such

that species are attracted to these states and evolve away from states where the morphology

and behaviour are mismatched.

Correlations between the frequency and the duration of mating behaviours: Among the

species that exhibited reciprocal mating (n=30), the average mating frequency was 0.84 hr^{-1}

(range: 0.02-7.82 hr^{-1}, Figure 6A) and the average mating duration was 283.7 s (range: 5.2-

4609 s, Figure 6B), with some sibling species showing fairly divergent values. Moreover,

among the species that showed the suck behaviour (n=31), the average suck frequency was

0.54 hr^{-1} (range: 0.01-3.7 hr^{-1}, Figure 6A) and the average suck duration was 9.6 s (range: 4.7-

16.1 s, Figure 6C).

In line with our predictions, we found significant positive relationships between both

reciprocal mating frequency and suck frequency (Figure 7A), and reciprocal mating duration

and suck duration (Figure 7B); while there was no significant relationship between reciprocal

mating frequency and reciprocal mating duration (Figure 7C), and suck frequency and suck

duration (Figure 7D). The reduced dataset also gave qualitatively similar results for all

analysis (Supplementary Table S5).

Discussion

Sexual conflict can give rise to antagonistic coevolution in all sexual systems (Charnov, 1979;

Bedhomme *et al.*, 2009). Here we documented the widespread occurrence of a putative

female resistance trait, the suck behaviour, in >30 species in the hermaphroditic flatworm

genus *Macrostomum*. Moreover, the direct observation of ejaculate removal in one species,

M. hamatum, corroborates the hypothesis that the suck functions as a female resistance trait to

remove received ejaculate (Schärer et al. 2004; Vizoso et al. 2010; Schärer et al. 2011), and

this interpretation is also supported by significant evolutionary correlations between different

aspects of reciprocal mating and suck behaviour. Finally, we could also show that the

reproductive morphology is a good proxy for inferring the mating strategy of a species,

presumably also as a result of coevolution. In the following we discuss these findings in more

detail.

Sperm deposition and removal during mating and suck behaviour in Macrostomum

hamatum

While multiple studies in *Macrostomum* have examined aspects of the suck behaviour

(Schärer *et al.*, 2004, 2011, 2020; Marie-Orleach *et al.*, 2013, 2017; Patlar *et al.*, 2020; Singh

et al., 2020a), its involvement in removing received ejaculate components has so far only

been hypothesized. Our detailed observations of mating interactions in *M. hamatum* provide

the first direct evidence that ejaculate is indeed removed during this postcopulatory

behaviour. Interestingly, compared to *M. lignano* (Schärer *et al.*, 2004), *M. hamatum* has a

more rectangular mating posture (possibly due to the angular position of the tail plate), a

larger erection around the stylet, and the worms prominently evert the female antrum just

before the suck behaviour, likely as a result of muscular contractions. This could result from

differences in the female antrum morphology: while *M. hamatum* has a strong musculature

and an inner second chamber connecting to the main female antrum (Luther, 1947),

M. lignano has a somewhat simpler female antrum with a single chamber (Ladurner *et al.*,

2005; Vizoso *et al.*, 2010). Similarly, the prominent erection of the male antrum could result

from a muscular morphology that is similar to the muscular cirrus seen in species of the sister

genus, *Psammomacrostomum* (Ax, 1966; Janssen *et al.*, 2015). The combination of a rather

prominent female antrum and the relatively transparent specimens may have helped us

visualise the function of the suck behaviour better in *M. hamatum* than in other *Macrostomum*

species observed to date.

While we see ejaculate being removed during the suck behaviour, we cannot clearly

determine whether it is ingested. Although sperm digestion is widespread in hermaphrodites

(Charnov, 1979; Baur, 1998; Dillen *et al.*, 2009; Koene *et al.*, 2009), it usually occurs inside

an organ connected to the individual's reproductive system, unlike in the case of the suck

behaviour. To our knowledge, there have been only two earlier reports of sperm being orally

taken up in hermaphrodites, one in the arrow worm *Spadella cephaloptera* (John, 1933) and

the other in the leech *Placobdella parasitica* (Myers, 1935). Thus, the suck behaviour seems

to be a novel trait, which to date has only been observed in species of the Macrostomidae

(including a member of the sister genus *Psammomacrostomum*; P. Singh, pers. obs.). Similar

to the suck behaviour, females of the ladybird beetle, *Adalia bipunctata*, consume a

spermatophore after mating (Perry & Rowe, 2008). Moreover, there is also sperm dumping in

many separate-sexed species, in which the female physically ejects received sperm from her

reproductive tract, and this, at least in some cases, is thought to be a mechanism of cryptic

female choice (Snook & Hosken, 2004; Peretti & Eberhard, 2010; Firman *et al.*, 2017). If the

suck behaviour also functioned in cryptic female choice, we might expect individuals to

remove or retain sperm of certain partners more frequently (e.g. Pizzari & Birkhead, 2000).

This has also been observed in *M. lignano*, where the propensity of the recipient to suck is

affected by the mating status (Marie-Orleach *et al.*, 2013) and the genotype of its partners

(Marie-Orleach *et al.*, 2017). However, it is difficult to ascertain whether the suck implies an

active choice by the recipient or whether it is sometimes prevented as a result of a

manipulation by the donor (Patlar *et al.*, 2020). Moreover, our study documents in detail the

reciprocal transfer and deposition of sperm by both mating individuals during a reciprocal

mating in *Macrostomum* (but see Ax & Borkott, 1968 which documents mating and unilateral

sperm transfer in *M. salinum*, now considered to be *M. romanicum*).

Evolution of the mating and suck behaviour across the genus Macrostomum

We found a significant evolutionary correlation between the presence of reciprocal mating

and the suck behaviour (Figure 5a). In reciprocally-mating species, ejaculate is deposited in

the female antrum allowing its removal during the suck behaviour, while in hypodermically-

inseminating species, sperm is injected potentially anywhere in the body (Schärer *et al.*, 2011;

Brand *et al.*, 2022b). Given that the function of the suck behaviour indeed appears to be the

removal of ejaculate, we do not expect to see the suck behaviour in hypodermically-

inseminating species. Performing a suck at a site of hypodermic insemination might not

permit effective ejaculate removal (particularly also given the above-mentioned active

participation of the female antrum musculature), but instead would more likely lead to

additional tissue damage. Interestingly, the transition rates showed that while it is unlikely for

a species that lacks both the reciprocal mating and suck behaviour to gain either of these

traits, the loss of either reciprocal mating or the suck behaviour was estimated as being more

likely. These transitions could represent transitional steps towards hypodermic insemination,

which might arise as a means to bypass the female control and allow access to the eggs

(Charnov, 1979; Brand *et al.*, 2022b). Moreover, this interpretation is also supported by the

finding that there are multiple origins of hypodermic insemination in the genus *Macrostomum*

(Brand *et al.*, 2022b; Singh & Schärer, 2021). There are at least nine independent shifts from

reciprocal mating to hypodermic insemination in *Macrostomum*, while no transition is

observed in the converse direction (Brand *et al.*, 2022b).

However, it is important to point out that some of these findings could also have resulted from

a lack of observations of either the reciprocal mating and/or suck behaviour (despite being

present in a species), leading to an overestimation of these transition rates. Specifically, there

were six species that showed only the reciprocal mating behaviour and five species that

showed only the suck behaviour (Supplementary Table S1). These mismatches usually

appeared in species for which we had comparatively few observation hours (for more detail

see Supplementary Figure S3), suggesting that additional observations could help to further

ascertain the actual presence/absence of reciprocal mating or the suck behaviour, respectively.

Moreover, mismatches could result from a species not exhibiting some behaviours under our

laboratory conditions, or they might indicate that a species indeed lacks a behaviour. In

addition, if a species mates only rarely, individuals might be less inclined to remove the

sperm they receive, and in our study the species that showed reciprocal mating but did not

suck, had low or intermediate mating frequencies (see *Macrostomum* sp. 43, *Macrostomum*

sp. 67, *M. distinguendum, M. gieysztori,* and *M. poznaniense* in Figure 6A). Alternatively,

species might actually lack reciprocal mating, but losing a resistance trait like the suck

behaviour might take longer, particularly if the suck behaviour does not impose costs on the

fecundity. Moreover, the suck behaviour could have additional functions, such as possibly

removing egg material that remains in the antrum after egg laying. Species are predicted to

lose defensive or resistance traits only after the persistence traits have become substantially

less harmful, leading to a time lag (Parker, 1979). A study on the seed beetle, *Callosobruchus*

maculatus, showed that, while large males evolved relatively reduced length of genital spines

under monogamy, there was no detectable evolution in female genitalia within the same time

period (Cayetano *et al.*, 2011). And finally, since the worms we observed may often have

mated before we placed them into the mating chambers, some of the observed sucks might

have occurred in response to unobserved earlier matings, since sucks do not only occur

immediately after mating (Schärer et al. 2004).

The significant evolutionary correlation between the presence of reciprocal mating and the

purely morphologically-derived reciprocal inferred mating syndrome (Figure 5b) confirms

previous findings (Schärer *et al.*, 2011). It shows that persistence and resistance are not

generally limited to single traits, but are often composite suites of behavioural, morphological

and physiological traits acting together (Arnqvist & Rowe, 2005). For example, the thickened

female antrum wall and the suck behaviour might be different components of female

resistance. While the former might prevent injury resulting from the male genitalia when

mating reciprocally, the suck behaviour serves to remove unwanted ejaculate received during

mating. Similar adaptations of the female reproductive tract are also seen in the seed beetle

C. maculatus, where a thicker female tract lining serves as a resistance trait against harm by

male genitalia (Dougherty *et al.*, 2017). Moreover, resistance and persistence traits can also

occur at the proteomic level. A study in *M. lignano* identified two seminal fluid transcripts,

experimental knock-down of which caused mating partners to suck more often (Patlar *et al.*,

2020). This suggests that the seminal fluid proteins derived from these transcripts might be

counter adaptations by the donor to prevent the suck behaviour by the recipient.

In our dataset, there was one species each that exhibited the hypodermic inferred mating

syndrome morphology and showed both reciprocal mating and the suck behaviour

(*M. rostratum*), only reciprocal mating (*M. distinguendum*), or only the suck behaviour

(*M. finlandense*) (Supplementary Table S1). Interestingly, the three species represent at least

two, but possibly three, of the above-mentioned multiple independent origins of the

hypodermic inferred mating syndrome (Brand *et al.*, 2022b). Conversely, there were 12

species that exhibited the reciprocal inferred mating syndrome, but in which neither reciprocal

mating, nor the suck behaviour was observed. As above, this mismatch occurred mainly in

species for which we had relatively few observation hours (Supplementary Figure S3),

suggesting that, if these species have a low mating frequency, then more observation time

may be needed to avoid falsely inferring the absence of the mating and suck behaviour. And

finally, for many of the species that showed the hypodermic inferred mating syndrome we had

considerable amounts of observation hours (Supplementary Figure S3), so that it seems

unlikely that the absence of mating and suck observations in these species were due to a lack

of effort.

Macrostomum species showed large interspecific variation in behaviour, with a nearly 900-

fold variation in mating duration, a 3-fold variation in the suck duration, and a nearly 400-fold

variation in the mating and suck frequency across the genus (Figure 6). Remarkably, despite

this extensive interspecific variation in behavioural traits, we see clear correlations between

both the mating and suck duration, as well as the mating and suck frequency, suggesting that

the mating and suck behaviour have coevolved. If a longer mating duration or more frequent

mating implies more sperm transfer, then we expect selection for a longer suck duration

and/or a more frequent suck behaviour (particularly if ejaculate receipt is associated with

fitness costs). In some species, at least, a longer mating duration does imply more ejaculate

transferred (Engqvist & Sauer, 2003), and is often used as a proxy for ejaculate size (Kelly &

Jennions, 2011). Alternatively, such a correlation could also emerge as a result of variation in

genital complexity, e.g. if it takes longer to insert and remove more complex male genitalia,

and to suck out ejaculate from more complex female antra. Interestingly in *Macrostomum*,

male and female genital complexity are indeed correlated (Brand *et al.*, 2022b). Moreover, a

positive correlation between reciprocal mating and suck could also appear, if some species do

not do well under our laboratory conditions, leading to an overall low behaviour frequency.

Note, however, that we confirmed that the individuals we used for mating movies were adults

with visible testes and ovaries, and we also established the robustness of the observed

correlations by excluding species in which mating or suck had only been observed in one

replicate (Supplementary Table S5). We did not find any correlations between the frequency

and duration for either reciprocal mating or suck. While we might have expected mating

duration to trade off with mating frequency, mating duration only made up a relatively small

percentage of total tine, potentially posing no trade-off. Similarly, if sucking is not very

costly, the suck duration and frequency may not trade-off; and could even be positively

correlated, since both help to remove ejaculate.

Finally, mating frequency (and possibly mating duration) could be positively correlated with

allocation towards the male function (e.g., testes). Indeed, studies in *Macrostomum* have

shown interspecific variation in sex allocation towards the male and female functions, such as

testes and ovaries (Singh *et al.*, 2020b; Brand *et al.*, 2022a; Singh & Schärer, 2021). This

interspecific variation could potentially relate to the mating behaviour, as we can expect

species that have a longer mating duration or higher mating frequency to have larger testes, if

longer and/or more frequent mating implies that more sperm are transferred (Janicke &

Schärer, 2009). Mating duration could also correlate with the complexity of genitalia, such

that more complex genitalia might require longer mating duration (King *et al.*, 2009), and

future studies should investigate the correlations between different aspects of reproductive

behaviour and reproductive morphology in *Macrostomum*.

Conclusions

Our study provides direct observational evidence for ejaculate removal during the

postcopulatory suck behaviour in the species *M. hamatum*, compelling support for the

coevolution between the reciprocal mating and suck behaviour, and detailed information in a

phylogenetic context on the occurrence and interspecific variation of the suck behaviour.

Moreover, we show that reproductive morphology can be a good proxy to infer the

behavioural mating strategy. Taken together our study shows the presence of a postcopulatory

female behavioural resistance trait that co-evolves with mating strategy and allows

manipulation of received ejaculate in a simultaneously hermaphroditic sexual system. Thus,

our study adds to the repertoire of information on traits involved in sexual conflict in

Macrostomum genus and demonstrates the genus as an excellent model system for

understanding sexual antagonistic coevolution by allowing us to examine the evolution of

diverse female resistance and male persistence traits, spanning behavioural and morphological

traits, simultaneously.

Acknowledgements

We would like to thank Gudrun Viktorin, Lukas Zimmermann, Jürgen Hottinger, Urs Stiefel

and Daniel Lüscher for technical support. The help from many people that have been involved

in the collection and maintenance of the worms we have used here is acknowledged in the

study that presents the large-scale *Macrostomum* phylogeny, but here we here would like to

specifically thank Philipp Kaufmann, Gregor Schulte, Dita Vizoso, and Michaela Zwyer for

contributing several of the analysed mating movies. This research was supported by grants

31003A_162543 and 310030_184916 of the Swiss National Science Foundation (SNSF).

References

Abbott, J.K. (2011) Intra-locus sexual conflict and sexually antagonistic genetic variation in
hermaphroditic animals. *Proceedings of the Royal Society B: Biological Sciences*, **278**,
161–169.
Anthes, N., Schulenburg, H. & Michiels, N.K. (2008) Evolutionary links between
reproductive morphology, ecology and mating behavior in opisthobranch gastropods.
Evolution, **62**, 900–916.
Apelt, G. (1969) Fortpflanzungsbiologie, entwicklungszyklen und vergleichende
frühentwicklung acoeler Turbellarien. *Marine Biology*, **4**, 267-325.

Arbuthnott, D., Crespi, B.J. & Schwander, T. (2015) Female stick insects mate multiply to
find compatible mates. *The American Naturalist*, **186**, 519–530.
Arnqvist, G. & Rowe, L. (2005) *Sexual conflict*. Princeton University Press, Princeton, NJ.
Ax, P. (1966) Die bedeutung der interstitiellen sandfauna für allgemeine probleme der
systematik, ökologie und biologie. *Verhandlungen des Institutes für Meeresforschung*
Bremerhaven, **Sonderband**, 15–65.
Ax, P. & Borkott, H. (1968) Organisation und fortpflanzung von *Macrostomum romanicum*
(Turbellaria, Macrostomida). *Verh Dtsch Zool Ges Innsbruck*, **30**, 344–347.
Bateman, A.J. (1948) Intra-sexual selection in *Drosophila*. *Heredity*, **2**, 349–368.
Baur, B. (1998) Sperm competition in molluscs. In *Sperm Competition and Sexual Selection*.
Bedhomme, S., Bernasconi, G., Koene, J.M., Lankinen, Å., Arathi, H.S., Michiels, N.K., *et*
al. (2009) How does breeding system variation modulate sexual antagonism? *Biology*
Letters, **5**, 717–720.
Beese, K., Armbruster, G.F.J., Beier, K. & Baur, B. (2009) Evolution of female sperm-storage
organs in the carrefour of stylommatophoran gastropods. *Journal of Zoological*
Systematics and Evolutionary Research, **47**, 49–60.
Beese, K., Beier, K. & Baur, B. (2006) Coevolution of male and female reproductive traits in
a simultaneously hermaphroditic land snail. *Journal of Evolutionary Biology*, **19**, 410–
418.
Brand, J.N., Harmon, L.J. & Schärer, L. (2022a) Mating behavior and reproductive
morphology predict macroevolution of sex allocation in hermaphroditic flatworms.
BMC Biology, **20**, 35.
Brand, J.N., Harmon, L.J. & Schärer, L. (2022b) Frequent origins of traumatic insemination
involve convergent shifts in sperm and genital morphology. *Evolution Letters*, **20**,
evl3.268.
Brand, J.N., Viktorin, G., Wiberg, R.A.W., Beisel, C. & Schärer, L. (2022c) Large-scale
phylogenomics of the genus *Macrostomum* (Platyhelminthes) reveals cryptic diversity
and novel sexual traits. *Molecular Phylogenetics and Evolution*, **166**, 107296.
Brooks, S.P. & Gelman, A. (1998) General methods for monitoring convergence of iterative
simulations. *Journal of Computational and Graphical Statistics*, **7**, 434–455.
Cayetano, L., Maklakov, A.A., Brooks, R.C. & Bonduriansky, R. (2011) Evolution of male
and female genitalia following release from sexual selection. *Evolution*, **65**, 2171–
2183.
Charnov, E.L. (1979) Simultaneous hermaphroditism and sexual selection. *Proceedings of the*
National Academy of Sciences, **76**, 2480–2484.
Dillen, L., Jordaens, K. & Backeljau, T. (2009) Sperm transfer, sperm storage, and sperm
digestion in the hermaphroditic land snail *Succinea putris* (Gastropoda, Pulmonata).
Invertebrate Biology, **128**, 97–106.
Dougherty, L.R., Lieshout, E. van, McNamara, K.B., Moschilla, J.A., Arnqvist, G. &
Simmons, L.W. (2017) Sexual conflict and correlated evolution between male
persistence and female resistance traits in the seed beetle *Callosobruchus maculatus*.
Proceedings of the Royal Society B: Biological Sciences, **284**, 20170132.
Eberhard, W. (1996) *Female control: sexual selection by cryptic female choice*. Princeton
University Press.
Engqvist, L. & Sauer, K.P. (2003) Determinants of sperm transfer in the scorpionfly *Panorpa*
cognata: male variation, female condition and copulation duration. *Journal of*
Evolutionary Biology, **16**, 1196–1204.
Firman, R.C., Gasparini, C., Manier, M.K. & Pizzari, T. (2017) Postmating female control: 20
years of cryptic female choice. *Trends in Ecology & Evolution*, **32**, 368–382.

Frank, S.A. (2000) Sperm competition and female avoidance of polyspermy mediated by sperm-egg biochemistry. *Evolutionary Ecology Research*, **2**, 613–625.

Gelman, A. & Rubin, D.B. (1992) Inference from iterative simulation using multiple sequences. *Statistical Science*, **7**, 457–472.

Harmon, L.J., Weir, J.T., Brock, C.D., Glor, R.E. & Challenger, W. (2008) GEIGER: investigating evolutionary radiations. *Bioinformatics*, **24**, 129–131.

Hemmings, N. & Birkhead, T. (2017) Differential sperm storage by female zebra finches *Taeniopygia guttata*. *Proceedings of the Royal Society B: Biological Sciences*.

Holland, B. & Rice, W.R. (1998) Perspective: chase‐away sexual selection: antagonistic seduction versus resistance. *Evolution*, **52**, 1–7.

Janicke, T. & Schärer, L. (2009) Sex allocation predicts mating rate in a simultaneous hermaphrodite. *Proceedings of the Royal Society B: Biological Sciences*, **276**, 4247–4253.

Janssen, T., Vizoso, D.B., Schulte, G., Littlewood, D.T.J., Waeschenbach, A. & Schärer, L. (2015) The first multi-gene phylogeny of the *Macrostomorpha* sheds light on the evolution of sexual and asexual reproduction in basal Platyhelminthes. *Molecular Phylogenetics and Evolution*, **92**, 82–107.

Jennions, M.D. & Petrie, M. (2007) Why do females mate multiply? A review of the genetic benefits. *Biological Reviews*, **75**, 21–64.

John, C. (1933) Habits, structure and development of *Spadella cephaloptera*. *Q J Microsc Sci*, **75**, 625–696.

Kelly, C.D. & Jennions, M.D. (2011) Sexual selection and sperm quantity: Meta-analyses of strategic ejaculation. *Biological Reviews*, **86**, 863–884.

King, R.B., Jadin, R.C., Grue, M. & Walley, H.D. (2009) Behavioural correlates with hemipenis morphology in New World natricine snakes. *Biological Journal of the Linnean Society*, **98**, 110–120.

Koene, J.M., Montagne-Wajer, K., Roelofs, D. & Maat, A. Ter. (2009) The fate of received sperm in the reproductive tract of a hermaphroditic snail and its implications for fertilisation. *Evolutionary Ecology*, **23**, 533–543.

Koene, J.M. & Schulenburg, H. (2005) Shooting darts: co-evolution and counter-adaptation in hermaphroditic snails. *BMC Evolutionary Biology*, **5**, 25.

Kokko, H. & Mappes, J. (2013) Multiple mating by females is a natural outcome of a null model of mate encounters. *Entomologia Experimentalis et Applicata*, **146**, 26–37.

Ladurner, P., Schärer, L., Salvenmoser, W. & Rieger, R.M. (2005) A new model organism among the lower Bilateria and the use of digital microscopy in taxonomy of meiobenthic Platyhelminthes: *Macrostomum lignano*, n. sp. (Rhabditophora, Macrostomorpha). *Journal of Zoological Systematics and Evolutionary Research*, **43**, 114–126.

Lange, R., Reinhardt, K., Michiels, N.K. & Anthes, N. (2013) Functions, diversity, and evolution of traumatic mating. *Biological Reviews*, **88**, 585–601.

Lehtonen, J., Parker, G.A. & Schärer, L. (2016) Why anisogamy drives ancestral sex roles: Brief Communication. *Evolution*, **70**, 1129–1135.

Leonard, J.L. (1991) Sexual conflict and the mating systems of simultaneously hermaphroditic gastropods. *American Malacological Bulletin*, **9**, 45–58.

Luther, A. (1947) Untersuchungen an rhabdocoelen Turbellarien VI. Macrostomiden aus Finnland. *Acta Zoologlca Fennica*, **49**, 1–38.

Marie-Orleach, L., Janicke, T. & Schärer, L. (2013) Effects of mating status on copulatory and postcopulatory behaviour in a simultaneous hermaphrodite. *Animal Behaviour*, **85**, 453–461.

746   Marie-Orleach, L., Vogt-Burri, N., Mouginot, P., Schlatter, A., Vizoso, D.B., Bailey, N.W., *et*
747   *al.* (2017) Indirect genetic effects and sexual conflicts: Partner genotype influences
748   multiple morphological and behavioral reproductive traits in a flatworm. *Evolution*,
749   **71**, 1232–1245.
750   Meade, A. & Pagel, M. (2016) Bayes Traits. http://www.evolution.rdg.ac.uk/BayesTraitsV3
751   Michiels, N.K. (1998) Mating conflicts and sperm competition in simultaneous
752   hermaphrodites. In *Sperm Competition and Sexual Selection*. Elsevier, pp. 219–254.
753   Mundry, R. (2014) Statistical issues and assumptions of phylogenetic generalized least
754   squares. In *Modern Phylogenetic Comparative Methods and their Application in*
755   *Evolutionary Biology*.
756   Myers, R.J. (1935) Behavior and morphological changes in the leech, Placobdella parasitica,
757   during hypodermic insemination. *Journal of Morphology*, **57**, 617–653.
758   Orme, D., Freckleton, R., Thomas, G., Petzoldt, T., Fritz, S., Isaac, N., *et al.* (2014) Caper:
759   Comparative analyses of phylogenetics and evolution in R. *R package version 0.5.2/*
760   *r121*.
761   Pagel, M. (1994) Detecting correlated evolution on phylogenies: a general method for the
762   comparative analysis of discrete characters. *Proceedings of the Royal Society of*
763   *London. Series B: Biological Sciences*, **255**, 37–45.
764   Pagel, M. (1999) Inferring the historical patterns of biological evolution. *Nature*, **401**, 877–
765   884.
766   Pagel, M. & Meade, A. (2006) Bayesian analysis of correlated evolution of discrete characters
767   by reversible‐jump markov chain monte carlo. *The American Naturalist*, **167**, 808–
768   825.
769   Parker, G.A. (1979) Sexual selection and sexual conflict. In *Sexual Selection and*
770   *Reproductive Competition in Insects*. Elsevier, pp. 123–166.
771   Parker, G.A. (2011) The origin and maintenance of two sexes (anisogamy), and their gamete
772   sizes by gamete competition. In *The Evolution of Anisogamy* (ed. by Togashi, T. &
773   Cox, P.A.). Cambridge University Press, Cambridge, pp. 17–74.
774   Patlar, B., Weber, M., Temizyürek, T. & Ramm, S.A. (2020) Seminal fluid-mediated
775   manipulation of post-mating behavior in a simultaneous hermaphrodite. *Current*
776   *Biology*, **30**, 143-149.e4.
777   Peretti, A. V. & Eberhard, W.G. (2010) Cryptic female choice via sperm dumping favours
778   male copulatory courtship in a spider. *Journal of Evolutionary Biology*, **23**, 271–281.
779   Perry, J.C. & Rowe, L. (2008) Ingested spermatophores accelerate reproduction and increase
780   mating resistance but are not a source of sexual conflict. *Animal Behaviour*.
781   Pizzari, T. & Birkhead, T.R. (2000) Female feral fowl eject sperm of subdominant males.
782   *Nature*, **405**, 787–789.
783   Plummer, M., Best, N., Cowles, K. & Vines, K. (2006) {CODA}: Convergence Diagnosis
784   and Output Analysis for {MCMC}. *R News*.
785   R Core Team. (2019) R: A language and environment for statistical computing.
786   http://www.R-project.org/. [WWW Document]. *R Foundation for Statistical*
787   *Computing, Vienna, Austria*. URL [accessed on 2019].
788   Reinhardt, K., Anthes, N. & Lange, R. (2015) Copulatory wounding and traumatic
789   insemination. *Cold Spring Harbor Perspectives in Biology*, **7**, a017582.
790   Revell, L.J. (2012) phytools: an R package for phylogenetic comparative biology (and other
791   things). *Methods in Ecology and Evolution*, **3**, 217–223.
792   Sauer, J. & Hausdorf, B. (2009) Sexual selection is involved in speciation in a land snail
793   radiation on Crete. *Evolution*, **63**, 2535–2546.
794   Schärer, L., Brand, J.N., Singh, P., Zadesenets, K.S., Stelzer, C.-P. & Viktorin, G. (2020) A
795   phylogenetically informed search for an alternative *Macrostomum* model species, with

notes on taxonomy, mating behavior, karyology, and genome size. *Journal of*
Zoological Systematics and Evolutionary Research, **58**, 41–65.

Schärer, L., Janicke, T. & Ramm, S.A. (2015) Sexual conflict in hermaphrodites. *Cold Spring*
Harbor Perspectives in Biology, **7**, a017673.

Schärer, L., Joss, G. & Sandner, P. (2004) Mating behaviour of the marine turbellarian
Macrostomum sp.: these worms suck. *Marine Biology*, **145**, 373–380.

Schärer, L., Littlewood, D.T.J., Waeschenbach, A., Yoshida, W. & Vizoso, D.B. (2011)
Mating behavior and the evolution of sperm design. *Proceedings of the National*
Academy of Sciences, **108**, 1490–1495.

Schärer, L. & Pen, I. (2013) Sex allocation and investment into pre- and post-copulatory traits
in simultaneous hermaphrodites: the role of polyandry and local sperm competition.
Philosophical Transactions of the Royal Society B: Biological Sciences, **368**,
20120052.

Singh, P., Ballmer, D.N., Laubscher, M. & Schärer, L. (2020a) Successful mating and
hybridisation in two closely related flatworm species despite significant differences in
reproductive morphology and behaviour. *Scientific Reports*, **10**, 12830.

Singh, P. & Schärer, L. (2021) Self-fertilization, but not mating strategy, predicts the
evolution of sex allocation plasticity in a hermaphroditic flatworm genus. BioRxiv.
https://doi.org/10.1101/2020.06.12.149351

Singh, P., Vellnow, N. & Schärer, L. (2020b) Variation in sex allocation plasticity in three
closely related flatworm species. *Ecology and Evolution*, **10**, 26–37.

Snook, R.R. & Hosken, D.J. (2004) Sperm death and dumping in *Drosophila*. *Nature*, **428**,
939–941.

Velzen, E. van, Schärer, L. & Pen, I. (2009) The effect of cryptic female choice on sex
allocation in simultaneous hermaphrodites. *Proceedings of the Royal Society B:*
Biological Sciences, **276**, 3123–3131.

Vizoso, D.B., Rieger, G. & Schärer, L. (2010) Goings-on inside a worm: Functional
hypotheses derived from sexual conflict thinking. *Biological Journal of the Linnean*
Society, **99**, 370–383.

Zhang, S., Shi, Y., Zeng, Z., Xin, F., Deng, L. & Wang, A. (2021) Two new brackish-water
species of *Macrostomum* (Platyhelminthes: Macrostomorpha) from China and their
phylogenetic positions. *Zoological Science*, **38**.

Figures

Figure 1. (A) Photograph and line drawing of an adult *Macrostomum cliftonense* (previously

M. cliftonensis, name updated following Zhang *et al.*, 2021), showing some of the

components of the reproductive system to help understand the mating behaviour observations

(total length ~1.2 mm). (B) Schematic drawings of the typical morphology of the antrum

(female reproductive organ), sperm, and stylet (male intromittent organ) of *Macrostomum*

species with reciprocal mating (i.e. complex antrum and sperm, and stylet with a blunt distal

end) and hypodermic mating (i.e. simple antrum and sperm, and a needle-like stylet) (see also

Brand *et al.*, 2022b).

Figure 2. Presence (green) or absence (yellow) of reciprocal mating, the suck behaviour, and

the reciprocal inferred mating syndrome across the *Macrostomum* phylogeny (for a total of 64

Macrostomum species, see Brand *et al.*, 2022c for full phylogeny). Note that for the

behaviourally-inferred traits an absence may be due to a lack of sufficient data for observing

the behaviour, and that for the reciprocal inferred mating syndrome the absence represents the

hypodermic inferred mating syndrome (except for *Macrostomum* sp. 101, which showed an

intermediate inferred mating syndrome, grey). Branch supports are indicated by ultrafast

bootstrap (first number) and approximate likelihood ratio tests (second number), respectively

(from Brand *et al.*, 2022c).

Figure 3. Reciprocal mating followed by a postcopulatory suck in *Macrostomum hamatum*,

including ejaculate deposition by Grey (the worm on the left at 0 s) and its subsequent

removal during the suck behaviour by Orange (the worm on the right at 0 s). Before transfer,

the sperm is stored in the seminal vesicle of Grey (blue arrow in first frame), which is

connected to its stylet. Ejaculate (dark mass indicated by red arrow) can be seen being

deposited by Grey from the seminal vesicle starting from 3.7 s in the female antrum of

Orange, followed by Orange pushing its female antrum region out (at 27.4 s) and sucking

(note that Orange is also depositing ejaculate in Grey from 23.3 s). There is a visible

reduction in the quantity of received ejaculate in the female antrum of Orange after the suck

ends. Note that we call the frame from where we start describing the movie as $t = 0$ s, but the

mating had already started before that timepoint. In some frames, parts of the worms are not

visible on the video, and the presumed outlines are drawn using stippled lines. A high-

resolution version is provided in Supplementary Figure S1A.

Figure 4. A postcopulatory suck following the reciprocal mating shown in Figure 3 (a

continuation of the same movie), performed by Grey (i.e. the individual that had not yet

sucked). Grey completely pushes out its female antrum region at t = 78 s (which leads to

some sperm appearing near the female genital opening at t = 78.5 s), puts its pharynx over the

female genital opening, and then sucks out most of the previously deposited ejaculate over a

period of 10 s (from t = 81 s). Moreover, some sperm can be seen sticking out of the female

genital opening after the suck ends. In some frames, part of the worm is not visible on the

video, and the presumed outline is thus drawn using stippled lines. A high-resolution version

is provided in Supplementary Figure S1B.

Figure 5. Correlated evolution of behavioural character states. The panels show the transition

rates and the Z values (in brackets, expressed as %) for transitions between (A) the presence

or absence of reciprocal mating and the suck behaviour (crossed out when absent), and (B) the

presence or absence of reciprocal mating (crossed out when absent) and the inferred mating

syndrome (from Brand *et al.*, 2022c). For the transition rates, the mean of the posterior

distributions across all runs is given. The Z value can be understood as the percentage of

times the transition rate was set to zero, amongst all the sampled parameters. The different

arrows represent different probabilities of transitions between the states: high probability

(strong black arrows, Z value < 15%), moderate probability (thin black arrows, Z value 20-

55%), and low probability (dashed black arrows, Z value > 85%). The posterior distributions

of the transition rate parameters are given in Supplementary Figure S2.

Figure 6. Trimmed phylogeny of the 36 *Macrostomum* species that showed reciprocal mating and/or the suck behaviour alongside data on means

and standard errors of (A) reciprocal mating and suck frequency, (B) reciprocal mating duration (log-transformed), and (C) suck duration (log-

transformed). Note that some species exhibited either only reciprocal mating or only the suck behaviour. Also note that for the species in which a

896   behaviour had been observed in only 1 replicate, we report only that single value. The branch support values are indicated by ultrafast bootstrap

897   (first number) and approximate likelihood ratio tests (second number), respectively (from Brand *et al.*, 2022c).

Figure 7. Relationships between (A) reciprocal mating frequency and suck frequency, (B) reciprocal mating duration and suck duration, (C) reciprocal mating duration and frequency, and (D) suck duration and frequency for *Macrostomum* species. Note that (A-D) show values plotted on log-transformed axes with PGLS results.