CRISPR screens reveal genetic determinants of PARP inhibitor sensitivity and resistance in prostate cancer

Takuya Tsujino[1,2,11], Tomoaki Takai[1,2,11], Kunihiko Hinohara[3,4,11], Fu Gui[1], Takeshi Tsutsumi[1,2], Xiao Bai[1], Chenkui Miao[1], Chao Feng[1], Bin Gui[1], Zsofia Sztupinszki[5,6], Antoine Simoneau[7], Ning Xie[8], Ladan Fazli[8], Xuesen Dong[8,9], Haruhito Azuma[2], Atish D. Choudhury[3], Kent W. Mouw[10], Zoltan Szallasi[5,6], Lee Zou[7], Adam S. Kibel[1], Li Jia[1, *]

[1]Division of Urology, Department of Surgery, Brigham and Women's Hospital & Harvard Medical School, Boston, MA, USA

[2]Department of Urology, Osaka Medical and Pharmaceutical University, Osaka, Japan

[3]Department of Medical Oncology, Dana-Farber Cancer Institute & Harvard Medical School, Boston, MA, USA

[4]Department of Immunology, Nagoya University Graduate School of Medicine, Nagoya, Japan

[5]Computational Health Informatics Program, Boston Children's Hospital, Boston, MA, USA

[6]Danish Cancer Society Research Center, Copenhagen, Denmark

[7]Department of Pathology, Massachusetts General Hospital & Harvard Medical School, Boston, MA, USA

[8]Vancouver Prostate Centre, Vancouver General Hospital, Vancouver, British Columbia, Canada

[9]Department of Urologic Sciences, University of British Columbia, Vancouver, British Columbia, Canada

[10]Department of Radiation Oncology, Dana-Farber Cancer Institute & Brigham and Women's Hospital & Harvard Medical School, Boston, MA, USA

[11]These authors contributed equally to this article.

* Correspondence author: ljia@bwh.harvard.edu

ABSTRACT

Prostate cancer (PCa) harboring BRCA1/2 mutations is often exquisitely sensitive to PARP inhibition. However, genomic alterations in other DNA damage response genes have not been consistently predictive of clinical response to PARP inhibitors (PARPis). Here, we perform genome-wide CRISPR-Cas9 knockout screens in BRCA1/2-proficient PCa cell lines and identify novel genes whose loss has a profound impact on PARPi sensitivity and resistance. Specifically, MMS22L deletion, frequently observed (up to 14%) in PCa, renders cells hypersensitive to PARPis by disrupting RAD51 loading required for homologous recombination repair, although this response is TP53-dependent. Unexpectedly, loss of CHEK2 confers resistance rather than sensitivity to PARPis in PCa cells through increased expression of BRCA2, a target of CHEK2-TP53-E2F7-mediated transcriptional repression. Combined PARP and ATR inhibition overcomes PARPi resistance caused by CHEK2 loss. Our findings may inform the use of PARPis beyond BRCA1/2-deficient tumors and support reevaluation of currently used biomarkers for PARPi treatment in PCa.

INTRODUCTION

Despite treatment advances, metastatic castration-resistant prostate cancer (mCRPC) remains a lethal disease. Genomic sequencing studies reveal that approximately 90% of mCRPC patients carry clinically actionable genomic alterations [1]. Alterations in genes involved in the DNA damage response (DDR) are among the most common genetic events. Approximately 10% of primary and 25% of metastatic PCa patients have an alteration in at least one gene involved in DDR [2], which represents a potential therapeutic vulnerability. In particular, defects in homologous recombination repair (HRR) render cells highly sensitive to inhibition of Poly (ADP-ribose) polymerase (PARP). As a targeted therapy, PARP inhibitors (PARPis) prevent PARP1 and PARP2 from repairing DNA single-strand breaks (SSBs) and leads to stalled and collapsed replication forks by trapping PARP1/2 on the DNA breaks [3]. Subsequently, SSBs are converted to double-strand breaks (DSBs) that HRR-deficient cells cannot repair effectively, leading to overwhelming DNA damage, cell cycle arrest, and cell death. The BRCA1 and BRCA2 genes encode proteins essential for HRR, a pathway that repair DNA DSBs. PARPis selectively induce synthetic lethality in cancer cells harboring mutations in the BRCA1/2 genes [4,5].

Several PARPis are presently under clinical investigation in PCa, including olaparib, rucaparib, niraparib, and talazoparib, as a single agent. Two of them (olaparib and rucaparib) have been approved by the U.S. Food and Drug Administration (FDA) for the treatment of mCRPC patients with deleterious germline and/or somatic mutations in BRCA1/2, and the olaparib indication includes mutations in 12 additional HRR genes [6-11]. It is clear that BRCA1/2-mutant tumors exhibit high sensitivity and improved outcome to PARP inhibition based on results from these trials, as measured by standard radiographic criteria, 50% decrease in prostate-specific antigen (PSA), circulating tumor-cell counts, and progression-free survival or overall survival. However, whether and to what extent PARPis can be used to treat tumors with

non-BRCA1/2 alterations remains controversial after gene-by-gene analysis. Furthermore,

HRR-deficient tumors are not always sensitive to PARPis. Both intrinsic and acquired resistance

to PARP inhibition represents a formidable clinical problem. Therefore, one of the major barriers

to effective treatment using PARPis is how to distinguish patients who may and may not benefit

from PARP inhibition.

The genome-wide CRISPR-Cas9 (clustered regularly interspersed short palindromic repeat-

CRISPR associated nuclease 9) knockout (KO) screen is a powerful and unbiased approach to

identify genes that, when deleted, confer PARPi sensitivity or resistance. Previous studies using

CRISPR screens in non-PCa cell lines have identified genes whose loss impacts PARPi

response [12-14]. In this study, we carried out CRISPR screens in four BRCA1/2-proficient PCa cell

lines with the goal of expanding the use of PARPis beyond BRCA1/2 mutations and finding new

synthetically lethal interactions. Our screens led to the identification of MMS22L, a frequently

deleted HRR gene in PCa, as a new predictive biomarker for PARPis. Unexpectedly, we found

that loss of CHEK2 caused PARPi resistance, which we explored a therapeutic approach

through ATR inhibition to overcome in pre-clinical models.

RESULTS

Genome-wide CRISPR KO screens identify genes that modulate PARPi response in PCa

cells.

To identify genes whose loss increases or decreases the sensitivity of PCa cells to PARP

inhibition, we performed genome-wide CRISPR KO screens in the PCa LNCaP, C4-2B, 22Rv1

and DU145 cells in the presence of olaparib or DMSO vehicle. These four cell lines reflect

different aspects of PCa progression to castration resistance and have no deleterious biallelic

mutations in BRCA1/2 and other canonical HRR genes determined by whole exome sequencing

(Supplementary Table 1). LNCaP and C4-2B cells are relatively more sensitive to olaparib in

contrast to 22Rv1 and DU145 cells (Supplementary Fig. 1), but less sensitive when compared

to BRCA1-null ovarian cancer UWB1 cells. Cells were transduced with the lentiviral-based

CRISPR-Cas9 KO libraries, targeting over 18,000 protein-coding genes as previously described

[15,16], followed by 28 days of treatment with olaparib or DMSO (Fig. 1a; Supplementary Fig.

2a,b). The abundance of single guide RNA (sgRNA) for each gene was assessed by β-score

and the differential β-score was further calculated using the MAGeCKFlute pipeline by

comparing olaparib treatment to DMSO treatment [17,18]. We identified 216, 243, 153, and 211

negatively selected genes (with a stringent cutoff of p-value < 0.01, σ = 2.58), loss of which

sensitizes cells to olaparib in LNCaP, C4-2B, 22Rv1, and DU145 cells, respectively (Fig. 1b;

Supplementary Table 2). There were 67 genes shared by at least two cell lines, considered as

common hits (Supplementary Table 3). Gene Ontology (GO) analyses revealed that these

common hits were enriched for DNA repair and replication functions (Fig. 1c). Further analysis

of each individual cell line showed that the mitochondrial complex I assembly was the most

enriched function in DU145 (Fig. 1d; Supplementary Fig. 3). Dysfunction of mitochondrial

complex I causes a decline of NAD^+ and ATP supply required for PARP-mediated DNA repair [19].

The cell-type specific mechanism may provide a unique therapeutic vulnerability.

We identified 280, 347, 222, and 281 positively selected genes (p < 0.01, σ = 2.58), loss of

which renders cells resistant to olaparib in LNCaP, C4-2B, 22Rv1, and DU145 cells,

respectively (Fig. 1e; Supplementary Table 4). Given that cells were cultured in the presence of

olaparib for only 28 days, these genes are more likely related to intrinsic resistance rather than

acquired resistance arising after prolonged treatment with PARPis. There were 103 genes

shared in at least two cell lines (Supplementary Table 5). These common hits were enriched in

cell cycle phase transition and positive regulation of transcription (Fig. 1f). Analyses of positively

selected genes in each individual cell line showed that cell cycle genes are largely enriched in

CRPC C4-2B, 22Rv1, and DU145 cells, but not in androgen-dependent LNCaP cells (Fig. 1d;

Supplementary Fig. 3), reflecting the role of cell cycle transition in mediating PARPi resistance

in CRPC [20].

To further dissect the identified genes implicated in PARPi sensitivity and resistance, we

analyzed the common hits using the database in STRING protein-protein association network [21].

We found that 50% of the common negatively selected genes could be assigned to well-defined

functions in DDR, including HRR, DNA replication, Fanconi anemia, helicase, and cohesion

(Fig. 1g). In contrast, approximately 50% of the common positively selected genes were

involved specifically in cell cycle, ADP-ribosylation, transcriptional regulation (Fig. 1h). Finally,

we ranked all genes based on the average of differential β-scores from all four cell lines.

Representative top-ranked negatively and positively selected genes are shown in Fig. 1i,

including ones studied in this work. Notably, the BRCA1 and BRCA2 genes were highly ranked

in negative selection but reached the cutoff only in C4-2B and DU145 cells, respectively. This is

likely because acute inactivation of BRCA1 and BRCA2 is lethal without adaptive mechanisms

in cells [22-24], as reflected by the depletion of sgRNAs targeting BRCA1/2 genes even in the

absence of olaparib treatment in most cases, thereby failing to reach statistically significant

depletion in the olaparib condition (Supplementary Fig. 4). Together, our screen results provide

a global view of genetic determinants of PARPi response in PCa.

Validation of negatively selected genes related to PARPi sensitivity.

Next, we examined the alteration frequency (deletion and mutation) of the 67 common

negatively selected genes in the TCGA (primary tumors) and SU2C/PCF (metastatic tumors)

cohorts (Fig. 2a). We selected ten genes, based on their alteration frequency as well as their

important functions in DDR, and individually deleted these genes in C4-2B cells (Supplementary

Fig. 5). We observed markedly increased sensitivity to olaparib following deletion (Fig. 2b).

Among these genes, HELLS and WDR76 are the only two hits in all four cell lines. HELLS is a

helicase and chromatin remodeling enzyme that promotes the initiation of HRR and contributes

to DSB repair [25], while WDR76 is a specific reader of 5-hydroxymethylcytosine (5hmC) with

uncharacterized functions in DNA repair [26,27]. Indeed, WDR76 has a large interaction network,

including HELLS and PARP1 [28], supporting synthetically lethal interaction with PARP inhibition.

RNASEH2B and MMS22L are located on chromosomes 13q14 and 6q16, respectively, which

are two genomic regions frequently deleted in PCa [29-35]. While loss of RNASEH2B might cause

ribonucleotide excision repair deficiency and PARP-trapping lesions as previously reported [13],

we recently reported that co-loss of RB1, a closely located tumor suppressor gene, was

antagonistic in PCa cells [36]. MMS22L is known to form an obligate heterodimer complex with

TONSL, which reads histone H4 unmethylated lysine 20 [37] and promotes HRR of stalled or

collapsed replication forks and attendant DSBs by aiding RAD51 loading [38-40]. Cells with

MMS22L deletion are highly sensitive to the topoisomerase I inhibitor camptothecin. However,

the impact of MMS22L loss on PARP inhibition has not been studied.

Loss of MMS22L increases PARPi response due to impaired HRR function in PCa cells

Since MMS22L always forms a complex with TONSL for replication-associated DNA

damage repair, TONSL was unsurprisingly identified as one of the top hits as well (Fig. 1I;

Supplementary Table 2). Nevertheless, the frequency of MMS22L homozygous deletions (14%

in primary and 5% in metastatic prostate tumors) makes it an attractive biomarker to predict

PARPi response in PCa. To further validate this finding, we deleted MMS22L and TONSL in five

PCa cell lines - LNCaP, C4-2B, 22Rv1, PC-3, DU145 and MDAPCa2b (Fig. 3a). We showed

that deletion of either MMS22L or TONSL resulted in significantly increased sensitivity to

olaparib in LNCaP, C4-2B, MDAPCa2b and 22Rv1 cells, but not in DU145 or PC-3 cells. In a

growth competition experiment, we observed that olaparib treatment significantly reduced the

fraction of MMS22L-KO cells in contrast to corresponding control cells when they were equally

pre-mixed and grown under the same treatment condition (Fig. 3b). Notably, MMS22L-deleted

C4-2B cells were also sensitive to other PARPis (rucaparib, talazoparib, and veliparib), but

displayed lower sensitivity to carboplatin, a DNA crosslinking agent (Fig. 3c). In addition, we

generated MMS22L-KO C4-2B cell clones through single cell sorting and confirmed either

complete (sg1, clone #1-4) or partial MMS22L deletion (sg1, clone #5-6; sg2, clone #1-6) using

immunoblot (Fig. 3d). All MMS22L-KO clones exhibited high sensitivity to olaparib to a similar

extent without a gene dose effect, supporting MMS22L haploinsufficiency for its function.

Restoration of MMS22L expression completely abolished the sensitivity of MMS22L-KO cells to

olaparib (Fig. 3e), confirming that the phenotype was specifically due to MMS22L loss.

Next, we sought to investigate the mechanism by which loss of MMS22L increased PARPi

sensitivity. We recently reported that PARPi induces replication fork collapse and DSBs in a

trans cell cycle manner and BRCA1/2-deficient cells cannot recruit RAD51 to repair them,

leading to cell death [20]. Similarly, we observed increased expression of γ-H2AX and cleaved-

PARP in MMS22L-KO C4-2B cells following 72h olaparib treatment (Fig. 4a), reflecting PARPi-

induced DNA DSBs and apoptosis. Increased γ-H2AX foci were detected by

immunofluorescence staining (Fig. 4b). Cell cycle analysis showed a substantial accumulation

of cells in G2/M phase after deletion of MMS22L or TONSL (Fig. 4c). Given the function of the

MMS22L-TONSL complex in loading RAD51 at DSB sites and promoting HRR after replication

fork collapse independent of BRCA2 [38], we postulated that loss of MMS22L might impair HRR

that occurred predominantly in the late S and G2 phases of the cell cycle, leading to

overwhelming DSBs and mitotic catastrophe. Indeed, using immunofluorescence staining, we

found significantly reduced RAD51 foci in MMS22L-KO C4-2B cells after olaparib treatment for

24 hours in comparison with control cells (Fig. 4d). RAD51 loading was enhanced after

MMS22L expression was restored. From these observations, we conclude that MMS22L loss

confers high sensitivity to PARPis due to compromised RAD51 recruitment to PARPi-induced

DSBs, causing homologous recombination deficiency (HRD).

Tumors with HRD have been found to display specific patterns of genomic alterations,

which can be quantified using a HRD score based on genome-wide measuring three patterns of

genomic instability (loss of heterozygosity, telomeric allelic imbalance, and large-scale state

transitions) [41]. A high HRD score (≥42) has been shown to be predictive of clinical benefit with

PARPi or platinum therapy in breast and ovarian cancer patients [42,43]. Using whole-genome

sequencing data, we previously showed that a subset of PCa patients with a high HRD score

did not harbor germline or somatic mutations in BRCA1/2 and other HRR genes [44]. Further

analysis revealed that tumors with MMS22L heterozygous or homozygous deletion displayed

significantly higher HRD scores (Fig. 4e), although much higher scores were observed in tumors

with homozygous deletion. Tumors with both BRCA and MMS22L deficiency had the highest

HRD score. The MMS22L loss-mediated genomic instability was further confirmed using

HRDetect (Supplementary Fig. 6), a model that quantitatively aggregates six HRD-associated

signatures [45]. These results may partially explain previously uncharacterized cause of HRD in

PCa. In addition, using publicly available PCa clinical data, we found MMS22L deletion is

correlated with decreased MMS22L transcript levels (Fig. 4f). Tumors with heterozygous or

homozygous deletion had significantly lower MMS22L mRNA levels compared to those with

wild-type MMS22L. Patients with lower MMS22L expression had a shorter survival (Fig. 4g),

suggesting tumor progression driven by HRD-mediated genomic instability.

The response to PARP inhibition after MMS22L loss is p53-dependent

Next, we asked why deletion of MMS22L did not increase PARPi response in PC-3 and

DU145 cells (Fig. 3a). CRISPR screens revealed that TP53 was one of the top resistance genes

in LNCaP, C4-2B, and 22Rv1 cells (Fig. 1i; Supplementary Table 4). This was validated by

deletion of TP53 in these cell lines (Fig. 5a). LNCaP and C4-2B cells possess wild-type TP53,

while 22Rv1 cells harbor a mono-allelic TP53 mutation and have a relatively low p53 protein

level (Supplementary Table 1 and Supplementary Fig. 7). In contrast, PC-3 cells have TP53

truncating mutations and do not express p53 (p53-null), and DU145 cells harbor dominant

negative TP53 mutations [46]. Notably, PCa cell lines with TP53 mutations are generally less

sensitive to olaparib compared to cell lines with wild-type TP53 (Supplementary Fig. 1).

Furthermore, RNA-sequencing (RNA-seq) analysis revealed 271 commonly upregulated genes

in LNCaP and C4-2B cells after olaparib treatment, in which the p53 pathway is significantly

enriched (Fig. 5b,c; Supplementary Table 6), supporting the role of p53 activation in modulating

response to PARP inhibition. We reasoned that TP53 status might impact PARPi response of

MMS22L-deleted cells. Using an RNA interference approach, we showed that knockdown of

MMS22L expression significantly increased PARPi sensitivity in TP53 wild-type C4-2B cells,

whereas TP53-KO cells remained insensitive (Fig. 5d). Using a Dox-inducible system, we

reintroduced the TP53 gene into p53-null PC-3 cells. Restoration of p53 expression re-

sensitized MMS22L-KO PC-3 cells to olaparib *in vitro* and *in vivo* (Fig. 5e, f). We observed that

the level of γ-H2AX protein expression and foci were increased after restoration of p53

expression in MMS22L-KO PC-3 cells, indicating more DNA DSBs (Fig. 5g,h). Accordingly, cell

apoptosis was increased as determined by cleaved PARP levels. These results suggest that

MMS22L deletion increases PARPi sensitivity only when p53 remains functional. While 6% of

primary and 15.5% of metastatic tumors contain concurrent genomic alterations of MMS22L and

TP53, approximately 27% of primary and 20.5% of metastatic prostate tumors harbor wild-type

TP53 and MMS22L deletion when heterozygous deletion is counted (Fig. 5i). To further identify

MMS22L deletion in clinical samples, we utilized DNAscope assay, a new chromogenic DNA *in*

situ hybridization technique, on a tissue microarray (TMA) with 146 primary PCa tissue cores.

We detected 4.1% homozygous deletion and 26% heterozygous deletion of MMS22L in primary

PCa patients (Fig. 5j; Supplementary Fig. 8). Together, our results suggest that loss of MMS22L

occurs in a considerable fraction of PCa patients, who may therefore benefit from PARP

inhibition.

Validation of positively selected genes related to PARPi resistance

Next, we set out to investigate PARPi resistance. We found that the top positively selected

genes include those previously identified as conferring PARPi resistance in other contexts,

including PARP1, PARP2, PARG, ADPRHL2 (also known as ARH3), and TP53BP1 (Fig. 1i) [3,47-

50]. Although not altered as frequently as TP53 in PCa, downregulation of these genes could

influence PARPi response. PPP2R2A was previously included in the HRR gene list used to

select patients for the PROfound trial (NCT02987543). However, the FDA subsequently

removed PPP2R2A from the approved HRR gene panel because outcomes appeared worse for

patients with PPP2R2A alterations with olaparib compared to the control [6]. In accordance with

the clinical result, we identified PPP2R2A as one of the common hits from positive selection

(Fig. 1i; Supplementary Table 4). PPP2R2A is located on chromosome 8p and frequently

deleted in PCa (Supplementary Fig. 9). Indeed, chromosome 8p is the most frequent lost

chromosomal arm of all tumors (36%) in the TCGA cohort [35,51,52], suggesting a potentially

common intrinsic resistance mechanism in tumors with 8p loss.

Surprisingly, we identified Checkpoint kinase 2 (CHEK2) as one of the top resistance genes

in LNCaP, C4-2B and 22Rv1 cells. This is an unexpected result since CHEK2 is one of the

previously reported BRCAness genes [53] and an FDA-approved biomarker for olaparib in the

treatment of mCRPC patients. CHEK2 has been used as a biomarker of PARPi sensitivity in

several PARPi clinical trials [7-10,54,55] because of its known function in promoting HRR through

phosphorylation of BRCA1 [56,57]. We deleted CHEK2 in five PCa cell lines and observed

significantly increased resistance in LNCaP, C4-2B, 22Rv1 cells, but not in TP53-mutant PC-3

and DU145 cells using cell viability and colony formation assays (Fig. 6a,b). We further

generated CHEK2-KO C4-2B single cell clones, which all displayed resistance to olaparib (Fig.

6c). The resistance was also observed with other PARPis (rucaparib, talazoparib and veliparib)

and carboplatin using C4-2B and 22Rv1 cell models (Fig. 6d). We conclude that CHEK2 loss is

associated with PARPi resistance rather than sensitivity in PCa with functional TP53.

Loss of CHEK2/TP53 enhances HRR function through E2F7-controlled BRCA2

expression

CHEK2 regulates multiple proteins beyond BRCA1 in response to DNA damage. CHEK2

phosphorylates p53 on serine 20 and stabilizes p53, leading to cell cycle arrest in G1 phase

[58,59]. We therefore asked whether PARPi resistance caused by CHEK2 loss is mediated through

p53 inactivation. In line with activation of p53 pathway after PARP inhibition (Fig. 5c), we found

that olaparib treatment increased phosphorylated CHEK2 and p53 protein expression levels in

CHEK2-intact C4-2B and 22Rv1 cells, but not in CHEK2-KO cells (Fig. 7a). The extent of DNA

DSBs was also greater in CHEK2-intact cells compared to CHEK2-KO cells following olaparib

treatment as measured by γ-H2AX protein expression and foci (Fig. 7a,b). Accordingly, there

was increased olaparib-induced apoptosis in CHEK2-intact cells as measured by cleaved-PARP

levels (Fig. 7a). Deletion of TP53 had no effect on olaparib-induced CHK2 phosphorylation but

reduced DNA DSBs (γ-H2AX foci) and cell apoptosis (cleaved-PARP) as similarly observed in

CHEK2-KO cells (Fig. 7c).

Next, we sought to investigate the CHEK2-TP53 downstream target genes that may

contribute to PARPi resistance in CHEK2-KO cells. Previous studies have shown that E2F7 is a

TP53-reulated gene [60], and that deletion of E2F7 renders cells resistant to PARP inhibition

through upregulation of RAD51 in BRCA2-deficient cells [61]. E2F7 was also one of the top

positively selected genes in our CRISPR screens (Fig. 1i). E2F7 is an atypical member of the

E2F transcription factor family, which regulates its target genes, including HRR genes, through

transcription repression rather than activation. We reasoned that CHEK2 loss might derepress

HRR gene expression via the TP53-E2F7 axis. To investigate TP53-mediated regulation of

E2F7 expression, we analyzed publicly available p53 ChIP-seq data in multiple cell lines and

found highly enriched p53 binding immediately upstream of the E2F7 transcription start site

(TSS) (Supplementary Fig. 10a), indicating a conserved and direct transcriptional regulation

mechanism. Using p53 ChIP-qPCR, we observed baseline p53 binding at the E2F7 promoter,

which was significantly increased after olaparib treatment in CHEK2-intact C4-2B and 22Rv1

cells (Fig. 7d). However, olaparib-induced p53 binding was not observed after CHEK2 loss.

Accordingly, E2F7 expression was increased after olaparib treatment in CHEK2-intact C4-2B

cells within a few hours, but not in CHEK2-KO cells (Fig. 7e). Furthermore, the MDM2 inhibitor

(MDM2i) nutlin, an agent that blocks MDM2-mediated ubiquitination and promotes p53

stabilization and activity, increased p53 expression and p53 binding at the E2F7 promoter as

well (Supplementary Fig. 10b,c), supporting p53-regulated E2F7 expression. Indeed, MDM2i

treatment overcame olaparib resistance as a result of CHEK2 deletion in line with the role of

TP53-E2F7 axis in PARPi resistance (Supplementary Fig. 10d).

To investigate E2F7-controlled HRR gene expression, we analyzed publicly available E2F7

ChIP-seq data in multiple cell lines and observed strong E2F7 binding immediately upstream of

the TSSs of BRCA1/2 and RAD51 genes (Supplementary Fig. 10e). Using E2F7 ChIP-qPCR,

we demonstrated strong E2F7 binding at these genomic regions in both CHEK2-intact and

CHEK2-KO C4-2B and 22Rv1 cells (Fig. 7f). Importantly, E2F7 binding was significantly

increased in CHEK2-intact control cells after olaparib treatment but remained unchanged or

significantly decreased in CHEK2-KO cells. In agreement with the E2F7 ChIP results, we found

that mRNA levels of BRCA1/2 and RAD51 were significantly decreased after olaparib treatment

in CHEK2-intact C4-2B cells but increased in CHEK2-KO cells (Fig. 7g), indicating E2F7-

mediated transcriptional suppression on HRR genes. It should be noted that no significant cell

cycle alterations were observed after CHEK2 or TP53 deletion in C4-2B cells (Supplementary

Fig. 11a). Therefore, upregulation of BRCA1/2 and RAD51 gene expression is largely due to

transcriptional regulation rather than cell cycle alteration, although the expression of HRR genes

is cell cycle-dependent. The protein levels of BRCA1/2 and RAD51 were decreased as well

after olaparib treatment in CHEK2-inatct cells (Fig. 7h). However, the protein expression

changes were not all consistent with the mRNA expression levels in CHEK2-KO cells. While the

BRCA2 and RAD51 protein levels were increased or remained at a high level following olaparib

treatment, the BRCA1 protein level was decreased. This is likely because BRCA1 is directly

regulated by CHEK2 through posttranslational modification [56]. On the other hand, the reduction

of BRCA1 protein expression was not observed in TP53-KO cells since p53 is a downstream

effector of CHEK2 (Supplementary Fig. 11b). These results suggest that the CHEK2-TP53-

E2F7 axis is activated after olaparib treatment, leading to suppression of HRR gene expression,

and that this axis is disrupted when either CHEK2 or TP53 is lost. Specifically, the protein level

of the BRCA2 gene was significantly upregulated after CHEK2 or TP53 loss. In addition, we

found that olaparib-induced BRCA2 suppression was rescued after knockdown of E2F7,

rendering cell resistant to olaparib (Fig. 7i; Supplementary Fig. 11c).

To determine whether HRR capacity was enhanced after CHEK2 loss, we performed

RAD51 foci formation assay and found significantly increased olaparib-induced RAD51 foci in

CHEK2-KO C4-2B and 22Rv1 cells (Fig. 7j). These results suggest that the PARPi resistance

arising from CHEK2 loss is, at least in part, dependent on enhanced HRR function. To further

determine whether increased BRCA2 expression is responsible for PARPi resistance after

CHEK2 loss, we knocked down BRCA2 in CHEK2-KO C4-2B and 22Rv1 cells and found that

these cells were re-sensitized to olaparib (Fig. 7k). Similarly, knockdown of BRCA2 in TP53-KO

cells increased olaparib sensitivity (Supplementary Fig. 11d). In agreement with these results,

we found that there was a negative correlation between CHEK2 and BRCA2 protein expression

in the TCGA cohort (Fig. 7l). Together, our results support the notion that the CHEK2-TP53-

E2F7-BRCA2 pathway is one of the primary resistance mechanisms for PARP inhibition after

CHEK2 loss (Fig. 7m).

ATR inhibition overcomes PARPi resistance in CHEK2-deficient PCa cells.

Finally, we sought to explore therapeutic strategies to overcome PARPi resistance after

CHEK2 loss. Emerging evidence has shown that PARPi resistance is often accompanied by

increased ATR activity, which coordinates cell cycle checkpoint response through

phosphorylation of CHK1 and allows cells to survive PARPi-induced replication stress [62]. We

found that ATR activity was elevated in CHEK2-KO cells after olaparib treatment, as evidenced

by increased CHK1 phosphorylation in a dose-dependent manner (Supplementary Fig. 12). We

treated CHEK2-KO cells with olaparib in combination with M6620, a clinically used ATR inhibitor

(ATRi) [63]. M6620 was remarkably more effective when combined with olaparib in cell viability

assays (Fig. 8a). The combination therapy profoundly suppressed cell colony formation of

CHEK2-KO cells in a synergistic manner, as demonstrated by the HSA and Bliss synergy

scores [64,65] (Fig. 8b). Using *in vivo* xenograft models, we observed that CHEK2-KO C4-2B and

22Rv1 tumors did not respond to olaparib or M6620 as a single agent. However, the

combination treatment abolished the tumor growth (Fig. 8c and Supplementary 13). We did not

observe any weight loss in each group, indicating the combination therapy was well-tolerated.

Mechanistically, M6620 inhibited CHK1 phosphorylation as expected (Fig. 8d). The combination

treatment significantly inhibited BRCA1/2 and RAD51 expression in CHEK2-KO C4-2B and

22Rv1 cells, leading to more DNA damage and cell apoptosis as determined by γ-H2AX and

cleaved-PARP expression. Functionally, olaparib-induced RAD51 formation was completely

abolished in CHEK2-KO cells when they were treated with M6620 in combination with olaparib

(Fig. 8e).

DISCUSSION

In this study, we provided a systematic view of the genetic determinants and mechanisms

underlying the sensitivity and resistance of PCa to PARP inhibition beyond BRCA1/2 alterations.

We identified genes, such as RAD51B, RAD54L, FANCL, that have already been used as

biomarkers for PARP inhibition in the FDA-approved genetic test for mCRPC patients, as well

as previously unknown genes. These new genes could serve as predictive biomarkers for PARP

inhibition if mutated or deleted in PCa or therapeutic targets through pharmacologic inhibition in

combination with PARPis. One such novel hit is MMS22L, a gene located in a genomic region

frequently deleted in PCa, loss of which significantly increases the response to PARPis.

Mechanistically, MMS22L forms a complex with TONSL, which accumulates at distressed

replication forks and is required for the HRR of replication fork-associated DSBs through

promoting RAD51 loading [38,39,66]. When MMS22L is deleted, cells fail to load RAD51 to PARPi-

induced collapsed replication forks, leading to accumulation of DSBs, cell cycle arrest at G2/M

phase, and apoptotic cell death. The HRR function of MMS22L-TONSL seems specific to the

recovery of collapsed replication forks and does not act through BRCA2 [38]. Interestingly,

MMS22L deletions appear to be an early event in prostate tumorigenesis since they are more

frequently detected in primary tumors. If the use of PARPis in localized PCa (e.g., in the

neoadjuvant/adjuvant setting) is found to confer clinical benefit as has been suggested in breast

cancer with BRCA mutations [67], detection of MMS22L deletion would identify a much larger

population of patients who may benefit. In addition, our data suggest that cells with partial loss

of MMS22L (i.e., monoallelic loss) are equally sensitive to PARP inhibition compared to

complete loss (i.e., biallelic loss) likely due to insufficient MMS22L-TONSL complex formation.

This is further supported by the evidence from clinical sample analyses, showing significantly

lower mRNA levels and higher HRD scores in both homozygous and heterozygous MMS22L

deletion tumors. Identification of MMS22L as a new BRCAness gene explains a significant part

of patients that have high HRD scores but without HRR gene alterations per the olaparib label.

Given a considerable number of PCa patients with MMS22L deletion detected by next

generation sequencing- or DNA *in situ* hybridization-based assays, this genomic alteration may

be a valuable biomarker for PARP inhibition despite the fact that the response is p53-

dependent.

Previous studies of PARPi resistance have been focusing on acquired resistance

mechanisms in BRCA1/2-deficient cancers, showing that residual or restored HRR activity is the

most commonly observed resistance mechanism. This can be achieved by secondary or

reversion mutations of BRCA1, BRCA2 and RAD51 isoforms [68-70], loss of BRCA1 promoter

methylation [71], increased expression of hypomorphic isoforms of BRCA1 [72,73], loss of TP53BP1

and resection-associated factors RIF1, REV7, and Shieldin [49,74-76]. In contrast, our work

uncovers an intrinsic resistance mechanism in BRCA1/2-sufficient PCa cells, involving two

frequently mutated tumor suppressor genes TP53 and CHEK2. The role of TP53 and its

upstream activator CHEK2 in a set of tightly regulated cell cycle checkpoints and DDR events

has been extensively studied. In response to DNA DSBs, cells activate CHEK2 that

phosphorylates and stabilizes p53, leading to cell cycle arrest to allow for DNA repair or

inducing apoptosis after genotoxic damage. However, the PARPi resistance caused by loss of

CHEK2 or TP53 cannot be fully explained by impaired p53-mediated apoptosis. Indeed, PARPi

can trigger apoptotic cell death in TP53-deficient PCa cells [77]. High-grade serous ovarian cancer

(HGSOC) patients with defects in HRR are highly sensitive to PARP inhibition despite the fact

that TP53 mutations are detected in 96% HGSOC tumors [78]. Here, we propose an as yet

uncharacterized pathway that underlies the resistance of PCa cells to PARP inhibition. We show

that PARP inhibition activates the CHEK2-TP53-E2F7 pathway that suppresses the expression

of HRR genes and potentiates the cytotoxicity of PARPis. Loss of CHEK2 or TP53, however,

markedly reduces the expression of their downstream target E2F7, leading to increased HRR

gene (largely BRCA2) expression due to lack of E2F7-mediated transcriptional suppression.

Upregulation of BRCA2 enhances HRR capacity sufficient for the repair of PARPi-induced

DSBs and cell survival. Interestingly, BRCA1 is another direct target of the CHEK2 kinase. The

fact that loss of CHEK2 leads to PARPi resistance instead of sensitivity is unexpected. We show

that loss of CHEK2 may compromise BRCA1 regulation; after all, the BRCA1 protein expression

is not completely abolished. On the other hand, the CHEK2-TP53-E2F7 pathway has

predominantly emerged after PARP inhibition. Our finding is consistent with the results from

recent clinical trials which show little benefit in patients with mutations in CHEK2 [8,10,79], providing

a rationale to revisit FDA-approved clinically-used biomarkers for the use of olaparib. Our data

support the notion that HRR deficiency resulting from alterations in non-BRCA genes is unlikely

to be of similar therapeutic relevance in comparison to BRCA2 deficiency, considering that

mutations in the TP53 gene occur in more than 50% of all cancers. Conversely, PCa harboring

BRCA2 deficiency more likely respond to PARP inhibition regardless of TP53 status. This is

consistent with our CRISPR screen results, showing that BRCA2 was identified as one of the

top hits in TP53-mutated DU145 cells, while many of the canonical HRR genes (including

Fanconi anemia genes) were not negatively selected. Further investigation is needed to

determine which genomic alterations in the HRR pathway can or cannot be rescued by

upregulation of BRCA2 after loss of CHEK2 or TP53. We recently reported that RB1, another

frequently mutated tumor suppressor gene, may also confer resistance to PARP inhibition

through E2F1-mediated upregulation of HRR genes [36], indicating the importance and complexity

of E2F transcriptional network through activation and repression in the context of DNA repair.

The finding of the TP53/E2F7- and RB1/E2F1-mediated resistance mechanisms is clinically

relevant since TP53 and RB1 are concurrently altered in 39% of mCRPC tumors with

adenocarcinoma histology and 74% of mCRPC tumors with neuroendocrine features [80,81].

Current clinical use of PARPis is guided by mutations of a single gene and overlooks common

concurrent genomic alterations that may have an antagonistic effect.

Finally, we propose a therapeutic approach with combined PARP and ATR inhibition to

overcome PARPi resistance. Mechanistically, in order to survive PARP inhibition, cancer cells

rely more on ATR checkpoint to slow down cell cycle and reduce replication stress for repairing

DSBs. ATR inhibition increases replication origin firing and DNA synthesis despite the presence

of PARPi-induced replication fork gaps, leading to progressive accumulation of DSBs [20].

Moreover, ATR inhibition can disrupt RAD51 loading to stalled replication forks and DSBs in

PARPi-resistant BRCA1-deficient cells [82]. Our results further demonstrate that blocking ATR

activity inhibits HRR gene expression and RAD51 foci formation, leading to a high number of

stalled replication forks collapsing into toxic, irreparable DSBs. Indeed, the combination of

PARPi with ATRi was synergistic in many PARPi-resistant models regardless of their resistance

mechanisms [62]. Simultaneous suppression of ATR and PARP may be a preferred strategy to

overcome PARPi resistance.

METHODS

Cell Lines and Materials

Prostate cancer cell lines LNCaP, C4-2B, 22Rv1, PC-3, DU145, and MDA PCa 2b were

obtained from American Type Culture collection (ATCC). Cells were cultured following the

provider's recommendations. The identity of the cell lines was confirmed based on high-

resolution small tandem repeats (STR) profiling at Dana-Farber Cancer Institute (DFCI)

Molecular Diagnostics Core Laboratory and exome-sequencing analyses performed by LC

Sciences. Cells were regularly tested for mycoplasma. All reagents (including antibodies, small

molecule inhibitors, oligonucleotides, and plasmids) used in this study are listed in

Supplementary Table 7.

Cellular Viability Assay

PCa cells were seeded in 96-well plates with 1000-3000 cells/well (n = 6) and treated as

indicated. Cellular viability assays were performed using alamarBlue (ThermoFisher Scientific)

5-10 days after treatments and repeated 2–3 times.

Clonogenic Assay

PCa cells were seeded in 12-well plates at low density to avoid contact between clones.

Subsequently, cells were treated as indicated and allowed to grow for 10-14 days. Colonies

were fixed with paraformaldehyde (4%) for 20 minutes and stained with crystal violet (0.1%) for

15 minutes. Colony images were quantified using ImageJ software (National Institutes of

Health).

Cell Growth Competition Assay

MMS22L-KO and AAVS1 control cells were infected with lentiviruses generated using LV-GFP

(Addgene #25999) or LV-RFP (Addgene #26001) plasmid, respectively. MMS22L-KO LNCaP

and C4-2B cells stably express GFP, whereas control cells stably express RFP. GFP or RFP

positive cells were collected by the BD FACSAria cell sorter (BD Biosciences) and mixed in a

1:1 ratio. The mixed cell populations were incubated for 7 days with 3 uM of olaparib or DMSO

treatment, followed by collection and calculation of GFP and RFP positive cells.

Drug Synergy analysis

Cells were seeded in 12-well plates and treated with vehicle and 4 doses of olaparib (200, 300,

500 and 1000 nM) and M6620 (20, 30, 50 and 100 nM) in a matrix format to include 25 different

dose combinations. Colonies were quantified on day 14 using imageJ. Drug synergy scores

were calculated based on the HSA and Bliss model using the SynergyFinder2.0 [64].

RNA Interference

E2F7 and BRCA2 siRNAs and negative control siRNA were purchased from Sigma-Aldrich.

Cells were seeded onto 6-well or 96-well plates for 24 hours and followed by siRNA (10 nM)

transfection using Lipofectamine RNAiMAX (Invitrogen) according to the manufacturer's

protocol. The siRNA sequences are listed in Supplementary Table 7.

Cell Cycle Analysis

Cells were plated in 6-cm dishes with or without olaparib treatment for 72 hours. Collected cells

were labelled using Click-iT® EdU Flow Cytometry Assay Kit (Invitrogen) according to the

manufacturer's protocol and followed by flow cytometry using BD LSR Fortessa (BD

Biosciences).

Chromatin Immunoprecipitation Quantitative PCR (ChIP-qPCR)

ChIP experiments were performed as previously described [83,84]. Briefly, PCa cells were grown in

regular media and treated with olaparib (10 μM) for 24 hours prior to ChIP. Cells were cross-

linked by formaldehyde (1%) at room temperature (RT) for 10 minutes. After washing with ice-

cold PBS, cells were collected and lysed. The soluble chromatin was purified and fragmented by

sonication. Immunoprecipitation (IP) was performed using normal IgG or p53 or E2F7 antibody

(2 μg/IP). ChIP DNA was extracted and analyzed by qPCR using iTaq Universal SYBR Green

Supermix (Bio-Rad). The primer sequences are listed in Supplementary Table 7.

Quantitative RT-PCR (RT-qPCR)

Total RNA was extracted using RNeasy Mini Kit (Qiagen) according to the manufacturer's

protocol. RT-qPCR was performed as previously described [85]. The primer sequences are listed

in Supplementary Table 7.

RNA Sequencing (RNA-seq)

LNCaP and C4-2B cells were treated with olaparib (3 μM) or DMSO in biological duplicates.

Total RNAs were extracted using RNeasy Mini Kit (Qiagen). RNA-Seq library preparation and

next generation sequencing by Illumina HiSeq were conducted using GENEWIZ services.

***In vivo* Xenograft studies**

All mice were maintained in compliance with the guidelines approved by the Institutional Animal

Care and Use Committee (IACUC) at the Brigham and Women's Hospital. For xenograft

studies, male ICR/SCID mice and male NCG mice (NOD-$Prkdc^{em26Cd52}Il2rg^{em26Cd22}$/NjuCrl) at 4-5

weeks of age were purchased from Tconic Bioscience and Charles River Laboratories,

respectively. For the PC-3 xenograft assay, 2.5 million cells were prepared in 50 µl PBS and

mixed with 50 µl Matrigel (Corning Matrigel Matrix High Concentration #354262) to form a total

of 100 µl cell suspension and followed by the subcutaneous inoculation in male ICR/SCID mice.

Tumor volume was calculated using the modified ellipsoid formula: length x width^2/2. After

tumors developed to a volume of approximately150 mm^3, mice were randomized into 4 groups

(vehicle: n = 9, doxycycline: n = 5, olaparib: n = 5, and doxycycline plus olaparib: n = 5). Mice

were treated with vehicle (5% DMSO/10% d-a-tocopherol polyethylene glycol 1000 succinate;

oral gavage, OG), doxycycline (10 mg/kg in 10% d-a-tocopherol polyethylene glycol 1000

succinate; intraperitoneal, IP), olaparib (50 mg/kg in 5%DMSO/10% d-a-tocopherol polyethylene

glycol 1000 succinate; OG) or doxycycline (IP) plus olaparib (OG). Treatment was administered

every day for doxycycline and 5 days a week for vehicle and olaparib.

For the C4-2B and 22Rv1 xenograft assays, 5 million cells of C4-2B or 2.5 million cells of 22Rv1

were prepared as above. After tumors developed to a volume of approximately150 mm^3, mice

were randomized into 4 groups (vehicle, olaparib, M6620, and combination treatment) with 4

mice in each group. Mice were treated with vehicle, olaparib, M6620 (25 mg/kg in

5%DMSO/10% d-a-tocopherol polyethylene glycol 1000 succinate) or combination treatment

(olaparib 50 mg/kg and M6620 25 mg/kg in 5%DMSO/10% d-a-tocopherol polyethylene glycol

1000 succinate) through OG. Treatment was administered 5 days a week for single agent, and

5 days a week of olaparib plus 4 days a week of M6620 for combination treatment. Tumor

volume was manually measured every 3 days.

Immunoblot analysis

Cells were lysed in RIPA buffer supplemented with Halt™ protease and phosphatase inhibitor

cocktail (ThermoFisher Scientific). The protein concentration was determined using Pierce BCA

Protein Assay Kit (ThermoFisher Scientific). Proteins were resolved in SDS-polyacrylamide gels

(4–12%) and transferred to PVDF membranes using a Tris-glycine buffer system. Membranes

were blocked with 5% non-fat milk in 0.1% Tween20 in TBS (TBS-T) for 1 hour at room

temperature followed by incubation with primary antibodies and subsequently corresponding

secondary antibodies in 5% milk TBS-T. The membranes were developed with Immobilon

substrate (EMD Millipore).

PAR immunoblotting

The assay was performed according to the protocol as previously described [48]. Briefly, cells

were grown on 10 cm dishes for 24 hours to achieve 90% confluency. Cells were treated with

olaparib and/or PDDX-001 in the presence or absence of 0.01% MMS for a short period of time

as indicated. For total protein extraction, cells were lysed in RIPA buffer supplemented with

protease and phosphatase inhibitor cocktail and 1 μM PARG inhibitor ADP-HPD (Merck). For

chromatin fraction, cells were lysed and each fraction was collected using Subcellular Protein

Fractionation Kit (ThermoFisher Scientific) according to the manufacturer's protocol.

Immunoblotting was carried out as described in above.

Immunofluorescence Analyses

Cells were grown in a poly-L lysine-coated 4 well Millicell EZ slides (EMD Millipore) for 48 hours

and pre-extracted with 0.2% Triton-X on ice 90 seconds before fixation with 4%

paraformaldehyde in PBS for 20 minutes at room temperature (RT) and then permeabilized by

incubation with ice-cold methanol. After permeabilization, cells were blocked with 5% BSA in

PBS for 1 hour at RT, followed by incubation with 5% BSA in TBS-T containing primary

antibodies at a ratio of 1:200 overnight at 4°C. Cells were washed and incubated with secondary

fluorescent antibodies for 1 hour at RT. Alexa Fluor 488 anti-rabbit and Alexa Fluor 488 anti–

mouse antibodies were purchased from Invitrogen. After washing, the nuclear content was

stained with Mounting Medium with DAPI (Abcam) overnight at 4°C. Images were obtained with

fluorescence microscope BX53 (Olympus). Images were quantitatively assessed using ImageJ

software (NIH, Bethesda, MD) with 'Find maxima' function. More than 100 cells were analyzed

per condition in each experiment in a blinded manner.

Generating Knockout Cell Lines Using CRISPR/Cas9 Gene Editing

CRISPR guides targeting each gene were cloned into lentiGuide-Puro vector (#52963;

Addgene). The sgRNA sequences are listed in the Supplementary Table 7. The lentiCas9-Blast

vector that expresses Cas9 was obtained from Addgene (#52962). Lentiviruses were generated

using packaging vectors pMD2.G (#12259; Addgene) and psPAX2 (#12260; Addgene) with

Lipofectamine™3000 transfection reagents (#L3000015; Invitrogen) in 293FT cells. PCa cells

were infected with lentiviruses expressing Cas9 and selected with Blasticidin (10 µg/ml) to

establish stable Cas9-expressing cell lines. Polybrene was added at a final concentration of

8ug/ml to increase infecting efficiency. To generate KO cells, PCa cells were infected with

lentiviruses containing specific sgRNA and selected with puromycin (3 µg/ml). KO efficacy was

determined by immunoblot analyses. Single cell clones of MMS22L-KO and CHEK2-KO were

generated using BD FACSAria cell sorter (BD Biosciences) and followed by expansion.

Generation of TET-inducible Exogenous MMS22L- or P53-expressing Cells

MMS22L-KO and control C4-2B clone cells were infected with viruses containing TET-inducible

MMS22L gene. The MMS22L cDNA was created to be resistant to MMS22L sgRNA1 by the

introduction of silent mutations in the crispr RNA (crRNA) recognition sequence (change

CTTGGCAGGAATATAGCACAA to **CTAGGTAGA**AATATAGCACAA).

After neomycin (500 mg/ml) selection, doxycycline (0.15 μg/ml) treatment induces the MMS22L

expression confirmed by immunoblot. In addition, MMS22L-KO PC-3 cells were infected with

viruses containing TET-inducible WT TP53. Neomycin selection and the confirmation of p53

expression were performed as above. The MMS22L and TP53 cDNAs were cloned into the

Lenti-TRE3G-ORF-IRES-tRFP-PGK-Tet3G-neo vector, respectively. The plasmids were

custom-designed and synthesized at Transomic Technologies, Inc.

Genome-wide CRISPR-Cas9 Knockout Screen

We used genome-wide CRISPR-Cas9 KO H1 and H2 libraries obtained from Drs. Myles Brown

and X. Shirley Liu's laboratories [15], consisting of over 180,000 sgRNAs (10sgRNAs per gene).

For the CRISPR-Cas9 KO screens, 200 million PCa cells were infected with the lentiviral

CRISPR-Cas9 KO H1 and H2 libraries at a low multiplicity of infection (~0.3) to to maximize the

number of cells that have only one sgRNA integration. After 5 days of puromycin selection, the

surviving cells were expanded. Following preparation of 60 million cells for each condition to

achieve a representation of at least 300 cells per sgRNA, cells were divided into day 0 control

cells and cells cultured for 28 days (nine passages) treated with DMSO or olaparib before

genomic DNA extraction and library preparation. We used olaparib at the concentration of 5 μM

for LNCaP and C4-2B cells and 10 μM for 22Rv1 and DU145 cells. These concentrations were

close to the half maximal inhibitory concentration (IC50) for each cell line and allowed us to

identify both negatively (depleted) and positively (enriched) selected sgRNAs corresponding to

gene knockouts that increase and decrease olaparib response respectively. PCR was

performed using genomic DNA to construct the sequencing libraries. Each library was

sequenced at 30-40 million reads to achieve ~300 x average coverage over the CRISPR library.

The library from day 0 sample of each screen served as controls to identify positively or

negatively selected genes.

CRISPR-Cas9 KO Screen Data Analysis

CRISPR-Cas9 KO screen data was analyzed using the MAGeCK and MAGeCK-VISPR

algorithms [86,87]. MAGeCK calculated the read counts for each sgRNA. MAGeCK-VISPR

calculated the β-score for each gene by using AAVS1 gene as control. Comparison of the

differential β-score between olaparib treatment and DMSO treatment was performed using

MAGeCKFlute [17,18]. We ranked genes by differential β-score and robustly estimated σ, which is

the standard deviation of the differential β-score by a "quantile matching" approach. A cut-off

value was set to correspond 99 % of the data falling within 2σ, which defined genes with lower

β-scores than minus σ and higher β-scores than σ as negatively and positively selected genes,

respectively. Identified genes were further analyzed using STRING protein interaction analysis

[21]. GO analysis was performed using DAVID bioinformatics resources (https://david.ncifcrf.gov).

DNAscope Assay Using Tissue Microarray (TMA)

The TMA was constructed using primary prostate tumors retrieved from the Vancouver Prostate

Centre Tissue Bank as previously reported as approved by the institutional Review Board [88]. A

total of 146 tissue cores from 73 patients who had undergone radical prostatectomy were

included in this study. The DNAscope assay (Advanced Cell Diagnostics, Newark, CA) is a

chromogenic DNA in situ hybridization assay using two sets of target specific probes. Two

custom-designed probes were synthesized by Advanced Cell Diagnostics. DS-Hs-MMS22L-C1

is the probe targeting 29194-49180 of hg38 DNA range = chr 6:97142161-97283437. DS-Hs-

CEP6p-C2 is the control probe targeting 20891-35730 of hd38 DNA range = chr 6: 57224489-

57312704. The assay was performed on the TMA using DNAscope™ HD Duplex Reagent Kit

(Advanced Cell Diagnostics) according to the manufacturer's protocol. Briefly, the tissue section

was pretreated to allow access to target DNAs and followed by hybridization with two sets of

probes. Two independent signal amplification systems were used to detect both target DNAs.

Probes were hybridized to a cascade of signal amplification molecules, culminating in binding of

enzyme-labeled probes. Two chromogenic substrates were used and probe targeted regions

were visualized in red for the MMS22L locus and blue for the control region (i.e., the

centromeric region of chromosome 6p). An increase in the number of red dots relative to blue

dots indicates a copy number gain or amplification, while a decrease in the number of red dots

or no red dots indicates a copy number loss or deletion. The images were evaluated and scored

both digitally by the software (Aperio ImageScope, Leica Biosystems) and visually by a

pathologist (L. Fazli, Vancouver Prostate Centre). The MMS22L wild-type was defined as the

ratio of the total number of red dots divided by the total number of blue dots (red/blue) greater

than 0.5. The heterozygous deletion was defined as the ratio of red/blue is between 0.1-0.5,

while the homozygous deletion was defined as the ratio of red/blue is below 0.1.

RNA Sequencing Analysis

Raw RNA-seq reads are aligned to the human genome version hg38 by using STAR aligner [89].

Gene counts are quantified by using HT-seq [90] with REFSEQ annotation. Differentially

expressed genes are identified by using DESeq2 [91] with cutoff of FDR < 0.01, and ranked by the

statistics. The GSEA software was used for determining the KEGG pathways [92].

HRD and HRDetect Scores

HRD and HRdetect scores were determined as previously described [44]. A total of 267 PCa

cases with whole genome sequencing data were analyzed using the scarHRD R package and

HRDetect algorithm [45,93].

Statistical Analysis

Statistical analyses were performed using the unpaired 2-tailed Student's *t* test or 2-way

ANOVA with a post hoc Tukey's honest significant difference (HSD) test when comparing at

least three conditions using the Prism software (GraphPad). *P*-values of less than 0.05 were

considered as statistically significant.

Public Data Analysis

Clinical data sets were analyzed using the cBio Portal for Cancer Genomics (cBioPortal;

www.cbioportal.org) and PCaDB [94] (http://bioinfo.jialab-ucr.org/PCaDB/). Integrated Genome

Viewer (https://www.broadinstitute.org/igv/) was used for visualization of ChIP-seq data

analysis. ChIP-seq data were obtained from ChIP-Atlas [95] (http://chip-atlas.org).

Data Availability

RNA-seq data are available under GEO accession number GSE189186.

References

1   Robinson, D. *et al.* Integrative Clinical Genomics of Advanced Prostate Cancer. *Cell* **162**,
454, doi:10.1016/j.cell.2015.06.053 (2015).
2   Armenia, J. *et al.* The long tail of oncogenic drivers in prostate cancer. *Nat Genet* **50**,
645-651, doi:10.1038/s41588-018-0078-z (2018).
3   Murai, J. *et al.* Trapping of PARP1 and PARP2 by Clinical PARP Inhibitors. *Cancer Res*
72, 5588-5599, doi:10.1158/0008-5472.CAN-12-2753 (2012).
4   Farmer, H. *et al.* Targeting the DNA repair defect in BRCA mutant cells as a therapeutic
strategy. *Nature* **434**, 917-921, doi:10.1038/nature03445 (2005).
5   Bryant, H. E. *et al.* Specific killing of BRCA2-deficient tumours with inhibitors of
poly(ADP-ribose) polymerase. *Nature* **434**, 913-917, doi:10.1038/nature03443 (2005).
6   Hussain, M. *et al.* Survival with Olaparib in Metastatic Castration-Resistant Prostate
Cancer. *N Engl J Med* **383**, 2345-2357, doi:10.1056/NEJMoa2022485 (2020).
7   de Bono, J. *et al.* Olaparib for Metastatic Castration-Resistant Prostate Cancer. *N Engl J
Med* **382**, 2091-2102, doi:10.1056/NEJMoa1911440 (2020).
8   de Bono, J. S. *et al.* Talazoparib monotherapy in metastatic castration-resistant prostate
cancer with DNA repair alterations (TALAPRO-1): an open-label, phase 2 trial. *Lancet
Oncol* **22**, 1250-1264, doi:10.1016/S1470-2045(21)00376-4 (2021).
9   Abida, W. *et al.* Rucaparib in Men With Metastatic Castration-Resistant Prostate Cancer
Harboring a BRCA1 or BRCA2 Gene Alteration. *J Clin Oncol* **38**, 3763-3772,
doi:10.1200/JCO.20.01035 (2020).
10   Abida, W. *et al.* Non-BRCA DNA Damage Repair Gene Alterations and Response to the
PARP Inhibitor Rucaparib in Metastatic Castration-Resistant Prostate Cancer: Analysis
From the Phase II TRITON2 Study. *Clin Cancer Res* **26**, 2487-2496, doi:10.1158/1078-
0432.CCR-20-0394 (2020).
11   Mateo, J. *et al.* Olaparib in patients with metastatic castration-resistant prostate cancer
with DNA repair gene aberrations (TOPARP-B): a multicentre, open-label, randomised,
phase 2 trial. *Lancet Oncol* **21**, 162-174, doi:10.1016/S1470-2045(19)30684-9 (2020).
12   Olivieri, M. *et al.* A Genetic Map of the Response to DNA Damage in Human Cells. *Cell*
182, 481-496 e421, doi:10.1016/j.cell.2020.05.040 (2020).
13   Zimmermann, M. *et al.* CRISPR screens identify genomic ribonucleotides as a source of
PARP-trapping lesions. *Nature* **559**, 285-289, doi:10.1038/s41586-018-0291-z (2018).
14   Clements, K. E. *et al.* Identification of regulators of poly-ADP-ribose polymerase inhibitor
response through complementary CRISPR knockout and activation screens. *Nat
Commun* **11**, 6118, doi:10.1038/s41467-020-19961-w (2020).
15   Jeselsohn, R. *et al.* Allele-Specific Chromatin Recruitment and Therapeutic
Vulnerabilities of ESR1 Activating Mutations. *Cancer Cell* **33**, 173-186 e175,
doi:10.1016/j.ccell.2018.01.004 (2018).
16   Xu, H. *et al.* Sequence determinants of improved CRISPR sgRNA design. *Genome Res*
25, 1147-1157, doi:10.1101/gr.191452.115 (2015).
17   Wang, B. *et al.* Integrative analysis of pooled CRISPR genetic screens using
MAGeCKFlute. *Nat Protoc* **14**, 756-780, doi:10.1038/s41596-018-0113-7 (2019).
18   Li, Z. *et al.* CRISPR Screens Identify Essential Cell Growth Mediators in BRAF Inhibitor-
resistant Melanoma. *Genomics Proteomics Bioinformatics* **18**, 26-40,
doi:10.1016/j.gpb.2020.02.002 (2020).
19   Gupte, R., Liu, Z. & Kraus, W. L. PARPs and ADP-ribosylation: recent advances linking
molecular functions to biological outcomes. *Genes Dev* **31**, 101-126,
doi:10.1101/gad.291518.116 (2017).

20   Simoneau, A., Xiong, R. & Zou, L. The trans cell cycle effects of PARP inhibitors
underlie their selectivity toward BRCA1/2-deficient cells. *Genes Dev* **35**, 1271-1289,
doi:10.1101/gad.348479.121 (2021).
21   Szklarczyk, D. *et al.* The STRING database in 2021: customizable protein-protein
networks, and functional characterization of user-uploaded gene/measurement sets.
Nucleic Acids Res **49**, D605-D612, doi:10.1093/nar/gkaa1074 (2021).
22   Hakem, R., de la Pompa, J. L., Elia, A., Potter, J. & Mak, T. W. Partial rescue of Brca1
(5-6) early embryonic lethality by p53 or p21 null mutation. *Nat Genet* **16**, 298-302,
doi:10.1038/ng0797-298 (1997).
23   Ludwig, T., Chapman, D. L., Papaioannou, V. E. & Efstratiadis, A. Targeted mutations of
breast cancer susceptibility gene homologs in mice: lethal phenotypes of Brca1, Brca2,
Brca1/Brca2, Brca1/p53, and Brca2/p53 nullizygous embryos. *Genes Dev* **11**, 1226-
1241, doi:10.1101/gad.11.10.1226 (1997).
24   Adam, S. *et al.* The CIP2A–TOPBP1 axis safeguards chromosome stability and is a
synthetic lethal target for BRCA-mutated cancer. *Nat. Cancer* **2**, 1357-1371,
doi:https://doi.org/10.1038/s43018-021-00266-w (2021).
25   Kollarovic, G., Topping, C. E., Shaw, E. P. & Chambers, A. L. The human HELLS
chromatin remodelling protein promotes end resection to facilitate homologous
recombination and contributes to DSB repair within heterochromatin. *Nucleic Acids Res*
48, 1872-1885, doi:10.1093/nar/gkz1146 (2020).
26   Spruijt, C. G. *et al.* Dynamic readers for 5-(hydroxy)methylcytosine and its oxidized
derivatives. *Cell* **152**, 1146-1159, doi:10.1016/j.cell.2013.02.004 (2013).
27   Gilmore, J. M. *et al.* WDR76 Co-Localizes with Heterochromatin Related Proteins and
Rapidly Responds to DNA Damage. *PLoS One* **11**, e0155492,
doi:10.1371/journal.pone.0155492 (2016).
28   Dayebgadoh, G., Sardiu, M. E., Florens, L. & Washburn, M. P. Biochemical Reduction of
the Topology of the Diverse WDR76 Protein Interactome. *J Proteome Res* **18**, 3479-
3491, doi:10.1021/acs.jproteome.9b00373 (2019).
29   Kluth, M. *et al.* 13q deletion is linked to an adverse phenotype and poor prognosis in
prostate cancer. *Genes Chromosomes Cancer* **57**, 504-512, doi:10.1002/gcc.22645
(2018).
30   Brookman-Amissah, N. *et al.* Allelic imbalance at 13q14.2 approximately q14.3 in
localized prostate cancer is associated with early biochemical relapse. *Cancer Genet
Cytogenet* **179**, 118-126, doi:10.1016/j.cancergencyto.2007.08.017 (2007).
31   Dong, J. T., Chen, C., Stultz, B. G., Isaacs, J. T. & Frierson, H. F., Jr. Deletion at 13q21
is associated with aggressive prostate cancers. *Cancer Res* **60**, 3880-3883 (2000).
32   Lu, T. & Hano, H. Deletion at chromosome arms 6q16-22 and 10q22.3-23.1 associated
with initiation of prostate cancer. *Prostate Cancer Prostatic Dis* **11**, 357-361,
doi:10.1038/pcan.2008.4 (2008).
33   Hyytinen, E. R. *et al.* Defining the region(s) of deletion at 6q16-q22 in human prostate
cancer. *Genes Chromosomes Cancer* **34**, 306-312, doi:10.1002/gcc.10065 (2002).
34   Kluth, M. *et al.* Deletion lengthening at chromosomes 6q and 16q targets multiple tumor
suppressor genes and is associated with an increasingly poor prognosis in prostate
cancer. *Oncotarget* **8**, 108923-108935, doi:10.18632/oncotarget.22408 (2017).
35   Stopsack, K. H. *et al.* Aneuploidy drives lethal progression in prostate cancer. *Proc Natl
Acad Sci U S A* **116**, 11390-11395, doi:10.1073/pnas.1902645116 (2019).
36   Miao, C. *et al.* RB1 loss overrides PARP inhibitor sensitivity driven by RNASEH2B loss
in prostate cancer. *Sci Adv* **(in press)** (2022).
37   Saredi, G. *et al.* H4K20me0 marks post-replicative chromatin and recruits the TONSL-
MMS22L DNA repair complex. *Nature* **534**, 714-718, doi:10.1038/nature18312 (2016).

38 O'Donnell, L. *et al.* The MMS22L-TONSL complex mediates recovery from replication
stress and homologous recombination. *Mol Cell* **40**, 619-631,
doi:10.1016/j.molcel.2010.10.024 (2010).
39 Duro, E. *et al.* Identification of the MMS22L-TONSL complex that promotes homologous
recombination. *Mol Cell* **40**, 632-644, doi:10.1016/j.molcel.2010.10.023 (2010).
40 Piwko, W. *et al.* The MMS22L-TONSL heterodimer directly promotes RAD51-dependent
recombination upon replication stress. *EMBO J* **35**, 2584-2601,
doi:10.15252/embj.201593132 (2016).
41 How, J. A. *et al.* Modification of Homologous Recombination Deficiency Score Threshold
and Association with Long-Term Survival in Epithelial Ovarian Cancer. *Cancers (Basel)*
13, doi:10.3390/cancers13050946 (2021).
42 Telli, M. L. *et al.* Homologous Recombination Deficiency (HRD) Score Predicts
Response to Platinum-Containing Neoadjuvant Chemotherapy in Patients with Triple-
Negative Breast Cancer. *Clin Cancer Res* **22**, 3764-3773, doi:10.1158/1078-0432.CCR-
15-2477 (2016).
43 Ray-Coquard, I. *et al.* Olaparib plus Bevacizumab as First-Line Maintenance in Ovarian
Cancer. *N Engl J Med* **381**, 2416-2428, doi:10.1056/NEJMoa1911361 (2019).
44 Sztupinszki, Z. *et al.* Detection of Molecular Signatures of Homologous Recombination
Deficiency in Prostate Cancer with or without BRCA1/2 Mutations. *Clin Cancer Res* **26**,
2673-2680, doi:10.1158/1078-0432.CCR-19-2135 (2020).
45 Davies, H. *et al.* HRDetect is a predictor of BRCA1 and BRCA2 deficiency based on
mutational signatures. *Nat Med* **23**, 517-525, doi:10.1038/nm.4292 (2017).
46 Chappell, W. H. *et al.* p53 expression controls prostate cancer sensitivity to
chemotherapy and the MDM2 inhibitor Nutlin-3. *Cell Cycle* **11**, 4579-4588,
doi:10.4161/cc.22852 (2012).
47 Makvandi, M. *et al.* A PET imaging agent for evaluating PARP-1 expression in ovarian
cancer. *J Clin Invest* **128**, 2116-2126, doi:10.1172/JCI97992 (2018).
48 Gogola, E. *et al.* Selective Loss of PARG Restores PARylation and Counteracts PARP
Inhibitor-Mediated Synthetic Lethality. *Cancer Cell* **33**, 1078-1093 e1012,
doi:10.1016/j.ccell.2018.05.008 (2018).
49 Jaspers, J. E. *et al.* Loss of 53BP1 causes PARP inhibitor resistance in Brca1-mutated
mouse mammary tumors. *Cancer Discov* **3**, 68-81, doi:10.1158/2159-8290.CD-12-0049
(2013).
50 Prokhorova, E. *et al.* Unrestrained poly-ADP-ribosylation provides insights into chromatin
regulation and human disease. *Mol Cell*, doi:10.1016/j.molcel.2021.04.028 (2021).
51 El Gammal, A. T. *et al.* Chromosome 8p deletions and 8q gains are associated with
tumor progression and poor prognosis in prostate cancer. *Clin Cancer Res* **16**, 56-64,
doi:10.1158/1078-0432.CCR-09-1423 (2010).
52 Cai, Y. *et al.* Loss of Chromosome 8p Governs Tumor Progression and Drug Response
by Altering Lipid Metabolism. *Cancer Cell* **29**, 751-766, doi:10.1016/j.ccell.2016.04.003
(2016).
53 Lord, C. J. & Ashworth, A. BRCAness revisited. *Nat Rev Cancer* **16**, 110-120,
doi:10.1038/nrc.2015.21 (2016).
54 Stopsack, K. H. Efficacy of PARP Inhibition in Metastatic Castration-resistant Prostate
Cancer is Very Different with Non-BRCA DNA Repair Alterations: Reconstructing
Prespecified Endpoints for Cohort B from the Phase 3 PROfound Trial of Olaparib. *Eur*
Urol **79**, 442-445, doi:10.1016/j.eururo.2020.09.024 (2021).
55 Tung, N. M. *et al.* TBCRC 048: Phase II Study of Olaparib for Metastatic Breast Cancer
and Mutations in Homologous Recombination-Related Genes. *J Clin Oncol* **38**, 4274-
4282, doi:10.1200/JCO.20.02151 (2020).

56 Lee, J. S., Collins, K. M., Brown, A. L., Lee, C. H. & Chung, J. H. hCds1-mediated
phosphorylation of BRCA1 regulates the DNA damage response. *Nature* **404**, 201-204,
doi:10.1038/35004614 (2000).
57 Zhang, J. *et al.* Chk2 phosphorylation of BRCA1 regulates DNA double-strand break
repair. *Mol Cell Biol* **24**, 708-718, doi:10.1128/MCB.24.2.708-718.2004 (2004).
58 Hirao, A. *et al.* DNA damage-induced activation of p53 by the checkpoint kinase Chk2.
Science **287**, 1824-1827, doi:10.1126/science.287.5459.1824 (2000).
59 Takai, H. *et al.* Chk2-deficient mice exhibit radioresistance and defective p53-mediated
transcription. *EMBO J* **21**, 5195-5205, doi:10.1093/emboj/cdf506 (2002).
60 Aksoy, O. *et al.* The atypical E2F family member E2F7 couples the p53 and RB
pathways during cellular senescence. *Genes Dev* **26**, 1546-1557,
doi:10.1101/gad.196238.112 (2012).
61 Clements, K. E. *et al.* Loss of E2F7 confers resistance to poly-ADP-ribose polymerase
(PARP) inhibitors in BRCA2-deficient cells. *Nucleic Acids Res* **46**, 8898-8907,
doi:10.1093/nar/gky657 (2018).
62 Kim, H. *et al.* Combining PARP with ATR inhibition overcomes PARP inhibitor and
platinum resistance in ovarian cancer models. *Nat Commun* **11**, 3726,
doi:10.1038/s41467-020-17127-2 (2020).
63 Yap, T. A. *et al.* Phase I Trial of First-in-Class ATR Inhibitor M6620 (VX-970) as
Monotherapy or in Combination With Carboplatin in Patients With Advanced Solid
Tumors. *J Clin Oncol* **38**, 3195-3204, doi:10.1200/JCO.19.02404 (2020).
64 Ianevski, A., Giri, A. K. & Aittokallio, T. SynergyFinder 2.0: visual analytics of multi-drug
combination synergies. *Nucleic Acids Res* **48**, W488-W493, doi:10.1093/nar/gkaa216
(2020).
65 Ianevski, A., He, L., Aittokallio, T. & Tang, J. SynergyFinder: a web application for
analyzing drug combination dose-response matrix data. *Bioinformatics* **33**, 2413-2415,
doi:10.1093/bioinformatics/btx162 (2017).
66 Huang, T. H. *et al.* The Histone Chaperones ASF1 and CAF-1 Promote MMS22L-
TONSL-Mediated Rad51 Loading onto ssDNA during Homologous Recombination in
Human Cells. *Mol Cell* **69**, 879-892 e875, doi:10.1016/j.molcel.2018.01.031 (2018).
67 Tutt, A. N. J. *et al.* Adjuvant Olaparib for Patients with BRCA1- or BRCA2-Mutated
Breast Cancer. *N Engl J Med* **384**, 2394-2405, doi:10.1056/NEJMoa2105215 (2021).
68 Kondrashova, O. *et al.* Secondary Somatic Mutations Restoring RAD51C and RAD51D
Associated with Acquired Resistance to the PARP Inhibitor Rucaparib in High-Grade
Ovarian Carcinoma. *Cancer Discov* **7**, 984-998, doi:10.1158/2159-8290.CD-17-0419
(2017).
69 Lin, K. K. *et al.* BRCA Reversion Mutations in Circulating Tumor DNA Predict Primary
and Acquired Resistance to the PARP Inhibitor Rucaparib in High-Grade Ovarian
Carcinoma. *Cancer Discov* **9**, 210-219, doi:10.1158/2159-8290.CD-18-0715 (2019).
70 Pettitt, S. J. *et al.* Clinical BRCA1/2 Reversion Analysis Identifies Hotspot Mutations and
Predicted Neoantigens Associated with Therapy Resistance. *Cancer Discov* **10**, 1475-
1488, doi:10.1158/2159-8290.CD-19-1485 (2020).
71 Ter Brugge, P. *et al.* Mechanisms of Therapy Resistance in Patient-Derived Xenograft
Models of BRCA1-Deficient Breast Cancer. *J Natl Cancer Inst* **108**,
doi:10.1093/jnci/djw148 (2016).
72 Wang, Y. *et al.* BRCA1 intronic Alu elements drive gene rearrangements and PARP
inhibitor resistance. *Nat Commun* **10**, 5661, doi:10.1038/s41467-019-13530-6 (2019).
73 Wang, Y. *et al.* The BRCA1-Delta11q Alternative Splice Isoform Bypasses Germline
Mutations and Promotes Therapeutic Resistance to PARP Inhibition and Cisplatin.
Cancer Res **76**, 2778-2790, doi:10.1158/0008-5472.CAN-16-0186 (2016).

74 Xu, G. *et al.* REV7 counteracts DNA double-strand break resection and affects PARP
inhibition. *Nature* **521**, 541-544, doi:10.1038/nature14328 (2015).
75 Noordermeer, S. M. *et al.* The shieldin complex mediates 53BP1-dependent DNA repair.
Nature **560**, 117-121, doi:10.1038/s41586-018-0340-7 (2018).
76 Gupta, R. *et al.* DNA Repair Network Analysis Reveals Shieldin as a Key Regulator of
NHEJ and PARP Inhibitor Sensitivity. *Cell* **173**, 972-988 e923,
doi:10.1016/j.cell.2018.03.050 (2018).
77 Gonzalez-Billalabeitia, E. *et al.* Vulnerabilities of PTEN-TP53-deficient prostate cancers
to compound PARP-PI3K inhibition. *Cancer Discov* **4**, 896-904, doi:10.1158/2159-
8290.CD-13-0230 (2014).
78 Cancer Genome Atlas Research, N. Integrated genomic analyses of ovarian carcinoma.
Nature **474**, 609-615, doi:10.1038/nature10166 (2011).
79 de Bono, J., Kang, J. & Hussain, M. Olaparib for Metastatic Castration-Resistant
Prostate Cancer. Reply. *N Engl J Med* **383**, 891, doi:10.1056/NEJMc2023199 (2020).
80 Mu, P. *et al.* SOX2 promotes lineage plasticity and antiandrogen resistance in TP53- and
RB1-deficient prostate cancer. *Science* **355**, 84-88, doi:10.1126/science.aah4307 (2017).
81 Ku, S. Y. *et al.* Rb1 and Trp53 cooperate to suppress prostate cancer lineage plasticity,
metastasis, and antiandrogen resistance. *Science* **355**, 78-83,
doi:10.1126/science.aah4199 (2017).
82 Yazinski, S. A. *et al.* ATR inhibition disrupts rewired homologous recombination and fork
protection pathways in PARP inhibitor-resistant BRCA-deficient cancer cells. *Genes Dev*
31, 318-332, doi:10.1101/gad.290957.116 (2017).
83 Decker, K. F. *et al.* Persistent androgen receptor-mediated transcription in castration-
resistant prostate cancer under androgen-deprived conditions. *Nucleic Acids Res* **40**,
10765-10779, doi:10.1093/nar/gks888 (2012).
84 Gui, B. *et al.* Selective targeting of PARP-2 inhibits androgen receptor signaling and
prostate cancer growth through disruption of FOXA1 function. *Proc Natl Acad Sci U S A*
116, 14573-14582, doi:10.1073/pnas.1908547116 (2019).
85 Zheng, D. *et al.* Secretory leukocyte protease inhibitor is a survival and proliferation
factor for castration-resistant prostate cancer. *Oncogene* **35**, 4807-4815,
doi:10.1038/onc.2016.13 (2016).
86 Li, W. *et al.* MAGeCK enables robust identification of essential genes from genome-
scale CRISPR/Cas9 knockout screens. *Genome Biol* **15**, 554, doi:10.1186/s13059-014-
0554-4 (2014).
87 Li, W. *et al.* Quality control, modeling, and visualization of CRISPR screens with
MAGeCK-VISPR. *Genome Biol* **16**, 281, doi:10.1186/s13059-015-0843-6 (2015).
88 Li, Y. *et al.* SRRM4 gene expression correlates with neuroendocrine prostate cancer.
Prostate **79**, 96-104, doi:10.1002/pros.23715 (2019).
89 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
doi:10.1093/bioinformatics/bts635 (2013).
90 Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31**, 166-169,
doi:10.1093/bioinformatics/btu638 (2015).
91 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8
(2014).
92 Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-
15550, doi:10.1073/pnas.0506580102 (2005).

93　Sztupinszki, Z. *et al.* Migrating the SNP array-based homologous recombination
deficiency measures to next generation sequencing data of breast cancer. *NPJ Breast*
Cancer **4**, 16, doi:10.1038/s41523-018-0066-6 (2018).
94　Li, R. & Jia, Z., doi:10.1101/2021.06.29.449134 (2021).
95　Oki, S. *et al.* ChIP-Atlas: a data-mining suite powered by full integration of public ChIP-
seq data. *EMBO Rep* **19**, doi:10.15252/embr.201846255 (2018).
96　Conway, J. R., Lex, A. & Gehlenborg, N. UpSetR: an R package for the visualization of
intersecting sets and their properties. *Bioinformatics* **33**, 2938-2940,
doi:10.1093/bioinformatics/btx364 (2017).
97　Long, Q. *et al.* Global transcriptome analysis of formalin-fixed prostate cancer
specimens identifies biomarkers of disease recurrence. *Cancer Res* **74**, 3228-3237,
doi:10.1158/0008-5472.CAN-13-2699 (2014).
98　Ross-Adams, H. *et al.* Integration of copy number and transcriptomics provides risk
stratification in prostate cancer: A discovery and validation cohort study. *EBioMedicine* **2**,
1133-1144, doi:10.1016/j.ebiom.2015.07.017 (2015).
928
929

Author contribution

T.Tsujino and T.Takai performed the majority of experiments and analyses. T.Tsujino and

T.Takai, and K.H undertook the CRISPR screens. T.Tsujino and L.J. wrote the manuscript and

prepared figures. F.G. performed animal studies. T.Tsutsumi, X.B., C.M, C.F., B.G., and A.S.

performed cell studies and provided technical assistance. Z.Sztupinszki performed HRD

analysis. N.X., L.F. and X.D. performed DNA scope assay. H.A. and A.S.K. provided

administrative support. T.Tsujino, K.H, A.D.C., K.W.M., Z. Szallasi, L.Z., A.S.K., L.J. analyzed

the data and revised manuscript. L.J. conceptualized, designed the study, and acquired funding

for the present study.

Figure 1.

Fig. 1 CRISPR screens identify genes that modulate PARPi response in PCa cells. a,

Schematic of genome-wide CRISPR/Cas9 screens. **b,** UpSet plot [96] of negatively selected

genes in four PCa cell lines as indicated. Blue bars indicate the number of common hits in at

least two screens. **c,** Top GO terms enriched in 67 common hits from negative selection. **d,** Top

GO terms enriched in negatively (upper panel) and positively (lower panel) selected genes in

each individual cell line. **e,** UpSet plot of positively selected genes in four PCa cell lines as

indicated. Red bars indicate the number of common hits in at least two screens. **f,** Top GO

terms enriched in 103 common hits from positive selection. **g,** The networks of common hits

from negative selection, grouped according to their roles in specific pathways, and their genetic

and physical interactions (gray lines) based on STRING analyses. **h,** The networks of common

hits from positive selection as described in (**g**). **i,** Top-ranked genes in CRISPR screens

determined by comparing olaparib treatment to DMSO treatment. Genes are ranked by the

average of differential β-scores from all four cell lines. Negatively and positively selected genes

are marked in blue and red, respectively.

Figure 2.

Fig. 2 Validation of negatively selected genes with frequent genomic alterations. a, The

frequency of mutations and homozygous (Homo) deletions in 67 common negatively selected

962   genes from the TCGA Firehose Legacy cohort (Upper panel, n = 492) and the SU2C/PCF

963   cohort (Lower panel, n = 444). **b,** Dose response curves after treatment with olaparib in C4-2B

964   cells with knockout (KO) of the indicated genes. The *p*-values were determined by comparing

965   two gene-specific sgRNAs to a control AAVS1 sgRNA using two-way ANOVA. Error bars

966   represent SD, n = 6.

Figure 3.

Fig. 3 Loss of MMS22L increases PARPi response in PCa cells. a, Dose response curves

after treatment with olaparib in MMS22L- or TONSL-KO cells versus corresponding AAVS1

control cells of C4-2B, LNCaP, MDAPCa2b, 22Rv1, DU145 and PC-3. Upper right panels in

973 each cell line are immunoblot analysis of MMS22L or TONSL KO efficiency. ACTB (β-actin) is a

974 loading control. The olaparib response in MMS22L-KO C4-2B cells is presented in (**2b**). **b,** Flow

975 cytometry analyses of GFP and RFP positive cells. MMS22L-KO LNCaP or C4-2B cells (with

976 GFP) were co-cultured with corresponding AAVS1 control cells (with RFP) in a 1:1 ratio in the

977 presence of DMSO or olaparib (3 μM). Two MMS22L-KO cell lines (sg1 and sg2) and one

978 control cell line (sg1) were used. Cells were collected and analyzed using flow cytometry 7 days

979 after the treatment. The percentage of each cell population is presented in each panel. **c,** Dose

980 response curves after treatment with rucaparib, talazoparib, veliparib and carboplatin in two

981 MMS22L-KO C4-2B cell lines (sg1 and sg2) versus two AAVS1 control cell lines (sg1 and sg2).

982 **d,** Dose response curves (upper panel) after treatment with olaparib in control and MMS22L-KO

983 C4-2B cell clones. Immunoblot analysis (lower panel) of the MMS22L protein level in AAVS1

984 control and MMS22L-KO cell clones. **e,** Immunoblot analysis (left panel) of the MMS22L protein

985 level in C4-2B AAVS1 control sg1 clone 1 (cl1) and MMS22L-KO sg1 clone 1 (cl1), stably

986 infected with TET-inducible sgRNA-resistant MMS22L gene, after treatment with or without

987 doxycycline (0.15 μg/ml) for 3 days. Dose response curves (right panel) after olaparib treatment

988 with or without doxycycline (0.15 μg/ml) treatment in the same C4-2B cell clones. The *p*-value

989 was determined by comparison between the MMS22L-KO cell clone with and without

990 doxycycline treatment using two-way ANOVA. Error bars represent SD, n = 6. In (**a**), (**c**) and (**d**),

991 the *p*-values were determined by comparing MMS22L- or TONSL-KO cells to corresponding

992 control cells using two-way ANOVA. Error bars represent SD, n = 6.

Figure 4.

Fig. 4 Loss of MMS22L impairs HRR function in PCa cells. a, Immunoblot analysis of

cleaved PARP and γ-H2AX in AAVS1 control and MMS22L-KO C4-2B cells after olaparib

treatment for 72 hours with the indicated concentrations. **b,** Representative images and

quantification of γ-H2AX foci in AAVS1 control and MMS22L-KO C4-2B cells after olaparib

treatment (1 μM) for 24 hours. More than 100 cells were analyzed per condition. Solid lines

inside the violin indicate the median. The *p*-value was determined using unpaired *t*-test. Scale

bar = 5 μm. **c,** Cell cycle analysis (upper panel) in AAVS1 control, MMS22L-KO and TONSL-KO

LNCaP and C4-2B cells after treatment with DMSO or olaparib (5 μM) for 72 hours. Percentage

of cells (lower panel) in each phase of the cell cycle in these cells. **d,** Representative images

and quantification of RAD51 foci in AAVS1 control and MMS22L-KO C4-2B cells stably infected

with TET-inducible MMS22L gene after olaparib (5 μM) treatment in the presence or absence of

doxycycline (0.15 μg/ml) for 24 h. Dots indicate each replicate with more than 100 cells

analyzed. The *p*-values were determined using unpaired *t*-test. Error bars represent SD, n = 3.

ns = not significant. Scale bar = 5 μm. **e,** Ranked HRD scores (Upper panel) in PCa tumors with

BRCA and/or MMS22L genomic alterations as indicated. Comparison of HRD scores (lower

panel) between patients groups classified based on BRCA and MMS22L status. HRD score is

calculated using whole-genome sequencing datasets as previously described [44]. The *p*-values

were determined using unpaired *t*-test. **f,** The mRNA levels of MMS22L in PCa tumors with

Intact, heterozygous (Hetero) deletion and homozygous (Homo) deletion of MMS22L in the

TCGA cohort (Firehose Legacy; n = 487) and the SU2C/PCF cohort (n = 196). The *p*-values

were determined using unpaired *t*-test. **g,** Kaplan-Meier survival curves in the Long PCa cohort

[97] and the Cambridge PCa cohort [98] based on the MMS22L mRNA expression level (lower

versus higher). A log-rank test was carried out to examine the survival difference.

1019

Figure 5.

1021

Fig. 5 TP53 status impacts PARPi response in MMS22L-depleted PCa cells. a, Dose

response curves after treatment with olaparib in LNCaP, C4-2B, and 22Rv1 cells after TP53

deletion. The *p*-values were determined by comparing two TP53-KO cell lines (sg1 and sg2) to

two corresponding AAVS1 control cell lines (sg1 and sg2) using two-way ANOVA. Error bars

represent SD, n = 6. **b,** Venn diagram of up-regulated genes in LNCaP and C4-2B cells after

olaparib treatment. **c,** GSEA (upper panel) of RNA-seq data showing enrichment of p53

signaling pathway in up-regulated genes in LNCaP and C4-2B after olaparib treatment. Top

KEGG pathways (lower panel) enriched in these up-regulated genes. Normalized enrichment

score (NES) and false discover rate (FDR) in each cell line are shown. **d,** Immunoblot analysis

(upper panel) of the MMS22L protein level after siRNA knockdown. Dose response curves

(lower panel) after olaparib treatment in TP53-KO and AAVS1 control C4-2B cells with or

without MMS22L siRNA knockdown. Two different MMS22L siRNAs (#1 and #2) and one

control siRNA were transfected into TP53-KO and control C4-2B cells for 2 days followed by

olaparib treatment for 7 days. The *p*-values were determined by comparing siMMS22L to

siControl using two-way ANOVA. Error bars represent SD, n = 6. **e,** Immunoblot analysis (upper

panel) of p53 in AAVS1 control and MMS22L-KO PC-3 cells stably infected with TET-inducible

TP53 gene in the presence or absence of doxycycline (0.15 μg/ml) for 3 days. Dose response

curves (lower panel) after olaparib treatment for 7 days in the presence or absence of

doxycycline (0.15 μg/ml) in these cells. The *p*-values were determined by comparing between

cells treated with and without doxycycline using two-way ANOVA. Error bars represent SD, n =

6. **f,** Schematic of the procedure (upper panel) in the xenograft model using MMS22L-KO PC-3

cells containing TET-inducible TP53 gene. Tumor growth (lower panel) in ICR-SCID mice after

treatment with olaparib (50 mg/kg) or vehicle with or without doxycycline induction for 4 weeks.

The *p*-values were determined by comparing between treatment with and without doxycycline

using two-way ANOVA. Error bars represent SD, n = 5. **g,** Immunoblot analysis of cleaved

PARP and γ-H2AX in MMS22L-KO PC-3 cells containing TET-inducible TP53 gene after the

treatment as indicated for 3 days. **h,** Representative images and quantification of γ-H2AX foci in

AAVS1 control and MMS22L-KO PC-3 cells containing TET-inducible TP53 gene after olaparib

(5 μM) treatment for 24 h in the presence or absence of doxycycline (0.15 μg/ml). More than

100 cells were analyzed per condition. Solid lines inside the violin indicate the median. The *p-*

value was determined by unpaired *t*-test. Scale bar = 5 μm. **i,** Frequency of genomic alterations

in the TP53 and MMS22L genes in the TCGA cohort (left panel; Firehose Legacy, n = 492) and

the SU2C/PCF cohort (right panel; n = 429). **j,** Representative images (40X) of MMS22L wild-

type (WT), heterozygous (Hetero) deletion, and homozygous (Homo) deletion determined by

DNAscope using a tissue microarray (TMA) containing 146 primary prostate tumor samples.

Red signals (red arrow) indicate a probe targeting the MMS22L gene on chromosome 6q. Blue

signals (blue arrow) indicate a chromosome enumeration control probe targeting the

centromeric region of chromosome 6p (CEP6p). The number of red and blue dots were counted

and the ratio of red/blue was calculated. WT, red/blue > 0.5; Hetero deletion, red/blue between

0.1-0.5; Homo deletion, red/blue < 0.1. Scale bar = 20 μm.

1062

1063

1064

Figure 6.

 Fig. 6 Loss of CHEK2 renders PCa cells resistant to PARP inhibition. a, Dose response

curves after treatment with olaparib in two CHEK2-KO (sg1 and sg2) and two AAVS1 control

(sg1 and sg2) LNCaP, C4-2B, 22Rv1, DU145, and PC-3 cell lines. Upper right panel in each cell

line is Immunoblot analysis of the CHEK2 protein level in CHEK2-KO versus control cells. **b,**

Representative colony growth images (upper panel) and quantification (lower panel) after

treatment with olaparib in CHEK2-KO and AAVS1 control PCa cell lines as indicated. The *p*-

values were determined using one-way ANOVA. Error bars represent SD, n = 3. **c,** Dose

response curves (left panel) after treatment with olaparib in CHEK2-KO and AAVS1 control C4-

2B cell clones. Immunoblot analysis (right panel) of the CHEK2 protein level in CHEK2-KO and

control cell clones. **d,** Dose response curves after treatment with rucaparib, talazoparib,

veliparib and carboplatin in CHEK2-KO and AAVS1 control C4-2B and 22Rv1 cells. In (**a**), (**c**)

and (**d**), the *p*-values were determined by comparing CHEK2-KO to AAVS1 control cells using

two-way ANOVA. Error bars represent SD, n = 6.

1080

1081

Figure 7.

1083

Fig. 7 Loss of CHEK2/TP53 enhances HRR function through E2F7-controlled BRCA2

expression. a, Immunoblot analysis of the indicated proteins in CHEK2-KO and AAVS1 control

C4-2B and 22Rv1 cells after treatment with and without olaparib (3 μM for C4-2B and 5 μM for

22Rv1) for 72 h. **b,** Representative images (left panel) and quantification (right panel) of γ-H2AX

foci in CHEK2-KO and AAVS1 control C4-2B and 22Rv1 cells after treatment with olaparib (5

μM for C4-2B and 7 μM for 22Rv1) for 24 h. More than 100 cells were analyzed per condition.

Solid lines inside the violin indicate the median. The *p*-values were determined using unpaired *t*-

test. Scale bar = 5 μm. **c,** Immunoblot analysis of the indicated proteins in TP53-KO and AAVS1

control C4-2B cells after treatment with and without olaparib (3 μM) for 72 h. **d,** p53 ChIP-qPCR

at the E2F7 promoter in CHEK2-KO and AAVS1 control C4-2B and 22Rv1 cells after treatment

with or without olaparib (10 μM) for 24 h. **e,** Immunoblot analysis (left panel) of the E2F7 protein

level in CHEK2-KO and AAVS1 control C4-2B cells after treatment with olaparib (10 μM) for the

indicated time. Quantitative evaluation of integrated optical density (IOD) (right panel) was

performed from three independent experiments. The E2F7 IOD values were normalized by

ACTB IOD. The results are shown as mean of each time point. The *p*-values were determined

by comparing CHEK2-KO to control cells using two-way ANOVA. Error bars represent SD, n=3.

f, E2F7 ChIP-qPCR at the promoter regions of BRCA1/2 and RAD51 genes in CHEK2-KO and

AAVS1 control C4-2B and 22Rv1 cells after treatment with or without olaparib (10 μM) for 24 h.

g, The mRNA expression of BRCA1/2 and RAD51 genes determined by RT-qPCR in CHEK2-

KO and AAVS1 control C4-2B cells after treatment with and without olaparib (10 μM) for 8 h. **h,**

Immunoblot analysis of the indicated proteins in CHEK2-KO and AAVS1 control C4-2B and

22Rv1 cells after treatment with and without olaparib (10 μM) for 72 h. The IOD values of the

indicated proteins normalized by ACTB IOD are shown. **i,** Immunoblot analysis (upper panel) of

BRCA2 expression in C4-2B and 22Rv1 cells after knockdown of E2F7 through RNA

interference with or without olaparib (3 μM) treatment for 72 h. Two E2F7 siRNAs and one

negative control (Ctrl) siRNA were used (also see Fig. 16). Dose response curves (lower panel)

after treatment with olaparib for 7 days in C4-2B and 22Rv1 cells transfected with siRNAs

against E2F7 or control. The *p*-values were determined by comparing siE2F7 to siControl using

two-way ANOVA. Error bars represent SD, n = 6. **j,** Representative images and quantification of

RAD51 foci in CHEK2-KO and AAVS1 control C4-2B cells after treatment with olaparib (3 μM)

for 24 h. Dots indicate each replicate with more than 100 cells analyzed. n = 3. Scale bar = 5

μm. **k,** Cell viability after treatment with olaparib (1 μM) for 7 days in CHEK2-KO C4-2B and

22Rv1 cells transfected with siRNAs against BRCA2 or control. n = 6. Immunoblot analysis of

the BRCA2 protein level after siRNA knockdown (upper panel). **l,** The correlation between

CHEK2 and BRCA2 protein levels in the TCGA cohort (Firehose Legacy; n = 352). Spearman

correlation is calculated. **m,** Schematic model of HRR function regulated by the CHEK2-TP53-

E2F7-BRCA2 pathway. In (**d**), (**f**), (**g**), (**j**) and (**k**), the *p*-values were determined using unpaired

t-test. Error bars represent SD. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, ns= not significant.

1122

1123

Figure 8.

1125

Fig. 8 ATR inhibition overcomes PARPi resistance caused by CHEK2 loss in PCa cells. a,

Cell viability in CHEK2-KO and AAVS1 control C4-2B and 22Rv1 cells after treatment with

DMSO, olaparib (3 μM), M6620 (100 nM) or combination of olaparib and M6620. The *p*-values

were determined using one-way ANOVA. Error bars represent SD, n = 6. **b,** Representative

colony growth images (left panel) after combination treatment of olaparib and M6620 in CHEK2-

KO and AAVS1 control C4-2B cells. 3D synergy maps (right panel) of HSA and Bliss scores

between olaparib and M6620. **c,** Schematic of the procedure (left panel) in xenograft mouse

models using CHEK2-KO C4-2B and 22Rv1 cells. After tumors developed to a volume of

around 150 mm^3, mice were randomized into 4 treatment groups as vehicle, olaparib (50

mg/kg), M6620 (25 mg/kg) and olaparib plus M6620. The single agent was administered 5

days/week, while the combination treatment was given 5 days/week for olaparib plus 4

days/week for M6620. Tumor growth of CHEK2-KO C4-2B (middle panel) and 22Rv1 (right

panel) cells in ICR-SCID mice treated as indicated are shown with representative images. * p <

0.0001 was determined using two-way ANOVA. Error bars represent SD, n = 4. **d,** Immunoblot

analyses of the indicated proteins in CHEK2-KO C4-2B and 22Rv1 cells after treatment as

indicated (olaparib, 3 μM; M6620, 100 nM) for 72 h. The IOD values of the indicated proteins

normalized by ACTB IOD were shown. **e,** Representative images (upper panel) and

quantification (lower panel) of RAD51 foci in CHEK2-KO C4-2B and 22Rv1 cells after treatment

as indicated (olaparib, 10 μM; M6620, 200 nM) for 24 h. Dots indicate each replicate with more

than 100 cells analyzed. The *p*-values were determined using unpaired *t*-test. Error bars

represent SD, n = 3. Scale bar = 5 μm.

1147