Proteomic study of differentially expressed proteins in seeds between parents and offspring of castor bean (*Ricinus communis* L.)

Xiaotian Liang[1]; Rui Luo[1]; Yanxin Zhang[1]; Mingda Yin[1]; Yanpeng Wen[1]; Xuemei Hu[1]; Zhiyan Wang[1]; Yumiao Huo[1]; Fenglan Huang[1,2,3,4,5,6]*

[1] College of Life Science and Food Engineering, Inner Mongolia Minzu University, Tongliao, Inner Mongolia, China

[2] Key Laboratory of Castor Breeding of State Ethnic Affairs Commission, Tongliao, Inner Mongolia, China

[3] Research Center for Engineering and Technology of Castor Industry by Universities in Inner Mongolia Autonomous Region, Tongliao, Inner Mongolia, China

[4] Key Laboratory of Castor Breeding of Inner Mongolia Autonomous Region, Tongliao, Inner Mongolia, China

[5] Collaborative Innovation Center of Castor Industry of Inner Mongolia Autonomous Region, Tongliao, Inner Mongolia, China

[6] Engineering Research Center of Castor Industry Technology Innovation, Inner Mongolia Autonomous Region, Tongliao, Inner Mongolia, China

* Corresponding author

E-mail: huangfenglan@imun.edu.cn (FH)

Abstract

Castor bean (*Ricinus communis L.*), one of the top 10 oilseed crops in the world, has high economic value. Hybridization is the most direct and effective method to breed new varieties with high yield, high oil content, and strong stress resistance. Therefore, prediction of desired traits in castor hybrid offspring is particularly important. In this study, proteomic analysis was performed to identify differentially expressed proteins (DEPs) in seeds between castor hybrid offspring and their female (Lm female line aLmAB2) and male parents (CSR•181). Among the DEPs upregulated in the seeds of hybrid offspring, the majority were related to seed yield and stress resistance, while some were related to oil synthesis and fatty acid synthesis and metabolism in seeds. In other words, the hybrid offspring showed heterosis for seed yield, stress resistance, oil synthesis, and fatty acid synthesis and metabolism when compared with their parents. Further, real-time quantitative polymerase chain reaction assays were performed on 12 genes encoding DEPs involved in oil synthesis, pollen abortion, yield, and stress resistance of seeds. The results showed that the expression levels of the 12 genes were consistent with those of the DEPs.

Introduction

Castor bean (*Ricinus communis* L.) belongs to the genus *Ricinus* in the family Euphorbiaceae. Due to its strong environmental adaptability, castor bean can grow in harsh conditions such as saline-alkali soil and heavy metal–contaminated soil and is considered a valuable plant resource for phytoremediation. Therefore, it has been widely planted in tropical, subtropical, and warm temperate countries, especially in India, China, and Brazil [1,2]. China has more than 2300 germplasm resources of castor collected both at home and abroad, and in-depth studies have been conducted to further explore the application of castor bean. In Inner Mongolia, castor bean is a characteristic oilseed crop pivotal to local economic and social development [3]. It is a special industrial oil crop with prominent intraspecific heterosis.

Seed storage substances gradually accumulate during seed development and maturation. Protein, one of the seed storage substances, can help a seed to form cellular structures and provide a nitrogen source for normal metabolism and development of seeds, thereby playing a vital role in cellular structure and function [4]. In 1994, Wilkins and Williams first proposed the concept of the proteome, which refers to all proteins expressed in a cell, tissue, or organism [5]. The methods used for proteomic analysis of seeds mainly include two categories: gel and nongel analysis. Gel analysis is most commonly used, including two-dimensional gel electrophoresis (2-DE) and two-dimensional difference gel electrophoresis (2D-DIGE) [6]. After the differentially expressed spots are determined by using 2-DE and 2D-DIGE, a complete protein atlas can be obtained by mass spectrometry (MS), and this can better

reflect the changes in protein expression and posttranslational modification [7,8]. Proteomic studies of plant seeds have attracted an increasing amount of attention because they can provide information on substance changes in seeds during their growth and development and help explore the adaptative mechanism of plants to environmental stress. Xia et al. [9] conducted proteomic analysis to investigate the impact of temperature on seed metabolism in dormant and germinating seeds and deciphered the regulation of central metabolism activity during seed germination. Gu et al. [10] studied seed germination vigor in *Brassica napus* seeds differing in oil content by an integrated proteomic and genomic approach, and their research laid a foundation for studying the regulatory mechanism of seed germination vigor in *Brassica napus*.

In this study, the seeds from castor female parents (the Lm female line aLmAB2), male parents (CSR•181), and their hybrid offspring were used as the study materials, and the heterotic traits of offspring compared with their parents were identified by proteomic techniques. Further, genes associated with these traits were determined and tested with reverse transcription–quantitative PCR (RT-qPCR). The findings of this study can lay a foundation for guiding the production practice of castor bean and cultivating new castor varieties with high yield, high oil content, and strong stress resistance.

Materials and methods

Experimental materials

Plant materials

The plant materials used in this study were the castor seeds from female parent (aLmAB2), male parent (CSR•181), and their hybrid offspring F_0 and F_1. Castor plants at the flowering stage were bagged and pollinated manually. Subsequently, the seeds that had a developmental time of 20 days (Fig 1A), 40 days (Fig 1B), and 60 days post pollination (DPP) (Fig 1C) were collected. After their testa were removed, the seeds were wrapped in tinfoil paper and stored in an ultralow-temperature freezer at -80°C for subsequent use. The experimental materials were provided by Tongliao Research Institute of Agriculture and Animal Husbandry.

Fig 1. Castor Seeds at Different Developmental Stages.

110   Note: A: 20 DPP; B: 40 DPP; C: 60 DPP

Reagents and instruments

The reagents used in this experiment included Tris-HCl solution, ethylene

diamine tetraacetic acid stock solution, 0.4% β-mercaptoethanol solution, 0.9 M

sucrose, balanced phenol solution, extraction solution, ammonium acetate/methanol

solution, acetone solution, lysis solution, NaOH solution, HCl solution, lysine

solution, Amersham CyDye DIGE Fluors (minimal dyes) for the Ettan DIGE kit

(GE), the 2-D Clean-up kit (GE), the 2-D Quant kit (GE), hydration solution, 87%

glycerol, covering oil, monomer storage solution, 4× separating gel buffer, 10%

sodium dodecyl sulfate (SDS) solution, N,N,N′,N′-tetramethylethylenediamine

(TEMED), 10% ammonium persulfate solution, 75% alcohol, gel-sealing liquid,

equilibrium solution, bromophenol blue dye, 1,4-dithiothreitol (DTT), iodoacetamide

(IAA), premixed protein marker, SDS electrophoresis buffer, stationary liquid,

amplification solution, Coomassie brilliant blue dye, decolorizing solution, RNA

Rapid Extraction Kit for plants (ZOMANBIO), GoTaq® qPCR and RT-qPCR

Systems (Promega), and reverse transcription system (Promega). All reagent formulas

are shown in S1 Table. The instruments used in this study included high-speed and

low-speed refrigerated centrifuges (Thermo), an Infinite M200 Pro multifunctional

microplate reader (TECAN), a gel imaging system and electrophoresis apparatus

(Bio-Rad), the Ettan IPGphor 3 for electrophoresis by isoelectric focusing (IEF) in the

first dimension, a MultiTemp IV water bath, a FLA9500 multifunctional biomolecular

imager (GE), and a real-time fluorescence-based quantitative PCR system (ABI 7500,

ABI company).

2-DE

Sample preparation

An optimized phenol extraction method was used to extract protein from

castor seeds. The obtained dry protein powder was aliquoted into 1.5-mL centrifuge

tubes. Each tube was added with a 20-fold volume of lysis solution (each 1 mL of

lysis solution was added to 0.0062 g DTT, 20 μL of immobilized pH gradient (IPG)

buffer, and 40 μL of protease inhibitor before use). Then, the tube was vortexed at a

low temperature for 30 minutes, followed by an ultrasound ice-bath for 20 minutes.

The process was repeated until dry protein powder was fully dissolved. After

centrifugation at $13,000g$ for 1 hour at 4 °C, the supernatant was transferred to a

1.5-mL centrifuge tube. The 2-D Clean-up kit was used to remove impurity from the

sample. The 2-D Quant kit was used for sample quantitation, which derived a

standard curve: $y = -0.0041x + 0.3751$ ($R^2 = 0.999$), where y was the absorbance

value of the postquantitation sample at 480 nm and x was the protein concentration

(μg/μL).

2-DE procedures

153    For the first dimension, 450 μL of the sample solution containing 800 μg of

154 protein was loaded for electrophoresis by IEF. First, 800 μg of the lysed sample was

155 transferred into a centrifuge tube, which was then added to hydration solution to

156 achieve a total volume of 450 μL (each 1 mL of hydration solution was added with 5

157 μL of NL3-10 IPG buffer and 0.0028 g of DTT). Then, the sample solution was

158 vacuum-centrifuged at 13,000 rpm for 1 hour at 4 °C. After the centrifuged sample

159 solution was transferred to the strip groove, a 24-cm-long dry gel strip (pH 3-10 NL)

160 was placed into the groove; after ensuring that the gel surface was fully in contact

161 with the sample, the covering oil was added, and then a lid was placed to cover the gel

162 strip groove. Finally, after all gel strips of samples had been placed into a PROTEAN

163 IEF instrument, the following IEF protocol was run the Table 1 at a constant

164 temperature of 20 °C:

165 **Table 1. IEF Program.**

Program	Voltage/V	Time/h	Purpose
S1	30	8	Hydration
S2	50	4	Desalination
S3	100	1	Desalination
S4	300	1	Desalination
S5	600	1	Desalination
S6	1000	2	Desalination
S7	8000	12	Focusing
S8	1000	∞	Maintaining

166    For the second-dimension SDS-PAGE, 12.5%T polyacrylamide gel was used.

167 First, the gel strips after the first-dimension IEF were equilibrated for 15 minutes in

each of 2% (w/v) DTT and 2.5% (w/v) IAA equilibration buffer. Then the

well-equilibrated strips were sealed with sealing liquid in a gel plate, which was then

placed in the Ettan DALT 12 unit (12 gels per run) (GE-061-TY7119, GE

Healthcare). Electrophoresis was performed at 2 W per gel for 1 hour and then 16 W

per gel until the bromophenol blue band was 0.5-1 cm away from the bottom of the

gel.

After electrophoresis, the gel was taken out and sequentially submerged in

stationary liquid for 30 minutes, amplification solution for 20 minutes, Coomassie

brilliant blue dye for 10 minutes, and finally the decolorizing solution until only

protein spots appeared on the gel, without other, interfering dye residues. The Image

Scanner III image scanning system (28-9076-07 GE Healthcare, USA) was used for

gel imaging, and the Image Master 2-D Platinum 7.0 software was used for gel image

analysis.

2D-DIGE

Sample preparation

185   The sample was lysed using the lysis solution without DTT or IPG buffer.

186   After lysis, the sample solution was adjusted to pH 8.0-9.0 using 100 mM NaOH and

187   1 M HCl, with a target pH of 8.5. The subsequent procedures were the same as those

188   described in 1.2.1.

Preparation of CyDye DIGE fluor minimal dyes

Preparation of 1 nmol/µL fluorescent dye stock solution: After the dye-storing tube was taken out of the -80°C refrigerator, each tube received 5 µL of *N,N*-dimethylformamide solution and then vortexed for 30 seconds to fully dissolve the dye. Next, it was centrifuged at 12,000*g* for 30 seconds to push the dye stock solution to the bottom of the tube. Preparation of 400 pmol/µL fluorescent dye working solution: Five microliters of fluorescent dye stock solution was diluted with 7.5 µL of *N,N*-dimethylformamide solution to achieve a final concentration of 400 pmol/µL, followed by centrifugation at 12,000*g* for 30 seconds to push the solution to the bottom of the tube. All the above procedures were performed at 4 °C in the dark.

A certain volume of the sample solution (containing 50 µg of protein) was taken out and mixed with 1 µL of the dye working solution through vortexing, followed by a brief centrifugation. Then, as a quench-labeling reaction, the sample was kept on ice in the dark for 30 minutes and then mixed with 1 µL of 10 mM lysine solution through vortexing. After a brief centrifugation, the sample was placed on ice in the dark for 10 minutes.

2D-DIGE procedures

According to the experimental design, the internal standard sample and the target samples labeled with Cy2, Cy3, or Cy5 were mixed in a centrifuge tube and then added to hydration solution (every 1 mL of hydration solution had been added with 5 µL of NL3-10 IPG buffer and 0.0028 g of DTT) up to a total volume of 450

μL. After being placed on ice in the dark for 10 minutes, the sample solution was treated as in 1.2.2 for the first-dimension IEF.

For the second-dimension SDS-PAGE, 12.5% polyacrylamide gel was prepared using low-fluorescence glass plates, followed by the same procedures as in 1.2.2. Light exposure was avoided during the entire process. The Typhoon FLA9500 multifunctional biomolecular imager was used for gel imaging, and the DeCyder software for 2D-DIGE analysis was used to identify and annotate the differentially expressed spots (ratio $\geq$ 1.5 and ANOVA $\leq$ 0.05).

MS identification and analysis of the DEP spots

MS identification

The DEP spots in the gel were recovered and sent to Shanghai Bioclouds Biological Tech Co. Ltd. for MS identification.

Database search and functional classification of differentially expressed proteins (DEPs)

The information of the DEPs identified by MS were obtained by searching the database of castor bean proteins (https://www.ncbi.nlm.nih.gov/). The proteins were functionally classified into Gene Ontology (GO) categories according to UniProt (http://www.UniProt.org/) and QuickGO (https://www.ebi.ac.uk/QuickGO/), and their

233 roles in metabolic pathways were categorized according to the Kyoto Encyclopedia of

234 Genes and Genomes (KEGG) Pathway Database (https://www.kegg.jp/kegg/pathway.

235 html). The histograms of GO classification were generated on the Omics Cloud

236 platform established by GENE DENOVO (http://www.omicshare.com/index.php).

237 The heatmaps were drawn using GraphPad Prism7.04 software.

RT-qPCR verification

240   The RNA Rapid Extraction Kit for plants was used to extract RNA from

241 castor seeds. The extracted RNA was reverse-transcribed into cDNA. SnapGene 2.8.0

242 software was used to design primers (Table 2).

243 **Table 2. RT-qPCR Primer Sequences.**

Spot No.	Name	Primer sequence (5′-3′)	Tm (°C)
A2	A2F	CAGGCTATCGCAAATGGTGAC	56.6
	A2R	ACAGGTGCAAGATTCTTCCAG	54.9
A9	A9F	CAAGTCAGGCCAATCAAGCAC	56.4
	A9R	ACAAGCTCGCCAACTTTGAC	56.8
A10	A10F	GCATCCTCAAGGAAGAGTTGC	56.4
	A10R	TGTTGCCGACAAGGTATGGT	56.7
A13	A13F	CAGGCCATCTTGCAGCATTG	57.6
	A13R	AGAGACAGTCATGACACGGTG	56.8
J1	J1F	TTCCACCAATGGCCAGTACC	57.8
	J1R	GGAGGCATGAAAGGCTTCTTG	56.7
B5	B5F	TCAGAGCTTCAGAGCTGGTG	57.1
	B5R	CGTGTTGGCCTGTTTCATAGG	56.6
B9	B9F	TCATGGAGTTTGCTTGTACTCG	54.8
	B9R	AGATAGGAGCAGCATTGTTTCC	54.8

B12	B12F	TGAGGCGAAACGTGATGTCAG	57.8
	B12R	CGGTTCGAGGTCTAGTCTTCTC	56.8
K15	K15F	GCCACAGGCTAATGAAGTTCG	56.6
	K15R	ACACCAGGACCAACACTCTC	57.1
K4	K4F	GGTGCACAGATGTTGCAGTC	57.1
	K4R	TCCTGGAGACAAAGTGGTGAG	56.7
K13	K13F	GAGGAAGGACTTGAGCTCCTG	57.7
	K13R	ATTGCTGGAGTTGCTGGC	56.4
N5	N5F	AGCATAATGCCATGCATGTCC	56.0
	N5R	AGGTTCCATTTCCTCGTGGC	58.0

The GoTaq® qPCR systems was used for qPCR. The reaction system

consisted of 10 μL of SYBR Premix Ex *Taq*, 0.4 μL of Rox reference dye II, 1.0 μL

of each of the upstream and downstream primers, 2.0 μL of template DNA, and

ddH$_2$O up to 20 μL. The amplification program was by 45 cycles of 95°C for 30

seconds (s), 95°C 10 s, melting temperature for 45 s, and 72 °C for 10 s. The results

were calculated using the 2$^{-\Delta\Delta Ct}$ method.

One-way ANOVA was performed with SPSS version 19.0 software, and the

relevant histograms were generated in Excel 2019 software.

Results

Optimization of the 2-DE system

Optimization of conditions for 2-DE

To determine whether it is necessary to remove impurities from protein

samples in this experiment, the same protein samples were divided into two groups,

one with impurity removal and the other without impurity removal, before 2-DE

(24-cm dry gel strips, pH 3-10 NL, protein loading amount 800 μg). The results are

presented in S1 Fig. To determine the optimal pH range for this experiment, the

protein samples with the same treatment (with or without impurity removal) were

subjected to IEF and SDS-PAGE using the 24-cm dry gel strips with pH 3-10 NL or

pH 4-7. The results are shown in S2 Fig. To determine the optimal protein loading

amount, different amounts of protein (800 μg, 1000 μg, and 1200 μg) were loaded for

2-DE (24-cm dry gel strips; pH 3-10 NL) under the same conditions. The comparison

results are shown in S3 Fig. Based on the results of these preliminary experiments, the

optimal experimental conditions were impurity removal, 24-cm dry gel strips with pH

3-10 NL, and a loading amount of 800 μg. These conditions were used in the

following experiments.

273   2D-DIGE results

274   The castor seeds at different developmental stages from parents and offspring

275   were subjected to 2D-DIGE. The results are shown in Fig 2.

Fig 2. 2D-DIGE Results of Castor Parents and their Offspring

Note: A: Male, F_0, 20 d. B: Female, F_1, 20 d, A&B when they had Developed to 20

DPP. C: Male, F_0, 40 d. D: Female, F_1, 40 d, C&D when they had Developed to 40

DPP. E: Male, F_0, 60 d. F: Female, F_1, 60 d, E&F when they had Developed to 40

DPP.

In the figures, the blue protein spots denote the internal standard samples

labeled with Cy2; the green protein spots are the Cy3-labeled samples from male and

284 female parents; and the red protein spots are the Cy5-labeled samples from castor

285 offspring. In Fig 2, the 2D-DIGE protein spots for the seeds at different development

286 stages from both castor parents and offspring indicate that the dye labeling of all three

287 samples was successful and met the requirements for experimental analysis.

MS identification and functional classification results of DEPs

MS identification results

The 2D-DIGE results of castor seeds at different development stages from

parents and their offspring were analyzed for differences using DeCyder 2-D

software. A total of 122 differentially expressed spots (ratio $\geq$ 1.5 and ANOVA $\leq$

0.05) were determined and recovered, of which 121 spots containing 93 proteins were

successfully identified by matrix-assisted laser desorption/ionization mass

spectrometry (MALDI-TOF-MS). The results are shown in S2 Table. The

sources/names and classification of these DEPs are listed in S3 Table. Among the

DEPs between the castor parents and their offspring, 69 were present in seeds at 20

DPP, 46 in seeds at 40 DPP, and 20 in seeds at 60 DPP. Some proteins had a

significantly different expression in the castor seeds at two or all three development

stages between castor parents and their offspring, such as starch synthase, D-7 late

embryogenesis abundant (LEA) protein, superoxide dismutase (SOD), and aspartic

proteases (APs).

Functional classification results of the DEPs

The 93 DEPs identified were classified to five categories (Fig 3) related to stress responses (61.29%), seed nutrient storage (17.20%), fatty acid synthesis and metabolism (12.90%), amino acid metabolism (6.45%), and others (2.15%). The DEPs were most often related to stress responses. These included SOD, APs, D-7 LEA protein, and glutathione peroxidase. The DEPs associated with nutrient storage in seeds included starch synthase (chloroplast/starch), 2S albumin precursor, and vicilin GC72-A. The DEPs associated with fatty acid synthesis and metabolism included stearoyl-ACP desaturase, oil body–associated protein 2C (OBAP2C), and acetyl-CoA carboxylase. The DEPs associated with amino acid metabolism included transaldolase, transketolase (chloroplast), S-adenosylmethionine synthase (SAMS), and others.

Fig 3. Functional Classification of the Differentially Expressed Proteins between Castor Parents and their Offspring.

Functional classification results of the DEPs in seeds at 20 DPP between castor parents and their offspring

A total of 69 DEPs were identified in seeds at 20 DPP between castor parents

and their offspring. The seeds from F_1 offspring had 16 upregulated DEPs (e.g.,

OBAP2C, acetyl transferase NATA1, and 2S albumin precursor) when compared with

their male parent and 13 upregulated DEPs (e.g., protein disulfide isomerase (PDI),

luminal-binding protein 5, and triphosphate isomerase) when compared with their

female parent. The seeds from F_1 offspring had 23 downregulated DEPs (e.g.,

dihydrolipoyl dehydrogenase 1, MLP-like protein 34, and salicylic acid-binding

protein 2) when compared with their male parent and nine downregulated DEPs (e.g.,

l-ascorbate peroxidase, adenosine kinase 2, and the 36-kDa porin in mitochondrial

outer membrane) when compared with their female parent. In the F_0 seeds, 11 DEPs

(e.g., legume protein A, SOD, and hydroxyl-ACP dehydrase) were upregulated when

compared with their male parent, and five DEPs (i.e., transaldolase, stearoyl-ACP

desaturase, 60S acidic ribosomal protein P2, SOD, and uncharacterized protein) were

upregulated when compared with their female parent. The F_0 seeds had 11

downregulated DEPs (e.g., the major latex allergen Hev b5, MLP-like protein 31, and

GDP-mannose 3,5-epimerase 1) when compared with their male parent and four

downregulated DEPs (i.e., adenosyl homocysteinase, xylem serine proteinase 1

precursor, cytochrome c oxidase subunit 6b-1, and lactoylglutathione lyase GLX1)

when compared with their female parent. The relative expression levels of DEPs are

shown in S4 Fig, and the results regarding their involvement in KEGG metabolic

pathways are shown in S5 Fig.

Fig 4 shows the GO functional annotations of the 69 DEPs in the three

categories of biological process, cellular component, and molecular function. The

DEPs between F_1 and parents and DEPs between F_0 and parents were mostly

associated with biological processes, including starch synthase, ribulose bisphosphate

carboxylase small chain, and glutathione peroxidase. A smaller number of DEPs were

associated with molecular functions, including glutathione peroxidase, APs, and

acetyltransferase NATA1. The fewest DEPs were associated with cellular

components, including 60S acidic ribosomal protein P2, stearoyl-ACP desaturase, and

PDI. The analysis of KEGG metabolic pathways showed that the DEPs between F_1

and parents and the DEPs between F_0 and parents were mostly involved in rcu01100

(metabolic pathways) and rcu01110 (biosynthesis of secondary metabolites), such as

alcohol dehydrogenase 1 (ADH1), starch synthase, and hydroxyl-ACP dehydrase. A

number of DEPs were involved in rcu01230 (biosynthesis of amino acids), rcu00710

(carbon fixation in photosynthetic organisms), and rcu00061(fatty acid biosynthesis).

Fig 4. GO Functional Annotation of Differentially Expressed Proteins in Seeds at 20 DPP between Castor Parents and their Offspring.

Note: A: Male, F_1, 20 DPP; B: Female, F_1, 20 DPP, C: Male, F_0, 20 DPP; D: Female, F_0, 20 DPP.

Functional classification results of the DEPs in seeds at 40 DPP between castor parents and their offspring

A total of 46 DEPs were identified in seeds at 40 DPP between castor parents and their offspring. The F_1 seeds had three upregulated DEPs when compared with their male parent, including D-7 LEA protein, legume protein A, and heat shock protein (HSP); and they had 10 upregulated DEPs when compared with their female parent, including the large subunit of 2S sulfur-rich seed storage protein, nucleolar

protein 56 (NOP56), and reduced nicotinamide adenine dinucleotide phosphate

(NADPH)-dependent aldehyde reductase 1. Compared with their male parents, four

DEPs were downregulated in the F_1 seeds, including ATP synthase subunit β, HSP,

ADH1, and plastid 3-keto-acyl-ACP synthase; when compared with female parents,

six DEPs were downregulated, including starch synthase, fructose-bisphosphate

aldolase, GroES chaperonin, etc. In the F_0 seeds, seven DEPs (e.g., APs, large

subunits of 2S sulfur-rich seed storage protein, and nucleoside diphosphate kinase

(NDPK)) and nine DEPs (e.g., SOD, D-7 LEA protein, and glutelin type-A precursor)

were upregulated when compared with the male and female parents, respectively. One

DEP (enoyl-ACP reductase (EAR) precursor) and three DEPs (i.e.,

translation-controlled tumor protein homolog, pre-pro-ricin, and GroES chaperonin)

were downregulated when compared with the male and female parents, respectively.

Compared with the F_0 generation, the seeds of the F_1 generation had seven

upregulated DEPs (e.g., legume protein A, HSP, and 14-3-3-like protein D isoform

X1) and seven downregulated DEPs (e.g., fructose-bisphosphate aldolase 1, adenosine

kinase 2, and adenosyl homocysteinase). The relative expression levels of the DEPs

mentioned above are depicted in S6 Fig, and the results regarding their involvement in

the KEGG metabolic pathways are presented in S7 Fig.

The GO functional annotations for the 46 DEPs are presented in Fig 5. The

majority of DEPs in seeds between F_1 and male parent were related to biological

processes and molecular functions; compared with the male parent, only legume

protein A was upregulated while all other DEPs were downregulated in the F_1 seeds.

Most of the DEPs in seeds between F_1 and the female parent (e.g., NADPH-dependent

aldehyde reductase 1 and rRNA N-glycosidase) were associated with biological

processes and molecular functions. Similarly, most of the DEPs between F_0 and

parents were related to biological processes and molecular functions, including

cysteine protease inhibitor and glutelin type-A precursor. The DEPs between F_1 and

F_0 were mostly related to biological processes, including SAMS and adenosyl

homocysteinase; the DEP categorized under cellular component was enolase, and

those categorized under molecular function included HSP and 14-3-3-like protein D

isoform X1. According to the KEGG metabolic pathways, some HSP family members

involved in rcu04141 (protein processing in endoplasmic reticulum), including HSP,

were differentially expressed in seeds between F_1 and parents, between F_0 and

parents, and between F_1 and F_0. In addition, among the upregulated DEPs in the F_0

seeds vs. the male parent, NDPK was involved in rcu04016 (MAPK signaling

pathway) and betaine aldehyde dehydrogenase 1 (BADH1) was involved in rcu00260

(glycine, serine, and threonine metabolism). SOD, an upregulated DEP in the F_0 seeds

compared with their female parent, was involved in rcu04146 (peroxisome); and

chaperonin, an upregulated DEP in the F_1 generation compared with F_0, was involved

in rcu03018 (RNA degradation).

Fig 5. GO Functional Annotation of Differentially Expressed Proteins in Seeds at

40 DPP between Castor Parents and their Offspring.

Note: A: Male, F_1, 40 DPP; B: Female, F_1, 40 DPP; C: Male, F_0, 40 DPP; D: Female,

F_0, 40 DPP; E: F_1 vs. F_0, 40 DPP.

Functional classification results of the DEPs in seeds at 60 DPP

between castor parents and their offspring

A total of 25 DEPs were identified in seeds at 60 DPP between castor parents

and their offspring. The F_1 seeds had two upregulated DEPs when compared with

their male parent, NOP56 and 5-methyltetrahydroproylglutamate-homocysteine

methyltransferase; and they had six upregulated DEPs when compared with their

female parent, including D-7 LEA protein, NDPK, and large subunits of

ribulose-1,5-bisphosphate carboxylase/oxygenase. Five DEPs were downregulated in

the F_1 seeds compared with their male parent, including malate dehydrogenase,

phospho-glucosidase, and enolase, and no DEP showed detectable downregulation in

expression in the F_1 seeds compared to their female parent. In the seeds of the F_0

generation, three DEPs (i.e., vicilin GC72-A, legume protein A, and seed storage

protein) and eight DEPs (e.g., 6-phosphogluconate dehydrogenase, large subunits of

2S sulfur-rich seed storage protein, and D-7 LEA protein) were upregulated compared

to the male and female parents, respectively. No DEP showed detectable

downregulation in the F_0 seeds compared with either of their parents. Compared with

F_0, the F_1 seeds had five upregulated DEPs (e.g., 2S albumin precursor,

phospho-glucosidase, and malate dehydrogenase) but no downregulated DEP. The

relative expression levels of the abovementioned DEPs are shown in S8 Fig, and the

results on their involvement in KEGG metabolic pathways are shown in S9 Fig. The

DEPs between F_0 and the male parent were not involved in any metabolic pathways.

The GO functional annotations of the 25 DEPs are shown in Fig 6. Most of the

DEPs between F_1 and parents and between F_0 and parents (e.g., ADH1, NOP56, and

NDPK) were related to biological processes and molecular functions. Most of the

DEPs between F_1 and F_0 (e.g., phospho-glucosidase, malate dehydrogenase, and

enolase) were related to biological processes. From the KEGG metabolic pathway

analysis, the DEPs between F_1 and parents, between F_0 and parents, and between F_1

and F_0 were mostly involved in rcu01100 (metabolic pathways) and rcu01110

(biosynthesis of secondary metabolites), including 6-phosphogluconate

dehydrogenase, ADH1, and 5-methyltetrahydroproylglutamate-homocysteine

methyltransferase. In addition, the DEPs upregulated in F_1 and F_0 over their parents

were also involved in the pathways of rcu00270 (cysteine and methionine

metabolism), rcu00592 (alpha-Linolenic acid metabolism), rcu00230 (purine

metabolism), etc. The DEPs upregulated in the F1 generation compared with F0 also

participated in the pathways related to rcu01200 (carbon metabolism) and rcu01230

(biosynthesis of amino acids).

Fig. 8-1. GO Ffunctional Aannotation of Ddifferentially Eexpressed Pproteins in

Sseeds at 60 DPP between Ccastor Pparents and their Ooffspring.

Note: A: Male, F1, 60 DPP; B: Female, F1, 60 DPP; C: Male, F0, 60 DPP; D: Female,

F0, 60d; E: F1 vs. F0, 60 DPP.

RT-qPCR results of the DEPs

Twelve DEPs in the seeds between castor parents and their offspring (Table 3)

were selected for RT-qPCR analysis, and the results are shown in Fig 9. The

463 expression of the genes corresponding to the 12 DEPs all showed differences

464 consistent with those of the DEPs.

465 **Table 3. The 12 Differentially Expressed Proteins Chosen for RT-qPCR**

466 **Analysis.**

Spot No.	Protein Name	Source	Time (d)	Related Traits
J1	Stearoyl ACP desaturase	F_0 Male	20	Oil synthesis Amino acid metabolism
K4	Oil body associated protein 2C	F_1 Male	20	Oil synthesis Amino acid metabolism
B12	Acetyl-CoA carboxylase	F_1 Female	20	Oil synthesis Amino acid metabolism
A10	26S protease regulatory subunit 6a	F_1 Male	20	Pollen abortion
N5	Cysteine protease	F_1 Male	20	Pollen abortion
A2	Superoxide dismutase	F_0 Female	40	Yield and stress resistance
A13	Amylosynthetase	F_1 Male	20	Yield and stress resistance
B5	Abundant protein D-7 in late embryonic development	F_1 Male	20	Yield and stress resistance
B9	Glutathione peroxidase	F_1 Male	20	Yield and stress resistance
K15	Alcohol dehydrogenase 1	F_1 Female	60	Yield and stress resistance
K13	2S albumin precursor	F_1 Male	20	Yield and stress resistance
A9	Aspartic protease	F_1 Male	20	Yield and stress resistance

Fig 7. The RT-qPCR Results of Partially Differentially Expressed Proteins

between Castor Parents and their Offspring.

Note: *: 0.01<P≤0.05, **: 0<P≤0.01.

Discussion

Through comparative proteomic analysis of seeds at different developmental

stages between castor parents and their offspring, 122 DEP spots were detected by

MS, though they only contained 93 DEPs. This is because a number of proteins were

significantly differentially expressed at all developmental stages of seeds, such as

starch synthase, D-7 LEA protein, SOD, and APs.

Most of the DEPs were associated with stress responses (61.29%) and they

were mostly upregulated in the seeds of offspring, which was consistent with the

results of physiological and biochemical measurements. SOD is an antioxidant

enzyme in the reactive oxygen species scavenging system that effectively prevents

peroxides (e.g., H_2O_2) from damaging the structure and function of plant cells. This

enzyme participates in the rcu04146 pathway (peroxisome). The peroxisome, an

essential organelle of plants, plays a key role in redox signal transduction and lipid

homeostasis [11-15]. APs are mainly involved in such processes as metabolism,

precursor protein processing, protein degradation, and programmed cell death, as well

as pollen germination and pollen-tube growth in plants [16-18]. D-7 LEA is a

stress-responding protein that massively accumulates in the late embryonic stage and

is generated under adverse conditions such as drought. It regulates the metabolic

balance of plants through mechanisms such as stabilizing the cell membrane and

binding metal ions [19-21]. Glutathione peroxidase contains sulfhydryl groups and

can effectively block the free radicals of reactive oxygen species from further

damaging the body by scavenging H_2O_2 and lipid peroxides [22,23]. ADH1, a

medium-chain zinc-containing enzyme widely found in various tissues in plants,

binds zinc ions. Zinc plays an essential role in the synthesis of indole-3-acetic acid

(IAA), an auxin in plants. Zinc deficiency can lead to decreased tryptophan and IAA

content, hindering the growth of plant roots. Therefore, ADHs are is closely related to

plant growth and development [24-26]. In addition to those five proteins, several

DEPs between offspring and parents were associated with stress responses, including

14-3-3-like protein, BADH1 (chloroplast), and l-ascorbate peroxidase.

Among the DEPs between castor offspring and parents, 17.20% were involved

in seed nutrient storage, most of which were upregulated in the offspring seeds. Starch

synthase is a key enzyme involved in starch synthesis and metabolism. Starch, the

main storage carbohydrate, may account for 50% of the dry matter in storage organs

(tubers or seeds), and it is an important energy source for plant growth and

development, so its content directly determines the yield of crops [27-29]. 2S albumin

precursor, a storage protein, can gradually be degraded to produce various amino

acids and thereby provide essential nitrogen, carbon, and sulfur sources for plant

growth during the seed germination and early growth stages, which is of great

significance for plant growth and development [30]. Vicilin GC72-A belongs to the

family of antimicrobial peptides, a class of small-molecule polypeptides synthesized

by plants to defend against pathogen invasion; these antimicrobial peptides can inhibit

or kill a variety of microorganisms and fungi [31,32]. Other DEPs associated with

seed nutrient storage of plants included adenosyl homocysteinase, NOP56, and

salicylic acid-binding protein 2.

Among the DEPs between castor offspring and parents, 12.90% was related to

plant fatty acid synthesis and metabolism, most of which were upregulated in the

offspring seeds. This result was consistent with the results of trait and quality

assessment. Stearoyl-ACP desaturase is a key enzyme in the conversion between

stearic acid and unsaturated fatty acids, so its upregulation may increase the

unsaturated fatty acid content in plants, thereby improving their resistance to low

temperature [33]. OBAP2C plays an essential role in the seed development of plants.

The oil body, a lipid-storing subcellular organellar particle in plant seeds, provides

nutrients to plants and is critical for plant growth and development [34-36].

Acetyl-CoA carboxylase is a biotinidase catalyzing the carboxylation of acetyl-CoA

to form malonyl-CoA in the initial step of fatty acid synthesis. Fatty acids re

membrane lipid components of all cells and an important energy reserve of

multicellular organisms. Hence, acetyl-CoA carboxylase, the catalyzer of the first step

of fatty acid biosynthesis, plays a critical role in plant growth and development

[37-39]. Other DEPs associated with fatty acid synthesis and metabolism included

EAR precursor, plastid 3-keto-acyl-ACP synthase I, and hydroxyl-ACP dehydrase.

Only 6.45% of the DEPs between castor offspring and parents were related to

amino acid metabolism, most of which were upregulated in the offspring seeds. Both

transaldolase and transketolase (chloroplast) are enzymes involved in the

nonoxidative phase of the pentose phosphate pathway. Transaldolase catalyzes the

transfer of the dihydroxyacetone moiety from sedoheptulose-7-phosphate to

glyceraldehyde-3-phosphate; as a result, it can regulate the production of

ribose-5-phosphate and NADPH by the pentose phosphate pathway, through which it

links the pentose phosphate shunt and glycolysis pathways and regulates the balance

between the oxidative and nonoxidative phases of the pentose phosphate pathway.

Transketolase is a "nonregulatory" enzyme that catalyzes the reversible transfer of

aldehyde group from activated ketose (donor) to activated aldose (receiver). The

upregulation of transketolase expression facilitates carbon fixation in plants, as well

as plant growth and development [40-44]. SAMS is the only enzyme synthesizing

SAM in organisms. SAM, a metabolite from the interaction of methionine and

adenosine triphosphate, has been widely found in animals, plants, and

microorganisms. It is involved in up to 40 types of biochemical reactions and plays a

significant role in improving the drought resistance of plants [45-48]. All three

enzymes mentioned above were involved in rcu01230 (biosynthesis of amino acids).

In addition, NADP-dependent isocitrate dehydrogenase, l,l-diaminopimelate

aminotransferase (chloroplast), and 5-methyltetrahydroproylglutamate-homocysteine

methyltransferase in the amino acid biosynthesis pathway were also differentially

expressed in seeds between castor offspring and parents.

The high expression of cysteine protease and 26S protease regulatory subunit

6a can lead to abnormal degradation of some regulatory proteins in anther tissue and

cause pollen abortion. In this study, these two proteins were downregulated in F_1

compared with their male parent [49]. Importantly, we found that cysteine protease

inhibitor was upregulated in the offspring seeds compared with their parents. This

enzyme can bind to cysteine protease to inhibit its activity and reduce the risk of

pollen abortion, and it is involved in the stress responses of plants under adverse

conditions and participates in physiological processes such as senescence of plants

[50].

Conclusions

To conduct a proteomic analysis to compare the differences in castor seeds at

different developmental stages between parents and their offspring, we first optimized

our two-dimensional electrophoresis system. The protein atlas obtained from

SDS-PAGE following IEF was the best when using samples after impurity removal,

24-cm dry gel strips (pH 3-10 NL), and a protein loading amount of 800 μg.

By 2D-DIGE analysis under these optimal two-dimensional electrophoresis,

93 DEPs were identified in the seeds at different developmental stages between castor

parents and offspring. They belonged to five categories: proteins related to stress

response (61.29%), seed nutrient storage (17.20%), fatty acid synthesis and

metabolism (12.90%), amino acid metabolism (6.45%), and others (2.15%).

RT-qPCR was performed on 12 DEPs, including 1) stearoyl-ACP desaturase,

OBAP2C, and acetyl-CoA carboxylase, which were upregulated in the offspring seeds

compared with their parents and are involved in oil synthesis and fatty acid synthesis

and metabolism; 2) 26S protease regulatory subunit 6a and cysteine protease, which

were downregulated in the offspring seeds compared with their parents and are

involved in pollen abortion; and 3) SOD, starch synthase, D-7 LEA protein,

glutathione peroxidase, ADH1, 2S albumin precursor, and APs, which were

upregulated in the offspring seeds compared with their parents and are involved in

plant resistance to adverse conditions. The RT-qPCR results showed that the

expression levels of the genes encoding these 12 DEPs were consistent with the

expression of the DEPs.

In summary, this study can help guide the production of castor beans and lays

the foundation for cultivating new castor varieties with high yield, high oil content,

and strong stress resistance.

References

1. Yu A, Li F, Liu A. Comparative proteomic and transcriptomic analyses provide new insight into the formation of seed size in castor bean. BMC Plant Biol. 2020;20: 48.

2. Olivares AR, Carrillo-González R, González-Chávez MdCA, Soto Hernández RM. Potential of castor bean (Ricinus communis L.) for phytoremediation of mine tailings and oil production. J Environ Management. 2013;114: 316-323.

3. Peng M, Xiao-Hua Z, Feng-Lan H, et al. Effect of different treatments on seeds germination of castor. Heilongjiang Agricultural Sciences. 2012.

4. Zhu MM, Liu EQ, Bao Y, Duan SL, She J, Liu H, et al. Low concentration of corn steep liquor promotes seed germination, plant growth, biomass production and flowering in soybean. Plant Growth Regul. 2019;87: 29-37.

5. Iimure T, Sato K. Beer proteomics analysis for beer quality control and malting barley breeding. Food Res Int. 2013;54: 1013-1020.

6.   Kim ST, Kim SG, Agrawal GK, Kikuchi S, Rakwal R. Rice proteomics: a

model system for crop improvement and food security. Proteomics. 2014;14:

593-610.

7.   Sghaier-Hammami B, Castillejo MÁ, Baazaoui N, Jorrín-Novo JV, Escandón

M. GeLC-Orbitrap/MS and 2-DE-MALDI-TOF/TOF comparative proteomics

analysis of seed cotyledons from the non-orthodox Quercus ilex tree species. J

Proteom. 2021;233: 104087.

8.   Jian M, Zhang D, Wang X, Wei S, Zhao Y, Ding Q, et al. Differential

expression pattern of the proteome in response to cadmium stress based on

proteomics analysis of wheat roots. BMC Genom. 2020;21: 343-343.

9.   Xia Q, Ponnaiah M, Cueff G, Rajjou L, Prodhomme D, Gibon Y, et al.

Integrating proteomics and enzymatic profiling to decipher seed metabolism

affected by temperature in seed dormancy and germination. Plant Sci.

2018;269: 118-125.

10.   Gu J, Hou D, Li Y, Chao H, Zhang K, Wang H, et al. Integration of proteomic

and genomic approaches to dissect seed germination vigor in Brassica napus

seeds differing in oil content. BMC Plant Biol. 2019;19: 21.

11.   Azarabadi S, Abdollahi H, Torabi M, Salehi Z, Nasiri J. ROS generation,

oxidative burst and dynamic expression profiles of ROS-scavenging enzymes

of superoxide dismutase (SOD), catalase (CAT) and ascorbate peroxidase

(APX) in response to Erwinia amylovora in pear (Pyrus communis L). Eur J

Plant Pathol. 2016;147: 279-294.

12.   Lu BW, An FX, Cao LJ, Yang YJ, Liu PM, Wang X, et al. Proteomic profiling

627   uncovered the cytosolic superoxide dismutase BsSOD1 associated with plant

628   defence in the herbal orchid Bletilla striata. Funct Plant Biol. 2020;47: 937.

629   13.   Jafari S, Garmdareh SEH, Azadegan B. Effects of drought stress on

630   morphological, physiological, and biochemical characteristics of stock plant

631   (Matthiola incana L.). Sci Hortic. 2019;253: 128-133.

632   14.   Pour-Aboughadareh A, Etminan A, Abdelrahman M, Siddique KHM, Tran

633   LSP. Assessment of biochemical and physiological parameters of durum

634   wheat genotypes at the seedling stage during polyethylene glycol-induced

635   water stress. Plant Growth Regul. 2020;92: 81-93.

636   15.   Che Y, Zhang N, Zhu X, Li S, Wang S, Si H. Enhanced tolerance of the

637   transgenic potato plants overexpressing Cu/Zn superoxide dismutase to low

638   temperature. Sci Hortic. 2020;261: 108949.

639   16.   Soares A, Carlton SMR, Simões I. Atypical and nucellin-like aspartic

640   proteases: emerging players in plant developmental processes and stress

641   responses. J Exp Bot. 2019;70: 2059-2076.

642   17.   Yang Y, Feng D. Genome-wide identification of the aspartic protease gene

643   family and their response under powdery mildew stress in wheat. Mol Biol

644   Rep. 2020;47: 8949-8961.

645   18.   Sebastián DI, Fernando FD, Raúl DG, Gabriela GM. Overexpression of

646   Arabidopsis aspartic protease APA1 gene confers drought tolerance. Plant Sci.

647   2020;292: 110406.

648   19.   Park SC, Kim YH, Jeong JC, Kim CY, Lee HS, Bang JW, et al. Sweetpotato

649   late embryogenesis abundant 14 (IbLEA14) gene influences lignification and

650   increases osmotic- and salt stress-tolerance of transgenic calli. Planta.

651   2011;233: 621-634.

652   20.   Li N, Zhang S, Liang Y, Qi Y, Chen J, Zhu W, et al. Label-free quantitative

653   proteomic analysis of drought stress-responsive late embryogenesis abundant

654   proteins in the seedling leaves of two wheat (Triticum aestivum L.) genotypes.

655   J Proteom. 2018;172: 122-142.

656   21.   Zhao W, Yao F, Zhang M, Jing T, Zhang S, Hou L, et al. The potential roles

657   of the G1LEA and G3LEA proteins in early embryo development and in

658   response to low temperature and high salinity in artemia sinica. PLoS One.

659   2016;11: e0162272.

660   22.   Eshdat Y, Holland D, Faltin Z, Ben-Hayyim G. Plant glutathione peroxidases.

661   Physiol Plant. 1997;100: 234-240.

662   23.   Kammoun M, Essid MF, Ksouri F, Rokka V-M, Charfeddine M,

663   Gargouri-Bouzid R, et al. Assessment of physiological age and antioxidant

664   status of new somatic hybrid potato seeds during extended cold storage. J

665   Plant Physiol. 2020;254: 153279.

666   24.   Kim YH, Huh GH. Overexpression of cinnamyl alcohol dehydrogenase gene

667   from sweetpotato enhances oxidative stress tolerance in transgenic

668   Arabidopsis. Vitro Cell Dev Biol Plant. 2019;55: 172-179.

25.   Zhuang M, Lam SK, Zhang J, Li H, Shan N, Yuan Y, et al. Effect of full

substituting compound fertilizer with different organic manure on reactive

nitrogen losses and crop productivity in intensive vegetable production system

of China. J Environ Manag. 2019;243: 381-384.

26.   Rattanapolsan L, Nakbanpote W, Sangdee A. Zinc- and cadmium-tolerant

endophytic bacteria from Murdannia spectabilis (Kurz) Faden. studied for

plant growth-promoting properties, in vitro inoculation, and antagonism. Arch

Microbiol. 2020;203: 1131-1148.

27.   Wang CL, Zhang ZP, Miao MM. SNF1-related protein kinase (SnRK) 1

involved in the regulation of raffinose family oligosaccharide metabolism in

cucumber (Cucumis sativus L.) Calli. J Plant Growth Regul. 2016;35:

851-864.

28.   Mehrpouyan S, Menon U, Tetlow IJ, Emes MJ. Protein phosphorylation

regulates maize endosperm starch synthase IIa activity and protein−protein

interactions. Plant J. 2021;105: 1098-1112.

29.   do Carmo CD, Sousa MB, Brito AC, de Oliveira EJ. Genome-wide association

studies for waxy starch in cassava. Euphytica. 2020;216: 82.

30.   Nguyen TP, Cueff G, Hegedus DD, Rajjou L, Bentsink L. A role for seed

storage proteins in Arabidopsis seed longevity. J Exp Bot. 2015;66:

6399-6413.

31.   Lopes FES, da Costa HPS, Souza PFN, Oliveira JPB, Ramos MV, Freire JEC,

et al. Peptide from thaumatin plant protein exhibits selective anticandidal

activity by inducing apoptosis via membrane receptor. Phytochemistry. 2019;159: 46-55.

32. Wang C, Yang C, Gao C, Wang Y. Cloning and expression analysis of 14 lipid transfer protein genes from Tamarix hispida responding to different abiotic stresses. Tree Physiol. 2009;29: 1607-1619.

33. Whittle EJ, Cai Y, Keereetaweep J, Chai J, Buist PH, Shanklin J. Castor stearoyl-ACP desaturase can synthesize a vicinal diol by dioxygenase chemistry. Plant Physiol. 2020;182: 730-738.

34. de Oliveira TS, Freitas-Silva O, Kluczkovski AM, Campelo PH. Potential use of vegetable proteins to reduce Brazil nut oil oxidation in microparticle systems. Food Res Int. 2020;137: 109526.

35. Zhou LZ, Chen FS, Hao LH, Du Y, Liu C. Peanut oil body composition and stability. J Food Sci. 2019;84: 2812-2819.

36. Jolivet P, Acevedo F, Boulard C, d'Andréa S, Faure J-D, Kohli A, et al. Crop seed oil bodies: from challenges in protein identification to an emerging picture of the oil body proteome. Proteomics. 2013;13: 1836-1849.

37. Parvy JP, Napal L, Rubin T, Poidevin M, Perrin L, Wicker-Thomas C, et al. Drosophila melanogaster Acetyl-CoA-carboxylase sustains a fatty acid-dependent remote signal to waterproof the respiratory system. PLoS Genet. 2012;8: e1002925.

38. Vijayakumar A, Olson I, Wang T, Hollenback D, Toteva M, Trevaskis J, et al. Efficacy and safety of an acetyl CoA carboxylase inhibitor are improved in

combination with PPAR agonists in a dyslipidemic rat model. J Hepatol. 2020;73: S656.

39. Thelen JJ, Ohlrogge JB. Both antisense and sense expression of biotin carboxyl carrier protein isoform 2 inactivates the plastid acetyl-coenzyme A carboxylase inArabidopsis thaliana. Plant J. 2002;32: 419-431.

40. Oaks Z, Jimah J, Grossman CC, Beckford M, Kelly R, Banerjee S, et al. Transaldolase haploinsufficiency in subjects with acetaminophen-induced liver failure. J Inherit Metab Dis. 2020;43: 496-506.

41. Zheng M, Zhu C, Yang T, Qian J, Hsu Y-F. GSM2, a transaldolase, contributes to reactive oxygen species homeostasis in Arabidopsis. Plant Mol Biol. 2020;104: 39-53.

42. Tylki-Szymanska A, Wamelink MMC, Stradomska TJ, Salomons GS, Taybert J, Dąbrowska-Leonik N, et al. Clinical and molecular characteristics of two transaldolase-deficient patients. Eur J Pediatr. 2014;173: 1679-1682.

43. Li M, Zhang X, Lu Y, Meng S, Quan H, Hou P, et al. The nuclear translocation of transketolase inhibits the farnesoid receptor expression by promoting the binding of HDAC3 to FXR promoter in hepatocellular carcinoma cell lines. Cell Death Dis. 2020;11: 31.

44. Bi H, Li F, Wang H, Ai X. Overexpression of transketolase gene promotes chilling tolerance by increasing the activities of photosynthetic enzymes, alleviating oxidative damage and stabilizing cell structure in Cucumis sativus L. Physiol Plant. 2019;167: 502-515.

45. Seong ES, Jeon MR, Choi JH, Yoo JH, Lee JG, Na JK, et al. Overexpression of S-Adenosylmethionine synthetase enhances tolerance to cold stress in Tobacco. Russ J Plant Physiol. 2020;67: 242-249.

46. Mo HJ, Sun YX, Zhu XL, Wang XF, Zhang Y, Yang J, et al. Cotton S-adenosylmethionine decarboxylase-mediated spermine biosynthesis is required for salicylic acid- and leucine-correlated signaling in the defense response to Verticillium dahliae. Planta. 2016;243: 1023-1039.

47. Elango R. Methionine nutrition and metabolism: insights from animal studies to inform human nutrition. J Nutr. 2020;150: 2518S-2523S.

48. Kovács L, Mendel Á, Szentgyörgyi A, Fekete S, Söre F, Posta K, et al. Comparative analysis of overexpressed Fragaria vesca S-adenosyl-l-methionine synthase (FvSAMS) and decarboxylase (FvSAMDC) during salt stress in transgenic Nicotiana benthamiana. Plant Growth Regul. 2020;91: 53-73.

49. Passarge A, Demir F, Green K, Depotter JRL, Scott B, Huesgen PF, et al. Host apoplastic cysteine protease activity is suppressed during the mutualistic association of Lolium perenne and Epichloë festucae. J Exp Bot. 2021;72: 3410-3426.

50. Kaur H, Gill RS, Kaur S. Rice bean (Vigna umbellata Thunb. Ohwi and Ohashi) protection against Callosobruchus maculatus F." by presence of protein profile. J Stored Prod Res. 2020;86: 101574.