Individual-based quantitative temporal beta-diversity:

Novel indices for assessing community compositional shifts accounting for

changes in the properties of individuals

Running title: Individual-based quantitative beta-diversity

Ryosuke Nakadai[*1]

ORCID: 0000-0002-9512-8511

[1] Biodiversity Division, National Institute for Environmental Studies, Onogawa 16-2,

Tsukuba, Ibaraki, 305-8506, Japan

* Corresponding author: R. Nakadai

E-mail: r.nakadai66@gmai.com, nakadai.ryosuke@nies.go.jp

Biodiversity Division, National Institute for Environmental Studies, Onogawa 16-2,

Tsukuba, Ibaraki, 305-8506, Japan

Author contributions

R.N. conceived the study, designed the methods described herein, analysed the

empirical data, and wrote the manuscript.

Acknowledgments

Financial support was provided by the Japan Society for the Promotion of Science

and a Program of the National Institute for the Environmental Studies, Japan.

Conflicts of interest

The author declares no conflicts of interest.

Data Availability

Data are available from the package rFIA (Stanke et al., 2020) in R, and the code

used to calculate the indices introduced in the present study is provided in

Supplementary File 1.

Biosketch

Ryosuke Nakadai is a Research Associate at the National Institute for Environmental

Studies, Japan, with a broad interest in biodiversity, biological interactions, and

ecosystem services across temporal and spatial scales. His work includes analyses of

biodiversity patterns and the development of both statistical and molecular tools for a

wide range of taxa.

Individual-based quantitative temporal beta-diversity:

Novel indices for assessing community compositional shifts explicitly accounting

for changes in the properties of individuals

Running title: Individual-based quantitative beta-diversity

Abstract

Aim

Revealing patterns of temporal biodiversity change is an urgent issue in light of the

effects of global climate change. In addition, climate change is known to modify not

only biodiversity but also individual life-history. Nevertheless, although the

connection between individual physiology and large-scale ecological patterns is a

major challenge for the mechanistic understanding of macroecology, large gaps exist

between individual life-history and biodiversity. In this context, novel methods can be

used to evaluate changes in individual properties and their effects on biodiversity.

Innovation

Novel indices that can evaluate compositional shifts across time by explicitly

considering the contributions of the life-histories (i.e., growth, mortality, and

recruitment) of individuals in a community were successfully developed. By applying

subset data of the Forest Inventory and Analysis database for the state of Rhode

Island in the USA, changes in the contribution of individual life-histories to

biodiversity change were successfully identified.

Main conclusions

The development of the proposed novel indices not only represents methodological

progress in community ecology and macroecology but also a conceptual advance,

bridging studies of biodiversity and individual life-history and physiology. The

individual-based diversity index opens the door to the possibility of individual-based

biodiversity science, which could facilitate understanding the comprehensive effects

of climate change across different hierarchies of biological organisation.

Keywords: climate change, compositional shift, life-history, growth, temporal beta

diversity

Introduction

Elucidating patterns of biodiversity change over time in light of the effects of global climate change and anthropogenic disturbance has been the focus of much attention (Dornelas et al., 2014; García et al., 2016; Vellend et al., 2013). Classical studies have focused on biodiversity loss (i.e., alpha and gamma diversity), and a lack of knowledge on compositional shifts (i.e., beta diversity) has been highlighted until relatively recently (Dornelas et al., 2014; Magurran et al., 2015, 2019). Within this context, several studies have attempted to quantify compositional shifts in biodiversity, termed temporal beta diversity (Gotelli et al., 2017; Magurran, 2011). Recent studies have developed unique methodologies and concepts for temporal beta diversity based on classical spatial beta diversity research (Gotelli et al., 2021; Legendre, 2019; Shimadzu, et al., 2015; Tatsumi et al., 2021).

Because an individual cannot exist in two places at once, it can persist in the same habitat over time (Magurran et al., 2019). Therefore, individual persistence and turnover are important factors for temporal beta diversity. In fact, some individuals of long-lived species (e.g., trees and vertebrates) persist over the long term, whereas individuals of short-lived species are replaced by conspecifics or individuals of other species (Nakadai, 2020, 2021). In this context, Nakadai (2020) recently proposed the "individual-based beta-diversity" concept to bridge the gap between biodiversity and individual information, and provided novel indices to evaluate individual turnover and compositional shifts based on comparisons between two periods at a given site. Specifically, I developed an individual-based temporal beta-diversity approach for the Bray-Curtis dissimilarity index (i.e., an abundance-based temporal beta-diversity index) (Nakadai, 2020). Subsequently, Nakadai (2020, 2021) added both persistence and turnover of individuals to the compositional shifts in biodiversity across time.

Climate change is known to modify not only biodiversity directly but also the individual life-histories of organisms (Brienen et al., 2020; Searle & Chen, 2018).

Recent studies have shown that climate warming will facilitate faster growth and

shorter life spans for individual trees as a physiological process within individuals

(Körner, 2017; Munné-Bosch, 2018; Searle & Chen, 2018) and increase the amount of

carbon stocks in terrestrial ecosystems (Brienen et al., 2015). As a result, these changes

in individual life-history potentially affect the compositional shifts of local

communities and, consequently, global biodiversity. Nevertheless, although the

connection between individual physiology and large-scale ecological patterns is a

major challenge for mechanistic understanding in macroecology (Brown, 1995, 1999;

Brown et al. 2004; Hubbell, 2001; e.g., metabolic theory), large gaps still exist between

individual life-history and biodiversity. To comprehensively understand the effects of

climate change, it is essential to simultaneously capture all the property changes of

individuals, individual turnover, and changes in biodiversity.

Despite these needs, the approaches of previous individual-based temporal beta-

diversity (Nakadai, 2020, 2021) only take into account the presence or absence of each

individual (i.e., qualitative properties such as 'dead' or 'alive'), and the influences of

the property changes of each individual (e.g., individual growth) over time on

biodiversity have been neglected, which is further influenced by global climate change.

Therefore, the purpose of this study was to develop new individual-based temporal

beta-diversity indices that can assess compositional shifts across time by explicitly

including the property changes of each individual in a community. Hereafter, I call the

indices developed in Nakadai (2020) and the novel indices introduced in the present

study individual-based "qualitative (QL)" indices and individual-based "quantitative

(QT)" indices, respectively. By introducing quantitative individual information (e.g.,

size and weight), these novel indices can evaluate the contribution of individual growth

and the differences between large and small individuals. Thus, after briefly reviewing

the history of pairwise beta-diversity indices, I extend individual-based QL temporal

beta-diversity indices to QT indices. I then apply these indices to a subset of the Forest

Inventory and Analysis (FIA) database for the state of Rhode Island in the USA (Stanke

et al., 2020, Burrill et al., 2021) as a case study to demonstrate their application.

Materials and Methods

Conventional incidence-based and abundance-based indices and recently developed

individual-based qualitative (QL) temporal beta-diversity indices

Incidence-based and abundance-based beta-diversity indices have been previously

proposed (Baselga, 2013). For example, the widely used Bray-Curtis dissimilarity

index (Odum, 1950) is an abundance-based extension of the Sørensen index (Baselga,

2013; Legendre & Legendre, 2012; Sørensen, 1948). Incidence-based methods target

the number of species, whereas abundance-based methods mainly target the number

of individuals. It should be noted that abundance-based methods sometimes target

area, biomass, weight, etc. (e.g., Li et al., 2016). I describe this approach in more

detail in the present paper. Conventional beta-diversity indices are based on

intersection (A, a) and relative complement (B, b, C, c) components. Hereafter, here I

explain only the context of temporal beta diversity. Specifically, the Sørensen

dissimilarity index (d_{sor}; Sørensen, 1948) is formulated as follows:

$$d_{sor} = \frac{b+c}{2a+b+c} \qquad\qquad \text{equation 1}$$

where a is the number of species common to both temporal units; b is the number of

species that occur in the first but not the second unit; and c is the number of species

that occur in the second unit but not the first unit. Analogously, the Bray-Curtis

dissimilarity index (Odum, 1950) is formulated as follows:

$$d_{BC} = \frac{B+C}{2A+B+C}. \qquad\qquad \text{equation 2}$$

As a premise for explanation, assume that there are two sets of community data along the

time axis—communities at time j and time k (abbreviated as Tj and Tk: $j < k$), respectively.

The abundance-based components are as follows.

$A = \sum_i \min(x_{ij}, x_{ik})$ equation 3a

$B = \sum_i [x_{ij} - \min(x_{ij}, x_{ik})]$ equation 3b

$C = \sum_i [x_{ik} - \min(x_{ij}, x_{ik})]$ equation 3c

where x_{ij} is the abundance of species i at temporal unit j, and x_{ik} is the abundance of

species i at temporal unit k. Therefore, A is the number of individuals of each species

shared by units j and k, and B and C are the numbers of individuals unique to units j

and k, respectively.

Nakadai (2020) introduced the categorisation of the community dynamics

component into persistence (p), mortality (m), and recruitment (r) in individual-based

temporal beta diversity. m refers to individuals occurring at Tj that die before Tk,

thereby representing the total number of dead individuals during the time between Tj

and Tk. r identifies individuals that were not alive at Tj (or were not yet in the size class

counted) but were subsequently alive and counted at Tk, thereby representing the total

number of recruited individuals between Tj and Tk. Component p refers to the

persistence of individuals from Tj to Tk. Note that, unlike previous studies (Nakadai,

2020, 2021: Fig. 1a), I use lower-case letters for the components of individual-based

qualitative indices (i.e., p, m, r) to distinguish the components of the individual

quantitative indices described later. Thus, the abundances are calculated as follows:

$p = \sum_o \min(z_{oj}, z_{ok})$ equation 4a

$m = \sum_o [z_{oj} - \min(z_{oj}, z_{ok})]$ equation 4b

$r = \sum_o [z_{ok} - \min(z_{oj}, z_{ok})]$, equation 4c

where z_{oj} is individual o at time Tj, and z_{ok} is individual o at time Tk. Both z_{oj} and z_{ok}

have values of one or zero (presence and absence, respectively). Therefore, p is the

total number of individuals present at both Tj and Tk, and m and r are the numbers of

individuals unique to Tj and Tk, respectively (Nakadai, 2020, 2021: Fig. 1a).

This formulation of the individual-based temporal beta-diversity index for the Bray-

Curtis (d_{mr}) dissimilarity can be expressed as follows:

$d_{mr} = \dfrac{m+r}{2p+m+r}$ equation 5

In some cases, there is a turnover of different individuals belonging to the same

species, even when the community composition is stable over time. Therefore, the

components that change over time (m and r) are reorganised into two further

components, namely compositional equilibrium (e) and shift (B and C), respectively

(Nakadai, 2020, 2021). Component m can be partitioned into lost individuals that

contribute to compositional shifts in a community (B) and lost individuals replaced by

conspecifics that contribute to equilibrium in community composition (e_{loss}) (Fig. 1a).

Similarly, component r can be partitioned into gained individuals that contribute to

compositional shifts in a community (C) and gained individuals that replace

conspecifics, thereby contributing to equilibrium in community composition (e_{gain})

(Fig. 1a). By definition, the number of e_{loss} and e_{gain} individuals is identical; thus, I

replaced both e_{loss} and e_{gain} with the coefficient e (Nakadai, 2020, 2021). Furthermore,

components m and r can be analysed in the same manner as the community data

(Nakadai, 2020). Therefore, the Bray-Curtis dissimilarity between communities can be

calculated based on m and r as follows:

$v_{s.QL} = \dfrac{B+C}{2e+B+C} = \dfrac{B+C}{m+r}$ equation 6

where $v_{s.QL}$ indicates the speed of compositional shifts in a community relative to the total

speed of individual turnover associated with m and r. It is also possible to calculate $v_{s.QL}$

as d_{BC} divided by d_{mr}, as follows:

$$d_{BC} \div d_{mr} = \frac{B+C}{2A+B+C} \div \frac{m+r}{2p+m+r} = \frac{B+C}{2(p+e)+B+C} \div \frac{m+r}{2p+m+r} = \frac{B+C}{2o+m+r} \div \frac{m+r}{2p+m+r} =$$

$$\frac{B+C}{2e+B+C} = v_{s.QL}. \quad \text{equation 7}$$

Novel individual-based "qualitative (QT)" beta-diversity indices

I developed individual-based QT components and novel indices for community

datasets over time using individual property information (e.g., size and weight). The

relationship between the QL and QT indices is similar to that between the incident-

based and abundance-based indices. In other words, individual-based QT components

were calculated as individual-based QL weighted by individual properties. As

previously noted, the abundance targeted by abundance-based indicators is not

necessarily the number of individuals but can also include area, volume, etc.

Abundance except for the number of individuals for community datasets has been

used in cases of large size variation among individuals within a community or where

it is difficult to recognise proper individuals (Li et al., 2016). Hereafter, the Bray-

Curtis indices targeting measures except the number of individuals (e.g., sizes and

weights) are described as $d_{BC'}$ and their components are described with apostrophes

(A', B', and C') to distinguish the unweighted Bray-Curtis index as follows:

$$d_{BC'} = \frac{B'+C'}{2A'+B'+C'}. \qquad \text{equation 8}$$

This weighting indicates that abundance-based (number of individual-based)

diversity indices were weighted by total measures of individuals for each species,

thus calculating diversity indices by considering the variation of measures of each

individual implicitly. Because individuality is rarely clearly identifiable in weighted

indices, this relationship is not well recognised.

In addition to individual-based QL indices, individual-based QT indices comprise

components of mortality and recruitment (M and R, respectively). In the QT indices,

M refers to the total number of measures of individuals occurring at Tj that die before

Tk, thereby representing the total number of measures of the mortal individuals

between Tj and Tk (Fig. 1b). R identifies the total number of measures of individuals

that were not alive at Tj (or were not yet in the size class counted) but were

subsequently alive and counted at Tk, thereby representing the total number of

measures of the recruited individuals between Tj and Tk (Fig. 1b).

From this point on, the major differences between the QL and QT indices are as

follows. The persistent individuals are qualitatively identical, thus keeping their alive

(or present) status between Tj and Tk, but rarely maintain the same status in terms of

size and weight; thus, there is a change in individual properties between Tj and Tk.

For example, in tree communities, one newly recruited small tree can be a large tree

after the persistence of half a century. In the p component of QL indices, this means

that an individual is persistent, while it is impossible to evaluate changes in the

features (i.e., basal area, biomass, and/or weight, etc.) of the individual. The indices

introduced here overcome this problem by considering the contribution of changes in

individual properties, mainly targeting individual growth. Specifically, in the case of

persistent individuals, the original and altered total number of measures are

partitioned into persistent (P) and growth (G) components, respectively. Thus, the

respective total number of measures of the components were calculated as follows:

$M = \sum_o \left[w_{oj} - \min(w_{oj}, w_{ok}) \times \left[z_{oj} - \min(z_{oj}, z_{ok}) \right] \right]$ equation 9a

$R = \sum_o \left[\left[w_{ok} - \min(w_{oj}, w_{ok}) \right] \times \left[z_{ok} - \min(z_{oj}, z_{ok}) \right] \right]$ equation 9b

$P = \sum_o \min(w_{oj}, w_{ok})$ equation 9c

$$G = \sum_o \left[\left[\max(w_{oj}, w_{ok}) - \min(w_{oj}, w_{ok}) \right] \times \min(z_{oj}, z_{ok}) \right]$$ equation

9d,

where w_{oj} is the measure of individual o at time Tj, and w_{o2} is the measure of

individual o at time Tk. M and R are the total number of measures of mortal and

recruited individuals, respectively. The persistent and growth parts represent the

unchanged and changed parts of measures within an individual between Tj and Tk,

respectively. Thus, P and G represent the total amount of these components within a

community. Equation 9d also considers decreasing changes in the measures of each

individual. Therefore, the calculation of the G component can be partitioned into

increasing and decreasing parts, as follows:

$$G = \sum_o \left[\left[\max(w_{oj}, w_{ok}) - \min(w_{oj}, w_{ok}) \right] \times \min(z_{oj}, z_{ok}) \right] = G_{in} + G_{de}$$

equation 10,

where G_{in} and G_{de} are the total increased and decreased parts of the targeted

measures, respectively. Specifically, the formulas for G_{in} and G_{de} are as follows:

$$G_{in} = \sum_o \left[\left[w_{ok} - \min(w_{oj}, w_{ok}) \right] \times \min(z_{oj}, z_{ok}) \times \frac{1}{2} \left[1 + \text{sign}(w_{ok} - w_{oj}) \right] \right]$$

equation 11a

$$G_{de} = \sum_o \left[\left[w_{oj} - \min(w_{oj}, w_{ok}) \right] \times \min(z_{oj}, z_{ok}) \times \frac{1}{2} \left[1 - \text{sign}(w_{ok} - w_{oj}) \right] \right]$$

equation 11b,

where $\frac{1}{2} \left[1 + \text{sign}(w_{ok} - w_{oj}) \right]$ and $\frac{1}{2} \left[1 - \text{sign}(w_{ok} - w_{oj}) \right]$ distinguish the large

and small relationships between w_{ok} and w_{oj} by returning 1 or 0.

This formulation of the individual-based QT temporal beta-diversity index for

Bray-Curtis dissimilarity (d_{MRG}) can be expressed as follows:

$$d_{MRG} = \frac{M+R+G}{2P+M+R+G}.$$ equation 12

Furthermore, this individual-based QT temporal beta diversity can be partitioned

into the relative contributions of mortality (d_M), recruitment (d_R), and growth (d_G) to

the target weights or measurements, as follows:

$$d_{MRG} = \frac{M+R+G}{2P+M+R+G}$$

$$= \frac{M}{2P+M+R+G} + \frac{R}{2P+M+R+G} + \frac{G}{2P+M+R+G} = d_M + d_R + d_G.$$ equation 13

In addition, as shown in equations 11a and b, the G component can be partitioned into

G_{de} and $G_{in,}$ which are the total decrease and increase, respectively, in the targeted

measures. Equation 13 can then be partitioned into two components, as follows:

$$d_{MRG} = \frac{M+R+G}{2P+M+R+G}$$

$$= \frac{M+G_{de}}{2P+M+R+G} + \frac{R+G_{in}}{2P+M+R+G} = d_{de} + d_{in}.$$ equation 14

where d_{de} and d_{in} indicate the contributions of the decrease and increase, respectively,

to the total changes in the targeted measures in Tj and Tk. In addition to the QL indices

(Nakadai, 2020, 2021), the components that change over time (M, R, and G) can be

reorganised into two components—compositional equilibrium (E) and shift (B' and C'),

respectively. Components $M + G_{de}$ can be reorganised as the total number of lost

measures that contribute to compositional shifts in a community (B') and the total

number of lost measures replaced by conspecifics, contributing to the equilibrium in

community composition (E_{loss}) (Fig. 1b). Similarly, component $R + G_{in}$ can be

reorganised as the total number of measures that contribute to compositional shifts in

a community (C') and the total number of gained measures replacing conspecifics,

contributing to equilibrium in community composition (E_{gain}) (Fig. 1b). By definition,

the total amount of E_{loss} and E_{gain} are identical; hence, I replaced both E_{loss} and E_{gain}

with coefficient E.

$$d_{MRG} = \frac{M+R+G}{2P+M+R+G} = \frac{M+G_{de}}{2P+M+R+G} + \frac{R+G_{in}}{2P+M+R+G}$$

$$= \frac{E+B}{2P+M+R+G} + \frac{E+C}{2P+M+R+G} = \frac{2E}{2P+M+R+G} + \frac{B+C}{2P+M+R+G} = d_S + d_E.$$

equation 15

Furthermore, components M, R, and G (i.e., $M + G_{de}$ and $R + G_{in}$) can be analysed in

the same manner as the community data. Therefore, the Bray-Curtis dissimilarity

between communities (i.e., the total amount of lost and gained measures) can be

calculated based on M, R, and G, as follows:

$$v_{s.QT} = \frac{B'+C'}{2E+B'+C'} = \frac{B'+C'}{M+R+G} \qquad\qquad \text{equation 16}$$

where $v_{s.QT}$ indicates the relative speed of compositional shifts in a community against

the total turnover of targeted measures associated with M, R, and G. If $v_{s.QT}$ is 0, then the

turnover of the targeted measures would not contribute to compositional shifts in a

community. If $v_{s.QT}$ is 1, all turnover of the targeted measures contributes to the

compositional shifts. It is also possible to calculate $v_{s.QT}$ as $d_{BC'}$ divided by d_{MRG}, as

follows:

$$d_{BC'} \div d_{MRG} = \frac{B'+C'}{2A'+B'+C'} \div \frac{M+R+G}{2P+M+R+G} = \frac{B'+C'}{2(P+E)+B'+C'} \div \frac{M+R+G}{2P+M+R+G} = \frac{B'+C'}{2P+M+R+G} \div$$

$$\frac{M+R+G}{2P+M+R+G} = \frac{B'+C'}{M+R+G} = v_{s.QT}. \quad \text{equation 17}$$

Here, I have only focused on an extension of Bray-Curtis dissimilarity. In

Supplementary Text 1, I also summarise individual-based beta diversity using the

Ružička dissimilarity index (Ružička, 1958), which was identified by Podani et al.

(2013) and Legendre (2014) as an appropriate index for beta diversity studies. The

Ružička dissimilarity index is an abundance-based extension of the Jaccard

dissimilarity index (Jaccard, 1901). Nevertheless, I recommend using the Bray-Curtis

family of indices in individual-based temporal beta diversity because the formulations

based on these indices are simpler than the Ružička indices, especially with respect to

the derivation $v_{s.QL}$ and $v_{s.QT}$.

Case study: Subset of the Forest Inventory and Analysis (FIA) database for the state of

Rhode Island in the USA

As a case study application of the new individual-based QT indices based on publicly

available data, here I focus on a subset of data from the United States Department of

Agriculture Forest Service's FIA Program for the state of Rhode Island using the R

package rFIA (Stanke et al., 2020). The FIA database is a well-known dataset that

includes individual tree tracking and size information (i.e., basal area and bole

biomass). A comprehensive summary of the associated data and sampling methodology

is provided by Burrill et al. (2021). Briefly, each FIA plot comprises four 0.017-ha

circular subplots, each located within a circular 0.10 ha macroplot. All free-standing

woody stems (alive and dead) with a diameter of ≥ 12.7 cm were sampled within each

subplot (Burrill et al., 2021).

The original dataset included 10,093 individual-time records for 58 species,

including multiple species (spp.) in 140 plots from 2004 to 2018. Some plots included

more than two surveys, for which it was possible to calculate more than one value of

temporal beta diversity. I conducted plot-level filtering of the dataset using the

following criteria: (1) those with unidentified individuals at the species level were

excluded; (2) those with ten individuals or more were included; and (3) a survey

interval between two surveys of 5 ± 1 years was targeted. In addition, individual trees

with a 12.7-cm diameter at breast height (DBH) and above were targeted because the

bole biomass was also estimated. The final dataset included 5,705 individual-time

records comprising 2,541 individual trees of 32 species across 126 survey intervals (86

unique plots). The latitudinal and longitudinal ranges in the dataset are N 41.39446°–

42.00797° and 71.79635°–71.12814°W, respectively. The elevation range was 10–680

m. The geographical location of each plot is shown in Fig. S1.

I focused on the basal area and bole biomass of each tree as changes in individual

properties that can be considered in the novel indices developed in the previous

sections. The basal areas of live trees were calculated as $\frac{DIA}{2} \times \frac{DIA}{2} \times \pi$. The gross

cubic-foot volume of the live trees was estimated at ≥ 12.7 cm DBH, which defined

the total volume of wood in the central stem of the inventoried trees (bole biomass)

assuming a 30.48-cm mean stump height and a minimum 10.16 cm top diameter (for

detailed estimation method see Domke et al., 2013).

To demonstrate the application of the novel indices, I focused mainly on the

following 11 indices: d_{BC}, d_{mr}, d_m, d_r, $v_{s.QL}$, $d_{BC'}$, d_{MRG}, d_M, d_R, d_G, and $v_{s.QT}$. Formulas

of the base indices (i.e., d_{BC}, d_{mr}, $v_{s.QL}$, $d_{BC'}$, d_{MRG}, and $v_{s.QT}$) are summarised in Table

1. All 11 indices were calculated for each survey interval for each plot. For the latter

six indices (i.e., $d_{BC'}$, d_{MRG}, d_M, d_R, d_G, and $v_{s.QT}$) were calculated for both basal area and

bole biomass, and I added "_area" or "_biomass" to distinguish them, respectively.

First, I tested the relationships among d_{BC}, d_{mr}, $v_{s.QL}$, $d_{BC'}$, d_{MRG}, and $v_{s.QT}$ using simple

linear regression. Second, to reveal the geographical and temporal components of the

indices, I constructed models using linear mixed models and linear models for each

index. As explanatory variables, geographic factors (latitude and elevation), a temporal

factor (mean calendar year between the first and second surveys), and a sampling-

design-related factor (length of survey interval) were used. Plot identity was considered

a random effect in the model comparisons because 40 plots included two survey

intervals. However, the model comparisons showed that the linear mixed models

always had higher Akaike information criterion (AIC; Akaike, 1998) values than the

linear model (Table S1). Thus, all of the results shown in the following discussion are

based on linear models.

All analyses were conducted using R software (ver. 4.1.0; R Development Core

Team, 2021). The R packages *vegan* (Oksanen et al., 2020), *nlme* (Pinheiro et al., 2022),

and *MuMIn* (Barton, 2022) were used to convert the abundance data to presence-

absence data, construct a linear mixed model, and model comparisons based on AIC.

The R scripts used to calculate the new indices are provided in Supplementary File 1.

Results

Among 126 survey intervals, the individual turnover of 10 survey intervals was zero,

which were subsequently excluded from the targets when calculating the $v_{s.QL}$ indices.

The distribution of the novel indices, except $v_{s.QT}$, is shown in Fig. 2. The d_{MRG} indices

based on both basal area and bole biomass were strongly and positively correlated with

d_{BC}, d_{mr}, area-based $d_{BC'}$, and biomass-based $d_{BC'}$ ($\geq$ 0.87; Fig. S3). On the other hand,

the $v_{s.QT}$ indices based on both basal area and bole biomass were only weakly correlated

with the $d_{BC'}$ indices based on both basal area and bole biomass (r = 0.23–0.28, Fig.

S3) and showed no correlation with $v_{s.QL}$ (r = 0.16 and 0.08, Fig. S3). For both d_{MRG}

and $v_{s.QT}$, strong correlations existed between the indices based on basal area and bole

biomass (r = 0.99 and 0.91, Fig. S3). All the Pearson correlation coefficients are

summarised in Fig. S3.

As a result of AIC comparison, null models were chosen for the three indices, i.e.,

the indices for compositional shift based on number of individuals and basal area (d_{BC},

$d_{BC'\_area}$) and the relative speed of compositional shift against total changes in basal

area ($v_{s.QT\_area}$) (Table 2). The index of compositional shift based on bole biomass

($d_{BC'\_biomass}$) tended to increase with longer survey intervals as a result of model

comparison (Table 2). In the index for the relative speed of compositional shift against

total individual turnover ($v_{s.QL}$), only elevation was chosen as an explanatory variable

based on the AIC comparison; thus, the index increased toward higher elevations

(Table 2). For the relative speed of compositional shift against total changes in basal

area ($v_{s.QT\_biomass}$), a model including calendar year and survey interval was chosen by

the AIC comparison (Table 2); specifically, the index increased toward older calendar

years and longer survey intervals. In both indices of the total amount of change in basal

area and bole biomass (d_{MRG_area}, $d_{MRG\_biomass}$), models with latitude and survey interval

were chosen as the best models in the AIC comparisons (Table 2); specifically, the

indices tended to be higher toward lower latitudes and with longer survey intervals. On

the other hand, in the index of total individual turnover (d_{mr}), the model with only the

survey interval was chosen as the best model (Table 1), increasing toward longer survey

intervals. In the three indices related to the contribution of mortality to the total amount

of change (d_m, d_{M_area}, $d_{M\_biomass}$), models with latitude and calendar year were chosen

as the best models (Table 2), tending to be higher toward lower latitudes and more

recent years. In the indices related to the contribution of recruitment to the total amount

of change (d_{R_area}, $d_{R\_biomass}$), models with calendar year were selected as the best

models (Table 2), tending to be higher toward less recent years. In addition, in the index

related to the contribution of recruitment to the total individual turnover (d_r), the model

with calendar year and survey interval was chosen as the best model (Table 2), with the

index tending to be higher toward less recent years and at longer survey intervals. For

both indices related to the contribution of growth to the total amount of change (d_{G_area},

$d_{G\_biomass}$), models were chosen that included the survey interval and, in the case of

basal area, calendar year.

Discussion

The potential for evaluating the contributions of individuals' persistence to

compositional shifts in a community was first proposed by Nakadai (2020). Here, I

extended the previously established individual-based beta-diversity approaches

(Nakadai, 2020, 2021) to take account of individual property changes—even for

persistent individuals—that are not currently considered in qualitative indices.

Specifically, in addition to the components of mortality and recruitment as considered

in Nakadai (2020), I introduced a growth component to account for the property

changes individuals in the context of compositional shifts for the first time. Thus, the

novel indices proposed here not only represent methodological progress in

community ecology and macroecology but also provide a conceptual advance

bridging biodiversity and individual life-history and physiology studies.

To illustrate the application of the novel indices, I applied these to a subset of the FIA

database for the state of Rhode Island in the USA. This analysis demonstrated that the

contribution of tree growth to the total turnover of both basal area and bole biomass

was generally larger than that of mortality and recruitment (Fig. 2a, b: Fig. S2). This

suggests the importance of previously overlooked growth factors in forest turnover

and dynamics as well as subsequent compositional shifts. In the case study dataset,

the calculated abundance-based and individual-based indices (i.e., d_{BC}, d_{mr}, $d_{BC'}$, and

d_{MRG}) were strongly correlated. Thus, iff the contribution of individual tree growth is

high, the strength of the correlation between d_{mr} and d_{MRG} should decrease, yet this

was not the case. On the other hand, I found no clear relationships between the $v_{s.QL}$

index and $v_{s.QT}$ indices based on both basal area and bole biomass and the relative

speed of compositional shift. Generally, the values of $v_{s.QL}$ for the FIA dataset are

high; thus, individual turnover generally contributed to compositional shift, although

both $v_{s.QT\_area}$ and $v_{s.QT\_biomass}$ were lower and more variable than $v_{s.QL}$. These results

reflect the differences in the characteristics of the novel quantitative indices

developed here and the conventional qualitative index. Indeed, $v_{s.QL}$ could not be

applied to 10 plots that had no individual turnover, whereas, in such cases, the $v_{s.QT}$

index is more sensitive than the $v_{s.QL}$ index to temporal changes in the community.

In the driver analysis of the calculated indices, I found that various explanatory

variables affected the results. First, focusing on the growth component specific to the

new quantitative index, both area- and biomass-based d_G indices, which indicate the

contribution of growth, were responsive to the survey interval. This strongly suggests

an accurate evaluation of the contribution of growth was achieved. In addition, the

area-based d_G index tended to decrease toward a larger (i.e., more recent) calendar

year. In addition, overall, the contributions related to recruitment decreased, and those

related to mortality increased (Table 2), which may be indicative of succession (i.e.,

aging forests) and/or climate change.

The $v_{s.QT\_area}$ index showed no clear pattern due to the null model selected, whereas

the $v_{s.QL}$ and $v_{s.QT\_biomass}$ indices varied with elevation and calendar year, respectively.

The v_s family of indices is designed to capture changes that take account of stochastic

changes, such as individual, basal area, or bole volume turnover. This suggests that

along with the explanatory variables, niche processes drive compositional shifts,

which cannot be explained by stochastic processes alone. For example, Nakadai

(2020) showed that $v_{s.QL}$ values vary along climate gradients (i.e., the change ratio of

annual mean temperature) for both deciduous and evergreen broadleaf forests.

Although the indices I have introduced here are mainly applicable to datasets

containing both individual identity and property information, a method that

accommodates various types of quantitative individual information may become

available through future technological developments. Generally, information for

individuals remains limited in community ecology and macroecology, yet in the era

of big data, this is likely to be addressed in the near future. In addition, if the

contributions of demographic factors (i.e., growth, mortality, and recruitment) can be

estimated using hierarchical modelling approaches (e.g., Hill et al., 2004, Kanamori

et al., 2017), these indices could be calculated even in the absence of detailed

individual-tracking information.

Notably, climate change affects various ecological hierarchies, from global

biodiversity to individuals and even intra-individual phenomena (Arneth et al., 2020;

Körner, 2017). Applying the individual-based beta-diversity indices to global

datasets, especially forest inventory data (e.g., ForestGEO; https://forestgeo.si.edu/),

is, therefore, highly promising for assessing the impacts of global climate change.

Indeed, individual-based diversity indices able to account for previously overlooked

elements of biodiversity, such as individual persistence and growth, open the door to

innovative individual-based biodiversity science. This should facilitate a greater

understanding of the effects of climate change across multiple hierarchies of

biological organisation.

Data Availability Statement

Data are available from the package rFIA (Stanke et al., 2020) in R, and the code

used to calculate the indices introduced in the present study is provided in

Supplementary File 1.

References

Akaike, H. (1998). Information theory and an extension of the maximum likelihood

principle. In E. Parzen, K. Tanabe, and G. Kitagawa (Eds.), *Selected Papers of*

Hirotugu Akaike (pp. 199–213). New York, NY. doi:10.1007/978-1-4612-1694-

0_15

Arneth, A., Shin, Y. J., Leadley, P., Rondinini, C., Bukvareva, E., Kolb, M., . . . Saito,

O. (2020). Post-2020 biodiversity targets need to embrace climate change.

Proceedings of the National Academy of Sciences, *117*(49), 30882–30891.

doi:10.1073/pnas.2009584117

Bartoń, K. (2020). MuMIn: Multi-model inference. R package version 1.43.17.

Retrieved from https://CRAN.R-project.org/package=MuMIn

Baselga, A. (2013). Separating the two components of abundance-based dissimilarity:

balanced changes in abundance vs. abundance gradients. *Methods in Ecology*

and Evolution, *4*, 552–557. doi:10.1111/2041-210X.12029

Brienen, R. J. W., Caldwell, L., Duchesne, L., Voelker, S., Barichivich, J., Baliva,

M., . . . Gloor, E. (2020). Forest carbon sink neutralized by pervasive growth-

lifespan trade-offs. *Nature Communications, 11*, 4241.

https://doi.org/10.1038/s41467-020-17966-z

Brienen, R., Phillips, O., Feldpausch, T. Gloor, E., Baker, T., Lloyd, J., . . . Zagt, R. J.

(2015). Long-term decline of the Amazon carbon sink. *Nature*, *519*, 344–348.

https://doi.org/10.1038/nature14283

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004).

Toward a metabolic theory of ecology. *Ecology, 85*(7), 1771–1789.

doi:10.1890/03-9000

Brown, J. H. (1995). *Macroecology*. University of Chicago Press.

Brown, J. H. (1999). Macroecology: Progress and prospect. *Oikos*, 87, 3–14.

doi:10.2307/3546991

Burrill, E. A., DiTommaso A. M., Turner, J. A., Pugh, S. A., Menlove, J.,

Christiansen, G., Perry, C. J., & Conkling, B. L. (2021). *The forest inventory and*

analysis database: Database description and user guide version 9.0.1 for Phase

2. U.S. Department of Agriculture, Forest Service (p. 1026). Retrieved from

http://www.fia.fs.fed.us/library/database-documentation/

Domke, G. M., Oswalt, C. M., Woodall, C. W., & Turner, J. A. (2013). Estimation of

merchantable bole volume and biomass above sawlog top in the National Forest

Inventory of the United States. *Journal of Forestry*, *111*(6), 383–387.

doi:10.5849/jof.13-042

Dornelas, M., Gotelli, N. J., Mcgill, B., Shimadzu, H., Moyes, F., Sievers, C., &

Magurran, A. E. (2014). Assemblage time series reveal biodiversity change but

not systematic loss. *Science*, *344*, 296–299. doi:10.1126/science.1248484

García Molinos, J., Halpern, B. S., Schoeman, D. S., Brown, C. J., Kiessling, W.,

Moore, P. J., . . . Burrows, M. T. (2016). Climate velocity and the future global

redistribution of marine biodiversity. *Nature Climate Change, 6*, 83–88.

doi:10.1038/nclimate2769

Gotelli, N. J., Shimadzu, H., Dornelas, M., McGill. B., Moyes. F., & Magurran, A. E.

(2017). Community-level regulation of temporal trends in biodiversity. *Science*

Advances, 3, e1700315. doi:10.1126/sciadv.1700315

Hill, M. F., Witman, J. D., Caswell, H. (2004). Markov chain analysis of succession

in a rocky subtidal community. *The American Naturalist, 164*, 46–61.

doi:10.1086/422340

Hubbell, S. P. (2001). *The unified neutral theory of biodiversity and biogeography*.

Princeton University Press.

Kanamori Y., Fukaya, K., & Noda, T. (2017). Seasonal changes in community

structure along a vertical gradient: Patterns and processes in rocky intertidal

sessile assemblages. *Population Ecology, 59*, 301–313. doi:10.1007/s10144-

017-0596-z

Körner, C. (2017). A matter of tree longevity. *Science, 355*, 130–131.

doi:10.1126/science.aal2449

Legendre, P., & Legendre, L. (2012). Numerical ecology (3rd ed.). Amsterdam:

Elsevier.

Legendre, P. (2019). A temporal beta-diversity index to identify sites that have

changed in exceptional ways in space–time surveys. *Ecology and Evolution, 9*,

3500–3514. doi:10.1002/ece3.4984

Li, S. P., Cadotte, M. W., Meiners, S. J., Pu, Z., Fukami, T., & Jiang, L. (2016).

Convergence and divergence in a long-term old-field succession: The

importance of spatial scale and species abundance. *Ecology Letters, 19*(9),

1101–1109. doi:10.1111/ele.12647

Magurran, A. E. (2011). Measuring biological diversity in time (and space). In A.

Magurran & B. J. McGill (Eds.), *Biological diversity: Frontiers in measurement*

and assessment (pp. 85–104). Oxford: Oxford University Press.

Magurran, A. E., Dornelas, M., Moyes, F., & Henderson, P. A. (2019). Temporal β

diversity: A macroecological perspective. *Global Ecology and Biogeography,*

28, 1949–1960. doi:10.1111/geb.13026

Magurran, A. E., Dornelas, M., Moyes, F., Gotelli, N. J., & McGill, B. (2015). Rapid

biotic homogenization of marine fish assemblages. *Nature Communication, 6*,

8405. doi:10.1038/ncomms9405

Munné-Bosch, S. (2018). Limits to tree growth and longevity. *Trends in Plant*

Science, 23, 985–993. doi:10.1016/j.tplants.2018.08.001

Nakadai, R. (2020). Degrees of compositional shift in tree communities vary along a

gradient of temperature change rates over one decade: Application of an

individual-based temporal beta-diversity concept. *Ecology and Evolution,*

10(24), 13613–13623. doi:10.1002/ece3.6579

Nakadai, R. (2021). Individual-based multiple-unit dissimilarity: novel indices and

null model for assessing temporal variability in community composition.

Oecologia, 197(2), 353–364. doi:10.1007/s00442-021-05025-3

Odum, E. P. (1950). Bird populations of the Highlands (North Carolina) Plateau in

relation to plant succession and avian invasion. *Ecology, 31*, 587–605.

doi:10.2307/1931577

Oksanen, J., Guillaume, B. F., Michael, F., Roeland, K., Pierre, L., Dan, M., . . .

Helene, W. (2020). Vegan: Community ecology package. R package version 2.5-

7. Retrieved from https://CRAN.R-project.org/package=vegan

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R Core Team. (2022). Nlme: Linear

and nonlinear mixed effects models. R package version 3.1-155. Retrieved from

https://CRAN.R-project.org/package=nlme

R Development Core Team. (2021). R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria. (ver.

4.1.0). Retrieved from http://www.R-project.org/

Searle, E. B., & Chen, H. Y. (2018). Temporal declines in tree longevity associated

with faster lifetime growth rates in boreal forests. *Environmental Research*

Letters, 13(12), 125003. doi:10.1088/1748-9326/aaea9e

Shimadzu, H. et al. (2015). Measuring temporal turnover in ecological communities.

Methods in Ecology and Evolution, 6, 1384–1394. doi:10.1111/2041-

210X.12438

Sørensen, T. A. (1948). A method of establishing groups of equal amplitude in plant

sociology based on similarity of species content and its application to analyses

of the vegetation on Danish commons. *Biologiske Skrifter, 5(4)*, 1–34.

Stanke, H., Finley, A. O., Weed, A. S., Walters, B. F., & Domke, G. M. (2020). rFIA:

An R package for estimation of forest attributes with the US Forest Inventory

and Analysis database. *Environmental Modelling & Software, 127*, 104664.

doi:10.1016/j.envsoft.2020.104664

Tatsumi, S., Iritani, R., & Cadotte, M. W. (2021). Temporal changes in spatial

variation: Partitioning the extinction and colonisation components of beta

diversity. *Ecology Letters, 24*(5), 1063–1072. doi:10.1111/ele.13720

Vellend, M., Baeten, L., Myers-Smith I. H., Elmendorf, S. C., Beauséjour, R., Brown

C. D., . . . Wipf, S. (2013). Global meta-analysis reveals no net change in local-

scale plant biodiversity over time. *PNAS, 110*, 19456–19459.

doi:10.1073/pnas.1312779110

Legends

Figure 1 Schematic representation of individual-based temporal beta-diversity for

both qualitative (a) and quantitative (b) components. T1 and T2 refer to the

communities at times 1 and 2, respectively. Both communities T1 and T2 comprise

four species (sp. 1, 2, 3, and 4) and all individuals are identified separately (Z_o) in

both cases. The arrows indicate the temporal trajectory for each individual. In the

qualitative component (a), m refers to individuals in existence at T1 but dead before

T2, thereby separately identifying the deaths of individuals during the period between

T1 and T2; r identifies individuals that were not alive at T1 but found at T2, thereby

identifying the recruitment of individuals during the period between T1 and T2; p

refers to individuals that persisted from T1 to T2, thus both T1 and T2 = 1. In the

qualitative component (b), M refers to the total amount of measures of individual

occurring at T1 that die before T2, thereby representing the total measures of the

mortal individuals during the time between T1 and T2; R identifies the total measures

of individuals that were not alive at T1 but were subsequently alive and counted at

T2, thereby representing the recruit individuals between T1 and T2. Persistent

individuals are qualitatively identical, having alive (or present) status between T1 and

T2, but rarely keep the same status with respect to size and weight. Therefore, change

in individual properties occurs between T1 and T2. The quantitative indices consider

the contribution of changes of individuals' properties, mainly targeting individual

growth. Specifically, for persistent individuals, the original and altered total measures

are partitioned into persistent (P) and growth (G) components, respectively.

Figure 2

Density plots representing the distribution of individual-based quantitative beta-

diversity indices across 126 survey intervals for 86 plots (data from Stanke et al.

2020). In all four plots, the values of the indices are limited to the range 0–0.15 to

enable comparison (see Fig. S2 for a plot of the entire range). The plots (a) and (c)

are based on basal area, and plots (b) and (d) are based on bole biomass.

Figure 1

Figure 2

Table 1. Summary of indices, formulae, and references citing indices developed in previous studies and proposed in this study. y_{ij} is

abundance excluding individuals (i.e., measures or weights) of species i at time Tj at a site, and y_{ik} is abundance excluding individuals

of species i at time Tk at a site.

Index	Formula	References
d_{BC}	$$\frac{B+C}{2A+B+C} = \frac{\sum_i[x_{ij}-\min(x_{ij},x_{ik})]+\sum_i[x_{ik}-\min(x_{ij},x_{ik})]}{2[\sum_i\min(x_{ij},x_{ik})]+\sum_i[x_{ij}-\min(x_{ij},x_{ik})]+\sum_i[x_{ik}-\min(x_{ij},x_{ik})]}$$	Odum (1950)
d_{mr}	$$\frac{m+r}{2p+m+r} = \frac{\sum_o[z_{oj}-\min(z_{oj},z_{ok})]+\sum_o[z_{ok}-\min(z_{oj},z_{ok})]}{2[\sum_o\min(z_{oj},z_{ok})]+\sum_o[z_{oj}-\min(z_{oj},z_{ok})]+\sum_o[z_{ok}-\min(z_{oj},z_{ok})]}$$	Nakadai (2020)
$v_{s.QL}$	$$\frac{B+C}{2e+B+C}=\frac{B+C}{m+r} = \frac{\sum_i[x_{ij}-\min(x_{ij},x_{ik})]+\sum_i[x_{ik}-\min(x_{ij},x_{ik})]}{2[\sum_i\min(x_{ij},x_{ik})-\sum_o\min(z_{oj},z_{ok})]+\sum_i[x_{ij}-\min(x_{ij},x_{ik})]+\sum_i[x_{ik}-\min(x_{ij},x_{ik})]}$$	Nakadai (2020)
$d_{BC'}$	$$\frac{B'+C'}{2A'+B'+C'} = \frac{\sum_i[y_{ij}-\min(y_{ij},y_{ik})]+\sum_i[y_{ik}-\min(y_{ij},y_{ik})]}{2[\sum_i\min(y_{ij},y_{ik})]+\sum_i[y_{ij}-\min(y_{ij},y_{ik})]+\sum_i[y_{ik}-\min(y_{ij},y_{ik})]}$$	Odum (1950)

$$d_{MRG} = \frac{\frac{M+R+G}{2P+M+R+G}}{\sum_o\left[w_{oj}-\min(w_{oj},w_{ok})\times[z_{oj}-\min(z_{oj},z_{ok})]\right]+\sum_o\left[[w_{ok}-\min(w_{oj},w_{ok})]\times[z_{ok}-\min(z_{oj},z_{ok})]\right]+\sum_o\left[[\max(w_{oj},w_{ok})-\min(w_{oj},w_{ok})]\times\min(z_{oj},z_{ok})\right]}{2[\sum_o\min(w_{oj},w_{ok})+\sum_o\left[w_{oj}-\min(w_{oj},w_{ok})\times[z_{oj}-\min(z_{oj},z_{ok})]\right]+\sum_o\left[[w_{ok}-\min(w_{oj},w_{ok})]\times[z_{ok}-\min(z_{oj},z_{ok})]\right]+\sum_o\left[[\max(w_{oj},w_{ok})-\min(w_{oj},w_{ok})]\times\min(z_{oj},z_{ok})\right]}$$

This study

$$v_{s.QT} \quad \frac{B'+C'}{2E+B'+C'} = \frac{\sum_i[y_{ij}-\min(y_{ij},y_{ik})]+\sum_i[y_{ik}-\min(y_{ij},y_{ik})]}{2[\sum_i\min(y_{ij},y_{ik})-\sum_o\min(w_{oj},w_{ok})]+\sum_i[y_{ij}-\min(y_{ij},y_{ik})]+\sum_i[y_{ik}-\min(y_{ij},y_{ik})]}$$

This study

681   Table 2. Results of a multi-regression model based on the best models selected using an Akaike information criterion (AIC)

682   comparison for individual-based qualitative and quantitative beta-diversity indices. β and *parR2* indicate the regression coefficients

683   and partial adjusted R-squared values of each explainatory variable, respectively. *Adj R2* is the adjusted R-squared value for the

684   complete model. Note: Significance: .<0.1, *<0.05, **<0.01, ***<0.001. When a null model was chosen, † is added into the column

685   of p-value for whole model.

	Latitude				Elevation				Calendar year				Survey interval				Adj R2	P-values	
	β	parR2	P-values		β	parR2	P-values		β	parR2	P-values		β	parR2	P-values				
(a) Qualitative-family indices																			
d_{BC}	–	–	–		–	–	–		–	–	–		–	–	–			†	
d_{mr}	–	–	–		–	–	–		–	–	–		0.0186	0.0089	0.1473		0.0089	0.1473	
d_m	-0.0807	0.0333	0.0233	*	–	–	–		0.0098	0.0953	0.0003	***	–	–	–		0.1211	0.0001	***
d_r	–	–	–		–	–	–		-0.0129	0.0513	0.0064	**	0.0277	0.0146	0.0944	.	0.0554	0.0111	*
$v_{s.QL}$	–	–	–		0.0002	0.0283	0.0393	*	–	–	–		–	–	–		0.0283	0.0393	*
(b) Qantitative-family indices																			
(i) Basal area																			
$d_{BC'}$	–	–	–		–	–	–		–	–	–		–	–	–			†	
d_{MRG}	-0.0705	0.0095	0.1413		–	–	–		–	–	–		0.0231	0.0192	0.0663	.	0.0269	0.0695	.
d_M	-0.0768	0.0253	0.0421	*	–	–	–		0.0094	0.0770	0.0010	**	–	–	–		0.0972	0.0007	***
d_R	–	–	–		–	–	–		-0.0044	0.0259	0.0398	*	–	–	–		0.0259	0.0398	*
d_G	–	–	–		–	–	–		-0.0028	0.0236	0.0477	*	0.0122	0.0381	0.0165	*	0.0305	0.0553	.
$v_{s.QT}$	–	–	–		–	–	–		–	–	–		–	–	–			†	
(ii) Bole biomass																			
$d_{BC'}$	–	–	–		–	–	–		–	–	–		0.0205	0.0114	0.1210		0.0114	0.1210	
d_{MRG}	-0.0748	0.0094	0.1429		–	–	–		–	–	–		0.0278	0.0264	0.0387	*	0.0336	0.0454	*
d_M	-0.0766	0.0209	0.0586	.	–	–	–		0.0092	0.0641	0.0026	***	–	–	–		0.0812	0.0020	**
d_R	–	–	–		–	–	–		-0.0039	0.0187	0.0684	.	–	–	–		0.0187	0.0685	.
d_G	–	–	–		–	–	–		–	–	–		0.0089	0.0227	0.0503	.	0.0227	0.0503	.
$v_{s.QT}$	–	–	–		–	–	–		-0.0301	0.0272	0.0366	*	0.0717	0.0082	0.1569		0.0232	0.0877	.

Appendices

Table S1. Comparison using a linear and linear mixed model for each index. The best models chosen from the comparison are shown in Table 2

Figure S1. Map of geographic locations of plots included in the subset of the Forest Inventory and Analysis (FIA) database for the state of Rhode Island in USA (Stanke et al., 2020). Sites with darker colour points have multiple survey intervals.

Figure S2. Density plots representing the distribution of individual-based quantitative beta-diversity indices across 126 survey intervals of 86 plots (data from Stanke et al., 2020). In the right four plots (b, d, f, and h), the index values are limited to the range 0–0.15 to enable comparison (also see Fig. 2). In the left four plots, the entire value is shown. Plots (a), (b), (e), and (f) are based on basal area, and plots (c), (d), (g), and (h) are based on bole biomass.

Figure S3. Correlations among the calculated indices.

Supplementary text 1. Explanation of individual-based temporal beta diversity employing the Ružička dissimilarity index.

Supplementary file 1. R software script for individual-based temporal beta diversity calculations using sample data.