Development of novel temporal beta-diversity indices for assessing community

compositional shifts accounting for changes in the properties of individuals

Ryosuke Nakadai

ORCID: 0000-0002-9512-8511

Biodiversity Division, National Institute for Environmental Studies, Onogawa 16-2, Tsukuba,

Ibaraki 305-8506, Japan

E-mail address: r.nakadai66@gmai.com, nakadai.ryosuke@nies.go.jp

[1]

---

[1] AIC, Akaike information criterion; DBH, Diameter at breast height; FIA, Forest Inventory and Analysis

Abstract

Revealing the patterns of temporal biodiversity changes and deciphering the connection

between individual life histories and large-scale ecological patterns is essential for

improving the mechanistic understanding of macroecology. However, this is challenging

because the relationship between individual life history and biodiversity remains unclear.

In the present study, in order to link the individual life history and community-level

phenomena, I developed novel indices that allow the evaluation of community

compositional shifts over time by explicitly considering the contributions of the life

histories of individuals (i.e. growth, mortality, and recruitment) in a community. These

novel indices are quantitative extensions of the individual-based temporal beta-diversity

indices which can include information on individual sizes. The indices were applied to a

subset of data from the US Forest Inventory and Analysis database for the state of Rhode

Island, USA, to identify changes in the contribution of individual life histories to

biodiversity change. The results of this study represent methodological progress in

community ecology and macroecology, as well as a conceptual advancement in bridging

studies on biodiversity with those on individual life history and physiology. The

individual-based diversity indices developed here pave the way for individual-based

biodiversity science, which may facilitate the understanding of the effects of climate

change across different hierarchies of biological organisation.

Keywords: climate change, size, growth, forest dynamics, life history

1. Introduction

An individual cannot exist in two places simultaneously, but it can persist in the same habitat over time (Magurran et al., 2019). This feature of individuals is a major difference between time and space in community ecology. In a temporal context, individual turnover between different species is directly linked to compositional shifts in a community (Magurran et al., 2019, Nakadai 2020). Therefore, individual persistence and turnover are important factors for measuring temporal compositional shifts in biodiversity, termed temporal beta diversity (Gotelli et al., 2017; Magurran, 2011, Tatsumi et al., 2021). For example, individuals of long-lived species (e.g. trees and vertebrates) can persist in the long term, whereas individuals of short-lived species are replaced by conspecifics or individuals of other species (Nakadai, 2020, 2021). Nakadai (2020) proposed the concept of 'individual-based beta diversity' to bridge the gap between biodiversity and individual information and provided novel indices to evaluate individual turnover and compositional shifts based on comparisons between two periods at a given site. Specifically, Nakadai (2020) developed an individual-based temporal beta-diversity approach for the Bray–Curtis dissimilarity index (i.e. an abundance-based temporal beta-diversity index) and accounted for the contribution of both individual persistence and turnover to the compositional shifts in biodiversity over time (Nakadai, 2020, 2021).

Climate change is known to directly affect not only biodiversity, but also the individual life histories of organisms (Brienen et al., 2020; Searle & Chen, 2018). Recent studies have shown that climate warming will accelerate the growth and shorten the life spans—which are determined by physiological processes—of individual trees (Körner, 2017; Munné-Bosch, 2018; Searle & Chen, 2018) and increase the amount of carbon stocks in terrestrial ecosystems (Brienen et al., 2015). Such changes in individual life histories potentially affect the compositional shifts of local communities and, consequently, global biodiversity. While understanding the connection between individual physiology and large-scale ecological patterns is essential for a mechanistic understanding of macroecology (Brown, 1995, 1999; Brown et al. 2004; e.g. metabolic theory), this is challenging because the connection between individual life history and biodiversity remains unknown.

Nevertheless, to comprehensively understand the effects of climate change, it is essential to

simultaneously capture changes in individual properties, individual turnover, and

compositional shifts in biodiversity.

Despite these requirements, previous approaches to individual-based temporal beta

diversity (Nakadai, 2020, 2021) have only considered the presence or absence of each

individual (e.g. qualitative properties such as 'dead' or 'alive'), and the effects of property

changes in each individual (e.g. individual growth) on biodiversity over time have been

neglected. Therefore, the aim of this study was to develop new individual-based temporal

beta-diversity indices that can assess compositional shifts over time by explicitly including

property changes in each individual in a community. Hereafter, I refer to the indices

developed by Nakadai (2020) and the novel indices introduced in the present study

individual-based 'qualitative' ('QL') indices and individual-based 'quantitative' ('QT')

indices, respectively. By introducing quantitative individual information (e.g. size and

weight), these novel indices can evaluate the contribution of individual growth and the

differences between large and small individuals. After briefly reviewing the history of

pairwise beta-diversity indices, I extended individual-based QL temporal beta-diversity

indices to QT indices. I then applied these indices to subset data from the US Forest

Inventory and Analysis (FIA) database for the state of Rhode Island in the USA (Stanke et

al., 2020, Burrill et al., 2021) as a case study to demonstrate their application.

2. Materials and methods

2.1. Conventional incidence- and abundance-based indices and recently developed

individual-based qualitative (QL) temporal beta-diversity indices

Incidence-based and abundance-based beta-diversity indices have been previously

proposed (Baselga, 2013). For example, the widely used Bray–Curtis dissimilarity index

(Odum, 1950) is an abundance-based extension of the Sørensen index (Baselga, 2013;

Legendre & Legendre, 2012; Sørensen, 1948). Incidence-based methods target the number

of species, whereas abundance-based methods mainly target the number of individuals and,

sometimes, area, biomass, and weight (e.g. Li et al., 2016). Conventional beta-diversity

indices are based on intersection (A, a) and relative complement (B, b, C, c) components.

For context, the Sørensen dissimilarity index (d_{sor}; Sørensen, 1948) is formulated as

follows:

$$d_{sor} = \frac{b+c}{2a+b+c}$$ Eqn. 1

where a is the number of species common to both temporal units; b is the number of

species that occur in the first but not the second temporal unit; and c is the number of

species that occur in the second but not the first temporal unit (Legendre, 2019).

Analogously, the Bray–Curtis dissimilarity index (d_{BC}; Odum, 1950) is formulated as

follows:

$$d_{BC} = \frac{B+C}{2A+B+C}$$ Eqn. 2

As a premise for explanation, assume that there are two sets of community data along the

time axis—communities at time j and time k (abbreviated as Tj and Tk: $j < k$), respectively.

The abundance-based components are as follows:

$A = \sum_i \min(x_{ij}, x_{ik})$ Eqn. 3a

$B = \sum_i [x_{ij} - \min(x_{ij}, x_{ik})]$ Eqn. 3b

$C = \sum_i [x_{ik} - \min(x_{ij}, x_{ik})]$ Eqn. 3c

where x_{ij} is the abundance of species i at temporal unit j, and x_{ik} is the abundance of species

i at temporal unit k. Therefore, A is the total number of individuals of each species shared

by units j and k, and B and C are the total numbers of individuals of each species unique to

units j and k, respectively (Odum, 1950, Legendre, 2019).

Nakadai (2020) introduced the categorisation of the community dynamics component

into persistence (p), mortality (m), and recruitment (r) in individual-based temporal beta

diversity. m refers to the number of individuals occurring at Tj that die before Tk,

representing the total number of individuals that died during the time between Tj and Tk. r

identifies the number of individuals that were not yet alive at Tj (or were not yet in the size

class counted, e.g. diameter at breast height (DBH) $\geq$ 1 cm), but that were alive and

counted at Tk, representing the total number of individuals recruited between Tj and Tk.

Component p refers to the persistence of individuals from Tj to Tk. Note that, unlike

previous studies (Nakadai, 2020, 2021; Fig. 1a), I use lower-case letters for the

components of individual-based qualitative indices (i.e. p, m, and r) to distinguish the

components of the individual quantitative indices described later. Thus, the abundances are

calculated as follows:

$\quad p = \sum_o \min(z_{oj}, z_{ok})$ $\qquad$ Eqn. 4a

$\quad m = \sum_o [z_{oj} - \min(z_{oj}, z_{ok})]$ Eqn. 4b

$\quad r = \sum_o [z_{ok} - \min(z_{oj}, z_{ok})]$ Eqn. 4c

where z_{oj} is individual o at time Tj, and z_{ok} is individual o at time Tk. Both z_{oj} and z_{ok} have

values of one or zero (presence and absence, respectively). Therefore, p is the total number

of individuals present at both Tj and Tk, and m and r are the numbers of individuals unique

to Tj and Tk, respectively (Nakadai, 2020, 2021; Fig. 1a).

Fig. 1. Schematic representation of individual-based temporal beta diversity for both qualitative (a) and quantitative (b) components. T1 and T2 refer to the communities at times 1 and 2, respectively. Both communities T1 and T2 comprise four species (sp. 1, 2, 3, and 4), and all individuals are identified separately (Z_o) in both cases. The arrows indicate the temporal trajectory for each individual. In the qualitative components (a), m refers to the number of individuals in existence at T1 but dead before T2, thereby separately identifying the deaths of individuals during the period between T1 and T2; r refers to the number of individuals that were not yet alive at T1 but were present at T2, thereby identifying the recruitment of individuals during the period between T1 and T2; p refers to the number of individuals that persisted from T1 to T2, thus both T1 and T2 = 1; e refers to the number of individuals that did not contribute to the apparent compositional shift among mortality or recruitment of individuals, which is calculated as $m–B$ or $r–C$. In the qualitative components (b), M refers to the total measures of individuals occurring at T1 that died before T2, thereby representing the total measures of the mortal individuals during the time between T1 and T2; R refers to the total measures of individuals that were not yet alive at T1 but were alive and counted at T2, thereby representing the individuals recruited between T1 and T2. Persistent individuals are qualitatively identical,

152 having alive (or present) status between T1 and T2, but rarely maintain the same status with respect to size

153 and weight (e.g., no size change). Therefore, changes in individual sizes occur between T1 and T2. The

154 quantitative indices consider the contribution of changes in individuals' sizes, mainly targeting individual

155 growth. Specifically, for persistent individuals, the original and altered total measures are partitioned into

156 persistent (P) and growth (G) components, respectively. E refers to the total basal area that did not contribute

157 to the apparent compositional shift among mortality, recruitment of individuals, or growth of individuals,

158 which is calculated as $M+G_{de}-B'$ or $R+G_{in}-C'$.

160 The individual-based temporal beta-diversity index for Bray–Curtis dissimilarity (d_{mr})

161 can be formulated (Nakadai 2020) as follows:

163 $d_{mr} = \dfrac{m+r}{2p+m+r}$   Eqn. 5

165 In some cases, there is a turnover of different individuals belonging to the same species,

166 even when the community composition is stable over time. Therefore, the components that

167 change over time (m and r) are reorganised into two further components, namely

168 compositional equilibrium (e) and shift (B and C) (Nakadai, 2020, 2021). Here, I define

169 compositional equilibrium as a turnover component without apparent compositional shift

170 in two time steps, thereby contributing to the dynamic equilibrium. Component m can be

171 partitioned into two additive components—the lost individuals that contribute to

172 compositional shifts in a community (B) and the lost individuals replaced by conspecifics

173 that contribute to an equilibrium in community composition (e_{loss}) (Fig. 1a). Similarly,

174 component r can be partitioned into two additive components—gained individuals that

175 contribute to compositional shifts in a community (C) and gained individuals that replace

176 conspecifics, thereby contributing to an equilibrium in community composition (e_{gain}) (Fig.

177 1a). By definition, the numbers of e_{loss} and e_{gain} individuals are identical; thus, I replaced

178 both e_{loss} and e_{gain} with the coefficient e (Nakadai, 2020, 2021). The persistence component

179 (p) is the shared (apparent non-changed) component (A) minus the equilibrium (changed,

180 but apparent non-changed) component (e) of community composition. The above

descriptions are summarised below:

$e_{loss} = e_{gain} = e$ Eqn. 6

$m = B + e$ Eqn. 7

$r = C + e$ Eqn. 8

$p = A - e$ Eqn. 9

Furthermore, components m and r can be analysed in the same manner as the

community data (Nakadai, 2020). Therefore, the Bray–Curtis dissimilarity between

communities can be calculated based on m and r (Fig. S1d), as follows:

$v_{s.QL} = \frac{B+C}{2e+B+C} = \frac{B+C}{m+r}$ Eqn. 10

where $v_{s.QL}$ indicates the speed of compositional shifts in a community relative to the total

speed of individual turnover associated with m and r. In other words, $v_{s.QL}$ indicates total

changes in community composition divided by total changes in the number of individuals. It

is possible to calculate $v_{s.QL}$ as d_{BC} divided by d_{mr} as:

$d_{BC} \div d_{mr} = \frac{B+C}{2A+B+C} \div \frac{m+r}{2p+m+r} = \frac{B+C}{2(p+e)+B+C} \div \frac{m+r}{2p+m+r} = \frac{B+C}{2p+m+r} \div \frac{m+r}{2p+m+r} = \frac{B+C}{2e+B+C} =$

$v_{s.QL}$ Eqn. 11

2.2. Novel individual-based "quantitative (QT)" beta-diversity indices

I developed individual-based QT components and novel indices to analyse community

datasets over time using individual property information (e.g. size and weight). The

relationship between the QL and QT indices is similar to that between the incident-based

and abundance-based indices. In other words, individual-based QT components are

calculated as individual-based QL weighted by individual sizes. As previously noted, the

abundance targeted by abundance-based indicators is not necessarily the number of

individuals but can also include area and volume. Abundance measures excluding the

number of individuals for community datasets have been used in cases of large size

variation among individuals within a community or where it is difficult to recognise proper

individuals (Li et al., 2016). Hereafter, the Bray–Curtis indices targeting measures

excluding the number of individuals (e.g. sizes and weights) are described as $d_{BC'}$, and

their components are described with apostrophes (A', B', and C'; detailed equations of the

component see Supplementary Text 1) to distinguish them from the components in the

unweighted Bray–Curtis index, as follows:

$$d_{BC'} = \frac{B' + C'}{2A' + B' + C'}$$ Eqn. 12

This weighting indicates that abundance-based (number of individual-based) diversity

indices are weighted by total measures of individuals for each species, thus calculating

diversity indices by implicitly considering the variation in the measures of each individual.

As individuality is rarely clearly identifiable in weighted indices, this relationship is not

well recognised.

In addition to individual-based QL indices, individual-based QT indices comprise the

components of mortality and recruitment (M and R, respectively). In the QT indices, M

refers to the total measures of individuals occurring at Tj that died before Tk, representing

the total measures of the individuals that died between Tj and Tk (Fig. 1b). R identifies the

total measures of individuals that were not yet alive at Tj (or were not yet in the size class

counted, e.g. DBH $\geq$ 1 cm) but were alive and counted at Tk, representing the total

measures of the individuals recruited between Tj and Tk (Fig. 1b).

From this point on, the major differences between the QL and QT indices are as follows.

The persistent individuals are qualitatively identical, maintaining their alive (or present)

status between Tj and Tk, but rarely maintain the same status in terms of size and weight;

thus, there is a size change in an individual between Tj and Tk. For example, in tree

communities, a newly recruited small tree can be a large tree after persisting for half a

century. In the p component of QL indices, this implies that an individual is persistent;

however, evaluating changes in its features (e.g. basal area, biomass, and/or weight) is not

feasible. The indices introduced here overcome this problem by considering the

contribution of changes in individual sizes, mainly targeting individual growth.

Specifically, in the case of persistent individuals, the original and altered total measures are

partitioned into persistent (*P*) and growth (*G*) components, respectively. Thus, the

respective total measures of the components are calculated as follows:

$M = \sum_o \left[\left[w_{oj} - \min(w_{oj}, w_{ok}) \right] \times \left[z_{oj} - \min(z_{oj}, z_{ok}) \right] \right]$ Eqn. 13a

$R = \sum_o \left[\left[w_{ok} - \min(w_{oj}, w_{ok}) \right] \times \left[z_{ok} - \min(z_{oj}, z_{ok}) \right] \right]$ Eqn. 13b

$P = \sum_o \min(w_{oj}, w_{ok})$ Eqn. 13c

$G = \sum_o \left[\left[\max(w_{oj}, w_{ok}) - \min(w_{oj}, w_{ok}) \right] \times \min(z_{oj}, z_{ok}) \right]$ Eqn. 13d,

where w_{oj} is the measure of individual *o* at time T*j*, and w_{ok} is the measure of individual *o*

at time T*k*. In general, basal area and biomass of each individual stem can be applicable as

a measure, i.e., w_{oj} and w_{ok}. *M* and *R* are the total measures of demised and recruited

individuals, respectively. The latter parts of Equation 13a, b, and d identify each individual

based on QL information which is equal to Equation 4a, b, and c. The persistent and

growth components represent the unchanged and changed components of measures within

an individual between T*j* and T*k*, respectively. Thus, *P* and *G* represent the total values of

these components within a community. Equation 13d also considers decreases in the

measures of each individual, because decreases in individual size may occasionally be

described owing to errors, vertebrate foraging, and other unknown reasons in empirical

forest datasets. Therefore, the calculation of the *G* component can be partitioned into

increasing and decreasing components, as follows:

$G = \sum_o \left[\left[\max(w_{oj}, w_{ok}) - \min(w_{oj}, w_{ok}) \right] \times \min(z_{oj}, z_{ok}) \right] = G_{in} + G_{de}$

Eqn. 14,

where G_{in} and G_{de} are the total increased and decreased components of the targeted

measures, respectively. Specifically, the formulas for G_{in} and G_{de} are as follows:

$G_{in} = \sum_o \left[\left[w_{ok} - \min(w_{oj}, w_{ok}) \right] \times \min(z_{oj}, z_{ok}) \times \frac{1}{2} \left[1 + \text{sign}(w_{ok} - w_{oj}) \right] \right]$ Eqn.

15a

$G_{de} = \sum_o \left[\left[w_{oj} - \min(w_{oj}, w_{ok}) \right] \times \min(z_{oj}, z_{ok}) \times \frac{1}{2} \left[1 - \text{sign}(w_{ok} - w_{oj}) \right] \right]$ Eqn.

15b,

where $\frac{1}{2}\left[1 + \text{sign}(w_{ok} - w_{oj})\right]$ and $\frac{1}{2}\left[1 - \text{sign}(w_{ok} - w_{oj})\right]$ distinguish the large and

small relationships (i.e. increase changes or decreased changes) between w_{ok} and w_{oj} by

returning 1 only when $\left[w_{ok} - \min(w_{oj}, w_{ok})\right]$ and $\left[w_{oj} - \min(w_{oj}, w_{ok})\right]$ are positive

values, otherwise returning 0. Additionally, $\min(z_{oj}, z_{ok})$ identifies persistent individuals

between time k and j, similar to that in Equation 4a.

This formulation of the individual-based QT temporal beta-diversity index for Bray–

Curtis dissimilarity (d_{MRG}) can be expressed as follows:

$d_{MRG} = \frac{M+R+G}{2P+M+R+G}$ Eqn. 16

Further, this individual-based QT temporal beta diversity can be partitioned into the

relative contributions of mortality (d_M), recruitment (d_R), and growth (d_G) to the target

weights or measurements, as follows:

$d_{MRG} = \frac{M+R+G}{2P+M+R+G}$

$= \frac{M}{2P+M+R+G} + \frac{R}{2P+M+R+G} + \frac{G}{2P+M+R+G} = d_M + d_R + d_G$ Eqn. 17

In addition, as shown in Equations 15a and 15b, the G component can be partitioned into

G_{de} and $G_{in,}$ which are the total decrease and increase, respectively, in the targeted

measures. Equation 17 can be further partitioned into two components, as follows:

$d_{MRG} = \frac{M+R+G}{2P+M+R+G}$

$= \frac{M+G_{de}}{2P+M+R+G} + \frac{R+G_{in}}{2P+M+R+G} = d_{de} + d_{in}$ Eqn. 18

where d_{de} and d_{in} indicate the contributions of the decrease and increase, respectively, to

the total changes in the targeted measures in Tj and Tk. In addition to the QL indices

(Nakadai, 2020, 2021), the components that change over time (M, R, and G) can be

reorganised into two additive components—compositional equilibrium (E) and shift (B'

and C'). Component $M + G_{de}$ can be reorganised as the total values of lost measures that

contribute to compositional shifts in a community (B') and the total values of lost measures

replaced by conspecifics, contributing to the equilibrium in community composition (E_{loss})

(Fig. 1b). Similarly, component $R + G_{in}$ can be reorganised as the total measures that

contribute to compositional shifts in a community (C') and the total values of gained

measures replacing conspecifics, contributing to the equilibrium in community

composition (E_{gain}) (Fig. 1b). By definition, the values of E_{loss} and E_{gain} are identical; hence,

I replaced both E_{loss} and E_{gain} with the coefficient E. In addition, the persistence component

(P) is the shared (apparent non-changed) component (A') minus the community

composition equilibrium (changed, but apparent non-changed) component (E). The above

descriptions can be summarised as follows:

$\quad E_{loss} = E_{gain} = E$ Eqn. 19

$\quad M + G_{de} = B + E$ Eqn. 20

$\quad R + G_{in} = C + E$ Eqn. 21

$\quad P = A' - E$ Eqn. 22

The formulation of the individual-based QT beta-diversity index for Bray–Curtis (d_{MRG})

dissimilarity can be partitioned into two components—contribution to compositional shifts

(d_S) and equilibrium (d_E)—as follows:

$d_{MRG} = \dfrac{M+R+G}{2P+M+R+G} = \dfrac{M+G_{de}}{2P+M+R+G} + \dfrac{R+G_{in}}{2P+M+R+G}$

$= \dfrac{E+B}{2P+M+R+G} + \dfrac{E+C}{2P+M+R+G} = \dfrac{2E}{2P+M+R+G} + \dfrac{B+C}{2P+M+R+G} = d_S + d_E$ $\qquad$ Eqn. 23

Furthermore, components M, R, and G (i.e. $M + G_{de}$ and $R + G_{in}$) can be analysed in the

same manner as the community data. Therefore, the Bray–Curtis dissimilarity between

communities (i.e. the total values of lost and gained measures) can be calculated based on

M, R, and G (Fig. S1h), as follows:

$$v_{s.QT} = \frac{B'+C'}{2E+B'+C'} = \frac{B'+C'}{M+R+G}$$ Eqn. 24

where $v_{s.QT}$ indicates the relative speed of compositional shifts in a community against the

total turnover of targeted measures associated with M, R, and G. In other words, $v_{s.QT}$

indicates total changes in community composition divided by the total changes in targeted

measures. If $v_{s.QT}$ is 0, the turnover of the targeted measures does not contribute to

compositional shifts in a community, whereas if $v_{s.QT}$ is 1, all turnover of the targeted

measures contributes to the compositional shifts. $v_{s.QT}$ can be calculated as $d_{BC'}$ divided by

d_{MRG}, as follows:

$$d_{BC'} \div d_{MRG} = \frac{B'+C'}{2A'+B'+C'} \div \frac{M+R+G}{2P+M+R+G} = \frac{B'+C'}{2(P+E)+B'+C'} \div \frac{M+R+G}{2P+M+R+G} = \frac{B'+C'}{2P+M+R+G} \div$$

$$\frac{M+R+G}{2P+M+R+G} = \frac{B'+C'}{M+R+G} = v_{s.QT}$$ Eqn. 25

To facilitate a better understanding of the indices for potential users, I have summarised the

relationships among four types of index families based on their sizes in Table S1.

Here, I focus only on an extension of Bray–Curtis dissimilarity. In Supplementary Text 2,

I summarise individual-based beta diversity using the Ružička dissimilarity index (Ružička,

1958), which was identified by Podani et al. (2013) and Legendre (2014) as an appropriate

index for beta-diversity studies. The Ružička dissimilarity index is an abundance-based

extension of the Jaccard dissimilarity index (Jaccard, 1901).

It is noteworthy that length of the survey interval between two time periods (i.e. Tj and

Tk in the above equations) can affect the calculated value, as well as conventional beta

diversity indices (Hatosy et al., 2013), because longer intervals mean that individuals that

complete a life cycle between the two periods could be lost from the investigation. In cases

where a dataset with varying survey interval lengths is analysed, the length of the interval

should be a sampling design-related factor.

2.3. Hypothetical impacts of climate warming on compositional shifts in forest communities

The hypothetical impacts of climate warming on compositional shifts, especially in forest communities, are summarised in schematic diagrams for three simple scenarios of forest compositional dynamics in Fig. 2 ((a) no climate warming, (b) compositional dynamic changes due to warming, and (c) accelerated growth and shortened life span due to warming). In scenario (b), it is assumed that the species composition of recruited individuals (i.e., changes in the species pool) become different from species in a community during the previous time step. Hence, the proportions of compositional shift (B and C in $v_{s.QL}$ and B' and C' in $v_{s.QT}$) increase during transitional periods. However, the total growth of individuals that have persisted since before the climate warming acts as a factor decreasing the apparent compositional shift in the size-weighted indices (E in $v_{s.QT}$ increase). Scenarios (b) and (c) are represented separately to aid clarity, but they can simultaneously occur and facilitate compositional shifts in forests. In scenario (c), it is assumed that the speed of growth is twice that of scenarios (a) and (b); therefore, life spans become shorter and, thus, the turnover of both individuals and target measures increases. In contrast to scenario (b), I do not assume a change in the species pool in scenario (c), and thus, the values of both $v_{s.QL}$ and $v_{s.QT}$ do not change, but rather, the total turnover (d_{mr} and d_{MRG}) increases. Mathematically, if $v_{s.QL}$ and $v_{s.QT}$ are constant, the total turnover (d_{mr} and d_{MRG}) is directly linked to the apparent compositional shift (d_{BC} and $d_{BC'}$). Individual growth largely contributes to the fluctuation in compositional shift under accelerated growth as it is more variable than individual turnover, even if life spans become shorter. Although individuals which died due to causes other than having reached the natural life span (random death and death by environmental filtering in nature) are not considered because of the complexity of visualisation in Fig. 2, in scenarios (b) and (c), such deaths may facilitate compositional shifts, specifically by increasing $v_{s.QL}$ and $v_{s.QT}$ or d_{mr} and d_{MRG}.

ig. 2. Schematic diagrams of hypothetical impacts of climate warming on compositional shifts in forest communities. Three scenarios are shown: (a) no climate warming ('Constant', used as a comparator for the following two scenarios), (b) compositional dynamic changes due to warming, and (c) accelerated growth and shortened life span due to warming. Scenarios (b) and (c) separately show two types of potential impacts of climate warming on forest dynamics. Different symbols (both shape and fill) represent individuals of different species (i.e. eight tree species are represented here). The numbers under the symbols indicate individual size, and for simplicity, it is assumed that individuals that reach size 10 die. If an individual dies, another individual (size 2) is randomly recruited from the species pool. Two types of species pools are

considered alongside air temperature differences (i.e. AT1 [no warming, constant temperature] and AT2

[warming, increase in temperature]). It is assumed that for every time step, the number doubles (e.g. from 2

to 4). The symbols of individuals in the starting communities are coloured black, while other colours indicate

individuals recruited at time 2 (blue), time 3 (red), or time 4 (green). The coloured arrows with a circle (top

right in each time step) indicate the type of species pool (filled or unfilled symbols). The period between time

1 and time 2 is shared among the three scenarios, and differences occur only after this period. In scenario (a)

'constant', it is assumed that the air temperature is constant (i.e. AT1) throughout all time periods; thus, all

individuals are recruited from species indicated by unfilled symbols. Scenario (a) is included as a comparison

between conventional and warming conditions. In both scenarios (b) and (c), it is assumed that the

temperature increases from AT1 to AT2 after time 2. In scenario (b) 'compositional dynamic changes', the

species pools of recruited individuals shift from unfilled-symbol species to filled-symbol species. After time

2, all recruited individuals are filled-symbol species under scenario (b); thus, until the composition

completely shifts from unfilled-symbol species to filled-symbol species, the contribution of individual

turnover to compositional equilibrium becomes smaller than that in the conventional condition (scenario (a)).

In scenario (c) 'accelerated growth and shortened life span', the speed of growth increases (e.g. from 2 to 4),

life spans become shorter, and the turnover of both individuals and target measures increases as a result. Note

that for simplicity, deaths other than those due to having reached natural life spans are not considered.

2.4. Case study using subset data from the FIA database for the state of Rhode Island,

USA

For a case-study application of the new individual-based QT indices based on publicly

available data, I used a subset of data from the United States Department of Agriculture

Forest Service's FIA Program for the state of Rhode Island and the R package 'rFIA'

(Stanke et al., 2020). The FIA database includes individual tree-tracking and size

information (i.e. basal area and bole biomass). A comprehensive summary of the included

data and sampling methodology is provided by Burrill et al. (2021). Briefly, each FIA plot

comprises four 0.017-ha circular subplots, each located within a circular 0.10-ha macroplot.

All free-standing woody stems (alive and dead) with a diameter at breast height (DBH) of

≥12.7 cm were sampled within each subplot (Burrill et al., 2021).

The original dataset included 10,093 individual time records for 58 species, including

multiple species (spp.) in 140 plots between 2004 and 2018. For some plots, there were

more than two surveys; therefore, more than one value of temporal beta diversity could be

calculated. I conducted plot-level filtering of the datasets using the following criteria: (1)

datasets with unidentified individuals at the species level were excluded; (2) datasets with

10 individuals or more were included; and (3) an interval of 5 ± 1 years between two

surveys was targeted. In addition, individual trees with a DBH of ≥12.7 cm were targeted

because bole biomass was also estimated. The final dataset included 5,705 individual time

records comprising 2,541 individual trees of 32 species across 126 survey intervals (86

unique plots). The latitudinal and longitudinal ranges in the dataset were 41.39446°–

42.00797° N and 71.79635°–71.12814° W, respectively. The elevation range was 10–680

m. The geographical location of each plot is shown in Fig. S2.

I focussed on the basal area and bole biomass of each tree as changes in individual sizes

that can be considered in the novel indices developed in the previous sections. The basal

area of live trees was calculated as $\frac{DBH}{2} \times \frac{DBH}{2} \times \pi$. The gross cubic-foot volume of live

trees was estimated at ≥12.7 cm DBH, defining the total volume of wood in the central

stem of the inventoried trees (bole biomass) assuming a 30.48-centimetre mean stump

height and a minimum 10.16-centimetre top diameter (for a detailed estimation method,

see Domke et al., 2013).

 Table 1. Summary of indices and formulae developed in previous studies and used in this study.

Index	Formula	Reference	Ecological interpretation
d_{BC}	$\frac{B+C}{2A+B+C} = \frac{\sum_i[x_{ij} - \min(x_{ij}, x_{ik})] + \sum_i[x_{ik} - \min(x_{ij}, x_{ik})]}{2[\sum_i \min(x_{ij}, x_{ik})] + \sum_i[x_{ij} - \min(x_{ij}, x_{ik})] + \sum_i[x_{ik} - \min(x_{ij}, x_{ik})]}$	Odum (1950)	The compositional shift based on the change in the number of individuals for each species in a community across two time periods ($0 \leq d_{BC} \leq 1$).
d_{mr}	$\frac{m+r}{2p+m+r} = \frac{\sum_o[z_{oj} - \min(z_{oj}, z_{ok})] + \sum_o[z_{ok} - \min(z_{oj}, z_{ok})]}{2[\sum_o \min(z_{oj}, z_{ok})] + \sum_o[z_{oj} - \min(z_{oj}, z_{ok})] + \sum_o[z_{ok} - \min(z_{oj}, z_{ok})]}$	Nakadai	The

		(2020) individual turnover by mortality and recruitment in a community across two time periods $(0 \le d_{mr} \le 1)$.
$v_{s.QL}$	$\dfrac{B+C}{2e+B+C} = \dfrac{B+C}{m+r} = \dfrac{\sum_i \left[x_{ij} - \min(x_{ij}, x_{ik})\right] + \sum_i \left[x_{ik} - \min(x_{ij}, x_{ik})\right]}{2\left[\sum_i \min(x_{ij}, x_{ik}) - \sum_o \min(z_{oj}, z_{ok})\right] + \sum_i \left[x_{ij} - \min(x_{ij}, x_{ik})\right] + \sum_i \left[x_{ik} - \min(x_{ij}, x_{ik})\right]}$	Nakadai (2020) The relative speed of compositional shift in relation to individual turnover $(0 \le v_{s.QL} \le 1)$.

$d_{BC'}$

$$\frac{B'+C'}{2A'+B'+C'} = \frac{\sum_i[y_{ij}-\min(y_{ij},y_{ik})]+\sum_i[y_{ik}-\min(y_{ij},y_{ik})]}{2[\sum_i\min(y_{ij},y_{ik})]+\sum_i[y_{ij}-\min(y_{ij},y_{ik})]+\sum_i[y_{ik}-\min(y_{ij},y_{ik})]}$$

Odum (1950)

The compositional shift based on the changes in the targeted measures in a community across two time periods ($0 \le d_{BC'} \le 1$).

d_{MRG}

$$\frac{M+R+G}{2P+M+R+G}$$

$$=\frac{\sum_o\left[[w_{oj}-\min(w_{oj},w_{ok})]\times[z_{oj}-\min(z_{oj},z_{ok})]\right]+\sum_o\left[[w_{ok}-\min(w_{oj},w_{ok})]\times[z_{ok}-\min(z_{oj},z_{ok})]\right]+\sum_o\left[[\max(w_{oj},w_{ok})-\min(w_{oj},w_{ok})]\times\min(z_{oj},z_{ok})\right]}{2[\sum_o\min(w_{oj},w_{ok})]+\sum_o\left[[w_{oj}-\min(w_{oj},w_{ok})]\times[z_{oj}-\min(z_{oj},z_{ok})]\right]+\sum_o\left[[w_{ok}-\min(w_{oj},w_{ok})]\times[z_{ok}-\min(z_{oj},z_{ok})]\right]+\sum_o\left[[\max(w_{oj},w_{ok})-\min(w_{oj},w_{ok})]\times\min(z_{oj},z_{ok})\right]}$$

This study

The turnover of the targeted measures by growth,

mortality, and recruitment in a community across two time periods ($0 \leq d_{MRG} \leq 1$).

$v_{s.QT}$

$$\frac{B' + C'}{2E + B' + C'} = \frac{\sum_i \left[y_{ij} - \min(y_{ij}, y_{ik})\right] + \sum_i \left[y_{ik} - \min(y_{ij}, y_{ik})\right]}{2\left[\sum_i \min(y_{ij}, y_{ik}) - \sum_o \min(w_{oj}, w_{ok})\right] + \sum_i \left[y_{ij} - \min(y_{ij}, y_{ik})\right] + \sum_i \left[y_{ik} - \min(y_{ij}, y_{ik})\right]}$$

This study

The relative speed of compositional shift in relation to the turnover of the target measure (0

x_{ij} is the abundance of species i at temporal unit j, and x_{ik} is the abundance of species i at temporal unit k. y_{ij} is abundance excluding individuals (i.e. measures or

weights) of species i at time Tj at a site, and y_{ik} is abundance excluding individuals of species i at time Tk at a site. z_{oj} is individual o at time Tj, and z_{ok} is

individual o at time Tk. Both z_{oj} and z_{ok} have values of one or zero (presence and absence, respectively). w_{oj} is the measure of individual o at time Tj, and w_{ok} is

the measure of individual o at time Tk. The equations of d_{BC} and $d_{BC'}$ are mathematically the same, and only the target abundances are different.

To demonstrate the application of the novel indices, I focussed mainly on the

following 11 indices: d_{BC}, d_{mr}, d_m, d_r, $v_{s.QL}$, $d_{BC'}$, d_{MRG}, d_M, d_R, d_G, and $v_{s.QT}$. Formulas

of the base indices (i.e. d_{BC}, d_{mr}, $v_{s.QL}$, $d_{BC'}$, d_{MRG}, and $v_{s.QT}$) are summarised in Table 1.

All 11 indices were calculated for each survey interval for each plot. The latter six

indices (i.e. $d_{BC'}$, d_{MRG}, d_M, d_R, d_G, and $v_{s.QT}$) were calculated for both basal area and

bole biomass, distinguished by the suffixes '_area' or '_biomass', respectively.

First, to show the contribution of each component to individual-based QT

dissimilarity d_{MRG}, I estimated kernel density for each partitioned component (d_M, d_R,

d_G, d_S, and d_E) of the dissimilarity indices. Second, I evaluated the relationships

among d_{BC}, d_{mr}, $v_{s.QL}$, $d_{BC'}$, d_{MRG}, and $v_{s.QT}$ using simple linear regression. Third, to

reveal the geographical and temporal components of the indices, I ran linear mixed

models for each index. As explanatory variables, geographic factors (latitude and

elevation), a temporal factor (mean calendar year between the first and second

surveys), and a sampling design-related factor (length of survey interval) were used.

Plot identity was considered a random effect in the model comparisons because 40

plots included two survey intervals. However, model comparisons showed that the

linear mixed models consistently had higher Akaike information criterion (AIC;

Akaike, 1998) values than those of the linear models (Table S2). Thus, all results

discussed in the following sections are based on linear models.

All of the analyses were conducted using R software (ver. 4.1.0; R Development

Core Team, 2021). The R packages '*vegan*' (Oksanen et al., 2020), '*nlme*' (Pinheiro

et al., 2022), and '*MuMIn*' (Barton, 2022) were used to convert the abundance data to

presence-absence data, run linear mixed models, and perform model comparisons

based on AIC values. The R scripts used to calculate the new indices are provided in

Supplementary File 1.

For this case study, I did not use a dataset specifically constructed for climate

change assessment; hence, it is not possible to directly test the hypothetical impacts

of climate warming on compositional shifts in forest communities. However, this

approach helps understand the mechanisms of climate change impacts by considering

changes in individual factors. Although I focus on climate change here, the indices

introduced in this study can be useful in cases where individual growth rates vary

over time for intrinsic reasons (e.g. ecological succession).

3. Results

Among the 126 survey intervals, the individual turnover of 10 survey intervals was

zero; therefore, these intervals were excluded from the targets when calculating the

$v_{s.QL}$ indices. The distribution of the novel indices, except $v_{s.QT}$, is shown in Fig. 3.

The contribution of individual growth to the total turnover of both basal area and bole

biomass was generally larger than that of mortality and recruitment (Fig. 3a, b; Fig.

S3). The d_{MRG} indices based on basal area and bole biomass were strongly and

positively correlated with d_{BC}, d_{mr}, area-based $d_{BC'}$, and biomass-based $d_{BC'}$ (≥ 0.87;

Fig. S4). In contrast, the $v_{s.QT}$ indices based on basal area and bole biomass were only

weakly correlated with the $d_{BC'}$ indices based on basal area and bole biomass ($r =$

0.23–0.28, Fig. S4) and showed no correlation with $v_{s.QL}$ ($r = 0.16$ and 0.08, Fig. S4).

For both d_{MRG} and $v_{s.QT}$, strong correlations existed between the indices based on

basal area and bole biomass ($r = 0.99$ and 0.91, Fig. S4). All Pearson correlation

coefficients are summarised in Fig. S4.

Fig. 3. Density plots representing the distribution of individual-based QT beta-diversity indices

across 126 survey intervals for 86 plots (data from Stanke et al. 2020). In all four plots, the index

values are limited to the range 0–0.15 for comparison (see Fig. S3 for a plot of the entire range).

The plots in panels (a) and (c) are based on basal area, and those in (b) and (d) are based on bole

biomass.

Based on the AIC comparison, models including only intercept were selected for

the three indices, i.e. the indices for compositional shift based on the number of

individuals and basal area (d_{BC} and $d_{BC'\_area}$), and the relative speed of compositional

shift against total changes in basal area ($v_{s.QT\_area}$) (Table 2). The index of

compositional shift based on bole biomass ($d_{BC'\_biomass}$) tended to increase with longer

survey intervals, as indicated by model comparison (Table 2). As for the index of the

relative speed of compositional shift against total individual turnover ($v_{s.QL}$), only

elevation was selected as an explanatory variable based on the AIC comparison; thus,

the index increased with increasing elevation (Table 2). For the relative speed of

compositional shift against total changes in basal area ($v_{s.QT\_biomass}$), a model including

calendar year and survey interval was selected based on the AIC comparison (Table

2); specifically, the index increased with increasing calendar year and longer survey

intervals. In both indices of the total change in basal area and bole biomass (d_{MRG_area}

and $d_{MRG\_biomass}$), models with latitude and survey interval were selected as the best

models based on the AIC comparisons (Table 2); specifically, the indices tended to

increase with decreasing latitude and longer survey intervals. However, in the index

of total individual turnover (d_{mr}), the model including only the survey interval was

selected as the best model (Table 2), with the index increasing with longer survey

intervals. As for the three indices related to the contribution of mortality to the total

change (d_m, d_{M_area}, and $d_{M\_biomass}$), models including latitude and calendar year were

selected as the best models (Table 2), and index values tended to increase with

decreasing latitude and increasing calendar year. For the indices related to the

contribution of recruitment to the total change (d_{R_area}, $d_{R\_biomass}$), models including

calendar year were selected as the best models (Table 2), with index values tending to

increase with increasing calendar year. For the index related to the contribution of

recruitment to the total individual turnover (d_r), the model including calendar year

and survey interval was selected as the best model (Table 2), with the index tending

to increase with increasing calendar year and longer survey intervals. For both indices

related to the contribution of growth to the total change (d_{G_area} and $d_{G\_biomass}$), the

models that included the survey interval and, in the case of basal area, calendar year,

were selected.

Table 2. Results of a multi-regression model based on the best models selected using an Akaike information criterion (AIC) comparison for individual-based

qualitative (QL) and quantitative (QT) beta-diversity indices.

Latitude	Elevation	Calendar year	Survey interval	*Adj R2*	*P-value*

556   *β* and *parR2* indicate the regression coefficients and partially adjusted *R*-squared values of each explanatory variable, respectively. *Adj R2* is the adjusted *R*-

557   squared value for the entire model. Note: Significance: <0.1, *<0.05, **<0.01, ***<0.001. To indicate a model where only the intercept was selected, † was

558   added to the *P*-value column for the entire model.

	β	parR2	P-value		β	parR2	P-value		β	parR2	P-value		β	parR2	P-value		parR2	P-value	
(a) Qualitative family indices																				
d_{BC}	–	–	–		–	–	–		–	–	–		–	–	–					†
d_{mr}	–	–	–		–	–	–		–	–	–		0.0186	0.0089	0.1473		0.0089	0.1473	
d_{m}	-0.0807	0.0333	0.0233	*	–	–	–		0.0098	0.0953	0.0003	***	–	–	–		0.1211	0.0001	***
dr	–	–	–		–	–	–		-0.0129	0.0513	0.0064	**	0.0277	0.0146	0.0944	.	0.0554	0.0111	*
$v_{s.QL}$	–	–	–		0.0002	0.0283	0.0393	*	–	–	–		–	–	–		0.0283	0.0393	*
(b) Quantitative family indices																				
(i) Basal area																				
$d_{BC'}$	–	–	–		–	–	–		–	–	–		–	–	–					†
d_{MRG}	-0.0705	0.0095	0.1413		–	–	–		–	–	–		0.0231	0.0192	0.0663	.	0.0269	0.0695	.
d_{M}	-0.0768	0.0253	0.0421	*	–	–	–		0.0094	0.0770	0.0010	**	–	–	–		0.0972	0.0007	***
d_{R}	–	–	–		–	–	–		-0.0044	0.0259	0.0398	*	–	–	–		0.0259	0.0398	*
d_{G}	–	–	–		–	–	–		-0.0028	0.0236	0.0477	*	0.0122	0.0381	0.0165	*	0.0305	0.0553	.
$v_{s.QT}$	–	–	–		–	–	–		–	–	–		–	–	–					†
(ii) Bole biomass																				
$d_{BC'}$	–	–	–		–	–	–		–	–	–		0.0205	0.0114	0.1210		0.0114	0.1210	
d_{MRG}	-0.0748	0.0094	0.1429		–	–	–		–	–	–		0.0278	0.0264	0.0387	*	0.0336	0.0454	*
d_{M}	-0.0766	0.0209	0.0586	.	–	–	–		0.0092	0.0641	0.0026	***	–	–	–		0.0812	0.0020	**
d_{R}	–	–	–		–	–	–		-0.0039	0.0187	0.0684	.	–	–	–		0.0187	0.0685	.
d_{G}	–	–	–		–	–	–		–	–	–		0.0089	0.0227	0.0503	.	0.0227	0.0503	.
$v_{s.QT}$	–	–	–		–	–	–		-0.0301	0.0272	0.0366	*	0.0717	0.0082	0.1569		0.0232	0.0877	.

4. Discussion

The potential for evaluating the contributions of individuals' persistence to compositional shifts in a community was first proposed by Nakadai (2020). In this study, I extended previously established individual-based beta-diversity approaches (Nakadai, 2020, 2021) to account for individual property changes—even for persistent individuals—that are not currently considered in qualitative indices. Specifically, in addition to the components of mortality and recruitment as considered by Nakadai (2020), I introduced a growth component to account for the individual size changes in the context of compositional shifts for the first time (Fig. 1). Also, using hypothetical scenarios, I illustrated the biological meaning of the individual-based indices in the context of climate warming (Fig. 2). The novel indices proposed here not only represent methodological progress in community ecology and macroecology, but also aid conceptual advancement by bridging studies in biodiversity with those on individual life history and physiology.

To illustrate the application of the novel indices, I applied them to a subset of data from the FIA database for the state of Rhode Island, USA. The analysis demonstrated that the contribution of tree growth to the total turnover of both basal area and bole biomass was generally larger than that of mortality and recruitment (Fig. 3a, b; Fig. S3). This indicates the importance of previously overlooked growth factors in forest turnover and dynamics and the consequent compositional shifts. In the case-study dataset, the calculated abundance- and individual-based indices (i.e. d_{BC}, d_{mr}, $d_{BC'}$, and d_{MRG}) were strongly correlated (Fig. S4). Thus, if the contribution of individual tree growth is high, the strength of the correlation between d_{mr} and d_{MRG} should decrease, yet this was not the case. I found no clear relationships between the $v_{s.QL}$ and $v_{s.QT}$ indices based on both basal area and bole biomass and the relative speed of compositional shift. Generally, the $v_{s.QL}$ values for the FIA dataset were high; thus, individual turnover generally contributed to compositional shift, although both $v_{s.QT\_area}$ and $v_{s.QT\_biomass}$ were lower and more variable than $v_{s.QL}$. These results reflect the differences in the characteristics of the novel QT indices developed here and the conventional QL indices. Indeed, $v_{s.QL}$ could not be applied to 10 plots that had no individual turnover, whereas in such cases, the $v_{s.QT}$ index was more sensitive to temporal changes in the community than the $v_{s.QL}$ index.

In the driver analysis of the calculated indices, I found that various explanatory variables affected the results (Table 2). First, focussing on the growth component specific to the new QT index, both area- and biomass-based d_G indices, which indicate the contribution of growth, were responsive to the survey interval. This strongly suggests that an accurate evaluation of the contribution of growth was achieved. In addition, the area-based d_G index tended to decrease with increasing (i.e. more recent) calendar year. In addition, overall, the contributions related to recruitment decreased, and those related to mortality increased (Table 2), which may be indicative of succession (i.e. aging forests) and/or climate change.

The $v_{s.QT\_area}$ index showed no clear pattern because the model including only intercept was selected, whereas the $v_{s.QL}$ and $v_{s.QT\_biomass}$ indices varied with elevation and calendar year, respectively. The v_s family of indices is designed to capture changes to account for stochastic drifts, such as individual, basal area, or bole-volume turnover. The significant relationships between several v_s indices with environmental variables suggest that along with the explanatory variables, niche processes drive compositional shifts, which cannot be explained by stochastic processes alone. For example, Nakadai (2020) showed that $v_{s.QL}$ values vary along climate gradients (i.e. the change ratio of annual mean temperature) for both deciduous and evergreen broadleaf forests.

The indices developed in the present study are applicable to classical forest inventory surveys in which all stems are tagged and their DBH measured. Here, I describe some potential applications of the novel indices as examples. First, as a representative example, applying the individual-based temporal beta-diversity indices to global datasets, especially forest inventory data (e.g. ForestGEO; https://forestgeo.si.edu/), is highly promising for assessing the impacts of global climate change. Specifically, the responses of the novel indices to both the mean and variable ratios of air temperature are important for mechanistically understanding the compositional shifts in forest tree communities. Furthermore, the average length of an individual's life cycle (i.e. longevity) varies across the stages of ecological succession. As a base condition to study the impacts of climate change on forest tree communities, the patterns of the novel indices along the stages of ecological succession could be one of the next challenges.

Although the indices introduced in this study are mainly applicable to datasets
containing both individual identity and size information, a method that accommodates
various types of quantitative individual information may become available through future
theoretical developments. Generally, information for individuals is limited in community
ecology and macroecology, yet in the era of big data, this is likely to be addressed in the
near future. In addition, if the contributions of demographic factors (i.e. growth, mortality,
and recruitment) can be estimated using hierarchical modelling approaches (Hill et al.,
2004, Kanamori et al., 2017), these indices could be calculated even in the absence of
detailed individual-tracking information. Furthermore, although the survey interval length
is treated as a sampling design-related factor in the present study, revealing the temporal
patterns throughout a survey interval, and therefore the temporal scale dependency, which
is called temporal distance decay (Hatosy et al., 2013), is also a future challenge.

5. Conclusion

Notably, climate change affects various ecological hierarchies, from global biodiversity to
individuals and even intra-individual phenomena (Arneth et al., 2020; Körner, 2017).
Indeed, using the newly developed individual-based temporal QT beta-diversity indices,
this study showed that the growth components largely contributed to forest community
compositional dynamics compared to mortality and recruitment components in the case-
study dataset. The important contribution of the growth components to temporal
compositional shifts (i.e. temporal beta diversity) can be evaluated using the QT indices
developed in this study, thus conventional abundance-based and individual-based indices
(Odam, 1950; Nakadai, 2020) cannot evaluate changes in individual sizes. Additionally,
the individual-based indices can facilitate our understanding of the link between
demographic processes and community compositional shifts, which conventional
approaches have not achieved until now. As in the Rhode Island case study, individual-
based diversity indices that account for previously overlooked elements of biodiversity,
such as individual persistence and growth, pave the way for further innovative research on
individual-based biodiversity. This should facilitate a greater understanding of the effects
of climate change across multiple hierarchies of biological organisation.

Author contributions

Ryosuke Nakadai: Conceptualization, Methodology, Formal analysis, Writing – original

draft, review & editing.

Funding

RN was supported by the Japan Society for the Promotion of Science (grant No.

22K15188) and the National Institute for Environmental Studies, Japan.

Conflicts of interest

The author declares no conflicts of interest.

Data availability

Data are available from the R package rFIA (Stanke et al., 2020), and the code used to

calculate the indices in the present study is provided in Supplementary File 1.

References

Akaike, H., 1998. Information theory and an extension of the maximum likelihood

principle. In: E. Parzen, K. Tanabe, and G. Kitagawa (Eds.), Selected Papers of

Hirotugu Akaike. Springer, New York, pp. 199–213. https://doi.org/10.1007/978-1-

4612-1694-0_15.

Arneth, A., Shin, Y. J., Leadley, P., Rondinini, C., Bukvareva, E., Kolb, M., Midgley, G. F.,

Oberdorff T., Palomo, I., Saito, O., 2020. Post-2020 biodiversity targets need to

embrace climate change. Proc. Natl. Acad. Sci. U. S. A. 117, 30882–30891.

https://doi.org/10.1073/pnas.2009584117.

Bartoń, K., 2020. MuMIn: Multi-model inference. R package version 1.43.17.

https://CRAN.R-project.org/package=MuMIn.

Baselga, A., 2013. Separating the two components of abundance-based dissimilarity:

balanced changes in abundance vs. abundance gradients. Methods Ecol. Evol. 4, 552–

557. https://doi.org/10.1111/2041-210X.12029.

Brienen, R. J. W., Caldwell, L., Duchesne, L., Voelker, S., Barichivich, J., Baliva, M.,

Ceccanthini, G., Di Fillippo, A., Helama, S., Locosselli, G. M., Lopez, L. Piovesan, J.,

Villabla, R., Gloor, E., 2020. Forest carbon sink neutralized by pervasive growth-

lifespan trade-offs. Nat. Commun. 11, 4241. https://doi.org/10.1038/s41467-020-

17966-z.

Brienen, R., Phillips, O., Feldpausch, T. Gloor, E., Baker, T., Lloyd, J., Lopez-Gonzalez, G.,

Monteagudo-Mendoza, A., Malhi, Y., Lewis, S. L., Vásquez Martinez, R., Alexiades, M.,

Álvarez Dávila, E., Alvarez-Loayza, P., Andrade, A., Aragão, L. E. O. C., Araujo-

Murakami, A., Arets, E. J. M. M., Arroyo, L., Aymard C. G. A., Bánki, O. S., Baraloto,

C., Barroso, J., Bonal, D., Boot, R. G. A., Camargo, J. L. C., Castilho, C. V., Chama, V.,

Chao, K. J., Chave, J., Comiskey, J. A., Cornejo Valverde, F., da Costa, L., de Oliveira, E.

A., Di Fiore, A., Erwin, T. L., Fauset, S., Forsthofer, M., Galbraith, D. R., Grahame, E. S.,

Groot, N., Hérault, B., Higuchi, N., Honorio Coronado, E. N., Keeling, H., Killeen, T. J.,

Laurance, W. F., Laurance, S., Licona, J., Magnussen, W. E., Marimon, B. S., Marimon-

Junior, B. H., Mendoza, C., Neill, D. A., Nogueira, E. M., Núñez, P., Pallqui Camacho, N.

C., Parada, A., Pardo-Molina, G., Peacock, J., Peña-Claros, M., Pickavance, G. C.,

Pitman, N. C. A., Poorter, L., Prieto, A., Quesada, C. A., Ramírez, F., Ramírez-Angulo,

H., Restrepo, Z., Roopsind, A., Rudas, A., Salomão, R. P., Schwarz, M., Silva, N., Silva-

Espejo, J. E., Silveira, M., Stropp, J., Talbot, J., ter Steege, H., Teran-Aguilar, J.,

Terborgh, J., Thomas-Caesar, R., Toledo, M., Torello-Raventos, M., Umetsu, R. K., van

der Heijden, G. M. F., van der Hout, P., Guimarães Vieira, I. C., Vieira, S. A., Vilanova,

E., Vos, V. A., Zagt, R. J., 2015. Long-term decline of the Amazon carbon sink. Nature

519, 344–348. https://doi.org/10.1038/nature14283.

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., West, G. B., 2004. Toward a

metabolic theory of ecology. Ecology 85, 1771–1789. https://doi.org/10.1890/03-9000.

Brown, J. H., 1995. Macroecology. University of Chicago Press.

Brown, J. H., 1999. Macroecology: Progress and prospect. Oikos 87, 3–14.

https://doi.org/10.2307/3546991.

Burrill, E. A., DiTommaso A. M., Turner, J. A., Pugh, S. A., Menlove, J., Christiansen, G.,

Perry, C. J., Conkling, B. L. 2021. The forest inventory and analysis database:

Database description and user guide version 9.0.1 for Phase 2. U.S. Department of

Agriculture, Forest Service, p. 1026. http://www.fia.fs.fed.us/library/database-

documentation/.

Domke, G. M., Oswalt, C. M., Woodall, C. W., Turner, J. A., 2013. Estimation of

merchantable bole volume and biomass above sawlog top in the National Forest

Inventory of the United States. J. For. 111, 383–387. https://doi.org/10.5849/jof.13-
042

Gotelli, N. J., Shimadzu, H., Dornelas, M., McGill. B., Moyes. F., & Magurran, A. E. 2017.
Community-level regulation of temporal trends in biodiversity. Sci. Adv. 3, e1700315.
https://doi.org/10.1126/sciadv.1700315.

Hatosy, S. M., Martiny, J. B., Sachdeva, R., Steele, J., Fuhrman, J. A., Martiny, A. C.
(2013). Beta diversity of marine bacteria depends on temporal scale. Ecology 94(9),
1898–1904. https://doi.org/10.1890/12-2125.1

Hill, M. F., Witman, J. D., Caswell, H., 2004. Markov chain analysis of succession in a
rocky subtidal community. Am. Nat. 164, 46–61. https://doi.org/10.1086/422340.

Jaccard, P., 1901. Distribution de la flore alpine dans le bassin des Dranses et dans
quelques régions voisines. Bull. de la Soc. Vaud. des Sci. Nat. 37, 241–272.

Kanamori Y., Fukaya, K., Noda, T., 2017. Seasonal changes in community structure along
a vertical gradient: Patterns and processes in rocky intertidal sessile assemblages.
Popul. Ecol. 59, 301–313. https://doi.org/10.1007/s10144-017-0596-z.

Körner, C., 2017. A matter of tree longevity. Science 355, 130–131.
https://doi.org/10.1126/science.aal2449.

Legendre, P., Legendre, L., 2012. Numerical ecology (3rd ed.). Amsterdam: Elsevier.

Legendre, P., 2014. Interpreting the replacement and richness difference components of
beta diversity. Glob. Ecol. Biogeogr. 23, 1324–1334.

Legendre, P., 2019. A temporal beta-diversity index to identify sites that have changed in
exceptional ways in space–time surveys. Ecol. Evol. 9, 3500–3514.
https://doi.org/10.1002/ece3.4984.

Li, S. P., Cadotte, M. W., Meiners, S. J., Pu, Z., Fukami, T., Jiang, L., 2016. Convergence
and divergence in a long-term old-field succession: The importance of spatial scale
and species abundance. Ecol. Lett. 19, 1101–1109. https://doi.org/10.1111/ele.12647.

Magurran, A. E., Dornelas, M., Moyes, F., Henderson, P. A., 2019. Temporal β diversity: A
macroecological perspective. Glob. Ecol. Biogeogr. 28, 1949–1960.
https://doi.org/10.1111/geb.13026.

Munné-Bosch, S., 2018. Limits to tree growth and longevity. Trends Plant Sci. 23, 985–
993. https://doi.org/10.1016/j.tplants.2018.08.001.

Nakadai, R., 2020. Degrees of compositional shift in tree communities vary along a
gradient of temperature change rates over one decade: Application of an individual-
based temporal beta-diversity concept. Ecol. Evol. 10, 13613–13623.
https://doi.org/10.1002/ece3.6579.

Nakadai, R., 2021. Individual-based multiple-unit dissimilarity: novel indices and null
model for assessing temporal variability in community composition. Oecologia 197,
353–364. https://doi.org/10.1007/s00442-021-05025-3.

Odum, E. P., 1950. Bird populations of the Highlands (North Carolina) Plateau in relation
to plant succession and avian invasion. Ecology 31, 587–605.
https://doi.org/10.2307/1931577.

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin,
P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Scöcs, E.,
Wagner, H. H., 2020. Vegan: Community ecology package. R package version 2.5-7.
https://CRAN.R-project.org/package=vegan.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., R Core Team., 2022. Nlme: Linear and
nonlinear mixed effects models. R package version 3.1-155. https://CRAN.R-
project.org/package=nlme.

Podani, J., Ricotta, C., Schmera, D., 2013. A general framework for analyzing beta
diversity, nestedness and related community-level phenomena based on abundance
data. Ecol. Complex. 15, 52–61.

R Development Core Team., 2021. R: A language and environment for statistical
computing. R Foundation for Statistical Computing, Vienna, Austria. (ver. 4.1.0).
http://www.R-project.org/.

Ružička, M., 1958. Anwendung mathematisch–statisticher Methoden in der Geobotanik
(synthetische Bearbeitung von Aufnahmen). Biologia (Bratislava) 13, 647–661.

Searle, E. B., Chen, H. Y., 2018. Temporal declines in tree longevity associated with faster
lifetime growth rates in boreal forests. Environ. Res. Lett. 13, 125003.
https://doi.org/10.1088/1748-9326/aaea9e.

Shimadzu, H. Dornelas, M., Magurran, A. E., 2015. Measuring temporal turnover in
ecological communities. Methods Ecol. Evol. 6, 1384–1394.
https://doi.org/10.1111/2041-210X.12438.

Sørensen, T. A., 1948. A method of establishing groups of equal amplitude in plant
sociology based on similarity of species content and its application to analyses of the
vegetation on Danish commons. Biol. Skr. 5, 1–34.
Stanke, H., Finley, A. O., Weed, A. S., Walters, B. F., Domke, G. M., 2020. rFIA: An R
package for estimation of forest attributes with the US Forest Inventory and Analysis
database.   Environ.   Model.   Softw.   127,   104664.
https://doi.org/10.1016/j.envsoft.2020.104664.
Tatsumi, S., Iritani, R., & Cadotte, M. W., 2021. Temporal changes in spatial variation:
Partitioning the extinction and colonisation components of beta diversity. Ecol. Lett.
24(5), 1063–1072. https://doi.org/10.1111/ele.13720.
789
790

Supplementary files

Table S1. Table summary to identify the applicable temporal beta-diversity indices for the community dataset

Table S2. Comparison of linear and linear mixed models for each index. The best models selected from the comparison are shown in Table 2

Fig. S1. Venn diagrams related to the calculation of both conventional and novel indices. Specifically, the first four Venn diagrams (a–d) are based on the number of individuals, and latter four (e–h) are based on the total values of targeted measures excluding the number of individuals (e.g. sizes and weights).

Fig. S2. Map of geographic locations of plots included in the subset of data from the Forest Inventory and Analysis (FIA) database for the state of Rhode Island, USA (Stanke et al., 2020). Sites indicated with darker-coloured points have multiple survey intervals.

Fig. S3. Density plots representing the distribution of individual-based quantitative beta-diversity indices across 126 survey intervals in 86 plots (data from Stanke et al., 2020). In the right four plots (b, d, f, and h), the index values are limited to the range 0–0.15 for comparison (also see Fig. 2). In the left four plots, the entire value range is shown. Plots in panels (a), (b), (e), and (f) are based on the basal area, and those in panels (c), (d), (g), and (h) on bole biomass.

Fig. S4. Correlations among the calculated indices. Pearson coefficients of correlation are shown. Note: Significance: <0.1, *<0.05, **<0.01, ***<0.001.

Supplementary Text 1. Detailed explanation for the components in the weighted Bray–Curtis index

Supplementary Text 2. Explanation of individual-based temporal beta diversity employing the Ružička dissimilarity index.

Supplementary file 1. R software script for individual-based temporal beta-diversity calculations using sample data.