Comparative genomics in Orobanchaceae provides insight into the origin and evolution of plant parasitism

Yuxing Xu[a], Jingxiong Zhang[a], Canrong Ma[a, c], Yunting Lei[a], Guojing Shen[a], Jianjun Jin[d], Deren A. R. Eaton[d], Jianqiang Wu[a, b]

[a] Department of Economic Plants and Biotechnology, Yunnan Key Laboratory for Wild Plant Resources, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China; [b] CAS Center for Excellence in Biotic Interactions, University of Chinese Academy of Sciences, Beijing 100049, China; [c] University of Chinese Academy of Sciences, Beijing 100049, China; [d] Department of Ecology, Evolution and Environmental Biology, Columbia University, New York, NY 10027, USA

*Jianqiang Wu

Email: wujianqiang@mail.kib.ac.cn

Author Contributions: Y.X. and J.W. designed research; J.Z., C.M., and G.S. prepared plant tissues, DNA samples, and RNA samples; Y.X., J.W., Y.L., G.S., and J.J. analyzed data; Y.X., J.W., J.J., and D.E. wrote the paper.

Competing Interest Statement: The authors declare no competing interest.

Classification: Biological Sciences: Plant Biology and Evolution.

Keywords: Genomics, Orobanchaceae, Parasitism, Whole-Genome Duplication, Gene Loss, Gene family expansion.

Abstract

Orobanchaceae is the largest family of plant parasites comprising autotrophic and parasitic plants

with all degrees of parasitism, making it by far the best family for studying the origin and evolution

of plant parasitism. Here we provide three high-quality genomes of orobanchaceous plants, the

autotrophic *Lindenbergia luchunensis* and holoparasitic *Phelipanche aegyptiaca* and *Orobanche*

cumana. Phylogenomic analysis on these three genomes and the previously published genomes

and transcriptomes of other orobanchaceous species created a robust phylogenetic framework of

Orobanchaceae. An ancient whole-genome duplication (WGD; about 79.29 Mya) was detected in

the common ancestor of Orobanchaceae and *Mimulus guttatus*, and this event might have

contributed to the origin of parasitism; however, except for *Striga*, no WGD events occurred in

any lineages of orobanchaceous parasites after divergence from their autotrophic common

ancestor, and the expansion of the genomes of orobanchaceous parasitic plants was mainly due

to retrotransposons. We detected evident convergent gene loss in all parasites within

Orobanchaceae and between Orobanchaceae and dodder *Cuscuta australis*. The gene families

in the orobanchaceous parasites showed a clear pattern of recent gains and expansions, and the

expanded gene families are enriched in functions related to development of haustorium. Thus,

recent gene family expansions may have facilitated the adaptation of orobanchaceous parasites

to different hosts. These data well support the previously proposed model of parasitic plant

genome evolution (1). This work illustrates the trajectory of the evolution of orobanchaceous

parasites, and will facilitate future studies on parasitism and the control of parasitic plants in

agriculture.

Introduction

Most plants are autotrophs. However, about 4750 angiosperms are parasitic plants (2), which completely or partly depend on hosts to survive by extracting water and nutrients from the host plants. Some parasitic plants still have chlorophyll and photosynthesize, and they are named hemiparasitic plants, while some no longer photosynthesize and these are called holoparasites. According to whether hosts are needed for the completion of life cycles, parasitic plants can also be classified as the facultative ones and the obligate ones (3). In angiosperms, parasitic plants independently evolved 12 or 13 times from autotrophic plants (3). In eight parasitic plant lineages, hemiparasitic species became extinct, leaving only holoparasites remaining until today (3), whereas Orobanchaceae (Lamiales), is the largest family of parasitic plants, flourished with all types of parasitic lifestyles: facultative hemiparasites and obligate hemi- and holoparasites (Fig. 1) (3, 4). Orobanchaceae even has three genera of autotrophs: *Rehmannia*, *Triaenophora*, and *Lindenbergia*, and *Lindenbergia* is the closest relative to the parasitic lineages (5). Furthermore, some of the orobanchaceous parasites are economically important for being devastating pests in agriculture. For example, the annual economic loss caused by the witchweed (*Striga*) and broomrapes (*Orobanche* and *Phelipanche*) is over 10 billion US dollars in Asia and Africa (6). Orobanchaceae is thus the best extant system in which to study the evolutionary transition to plant parasitism, as well as to develop genomic resources for future agricultural applications (3).

Phylogenetic relationships among the major lineages of Orobanchaceae remain unclear. Previous research by McNeal et al. (7) split the family into six major clades based on three nuclear and two plastid markers, while a more recent study by Li et al. (5) analyzed five additional markers and recognized eight major clades, with incongruent relationships compared to the previous treatment. Following the Angiosperm Phylogeny Group nomenclature the Li et al. treatment recognized Rehmannieae, Lindenbergieae, Cymbarieae, Orobancheae, *Brandisia*, Rhinantheae, Buchnereae, and Pedicularideae. A surprising result of this study was the placement of the holoparasitic clade Orobancheae sister to all other parasitic Orobanchaceae, which suggests holoparasitism may have evolved earlier than other forms of parasitism. This conflicts with the proposed that the common ancestor of orobanchaceous parasitic plants, which was a facultative hemiparasite, evolved from an autotrophic plant, and later obligate hemiparasites appeared, and the latest ones were the obligate holoparasites, which no longer photosynthesize (3).

The evolution of parasitism requires major morphological and physiological innovations to perceive and recognize potential host plants and to form parasitic connections. In this respect, parasitic plants evolved a unique organ, haustorium, using which these parasites can attach and

penetrate host tissues (8). The haustoria of hemiparasites, such as *Triphysaria* and *Striga*, fuse

with the xylem of host plants, enabling these hemiparasites to obtain water and nutrients in the

host xylem sap. Holoparasites respectively fuse their haustorial xylem and phloem with the xylem

and phloem of hosts, and the fusion between parasite and host phloem tissues form symplastic

continuity by plasmodesmatal connections (8). In Orobanchaceae, the obligate hemiparasites and

holoparasites can perceive strigolactones released from host roots and activate the germination

process (9, 10). Recent studies revealed that in the orobanchaceous obligate parasites, the

karrikin receptor gene *KAI2* evolved *KAI2d* by duplication and neofunctionalization, and KAI2d

perceive the host root-released strigolactones, enabling these parasites to detect the existence of

host plants in the vicinity (11–13). After germination, some parasites also sense certain host

signals, named haustorium-inducing factors, such as DMBQ (2,6-dimethoxy-p-benzoquinone),

which is a cell wall-derived simple quinone (8), and thus develop haustoria.

Previous studies on *Cuscuta australis* (Convolvulaceae) (14), a shoot parasitic plant, *Striga*

asiatica (Orobanchaceae) (15), and an endoparasite *Sapria himalayana* (Rafflesiaceae) (16)

indicated that gene loss is common in parasitic plants. Wicke et al. (17) studied the plastomes of

various species of Orobanchaceae and found that the plastomes of orobanchaceous parasites

obviously experienced large-scale gene loss, especially the plastomes of holoparasites, in which

most of the photosynthesis-related genes are missing. Gene loss was detected even in the

orobanchaceous hemiparasites, such as the *ndh* genes (17). The degree of gene loss is

generally considered to be positively associated with the degree of parasitism (16), as a greater

degree of parasitism requires higher dependency on the host and fewer physiological processes

are needed by the parasite itself, and gene loss may enable the parasite to synchronize its own

physiology with that of the host, probably resulting in better fitness (18).

Thus far, four orobanchaceous species' genomes have been released, the autotrophic

Rehmannia glutinosa (19) and the obligate hemiparasite *Striga asiatica* (15) and facultative

hemiparasite *Phtheirospermum japonicum* (20) and *Pedicularis cranolopha* (21), but the genomes

of orobanchaceous holoparasites are still lacking. In this study, we sequenced and assembled

high-quality genomes of the autotroph *Lindenbergia luchunensis* and the holoparasites

Orobanche cumana and *Phelipanche aegyptiaca*, among which the latter two are notorious

parasitic pests in agriculture. Detailed comparative genomic analyses were performed using

these new genome data and the previously published genomes and transcriptomes. We

reconstructed a robust phylogeny and analyzed the trajectory of the genome evolution in

Orobanchaceae. Furthermore, our analysis on orobanchaceous genomes revealed the genomic

basis for parasitism evolution. This study fills the gap in our understanding of parasitism evolution

and provides new insight into the physiology and ecology of these parasites.

Results

Sequencing, assembly, and annotation of the genomes of *Lindenbergia luchunensis*, *Orobanche cumana*, and *Phelipanche aegyptiaca*

On a Nanopore platform, 36.47Gb data were generated for *L. luchunensis* (Llu), and on a PacBio Sequel II platform, 158.30 Gb and 396.00 Gb data were generated respectively for *O. cumana* (Ocu) and *P. aegyptiaca* (Pae). Using an MGI-SEQ 2000 platform, short reads (38.19 Gb, 141.20 Gb, and 329.33 Gb, respectively) were produced for Llu, Ocu, and Pae. Genome survey analysis using the short reads indicated low heterozygosities (SI Appendix, Table S1). Long reads were thus directly used for *de novo* assembling the genomes, and thereafter the short reads were used to correct the genome assemblies, forming contigs. The total length of the genome assembly of Pae is 3876.90 Mb (N50 9.97 Mb); using the Hi-C data, we obtained chromosome-level Llu and Ocu genome (Llu: 20 pseudochromosomes, 212.21 Mb; Ocu: 19 pseudochromosomes, 1417.94 Mb) (SI Appendix, Table S1). The accuracy of these three genomes all exceeded 99.9% (SI Appendix, Table S1). Thereafter, we first annotated repetitive sequences, including simple repeats, transposable elements, and RNA genes. In the genomes of Llu, Ocu, and Pae, the repetitive sequences respectively occupied 30.72%, 79.82%, and 84.40% (SI Appendix, Table S2). Mapping the RNA-seq data (SI Appendix, Table S3) from different organs to the respective genomes resulted in 29,669, 42,525, and 50,484 protein-coding genes in Llu, Ocu, and Pae. Benchmarking universal single-copy orthologs (BUSCO) analysis indicated that the completeness of these three genomes is 97.1% (Llu), 80.5% (Ocu), and 76.3% (Pae) (SI Appendix, Table S4). The low completeness values of the genes in Ocu and Pae genome is due to the large-scale gene loss in Ocu and Pae (see below for analysis), which is associated with their holoparasitic lifestyle.

Phylogenetic relationships in Orobanchaceae based on nuclear genes

Previously, it was established that there are eight major clades in Orobanchaceae; however, the phylogenetic relationships among the clades were still uncertain (5, 7, 22). The nuclear genes from all available genomes/transcriptomes of all the 19 Orobanchaceae species (SI Appendix, Table S5), covering seven out of the eight clades of Orobanchaceae, and three outgroup species (SI Appendix, Table S5) were used for the phylogenetic analysis pipeline (SI Appendix, Supplementary Methodology). Finally, after filtering out the orthogroups (OGs) which had low quality and/or high copy numbers, 907 OGs were selected for phylogenetic reconstruction. Coalescent and concatenation method yielded identical topologies with 100% support for all the nodes (Fig. 2). In this phylogenetic tree, consistent with previous findings (5, 7, 22, 23), the autotrophic lineages Rehmannieae and Lindenbergieae are the basal clades, and Lindenbergieae is the sister clade of all the parasitic plant lineages (Fig. 2). Within the group of parasitic plants,

Cymbarieae is the sister of all the rest lineages, a result different from Li et al. (5), which was

based on a few nuclear genes. In the rest four clades, Buchnereae and Orobancheae are sisters,

and so are Pedicularideae and Rhinantheae (Fig. 2); the topology of these four clades is distinct

from all previous studies (5, 7, 22, 23), except that it is similar to the topologies based on two

matK and *rps2* (7). Previously, several studies indicated that parasitic plants have higher

nucleotide substitution rates than typical autotrophic plants (24, 25); however, in our phylogenetic

tree, only the branch length of the holoparasitic plant *Aeginetia indica* in the clade Buchnereae is

exceptionally long, the branches of all holoparasitic plants in the clade *Orobancheae* were not

much longer than those of the autotrophic plants (SI Appendix, Fig. S1). We calculated the K_s

values of all the lineages in Orobanchaceae, but indeed no branches showed high substitution

rates (SI Appendix, Fig. S2).

Based on this tree, we estimated the divergence times for all the nodes using a Bayesian

molecular dating analysis. The estimated origin time of Orobanchaceae is about 46.51 million

years ago (Mya) (95% highest posterior density (HPD), 37.91-55.39 Mya), and the time

parasitism appeared was around 38.58 Mya (95% HPD, 31.27-45.81 Mya). Holoparasites

evolved independently at least three times in Orobanchaceae (Fig. 2 and (26)), among which,

Orobancheae appeared the earliest at ca. 33.95 Mya (95% HPD, 27.41-40.43 Mya).

Genome evolution in Orobanchaceae

Genome sizes and chromosome numbers vary in Orobanchaceae, and they seem to be positively

associated with the degrees of parasitism (4, 27). It is seemingly plausible that multiple rounds of

WGD events occurred during the evolution of the Orobanchaceae species (4, 27). To reveal the

WGD events in Orobanchaceae comprehensively, six species with reference genomes (Pae,

Ocu, and Llu from this study; *Striga asiatica* (Sas), *Phtheirospermum japonicum* (Pja), and

Pedicularis cranolopha (Pcr) from published/released data) and four outgroup species (*Mimulus*

guttatus (Mgu; Lamiales), *Solanum lycopersicum* (Solanales, lamiids), *Coffea canephora*

(Gentianales, lamiids), and *Arabidopsis thaliana* (Brassicales, rosids) were chosen for colinear

fragment comparisons. Instead of the commonly used methods based on colinear fragments

obtained by homology search, an analysis based on "orthologous colinear fragments (OCFs)"

was specifically designed for this study (see SI Appendix, Supplementary Methodology), which

enabled us to investigate the most recent WGD events in Orobanchaceae.

Our analysis based on OCFs from all species pairs (Fig. 3A) indicated that the Sas lineage

experienced an independent WGD event (named α^B event) (Fig. 3A), and this was consistent with

a previous study (15, 28). However, no WGD events were detected in any other orobanchaceous

lineages, including those containing Pae and Ocu (Fig. 3A). The ancestor of Orobanchaceae did

not experience any independent WGD events, even though there was a WGD event during the

evolution of the common ancestor of Orobanchaceae and *Mimulus*, which is named the β^L event

(Fig. 3A). Notably, the detected WGD events in the outgroup species are consistent with the

previous reports (29–31), supporting the reliability of our novel WGD analysis pipeline. Next, we

selected four nodes (S1-S4 in Fig. 3A) within the Lamiales clade, and all the orthologous genes

originated from these speciation events were selected for calculating the K_s values (SI Appendix,

Fig. S3). We found that the peak values of the K_s distributions had a linear relationship with the

divergence times of these nodes (R^2 = 0.9926, Fig. 3A). Therefore, the K_s values from the

paralogous genes resulted from the α^B and β^L events were used to infer the times of these two

events: α^B and β^L occurred at 31.00 ± 5.39 and 79.29 ± 5.39 Mya, respectively (Fig. 3A). The α^B

event likely occurred in Buchnereae soon after it diverged from Orobancheae at 35.78 Mya (95%

HPD, 28.97-42.52 Mya) (Fig. 2, 3A). The occurrence of β^L event happened to be close to the

Cretaceous-Paleocene boundary, when the well-known mass extinction event occurred (32) and

many plant lineages experienced WGD events (33, 34). We speculated that the β^L event may

have played an important role in enabling the common ancestor of Mgu and Orobanchaceae to

survive the extreme environmental changes.

Chromosome numbers is important for understanding the genome evolution of eukaryotes. The

chromosome numbers vary in Orobanchaceae (4, 27). Based on the chromosome-level genome

data of Ocu, Pcr, Llu and the outgroup species Mgu, we obtained 18 protochromosomes by

reconstructing the ancestral karyotype of Orobanchaceae. Analysis of the syntenic chromosomal

fragment indicated that Llu and Ocu exhibited good collinearity with the ancestral

protochromosomes (Fig. 3B, SI Appendix, Fig. S4). For example, the Chromosome 9 of Ocu and

the Chromosome 5 of Llu appeared to be very similar to the ancestral Chromosome 8. However,

Pcr has only eight chromosomes and its chromosomes experienced at least 11 fusion events.

Orobanchaceous species have variable genome sizes, ranging from 220 Mb to 8.7 Gb, and

parasites with greater degrees of parasitism generally have larger genome sizes (5). However,

our analysis suggests that most lineages in Orobanchaceae did not experience WGD events,

ruling out the possibility that the large genomes were resulted from ploidization. To gain insight

into the driving force of genome expansions, we compared the genome sizes of six

orobanchaceous species (Llu, Pja, Pcr, Sas, Ocu, and Pae) and the respective repetitive

elements (REs) in their genomes (Fig. 4A). Indeed, the large genome sizes were mainly resulted

from increased numbers of REs: in Ocu, Pja, and Pcr, which are about 1 Gb, 64-79% of their

genomes are REs; Pae has the largest genome (3.877 Gb), and 84% of the Pae gnome are REs,

among which the Long Terminal Repeats (LTRs) family is the most abundant. Notably, analysis

of the sequences of the LTRs in Llu, Pcr, Ocu, and Pae suggested that different subfamilies of

LTRs caused genome expansions in different species, especially specific LTR subfamilies
expanded to extreme extents very recently in Pae and Ocu (SI Appendix, Fig. S5). To determine
whether the massively increased numbers of LTRs could affect the coding regions in these
species, we compared the intron sizes in Pae and Llu, which have the largest and smallest
genomes in the orobanchaceous species of this study. We found that most introns retained their
lengths, regardless of genome sizes (Fig. 4B). Moreover, the contents of REs in the flanking
regions of genes became larger as the genome size increased, suggesting a possibility that the
REs may influence the expression of neighboring genes (Fig. 4C). We also noticed that even
though no WGD events were involved, Pja, Pcr, Pae, and Ocu with large genomes have many
more genes than does Llu, and this was mainly due to tandem or segmental duplication (30.96%-
56.33%), but not LTR-assisted "copy-paste" duplication (6.55%-17.40%) (SI Appendix, Table S6).
Given that the Chromosome 9 of Llu and Chromosome 5 of Ocu are homologous, we specifically
inspected the differences between these two chromosomes. Ocu's Chromosome 5 is eight times
larger than is Llu's Chromosome 9 (82 Mb vs 10 Mb). We found that many Class I
retrotransposons, including LTRs, in the proximal region of the Chromosome 5 of Ocu, where
gene densities are relatively low, supporting that expansion of LTRs is the main driving force of
increased genome sizes of orobanchaceous parasites (Fig. 4D). This finding is consistent with a
previous conclusion obtained by genome skimming (35).
226

Evolution of genes and gene families in Orobanchaceae
We analyzed all the gene families in these orobanchaceous species (Pae, Ocu, Sas, Pja, Pcr,
and Llu), *Cuscuta australis* (Cau), and their relatives (for orobanchaceous species: Mgu and
Sesamum indicum; for Cau: *Ipomoea nil* and *Solanum lycopersicum*), and two outgroup species,
Coffea canephora and *Arabidopsis thaliana* (Dataset S1). We found that, like Cau, all parasites of
Orobanchaceae experienced frequently gene family contractions; however, many gene families in
orobanchaceous parasites exhibited even more severe expansions, which is rare in Cau (Fig. 5).
Based on the phylogenetic tree (Fig. 2), we calculated the numbers of gene families that
underwent significant expansions and contractions across all branches (Fig. 5). Contractions
were frequent when parasitism and obligate parasitism first appeared in Orobanchaceae, and
thereafter, most lineages appeared to have independent gene family contractions. Most of the
gene family expansions happened independently in different species. Among the expanded gene
families, genes related to transcriptional regulation, transport, cell wall formation, phloem/xylem
histogenesis, and lateral root formation were enriched (Dataset S2). These gene may be involved
in the evolution of haustorium. Furthermore, we screened the transcriptomic data from Pja, Sas,
Ocu, and Pae for genes with high expression levels in haustoria (expression level of a given gene
is at least 1-fold greater in haustorium tissues than in other tissues), and found that at least
69.06% of these highly expressed genes in haustoria evolved through duplication (Fig. 6, SI

Appendix, Table S7). Next, these duplicated genes were used to determine the history of the

duplication events. Many of the duplication events happened in Mgu and Orobanchaceae's

MRCA, and according to our colinear fragment comparisons, 98.73% (545/552) of the duplication

events were caused by the β^L WGD event; the rest of the duplication events, most occurred

during the specific evolution of Pja, Sas, and Pae, but not in the common ancestor of

orobanchaceous parasites (Fig. 6, SI Appendix, Table S7).

Gene loss is common in parasitic plants (14, 16). We specifically split all the gene families into

OGs, and these OGs were thereafter used for gene loss analysis. Gene family contractions were

not additionally analyzed in other ways, as comparison of gene loss provides more detailed

information on the biological significance of the missing genes. A bioinformatic pipeline was also

employed to screen for false positives generated by errors from gene annotation and

orthogrouping (SI Appendix, Supplementary Methodology). As a comparison, Cau was included

in the gene loss analysis with five parasitic Orobanchaceae species, as *Cuscuta* and

Orobanchaceae are both Lamiids. *Sapria himalayana* (Rafflesiaceae) (16) was not included in our

analysis as it is phylogenetically very distant from Lamiids and is an endoparasite. We found that

the severity of gene loss is positively associated with the degree of parasitism: the facultative

hemiparasites Pja and Pcr respectively lost 2.41% and 2.95% of the conserved genes, the

obligate hemiparasite Sas lost 5.71%, and the obligate holoparasites Ocu, Pae, and Cau

respectively lost 13.11%, 13.85%, and 14.88% (Fig. 7A and B). The history of gene loss events in

orobanchaceous parasites was estimated: the common ancestor of all orobanchaceous parasites

lost only 74 genes, and 191 and 777 genes were lost respectively during the evolution of the

ancestors of obligate parasites and holoparasites (Fig. 7A). Notably, even though hundreds of

genes were lost during the species-specific evolution of these parasites (Fig. 7A), many genes

were convergently lost (Fig. 7A): by comparing the lost genes in Pedicularideae (Pja and Pcr) and

Orobancheae (Ocu and Pae), we found that 74-122 genes were convergently lost, which was well

supported by the hypergeometric test (Fig. 7A). Furthermore, in the closely related holoparasite

Ocu and Pae, 33.6% and 29.9% of the lost genes are species-specific, and selection analysis

indicated that significant portions of these Ocu- and Pae-retained genes, which had been

specifically lost in the other species, are under relaxed selection (SI Appendix, Fig. S6). It is

possible that Ocu and Pae are still experiencing genes loss, and these genes will be convergently

lost in the future due to relaxed selection.

We next compared the lost genes in the orobanchaceous parasites and Cau and it was found that

among the 1687 lost genes in Cau, more than half (889 genes, 57.14%) were also lost in Pae and

Ocu, and 75 genes were also lost in the hemiparasites Sas, Pja, and Pcr (Fig. 7B). It is evident

that all orobanchaceous parasites and Cau convergently lost various genes, suggesting that

some of these gene loss events could contribute to their adaptation to the parasitic lifestyle. To

gain further insight into the functions of the lost genes in Cau, Pae, and Ocu, which are all

holoparasites, we first performed gene ontology (GO) enrichment analysis on all their lost genes,

and the GO terms were next used for interspecies pairwise comparisons. The resulting common

and unique GO terms were further categorized by a network analysis (Fig. 7C). Pae and Ocu-lost

genes exhibited very similar GO terms, while the enriched GO terms from the Cau lost genes

appeared to be quite different from those of Pae or Ocu. In Cau, most of the lost genes' GO terms

are related to ion transport in root, ethylene biosynthesis, and response to nutrient and oxygen

levels, and this is consistent with the rootless morphology of Cau (Fig. 7C). The majority of the

lost genes in Pae and Ocu are involved in photosynthesis, chloroplast organization, and

chlorophyll biosynthesis; although Cau also lost genes related to these processes, the numbers

of GO terms under these processes are remarkably fewer than those from Pae or Ocu (Fig. 7C).

The regulation of flowering in some parasitic plants, especially in holoparasites, is considered to

be unique (18). We specifically inspected the flowering regulatory genes in Cau and all the

parasites in Orobanchaceae (Fig. 7D). Many important flowering regulatory genes are missing in

the holoparasites Pae, Ocu, and Cau, including *CO*, *SVP*, *ELF4*, *AGL19* and *AGL16*, and *CDF1*

and *CDF3* (Fig. 7D). In terms of the regulatory pathways of flowering, Cau lost more genes

related to the autonomous and hormone pathway than did Pae and Ocu, while Pae and Ocu lost

more genes involved in the circadian and photoperiod pathway than did Cau. We suppose that

this was due to the fact that Cau uses host plant's flowering signal FT to activate its own flowering

(18) and Pae and Ocu likely do not rely on light signaling to flower, as they spend a significant

amount of time living underground.

The evolution of KAI2 gene family

After perceiving the strigolactones (SLs) released from adjacent host roots, seeds of obligate

parasites in Orobanchaceae germinate (4, 9, 36). In normal autotrophic plants, D14, an

alpha/beta hydrolase protein, perceives SLs and regulates physiological processes of

development and stress resistance (37). In *Striga*, however, the receptor of host-released SLs

has been identified to be encoded by *KAI2d*, and importantly, *KAI2d* did not evolve from *D14*, but

from the *KAI2* gene, which encodes the receptor of karrikin, through tandem duplication, large

gene family expansion, and neofunctionalization (11, 15).

We analyzed the *KAI2* gene families in Orobanchaceae and a few autotrophic species from

Lamiids, and Arabidopsis was used as the outgroup species (Fig. 8A). Only a single *KAI2* was

identified in Arabidopsis, whereas in *Coffea canephora*, *Solanum lycopersicum*, and

Orobanchaceae at least two copies exist, suggesting that *KAI2* duplicated in the ancestor of

Lamiids resulting in the two paralogs, *KAI2c* and *KAI2i* (Fig. 8A). We specifically inspected the

KAI2 gene-containing chromosomal fragments in Llu and *Solanum lycopersicum*, and found that

in both species, *KAI2c* and *KAI2i* are either on different chromosomes (Llu) or distantly located on

the same chromosome (~ 20 Mb in *Solanum lycopersicum*). Collinearity analysis indicated that

the two *KAI2* gene-containing chromosomal fragments are obviously homologous with the

fragment containing *KAI2* gene in Arabidopsis (Fig. 8B). Given that no WGD events were

detected in the ancestor of Lamiids ((30) and this study, Fig. 3), the duplication of *KAI2* in Lamiids

is a result of segmental duplication. Thereafter, *KAI2d* is derived from duplication of *KAI2i* (Fig.

8A); given that in Pja, Pcr, and Sas, *KAI2d* and *KAI2i* are adjacently located (Fig. 8B), *KAI2i* gave

rise to *KAI2d* by tandem duplication. In Pja and Pcr, the chromosome fragments containing *KAI2i*

and *KAI2d* show good collinearity with the chromosome fragment containing *KAI2i* in Llu, while in

Sas, Ocu, and Pae, the collinearity was not detected (Fig. 8B). Thus, we inferred that this tandem

duplication event, in which *KAI2d* was born, happened in the common ancestor of

orobanchaceous parasitic plants, and in Pja and Pcr the locations of these duplicated genes

remained, whereas in Sas, Pae, and Ocu both genes translocated to other loci (Fig. 8B).

Discussion

Orobanchaceae is the only family of plants with all types of parasitic lifestyles, and it is preferred

for studying the origin and evolution of parasitism. The phylogeny of Orobanchaceae has been

studied using plastome fragments (23, 38), ITS (7, 23), and individual nuclear genes (5, 7). Even

though most of these studies identified eight monophyly clades, the phylogenetic relationship

among these clades was not completely resolved. In this study, we assembled three genomes of

orobanchaceous plants, Llu, Ocu, and Pae. Together with the published/released transcriptome

and genome data of other orobanchaceous plants, this study covered seven out of eight major

clades of Orobanchaceae. Using 907 nuclear OGs, we reconstructed a highly resolved and well-

supported phylogeny of Orobanchaceae, and the divergence times were estimated using a

Bayesian molecular dating approach (Fig. 2). Based on the phylogeny, we propose that 1) the

common ancestor of Orobanchaceae, an autotrophic plant, evolved parasitism (haustorium) at

38.58 Mya (95% HPD, 31.27-45.81 Mya); 2) the Cymbarieae, which is composed of only

facultative hemiparasites, was the earliest diverged parasitic lineage with low biodiversity; 3) by

Oligocene (33.90-23.03 Mya), a facultative hemiparasitic ancestor rapidly evolved into two

lineages; one diverged into Buchnereae (facultative hemiparasitic, obligate hemiparasitic, and

holoparasitic species) and Orobancheae (holoparasites), while the other evolved to be

Pedicularideae (facultative hemiparasitic species) and Rhinantheae (facultative hemiparasitic and

holoparasitic); these four clades have diverse species. Grasslands rapidly expanded during

Oligocene (39), and this event may have provided orobanchaceous parasites with opportunities

to exploit grasses and herbs, enabling the expansion of these parasites (7, 40). Notably, different

from a recent study (5), in our scenario, hemiparasites evolved earlier than did holoparasites, and

this is consistent with the progressive evolution of increasing dependency of hosts.

Most species in Orobanchaceae have 8 to 20 pairs of chromosomes, but the chromosome

number can reach 72 in some species (4, 27). Based on the chromosome number variation, Lyko

and Wicke (27) proposed at least three ancient WGD events during the evolution of

Orobanchaceae, including one WGD event in *Striga* and another WGD event in the Orobancheae

clade. The WGD event in *Striga* was confirmed by the recent genome analysis (15). However, our

collinearity analysis of the genome data of Llu, Pga, Pcr, Sas, Ocu, and Pae did not find other

WGD events in Orobanchaceae, except a single WGD event (α^B) that occurred during the

evolution of the ancestor of Buchnereae (31.00 ± 5.39 Mya) (Fig. 3A), a clade where Sas resides.

Moreover, the 1:1 collinearity of the Llu and Ocu's chromosomes (SI Appendix, Fig. S4) also

strongly supports that no ploidization happened after the divergence between Llu and Ocu. Even

though no WGD events were detected at the times of the emergence of parasitism and

holoparasitism, there was a very ancient WGD event (β^L, 79.29 ± 5.39 Mya) in the common

ancestor of Orobanchaceae and *Mimulus* (Fig. 3A). In this study, we reconstructed the 18

protochromosomes of Orobanchaceae ancestor using the genomes of species from *Orobanche*,

Pedicularis, and *Lindenbergia* and the outgroup *Mimulus*. The chromosome numbers and

collinearity are well retained in *Lindenbergia* and *Orobanche*; however, the chromosome-level

genome of Pcr indicates that Pcr has only eight chromosomes (Fig. 3B). Thus, Pcr must have

experienced large-scale chromosome recombination and fusion events. *Striga* has only 12

chromosomes (4), even though WGD happened during the evolution of *Striga*. We inferred that

Striga also experienced many chromosome recombinations and fusions.

The dodder Cau and the endoparasite *Sapria himalayana* are both non-orobanchaceous species.

Cau has only 19671 genes (14), and it is among the higher plants with the least number of

protein-coding genes. *Sapria himalayana* genome harbors 55179 genes; however, most of them

are transposon-like sequences, leaving only about 10000 low-copy genes (16). By contrast,

orobanchaceous parasites have many more genes than do Cau and *Sapria himalayana* (Fig. 5),

and this was due to species-specific gene family expansions in different orobanchaceous

parasites (Fig. 5). Furthermore, among these expanded gene families, genes likely related to

haustorium development and function were identified and many of the genes highly expressed in

haustoria were evolved from lineage-specific duplications (Fig. 6). We propose that species-

specifically expanded gene families may be the driving force underlying the adaptation of the

parasites to different hosts.

Similar to Cau and *Sapria himalayana*, large-scale gene loss was also detected in

orobanchaceous parasites, especially in the holoparasites Ocu and Pae. The reconstructed gene-

loss history indicated that only 74 genes were missing in the common ancestor of

orobanchaceous parasites, when parasitism evolved; by contrast, 777 genes were deleted from

the genome of the MRCA of Ocu and Pae. Thus, gene loss became more severe as the degree

of parasitism increased. Considering our data of gene loss in Orobanchaceae and the previous

published data from Orchidaceae (41) and *Sapria himalayana* (16), in parasitic plants and

mycoheterotrophic plants, which are all heterotrophs, the degree of gene loss and parasitism

seems to be well positively correlated as well. Accordingly, heterotrophic plants could be divided

into four classes with increasing levels of heterotrophy and degrees of gene loss: in the first class,

heterotrophs with 2-3% gene loss are facultative hemiparasites; in the second class, heterotrophs

with 6-7% gene loss require hosts to complete certain developmental stages, and these are

obligate hemiparasites and initial mycoheterotrophic plants, such as most species of

Orchidaceae; in the third class, heterotrophs lost 13-15% genes, and these are holoparasites or

full mycoheterotrophs with vegetative organs, such as *Cuscuta*, *Orobanche*, and *Gastrodia*; in the

fourth class, gene loss is over 30% and these are endoparasites with only reproductive organs,

such as *Sapria himalayana*. The lost genes in the heterotrophs of the first and second class seem

to be rather random, while the genes lost in the heterotrophs of the third class are related to

photosynthesis, transcriptional regulation, stress adaptation, and organogenesis; the lost genes in

endoparasite *Sapria himalayana*, from the fourth class, includes almost all those lost in Pae, Ocu,

Cau, and *Gastrodia*, representing the extreme form of parasitism.

Importantly, comparisons among orobanchaceous species and between Orobanchaceae and

Cau indicated convergent evolution of gene loss, suggesting that abandonment of certain

physiological functions is positive selected in parasitic plants. In *Striga*, the *protein phosphatase*

2C gene is mutated, leading to abscisic acid (ABA) insensitivity, and the impaired ABA signaling

pathway led to very high transpiration in *Striga*, enabling this parasite to absorb nutrients from

host xylem sap, even under drought conditions (15, 42). The leafless *Cuscuta australis* lost

almost all the flowering regulatory pathways, and its *FT* gene, which encodes an important

flowering signal, had pseudogenized; however, Cau uses the FT proteins from host plants to

activate its own flowering, enabling it to synchronize its flowering times with those of the hosts

(18). In this manner, Cau can parasitize a large number of host species with varying flowering

times, and thus, losing the flowering genes is an elegant strategy of adaptation in Cau (18). It

would be interesting to study what pathways are affected by gene loss in orobanchaceous

parasites and importantly the physiological and ecological impact incurred by gene loss.

Searcy and MacInnis (1) proposed a model of parasitic plant evolution: first, development of
haustorium which enables physical connections with host plants; after haustorium evolved, the
parasites start to lose genes that are no longer needed by the parasites and parasites evolve to
adapt to their specific hosts. The evolution of orobanchaceous parasites well supports this model.
The β^L WGD event in the common ancestor of Orobanchaceae and *Mimulus* generated many
duplicated genes which likely contributed to the evolutionary gain of a haustorium; thereafter,
gene loss events occurred in orobanchaceous parasites, especially in holoparasites, and in
parallel, expansions in specific gene families may have contributed to specialization and
adaptation of the parasites to the hosts.
438
In this study, our phylogenomic analysis constructed a robust phylogenetic framework of the
major lineages of Orobanchaceae. Furthermore, genome evolution and comparative genomics
analysis shed new light on the origin and evolution of parasitism in Orobanchaceae. These
genome resources and further molecular, genetic, and genomic studies on Orobanchaceae
species will provide many interesting insights into the evolution underlying the drastic
physiological and morphological changes in these fascinating plants.
445
446   **Materials and Methods**
447
Genome sequencing
Leaves of Llu and aboveground parts of stems of Ocu and Pae were harvested and cleaned with
tap water and used for DNA extraction. For short-read sequencing: DNA libraries were
constructed using a VAHTS Universal Plus DNA Library Prep Kit for MGI (Vazyme, Nanjing,
China) following the manufacturer's instructions, and the libraries were sequenced on an MGI-
SEQ 2000 (MGI Tech, Shenzhen, China) to generate 150-bp paired-end data. For long-read
sequencing: for Llu, we used the PromethION platform from Oxford Nanopore Technologies; for
Ocu and Pae, we used the PacBio Sequel II platform.
456
457   Tissues of Llu and Ocu were used for the Hi-C pipeline following a previously described method
458   (43), including crosslinking, chromatin digestion with MboI (New England BioLabs, MA), labeling
459   of DNA ends, ligation, purification, shearing, and biotin pull-down. The Hi-C libraries were
460   sequenced on an MGI-SEQ 2000 (MGI Tech, Shenzhen, China) platform in the 150-bp paired-
461   end mode.
462
To aid genome annotation, we obtained RNA-seq data from different tissues of orobanchaceous
plants (SI Appendix, Table S3). The total RNA was extracted from these tissues using the TRI
Reagent (Sigma). The MGIEasy RNA library preparation kit was used for library construction.

From a DNBSEQ-T7 platform (MGI Tech, Shenzhen, China), 150bp pair-end sequencing reads
were obtained.

Genome assembly

We first surveyed the genome of Llu, Ocu, and Pae based on k-mer (k=17) depth frequency
distribution analysis using the GCE software (v1.0.0) (44) with short-read data, then obtained the
genome size, repeat content, and heterozygosity for these species (SI Appendix, Table S1).

For Ocu and Pae, the subreads from the Pacbio sequencing were corrected, trimmed, and
assembled by CANU (v2.2) (45). Contigs were polished with the PacBio data using GCpp (v1.9.0)
(https://github.com/PacificBiosciences/gcpp) and then corrected with short-reads data using Pilon
(v1.23) (46). For Llu, the long reads of ONT were de novo assembled using NextDenovo (v2.3.1)
(https://github.com/Nextomics/NextDenovo), and the obtained contigs were corrected with
NextPolish (v1.1.0) (47) using both short reads and long reads.

To obtain chromosome-scale assemblies for Llu and Ocu, we mapped the Hi-C data to contigs
and constructed the Hi-C contact matrices, which were used to correct the assembly of contigs
and build scaffolds using the 3D-DNA pipeline (v180922) (48).

In order to evaluate the quality of the assembly, short reads were mapped to the assemblies
using HISAT2 (49), and the results from mapping were used for calling SNPs and indels using
SAMtools (v1.9) (50) and BCFtools (v1.9) (51). The homozygous SNPs and indels were
considered as assembly error, and thus the correct rates of assembly were calculated.

Genome annotation

We first used the EDTA pipeline (v1.8.3) (52) to annotate repetitive sequences in the genomes,
which integrates with the LTR_FINDER (53), LTRharvest (54), LTR_retriever (55), Generic
Repeat Finder (56), TIR-learner (57), HelitronScanner (58), and TEsorter (59). EDTA first used
these software to de novo annotate genome-wide transposons, and then the resulting
transposons were integrated with the Dfam database (https://dfam.org/) into RepeatMasker
(http://repeatmasker.org) to annotate repetitive elements.

To annotate the protein-coding genes, four steps were used to guide the annotation. 1) Augustus
(v3.4.0) (60) for gene model prediction was trained with the MAKER-P pipeline (61) using the
RNA-seq data and the protein-coding genes in the close species (*Striga asiatica*, *Mimulus*
guttata, *Arabidopsis thaliana*, *Solanum lycopersicum*, *Coffea canephora*, and *Cuscuta australis*).

After three rounds of training of Augustus, de novo annotation of protein-coding genes was done.
2) The RNA-seq data were mapped to unmasked genomes using TopHat2 (v2.1.1) (62), and the
mapped reads were assembled to gene structures by Cufflinks (v2.2.1) (63); TRINITY (v2.8.5)
(64) was used to de novo assemble the RNA-seq reads and thereafter PASA2 (v2.4.1) (65) was
used to map the assembled contigs to the genomes and annotate the gene structures. These
annotation results were merged. 3) The protein-coding genes in the closely related species were
aligned to the unmasked genomes using GenBlastA (v1.0.1) (66), and GeneWise2 (v2.4.1) (67)
was employed to generate gene structures. 4) The gene structures obtained from the above three
steps were combined by EVidenceModeler (v1.1.1) (68). Then, the gene annotation results were
compared with the repetitive element annotation results to remove the genes that overlap with
repetitive elements. The final annotations were achieved.
514
515 See SI Appendix, Supplementary Methodology for phylogenetic analysis and comparative
516 genomics analysis.
517

518 **Data availability**
519 The genome assembly, gene models, and sequence reads of Llu, Ocu, and Pae are available at
520 the National Genomics Data Center (https://ngdc.cncb.ac.cn/) under BioProject number
521 PRJCA008847. All data are also available from the corresponding author upon request.
522

523 **Acknowledgments**
524
We thank Dr. Xiaopeng Yun (Inner Mongolia Academy of Agricultural and Animal Husbandry
Sciences) for providing Ocu and Pae seeds, and the Germplasm Bank of Wild Species in
Southwest China for providing Llu seeds. Drs. Jun Zhao and Jian Zhang (Inner Mongolia
Agricultural University) are thanked for identification of Ocu. We are grateful to the High-
Performance Computing Facility of Department of Economic Plants and Biotechnology at
Kunming Institute of Botany for providing computing services. This study was supported the
National Natural Science Foundation of China (No. 32000179, Y.X.), Strategic Priority Research
Program of Chinese Academy of Sciences (No. XDPB16, J.W.), CAS "Light of West China"
Program (G.S.), the Special Research Assistant of Chinese Academy of Sciences (Y.X. and J.Z.),
and the Postdoctoral Directional Training Foundation of Yunnan Province (Y.X.).

References
537

1. D. G. Searcy, A. J. MacInnis, Measurements by DNA renaturation of the genetic basis of
parasitic reduction. *Evolution* **24**, 796–806 (1970).

2.   D. L. Nickrent, Parasitic angiosperms: How often and how many? *TAXON* **69**, 5–27 (2020).

3.   J. H. Westwood, J. I. Yoder, M. P. Timko, C. W. dePamphilis, The evolution of parasitism in
plants. *Trends in Plant Science* **15**, 227–235 (2010).

4.   D. M. Joel, J. Gressel, L. J. Musselman, Eds., *Parasitic Orobanchaceae* (Springer Berlin
Heidelberg, 2013) https:/doi.org/10.1007/978-3-642-38146-1 (October 17, 2021).

5.   X. Li, T. Feng, C. Randle, G. M. Schneeweiss, Phylogenetic relationships in Orobanchaceae
inferred from low-copy cuclear genes: Consolidation of major clades and identification of a
novel position of the non-photosynthetic orobanche clade sister to all other parasitic
Orobanchaceae. *Front. Plant Sci.* **10**, 902 (2019).

6.   A. D. Twyford, Parasitic plants. *Current Biology* **28**, R857–R859 (2018).

7.   J. R. McNeal, J. R. Bennett, A. D. Wolfe, S. Mathews, Phylogeny and origins of
holoparasitism in Orobanchaceae. *American Journal of Botany* **100**, 971–983 (2013).

8.   S. Yoshida, S. Cui, Y. Ichihashi, K. Shirasu, The haustorium, a specialized invasive organ in
parasitic plants. *Annu. Rev. Plant Biol.* **67**, 643–667 (2016).

9.   C. E. Cook, L. P. Whichard, B. Turner, M. E. Wall, G. H. Egley, Germination of witchweed
(Striga lutea Lour.): Isolation and properties of a potent stimulant. *Science* **154**, 1189–1190
(1966).

10.   C. E. Cook, *et al.*, Germination stimulants. II. Structure of strigol, a potent seed germination
stimulant for witchweed (Striga lutea). *J. Am. Chem. Soc.* **94**, 6198–6199 (1972).

11.   C. E. Conn, *et al.*, Convergent evolution of strigolactone perception enabled host detection in
parasitic plants. *Science* **349**, 540–543 (2015).

12.   S. Toh, *et al.*, Structure-function analysis identifies highly sensitive strigolactone receptors in
Striga. *Science* **350**, 203–207 (2015).

13.   Y. Tsuchiya, *et al.*, PARASITIC PLANTS. Probing strigolactone receptors in Striga
hermonthica with fluorescence. *Science* **349**, 864–868 (2015).

14.   G. Sun, *et al.*, Large-scale gene losses underlie the genome evolution of parasitic plant
Cuscuta australis. *Nat Commun* **9**, 2683 (2018).

15.   S. Yoshida, *et al.*, Genome sequence of Striga asiatica provides insight into the evolution of
plant parasitism. *Curr Biol* **29**, 3041-3052.e4 (2019).

16.   L. Cai, *et al.*, Deeply altered genome architecture in the endoparasitic flowering plant Sapria
himalayana Griff. (Rafflesiaceae). *Current Biology* **31**, 1002-1011.e9 (2021).

17.   S. Wicke, *et al.*, Mechanistic model of evolutionary rate variation en route to a
nonphotosynthetic lifestyle in plants. *PNAS* **113**, 9045–9050 (2016).

18.   G. Shen, *et al.*, *Cuscuta australis* (dodder) parasite eavesdrops on the host plants' FT
signals to flower. *Proc Natl Acad Sci USA* **117**, 23125–23130 (2020).

19. L. Ma, *et al.*, De novo genome assembly of the potent medicinal plant Rehmannia glutinosa
using nanopore technology. *Computational and Structural Biotechnology Journal* **19**, 3954–
3963 (2021).

20. S. Cui, *et al.*, Ethylene signaling mediates host invasion by parasitic plants. *Science*
Advances (2020) https:/doi.org/10.1126/sciadv.abc2385 (January 4, 2022).

21. J. Jin, Pedicularis cranolopha genome reference project.
https://www.ncbi.nlm.nih.gov/bioproject/805569

22. G. M. Schneeweiss, "Phylogenetic relationships and evolutionary trends in Orobanchaceae"
in *Parasitic Orobanchaceae*, D. M. Joel, J. Gressel, L. J. Musselman, Eds. (Springer Berlin
Heidelberg, 2013), pp. 243–265.

23. A. D. Wolfe, C. P. Randle, L. Liu, K. E. Steiner, Phylogeny and biogeography of
Orobanchaceae. *Folia Geobot* **40**, 115–134 (2005).

24. B. Lemaire, S. Huysmans, E. Smets, V. Merckx, Rate accelerations in nuclear 18S rDNA of
mycoheterotrophic and parasitic angiosperms. *J Plant Res* **124**, 561–576 (2011).

25. L. Bromham, P. F. Cowman, R. Lanfear, Parasitic plants have increased rates of molecular
evolution across all three genomes. *BMC Evolutionary Biology* **13**, 126 (2013).

26. W. Fu, *et al.*, Testing the hypothesis of multiple origins of holoparasitism in Orobanchaceae:
Phylogenetic evidence from the last two unplaced holoparasitic genera, Gleadovia and
Phacellanthus. *Frontiers in Plant Science* **8**, 1380 (2017).

27. P. Lyko, S. Wicke, Genomic reconfiguration in parasitic plants involves considerable gene
losses alongside global genome size inflation and gene births. *Plant Physiology* **186**, 1412–
1423 (2021).

28. N. J. Wickett, *et al.*, Transcriptomes of the parasitic plant family Orobanchaceae reveal
surprising conservation of chlorophyll synthesis. *Curr Biol* **21**, 2098–2104 (2011).

29. S. Sato, *et al.*, The tomato genome sequence provides insights into fleshy fruit evolution.
Nature **485**, 635–641 (2012).

30. F. Denoeud, *et al.*, The coffee genome provides insight into the convergent evolution of
caffeine biosynthesis. *Science* **345**, 1181–1184 (2014).

31. C. Simillion, K. Vandepoele, M. C. E. V. Montagu, M. Zabeau, Y. V. de Peer, The hidden
duplication past of Arabidopsis thaliana. *PNAS* **99**, 13627–13632 (2002).

32. P. Schulte, *et al.*, The Chicxulub asteroid impact and mass extinction at the Cretaceous-
Paleogene boundary. *Science* **327**, 1214–1218 (2010).

33. J. A. Fawcett, S. Maere, Y. V. de Peer, Plants with double genomes might have had a better
chance to survive the Cretaceous–Tertiary extinction event. *PNAS* **106**, 5737–5742 (2009).

34. S. Wu, B. Han, Y. Jiao, Genetic contribution of paleopolyploidy to adaptive evolution in
angiosperms. *Molecular Plant* **13**, 59–71 (2020).

35. M. Piednoël, *et al.*, Next-generation sequencing reveals the impact of repetitive DNA across
phylogenetically closely related genomes of Orobanchaceae. *Molecular Biology and*
Evolution **29**, 3601–3611 (2012).

36. T. Yokota, H. Sakai, K. Okuno, K. Yoneyama, Y. Takeuchi, Alectrol and orobanchol,
germination stimulants for Orobanche minor, from its host red clover. *Phytochemistry* **49**,
1967–1973 (1998).

37. M. T. Waters, C. Gutjahr, T. Bennett, D. C. Nelson, Strigolactone signaling and evolution.
Annu. Rev. Plant Biol. **68**, 291–322 (2017).

38. N. D. Young, K. E. Steiner, C. W. dePamphilis, The evolution of parasitism in
Scrophulariaceae/Orobanchaceae: Plastid gene sequences refute an evolutionary transition
series. *Annals of the Missouri Botanical Garden* **86**, 876 (1999).

39. T. H. Torsvik, L. R. M. Cocks, "Paleogene" in *Earth History and Palaeogeography*,
(Cambridge University Press, 2016), pp. 240–255.

40. O. Eriksson, K. Kainulainen, The evolutionary ecology of dust seeds. *Perspectives in Plant*
Ecology, Evolution and Systematics **13**, 73–87 (2011).

41. Y. Xu, *et al.*, A chromosome‐scale *Gastrodia elata* genome and large‐scale comparative
genomic analysis indicate convergent evolution by gene loss in mycoheterotrophic and
parasitic plants. *The Plant Journal* **108**, 1609–1623 (2021).

42. H. Fujioka, *et al.*, Aberrant protein phosphatase 2C leads to abscisic acid insensitivity and
high transpiration in parasitic Striga. *Nat. Plants* **5**, 258–262 (2019).

43. S. S. P. Rao, *et al.*, A 3D map of the human genome at kilobase resolution reveals principles
of chromatin looping. *Cell* **159**, 1665–1680 (2014).

44. B. Liu, *et al.*, Estimation of genomic characteristics by analyzing k-mer frequency in de novo
genome projects. *arXiv* (2020) (March 14, 2022).

45. S. Koren, *et al.*, Canu: scalable and accurate long-read assembly via adaptive *k*-mer
weighting and repeat separation. *Genome Res.* **27**, 722–736 (2017).

46. B. J. Walker, *et al.*, Pilon: an integrated tool for comprehensive microbial variant detection
and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

47. J. Hu, J. Fan, Z. Sun, S. Liu, NextPolish: a fast and efficient genome polishing tool for long-
read assembly. *Bioinformatics* **36**, 2253–2255 (2020).

48. O. Dudchenko, *et al.*, De novo assembly of the Aedes aegypti genome using Hi-C yields
chromosome-length scaffolds. *Science* **356**, 92–95 (2017).

49. D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome alignment
and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–915 (2019).

50. H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–
2079 (2009).

51. P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *GigaScience* **10**, giab008
(2021).

52. S. Ou, *et al.*, Benchmarking transposable element annotation methods for creation of a
streamlined, comprehensive pipeline. *Genome Biol* **20**, 275 (2019).

53. Z. Xu, H. Wang, LTR_FINDER: an efficient tool for the prediction of full-length LTR
retrotransposons. *Nucleic Acids Res* **35**, W265-268 (2007).

54. D. Ellinghaus, S. Kurtz, U. Willhoeft, LTRharvest, an efficient and flexible software for de
novo detection of LTR retrotransposons. *BMC Bioinformatics* **9**, 18 (2008).

55. S. Ou, N. Jiang, LTR_retriever: a highly accurate and sensitive program for identification of
Long Terminal Repeat retrotransposons. *Plant Physiol* **176**, 1410–1422 (2018).

56. J. Shi, C. Liang, Generic Repeat Finder: a high-sensitivity tool for genome-wide de novo
repeat detection. *Plant Physiol* **180**, 1803–1815 (2019).

57. W. Su, X. Gu, T. Peterson, TIR-Learner, a new ensemble method for TIR transposable
element annotation, provides evidence for abundant new transposable elements in the
maize genome. *Mol Plant* **12**, 447–460 (2019).

58. W. Xiong, L. He, J. Lai, H. K. Dooner, C. Du, HelitronScanner uncovers a large overlooked
cache of Helitron transposons in many plant genomes. *Proc Natl Acad Sci U S A* **111**,
10263–10268 (2014).

59. R.-G. Zhang, Z.-X. Wang, S. Ou, G.-Y. Li, TEsorter: lineage-level classification of
transposable elements using conserved protein domains. *bioRxiv*, 800177 (2019).

60. O. Keller, M. Kollmar, M. Stanke, S. Waack, A novel hybrid gene prediction method
employing protein multiple sequence alignments. *Bioinformatics* **27**, 757–763 (2011).

61. M. S. Campbell, *et al.*, MAKER-P: a tool kit for the rapid creation, management, and quality
control of plant genome annotations. *Plant Physiology* **164**, 513–524 (2014).

62. D. Kim, *et al.*, TopHat2: accurate alignment of transcriptomes in the presence of insertions,
deletions and gene fusions. *Genome Biology* **14**, R36 (2013).

63. C. Trapnell, *et al.*, Differential gene and transcript expression analysis of RNA-seq
experiments with TopHat and Cufflinks. *Nat Protoc* **7**, 562–578 (2012).

64. B. J. Haas, *et al.*, De novo transcript sequence reconstruction from RNA-seq using the
Trinity platform for reference generation and analysis. *Nat Protoc* **8**, 1494–1512 (2013).

65. B. J. Haas, *et al.*, Improving the Arabidopsis genome annotation using maximal transcript
alignment assemblies. *Nucleic Acids Res* **31**, 5654–5666 (2003).

66. R. She, J. S.-C. Chu, K. Wang, J. Pei, N. Chen, GenBlastA: enabling BLAST to identify
homologous gene sequences. *Genome Res* **19**, 143–149 (2009).

67. E. Birney, M. Clamp, R. Durbin, GeneWise and Genomewise. *Genome Res* **14**, 988–995
(2004).

68. B. J. Haas, *et al.*, Automated eukaryotic gene structure annotation using EVidenceModeler
and the Program to Assemble Spliced Alignments. *Genome Biol* **9**, R7 (2008).
685

Figures

Lindenbergia	***Pedicularis***	***Striga***	***Orobanche***
Autotrophic	Facultative Hemiparasitic	Obligate Hemiparasitic	Obligate Holoparasitic

Figure 1. Plant species in the family Orobanchaceae spanning from non-parasitic to increasing levels of parasitism.

Figure 2. Phylogenetic and molecular clock dating analysis of the major clades in Orobanchaceae.

The phylogenetic tree was constructed using low-copy OGs from 19 orobanchaceous species and three outgroup plants. The gray background indicates autotrophic outgroups; the light grayish-green background represents orobanchaceous autotrophic plant clades; the pale-yellow background represents orobanchaceous parasitic plants; the red branches indicate holoparasites. The red dashed line for *Lathraea* highlights that its position is from the work of Li et al. (5). The numbers near the internodes represent divergence times (Mya) based on molecular clocking dating, and the black bars at the internodes represent 95% HPD for divergence times. All internodes are 100% supported. Important evolutionary events are shown in the dashed boxes. The vertical bar on the left indicates five monophyletic clades (Cymbarieae, Orobancheae, Rhinantheae, Buchnereae, and Pedicularideae) and two autotrophic clades (Rehmannieae and Lindenbergieae, which are shown as "autotroph"). The bar on the bottom indicates the geologic time scale.

Figure 3. Genome evolution in Orobanchaceae.

(A) The palaeopolyploidization in Orobanchaceae and outgroup lineages. The right histogram matrix indicates the coverage depths of orthologous colinear fragments (OCFs) from pairwise species comparisons. The OCFs from a given species pair were respectively mapped to each species to create two histograms indicating the coverage depths of OCFs, and the max depths of OCFs mapped to species B can be used to infer the WGD events in species A after speciation between species A and B, and vice versa (see methods). In the matrix, each histogram's row and column indicate the two species used for creating the OCFs, and the species of the column is the

one that was the mapping-target species. The percentage in the upper right corner of each histogram represents the proportion of genes covered by OCF in the target species to the total genes. The WGD events inferred from the histogram matrix are marked with stars in the species tree on the left (blue for triplication, red for duplication), and the positions of the stars represent the times of WGD events, which were inferred from the peak value of K_s. The black circles on the species tree represent the four speciation events which were used to calculate the correlation between K_s values and divergence times (see SI Appendix, Supplementary Methodology). The correlation between the peak value of K_s and divergence times is shown in the upper left corner. (B) Eighteen ancestral Orobanchaceae protochromosomes were constructed using *Mimulus guttatus* as an outgroup.

Figure 4. Expansion of repetitive elements in the genomes of Orobanchaceae.

(A) Genome sizes and proportions of repetitive elements in six species of Orobanchaceae. The black dots and X-axis on top represent the genome sizes, and the bars and X-axis on bottom represent the percentages of different types of repetitive elements in the genomes. (B) Size distribution of introns that are conserved in Llu and Pae. (C) Repetitive element content in gene flanking regions in six species of Orobanchaceae. (D) Comparison of chromosomal repetitive elements and gene positions in chromosome 9 of Llu and chromosome 5 of Ocu. Black lines in

the boxes indicate the positions of coding genes on the chromosomes; area graphs indicate the proportion of different types of repeat sequences in the corresponding chromosomal regions; curved lines connect orthologous gene pairs. The length of the bin used for plotting is 1/1000 of the total chromosome length.

Figure 5. Gene family expansion and contraction in Orobanchaceae.

On each branch of the species tree, the green number indicates the numbers of expanded gene families, and the red number indicates the number of contracted gene families. Each density curve in the violin plot shows the F-index distribution of conserved gene families in each species indicated on the left. Species whose density curves filled with soft red, soft orange, and grayish green are holoparasitic, hemiparasitic, and autotrophic plants, respectively. The bar plot on the right shows the numbers of conserved gene families which are completely lost in each species (red bars with X-axis on bottom) and the numbers of species-specific gene families (green bars with X-axis on top).

Figure 6. Phylogenetic placement of gene duplications detected in the highly expressed genes in the haustorium.

The highly expressed genes in each species' haustorium were identified. Gene trees were constructed for all the gene families comprising these highly expressed genes. The duplication events were inferred from the topologies of the gene trees. The yellow areas of the pie charts respectively represent the proportions of genes that experienced duplication(s). The size of the yellow circle on each tree node represents the number of gene duplication events occurred during that time.

Figure 7. Gene loss in the orobanchaceous parasitic plants and Cau.

(A) History and convergent evolution of gene loss in orobanchaceous parasitic plants. Left: the red numbers on the species tree branches represent the numbers of lost OGs; the number and percentage in brackets under the species name respectively indicates the total number of lost OGs and the proportion of total number of lost OGs to the number of conserved OGs in autotrophic plants. Right: the numbers on top indicate the numbers of the same lost OGs between the two species indicated by the dumbbells after their divergence from the MRCA; the

size of soft red circle on each number indicates the *p*-value of hypergeometric test. (B) The numbers of commonly and specifically lost OGs in Cau and orobanchaceous hemiparasitic and holoparasitic plants. (C) Gene ontology (GO) enrichment analysis of lost OGs in Cau, Pae and Ocu. Each node in the networks plot represents a GO term of a biological process that was enriched (p-value < 0.02), and its size indicates the number of OGs associated with the GO term; the colors of nodes are assigned according to the species in which the gene loss occurred specifically or in common with other species; each edge represents the similarity between two GO terms based on the number of common OGs. (D) Loss of genes related to flowering time regulation in orobanchaceous parasitic plants and Cau. Grey and blue boxes respectively represent absence and presence of the specific gene. Pathway information was retrieved from FIOR-ID (http://www.phytosystems.ulg.ac.be/florid/).

Figure 8. Molecular evolution of *KAI2* gene family.

(A) The phylogenetic tree of *KAI2* family in 13 species. Blue and orange dots indicate duplication events, and the colors of branches represent the values of K_a/K_s. (B) The collinearity relationships of chromosomal fragments containing KAI2 gene in Orobanchaceae parasitic plants and closely related autotrophic species. Syntenic gene pairs are connected by curves. Curves connecting orthologs of *KAI2c*, *KAI2i*, and *KAI2d* genes are highlighted by different colors.