
COMBINED EFFECTS OF 12-WEEK YOGA WARM-UP ON ATHLETIC PERFORMANCE IN MALE HIGH SCHOOL TRACK AND FIELD ATHLETES

Danyang Wei¹

¹ Department of Sports Science, Xi'an Physical Education University, Xian, China

ABSTRACT

Background: Practicing yoga could improve balance and flexibility, but its positive significance as a long-term warm-up for formal training was uncertain. We hypothesized that practicing yoga during warm-up might positively affect balance, flexibility, and speed performance in male high school track and field athletes.

Methods: Over a 12-week period, athletes in a yoga group (YG) (n=10) practiced yoga for 15 min 4 times a week during warm-up, while athletes in a dynamic stretching group (DSG) (n=10) practiced 15 min of dynamic stretching. Except for the warm-up activities, the training content of the two groups of students was the same. We tested performance indicators immediately before and after the intervention, including lower extremity flexibility test (right hip active flexion range), lower extremity balance test [using surface electromyography (sEMG) to measure right leg tibialis anterior (TA) activation during one-leg stance (OLS) with eyes closed], and speed performance test (100-meter and 800-meter tests).

Results: We performed between-group and within-group comparisons for indicators of two groups by using SPSS (version 26.0). Within-group comparisons showed a significant improvement in flexibility (P=0.002) and balance (P=0.003) in YG, but no significant change in DSG, after the 12-week intervention. In addition, speed performance of both YG (100m, P=0.026; 800m, P=0.045) and DSG (100m, P=0.029; 800m, P=0.006) was significantly improved. Between-group comparison showed that YG had a significant advantage in 800m (P=0.045) and flexibility (P=0.031).

Conclusions: These data suggested that practicing yoga as a long-term warm-up could help male high school track and field athletes improve lower body flexibility and 800m speed. In addition, yoga had a certain positive effect on balance, but it was not significant overall.

Keywords performance · sEMG · yoga · stretching

1 Introduction

How to design a warm-up has been controversial. The prevailing view was that dynamic stretching as well as static stretching should be used as a pre-training warm-up [1, 2], and proponents of this view believed that this type of training could increase range of motion, reduce muscle resistance, and improve athletic performance [3, 4]. But now this mainstream view is being challenged.

Static stretching was used in warm-up to increase the range of motion [5]. However, now studies have shown that repetitive and sustained static stretching may impair muscle strength and speed performance, also increase the risk of sports injury during subsequent exercise due to joint instability [6, 7, 8, 9]. Due to the negative effects of static stretch, dynamic stretching was used more in warm-up activities, which showed positive effects on vertical jump height, maximal muscle strength, and sprint speed [10, 11, 12]. However, there were also some studies showing that dynamic stretching had no positive effect on these performances [13, 14].

We found some studies showing that stretching had a positive effect on athletic performance, while others found the opposite. We speculated the reason was that there were differences in the design of warm-up activities by different

researchers, including different forms of stretching combinations, movement design, intensity and duration [15, 16, 17], which would affect the effectiveness of warm-up activities. In order to design more effective warm-up activities, researchers have conducted various explorations. Lykesas et al. [18] designed a warm-up routine using Greek traditional dance, and found no difference in flexibility performance and better performance in agility tests compared to dynamic stretching in immediate response. Chen et al. [19] added a vibrating foam roller to a traditional warm-up routine and found that this combination significantly improved performance on the hexagon test and the frequency speed of kick test (FSKT). Pojskić et al. [20] used prolonged intermittent low-intensity isometric exercise as a warm-up program and found that when 30 % body weight resistance was applied, the improvements of speed and agility were similar to dynamic stretching.

In the exploration of warm-up activity design, yoga was introduced [21, 22], as practicing yoga could increase heart rate and body temperature [23] and stretch 97% of the muscles in the body [24]. Yoga practice could smooth the contraction of skeletal muscles [25], which allowed joints to be stretched to a greater extent [26]. In addition, yoga always choreographed asanas in sequence, and this dynamic pattern helped improve movement coordination and muscle recruitment efficiency [27]. By practicing yoga, trainers could be more effective at developing flexibility [28, 29] and balance [30]. However, we found that these studies did not comprehensively analyze flexibility, balance, and athletic performance.

The purpose of this paper was to address these limitations and to analyze the combined effects of practicing yoga as a long-term warm-up, including: flexibility, balance, and speed performance on high school male track and field athletes,

2 Method

2.1 Participant

The participants in this experiment were high school track and field athletes from Xi'an Soaring Sports Club, a total of 20 young males. The average age of these participants was 17 years old, and their training experience did not exceed 6 months. All players were free from any current or previous injuries and were not taking drugs that affected balance. In addition they had no prior experience with yoga and were able to make sure to train on time during the experiment. We randomly assigned those athletes to a yoga group (YG, N=10) and a dynamic stretching group (DSG, N=10), which was shown in Table 1.

Table 1: Participant Demographics by Group.

	DSG	YG	P
Height(cm)	175.5	176.9±4.2	0.439
Weight (kg)	65.0±2.7	63.9±5.0	0.547
Age (years)	16.8±0.6	17.0±0.5	0.433
Training experience(months)	3.7±0.8	4.2±1.1	0.274
BMI	21.1±0.6	20.4±1.2	0.124
Attendance over 12-week intervention	45.8±2.0	46.6±2.0	0.376

2.2 Experimental Procedures

Over a 12-week period, athletes of both groups completed exercise training sessions 4 times a week for a total of 48 sessions. YG athletes practiced yoga for 15 min after 10 min of jogging as a warm-up training, while DSG athletes practiced dynamic stretching for 15 min. The training content of the two groups was the same except for the warm-up training, and these athletes were asked not to participate in additional physical activity. Flexibility, balance, and speed performance were measured before and after the experiment.

Yoga Intervention is designed by certified yoga instructors, which was a simple but progressive 12-week program. Based on Suryanamaskar, this yoga program added several asanas focused on lower limb flexibility and balance, and all these asanas were performed in series. YG athletes needed to complete 5 times yoga procedure within 15 minutes, while stretching movements in DSG were mainly in the marching style, followed by some standing and sitting stretches.

2.3 Measurements

Flexibility: We measured the athlete's right hip active flexion range [31] by SYJDC200 angle ruler and averaged three measurements. During the measurement, the athlete lied flat on the mat, one tester immobilizing the pelvis, and then the athlete straightened the right leg and raised it up to the maximum angle.

Balance: We chose the one-leg standing test (OLS) [32] with subjects' eyes closed, and then recorded the surface electromyography (sEMG) signal of the right leg tibialis anterior (TA) during this period. Before the test, we used a razor to shave off the body hair of the test site and wiped it with alcohol. According to the suggestion of SENIAM [33], the electrode was pasted on the corresponding part from TA. We used a 6-channel sEMG sensor, setting sampling frequency to 1000Hz, and the bandpass filter frequency to 20-500Hz. Before the start of the experiment, we measured the root mean square (RMS) of the subjects' TA during the maximum voluntary isometric contraction (MVIC) in dorsiflexion for 5s. This test was repeated twice with 5 min intervals, and the maximum value was taken as the record. During the OLS test, subjects were briefly destabilized for the first few seconds after closing their eyes. To alleviate this interference, we recorded sEMG 5 s after the start of the test for a total of 10s, repeated twice, with 5 min intervals between each measurement. We calculated the RMS average of the two measurements from the TA, and divided it by the RMS during maximal voluntary isometric contractions during dorsiflexion, using %RMS for further discussion.

Speed performance: We measured the time taken by the athletes for the 100m and 800m tests twice and recorded the best time.

2.4 Data Analysis

Before all hypothesis testing, independent samples t-tests were performed using SPSS 26 for Windows (SPSS, Inc., Chicago, IL, USA) to examine whether YG and DSG were significantly different in demographics, flexibility, balance, and speed performance. We could not use ANCOVA as the test showed that the regression lines for the 100m and 800m speed data for the two groups were not parallel. The T test showed that the data of the two groups were basically the same at baseline, so we chose independent samples T test and paired samples T test for intra- and between-group comparisons to analyze changes in two groups. Significance (*) was set at the alpha level of $P = 0.05$, and data given represents the mean \pm SD unless stated otherwise.

3 Results

Table 2: Comparison of flexibility, balance and speed performance between YG and DSG.

	Pre			Post		
	DSG	YG	T-test (P Value)	DSG	YG	T-test (P Value)
Flexibility($^{\circ}$)	76.5 \pm 11.4	75.1 \pm 8.7	0.748	78.4 \pm 8.0	88.8 \pm 11.6	0.031*
Balance(%RMS)	56.2% \pm 13.7%	63.4% \pm 12.5%	0.233	58.5% \pm 12.7%	52.1% \pm 11.6%	0.253
100-meter sprint(s)	12.86 \pm 0.69	12.67 \pm 0.57	0.531	12.43 \pm 0.22	12.30 \pm 0.19	0.185
800-meter sprint(s)	148.3 \pm 9.4	145.9 \pm 11.4	0.613	142.9 \pm 4.8	138.9 \pm 2.3	0.030*

Table 3: Flexibility, balance, and speed performance changes in YG and DSG.

	DSG		YG	
	Post-pre change	T-test(P Value)	Post-pre change	T-test(P Value)
Flexibility($^{\circ}$)	1.9 \pm 4.6	0.235	13.7 \pm 10.4	0.002*
Balance(%RMS)	2.3% \pm 5.5%	0.221	-11.3% \pm 8.7%	0.003*
100-meter sprint(s)	-0.43 \pm 0.52	0.029*	-0.37 \pm 0.45	0.027*
800-meter sprint(s)	-5.4 \pm 4.8	0.008*	-7.0 \pm 9.4	0.043*

Table2 and Table3 were shown the information on the between-group and within-group comparison respectively. We could see that the indicators of two groups were basically the same at the baseline, as the T test results showed that there was no significant difference. Since these athletes were novices, their speed performance improved significantly through 12 weeks of training. The intra-group comparison showed that the time of DSG at 100m and 800m tests was shortened by 0.43s ($P < 0.05$) and 5.4s ($P < 0.01$) respectively, while YG was shortened by 0.37s ($P < 0.05$) and 7.0s ($P < 0.05$) in these

two items. The between-group comparison showed that the speed performance of YG at 800m was better than that of DSG after intervention ($P < 0.05$).

The flexibility and balance indicators in two groups improved significantly after the intervention. The active flexion angle of the right hip joint increased by 13.7° ($P < 0.01$), and the sEMG signals of the right TA during balance maintenance decreased by 11.3%RMS ($P < 0.01$), while DSG hardly changed on these two indicators. When comparing the data between groups after the intervention, YG showed a significant advantage in flexibility ($P < 0.05$), but no such advantage was observed in balance. The reason might be that there was a certain degree of individual differences in the balance indicators of two groups at baseline.

4 Discussion

In this study, both YG and DSG participated in a complete training program, including warm-up exercise, formal training, and finishing exercise. Athletes in both groups were novices and their training content was the same except for warm-up activities. Therefore, we expected an improvement in flexibility, balance, and speed performance in both groups, while the experimental results were in line with expectations to a certain extent.

Our findings showed that practicing yoga as a long-term warm-up improved flexibility in these male athletes, whereas dynamic stretching did not. Furthermore, although not significant overall, yoga practice improved the athlete's balance. In terms of speed performance, both YG and DSG have improved, but compared to DSG (-3.6%), YG (-4.8%) has a greater speed improvement at 800m, and the difference is significant.

4.1 Effects of practicing yoga as a long-term warm-up on flexibility

Previous studies found that yoga appeared to be more effective than dynamic stretching for improving flexibility in both immediate [34] and long-term [35] effects, but these studies did not practice yoga during warm-up, whereas this experiment demonstrated that yoga as a warm-up still valid. There was no change in DSG flexibility, which we speculated was due to the lack of flexibility exercises in the training programs of these athletes. A previous study showed that athletes improved strength but decreased shoulder flexibility after weightlifting [36].

We analyzed that yoga could effectively improve flexibility from two aspects. First, the practice of yoga was not likely to induce stretch reflexes [37]. The practice of yoga involved short active contractions of the stretched muscle before stretching, which may reduce the sensitivity of muscle spindles and allow for greater stretches. Second, the training content of yoga was highly structured, which seemed to be more efficient. The asanas were carried out in series during the practice, and it only took a short time to change from one movement to the next, allowing more content to be arranged in a limited time.

4.2 Effects of practicing yoga as a long-term warm-up on balance

This study found that practicing yoga during warm-up appeared to be more effective than dynamic stretching for improving balance performance. The effect of dynamic stretching on balance has been controversial [38, 39, 40, 41], while yoga has been shown to be positive for balance in different populations [42, 43, 44, 45]. The training of ankle flexibility and stability in yoga asana practice accounted for a large proportion [46, 47], which made it important for running, because ankle injuries accounted for 1/3 of running injuries [48] and ankle stability also affected the efficient in running [49].

We used sEMG to demonstrate the positive effects of yoga on balance, which was rare in previous studies. Previous studies showed that sEMG was able to accurately measure muscle activity through bioelectrical changes [50, 51, 52]. TA played an important role in the stabilization of OLS [53], and the stability of the sEMG signal amplitude during the process of maintaining balance meant the improvement of balance efficiency. So, the mechanism was using less muscle activation to accomplish the same task [54]. We noticed that YG significantly decreased the percentage of tibialis anterior muscle EMG when maintaining balance, which proved the positive effect of yoga on balance on a physiological level.

4.3 Effects of practicing yoga as a long-term warm-up on speed performance

A previous study showed that after 10 weeks of yoga intervention, the yoga group presented an advantage in flexibility compared to the control group, but there was no significant difference in 30m sprint performance [55]. Our conclusion was similar to this study, as YG also did not show an advantage in the speed performance of 100m. But unlike previous

studies, YG achieved a bigger improvement at 800m. We speculated that the reason for this result was that as flexibility and balance improve, the economy of running improved.

Practicing yoga improved flexibility, which reduced active stiffness and improved muscle performance in stretch shortening cycle (SSC) [56]. By increasing the elastic potential energy stored in the muscles during the SSC, athletes could generate greater force [57, 58]. In addition, the viscosity of the muscle decreased, reducing the energy loss during SSC [59]. Improvements in balance could reduce the proportion of stabilizing muscles involved in balance tasks [60], which helped reduce erratic movements [61] and enhanced the synergistic contraction of agonist and antagonist muscles [62]. Ultimately these factors all increased the efficiency of the exercise.

Practicing yoga as a long-term warm-up improved the efficiency of exercise. Compared to DSG, YG showed a significant advantage in speed performance at 800m, but no difference at 100m. We speculated that the effect of sport economy on speed performance needed more time to manifest, because it took a lot longer to complete 800m than 100m, which allowed the movement economy to accumulate.

The focus of this study was to discuss the combined effects of yoga as a warm-up activity on high school track and field athletes, so we did not design additional groups to differentiate between the effects of flexibility and balance on exercise economy, which prevented us from discussing the impact of these two qualities on speed performance separately.

5 Conclusion

After 12 weeks of training, male track and field athletes who practiced yoga as a warm-up has significantly improved flexibility compared to dynamic stretching. Additionally, although not significantly overall, yoga practice improved the athlete's balance. Finally, yoga practice boosted athlete's 800m speed, which we analyzed was because yoga improved running economy. Therefore, we recommended that novice male track and field athletes practiced yoga sequences in general warm-up activities, which could help athletes improve balance, flexibility, and middle-distance running performance.

6 Financial support and sponsorship

Nil

7 Conflicts of interest

There are no conflicts of interest.

8 Acknowledgment

We would like to thank the coaches and participants of the study for their time.

References

- [1] American College of Sports Medicine et al. *ACSM's guidelines for exercise testing and prescription*. Lippincott Williams & Wilkins, 2013.
- [2] David G Behm and Anis Chaouachi. A review of the acute effects of static and dynamic stretching on performance. *European journal of applied physiology*, 111(11):2633–2651, 2011.
- [3] Bruno L Franco, Gabriel R Signorelli, Gabriel S Trajano, Pablo B Costa, and Carlos G de Oliveira. Acute effects of three different stretching protocols on the wingate test performance. *Journal of sports science & medicine*, 11(1):1, 2012.
- [4] Malachy P McHugh and CH Cosgrave. To stretch or not to stretch: the role of stretching in injury prevention and performance. *Scandinavian journal of medicine & science in sports*, 20(2):169–181, 2010.
- [5] William D Bandy and Jean M Irion. The effect of time on static stretch on the flexibility of the hamstring muscles. *Physical therapy*, 74(9):845–850, 1994.
- [6] Brandon M Kistler, Mark S Walsh, Thelma S Horn, and Ronald H Cox. The acute effects of static stretching on the sprint performance of collegiate men in the 60-and 100-m dash after a dynamic warm-up. *The Journal of Strength & Conditioning Research*, 24(9):2280–2284, 2010.

- [7] Malachy P McHugh and CH Cosgrave. To stretch or not to stretch: the role of stretching in injury prevention and performance. *Scandinavian journal of medicine & science in sports*, 20(2):169–181, 2010.
- [8] Martin Loughran, Philip Glasgow, Chris Bleakley, and Joseph McVeigh. The effects of a combined static-dynamic stretching protocol on athletic performance in elite gaelic footballers: A randomised controlled crossover trial. *Physical Therapy in Sport*, 25:47–54, 2017.
- [9] Akinori Nagano, Shinsuke Yoshioka, Dean C Hay, Ryutaro Himeno, and Senshi Fukushima. Influence of vision and static stretch of the calf muscles on postural sway during quiet standing. *Human movement science*, 25(3):422–434, 2006.
- [10] Alan J Pearce, Dawson J Kidgell, James Zois, and John S Carlson. Effects of secondary warm up following stretching. *European journal of applied physiology*, 105(2):175–183, 2009.
- [11] Thomas Little and Alun G Williams. Effects of differential stretching protocols during warm-ups on high speed motor capacities in professional soccer players. *Journal of strength and conditioning research*, 20(1):203–7, 2006.
- [12] Taichi Yamaguchi and Kojiro Ishii. Effects of static stretching for 30 seconds and dynamic stretching on leg extension power. *The Journal of Strength and Conditioning Research*, 19(3):677–683, 2005.
- [13] Trent J Herda, Joel T Cramer, Eric D Ryan, Malachy P McHugh, and Jeffrey R Stout. Acute effects of static versus dynamic stretching on isometric peak torque, electromyography, and mechanomyography of the biceps femoris muscle. *The Journal of Strength & Conditioning Research*, 22(3):809–817, 2008.
- [14] Anis Chaouachi, Carlo Castagna, Mokhtar Chtara, Matt Brughelli, Olfa Turki, Oliver Galy, Karim Chamari, and David G Behm. Effect of warm-ups involving static or dynamic stretching on agility, sprinting, and jumping performance in trained individuals. *The Journal of Strength & Conditioning Research*, 24(8):2001–2011, 2010.
- [15] Yaser Alikhajeh, Nasser Mohamad Rahimi, Khadijeh Fazeli, and Hajar Fazeli. The effect of different warm up stretch protocols on 20m-sprint performance in trained soccer players. *Procedia-Social and Behavioral Sciences*, 46:2210–2214, 2012.
- [16] Ognjen Andrejić. An investigation into the effects of different warm-up protocols on flexibility and jumping performance in youth. *Facta universitatis-series: Physical Education and Sport*, 10(2):107–114, 2012.
- [17] Michael J Redd, Tristan M Starling-Smith, Chad H Herring, Matt S Stock, Adam J Wells, Jeffrey R Stout, and David H Fukuda. Tensiomyographic responses to warm-up protocols in collegiate male soccer athletes. *Journal of Functional Morphology and Kinesiology*, 6(4):80, 2021.
- [18] Georgios Lykesas, Ioannis Giossos, Dimitrios Chatzopoulos, Maria Koutsouba, Stella Douka, and Eugenia Nikolaki. Effects of several warm-up protocols (static, dynamic, no stretching, greek traditional dance) on motor skill performance in primary school students. *International Electronic Journal of Elementary Education*, 12(5):481–487, 2020.
- [19] An-Hsu Chen, Chih-Hui Chiu, Chin-Hsien Hsu, I-Lin Wang, Kuei-Ming Chou, Yung-Shen Tsai, Yu-Fang Lin, and Che-Hsiu Chen. Acute effects of vibration foam rolling warm-up on jump and flexibility asymmetry, agility and frequency speed of kick test performance in taekwondo athletes. *Symmetry*, 13(9):1664, 2021.
- [20] Haris Pojskić, Jeffrey C Pagaduan, F Babajić, E Užičanin, M Muratović, and M Tomljanović. Acute effects of prolonged intermittent low-intensity isometric warm-up schemes on jump, sprint, and agility performance in collegiate soccer players. *Biology of sport*, 32(2):129, 2015.
- [21] Saravanan Elumalai and D Pajanivel. Comparative effects of traditional warming up and suryanamaskar as warming up on explosive power of volleyball players.
- [22] Abhijit Thander. Comparative effects of suryanamaskar and dynamic stretching on cricket specific motor performance in fast bowling, 2021.
- [23] Marian E Papp, Petra Lindfors, Malin Nygren-Bonnier, Lennart Gullstrand, and Per E Wändell. Effects of high-intensity hatha yoga on cardiovascular fitness, adipocytokines, and apolipoproteins in healthy students: a randomized controlled study. *The Journal of Alternative and Complementary Medicine*, 22(1):81–87, 2016.
- [24] Marian E Papp, Malin Nygren-Bonnier, Lennart Gullstrand, Per E Wändell, and Petra Lindfors. A randomized controlled pilot study of the effects of 6-week high intensity hatha yoga protocol on health-related outcomes among students. *Journal of Bodywork and Movement Therapies*, 23(4):766–772, 2019.
- [25] Catherine Woodyard. Exploring the therapeutic effects of yoga and its ability to increase quality of life. *International journal of yoga*, 4(2):49, 2011.
- [26] Sivananda Yoga Vedanta Centre. *Yoga Your Home Practice Companion*. Dorling Kindersley Ltd, 2010.
- [27] Mara Carrico et al. *Yoga journal's yoga basics: the essential beginner's guide to yoga for a lifetime of health and fitness*. Macmillan, 1997.

- [28] R Meera, R Mohanakrishnan, and T Arun Prasanna. Effect of core training with and without yogic practices on elasticity among college female athletes. *EXECUTIVE EDITOR*, 11(01):511, 2020.
- [29] MV Ajayaghosh and V Mahadevan. Effect of functional strength training and vinyasa flow yoga on selected physical variables among men soccer players. 2018.
- [30] M Jay Polsgrove, Brandon M Eggleston, and Roch J Lockyer. Impact of 10-weeks of yoga practice on flexibility and balance of college athletes. *International journal of yoga*, 9(1):27, 2016.
- [31] American Academy of Orthopaedic Surgeons et al. Joint motion: Method of measuring and recording,(1965). *American Academy of Orthopaedic Surgeons, Chicago*, 1988.
- [32] Jaap H van Dieën, Marloes van Leeuwen, and Gert S Faber. Learning to balance on one leg: motor strategy and sensory weighting. *Journal of neurophysiology*, 114(5):2967–2982, 2015.
- [33] Hermie J Hermens, Bart Freriks, Catherine Disselhorst-Klug, and Günter Rau. Development of recommendations for semg sensors and sensor placement procedures. *Journal of electromyography and Kinesiology*, 10(5):361–374, 2000.
- [34] Barton Lee Schroeter. *A Study on the Acute Effects of Yoga and Dynamic Stretching*. PhD thesis, 2016.
- [35] Prosenjit Paul, Kallol Chatterjee, and Biswabandhu Nayek. Effect of selected yogic practices and dynamic stretching on flexibility of the school children.
- [36] M Jay Polsgrove, Brandon M Eggleston, and Roch J Lockyer. Impact of 10-weeks of yoga practice on flexibility and balance of college athletes. *International journal of yoga*, 9(1):27, 2016.
- [37] Kalyan B Bhattacharyya. The stretch reflex and the contributions of c david marsden. *Annals of Indian Academy of Neurology*, 20(1):1, 2017.
- [38] Dimitris Chatzopoulos, Christos Galazoulas, Dimitrios Patikas, and Christos Kotzamanidis. Acute effects of static and dynamic stretching on balance, agility, reaction time and movement time. *Journal of sports science & medicine*, 13(2):403, 2014.
- [39] Nevzad Denerel, Metin Ergün, Oğuz Yüksel, Cengizhan Özgürbüz, and Oğuz Karamızrak. The acute effects of static and dynamic stretching exercises on dynamic balance performance. *Spor Hekimligi Dergisi/Turkish Journal of Sports Medicine*, 54(3), 2019.
- [40] L Belkhiria-Turki, A Chaouachi, O Turki, R Hammami, M Chtara, M Amri, EJ Drinkwater, and DG Behm. Greater volumes of static and dynamic stretching within a warm-up do not impair star excursion balance performance. *J Sports Med Phys Fitness*, 54(3):279–88, 2014.
- [41] Niamh Morrin and Emma Redding. Acute effects of warm-up stretch protocols on balance, vertical jump height, and range of motion in dancers. *Journal of dance medicine & science*, 17(1):34–40, 2013.
- [42] Arlene A Schmid, Marieke Van Puymbroeck, and David M Kocejka. Effect of a 12-week yoga intervention on fear of falling and balance in older adults: a pilot study. *Archives of physical medicine and rehabilitation*, 91(4):576–583, 2010.
- [43] Dheli Kadachha, Neela Soni, and Ankur Parekh. Effects of yogasana on balance in geriatric population. *International Journal of Physiotherapy and Research*, 4(2):1401–1407, 2016.
- [44] Tilak Raj. *The effects of yoga on sports performance and health: A thesis submitted in partial fulfilment of the requirements for the Degree of Doctor of Philosophy at Lincoln University*. PhD thesis, Lincoln University, 2021.
- [45] Shah Noman Md Iftekher, Md Bakhtiar, and Kh Shafiur Rahaman. Effects of yoga on flexibility and balance: a quasi-experimental study. *Asian Journal of Medical and Biological Research*, 3(2):276–281, 2017.
- [46] David P Heller. *The Effect of a Specialized Yoga Intervention on Characteristics of Functional Ankle Instability*. PhD thesis, University of Kansas, 2020.
- [47] DP Heller, M Maienschein, and W Liu. Effect of a 4-week yoga intervention on characteristics of functional ankle instability: A pilot study. In *International Journal of Exercise Science: Conference Proceedings*, volume 11, page 38, 2018.
- [48] Adam S Tenforde, Amy Yin, and Kenneth J Hunt. Foot and ankle injuries in runners. *Physical Medicine and Rehabilitation Clinics*, 27(1):121–137, 2016.
- [49] Stefano Lazzer, Paolo Taboga, Desy Salvadego, Enrico Rejc, Bostjan Simunic, Marco V Narici, Antonio Buglione, Nicola Giovanelli, Guglielmo Antonutto, Bruno Grassi, et al. Factors affecting metabolic cost of transport during a multi-stage running race. *Journal of Experimental Biology*, 217(5):787–795, 2014.

- [50] Chiako Mokri, Mahdi Bamdad, and Vahid Abolghasemi. Muscle force estimation from lower limb emg signals using novel optimised machine learning techniques. *Medical & Biological Engineering & Computing*, pages 1–17, 2022.
- [51] Ines Chihi, Lilia Sidhom, and Ernest Nlandu Kamavuako. Hammerstein–wiener multimodel approach for fast and efficient muscle force estimation from emg signals. *Biosensors*, 12(2):117, 2022.
- [52] Florent Moissenet, Anne Tabard-Fougère, Stéphane Genevay, and Stéphane Armand. Normalisation of a biarticular muscle emg signal using a submaximal voluntary contraction: Choice of the standardised isometric task for the rectus femoris, a pilot study. *Gait & posture*, 91:161–164, 2022.
- [53] Yu Uchida and Shinichi Demura. Differences in leg muscle activity and body sway between elderly adults able and unable to maintain one-leg stance for 1 min: the effect of hand support. *Aging clinical and experimental research*, 28(4):669–677, 2016.
- [54] Dennis Brueckner, Beat Göpfert, Rainer Kiss, and Thomas Muehlbauer. Effects of motor practice on learning a dynamic balance task in healthy young adults: A wavelet-based time-frequency analysis. *Gait & posture*, 70:264–269, 2019.
- [55] Tilak Raj, Michael J Hamlin, and Catherine A Elliot. Association between hamstring flexibility and sprint speed after 8 weeks of yoga in male rugby players. *International Journal of Yoga*, 14(1):71, 2021.
- [56] GREGORY J Wilson, BRUCE C Elliott, and GRAEME A Wood. Stretch shorten cycle performance enhancement through flexibility training. *Medicine and Science in Sports and Exercise*, 24(1):116–123, 1992.
- [57] Martyn R Shorten. Muscle elasticity and human performance. In *Current research in sports biomechanics*, volume 25, pages 1–18. Karger Publishers, 1987.
- [58] Paavo Komi. *Strength and power in sport*, volume 3. John Wiley & Sons, 2008.
- [59] Keitaro Kubo. In vivo elastic properties of human tendon structures in lower limb. *International Journal of Sport and Health Science*, 3(Special_Issue_2005):143–151, 2005.
- [60] Crystal O Kean, David G Behm, and Warren B Young. Fixed foot balance training increases rectus femoris activation during landing and jump height in recreationally active women. *Journal of sports science & medicine*, 5(1):138, 2006.
- [61] Wolfgang Taube, Markus Gruber, and Albert Gollhofer. Spinal and supraspinal adaptations associated with balance training and their functional relevance. *Acta physiologica*, 193(2):101–116, 2008.
- [62] David G Lloyd. Rationale for training programs to reduce anterior cruciate ligament injuries in australian football. *Journal of Orthopaedic & Sports Physical Therapy*, 31(11):645–654, 2001.