Sustained plumage divergence despite weak genomic differentiation and broad sympatry in sister species of Australian woodswallows (*Artamus* spp.)

Joshua V. Peñalba[1], Jeffrey L. Peters[2], Leo Joseph[3]

[1] Museum für Naturkunde Berlin, Leibniz Institute for Evolution and Biodiversity Science, Center for Integrative Biodiversity Discovery, Invalidenstr. 43, D-10115, Berlin, Germany

[2] Department of Biological Sciences, Wright State University, Dayton, OH, USA

[3] Australian National Wildlife Collection, CSIRO National Research Collections Australia, Canberra, Australia

Abstract

Plumage divergence can function as a strong premating barrier when species come into secondary contact. When it fails to do so, the results are often genome homogenization and phenotypic hybrids at the zone of contact. This is not the case in the largely sympatric masked- and white-browed woodswallow species (Passeriformes: Artamidae: *Artamus* spp) complex in Australia where phenotypic integrity is sustained despite no discernible mitochondrial structure. This lack of structure may suggest recent divergence, ongoing gene flow or both, and phenotypic hybrids are reported albeit rarely. Here, we further assessed the population structure and differentiation across the species' nuclear genomes using ddRAD-seq. As found in the mitochondrial genome, no structure or divergence within or between the two species is detected in the nuclear genome. This coarse sampling of the genome nonetheless revealed peaks of differentiation around the genes *SOX5* and *Axin1*. Both are involved in the *Wnt/*β-catenin signaling pathway, which regulates feather development. Reconstruction of demographic history supports a scenario with ongoing gene flow during divergence but involving shifts in direction and magnitude. Our study informs how divergent plumage morphs may arise and be sustained despite whole-genome homogenization and reveals new candidate genes potentially involved in plumage divergence.

Introduction

Plumage divergence between sister lineages of birds often distinguishes different subspecies and species. Often, the more divergent populations are in plumage, especially in changes of pattern rather than of simple color replacement, the more likely they are to be classified as different taxonomic species (Patten & Unitt, 2002; Paxton, 2009). This is not only because it is easier for us to distinguish between forms but also because plumage is a secondary sexual trait playing a role in mate choice and premating isolation (Mason & Bowie, 2020; Price, 2008). Large plumage differences and corresponding mate preferences can accumulate in allopatry and can subsequently serve as a strong barrier to gene flow when populations come in secondary contact resulting in speciation (Cowles & Uy, 2019; Sætre et al., 1997). Plumage traits can also serve as a target for divergent ecological selection. If this divergence also results in differences in mate choice, plumage can serve as 'magic traits' in facilitating speciation (Servedio et al., 2011). Comparative studies on both plumage and genomic divergence have shown a positive association with plumage divergence and reproductive isolation (Winger & Bates, 2015).

Despite this association between plumage and genome divergence, there are also many documented cases of a lack of genomic differentiation despite strong plumage divergence. These unusual systems have been crucial in revealing the genetic architecture of complex plumage traits and correlative if not causative links between plumage divergence and particular genes involved. When most of the genome is undifferentiated or has been homogenized by gene flow, loci that remain strongly differentiated usually underlie various components of plumage traits. For example, the carrion and hooded crows (*Corvus corone*, *C. cornix*, respectively) in Europe display strong plumage integrity despite extensive gene flow in their hybrid zone. An epistatic interaction between a gene, *NDP*, and a large region in chromosome 18 explains the divergence in plumage and variation in hybrid morphs (Knief et al., 2019). This genetic architecture coupled with mate-choice dynamics can explain the narrow hybrid zone despite extensive introgression (Metzler et al., 2020). Hybridization between the golden-winged warbler (*Vermivora chrysoptera*) and blue-winged warbler (*V. cyanoptera*; F_{ST} ~ 0.0004) in North America has revealed just six divergent peaks in the genome, four of which were adjacent to genes known to be involved in feather development and pigmentation (Toews et al., 2016). They were also able to isolate the *agouti signaling protein* (*ASIP*) as responsible for the divergence in throat coloration in the two morphs. Asymmetric introgression of the head plumage trait in two subspecies of the white wagtail (*Motacilla alba*) has revealed a partial dominant and epistatic interaction between two small regions in chromosome 1A and 20 (Semenov et al., 2021). Extensive introgression between two subspecies of the northern flicker (*Colaptes auratus*) in North America (F_{ST} ~ 0.008) has

revealed the genomic basis underlying four plumage traits and the role of *CYPJ2J19* to cause a red to yellow transition (Aguillon et al., 2021).

Three common findings emerge from these studies: (1) extensive introgression can occur despite plumage divergence or, conversely, plumage divergence can be sustained despite extensive introgression; (2) one or more of many different genes can be involved in plumage divergence and so are not always shared among systems, and (3) the genes involved in plumage divergence (a secondary sexual trait) are usually found on autosomes rather than the sex chromosomes.

Another shared feature of these study systems is that they include hybrid zones between two populations. This means that phenotypic hybrids are likely geographically restricted and can serve as a buffer for contact between the parental plumage forms. It is also likely that populations diverged in allopatry and came into secondary contact in the hybrid zone or diverged with relatively restricted gene flow only occurring in the zone of contact.

Here, we present a genomic interrogation of an unusual case that confounds these emergent themes. It concerns two Australian birds never classified as anything other than two species (see Schodde & Mason, 1999), the masked woodswallow (*Artamus personatus*) and white-browed woodswallow (*Artamus superciliosus*) (Passeriformes: Artamidae; Figure 1).They are largely sympatric across much of the eastern half of the Australian continent but only the masked woodswallow occurs across much of the continent's western half (Carnaby, 1965; Higgins, 2006; Johnstone & Storr, 2004; Figure S1). The two are phenotypically very different but some plumage traits are reasonably inferred to be homologous (e.g., dark facial "mask" in both; white post-supraocular spot of masked and the complete white superciliar stripe of white-browed; Figure 1). Especially notable given that they are a pair of closely related oscine songbird taxa is that their vocalizations appear all but identical, no discernible difference ever having been rigorously described Higgins *et al*. 2006; *cf* Lowe and Lowe 1972). Unlike the systems cited above, however, the two are often found in mixed flocks of up to hundreds of birds and breed near one another, even in the same tree (Carnaby, 1965; Higgins, 2006; Lenz, 2019). Despite this, phenotypic hybrids are rare but not unknown (Barnard, 1944; McGill, 1944); we are aware of photographic records of seven clearly intermediate individuals of presumed hybrid descent (e.g., see images at https://ebird.org/checklist/S79795345; https://ebird.org/australia/checklist/S61566354; Figures S2-S8).

Figure 1. *Population structure.* The grey and maroon correspond to masked- and white-browed woodswallow, respectively. The assignment to species is done purely by plumage (A) Autosomal network generated from uncorrected pairwise distances among individuals (B) Z chromosome network generated from uncorrected pairwise distances among individuals (C) Mitochondrial ND2 haplotype network (note that one internal haplotype is shared due to addition of data since Joseph *et al.* (2006) (D) Population assignment from 20 iterations of STRUCTURE with K=2 (E) Principal components analysis using smartPCA

To date, available molecular data concern only their mitochondrial DNA diversity. As in the systems cited above, they show no discernible population structure in mtDNA despite their striking plumage divergences and they are paraphyletic with respect to each other but together form a sister clade to the rest of *Artamus* (Joseph et al., 2006). The observed mtDNA structure detected in that study is consistent with the later stage of incomplete lineage sorting termed allophyly by Omland et al., (2006; see Figure 1). From this brief summary, we infer that while these two woodswallows exhibit substantial but incomplete premating isolation this has not led to any signal of population structure in incompletely sorted mtDNA. Moreover, we see no clear evidence of the phenomenon of a phenotypic hybrid swarm (Cockayne & Allan, 1926) throughout their extensive range of sympatry in eastern Australia.

In this study, we extend the earlier work on mtDNA (Joseph et al., 2006) to test whether the nuclear genome similarly lacks genetic structure between the two species or whether the previous pattern is a result of extensive mitochondrial introgression. We used ddRAD sequencing to sample loci across the nuclear genome to test whether there is genome-wide divergence across the chromosomes, to describe variation in divergence between autosomes and sex chromosomes, and to test whether divergence is localized to a few regions that may underlie plumage-related genes of the sister species. We then inferred the demographic history that could have resulted in the two morphs we see today. Lastly, we simulated various evolutionary scenarios that could have resulted in the plumage divergence to see which best explains our empirical data.

Methods

Sampling

We obtained tissue samples from 19 masked woodswallows (*A. personatus*) and 12 white-browed woodswallows (*A. superciliosus*) from the Australian National Wildlife Collection, CSIRO, Canberra (Table S1). The majority of samples were collected from eastern Australia where the two species are broadly sympatric (supplementary material, Figure S1). Parenthetically, we note that the samples are derived from several decades of opportunistic sampling reflecting the difficulty of sampling species most frequently seen hundreds of meters into the sky and which have often unpredictable nomadic movements. We also included three individuals from each of the following species as outgroups: white-breasted woodswallow (*A. leucorhynchus*), little woodswallow (*A. minor*), black-faced woodswallow (*A. cinereus*), and dusky woodswallow (*A. cyanopterus*).

ddRAD-seq library preparation and bioinformatics processing

We generated DNA sequences using the double-digest restriction-site-associated DNA sequencing (ddRAD-seq) protocol of (DaCosta & Sorenson, 2014). DNA was extracted using a DNeasy Blood and Tissue Kit (Qiagen, Valencia, CA), and approximately 1 µg of DNA was digested using the restriction enzymes *SbfI* and *EcoRI*. Adapters compatible with Illumina TruSeq reagents and containing unique barcodes were ligated to the digested DNA. Ligated DNA fragments ranging in size from 300 to 450 base pairs (bp) were extracted from 2% low-melt agarose gels and purified using a MinElute gel extraction kit (Qiagen, Valencia, CA). The recovered fragments were amplified using PCR, purified using magnetic AMPure XP beads (Beckman Coulter Inc., Indianapolis, IN), and quantified using real-time PCR with an Illumina library quantification kit (KAPA Biosystems, Wilmington, MA). Uniquely barcoded

samples were pooled in equimolar concentrations and sequenced (150 bp reads) on an Illumina HiSeq 2500 at TUCF Genomics, Tufts University (Medford, MA, USA).

Raw Illumina reads were processed using the computational pipeline described by DaCosta and Sorenson (2014) [Scripts available at: http://github.com/BU-RAD-seq/ddRAD-seq-Pipeline]. For each sample, identical reads were combined into a single read and recoded with the number of reads and the highest quality score for each nucleotide position. Reads with an average Phred score of <20 were removed. Retained reads from all individuals were clustered into putative loci using USEARCH v. 5 (R. C. Edgar, 2010), with an –id setting of 0.85, and aligned using MUSCLE V. 3 (R. C. Edgar, 2004). Individuals were genotyped at each locus as described in DaCosta and Sorenson (2014): homozygotes were defined when ≥93% of the reads were identical, whereas heterozygotes were defined when a second sequence was represented by ≥29% of reads, or if a second sequence was represented by as few as 10% of reads and the haplotype was confirmed in other individuals. Genotypes were flagged if none of these criteria were met or more than two haplotypes met the criteria. From these flagged genotypes, we retained the allele represented by the majority of reads and scored the second allele as missing data. Similarly, a second allele was scored as missing when the locus was represented by ≤5 reads. We retained all loci that contained ≤10% missing genotypes and ≤5% flagged genotypes.

Genome placement

We used BLAST to assign the chromosome and location for each ddRAD locus (Altschul et al., 1990). The ddRAD loci were queried against the New Caledonian crow genome (*Corvus moneduloides;* GenBank Accession GCA_009650955.1), the closest relative with a chromosome-scale reference genome available. This was done to maximize the number of loci that can be placed along a chromosome and to minimize difference in marker order due to chromosomal rearrangements. To stay consistent with nomenclature used in other studies, we renamed the *C. moneduloides* chromosomes based on the zebra finch nomenclature (Table S2) for our study. The original *C. moneduloides* chromosomes were named by descending order of chromosome length.

Population structure

The population structure from the nuclear genome was inferred in various ways. First, we used smartPCA from EIGENSOFT (v7.2.1) (http://www.hsph.harvard.edu/alkes-price/software/) to infer the structure through a Principal Components Analysis (PCA). Prior to running the software, the SNPs were filtered to only include variants fixed in the outgroups, non-singletons, and one SNP per ddRAD locus. This was to minimize the effects of incomplete lineage sorting and linkage and to maximize signal. We then ran STRUCTURE

using the same set of filtered SNPs (Pritchard et al., 2000). We ran 20 iterations of STRUCTURE for K=1 and K=2 and merged the iterations within K=2 using CLUMPP (Jakobsson & Rosenberg, 2007). The population network was generated using uncorrected pairwise distances among individuals. The network for the autosomal and Z-linked loci were inferred separately. Unlike for the PCA, the SNPs were not filtered for this analysis. The population network was then visualized using SplitsTree (Huson, 1998). Finally, the mitochondrial ND2 haplotype network was inferred and visualized using the TCS network implemented in PopArt (Leigh & Bryant, 2015). The ND2 sequences were downloaded from GenBank (accession numbers in Table S1).

Population genetic parameters

All population genetic parameters were calculated using custom scripts and the following equations. Population differentiation was estimated using Hudson's F_{ST} which is less influenced by differences in effective population size (Bhatia et al., 2013). We used the following equation: $F_{ST} = \frac{\left[(p_1-p_2)^2 - \frac{p_1(1-p_1)}{n_1-1} - \frac{p_2(1-p_2)}{n_2-1}\right]}{p_1(1-p_2)+p_2(1-p_1)}$, with p_1 and p_2 being the frequency of the allele in population 1 and population 2, respectively, and n_1 and n_2 being the number of chromosomes sampled in each population. Average F_{ST} was calculated within each ddRAD locus and across the genome using the ratio of averages. Absolute population divergence (D_{XY}) was estimated using the following equation $D_{XY} = p_1(1 - p_2) + p_2(1 - p_1)$ divided by the total length in base pairs. Genetic diversity within each species was calculated using both Tajima's (π) and Watterson's estimator (θ_W) using the equations $\pi = 2p(1 - p)$ and

$\theta_W = S/(\sum_1^{k-1} \frac{1}{k})$ divided by the total length in base pairs with p being the allele frequency of either allele, S the number of segregating sites and *k* the number of chromosomes. Lastly, Tajima's D was calculated using $D = (\hat{k} - \frac{S}{a_1})/\sqrt{e_1 S + e_2 S(S - 1)}$. The definitions for the different variables can be found in Tajima (1989).

Demographic modeling

We used fastsimcoal2 (Excofffier et al., 2021) to test various demographic scenarios between *A. superciliosus* and *A. personatus* using the unfolded 2D joint site frequency spectrum (SFS). The SNPs were filtered to retain only those fixed for all four outgroup species and we subsampled a single SNP per ddRAD locus to allow for an unfolded joint frequency spectrum of unlinked sites. We assumed the allele fixed for all outgroup species to be the ancestral allele. We also omitted sex-linked loci and outlier loci to minimize effects of differences in effective population size and strong selection within the genome. This retained

4169 unlinked, autosomal SNPs. We tested models of panmixia, recent divergence, isolation-with-migration, secondary contact, and change in migration. In order to be comparable to the other models, panmixia was simulated as two populations with a migration rate m = 0.5 which corresponds to full sympatry (Coyne & Orr, 2004). The change in migration scenario includes two time periods with differing levels of migration to reflect a history of low migration during divergence followed by a period of higher migration which homogenized the divergence. We created bootstrapped SFS by choosing a single random SNP within each ddRAD locus so the model selection is not influenced by stochastic differences in SNP selection. For model selection, we ran each model 100 times for 10 bootstrapped SFS and the run with the best likelihood for SFS was chosen.

When performing parameter estimation after choosing the best model, we discovered two optimal parameter combinations where either masked- or white-browed woodswallow would have the larger effective population size and corresponding migration rate. To differentiate between these scenarios, we performed another test comparing a model where migration rate parameter space is bound such that: $m_{MAWS->WBWS} > m_{WBWS->MAWS}$ versus $m_{WBWS->MAWS} < m_{MAWS->WBWS}$. We fixed parameters that had low variance from our initial runs and let other parameters such as effective population size still vary. We then chose the scenario with the higher likelihood and restricted the migration rates while estimating the variance of the parameters.

To obtain the variance in the parameters for the best scenario, we bootstrapped the SNPs to generate 50 SFS and reran the parameter inference 50 times for each. The parameters that were previously fixed were allowed to vary for this final parameter estimate. The best run for each SFS was chosen and the variance was calculated across the 50 bootstrapped SFS.

F_{ST} outlier analysis and gene ontology search
Because both species are largely sympatric throughout the range of white-browed woodswallow and the sampling was mostly done in the same locations (Figure S1), we did not need to test whether the F_{ST} outliers are due to isolation-by-distance as you might in a classic parapatric scenario. Each F_{ST} value corresponds to a single ddRAD locus rather than a single SNP. We used a 99.5 percentile threshold to identify putative outliers. Of these outliers, we also investigated four loci which were substantially more differentiated than the rest. We then identified which genes were 40kbp upstream and downstream of these outliers using the genome annotation of *Corvus moneduloides*. We retrieved the names and molecular functions of these genes based on homology to the chicken (*Gallus gallus*) annotation. We also searched whether we had ddRAD loci within the following known plumage genes from other studies: *ASIP, FST, CYPJ19, MC1R, TYR, TYRP1, OCA2,*

SLC45A, KIT, DCT, and *Mitf* (Funk & Taylor, 2019; Kim et al., 2019; San-Jose et al., 2017; Toews et al., 2016; Toomey et al., 2018; Twyman et al., 2018)*.*

SLiM simulations

To investigate the potential underlying mechanism that resulted in the outlier loci, we simulated various evolutionary scenarios that could result in high F_{ST}. Using SLiM3 (Haller & Messer, 2019) we simulated a 500bp sequence to represent an independent ddRAD locus under different scenarios. We simulated five main scenarios: *neutral, background selection, balancing selection, ancient sweep,* and *recent sweep*. For *neutral, background selection,* and *balancing selection* we also had a scenario with and without gene flow to reflect freely moving and barrier loci, respectively. Within *ancient sweep* and *recent sweep*, we also tested combinations of *hard sweeps* vs. *soft sweeps* and *complete sweeps* vs. *partial sweeps*. This resulted in a total of 14 different scenarios. The *neutral* and *background selection* were our null models where divergence was not driven by differences in selection regimes (Comeron, 2017). Both mutation and recombination rates were kept constant and we used the parameters estimated by the demographic modeling for population sizes, divergence times, and gene flow. We simulated each scenario 100 times and extracted the F_{ST} and D_{XY} between populations and the θ and π within populations. Specific details of the simulations can be found in the supplementary methods.

Results

ddRAD sequencing

We obtained an average of 869,568reads per individual (median = 754,301; range = 351,718–4,669,735). After filtering and assembly, we recovered a total of 4664 ddRAD loci containing 76 336 SNPs in 582 775 bp. Of these, 50 913 SNPs in 4371 loci were fixed in all four outgroup species. We were able to locate the genomic position of 4428 loci in the *Corvus modenuloides* assembly (181 of which mapped to the Z chromosome).

Population structure, differentiation and genetic diversity

We found no population structure in the nuclear ddRAD loci between the two species (Figure 1A to E). Both the autosomal and Z-linked markers (Figure 1 A, B) showed a distinctive star-like pattern indicative of a fairly panmictic population despite divergence in many plumage traits. The STRUCTURE plot for K=2 (Figure 1D) also shows no discernible population structure. The mean Ln likelihood of the data across the 20 runs is -63104.35 (±23.99) for K=1 and -78574.17 (±13503.70) for K=2 suggesting a model of no population structure fits

better. The PCA (Figure 1E) also does not separate the two species into distinct clusters but rather reflects differences in variance within species.

Consistent with the lack of population structure, the genetic differentiation between the two species is very low, the average autosomal and Z chromosome F_{ST} being 0.003 and 0.0002, respectively. Although the average F_{ST} is an order of magnitude lower in the Z chromosome and contrary to expectation from differences in effective population size, the median F_{ST} is 0.0 for both autosomes and Z chromosome. This suggests that the elevated F_{ST} in the autosomes is driven by the few outliers (Figure 2). The global D_{XY} between the two species is D_{XY} = 0.012 per bp. The estimates for genetic diversity within the species were similar: π_{WBWS}=0.0118, π_{MAWS}=0.0120, θ_{WBWS} = 0.0176, θ_{MAWS}=0.0199. The estimates of Tajima's D within each species were D_{WBWS}= -1.026 and D_{MAWS}= -1.184. The comparisons among parameters of the per locus estimates can be found in Figure 2.

Figure 2. *Genome differentiation and outlier loci.* Top left: Genetic differentiation across the ddRAD loci by physical order on the *C. moneduloides* reference genome. The alternating colors represent different chromosomes. The red line corresponds to the 0.995 percentile threshold. The points with open circles represent the extreme outlier loci. Top right: Histogram of estimates of genetic differentiation across ddRAD loci. Bottom (from left to right): Comparison between absolute and relative genetic divergence among ddRAD loci, comparison of genetic diversity (Watterson's theta among the ddRAD loci between the white-browed and masked woodswallows, comparison of nucleotide diversity (pi) among the ddRAD loci between the white-browed and masked woodswallows, and comparison of Tajima's D among ddRAD loci between the white-browed and masked woodswallows.

Demographic modeling

Five different demographic scenarios were tested. Of the five, the change in migration scenario had the best support (Figure 3). Isolation-with-migration ($\Delta\overline{AIC}$ = 118.09) and secondary contact ($\Delta\overline{AIC}$ = 107.27) had slightly lower support. The scenarios of recent

divergence ($\Delta\overline{AIC}$ = 840.98) and panmixia ($\Delta\overline{AIC}$ = 3174.69) had the lowest support. The estimations of migration rates were independent from one another so the models were allowed to explore scenarios where initial migration was higher than contemporary migration. Only models where contemporary gene flow was higher than ancient gene flow were recovered in our iterations.

Figure 3. *Demographic modeling.* Demographic inference performed using fastsimcoal2. *Top:* comparison of AIC across different demographic scenarios among 10 bootstrap replicates of the SFS. *Bottom (from left to right)*: an illustration of the inferred demographic history and parameter estimate of the masked- and white-browed woodswallow complex, parameter estimates and variance using 50 bootstrap replicates of the SFS. Error bars reflect standard deviation from the mean.

A model with a higher contemporary migration rate from white-browed to masked woodswallow also consistently had higher likelihood in all ten bootstrap replicates. Parameter estimation suggests that the ancestral population size was 2N = 277,741 individuals, that of masked and white-browed woodswallows were 2N = 1,303,207 and 62,695 individuals, respectively. This suggests that the masked woodswallow experienced an expansion while the white-browed woodswallow had a much lower population size. This significant difference in effective population size was consistently recovered in the bootstrap iterations despite the species having similar levels of genetic diversity (θ). Ancient gene flow from masked to white-browed and white-browed to masked differed by orders of magnitude and were 2Nm = 42.09 migrants/generation and 0.17 migrants/generation, respectively. Contemporary estimates of the same pair of gene flow values, however, were of an equal order of magnitude though substantially different at 17.23 and 71.28 migrants/generation, respectively. While the estimates of the timing of population divergence and change in migration had high variances, the period of low ancient gene flow was less variable. The mean estimated time since divergence was 612,444 generations (±188,221 generations) and mean estimated time of increased gene flow was 595,543 generations (±182,143 generations). These convert to 1,090,150 and 1,060,067 years, respectively, assuming a generation time of 1.78 years (Higgins, 2006).

Outlier loci

Based on the 99.5 percentile cut-off we recovered 22 outlier loci. Of these 22, four had substantially higher F_{ST} than the others. Three of these loci were located at the distal end of chromosome 1A and the fourth was in chromosome 14. The three loci in chr1A were also adjacent to two other outliers above the 99.5 percentile cut off. This region encompasses at least 700kb at the distal end of the chr1A. The loci in chr1A also had high D_{XY} (Figure 4) suggesting they are potentially barrier loci whereas the locus in chr14 had low D_{XY}, which might suggest background selection. The genetic network generated for the 1.5 Mb region of chr1A (Figure 4) suggests higher derived diversity in the white-browed woodswallow and retained ancestral polymorphism for the masked woodswallow.

Figure 4. *Differentiation and diversity along chromosome 1A.* (A) Uncorrected genetic distance network of from the 5 outlier loci suggesting masked woodswallow has retained ancestral variants from the outgroups and white-browed woodswallow is likely the derived form. (B) Relative genetic differentiation across the length of chromosome 1A (C) Absolute genetic divergence (D_{XY}) across the length of chr1A suggesting both elevated relative and absolute measures of divergence (D) Nucleotide diversity (pi) across the length of chr1A suggesting higher diversity in the white-browed woodswallow (E) Genetic diversity (theta) across the length of chr1A suggesting slightly higher diversity in the white-browed woodswallow (F) Tajima's D across the length of chr1A showing overall negative values but not substantial difference between the two species

There were a total of 36 genes within 40kb on either side of the 22 outlier loci (Table S3). Of the outlier loci with the highest differentiation, only one (*SOX5*) is found within or adjacent to the 700kb in chr1A and two (*AXIN1* and *PDIA2*) are found adjacent to the chr14 outlier locus. The four outlier loci in chr1A span 696kb-1.36Mb while *SOX5* spans 737kb-1.2Mb and the outlier locus in chr14 falls directly within the *AXIN1* gene coordinate range. Both *SOX5* and *Axin1* are involved in negative regulation of the *Wnt/*β-catenin signaling pathway, which regulates feather development (Chang et al., 2004; Xie et al., 2020). Of the previously characterized plumage genes, only one had an associated ddRAD locus and it did not have high differentiation between species.

SLiM simulations

Loci under *balancing selection* and *partial recent sweeps* (both *hard* and *soft*) were qualitatively most similar to our outlier loci (Figure 5). Simulations under *neutrality* and *background selection* with gene flow were similar to non-outlier loci in our empirical data. Inconsistent with the empirical data, however, the neutral genetic diversity (both θ and π) is

different between the two populations. This is likely due to the estimated effective population size in the demographic model that we used for these simulations. All simulations under selective sweeps had high F_{ST} and D_{XY} as expected. Simulations with *historical sweeps* and *complete recent sweeps* had reduced genetic diversity and these were not consistent with the diversity of outlier loci.

Figure 5. *Simulations.* Evolutionary scenarios simulated using SLiM3 to test potential origin and maintenance of the outlier loci. Each scenario was run 100 times and the bars reflect the standard deviation of the estimates among the 100 iterations.

Discussion

Divergence in secondary sexual traits such as plumage is often associated with isolation via sexual selection. Counter examples are accruing in avian systems where plumage divergence is maintained despite genome-wide homogenization (e.g., Knief et al. 2019; Toews et al. 2016). Such systems often involve geographically localized hybridization or introgression that effectively homogenizes most of the genome. Here we studied Australia's masked and white-browed woodswallows *Artamus* spp in their broad sympatry over half of the Australian continent and between which phenotypic hybrids, though not unknown, are rare. We investigated the genomic basis to their high phenotypic divergence, which is sustained despite a lack of genetic divergence.

Our study using nuclear ddRAD markers further supports the lack of structure and divergence found in the mitochondrial DNA (Joseph et al., 2006). We used various methods to infer population structure and so capitalize on different attributes of the nuclear DNA data, yet still we found little evidence of allele frequency differences. The population network of the autosomal data, for example, shows a star-like pattern among the individuals of both species far more clearly than that of mtDNA (Figure 1). Next, the mtDNA data might suggest either no divergence has occurred or that the mitochondrial genome from one species has 'swamped' or replaced the other. The lack of shared haplotypes, however, is unexpected under a simple mtDNA capture. Although we still lack sampling of the masked woodswallow from Western Australia and may have missed within-species structure or unique alleles, a key new finding is that the nuclear genomes of the two species are largely indistinguishable throughout their sampled range (autosomal F_{ST} = 0.002, Z chromosome F_{ST} = 0.0003, Figure 2).

Despite being the only two species in *Artamus* exhibiting sexual dichromatism (Schodde & Mason, 1999), there is no elevated differentiation in the Z chromosome. This contrasts with other avian systems having high plumage differentiation coupled with low autosomal and high Z genetic differentiation (Dhami et al., 2016); the latter "large Z effect" is especially prevalent when adaptive differences are involved (Ellegren, 2009; Schield et al., 2021). Mank (2009) noted the complex connections between sex chromosomes and sexual dimorphism, however, and called for tests of assumptions such as we have presented here. The largely nomadic and in some years, strongly migratory, nature of these woodswallows likely allows for sympatry without strong competitive exclusion. This broad sympatry also means that occasional hybridization is not geographically restricted and likely happens throughout much of south-eastern Australia where the birds mostly are known to breed. This would facilitate introgression and genome homogenization across both species and concomitantly hinder the accumulation of genome-wide divergence.

Despite the observed lack of population structure, our demographic analyses have poor support for panmixia or recent divergence. Although all scenarios with divergence and some historical gene flow have good support, a scenario where migration rate increased in time had the highest support (Figure 3). Due to the stark difference in effective population size between the two species, we also see a shift in migration rate mostly from masked to white-browed in the past to its reciprocal, white-browed to masked, in more contemporary times. The estimated values of gene flow greatly exceed the number of migrants necessary to inhibit neutral divergence (Nm ~ 1; Barton, 2020). This is likely why we failed to detect population structure. The higher migration rate from white-browed into masked woodswallow is consistent with a scenario where the range of the masked woodswallow invaded the white-browed's range thereby retaining more of the native species' alleles in the invading species' genome (Joseph et al., 2006; McElroy et al., 2020; Rheindt & Edwards, 2011). This contrasts with the scenario where alleles of the population with the lower effective population size (i.e., white-browed woodswallow) are purged due to high deleterious load (Moran et al., 2021). That scenario would have resulted in higher migration in the opposite direction. The novel plumage morph then likely established rapidly in the period of lower gene flow but is evidently resistant to gene flow after the shift in migration rate. We acknowledge that the absolute parameter estimates should be interpreted cautiously. We tested 2-population models based on plumage rather than genetically defined populations. Rather, we emphasize the relative values not exact point estimates.

Extensive homogenization of genomes can reveal regions that may be involved in phenotypic divergence and speciation, and by extension the underlying evolutionary processes, which are usually masked by a 'noisy' landscape of divergence. Under the inferred demographic history, we simulated different evolutionary processes that could have led to the pattern we observed with the outlier loci. Our simulations suggest that the elevated F_{ST} and D_{XY} are likely due to the outliers in chr1A acting as barrier loci with restricted gene flow. The simulations that resulted in the elevated genetic diversity similar to the chr1A outliers are recent sweeps and balancing selection. Balancing selection would be more consistent with our demographic analysis, which had low support for recent divergence. In contrast to chr1A, the outlier in chr14 has elevated F_{ST} but low D_{XY}, which can result from long-term background selection resulting in reduced genetic diversity (see (Cruickshank & Hahn, 2014). Our simulations under different levels of background selection, however, did not result in low D_{XY} and high F_{ST} relative to neutral as predicted (Figure S9). A denser sampling of SNPs adjacent to the outliers would provide better signal from linkage to help distinguish between these different scenarios.

Importantly, we can begin to formulate hypotheses on the relevance of the genes associated with the outlier loci. Here, we focus on *SOX5* in chr1A and *Axin1* in chr14 where

the extreme outlier loci fall within the boundaries of the respective genes. Neither of these genes have been implicated in previous studies of plumage divergence but both are involved in negative regulation of the *Wnt/β-catenin* pathway, which is well-known in regulating feather development (Chang et al., 2004; Xie et al., 2020). There is no documented direct interaction between *SOX5* and *Axin1* but *SOX5* induces the production of *Axin2*, which is a negative regulator of the *Wnt*/β-catenin signaling pathway and functionally equivalent to *Axin1* (Dao et al., 2007; Quiroga et al., 2015). Overexpression of *Axin1* and *Axin2* reduces β-catenin levels and signaling. Another gene, *G3BP1*, found near an outlier locus in chr13 is also a negative regulator of the *Wnt*/β-catenin signaling pathway. *SOX5* additionally has been associated with strong phenotypic effects such as the pea-comb phenotype in chickens (Wright et al., 2009) and pigmentation through melanocyte development in mice and medaka fish (Nagao et al., 2018; Stolt et al., 2008). More precise delineation of boundaries of highly differentiated genomic regions and other potential candidate genes associated with phenotypic divergence requires denser sampling of SNPs, preferably through whole genome resequencing.

Future prospects

Having established that the nuclear genomes of white-browed and masked woodswallows show little divergence and no population structure, this species complex offers multiple avenues for investigation of the genetic architecture and maintenance of plumage divergence. Assembling and annotating a high-quality reference genome for the two species would place the variants in the correct physical order and may even reveal unique structural variation between the species. Furthermore, population-level whole genome resequencing would provide a more complete sampling of SNPs across the genome. This can validate the outliers presented in this study and potentially identify other candidate genes missed by the ddRAD sequencing. Additional sampling of masked woodswallow individuals from the part of their range where they do not overlap with the white-browed woodswallow may reveal lineage specific alleles that have not introgressed. This will be logistically non-trivial given the remoteness of their range in western parts of Australia, the often-nomadic movements of the birds and indeed that they are often seen high in the sky prohibiting sampling. Conversely, the species are increasingly kept in captivity where gene expression studies during feather development could test whether the genes found in this study play a direct role or interact with other candidate genes. Field studies on mate choice and sampling hybrids, again difficult logistically due to the birds' often unpredictable occurrences, would nonetheless disentangle the role of premating and postzygotic isolation in maintaining plumage divergence despite extensive introgression.

Systems such as we have examined here with striking phenotypic divergence yet little genomic divergence contrast with those having little to no discernible phenotypic divergence and high genomic divergence (i.e., "cryptic species" (Fišer et al., 2018)). Although these systems may be the exceptions to the rule, they bring light to the complexity of understanding speciation and species delimitation based on these metrics of divergence. Our study adds to those exemplifying divergence in only a few loci strongly influencing divergence of a secondary sexual trait such as plumage. We also show two further key findings: strong plumage divergence does not necessarily prevent whole-genome introgression and genes underlying a secondary sexual trait such as plumage are more likely to be found in autosomes rather than sex chromosomes. As we build a better understanding of the genetic architecture and evolution of these individual traits, we can piece together a more complete picture of the origin and evolution of biodiversity.

Acknowledgments

Specimens used in this study were collected under all appropriate ethics and scientific collecting permits over several decades and with the help of many landholders as acknowledged in an earlier paper (Joseph et al. 2006). For the present study, we would like to thank Sara Seibert for assisting with laboratory work and Christine Darwood, Graeme Chapman and Andrew Silcocks for kindly granting permission to use their photographs in the Supplementary Material.

Data accessibility

Raw Illumina reads are deposited in GenBank Accession [TBD]
Processed ddRAD fasta files are available in DRYAD doi [TBD].

References

Aguillon, S. M., Walsh, J., & Lovette, I. J. (2021). Extensive hybridization reveals multiple coloration genes underlying a complex plumage phenotype. *Proceedings. Biological Sciences / The Royal Society*, *288*(1943), 20201805.

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–410.

Barnard, H. G. (1944). Hybrid Wood-swallows. *The Emu*, *44*(2), 154–154.

Barton, N. H. (2020). On the completion of speciation. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *375*(1806), 20190530.

Bhatia, G., Patterson, N., Sankararaman, S., & Price, A. L. (2013). Estimating and interpreting FST: the impact of rare variants. *Genome Research*, *23*(9), 1514–1521.

Carnaby, I. C. (1965). The White-browed Wood-Swallow in Western Australia. *The Emu*. https://www.publish.csiro.au/mu/mu965074

Chang, C.-H., Jiang, T.-X., Lin, C.-M., Burrus, L. W., Chuong, C.-M., & Widelitz, R. (2004). Distinct Wnt members regulate the hierarchical morphogenesis of skin regions (spinal tract) and individual feathers. *Mechanisms of Development*, *121*(2), 157–171.

Cockayne, L., & Allan, H. H. (1926). The naming of wild hybrid swarms. *Nature*.

Comeron, J. M. (2017). Background selection as null hypothesis in population genomics: insights and challenges from Drosophila studies. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *372*(1736), 20160471.

Cowles, S. A., & Uy, J. A. C. (2019). Rapid, complete reproductive isolation in two closely related Zosterops White-eye bird species despite broadly overlapping ranges. *Evolution; International Journal of Organic Evolution*.

Coyne, J. A., & Orr, H. A. (2004). *Speciation* (Vol. 37). Sinauer Associates Sunderland, MA.

Cruickshank, T. E., & Hahn, M. W. (2014). Reanalysis suggests that genomic islands of speciation are due to reduced diversity, not reduced gene flow. *Molecular Ecology*, *23*(13), 3133–3157.

DaCosta, J. M., & Sorenson, M. D. (2014). Amplification biases and consistent recovery of loci in a double-digest RAD-seq protocol. *PloS One*, *9*(9), e106713.

Dao, D. Y., Yang, X., Chen, D., Zuscik, M., & O'Keefe, R. J. (2007). Axin1 and Axin2 are regulated by TGF- and mediate cross-talk between TGF- and Wnt signaling pathways. *Annals of the New York Academy of Sciences*, *1116*, 82–99.

Dhami, K. K., Joseph, L., Roshier, D. A., & Peters, J. L. (2016). Recent speciation and elevated Z-chromosome differentiation between sexually monochromatic and dichromatic species of Australian teals. *Journal of Avian Biology*, *47*(1), 92–102.

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* . https://academic.oup.com/bioinformatics/article-abstract/26/19/2460/230188

Edgar, Robert C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, *32*(5), 1792–1797.

Ellegren, H. (2009). The different levels of genetic diversity in sex chromosomes and autosomes. *Trends in Genetics: TIG*, *25*(6), 278–284.

Excofffier, L., Marchi, N., Marques, D. A., Matthey-Doret, R., Gouy, A., & Sousa, V. C. (2021). fastsimcoal2: demographic inference under complex evolutionary scenarios. *Bioinformatics* . https://doi.org/10.1093/bioinformatics/btab468

Fišer, C., Robinson, C. T., & Malard, F. (2018). Cryptic species as a window into the paradigm shift of the species concept. *Molecular Ecology*, *27*(3), 613–635.

Funk, E. R., & Taylor, S. A. (2019). High-throughput sequencing is revealing genetic associations with avian plumage color. *The Auk*, *136*(4). https://doi.org/10.1093/auk/ukz048

Haller, B. C., & Messer, P. W. (2019). SLiM 3: Forward Genetic Simulations Beyond the Wright-Fisher Model. *Molecular Biology and Evolution*, *36*(3), 632–637.

Higgins, P. (2006). *Handbook of Australian, New Zealand and Antarctic Birds: Volume 7. Part A. Boatbills to Larks*. Oxford University Press.

Huson, D. H. (1998). SplitsTree: analyzing and visualizing evolutionary data. *Bioinformatics* , *14*(1), 68–73.

Jakobsson, M., & Rosenberg, N. A. (2007). CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* , *23*(14), 1801–1806.

Johnstone, R., & Storr, G. M. (2004). *Handbook of Western Australian birds. Volume Two: Passerines*. Western Australia Museum, Perth.

Joseph, L., Wilke, T., Ten Have, J., & Terry Chesser, R. (2006). Implications of mitochondrial DNA polyphyly in two ecologically undifferentiated but morphologically distinct migratory birds, the masked and white-browed woodswallows Artamus spp. of inland Australia. *Journal of Avian Biology*, *37*(6), 625–636.

Kim, K.-W., Jackson, B. C., Zhang, H., Toews, D. P. L., Taylor, S. A., Greig, E. I., Lovette, I. J., Liu, M. M., Davison, A., Griffith, S. C., Zeng, K., & Burke, T. (2019). Genetics and evidence for balancing selection of a sex-linked colour polymorphism in a songbird. *Nature Communications*, *10*(1), 1852.

Knief, U., Bossu, C. M., Saino, N., Hansson, B., Poelstra, J., Vijay, N., Weissensteiner, M., & Wolf, J. B. W. (2019). Epistatic mutations under divergent selection govern phenotypic variation in the crow hybrid zone. *Nature Ecology & Evolution*, *3*(4), 570–576.

Leigh, J. W., & Bryant, D. (2015). popart: full-feature software for haplotype network construction. *Methods in Ecology and Evolution / British Ecological Society*, *6*(9), 1110–1116.

Lenz, M. (2019). Notes on breeding of masked and white-browed woodswallows in the 2018/19 season in the COG area of interest. *Canberra Bird Notes*, *44*, 152–162.

Mank, J. E. (2009). The evolution of heterochiasmy: the role of sexual selection and sperm competition in determining sex-specific recombination rates in eutherian mammals. *Genetics Research*, *91*(5), 355–363.

Mason, N. A., & Bowie, R. C. K. (2020). Plumage patterns: Ecological functions, evolutionary origins, and advances in quantification. *The Auk*, *137*(4). https://doi.org/10.1093/auk/ukaa060

McElroy, K., Black, A., Dolman, G., Horton, P., Pedler, L., Campbell, C. D., Drew, A., & Joseph, L. (2020). Robbery in progress: Historical museum collections bring to light a mitochondrial capture within a bird species widespread across southern Australia, the Copperback Quail-thrush Cinclosoma clarum. *Ecology and Evolution*, *10*(13), 6785–6793.

McGill, A. R. (1944). Hybridism in Wood-swallows. *The Emu*. https://www.publish.csiro.au/mu/MU944153a

Metzler, D., Knief, U., Penalba, J., & Wolf, J. B. W. (2020). Frequency dependent sexual selection, mating trait architecture and preference function govern spatio-temporal hybrid zone dynamics. *BioRxiv*. https://www.biorxiv.org/content/10.1101/2020.03.10.985333v4.abstract

Moran, B. M., Payne, C., Langdon, Q., Powell, D. L., Brandvain, Y., & Schumer, M. (2021). The genomic consequences of hybridization. *ELife*, *10*. https://doi.org/10.7554/eLife.69016

Nagao, Y., Takada, H., Miyadai, M., Adachi, T., Seki, R., Kamei, Y., Hara, I., Taniguchi, Y., Naruse, K., Hibi, M., Kelsh, R. N., & Hashimoto, H. (2018). Distinct interactions of Sox5 and Sox10 in fate specification of pigment cells in medaka and zebrafish. *PLoS Genetics*, *14*(4), e1007260.

Omland, K. E., Baker, J. M., & Peters, J. L. (2006). Genetic signatures of intermediate divergence: population history of Old and New World Holarctic ravens (Corvus corax). *Molecular Ecology*, *15*(3), 795–808.

Patten, M. A., & Unitt, P. (2002). Diagnosability Versus Mean Differences of Sage Sparrow Subspecies. *The Auk*, *119*(1), 26–35.

Paxton, E. H. (2009). The utility of plumage coloration for taxonomic and ecological studies. *The Open Ornithology Journal*, *2*(1), 17–23.

Price, T. (2008). *Speciation in birds*. Roberts and Co.

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, *155*(2), 945–959.

Quiroga, A. C., Stolt, C. C., Diez del Corral, R., Dimitrov, S., Pérez-Alcalá, S., Sock, E., Barbas, J. A., Wegner, M., & Morales, A. V. (2015). Sox5 controls dorsal progenitor and interneuron specification in the spinal cord. *Developmental Neurobiology*, *75*(5), 522–538.

Rheindt, F. E., & Edwards, S. V. (2011). Genetic introgression: an integral but neglected component of speciation in birds. *The Auk*, *128*(4), 620–632.

Sætre, G.-P., Moum, T., Bureš, S., Král, M., Adamjan, M., & Moreno, J. (1997). A sexually selected character displacement in flycatchers reinforces premating isolation. *Nature*, *387*(6633), 589–592.

San-Jose, L. M., Ducrest, A.-L., Ducret, V., Simon, C., Richter, H., Wakamatsu, K., & Roulin, A. (2017). MC1Rvariants affect the expression of melanocortin and melanogenic genes and the association between melanocortin genes and coloration. *Molecular Ecology*, *26*(1), 259–276.

Schield, D. R., Scordato, E. S. C., Smith, C. C. R., Carter, J. K., Cherkaoui, S. I., Gombobaatar, S., Hajib, S., Hanane, S., Hund, A. K., Koyama, K., Liang, W., Liu, Y., Magri, N., Rubtsov, A., Sheta, B., Turbek, S. P., Wilkins, M. R., Yu, L., & Safran, R. J. (2021). Sex-linked genetic diversity and differentiation in a globally distributed avian species complex. *Molecular Ecology*, *30*(10), 2313–2332.

Schodde, R., & Mason, I. J. (1999). *The Directory of Australian Birds: Passerines*. CSIRO Publishing.

Semenov, G. A., Linck, E., Enbody, E. D., Harris, R. B., Khaydarov, D. R., Alström, P., Andersson, L., & Taylor, S. A. (2021). Asymmetric introgression reveals the genetic architecture of a plumage trait. *Nature Communications*, *12*(1), 1019.

Servedio, M. R., Van Doorn, G. S., Kopp, M., Frame, A. M., & Nosil, P. (2011). Magic traits in speciation: "magic" but not rare? *Trends in Ecology & Evolution*, *26*(8), 389–397.

Stolt, C. C., Lommes, P., Hillgärtner, S., & Wegner, M. (2008). The transcription factor Sox5 modulates Sox10 function during melanocyte development. *Nucleic Acids Research*, *36*(17), 5427–5440.

Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, *123*(3), 585–595.

Toews, D. P. L., Taylor, S. A., Vallender, R., Brelsford, A., Butcher, B. G., Messer, P. W., & Lovette, I. J. (2016). Plumage Genes and Little Else Distinguish the Genomes of Hybridizing Warblers. *Current Biology: CB*, *26*(17), 2313–2318.

Toomey, M. B., Marques, C. I., Andrade, P., Araújo, P. M., Sabatino, S., Gazda, M. A., Afonso, S., Lopes, R. J., Corbo, J. C., & Carneiro, M. (2018). A non-coding region near Follistatin controls head colour polymorphism in the Gouldian finch. *Proceedings.*

    Biological Sciences / The Royal Society, *285*(1888).
    https://doi.org/10.1098/rspb.2018.1788

Twyman, H., Prager, M., Mundy, N. I., & Andersson, S. (2018). Expression of a carotenoid-modifying gene and evolution of red coloration in weaverbirds (Ploceidae). *Molecular Ecology*, *27*(2), 449–458.

Winger, B. M., & Bates, J. M. (2015). The tempo of trait divergence in geographic isolation: avian speciation across the Marañon Valley of Peru. *Evolution; International Journal of Organic Evolution*, *69*(3), 772–787.

Wright, D., Boije, H., Meadows, J. R. S., Bed'hom, B., Gourichon, D., Vieaud, A., Tixier-Boichard, M., Rubin, C.-J., Imsland, F., Hallböök, F., & Andersson, L. (2009). Copy number variation in intron 1 of SOX5 causes the Pea-comb phenotype in chickens. *PLoS Genetics*, *5*(6), e1000512.

Xie, W. Y., Chen, M. J., Jiang, S. G., Yan, H. C., Wang, X. Q., & Gao, C. Q. (2020). The Wnt/β-catenin signaling pathway is involved in regulating feather growth of embryonic chicks. *Poultry Science*, *99*(5), 2315–2323.