LRT: T Cell Trajectory Inference by Integrative Analysis of Single-Cell TCR-seq and RNA-seq data

Juan Xie[1], Gang Xin[2,3], Qin Ma[3,4] and Dongjun Chung[3,4, *]

1 The Interdisciplinary Program in Biostatistics, The Ohio State University, Columbus, Ohio, USA.

2 Department of Microbial Infection and Immunity, The Ohio State University, Columbus, Ohio, USA.

3 Pelotonia Institute for Immuno-Oncology, The James Comprehensive Cancer Center, The Ohio State University, Columbus, Ohio, USA.

4 Department of Biomedical Informatics, The Ohio State University, Columbus, Ohio, USA.

* chung.911@osu.edu

Abstract

Single-cell RNA sequencing (scRNA-seq) data has been widely used for cell trajectory inference, with the assumption that cells with similar expression profiles share the same differentiation state. However, the inferred trajectory may not reflect true clonal relationships among cells. Single-cell T cell receptor sequencing (scTCR-seq) data provides invaluable insights into the clonal relationship among cells, yet it lacks functional characteristics. Therefore, scRNA-seq and scTCR-seq data complement each other in improving trajectory inference, where a reliable computational tool is still missing. We developed LRT, a computational framework for the integrative analysis of scTCR-seq and scRNA-seq data for T cell trajectory inference. Specifically, LRT utilizes the TCR sequence information to identify clonally related cells and then uses the transcriptomics information from scRNA-seq data to construct clonotype-level cell trajectories. LRT provides a comprehensive analysis workflow, including preprocessing, cell trajectory clustering, pseudotime inference, and marker gene identification. We illustrated its utility using scRNA-

seq and scTCR-seq data of CD4$^+$ T cells with acute lymphocytic choriomeningitis virus infection, where

we could identify cell trajectories that cannot be revealed solely based on scRNA-seq data. Our downstream

analyses showed that (i) these trajectories are involved in distinct functional roles; (ii) the expression

patterns of their marker genes over the estimated pseudotime nicely coincide with the Th1/Tfh biology that

is well established for the CD4$^+$ T cell differentiation; and (iii) the higher level of TCR sequence similarities

was observed within each cluster, compared to between clusters. The LRT framework was implemented as

an R package 'LRT', and it is now publicly accessible at https://github.com/JuanXie19/LRT. In addition, it

provides two Shiny apps 'shinyClone' and 'shinyClust' that allow users to interactively explore

distributions of clonotypes, conduct repertoire analysis, implement clustering of cell trajectories, and

predict cell trajectory cluster marker genes.

Author Summary

Understanding the dynamic changes behind biological processes is important for determining molecular

mechanisms underlying normal tissue formulation, developmental disorders and pathologies. Usually, a

biological process can be characterized by identifying a trajectory, a path that goes through the various

cellular states associated with the process. Since cells in different states may express different sets of genes,

researchers often infer cell trajectory via capturing transcriptomics changes. Dozens of methods have been

developed for cell trajectory inference, and scRNA-seq data is predominantly utilized. However, methods

based only on scRNA-seq data cannot tell us if cells from the same trajectory come from the same clone or

not. T cells play a key role in the immune system, and their high antigen recognition specificity is largely

determined by their TCR sequences. Thanks to the advent of scTCR-seq technology, people can identify

the group of cells coming from the same clone. This paper describes our novel computational framework,

namely LRT, and demonstrates that by complementing scRNA-seq data with the clonal information from

scTCR-seq data using LRT, we are able to identify cell trajectories that cannot be revealed solely based on

scRNA-seq data.

Introduction

Cell trajectory inference aims to understand how cells differentiate, which is a long-standing problem in

developmental biology. Understanding how progenitor cells transform into specified functional cells can

provide valuable insights into the molecular mechanisms underlying normal tissue formulation, as well as

developmental disorders and pathologies [1]. Single-cell RNA-seq (scRNA-seq) data has been widely used

to investigate cell trajectories. Assuming that cells in different states express different sets of marker genes,

we may order cells along a differentiation trajectory via capturing differentiated transcriptional activities

[2]. Based on this rationale, various computational and statistical approaches have been proposed for this

purpose, namely trajectory/pseudotime analysis, where well known examples include Monocle [2] and

Slingshot [3] (please see Saelens et al. [4] for a comprehensive review). Specifically, Monocle builds a

minimum spanning tree (MST) on cells in a low-dimensional independent component space and uses the

longest path through the MST as the backbone for pseudotime ordering, while branches were handled by

the PQ tree. In the case of Slingshot, in its first stage, it also uses MST to determine the global cell lineage

structure. Yet instead of building MST on individual cells, Slingshot constructs an MST on cluster centers,

which makes it more robust to noises. Then, in its second stage, it uses a novel method called simultaneous

principal curves to obtain smooth branching trajectories. While these approaches have been shown to be

useful, they still suffer from intrinsic limitations. Specifically, it assumes that cells with similar expression

profiles share similar developmental stages and may come from the same lineage, which might not be

always true [1, 5]. Besides, in the absence of clonal information, the inferred trajectory may not reflect the

true clonal relationship between cells [1].

The recent advent of single-cell T cell receptor (TCR) sequencing (scTCR-seq) technology has

enabled the use of TCR sequences as unique 'barcodes' to identify clonally related cells. This is because the

cells with identical paired TCR sequences usually arise from the same T cell clone due to the high

dimensionality of TCR sequence space [6]. Examining the abundance/proportion of clonal cells and their

changes greatly benefits our understanding of adaptive immune response in health and disease [7, 8];

therefore, there has been a rapid accumulation of scTCR-seq data in recent years. Accordingly, several

computational and statistical approaches have also been developed, where examples include Immunarch [9]

and scRepertoire [10]. However, currently available approaches mostly focus on TCR repertoire analysis,

such as clonality, repertoire overlap, repertoire diversity, and gene usage analyses. The utilization of

scTCR-seq data for cell trajectory analysis is still limited. In addition, although TCR data provides

invaluable insights into the breadth of the antigenic response and clonal relationships between cells, it does

not offer information about the functional characteristics of the T cells. Hence, integrating scTCR-seq and

scRNA-seq can help elucidate the interplay between TCR sequences and T cell phenotypes. For instance,

Zhang et al. [11] observed that TCR similarity constrained the similarity of T cell phenotypes, and TCRs

that are more similar to the center TCRs have stronger antigen binding affinity. However, despite such

potential to improve T cell trajectory inference, to the best of our knowledge, a computational approach to

integrate scRNA-seq and scTCR-seq data for cell trajectory inference is missing in the literature. This is

partially due to the statistical and computational challenges related to the high dimensionality of the

clonotype space. Specifically, for a given human subject, 10^{13} TCR sequences are typically observed while

each clonotype may only span a very limited number of cells. Hence, for effective cell trajectory inference

and integration of scTCR-seq and scRNA-seq data, we need a cleverly designed computational framework

that can handle these challenges.

In this paper, we propose LRT (**L**ineage inference by integrative analysis of sc**R**NA-seq and

sc**T**CR-seq data), a rigorous computational framework for the integrative analysis of scTCR-seq and

scRNA-seq data for cell trajectory inference. LRT addresses the complexity and the high dimensionality of

scTCR-seq data through a trajectory clustering algorithm and effective information sharing between

scTCR-seq and scRNA-seq data. In addition, to support researchers in utilizing the proposed LRT

framework for their studies, we developed an R package 'LRT', which is now publicly accessible at

https://github.com/JuanXie19/LRT. We also developed two Shiny apps, namely 'shinyClone' and

'shinyClust', and they allow researchers to interactively implement the complete LRT analysis workflow,

including exploratory analysis, cell trajectory inference, and marker gene identification. These Shiny apps

are provided as part of the R package 'LRT'.

Materials and Methods

The workflow of the 'LRT' framework is shown in **Fig. 1**, which starts from the integration of scTCR-seq

and scRNA-seq data (**Fig. 1A**). Using this integrated dataset, LRT infers cell trajectory by implementing

the following four steps. In the first step, it estimates clonotype-level trajectories by building an MST based

on the cells sharing the same clonotype (**Fig. 1B**). In the second step, LRT identifies clusters of clonotype-

level trajectories using the stochastic block model (SBM) based on the dynamic time warping (DTW)

distances (**Fig. 1C**). In the third step, a principal curve is fitted using the cells belonging to the same

trajectory cluster, to get the final trajectories and infer the pseudotime (**Fig. 1D**). Finally, differential

expression (DE) analysis is implemented to identify genes characterizing each trajectory cluster, namely

trajectory cluster markers (**Fig. 1E**).

Figure 1. The framework of LRT. (A) The preprocessed scRNA-seq and scTCR-seq data are integrated

based on cell barcodes. **(B)** Clonotype-level trajectories are first obtained by constructing minimum

spanning trees (MST). **(C)** Similar trajectories are grouped using the stochastic block model (SBM), where

similarity is evaluated based on the dynamic time warping (DTW) distance. **(D)** For each trajectory cluster,

cell pseudotimes are estimated by fitting a principal curve on the Uniform Manifold Approximation and

Projection (UMAP) space. **(E)** The relationship between read count and pseudotime is modeled by negative

binomial generalized additive model (NB-GAM), and the Wald test is used to determine marker genes.

Trajectory inference

122 The core of LRT lies in its trajectory clustering and characterization through the integration of scTCR-seq

123 and scRNA-seq data. It consists of four key steps, including clonotype-level trajectory estimation, trajectory

124   clustering, pseudotime inference, and marker identification (DE analysis). Each step is described in detail

125   below.

Clonotype-level trajectory estimation

Given the integrated scRNA-seq and scTCR-seq data (see the Section 'Software implementation' for more

details), LRT infers a trajectory for each clonotype. Specifically, a weighted graph is constructed for each

group of cells sharing the same TCR sequence. Here nodes are cell-cluster-specific centers of these cells,

while edge weights represent the Euclidean distances between the nodes on the Uniform Manifold

Approximation and Projection (UMAP) space [12]. Note that both cell clusters and UMAP in this step are

defined based on scRNA-seq data. An MST [13] is then constructed, and trajectories are interpreted as

ordered cell clusters by tracing paths through the MST.

Trajectory clustering

After obtaining the clonotype-level trajectories, LRT conducts a clustering analysis to group similar

trajectories, where similarity is defined based on DTW distances [5] between trajectories. DTW distance is

a measure to reflect the similarity between two temporal sequences, which is invariant to warping and thus

is particularly useful when the two sequences vary in speed. Here we treat each trajectory as time series

data, with cell clusters as time points and UMAP coordinates as measurements under each time point. Let's

denote two trajectories $S = (s_1, s_2, \ldots, s_i, \ldots, s_q)$ and $T = (t_1, t_2, \ldots, t_j, \ldots, t_r)$, where q and r are the

numbers of "time points", respectively. We first build the distance matrix $C \in \mathbb{R}^{q \times r}$ to represent pairwise

distances between S and T, where each element of C, $c(i, j) = \|s_i - t_j\|$, $i \in [1:q], j \in [1:r]$. Then, we

define a warping path as a sequence of points $p = (p_1, \ldots, p_Q)$ with $p_l = (i_l, j_l) \in [1:q] \times [1:r]$ for $l \in$

$[1:Q]$ to represent a possible alignment between S and T. The associated total cost/distance for p is

$$c_p(S, T) = \sum_{l=1}^{Q} c(s_{i_l}, t_{j_l}).$$

146 Finally, the DTW distance is defined as the cost associated with the optimal warping path, i.e.,

$$DTW(S,T) = c_{p^*}(S,T) = \min_{p \in P^{q \times r}} \{c_p(S,T)\},$$

where $P^{q \times r}$ is the set of all possible warping paths.

Based on this DTW distance matrix, LRT applies an SBM [14] to identify trajectory clusters (i.e.,

community detection). SBM is a generative model for random graphs, which is widely employed for

recovering (latent) community structures in network/graph data [15]. Let's denote n as the total number of

cells. To apply SBM in our case, we convert the DTW distance matrix to an adjacency matrix of a k-nearest

neighbor graph, denoted as A, where we set $k = \sqrt{n}$ by adopting the widely used heuristics [15, 16]. Given

$A = [A_{ij}] \in \mathbb{R}^{n \times n}$ denoting the binary adjacency matrix of a random graph, with elements A_{ij} indicating

the presence or absence of an undirected edge between nodes i and j, we assume that the absence or

presence of an edge between each pair of nodes i and j follows a Bernoulli distribution, i.e.,

$$A_{ij} \sim Bernoulli(p_{ij}), \; p_{ij} \in (0,1) \text{ for } 1 \leq i < j \leq n.$$

158 SBM assumes that there is a latent connectivity matrix $\Theta \in \mathbb{R}^{b \times b}$, $b \ll n$, and a map $Z: \{1, \dots, n\} \rightarrow$

159 $\{1, \dots, b\}$, such that

$$p_{ij} = \theta_{z(i),z(j)}.$$

161 SBM groups n nodes into b groups, with group labels given by Z, and we call these groups communities.

162 A variational Expectation-Maximization algorithm [17] is used for parameter estimation. After fitting

163 SBMs with a different number of clusters, the final number of clusters is determined using the maximum

164 integrated completed likelihood (ICL) criterion [18].

Pseudotime estimation

As the final step of trajectory inference, LRT obtains the final trajectory by fitting a principal curve [19] on

the UMAP coordinates of cells. Let us denote a random vector representing the UMAP coordinates for cells

from the same trajectory cluster as $X \in \mathbb{R}^2$. In addition, denote h as a one-dimensional smooth unit-speed

curve in $\mathbb{R}^2$ parameterized by a single variable t. In the principal curve estimation, we iterate the following

the projection step and the regularized expectation step:

(i) Projection step: Project all cells onto the curve $h^{(l-1)}$ and calculate the arc length from the beginning of

the curve to each point's projection, $t_i^{(l)} = t_{h^{(l-1)}}(x_i)$, where the projection $t_h \colon \mathbb{R}^2 \to \mathbb{R}^1$ is defined as

$$t_h(x) = \sup_t \{t \colon \| x - h(t) \| = \inf_u \| x - h(u) \|\}$$

174   (ii) Regularized expectation step: Update the curve h by minimizing the distance between data points x_i and

175   those projected onto the curve, $h(t_i^{(l)})$.

176   Here, we interpret the curve $h(x)$ as the final trajectory for the given cell trajectory cluster and the arc

177   length $t_i = t_h(x_i)$ as pseudotime for cell i.

Differential expression analysis

While the algorithms described above allow us to infer cell trajectories, it is critical to identify genes

characterizing these trajectories to fully understand the biological relevance of these trajectories. To

facilitate the interpretation of the inferred trajectories, LRT implements DE analysis for each trajectory

cluster. First, for each trajectory cluster, it infers a smooth function of the normalized gene read counts on

the pseudotime, using a negative binomial generalized additive model (NB-GAM) [20, 21]. To be specific,

for the v-th trajectory cluster with n_v cells, let Y_{gi} be the read counts for gene $g \in \{1, \dots, G\}$ and cell $i \in$

$\{1, \dots, n_v\}$, the NB-GAM assumes that

$$Y_{gi} \sim NB(\mu_{gi}, \phi_g)$$

$$\log(\mu_{gi}) = f_g(t_i) + \log(N_i)$$

189   where μ_{gi} are cell- and gene-specific means, ϕ_g are gene-specific dispersion parameters, N_i is cell-specific

190   offsets that account for sequencing depth, and f_g is the smoothing spline for gene g. Here we use a cubic

191   spline function with 6 knots for f_g to balance between the goodness of fit and running time. Finally, a Wald

192   test is used to determine DE genes [21], followed by Bonferroni correction for the multiple testing

193   adjustment.

Software implementation

The aforementioned LRT framework and the whole analysis workflow were implemented as an R package

'LRT' and it is now publicly accessible at https://github.com/JuanXie19/LRT. The LRT software takes both

scRNA-seq and scTCR-seq data as input. It accepts scRNA-seq data in the form of a Seurat object [22].

Such integration with Seurat not only allows LRT to be easily integrated into an existing scRNA-seq

analysis workflow, but also utilizes Seurat's powerful features, e.g., correcting batch effects and variations

due to technical factors (e.g., sequencing depth) [22, 23]. The Seurat object needs to contain clustering

labels and UMAP coordinates for each cell. In the case of scTCR-seq data, it requires a data frame with at

least two columns: one column with cell identifiers and another column with TCR sequence information,

for which users can provide amino acid or nucleotide sequences of TCR α, β, or paired α-β chain CDR3

region.

Data preprocessing

207   LRT integrates scTCR-seq data with scRNA-seq data by attaching the TCR sequence information to the

208   metadata of the Seurat object. LRT also provides some basic quality control functionalities, e.g., checking

209   whether the cell identifiers from the Seurat object match those in the TCR data frame. After the integration,

210   users will get an S4 object containing the updated Seurat object, which is the input for clonotype exploratory

211   analysis and trajectory inference.

Shiny apps for interactive visualization and analysis: shinyClone and shinyClust

We developed two Shiny apps, *shinyClone*, and *shinyClust*, for interactive and dynamic data exploration and integrative analysis of scRNA-seq and scTCR-seq data.

shinyClone

shinyClone is a Shiny app for exploratory analysis of clonotypes (**Fig. 2**). It provides commonly used repertoire analysis functionalities and lets users intuitively explore how each clonotype is distributed on the reduced dimensional space generated based on scRNA-seq data. *shinyClone* consists of two tabs, one for exploratory analysis of clonotype distribution (tab 'Clonotype') (**Fig. 2A**) and the other for repertoire analysis (tab 'Repertoire') (**Fig. 2B**). First, on the 'Clonotype' tab, users can find a table of clonotypes, along with the relevant information, including (i) how many cells are associated with each clonotype; and (ii) how these cells are distributed among groups (e.g., cell clusters, samples, conditions, or any other groups as available in the metadata of Seurat object) (**Fig. 2A**, left). Users can sort the table by clicking each column name, e.g., to check the most/least abundant clonotypes. By clicking each clonotype in this table, the user can check the distribution of cells associated with each clonotype on the reduced dimensional space, for which the user can choose among UMAP, t-SNE, or principal component analysis (PCA) for the dimension reduction (**Fig. 2A**, middle). A cell density plot is also displayed for clones spanning more than 10 cells (**Fig. 2A**, right). Both plots can be downloaded in the PDF file format by clicking the "Download" button. Second, the 'Repertoire' tab provides several common repertoire analysis functions using the R package 'scRepertoire' [10], where examples include clonal diversity, clonal homeostasis, clonal proportion, and clonal overlap analyses (**Fig. 2B**).

Figure 2. *shinyClone,* **a Shiny app for exploratory analysis of clonotypes. (A)** The 'Clonotype' tab allows users to implement various exploratory analyses of clonotype distributions. **(B)** The 'Repertoire' tab provides various TCR repertoire analysis functionalities.

shinyClust

shinyClust is another Shiny app that allows users to interactively and dynamically implement the trajectory clustering analysis and marker identification (**Fig. 3**). *shinyClust* consists of three tabs, 'Clustering exploration' (**Fig. 3A**), 'Trajectory clustering' (**Fig. 3B**), and 'DE analysis' (**Fig. 3C**). First, the 'Clustering

exploration' tab allows users to implement trajectory clustering interactively (**Fig. 3A**). This page shows

either ICL or silhouette scores under the current setting, to assist users in evaluating and determining the

optimal number of clusters. While SBM is set to be a default here, other clustering algorithms are also

provided for flexibility, including hierarchical and partitional clustering. We used the R package 'sbm' [24]

for the SBM implementation and the R package 'dtwclust' [25] for the hierarchical and partitional clustering

implementations. Users can also interactively tune parameters for each clustering algorithm. Second, on the

'Trajectory clustering' tab, upon choosing a particular cluster to display, *shinyClust* updates (i) a trajectory

plot, which shows clonotype trajectories corresponding to the trajectory cluster on the UMAP plot (**Fig. 3B**,

the plot on the left, titled 'Trajectories'); and (ii) a cell distribution plot, which shows the cells associated

with this trajectory cluster on the UMAP plot (**Fig. 3B**, the plot on the right, titled 'Cell distribution'). All

the plots can also be downloaded in the PDF file format by clicking the "Download" button. Finally, in the

'DE analysis' tab, users can implement DE analysis, which fits the NB-GAM model and conducts the Wald

test in the backend using the R package 'tradeSeq' [21] (**Fig. 3C**). Here, *shinyClust* shows a table of

predicted DE genes with both raw and Bonferroni-adjusted *p*-values, as well as a heatmap of DE gene

expression values. If users are interested in how the expression values of a particular gene vary along

pseudotime, they can click the corresponding row in the table and the page will show a scatter plot of log

read count versus pseudotime, along with the fitted smoothing spline curve.

Figure3. *shinyClust*, **a Shiny app for trajectory clustering and marker identification**. **(A)** The 'Clustering exploration' tab shows the summary of trajectory clustering. **(B)** The 'Trajectory clustering' tab

263   shows the trajectory clustering results. **(C)** The 'DE analysis' tab allows users to identify marker genes for

264   each trajectory cluster and their changes as a function of the pseudotime.

Results

Exploratory analysis of scRNA-seq and scTCR-seq data

To demonstrate the utility of LRT, we analyzed the scTCR-seq and scRNA-seq data generated from the

antigen-specific CD4$^+$ T cells obtained from five mice during acute lymphocytic choriomeningitis virus

infection [26] (GEO accession number: GSE158896). Seurat was used to preprocess scRNA-seq data and

implement UMAP dimension reduction and cell clustering. Specifically, for each sample, we filtered out

doublets and dead cells (those with a percentage of mitochondrial genes > 10%), calculated cell cycle scores

for the remaining cells and regressed out, and normalized using the *SCTransform* approach [23]. Then, the

processed datasets from five samples were integrated using the Seurat data integration workflow. Finally,

PCA and UMAP dimension reduction were conducted, and then cells were clustered using the Louvain

algorithm [27], for which the first 20 principal components and default resolution were used. We obtained

11 cell clusters and annotated them based on the expert knowledge (**Fig. 4**), including T central memory

precursor cells (Tcmp) and the cells corresponding to different differentiation stages leading to the type 1

helper T cells (Th1). Specifically, based on the expression of well-known T cell subtype marker genes [26],

we annotated cell clusters 0, 1, 4, 9, and 10 as Tcmp, cell clusters 5 and 6 as pre-Th1, cell clusters 2 and 8

as Th1-intermediate (Th1-inter), and cell clusters 3 and 7 as Th1.

For scTCR-seq data, only the chains that were annotated as full-length and functional were retained.

Besides, to avoid any contamination from sorting, only clones having at least two cells from a single sample

were considered. After filtering, the TCR sequences from five samples were combined into a single data

frame using the R package 'scRepertoire' [10]. The resulting scTCR-seq data was then integrated with the

above-mentioned Seurat object for the remaining data analyses. Exploratory analysis for the clonotypes

was conducted using the *shinyClone* app. There were 516 clonotypes in this dataset and, among those, 69

spanned all four annotated cell types, while 182 of them spanned only one cell type. The top two most

abundant clonotypes (defined by CDR3 β chain amino acid sequence) were 'CASSFAGGASYEQYF' and

'CAWDRDTEVFF', which appeared in 863 and 762 cells, respectively. Most of the cells with the

clonotype 'CASSFAGGASYEQYF' were Tcmp cells and located at the bottom of the UMAP plot. The

cells with the clonotype 'CAWDRDTEVFF' evenly spanned all four cell types and these cells mainly

occupied the upper part of the UMAP plot.

Cell trajectory inference

We first inferred cell trajectories based only on scRNA-seq data using the Slingshot algorithm [3] (**Fig. 4**).

When we used the Seurat cell clusters (using cluster 0 as the root), Slingshot predicted three trajectories

(**Fig. 4A**), including:

- Trajectory I: 0—10—9—5—2—8—7—6

- Trajectory II: 0—10—9—5—2—3

- Trajectory III: 0—10—1—4

When we inferred cell trajectories based only on scRNA-seq data but using four annotated cell types,

Slingshot identified a single trajectory of Tcmp – pre-Th1 – Th1-intermediate – Th1 (**Fig. 4B**), which

coincides with the known trajectory of Th1 cell differentiation.

Figure 4. Trajectories inferred based only on scRNA-seq data of the antigen-specific CD4[+] T cells visualized on the UMAP. (A) Cell trajectory inferred based only on scRNA-seq data using the Slingshot algorithm. Dots represent cells and curves represent the inferred trajectories. Here we considered Seurat cell clusters, which are denoted by colors and numbers (0 – 10) in the plot. Roman numbers within circles represent trajectory indices. **(B)** Cell trajectory inferred when we considered the expert-annotated cell types. Cells are colored and labeled accordingly in the plot (salmon for Tcmp, light violet for pre-Th1, green for Th1-inter, and Iris blue for Th1), and the black curve represents the inferred trajectory.

Next, we inferred cell trajectories by integrating scRNA-seq and scTCR-seq data using the LRT framework (with SBM as a clustering algorithm). In this case, we discovered five sub-groups of trajectories constituting this trajectory (**Fig. 5A**, **Fig. S1**). To evaluate whether this trajectory clustering makes sense from the perspective of TCR sequences, we first checked similarities in TCR sequences within vs. between the trajectory clusters. Specifically, we obtained numeric embeddings for TCR sequences using the tessa algorithm [11] and calculated the Euclidean distances between each pair of TCR sequences. We found that within-cluster distances were smaller than or comparable to between-cluster distances (**Fig. 5B**). Among those, within-cluster distances are significantly smaller than between-cluster distances in the case of the

trajectory clusters III and IV (p-value=7.38e-14 and 4.01e-7, respectively, one-sided Welch two-sample *t*-

test).

Interestingly we found that, among the five trajectory clusters, four of them started from the Tcmp

cluster but three of them started from different Tcmp sub-clusters. For instance, the trajectory clusters III,

I, and V started from Seurat cell clusters 0, 1, and 4, respectively (**Fig. S1**). First, to understand functional

differences among them, we identified marker genes for each cell cluster using Seurat and conducted a gene

set enrichment analysis using these marker genes. Specifically, hypergeometric tests were applied based on

a compilation of 160 immune signatures from the Immune Landscape of Cancer [28], followed by the

multiple testing adjustment using the Benjamini-Hochberg procedure [29]. Results indicate that these Tcmp

sub-clusters play disparate functional roles (**Fig. 5C**). Specifically, cell cluster 4 was enriched for cell cycle

mitotic and cell cycle checkpoints genes, cell cluster 1 was enriched for IL2 activation and signaling

pathways, and cell cluster 0 was enriched for immune response and T cell activation. This analysis suggests

that integration of scTCR-seq and scRNA-seq data using LRT can reveal the complexity of trajectories that

can be missed by the trajectory analysis based only on scRNA-seq data. Next, to understand differences

among trajectory clusters from a clonal perspective, we conducted the TCR repertoire analysis based on the

Seurat cell clusters. Specifically, we evaluated the overlap of trajectory clusters in terms of clonotype using

the Morisita index (**Fig. S2**). Morisita index ranges from 0 to 1 and the larger value indicates a higher extent

of overlapping [30]. Repertoire analysis indicates an extremely low overlap of clonotypes among these

Tcmp sub-clusters, which further confirmed our clustering results.

Figure 5. Trajectory inference using the LRT framework. (A) Trajectory clusters obtained using LRT. Dots represent cells, while colors and numbers (0 – 10) represent Seurat cell clusters. Each curve indicates the final trajectory for each trajectory cluster, and Roman numbers within circles (I – V) denote trajectory cluster indices. **(B)** Comparison of distances between TCR sequences, where 'W' and 'B' refer to within- and between-cluster distances, respectively (see the main text for calculation of these distances). **(C)** Enrichment analysis results for the Seurat cell clusters annotated as the Tcmp cell type. The colors represent -log10-transformed adjusted *p*-values.

Finally, we identified marker genes for each trajectory cluster using the NB-GAM approach (**Table**

S1) and evaluated their expressions as a function of the pseudotime. In general, we found that multiple

well-established Th1 markers showed increasing expressions while multiple well-known T follicular helper

cell (Tfh) markers showed decreasing expressions as a function of pseudotime (**Fig. 6**; **Fig. S3**). For instance,

it has been reported that enhanced inhibitor of DNA binding 2 (*Id2*) promotes Th1 cell differentiation while

suppressing *Id3* [31, 32]. Therefore, in the process of CD4$^+$ T cell differentiating towards Th1 cells, it is

expected that *Id2* is upregulated while *Id3* is downregulated, which coincides with our observation for the

identified trajectory clusters (**Fig. 6A** and **6B**; **Fig. S3**). Furthermore, chemokine receptor *Cxcr6* serves as

a useful marker for Th1 cells [26] while *Cxcr5* is a known marker for Tfh cells. When enhanced *Id2*

expression in CD4$^+$ T cell shifts the balance of Th1/Tfh towards Th1 cells, *Cxcr6* expression would be up-

regulated while *Cxcr5* expression would be suppressed [31]. **Fig. 6C** and **6D** showed patterns in line with

this point. Besides the above-mentioned markers, we also observed increasing trends for other Th1-

associated markers, such as the Th1 lineage-defining transcription factors *Tbx21*, *Gzmb*, *Ly6c2*, and *Cx3cr1*,

while observing decreasing trends for the Tfh-specific genes, such as *Ascl2*, *Il4*, *Il21*, and *Tox2* [33] (**Fig.**

S3). In summary, these expression patterns of Th1- and Tfh-associated markers nicely coincide with the

established Th1/Tfh biology, which further confirms the cell trajectories identified using LRT.

Fig. 6. Examples of the trajectory marker genes identified by LRT, which are involved in T cell differentiation. (A) and **(B)** Log-transformed read counts of *Id2* and *Id3* genes, respectively, as a function of pseudotime, for the trajectory cluster I. **(C)** and **(D)** Log-transformed read counts of *Cxcr6* and *Cxcr5* genes, respectively, as a function of pseudotime, for the trajectory cluster III. Curves represent smoothing splines fits and dots represent cells. The colors denote different cell types (green for Tcmp, salmon for pre-Th1, light violet for Th1-inter, and Iris blue for Th1). Arrows represent the direction of differentiation.

Discussion

In this paper, we proposed LRT, a novel computational framework for inferring T cell trajectory by integrating scTCR-seq and scRNA-seq data. LRT addresses the limitation of previous trajectory inference methods that solely rely on scRNA-seq data and neglects the clonal relationship between cells. Specifically, LRT utilizes both the functional information from scRNA-seq data and clonal information from scTCR-seq data. Such integrative analysis allows researchers to identify cell trajectories that cannot be revealed solely based on scRNA-seq data. In addition, LRT provides information about how clonotypes are clustered based on their DTW distances and MST, and this can help researchers decide which clonotypes need to be investigated jointly. Finally, LRT allows researchers to identify marker genes characterizing each of these trajectory clusters through DE analysis based on the principal curve and NB-GAM approaches, and this will help them understand potential drivers of cell differentiation, which is often the ultimate target.

Despite such strengths and power of LRT, it is also not free of limitations, which we plan to investigate in the future. First, while the current LRT framework assumes a single trajectory within each trajectory cluster, biologically various branching events can be possible within each trajectory cluster. We plan to address this limitation by considering more complex trajectory inference algorithms [3, 4]. Second, currently, LRT employs a simple association test evaluating which genes show significant expression changes along the pseudotime. However, researchers might also want to know whether these changes occur more in the earlier stage or in the later stage of the differentiation process. This limitation might be addressed by considering different types of the null hypothesis for the Wald test of NB-GAM. For example, we may modify the contrast matrix so that we can compare gene expression at starting points with those at ending points, as implemented in tradeSeq [21]. Third, currently, LRT implements DE analysis only to determine marker genes characterizing each trajectory cluster. However, researchers might also want to check which genes show DE between trajectory clusters. We plan to investigate approaches like those proposed by Van den Berge et al. [21] to address this. With the aforementioned strengths of the proposed

399 LRT framework and these future development plans, we believe that LRT can be a powerful tool for cell

400 trajectory inference and integrative analysis of scRNA-seq and scTCR-seq data.

Supporting information

Figure S1. Clonotype trajectories belonging to each of the trajectory clusters I - V.

Figure S2. TCR repertoire overlap between cell clusters. The numbers within cells show Morisita

indices. The numbers on the *x* and *y* axis denote cell cluster indices.

Figure S3. Additional examples of the trajectory marker genes identified by LRT, which are

involved in T cell differentiation. The left column shows the Th1-associated markers and the right

column shows the Tfh-associated markers. Curves represent smoothing splines fits and dots represent

cells. The colors denote different cell types (green for Tcmp, salmon for pre-Th1, light violet for Th1-

inter, and Iris blue for Th1).

Table S1. List of DE genes identified using LRT for the antigen-specific CD4$^+$ T cell data. Each tab

provides marker genes for each trajectory cluster. Wald statistics (waldStat), degree of freedom (df),

original *p*-value (pvalue), mean log fold change (meanLogFC), and Bonferroni-adjusted *p*-values (p.adj)

are reported for each gene.

Data Availability

417 The LRT framework was implemented as an R package 'LRT' and it is publicly available at

418 https://github.com/JuanXie19/LRT. The Shiny apps 'shinyClone' and 'shinyClust' are also provided as part

419 of this R package. The data used to produce the results presented in this manuscript is available at GEO

420 with accession number GSE158896.

Funding

This work has been supported through grant support from the National Human Genome Research Institute (R21 HG012482), National Institute on Aging (U54 AG075931), National Institute of General Medical Sciences (R01 GM122078 and R01 GM131399), National Institute on Drug Abuse (U01 DA045300), the National Science Foundation (NSF1945971), and the Pelotonia Institute of Immuno-Oncology (PIIO). The content is solely the responsibility of the authors and does not necessarily represent the official views of the funders.

Competing interests

The authors have declared that no competing interests exist.

References

1.   Wagner DE, Klein AM. Lineage tracing meets single-cell omics: opportunities and challenges. Nature Reviews Genetics. 2020;21(7):410-27.

2.   Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. Nature biotechnology. 2014;32(4):381-6.

3.   Street K, Risso D, Fletcher RB, Das D, Ngai J, Yosef N, et al. Slingshot: cell lineage and pseudotime inference for single-cell transcriptomics. BMC genomics. 2018;19(1):1-16.

4.   Saelens W, Cannoodt R, Todorov H, Saeys Y. A comparison of single-cell trajectory inference methods. Nature biotechnology. 2019;37(5):547-54.

5.   Zafar H, Lin C, Bar-Joseph Z. Single-cell lineage tracing by integrating CRISPR-Cas9 mutations with transcriptomic data. Nature communications. 2020;11(1):1-14.

6.   Stubbington MJ, Lönnberg T, Proserpio V, Clare S, Speak AO, Dougan G, et al. T cell fate and clonality inference from single-cell transcriptomes. Nature methods. 2016;13(4):329-32.

7.   Scheper W, Kelderman S, Fanchi LF, Linnemann C, Bendle G, de Rooij MA, et al. Low and variable tumor reactivity of the intratumoral TCR repertoire in human cancers. Nature medicine. 2019;25(1):89-94.

8.   Wang P, Jin X, Zhou W, Luo M, Xu Z, Xu C, et al. Comprehensive analysis of TCR repertoire in COVID-19 using single cell sequencing. Genomics. 2021;113(2):456-62.

9.   Team I. immunarch: an R package for painless bioinformatics analysis of T-cell and B-cell immune repertoires. Zenodo10. 2019;5281.

10.   Borcherding N, Bormann NL, Kraus G. scRepertoire: An R-based toolkit for single-cell immune
receptor analysis. F1000Research. 2020;9.
11.   Zhang Z, Xiong D, Wang X, Liu H, Wang T. Mapping the functional landscape of T cell receptor
repertoires by single-T cell transcriptomics. Nature methods. 2021;18(1):92-9.
12.   McInnes L, Healy J, Melville J. Umap: Uniform manifold approximation and projection for
dimension reduction. arXiv preprint arXiv:180203426. 2018.
13.   Graham RL, Hell P. On the history of the minimum spanning tree problem. Annals of the History
of Computing. 1985;7(1):43-57.
14.   Snijders TA, Nowicki K. Estimation and prediction for stochastic blockmodels for graphs with
latent block structure. Journal of classification. 1997;14(1):75-100.
15.   Allen C, Jung KJ, Chang Y, Ma Q, Chung D. Analysis of community connectivity in spatial
transcriptomics data. bioRxiv. 2022.
16.   Hart PE, Stork DG, Duda RO. Pattern classification: Wiley Hoboken; 2000.
17.   Daudin J-J, Picard F, Robin S. A mixture model for random graphs. Statistics and computing.
2008;18(2):173-83.
18.   Biernacki C, Celeux G, Govaert G. Assessing a mixture model for clustering with the integrated
completed likelihood. IEEE transactions on pattern analysis and machine intelligence. 2000;22(7):719-25.
19.   Hastie T, Stuetzle W. Principal curves. Journal of the American Statistical Association.
1989;84(406):502-16.
20.   Wood SN. Generalized additive models: an introduction with R: chapman and hall/CRC; 2006.
21.   Van den Berge K, Roux de Bézieux H, Street K, Saelens W, Cannoodt R, Saeys Y, et al.
Trajectory-based differential expression analysis for single-cell sequencing data. Nature communications.
2020;11(1):1-13.
22.   Hao Y, Hao S, Andersen-Nissen E, Mauck III WM, Zheng S, Butler A, et al. Integrated analysis
of multimodal single-cell data. Cell. 2021;184(13):3573-87. e29.
23.   Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq data
using regularized negative binomial regression. Genome biology. 2019;20(1):1-15.
24.   Chiquet J, Donnet S, Barbillon P. sbm:Stochastic Blockmodels. 2021.
25.   Sarda-Espinosa A. dtwclust: Time series clustering along with optimizations for the dynamic time
warping distance. R package version. 2019;5(6).
26.   Khatun A, Kasmani MY, Zander R, Schauder DM, Snook JP, Shen J, et al. Single-cell lineage
mapping of a diverse virus-specific naive CD4 T cell repertoire. Journal of Experimental Medicine.
2021;218(3).
27.   Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E. Fast unfolding of communities in large
networks. Journal of statistical mechanics: theory and experiment. 2008;2008(10):P10008.
28.   Thorsson V, Gibbs DL, Brown SD, Wolf D, Bortone DS, Yang T-HO, et al. The immune
landscape of cancer. Immunity. 2018;48(4):812-30. e14.
29.   Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful
approach to multiple testing. Journal of the Royal statistical society: series B (Methodological).
1995;57(1):289-300.
30.   Chiffelle J, Genolet R, Perez MA, Coukos G, Zoete V, Harari A. T-cell repertoire analysis and
metrics of diversity and clonality. Current Opinion in Biotechnology. 2020;65:284-95.
31.   Hwang S-M, Im S-H, Rudra D. Signaling networks controlling ID and E protein activity in T cell
differentiation and function. Frontiers in Immunology. 2022;13.
32.   Yang CY, Best JA, Knell J, Yang E, Sheridan AD, Jesionek AK, et al. The transcriptional
regulators Id2 and Id3 control the formation of distinct memory CD8+ T cell subsets. Nature
immunology. 2011;12(12):1221-9.
33.   Ruterbusch M, Pruner KB, Shehata L, Pepper M. In vivo CD4+ T cell differentiation and
function: revisiting the Th1/Th2 paradigm. Annual review of immunology. 2020;38:705-25.