Identification of microRNAs in the Lyme disease vector *Ixodes scapularis*

Deepak Kumar[1,2], Latoyia P. Downs[2], Monica Embers[3], Alex Flynt[1,2], and Shahid Karim[1,2]*

[1] Center for Molecular and Cellular Biosciences, University of Southern Mississippi, Hattiesburg,

MS 39406, USA

[2] School of Biological, Environmental, and Earth Sciences, The University of Southern

Mississippi, Hattiesburg, MS 39406, USA

[3] Division of Immunology, Tulane National Primate Research Center, 18703 Three Rivers Rd.

Covington, LA 70433

***Corresponding author:** shahid.karim@usm.edu

Abstract

MicroRNAs (miRNAs) are a class of small non-coding RNAs involved in many biological processes, including the immune pathways that control bacterial, parasitic, and viral infections. Pathogens probably modify host miRNAs to facilitate successful infection, so they might be useful targets for vaccination strategies. There are little data on differentially expressed miRNAs in the black-legged tick *Ixodes scapularis* after infection with *Borrelia burgdorferi*, the causative agent of Lyme disease in the United States. Small RNA sequencing and qRT-PCR analysis were used to identify and validate differentially-expressed *I. scapularis* salivary miRNAs. Small RNA-seq yielded 133,465,828 (≥18 nucleotides) and 163,852,135 (≥18 nucleotides) small RNA reads from *Borrelia*-infected and uninfected salivary glands for downstream analysis using the miRDeep2 algorithm. Two hundred and fifty-four miRNAs were identified across all datasets, 25 of which were high confidence and 51 low confidence known miRNAs. Twenty-three miRNAs were differentially expressed in uninfected and infected salivary glands: 11 were upregulated and 12 were downregulated upon pathogen infection. This study provides new insights into the miRNAs expressed in *I. scapularis* salivary glands and pave the way for their functional manipulation to prevent or treat *B. burgdorferi* infection.

Keywords: *Ixodes scapularis*, salivary glands, microRNA, RNA sequencing, *Borrelia burgdorferi*

Introduction

Small non-coding RNAs (sncRNAs) regulate genes at the post-transcriptional level in animals, plants, and arthropods including ticks [1-5]. MicroRNAs (miRNAs) are a class of sncRNA between 18 and 25 nucleotides in length, and they are now known to be important in arthropod immunity and host-pathogen interactions through their involvement in several cellular processes including development, immunity, and pathogen responses in arthropods [2, 6-10]. In animals, miRNAs regulate post-transcriptional gene expression, most often by binding to the 3'-untranslated region (3'-UTR) of target genes. While perfect complementarity of 2-8 nucleotides at the 5' end of the miRNA (seed region) is necessary for miRNA regulation, the remaining sequence can harbor mismatches or bulges [2, 11, 12]. miRNAs are transcribed as primary miRNA transcripts before processing by Drosha and Pasha proteins into a pre-miRNAs. These are then exported to the cytoplasm and processed by Dicer into mature miRNAs, which are loaded onto the microRNA-induced silencing complex (miRISC) before targeting complementary mRNA for degradation [13, 14].

Several proteomics and transcriptomics studies have identified differentially expressed transcripts and proteins in uninfected and infected ticks [1, 15-18], but few have investigated the role of pathogens in the differential modulation of the post-transcriptional tick and host machinery. Although there are over 800 tick species and despite their importance as vectors of human and animal diseases, ticks are underrepresented in available miRNA resources. For example, the miRbase database contains 49 *Ixodes scapularis* miRNAs and 24 *Rhipicephalus microplus* miRNAs, while MirGeneDB 2.1 contains 64 *Ixodes scapularis* miRNAs. However, small

RNA sequencing (RNA-seq) and computational approaches are accelerating the discovery of

miRNAs from species with incomplete genome sequencing, assembly, and annotation.

Ixodes scapularis is a primary vector of human pathogens including the Lyme disease

agent *Borrelia burgdorferi*, which infects vertebrates and ticks through evolved complex

mechanisms. There is now a good understanding of tick immune pathways and their interactions

with *B. burgdorferi* [19-21], but it is still uncertain how *B. burgdorferi* avoids clearance. *B.*

burgdorferi must traverse tick salivary glands during transmission [71]. Since saliva/salivary gland

proteins can enhance *B. burgdorferi* transmission into the vertebrate host, characterization of

molecular interactions at the tick bite site and the tick salivary glands is expected to facilitate

vaccine development [22, 23], since promoting immunity against tick salivary proteins could

neutralize tick bites and pathogen transmission. Once *B. burgdorferi* is acquired by ticks from

infected hosts, it resides in the tick gut and only migrates to the salivary gland during subsequent

blood feeding, which generally lasts for 3-7 days [72, 73, 74, 75] . While it is known that other

tick-borne pathogens such as *Anaplasma marginale* must replicate inside salivary glands for

efficient transmission [76], the details of *B. burgdorferi* replication are less well understood.

Indeed, borrelial spirochetes invade the tick salivary gland via an unknown mechanism [71] and

might be carried to the host dermis via tick saliva. Several salivary gland genes are upregulated

in *B. burgdorferi*-infected *Ixodes scapularis* nymphs compared with uninfected ones [24],

suggesting a significant role for salivary gland gene regulation in *B. burgdorferi* infection and

transmission.

To fill the knowledge gap on miRNA expression in *Ixodes scapularis* salivary glands, here

we performed miRNA profiling of partially fed *B. burgdorferi*-infected and uninfected tick salivary

glands to identify miRNAs that might play a role in *B. burgdorferi* survival, colonization,

transmission, and host immunomodulation. In doing so, we detected 254 miRNAs, of which 25

were high confidence miRNAs and 51 low confidence miRNAs. Forty-one of the identified miRNAs

were present as *I. scapularis* miRNAs in miRBase (v22.1). Gene ontology and network analysis of

target genes of differentially expressed miRNAs predicted roles in metabolic, cellular,

development, cellular component biogenesis, and biological regulation processes. Several KEGG

pathways including sphingolipid metabolism; valine, leucine and isoleucine degradation; lipid

transport and metabolism; exosome biogenesis and secretion; and phosphate-containing

compound metabolic processes were predicted as targets of differentially expressed miRNAs.

Materials and Methods

Ethics statement

All animal experiments were performed in strict accordance with the recommendations in the

NIH Guide for the Care and Use of Laboratory Animals. The Institutional Animal Care and Use

Committee of the University of Southern Mississippi approved the protocol for blood feeding of

field-collected ticks (protocol # 15101501.1).

Ticks and tissue dissections

Ticks were purchased from the Oklahoma State University Tick Rearing Facility. Adult male and

female *I. scapularis* were kept according to standard practices [25] and maintained in the

laboratory as described in our previously published work [26, 27]. Unfed female adult *I. scapularis*

were infected with laboratory grown *B. burgdorferi* strain B31.5A19 using the capillary feeding

method at Tulane National Primate Research Center [28]. *Borrelia* infected and uninfected ticks

(n=45 in each group) were placed on each ear of a rabbit host for tick blood-feeding. Blood-fed

adult female *I. scapularis* were dissected within 60 min of removal from the rabbit. Tick tissues

were dissected and washed in M-199 buffer as described previously [77]. Salivary glands and

midguts from individual *I. scapularis* were stored in RNAlater (Life Technologies, Carlsbad, CA,

USA) at −80°C until use.

RNA isolation, cDNA synthesis, and PCR-based *B. burgdorferi* detection in tick tissues

The TRIzol method was used for RNA extraction from individually-dissected midgut tissues, and

cDNA was synthesized as described previously [29, 78]. *B. burgdorferi* was detected in tick

midguts using the *flaB* gene in a PCR assay [29, 30]. After testing for *B. burgdorferi* infection in

tick midguts, the corresponding salivary gland tissues from the same uninfected/infected ticks

(n=10 salivary glands from each group) were pooled in separate tubes and RNA isolated using the

TRIzol method [78] .

Small RNA sequencing (RNA-seq)

Small RNA libraries were made using the Illumina TruSeq Kit following the manufacturer's

protocol (Illumina, San Diego, CA). Briefly, short adapter oligonucleotides were ligated to each

end of the small RNAs in the sample, cDNA made with reverse transcriptase, and PCR used to add

sample-specific barcodes and Illumina sequencing adapters. The final concentration of all

sequencing libraries was determined using a Qubit fluorometric assay (Thermo Fisher Scientific,

Waltham, MA), and the DNA fragment size of each library was assessed using a DNA 1000 high-

sensitivity chip on an Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA). After

purification by polyacrylamide gel electrophoresis, sample libraries were pooled and sequenced

on an Illumina NextSeq 500 (single end 36 base) using the TruSeq SBS kit v3 (Illumina) and

protocols defined by the manufacturer. Four small RNA libraries of clean and *B. burgdorferi*-

infected, partially fed, and pooled salivary glands were sequenced. RNA library preparation and

indexing were performed at the UMMC sequencing facility.

Data analysis

A schematic of the experimental plan and data analysis is shown in **Supplementary Figure S1**.

miRDeep2 v.2.0.0.8 [31, 32], was used to process RNA-seq data. To predict novel miRNAs, the

reads from all samples were combined. The mapper function of miRDeep2 first trims the adapter

sequences from the reads and converts the read files from FASTQ to FASTA format. Reads shorter

than 18 bases were discarded and the remaining reads mapped to the *I. scapularis* reference

genome using default miRDeep2 mapper function parameters. Reads mapping to the genome

were used to predict novel miRNAs. The *Drosophila melanogaster* genome was also used as

another reference genome, and mapped reads were aligned to available miRNAs of *D.*

melanogaster in miRBase (v22) and quantified. Reads were mapped to the reference genomes of

D. melanogaster and *I. scapularis* and locations of potential miRNA read accumulations

identified. The regions immediately surrounding the mapped reads were examined for miRNA

biogenesis features including mature miRNAs, star and precursor reads, and stem-loop folding

properties. miRDeep2 models the miRNA biogenesis pathway and uses a probabilistic algorithm

to score compatibility of the position and frequency of sequencing reads with the secondary

structure of the miRNA precursor.

140 For miRNA expression, a count table was generated using bedtools multicov, which counts

141 alignments from indexed BAM files that overlap intervals in BED files provided from the

142 miRDeep2 analysis.

143 **ISE6 cell culture, infection with *B. burgdorferi*, and validation of differentially expressed**

144 **miRNAs by qRT-PCR**

145 Differentially expressed miRNAs in RNA-seq data were validated by qRT-PCR in the ISE6 cell line

146 derived from *I. scapularis* embryos cultured and maintained as suggested by Munderloh and

147 Kurtti [33]. The *B. burgdorferi* (strain B31) isolate was a kind donation from Dr. Monica Embers

148 (Tulane University, Covington, LA). Cultures were grown and maintained as suggested previously

149 [28]. Once ISE6 cells reached confluency, they were infected with *B. burgdorferi* as described

150 previously [34]. Briefly, ISE6 cells were inoculated with the supernatant of a log phase *B.*

151 *burgdorferi* culture at a multiplicity of infection of 50 and incubated for 24 hours before

152 harvesting. RNA was isolated using the TRIzol method and the expression of miRNAs analyzed by

153 qRT-PCR with the Mir-X miRNA qRT-PCR TB Green kit (Takara Bio Inc, Kusatsu, Shiga, Japan;

154 catalog #638316). qRT-PCR conditions were an initial denaturation at 95°C for 10 mins then 40

155 cycles of 95°C for 5 secs, 60°C for 20 secs.

156 **Normalization, differential expression (DE), and statistical analysis of miRNAs between**

157 **uninfected and infected salivary glands**

158 Differential expression (DE) analysis of identified miRNAs was performed using the interactive

159 web interface DeApp [35]. Low expression genetic features were removed after alignment if the

160 counts per million (CPM) value was ≤1 in less than two samples. Sample normalization and a

multidimensional scaling (MDS) plot are shown in **Supplementary Fig. S2**. DE analysis was

performed with edgeR with a false discovery rate (FDR) adjusted *p*-value of 0.05 and minimum

fold-change of 1.5. DeApp displays a dispersion plot showing the overall DE analysis along with

statistical significance (*p*-value, FDR adjusted *p*-value) and a volcano plot corresponding to the

specified parameters and cutoff values.

In silico* mapping of *B. burgdorferi*-infected small RNA sequences to the *Borrelia burgdorferi

genome

In silico mapping of small RNA sequences of *B. burgdorferi*-infected salivary glands to the *Borrelia*

burgdorferi genome (GCF_000181575.2_ASM18157v2_genomic.fna) detected 18,165 *Borrelia*

burgdorferi sequences, but no miRNAs of *B. burgdorferi* origin were predicted by miRDeep2

analysis.

Prediction of target genes, proteome re-annotation, gene ontology (GO), and KEGG enrichment

analyses

TargetSpy [36], MIRANDA [37], and PITA [38] were used in miRNAconsTarget from sRNAtoolbox

to predict genes regulated by tick salivary gland miRNAs up- or downregulated in the *in silico*

analysis [39]. Targets common to all three programs were further considered. *In silico* target

prediction provided a high number of false positives, but cross-species comparisons and

combinatorial effects reduced this number [40]. Target gene networks and KEGG pathways

significantly enriched for target genes were extracted using the STRING [41] output. PANNZER2

[42] was used to functionally re-annotate the proteome of up- or downregulated genes (targets

of detected miRNAs), and WEGO [43] was used to analyze and plot gene ontology (GO)

annotations.

184   Results and Discussion

185   Profile characteristics of small RNA libraries

There were 216,292,174 raw small RNA reads from *B. burgdorferi*-infected salivary glands and

212,542,697 from uninfected salivary glands. After adapter trimming and removal of short reads

(≤18 nucleotides (nt)), 133,465,828 small RNA reads were available from *B. burgdorferi*-infected

samples and 163,852,135 from uninfected samples for downstream analysis. The read length

distribution shows the types of small RNAs present in *B. burgdorferi*-infected and uninfected

salivary gland samples. Two main peaks at 22 nt (miRNAs/siRNAs) at 29 nt (piRNAs) were

distinguishable in *B. burgdorferi*-infected and uninfected samples (**Figure 1A**). There were ~5 x

10^7 and ~2.5 x 10^7 22 nt miRNA sequences in uninfected and infected salivary gland samples,

respectively. The read length distribution in uninfected samples was comparable to infected

samples.

196   Other small RNA categories

A summary of reads matching various small RNAs categories from *B. burgdorferi*-infected and

uninfected salivary glands are shown in **Figure 1B**. Other small RNAs include signal recognition

particle rRNAs, ncRNAs, protein-coding RNAs, snoRNAs, snRNA SRP RNAs, and not annotated

small RNAs. Of the total small RNA reads from *Ixodes scapularis* salivary glands, 12.3% were

miRNAs, 2.2% rRNAs, 0.4% ncRNAs, 0.8% protein coding, 0.07% snoRNAs, 0.05% snRNAs, 0.04%

SRP RNAs, and 84.3% reads were not annotated.

Figure 1. (A) Small RNA sequence length distribution in uninfected (clean) and *Borrelia*
burgdorferi-infected *Ixodes scapularis* salivary glands. MicroRNAs are twenty-four (22)
nucleotides in length. Clean, uninfected salivary glands; infected, *Borrelia burgdorferi*-infected
salivary glands. (B) Summary of the abundance of combined small RNA reads from *Ixodes*
scapularis salivary glands matching various small RNA categories. Reads from uninfected and
Borrelia burgdorferi infected samples are combined together for abundance calculation.
210
MicroRNA profiling of infected and uninfected salivary glands and identification of novel tick

miRNAs

Fig. 2A shows the basic hairpin-loop structure of an miRNA and other parameters (Dicer cut

overhangs, total read count, mature read count, loop read count, total read count, randfold

score, and total score) used to determine whether a hairpin-loop structured RNA is an miRNA.

Using miRDeep2, 254 miRNAs were predicted in the salivary gland libraries. By adopting a

conservative approach, miRNAs were categorized as high (n=25) and low confidence (n=51)

miRNAs (**Fig. 2B**) based on standard criteria [79]. Of 49 *Ixodes scapularis* miRNAs present in

miRBase, 41 were detected in our data. Of the 254 *Ixodes scapularis* miRNAs, several had

homologs present in *D. melanogaster.* The details of the identified miRNAs and their miRDeep2

scores are presented in **Supplementary Table S1**.

Figure 2. (A) Basic stem-loop structures of predicted microRNAs. miRDeep2 was used to identify
potential miRNA precursors based on nucleotide length, star sequence, stem-loop folding, and
homology to the *Ixodes scapularis* reference genome. Shown are the predicted stem-loop
structures (yellow), star (pink), and mature sequences of predicted miRNAs (red) in the salivary
glands of *Ixodes scapularis* ticks. (B) Annotation of predicted miRNAs in partially fed uninfected
(clean) and *Borrelia burgdorferi*-infected *Ixodes scapularis* salivary glands. 254 microRNAs were
predicted in infected and uninfected salivary gland tissues. 25 were categorized as high
confidence (n = 25) and 51 as low confidence (n=51) based on standard criteria and a conservative
approach. Out of 49 *Ixodes scapularis* miRNAs available in miRbase (v22.1), 41 were detected in
this study.

***In silico* DE analysis of miRNAs in *B. burgdorferi*-infected salivary glands**

In DE analysis, 11 miRNAs were upregulated and 12 were downregulated in infected salivary
glands compared with uninfected salivary glands (**Figure 3A, B; Table 1**). Several of the
identified tick miRNAs were conserved in *D. melanogaster* (dme-miR-375-3p, dme-miR-993-5p,
dme-miR-12-5p, dme-bantam-3p, dme-miR-100-5p, dme-miR-8-3p, and dme-miR-304-5p).
miRNAs downregulated in *B. burgdorferi*-infected salivary glands relative to uninfected salivary
glands   were   isc-miR-153,   isc-miR-1   and   isc-miR-79,   nDS_002871802.1_376,
nDS_002633080.1_31953,   nDS_002549652.1_43448,   nDS_002537755.1_45711,
nDS_002537755.1_45739,   nDS_002861213.1_1750,   nDS_002763926.1_14582,   and
nDS_002548557.1_43749, while upregulated miRNAs were isc-miR-317, isc-miR5310, isc-miR-
2001,   isc-miR-5307,   isc-miR-71,   isc-miR-87,   nDS_002784743.1_12165,
nDS_002664372.1_28027,   nDS_002716687.1_21278,   nDS_002620414.1_33611,   and
nDS_002680650.1_26098.
246
247

Figure 3. *In silico*
differential expression of predicted miRNAs in *Borrelia burgdorferi*-infected, partially fed salivary
glands relative to partially fed clean salivary glands. EdgeR was used for differential expression
analysis. 12 predicted miRNAs were down-regulated, 11 were upregulated, while 39 were
unaffected. miRNAs with a log2 fold-change expression > |1| and FDR ≤ 0.1 were considered
significantly differentially expressed (see Table 1). Also, in Table 2, putative roles and targets of
these DE conserved miRNAs have been provided based on available studies in other arthropods.
275

Table 1. *In silico* differential expression of *Ixodes scapularis* miRNAs in salivary glands after *Borrelia burgdorferi* infection compared with uninfected salivary glands. miRNAs with a log2 fold-change expression > |1| and FDR ≤ 0.1 were considered significantly differentially expressed. Values highlighted in **red** indicate significant upregulation and values highlighted in **green** indicate significant downregulation.

Differentially expressed miRNA	Log$_2$FC	logCPM	LR	P-value	FDR
isc-miR-87	4.245604	11.83289	33.20241	8.30E-09	4.65E-08
nDS_002784743.1_12165	5.511035	11.48887	29.7299	4.97E-08	1.55E-07
isc-miR-71	5.226443	11.95849	48.94389	2.63E-12	3.69E-11
isc-miR-5307	2.570363	12.34313	24.71399	6.65E-07	1.55E-06
nDS_002664372.1_28027	5.362184	11.43227	26.80077	2.26E-07	5.84E-07
nDS_002716687.1_21278	5.511153	11.48887	29.72059	4.99E-08	1.55E-07
isc-miR-2001	3.146271	11.78845	19.75379	8.81E-06	1.64E-05
nDS_002620414.1_33611	5.361301	11.43227	26.76848	2.29E-07	5.84E-07
isc-miR-5310	5.009416	11.31195	20.86089	4.94E-06	1.06E-05
nDS_002680650.1_26098	1.567661	15.43032	103.3589	2.80E-24	7.83E-23
isc-miR-317	3.146271	11.78845	19.75379	8.81E-06	1.64E-05
nDS_002871802.1_376	-0.80359	15.73883	19.22785	1.16E-05	1.91E-05
nDS_002633080.1_31953	-4.7431	11.95812	10.11663	0.001469	0.001959
nDS_002549652.1_43448	-4.9974	12.11011	12.13051	0.000496	0.000731
nDS_002537755.1_45711	-6.34111	13.07343	30.24285	3.81E-08	1.52E-07
nDS_002537755.1_45739	-6.57147	13.2637	36.56262	1.48E-09	1.03E-08
isc-miR-79 dme-miR-9a-5p	-5.35587	12.34273	15.64359	7.65E-05	0.000119
isc-miR-1 dme-miR-1-3p	-2.44072	12.67077	10.50794	0.001189	0.001664
nDS_002861213.1_1750	-0.80359	15.73883	19.22785	1.16E-05	1.91E-05
nDS_002763926.1_14582	-6.57147	13.2637	36.56262	1.48E-09	1.03E-08
nDS_002548557.1_43749	-6.34111	13.07343	30.24285	3.81E-08	1.52E-07
isc-miR-153	-4.15091	11.64621	6.563386	0.01041	0.013249

miR-2001 was upregulated in *B. burgdorferi*-infected salivary glands compared with controls. miR-2001 is not present in the *Drosophila* genome but is an evolutionarily conserved

285   **Table 2.** List of differentially expressed microRNAs detected in *Borrelia burgdorferi* infected tick

286   salivary glands and their putative roles

microRNA	Role of ortholog microRNA	Target genes	Reference
isc-miR-153	Development, immune response		[82]
isc-miR-1	Stress response, immunity, development, facilitating infection	*HSP60, HSP70, GATA4*	[49, 50, 51, 52, 83, 84, 85, 86, 87]
isc-miR-79	Innate immunity, differentiation, apoptosis	*Roundabout protein 2 pathway (robo2), DRAPER, HP2 (Hemolymph protease), P38, pvr, puc*	[55, 56, 57, 88, 89, 90, 91]
isc-miR-317	Negative influencer of Toll pathway, host homeostatic response (targeting Gap junctions, TRP channels), mitotic regulation, Fecundity	*Dif-Rc, Thioredoxin peroxidase-2, STAT*	[45, 91, 92, 93, 94]
isc-miR-2001	Under extreme environmental conditions		[95]
isc-miR-5307	Propagation of Powassan virus inside vero cells		[96]
isc-miR-71	Stress response, development, facilitates WSSV infection	*cap-1 , riok1, Actin1, myD88, IML3*	[91, 97, 98, 99, 100]
isc-miR-87	Pathogen survival (disruption of Toll and IMD pathway)	*Serine/Threnine Kinase, Toll 1A, Putative TLR 5b, FADD*	[60, 62, 63, 91, 101, 102]
isc-miR-5310	Pathogen survival by modulating signaling pathways, Feeding behavior		[64-68, 81, 103]

miRNA in ticks [44]. It has previously been shown to play a role in host immunomodulation

required for pathogen survival. miR-2001 has been detected in *I. ricinus* tick saliva [45] and *H.*

longicornis tick saliva extracellular vesicles (EVs) [46] along with several other miRNAs, suggesting

that EVs containing these miRNAs could be transferred to the host to modulate host cellular

functions to facilitate tick and pathogen survival [45, 46]. EVs are involved in the intercellular

transfer of miRNAs, lipids, and proteins and the disposal of unnecessary cell contents [46, 47].

The discovery of EVs in the excretory-secretory products of ectoparasites suggests that EVs are

probably taken up by host cells, deliver their cargoes to the host, and favor immunomodulation,

pathogen survival, and disease progression [48, 80]

miR-1 was downregulated in *B. burgdorferi*-infected tick salivary glands compared with

uninfected salivary glands. miR-1 belongs to a family of miRNAs including miR-7 and miR-34

conserved across fruit flies, shrimps, and humans, where it modulates similar pathways

(development, apoptosis) and is upregulated during stress insults [49]. In mosquitoes, miR-1 is

upregulated during *Plasmodium* infection [50] and also facilitates or promotes West Nile virus

infection [51]. miR-1 has been shown to be generally upregulated in response to infection, while

we detected its downregulation in *B. burgdorferi*-infected SGs. In *Listeria*-infected macrophages,

miR-1 promotes IFN-γ-dependent activation of the innate immune response [52].

Tick saliva and salivary gland extracts reduce IFN-γ and IL-2 production in T cells and

inhibited T cell proliferation [53, 54] suggesting immune suppression, a possible survival

mechanism for tick pathogens. mir-79 was also downregulated in *Borrelia burgdorferi* infected

salivary glands, and mir-79 has been shown to participate in immunity and other processes such

as cellular differentiation, neurogenesis, and apoptosis. In *Rhipicephalus haemaphysaloides*, mir-

79 was downregulated upon lipopolysaccharide (LPS) induction in female and male ticks,

suggesting a role for mir-79 in LPS-mediated stimulation of the innate immune response [55].

The JNK pathway is an immune response pathway against Gram-negative bacteria [56], and mir-

79 is known to disrupt JNK signaling by targeting its component genes *pvr* (CG8222) and *puc*

(CG7850) [57]. Our detection of the downregulation of mir-79 in *B. burgdorferi*-infected salivary

glands is probably due to stimulation of the JNK pathway as a tick immune response against *B.*

burgdorferi. Although *B. burgdorferi* is described as an atypical Gram-negative bacterium due to

its double membrane, it lacks classical lipopolysaccharide (LPS). It also has a different cellular

organization and membrane composition to other diderms [58]. However, surprisingly in *A.*

phagocytophilum (an intracellular Gram-negative bacterial pathogen)-infected ticks, mir-79 was

upregulated to facilitate infection by targeting the Roundabout protein 2 pathway (Robo2) [59],

suggesting different roles for mir-79 in ticks when infected with different bacterial pathogens.

Borrelia burgdorferi (Bb) is extracellular and a Gram-negative bacterial pathogen, but *A.*

phagocytophilum is a well-known intracellular a Gram-negative bacterial pathogen. Do

extracellular and pseudo-Gram-negative status of *B. burgdorferi* make it a different bacterial

pathogen than Gram negative or positive? More clarity of tick immune response is required in

case of *B. burgdorferi* infection.

In *Drosophila [35]*, miR-317 negatively regulates Toll pathway signaling, and its

upregulation in *B. burgdorferi*-infected salivary glands may suggest a similar role to facilitate *B.*

burgdorferi survival inside tick salivary glands. *In silico*, miR-317 targets Dif-Rc, an important

transcription factor in the Toll pathway in *Drosophila* [35] and STAT in JAK-STAT signaling in

Manduca sexta [60]. An *in silico* study in *I. ricinus* suggested a combinatorial effect of tick salivary

miRNAs on host genes important for maintaining host homeostasis and tick-host interactions,

including miR-317 targeting gap junctions and TRP channels, which play significant roles in host

homeostatic responses [45]. miR-71 was upregulated in *B. burgdorferi*-infected salivary glands,

and its predicted targets include *MyD88*, which is activated when ligands bind to the Toll-like

receptor (TLR), interleukin 1 receptor (IL-1R), or IFN-γR1 and trigger MyD88-mediated signaling

and pro-inflammatory cytokine responses. Another miR-71 target is IML3, an arthropod

immunolectin that recognizes LPS on Gram-negative bacteria as a part of arthropod immune

defenses. Immunolectins are also predicted targets of miR-87, -276, -9a, and -71 [60]. We

hypothesize that miR-71 disrupts tick immune pathways and protects *B. burgdorferi* in the

salivary gland. It has also been shown to prolong the life and regulate stress responses in

nematodes, being upregulated in the Dauer larval stage when food or other life-sustaining

resources are scarce [61].

miR-87 was upregulated in *B. burgdorferi*-infected salivary glands. Previous studies in

other arthropods such as *Manduca sexta* and *Aedes albopictus* have suggested a role in

disrupting innate immunity, particularly via IMD and Toll signaling pathways [60, 62, 63]. In *Aedes*

albopictus, miR-87's predicted targets are Toll pathway signaling Ser/Thr kinase, Toll-like

receptor Toll1A, class A scavenger receptor with Ser-protease domain, galectin [63], and TLR5b

[62], while in *Manduca sexta* its predicted target is FADD, an adaptor protein involved in DISC

formation [60]. Our *in silico* data also showed upregulation of miR-5310 in *B. burgdorferi*-infected

salivary glands. miR-5310 is a tick-specific miRNA [44], and a recent study demonstrated its

downregulation in *Anaplasma phagocytophilum*-infected nymphs compared with unfed

uninfected nymphs [64]. Previous studies have also indicated modulation of signaling events via

miR-5310 upon *A*. *phagocytophilum* infection [64-68, 81] In *B. burgdorferi* infection, we speculate

that might modulate signaling events and protect *B. burgdorferi* in the tick salivary glands. miR-

5310 might also be involved in tick feeding, as it was found to be downregulated in *Rhipicephalus*

microplus tick larvae upon exposure to host odor but not being allowed to feed [44].

Prediction of target genes and gene ontology (GO) and functional enrichment analyses of the

target network

Target proteins were used to build a high-confidence interaction network (interaction scores

>0.9).

362
Figure 4. A network built exclusively from *Ixodes scapularis* proteins targeted by *in silico*
differentially expressed miRNAs in *Borrelia burgdorferi*-infected, partially-fed salivary glands
relative to partially fed clean salivary glands. Red boxes indicates the proteins involved in
significant KEGG pathways such as sphingolipid metabolism (EEC07251.1, KEGG:R02541);
valine, leucine, and isoleucine degradation (EEC10220.1, KEGG:R04188) ; lipid transport and
metabolism (EEC02614.1, KEGG:R01178) ; exosome biogenesis and secretion (EEC07251.1,
EEC17749.1, KEGG:R02541) ; and phosphate-containing compound metabolic process
(EEC02290.1, KEGG:R00004)/

STRING web analysis (**Figure 4**) showed that the target proteins of 23 DE miRNAs (11

upregulated and 12 downregulated) had similar interactions to those expected for a random set

of proteins of similar size sampled from the *I. scapularis* genome (nodes = 687, edges = 79,

average node degree = 0.23, average local clustering coefficient = 0.0966, expected number of

edges = 77, PPI enrichment *p*-value = 0.411). This does not necessarily mean that these selected

proteins are not biologically meaningful, rather that these tick proteins may not be very well

studied and their interactions might not yet be known to STRING.

Many target genes were identified for the DE miRNAs using the sRNAtoolbox

miRNAconsTarget program [39]. To minimize false-positive targets, we chose only those targets

predicted by all three miRNA target prediction algorithms (TargetSpy, MIRANDA, and PITA).

Forty-one KEGG pathways were enriched for target genes (proteins) of DE miRNAs

(**Supplementary Table S2**) and included sphingolipid metabolism; valine, leucine, and isoleucine

degradation; lipid transport and metabolism; exosome biogenesis and secretion; and phosphate-

containing compound metabolic process (**Figure 4**). Gene ontology (GO) analysis indicated that

most target genes of DE miRNAs play significant roles in cellular processes, metabolic processes,

biological regulation, developmental processes, and responses to stimuli (**Figure 5**). Surprisingly,

immune response genes were one of the least affected functions.

Figure 5. Gene ontology (GO)-derived biological processes related to genes targeted by differentially expressed miRNAs in partially fed *B. burgdorferi*-infected salivary glands relative to partially fed uninfected salivary glands from *Ixodes scapularis* ticks.

Lipid metabolism was one of the main KEGG pathways detected by Pannzer and STRING analyses and was predicted to be regulated by tick salivary gland miRNAs miR-1, miR-5310, miR-71, and miR-79. It has previously been shown that binding of *B. burgdorferi* to host glycosphingolipid can contribute to tissue-specific adhesion of *B. burgdorferi*, and the inflammatory process in Lyme borreliosis might be affected by interactions between *B. burgdorferi* and glycosphingolipid [69]. Therefore, we hypothesize that tick miRNAs (via saliva) promote sphingolipid synthesis inside hosts to promote *Borrelia* adhesion, and indeed there is evidence that its infection affects lipid metabolism in hosts [70].

Validation of DE miRNAs by qRT-PCR

The expression of DE miRNAs were validated in *B. burgdorferi*-infected and uninfected ISE6 cells

by qRT-PCR (**Figure 6; Supplementary Table S3**), which closely mirrored the RNA-seq data for

many targets, although some differences were not statistically significant by qRT-PCR and isc-

miR-317 was downregulated rather than upregulated. These differences could be due to the use

of different methodologies to quantify miRNA expression [10].

Figure 6. qRT-PCR validation of differentially expressed miRNAs detected in *Borrelia burgdorferi*-infected

,partially-fed salivary glands relative to partially-fed uninfected salivary glands from *Ixodes scapularis*

ticks. qPCR validation was performed in *Borrelia burgdorferi* infected and clean ISE6 cells. Expression of

miRNAs was normalized to clean ISE6 cells (indicated as 1 on the y-axis). Statistical significance for qRT-

PCR-based differential expression was determined by the 2-tailed Student's *t*-test, where * is $p<0.05$.

Conclusions

This is the first comprehensive miRNA profiling study of *Ixodes scapularis* salivary glands with and without *Borrelia burgdorferi* infection. Here we identified several potential miRNAs targets in tick salivary glands which might play a significant role in *Borrelia* colonization, survival, transmission, and host immunomodulation. Functional validation of these miRNAs is now required. Further characterization of tick salivary gland miRNAs would contribute to a better understanding of the mechanisms underpinning *Borrelia* transmission and propagation inside hosts, not least due to its special status as an extracellular spirochaete and atypical Gram-negative organism that might exploit different survival mechanisms. The impact of *B. burgdorferi* on miRNA expression must also be studied in other tick tissues and hosts to understand cues of its vector competence in ticks and immunomodulation in vertebrates.

References

1.   Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. Cell. 2004;116(2):281-97. 10.1016/s0092-8674(04)00045-5

2.   Bartel DP. MicroRNAs: target recognition and regulatory functions. Cell. 2009;136(2):215-33. 10.1016/j.cell.2009.01.002

3.   Carrington JC, Ambros V. Role of microRNAs in plant and animal development. Science. 2003;301(5631):336-8. 10.1126/science.1085242

4.   Griffiths-Jones S, Saini HK, van Dongen S, Enright AJ. miRBase: tools for microRNA genomics. Nucleic Acids Res. 2008;36(Database issue):D154-8. 10.1093/nar/gkm952

5.   Lai EC. Two decades of miRNA biology: lessons and challenges. RNA. 2015;21(4):675-7. 10.1261/rna.051193.115

6.   Bartel DP, Chen CZ. Micromanagers of gene expression: the potentially widespread

influence of metazoan microRNAs. Nat Rev Genet. 2004;5(5):396-400. 10.1038/nrg1328

7.   Alvarez-Garcia I, Miska EA. MicroRNA functions in animal development and human

disease. Development. 2005;132(21):4653-62. 10.1242/dev.02073

8.   Miesen P, Ivens A, Buck AH, van Rij RP. Small RNA Profiling in Dengue Virus 2-Infected

Aedes Mosquito Cells Reveals Viral piRNAs and Novel Host miRNAs. PLoS Negl Trop Dis.

2016;10(2):e0004452. 10.1371/journal.pntd.0004452

9.   Momen-Heravi F, Bala S. miRNA regulation of innate immunity. J Leukoc Biol.

2018;103(6):1205-17. 10.1002/JLB.3MIR1117-459R

10.   Saldana MA, Etebari K, Hart CE, Widen SG, Wood TG, Thangamani S, et al. Zika virus

alters the microRNA expression profile and elicits an RNAi response in Aedes aegypti

mosquitoes. PLoS Negl Trop Dis. 2017;11(7):e0005760. 10.1371/journal.pntd.0005760

11.   Rigoutsos I. New tricks for animal microRNAS: targeting of amino acid coding regions at

conserved and nonconserved sites. Cancer Res. 2009;69(8):3245-8.

12.   Schnall-Levin M, Zhao Y, Perrimon N, Berger B. Conserved microRNA targeting in

Drosophila is as widespread in coding regions as in 3' UTRs. Proceedings of the National

Academy of Sciences. 2010;107(36):15751-6.

13.   Asgari S. MicroRNAs as regulators of insect host–pathogen interactions and immunity.

Adv In Insect Phys. 55: Elsevier; 2018. p. 19-45.

14.   Flynt AS, Greimann JC, Chung WJ, Lima CD, Lai EC. MicroRNA biogenesis via splicing and

exosome-mediated trimming in Drosophila. Mol Cell. 2010;38(6):900-7.

10.1016/j.molcel.2010.06.014

15.   Anderson JM, Sonenshine DE, Valenzuela JG. Exploring the mialome of ticks: an

annotated catalogue of midgut transcripts from the hard tick, Dermacentor variabilis (Acari:

Ixodidae). BMC Genomics. 2008;9(1):552. 10.1186/1471-2164-9-552

16.   Antunes S, Couto J, Ferrolho J, Sanches GS, Merino Charrez JO, De la Cruz Hernandez N,

et al. Transcriptome and Proteome Response of Rhipicephalus annulatus Tick Vector to Babesia

bigemina Infection. Front Physiol. 2019;10:318. 10.3389/fphys.2019.00318

17.   Popara M, Villar M. Proteomics characterization of tick-host-pathogen interactions.

Veterinary Infection Biology: Molecular Diagnostics and High-Throughput Strategies: Springer;

2015. p. 513-27.

18.   Villar M, Ayllon N, Alberdi P, Moreno A, Moreno M, Tobes R, et al. Integrated

Metabolomics, Transcriptomics and Proteomics Identifies Metabolic Pathways Affected by

Anaplasma phagocytophilum Infection in Tick Cells. Mol Cell Proteomics. 2015;14(12):3154-72.

10.1074/mcp.M115.051938

19.   Bechsgaard J, Vanthournout B, Funch P, Vestbo S, Gibbs RA, Richards S, et al.

Comparative genomic study of arachnid immune systems indicates loss of beta-1,3-glucanase-

related proteins and the immune deficiency pathway. J Evol Biol. 2016;29(2):277-91.

10.1111/jeb.12780

20.   Gulia-Nuss M, Nuss AB, Meyer JM, Sonenshine DE, Roe RM, Waterhouse RM, et al.

Genomic insights into the Ixodes scapularis tick vector of Lyme disease. Nature

communications. 2016;7(1):1-13.

21.   Rosa RD, Capelli-Peixoto J, Mesquita RD, Kalil SP, Pohl PC, Braz GR, et al. Exploring the

immune signalling pathway-related genes of the cattle tick Rhipicephalus microplus: from

molecular characterization to transcriptional profile upon microbial challenge. Dev Comp

Immunol. 2016;59:1-14.

22.   Bensaci M, Bhattacharya D, Clark R, Hu LT. Oral vaccination with vaccinia virus

expressing the tick antigen subolesin inhibits tick feeding and transmission of Borrelia

burgdorferi. Vaccine. 2012;30(42):6040-6. 10.1016/j.vaccine.2012.07.053

23.   Narasimhan S, Kurokawa C, Diktas H, Strank NO, Černý J, Murfin K, et al. Ixodes

scapularis saliva components that elicit responses associated with acquired tick-resistance.

Ticks Tick Borne Dis. 2020;11(3):101369.

24.   Ribeiro JM, Alarcon-Chaidez F, Francischetti IM, Mans BJ, Mather TN, Valenzuela JG, et

al. An annotated catalog of salivary gland transcripts from Ixodes scapularis ticks. Insect

Biochem Mol Biol. 2006;36(2):111-29. 10.1016/j.ibmb.2005.11.005

25.   Patrick CD, Hair JA. Laboratory rearing procedures and equipment for multi-host ticks

(Acarina: Ixodidae). J Med Entomol. 1975;12(3):389-90. 10.1093/jmedent/12.3.389

26.   Karim S, Ramakrishnan VG, Tucker JS, Essenberg RC, Sauer JR. Amblyomma americanum

salivary glands: double-stranded RNA-mediated gene silencing of synaptobrevin homologue

and inhibition of PGE2 stimulated protein secretion. Insect Biochem Mol Biol. 2004;34(4):407-

13. 10.1016/j.ibmb.2004.01.005

27.   Kumar D, Budachetri K, Meyers VC, Karim S. Assessment of tick antioxidant responses to

exogenous oxidative stressors and insight into the role of catalase in the reproductive fitness of

the Gulf Coast tick, Amblyomma maculatum. Insect Mol Biol. 2016;25(3):283-94.

28.   Embers ME, Grasperge BJ, Jacobs MB, Philipp MT. Feeding of ticks on animals for

transmission and xenodiagnosis in Lyme disease research. J Vis Exp. 2013(78):e50617.

10.3791/50617

29.   Kumar D, Embers M, Mather TN, Karim S. Is selenoprotein K required for Borrelia

burgdorferi infection within the tick vector Ixodes scapularis? Parasites & vectors. 2019;12(1):1-

30.   Stone BL, Russart NM, Gaultney RA, Floden AM, Vaughan JA, Brissette CA. The Western

progression of lyme disease: infectious and Nonclonal Borrelia burgdorferi Sensu Lato

populations in Grand Forks County, North Dakota. Appl Environ Microbiol. 2015;81(1):48-58.

10.1128/AEM.02422-14

31.   Friedländer MR, Chen W, Adamidi C, Maaskola J, Einspanier R, Knespel S, et al.

Discovering microRNAs from deep sequencing data using miRDeep. Nat Biotechnol.

2008;26(4):407-15.

32.   Friedlander MR, Mackowiak SD, Li N, Chen W, Rajewsky N. miRDeep2 accurately

identifies known and hundreds of novel microRNA genes in seven animal clades. Nucleic Acids

Res. 2012;40(1):37-52. 10.1093/nar/gkr688

33.   Munderloh UG, Kurtti TJ. Formulation of medium for tick cell culture. Exp Appl Acarol.

1989;7(3):219-29. 10.1007/BF01194061

34.   Obonyo M, Munderloh UG, Fingerle V, Wilske B, Kurtti TJ. Borrelia burgdorferi in tick cell

culture modulates expression of outer surface proteins A and C in response to temperature. J

Clin Microbiol. 1999;37(7):2137-41. 10.1128/JCM.37.7.2137-2141.1999

35.   Li Y, Andrade J. DEApp: an interactive web interface for differential expression analysis

of next generation sequence data. Source Code Biol Med. 2017;12(1):2. 10.1186/s13029-017-

0063-4

36.   Sturm M, Hackenberg M, Langenberger D, Frishman D. TargetSpy: a supervised machine

learning approach for microRNA target prediction. BMC Bioinformatics. 2010;11(1):292.

10.1186/1471-2105-11-292

37.   John B, Enright AJ, Aravin A, Tuschl T, Sander C, Marks DS. Human MicroRNA targets.

PLoS Biol. 2004;2(11):e363. 10.1371/journal.pbio.0020363

38.   Kertesz M, Iovino N, Unnerstall U, Gaul U, Segal E. The role of site accessibility in

microRNA target recognition. Nat Genet. 2007;39(10):1278-84. 10.1038/ng2135

39.   Aparicio-Puerta E, Lebron R, Rueda A, Gomez-Martin C, Giannoukakos S, Jaspez D, et al.

sRNAbench and sRNAtoolbox 2019: intuitive fast small RNA profiling and differential

expression. Nucleic Acids Res. 2019;47(W1):W530-W5. 10.1093/nar/gkz415

40.   Min H, Yoon S. Got target?: computational methods for microRNA target prediction and

their extension. Exp Mol Med. 2010;42(4):233-44.

41.   Franceschini A, Szklarczyk D, Frankild S, Kuhn M, Simonovic M, Roth A, et al. STRING

v9.1: protein-protein interaction networks, with increased coverage and integration. Nucleic

Acids Res. 2013;41(Database issue):D808-15. 10.1093/nar/gks1094

42.   Toronen P, Medlar A, Holm L. PANNZER2: a rapid functional annotation web server.

Nucleic Acids Res. 2018;46(W1):W84-W8. 10.1093/nar/gky350

43.   Ye J, Zhang Y, Cui H, Liu J, Wu Y, Cheng Y, et al. WEGO 2.0: a web tool for analyzing and

plotting GO annotations, 2018 update. Nucleic Acids Res. 2018;46(W1):W71-W5.

10.1093/nar/gky400

44.   Barrero RA, Keeble-Gagnere G, Zhang B, Moolhuijzen P, Ikeo K, Tateno Y, et al.

Evolutionary conserved microRNAs are ubiquitously expressed compared to tick-specific

miRNAs in the cattle tick Rhipicephalus (Boophilus) microplus. BMC Genomics. 2011;12(1):328.

10.1186/1471-2164-12-328

45.   Hackenberg M, Langenberger D, Schwarz A, Erhart J, Kotsyfakis M. In silico target

network analysis of de novo-discovered, tick saliva-specific microRNAs reveals important

combinatorial effects in their interference with vertebrate host physiology. RNA.

2017;23(8):1259-69.

46.   Nawaz M, Malik MI, Zhang H, Gebremedhin MB, Cao J, Zhou Y, et al. miRNA profile of

extracellular vesicles isolated from saliva of Haemaphysalis longicornis tick. Acta Trop.

2020;212:105718. 10.1016/j.actatropica.2020.105718

47.   Colombo M, Raposo G, Thery C. Biogenesis, secretion, and intercellular interactions of

exosomes and other extracellular vesicles. Annu Rev Cell Dev Biol. 2014;30:255-89.

10.1146/annurev-cellbio-101512-122326

48.   Coakley G, Maizels RM, Buck AH. Exosomes and Other Extracellular Vesicles: The New

Communicators in Parasite Infections. Trends Parasitol. 2015;31(10):477-89.

10.1016/j.pt.2015.06.009

49.   Huang T, Xu D, Zhang X. Characterization of host microRNAs that respond to DNA virus

infection in a crustacean. BMC Genomics. 2012;13(1):1-10.

50.   Huang T, Zhang X. Functional analysis of a crustacean microRNA in host-virus

interactions. J Virol. 2012;86(23):12997-3004. 10.1128/JVI.01702-12

51.   Hussain M, Torres S, Schnettler E, Funk A, Grundhoff A, Pijlman GP, et al. West Nile virus

encodes a microRNA-like small RNA in the 3' untranslated region which up-regulates GATA4

mRNA and facilitates virus replication in mosquito cells. Nucleic Acids Res. 2012;40(5):2210-23.

52.   Xu H, Jiang Y, Xu X, Su X, Liu Y, Ma Y, et al. Inducible degradation of lncRNA Sros1

promotes IFN-gamma-mediated activation of innate immune responses by stabilizing Stat1

mRNA. Nat Immunol. 2019;20(12):1621-30. 10.1038/s41590-019-0542-7

53.   Ramachandra RN, Wikel SK. Effects of Dermacentor andersoni (Acari: Ixodidae) salivary

gland extracts on Bos indicus and B. taurus lymphocytes and macrophages: in vitro cytokine

elaboration and lymphocyte blastogenesis. J Med Entomol. 1995;32(3):338-45.

10.1093/jmedent/32.3.338

54.   Urioste S, Hall LR, Telford SR, 3rd, Titus RG. Saliva of the Lyme disease vector, Ixodes

dammini, blocks cell activation by a nonprostaglandin E2-dependent mechanism. J Exp Med.

1994;180(3):1077-85. 10.1084/jem.180.3.1077

55.   Wang F, Gong H, Zhang H, Zhou Y, Cao J, Zhou J. Lipopolysaccharide-Induced Differential

Expression of miRNAs in Male and Female Rhipicephalus haemaphysaloides Ticks. PLoS One.

2015;10(10):e0139241. 10.1371/journal.pone.0139241

56.   Bond D, Foley E. A quantitative RNAi screen for JNK modifiers identifies Pvr as a novel

regulator of Drosophila immune signaling. PLoS Pathog. 2009;5(11):e1000655.

57.   Fullaondo A, Lee SY. Identification of putative miRNA involved in Drosophila

melanogaster immune response. Dev Comp Immunol. 2012;36(2):267-73.

10.1016/j.dci.2011.03.034

58.   Kung F, Anguita J, Pal U. Borrelia burgdorferi and tick proteins supporting pathogen

persistence in the vector. Future Microbiol. 2013;8(1):41-56. 10.2217/fmb.12.121

59.   Artigas-Jeronimo S, Alberdi P, Villar Rayo M, Cabezas-Cruz A, Prados PJE, Mateos-

Hernandez L, et al. Anaplasma phagocytophilum modifies tick cell microRNA expression and

upregulates isc-mir-79 to facilitate infection by targeting the Roundabout protein 2 pathway.

Sci Rep. 2019;9(1):9073. 10.1038/s41598-019-45658-2

60.   Zhang X, Zheng Y, Jagadeeswaran G, Ren R, Sunkar R, Jiang H. Identification of conserved

and novel microRNAs in Manduca sexta and their possible roles in the expression regulation of

immunity-related genes. Insect Biochem Mol Biol. 2014;47:12-22.

61.   Boulias K, Horvitz HR. The C. elegans microRNA mir-71 acts in neurons to promote

germline-mediated longevity through regulation of DAF-16/FOXO. Cell Metab. 2012;15(4):439-

62.   Avila-Bonilla RG, Yocupicio-Monroy M, Marchat LA, De Nova-Ocampo MA, Del Angel

RM, Salas-Benito JS. Analysis of the miRNA profile in C6/36 cells persistently infected with

dengue virus type 2. Virus Res. 2017;232:139-51. 10.1016/j.virusres.2017.03.005

63.   Liu Y, Zhou Y, Wu J, Zheng P, Li Y, Zheng X, et al. The expression profile of Aedes

albopictus miRNAs is altered by dengue virus serotype-2 infection. Cell Biosci. 2015;5(1):1-11.

64.   Ramasamy E, Taank V, Anderson JF, Sultana H, Neelakanta G. Repression of tick

microRNA-133 induces organic anion transporting polypeptide expression critical for

Anaplasma phagocytophilum survival in the vector and transmission to the vertebrate host.

PLoS Genet. 2020;16(7):e1008856. 10.1371/journal.pgen.1008856

65.   Khanal S, Sultana H, Catravas JD, Carlyon JA, Neelakanta G. Anaplasma phagocytophilum

infection modulates expression of megakaryocyte cell cycle genes through phosphatidylinositol-

3-kinase signaling. PLoS One. 2017;12(8):e0182898.

66.   Sultana H, Neelakanta G, Kantor FS, Malawista SE, Fish D, Montgomery RR, et al.

Anaplasma phagocytophilum induces actin phosphorylation to selectively regulate gene

transcription in Ixodes scapularis ticks. J Exp Med. 2010;207(8):1727-43. 10.1084/jem.20100276

67.   Taank V, Dutta S, Dasgupta A, Steeves TK, Fish D, Anderson JF, et al. Human rickettsial

pathogen modulates arthropod organic anion transporting polypeptide and tryptophan

pathway for its survival in ticks. Sci Rep. 2017;7(1):13256. 10.1038/s41598-017-13559-x

68.   Turck JW, Taank V, Neelakanta G, Sultana H. Ixodes scapularis Src tyrosine kinase

facilitates Anaplasma phagocytophilum survival in its arthropod vector. Ticks Tick Borne Dis.

2019;10(4):838-47. 10.1016/j.ttbdis.2019.04.002

69.   Kaneda K, Masuzawa T, Yasugami K, Suzuki T, Suzuki Y, Yanagihara Y. Glycosphingolipid-

binding protein of Borrelia burgdorferi sensu lato. Infect Immun. 1997;65(8):3180-5.

10.1128/iai.65.8.3180-3185.1997

70.   O'Neal AJ, Butler LR, Rolandelli A, Gilk SD, Pedra JH. Lipid hijacking: a unifying theme in

vector-borne diseases. Elife. 2020;9:e61675. 10.7554/eLife.61675

71.   Pal U, Kitsou C, Drecktrah D, Yaş ÖB, Fikrig E. Interactions Between Ticks and Lyme

Disease Spirochetes. *Curr Issues Mol Biol*. 2021;42:113-144. doi:10.21775/cimb.042.113

72.   Piesman, J. (1993). Dynamics of Borrelia burgdorferi transmission by nymphal Ixodes

dammini ticks. J Infect Dis 167, 1082-1085. doi: 10.1093/infdis/167.5.1082

73.   Piesman, J., Schneider, B.S., and Zeidner, N.S. (2001). Use of quantitative PCR to measure

density of Borrelia burgdorferi in the midgut and salivary glands of feeding tick vectors. J Clin

Microbiol 39, 4145-4148. doi: 10.1128/JCM.39.11.4145-4148.2001.

74.   Pal, U., and Fikrig, E. (2003). Adaptation of Borrelia burgdorferi in the vector and

vertebrate host. Microbes Infect 5, 659-666. doi: 10.1016/ s1286-4579(03)00097-2.

75.   Fikrig, E., and Narasimhan, S. (2006). Borrelia burgdorferi--traveling incognito? Microbes

Infect 8, 1390-1399. https://doi.org/ S1286-4579(06)00046-3 [pii] 10.1016/j.micinf.2005.12.022

76. Ueti, M.W., Reagan, J.O., Jr., Knowles, D.P., Jr., Scoles, G.A., Shkap, V., and Palmer, G.H.

(2007). Identification of midgut and salivary glands as specific and distinct barriers to efficient

tick-borne transmission of Anaplasma marginale. Infect Immun 75, 2959-2964.

77. Karim S, Essenberg RC, Dillwith JW, Tucker JS, Bowman AS, Sauer JR. Identification of

SNARE and cell trafficking regulatory proteins in the salivary glands of the lone star tick,

Amblyomma americanum (L.). *Insect Biochem Mol Biol*. 2002;32(12):1711-1721.

doi:10.1016/s0965-1748(02)00111-x

78. Zia MF, Flynt AS. Detection and Verification of Mammalian Mirtrons by Northern

Blotting. *Methods Mol Biol*. 2018;1823:209-219. doi:10.1007/978-1-4939-8624-8_16

79. Bartel DP. Metazoan MicroRNAs. *Cell*. 2018;173(1):20-51. doi:10.1016/j.cell.2018.03.006

80. Lambertz U, Oviedo Ovando ME, Vasconcelos EJ, Unrau PJ, Myler PJ, Reiner NE. Small

RNAs derived from tRNAs and rRNAs are highly enriched in exosomes from both old and new

world Leishmania providing evidence for conserved exosomal RNA Packaging. *BMC Genomics*.

2015;16(1):151. Published 2015 Mar 5. doi:10.1186/s12864-015-1260-7

81. Neelakanta G, Sultana H, Fish D, Anderson JF, Fikrig E. Anaplasma phagocytophilum

induces Ixodes scapularis ticks to express an antifreeze glycoprotein gene that enhances their

survival in the cold. *J Clin Invest*. 2010;120(9):3179-3190. doi:10.1172/JCI42868

82. Wu Z, He B, He J, Mao X. Upregulation of miR-153 promotes cell proliferation via downregulation of

the PTEN tumor suppressor gene in human prostate cancer. *Prostate*. 2013. doi:10.1002/pros.22600

83. Huang T, Xu D, Zhang X. Characterization of host microRNAs that respond to DNA virus infection in

a crustacean. *BMC Genomics*. 2012. doi:10.1186/1471-2164-13-159

84. Huang T, Zhang X. Functional Analysis of a Crustacean MicroRNA in Host-Virus Interactions. *J Virol*.

2012. doi:10.1128/jvi.01702-12

85. Liu W, Hao Z, Huang L, et al. Comparative expression profile of microRNAs in Anopheles

anthropophagus midgut after blood-feeding and Plasmodium infection. *Parasites and Vectors*. 2017.

doi:10.1186/s13071-017-2027-6

86.   Hussain M, Torres S, Schnettler E, et al. West Nile virus encodes a microRNA-like small RNA in the

3' untranslated region which up-regulates GATA4 mRNA and facilitates virus replication in mosquito

cells. *Nucleic Acids Res*. 2012;40(5):2210-2223. doi:10.1093/nar/gkr848

87.   Xu H, Jiang Y, Xu X, et al. Inducible degradation of lncRNA Sros1 promotes IFN-γ-mediated

activation of innate immune responses by stabilizing Stat1 mRNA. *Nat Immunol*. 2019.

doi:10.1038/s41590-019-0542-

88.   Artigas-Jerónimo S, Alberdi P, Villar Rayo M, et al. Anaplasma phagocytophilum modifies tick cell

microRNA expression and upregulates isc-mir-79 to facilitate infection by targeting the Roundabout

protein 2 pathway. *Sci Rep*. 2019. doi:10.1038/s41598-019-45658-2

89.   Freitak D, Knorr E, Vogel H, Vilcinskas A. Gender- and stressor-specific microRNA expression in

Tribolium castaneum. *Biol Lett*. 2012. doi:10.1098/rsbl.2012.0273

90.   Wang F, Gong H, Zhang H, Zhou Y, Cao J, Zhou J. Lipopolysaccharide-induced differential expression

of miRNAs in male and female Rhipicephalus haemaphysaloides ticks. *PLoS One*. 2015.

doi:10.1371/journal.pone.0139241

91.   Zhang X, Zheng Y, Jagadeeswaran G, Ren R, Sunkar R, Jiang H. Identification of conserved and novel

microRNAs in Manduca sexta and their possible roles in the expression regulation of immunity-related

genes. *Insect Biochem Mol Biol*. 2014. doi:10.1016/j.ibmb.2014.01.008

92.   Li R, Huang Y, Zhang Q, Zhou H, Jin P, Ma F. The miR-317 functions as a negative regulator of Toll

immune response and influences Drosophila survival. *Dev Comp Immunol*. 2019.

doi:10.1016/j.dci.2019.01.012

93.   Pushpavalli SNCVL, Sarkar A, Bag I, et al. Argonaute-1 functions as a mitotic regulator by

controlling Cyclin B during Drosophila early embryogenesis. *FASEB J*. 2014. doi:10.1096/fj.13-231167

94.   Fricke C, Green D, Smith D, Dalmay T, Chapman T. MicroRNAs influence reproductive responses by

females to male sex peptide in drosophila melanogaster. *Genetics*. 2014.

95.   Zhou Y, He Y, Wang C, Zhang X. Characterization of miRNAs from hydrothermal vent 20 Rimicaris

exoculata. *Mar Genomics*. 2015. doi:10.1016/j.margen.2015.09.004

96.   Hermance ME, Widen SG, Wood TG, Thangamani S. Ixodes scapularis salivary gland microRNAs are

differentially expressed during Powassan virus transmission. *Sci Rep*. 2019. doi:10.1038/s41598-019-

49572-5

97.   Boulias K, Horvitz HR. The C. elegans MicroRNA mir-71 acts in neurons to promote germline-

mediated longevity through regulation of DAF-16/FOXO. *Cell Metab*. 2012.

doi:10.1016/j.cmet.2012.02.014

98.   Mukherjee K, Vilcinskas A. Development and immunity-related microRNAs of the

lepidopteran model host Galleria mellonella. *BMC Genomics*. 2014. doi:10.1186/1471-2164-15-705

99.   Labunskyy VM, Hatfield DL, Gladyshev VN. Selenoproteins: Molecular Pathways and

Physiological Roles. *Physiol Rev*. 2014;94(3):739-777. doi:10.1152/physrev.00039.2013

100.   He Y, Sun Y, Zhang X. Noncoding miRNAs bridge virus infection and host autophagy in shrimp in

vivo. *FASEB J*. 2017. doi:10.1096/fj.201601141RR

101.   Liu Y, Zhou Y, Wu J, et al. The expression profile of Aedes albopictus miRNAs is altered by dengue

virus serotype-2 infection. *Cell Biosci*. 2015. doi:10.1186/s13578-015-0009-y

102.   Avila-Bonilla RG, Yocupicio-Monroy M, Marchat LA, De Nova-Ocampo MA, del Ángel RM, Salas-

Benito JS. Analysis of the miRNA profile in C6/36 cells persistently infected with dengue virus type 2.

Virus Res. 2017;232:139-151. doi:10.1016/j.virusres.2017.03.005

103.   Barrero RA, Keeble-Gagnère G, Zhang B, et al. Evolutionary conserved microRNAs are

ubiquitously expressed compared to tick-specific miRNAs in the cattle tick Rhipicephalus (Boophilus)

microplus. *BMC Genomics*. 2011. doi:10.1186/1471-2164-12-328

712
713