**Combined nanometric and phylogenetic analysis of unique endocytic compartments in**

**Giardia lamblia sheds light on the evolution of endocytosis in Fornicata**

Rui Santos[1,8], Ásgeir Ástvaldsson[2,9], Shweta V. Pipaliya[3], Jon Paulin Zumthor[5], Joel B. Dacks[3,4],

Staffan Svärd[2], Adrian B. Hehl[1*], Carmen Faso[6,7*†]

[1]Institute of Parasitology, University of Zürich, Winterthurerstrasse 266a 8057 Zürich
[2]Department of Cell and Molecular Biology, University of Uppsala, Husargatan 3, 752 37
Uppsala
[3]Division of Infectious Diseases, Department of Medicine, University of Alberta, Edmonton,
Alberta, Canada
[4]Institute of Parasitology, Biology Centre, CAS, v.v.i. Branisovska 31, 370 05 Ceske Budejovice,
Czech Republic
[5]Amt für Lebensmittelsicherheit und Tiergesundheit Graubünden, Chur
[6]Institute of Cell Biology, University of Bern, Bern, Switzerland
[7]Multidisciplinary Center for Infectious Diseases, University of Bern, Bern, Switzerland
[8] Institute of Anatomy, University of Zürich, Winterthurerstrasse 190 8057 Zürich
[9] Department of Microbiology, National Veterinary Institute, 751 23 Uppsala
20

*These authors contributed equally to this work.

†Author for correspondence: Carmen Faso (carmen.faso@unibe.ch)

RS ORCID iD: 0000-0001-6003-9079

CF ORCID iD: 0000-0002-1831-9365

ABH ORCID iD: 0000-0002-2110-4445

SVP ORCID iD: 0000-0003-0630-4120

JBD ORCID iD: 0000-0003-4520-5694

SS ORCID iD: 0000-0002-7392-1746

**Keywords:** endocytosis, Super Resolution Microscopy (SRM), volumetric Electron Microscopy,

Peripheral Endocytic Compartments (PECs), convergent evolution, metamonada, *Giardia*,

*Tritrichomonas*, *Spironucleus,* peripheral vacuoles

**ABSTRACT**

*Giardia lamblia*, a parasitic protist of the metamonada supergroup, has evolved one of the most diverged endocytic compartment systems investigated so far. Peripheral endocytic compartments, currently known as peripheral vesicles or vacuoles (PVs), perform bulk uptake of fluid phase material which is then digested and sorted either to the cell cytosol or back to the extracellular space. Here, we present a quantitative morphological characterization of these organelles using volumetric electron microscopy and super-resolution microscopy (SRM). We defined a morphological classification for the heterogenous population of PVs and performed a comparative analysis of PVs and endosome-like organelles in representatives of phylogenetically-related taxa, *Spironucleus spp.* and *Tritrichomonas foetus*. To investigate the as-yet insufficiently understood connection between PVs and clathrin assemblies in *G. lamblia*, we further performed an in-depth search for two key elements of the endocytic machinery, clathrin heavy chain (CHC) and clathrin light chain (CLC) across different lineages in Metamonada. Our data point to the loss of a *bona fide* CLC in the last Fornicata common ancestor (LFCA) with the emergence of a protein analogous to CLC (*Gl*ACLC) in the *Giardia* genus. Taken together, this provides the first comprehensive nanometric view of *Giardia*'s endocytic system architecture and sheds light on the evolution of GLACLC analogues in the Fornicata supergroup and, specific to Giardia, as a possible adaptation to the formation and maintenance of stable clathrin assemblies at PVs.

**INTRODUCTION**

Endomembrane compartments, while present in a few prokaryotic lineages (Heimerl et al. 2017), have evolved and greatly diversified across eukaryotic lineages. A fundamental task performed by some membrane-bounded organelles is endocytosis – the controlled and directed uptake of nutrients and other materials from the extracellular space into the cell by membrane transport. Fluid phase or receptor-bound material at the cell surface is internalised via invaginations and formation of vesicles at the plasma membrane, mediated by clathrin-coated vesicles (CCVs) (Robinson 2015; Kaksonen and Roux 2018). In turn, CCVs fuse with early endosomes which mature into late endosomes upon lysosome fusion (Huotari and Helenius 2011; Naslavsky and Caplan 2018). Clathrin coats are also involved in protein secretion forming exocytic transport vesicles derived from the trans-Golgi compartment and play a role in Golgi apparatus reassembly after mitotic cell division (Radulescu et al. 2007; Jaiswal et al. 2009).

Evolutionary adaptations of endocytic pathways to specific environmental niches and nutrient sources are especially relevant to species adopting a fully parasitic or commensal lifestyle (Poulin and Randhawa 2015; Jackson et al. 2016; Dacks and Field 2018; Pipaliya et al. 2021).

Within the extant Metamonada supergroup (Hampl et al. 2009; Hug et al. 2016; Burki et al. 2020),

the parasitic protist *Giardia lamblia* (syn.: *intestinalis* or *duodenalis*) evolved a distinct endocytic

pathway, which reflects its adaptation to the host intestinal lumen environment. This unicellular

parasite is responsible for >300 million cases annually of water-borne infections causing

gastroenteritis – giardiasis – with higher incidence in low to middle income countries (Caccìo and

Ryan 2008) (Caccìo and Ryan 2008). *Giardia* is the etiological agent for symptomatic

gastroenteritis in 15% of children in developing countries, with 1-2% fatality in children with

severely compromised health status (Kotloff et al. 2013; Lanata et al. 2013). There is a strong

association of *Giardia* infections with chronic conditions such as irritable bowel syndrome or

inflammatory bowel disease as a result of intestinal barrier function disruption and microbiome

dysregulation (Allain et al. 2017; Fekete et al. 2021).

Cellular evolution of the *Giardia* genus as an obligate parasite adapted to the small

intestinal niche of vertebrates is characterised by a reduction in subcellular compartment diversity.

Peroxisomes, late endosomes and a permanent stacked Golgi complex have not been detected

in *Giardia* (Faso and Hehl 2011). Two nuclei (Benchimol 2005), an extensive endoplasmic

reticulum (ER) (Soltys et al. 1996), highly reduced mitochondria-derived organelles – the

mitosomes (Tovar et al. 2003) - and peripheral vesicles (PVs) (Lanfredi-Rangel et al. 1998) are

the only membrane-bounded organelles with conserved morphology and function documented in

the *Giardia* trophozoite (Marti, Regös, et al. 2003; Zumthor et al. 2016; Cernikova et al. 2020).

The complex array of PV organelles as the only documented endocytic membrane

compartment system in *Giardia* is responsible for uptake of fluid-phase and membrane-bound

material (Rivero et al. 2011; Zumthor et al. 2016; Frontera et al. 2018). These organelles acidify

and presumably serve as digestive compartments with capability for sorting after processing,

similar to early and late endosomes and lysosomes (Lanfredi-Rangel et al. 1998). The static

system of PV organelles (Abodeely et al. 2009; Zumthor et al. 2016) is restricted to the peripheral

cortex below the plasma membrane (PM) of the *Giardia* trophozoite. PV morphology was

investigated using high-resolution electron microscopy serial sectioning and three-dimensional

reconstruction (Zumthor et al. 2016). These organelles were resolved as tubular structures in

close proximity to funnel-shaped invaginations of the PM (Zumthor et al. 2016). In the same report,

the presence of focal accumulations of clathrin heavy chain (CHC) molecules and their main

interactors, collectively termed clathrin assemblies, were demonstrated at PM and PV membrane

interfaces.. The function of these stable focal assemblies as well as additional components at the

interface of the PV membranes and the PM, has proved elusive (Zumthor et al. 2016). However,

transient association of several members of the family of adaptor proteins (AP) suggests a role in

dynamic processes linked to uptake of fluid-phase and receptor-bound material into PVs (Zumthor
et al. 2016; Cernikova et al. 2020). Our current working model for bulk fluid-phase uptake of
extracellular material into PVs invokes a "kiss and flush" mechanism, whereby acidified PV
membranes and the PM transiently form channels at invaginations allowing exchange between
PV lumen content and the extracellular space at regular intervals. Endocytosed material is
digested in the sealed-off acidified PVs and transported towards the cell interior while residual
material and waste is flushed to the extracellular space in the next round of membrane fusion,
thus completing the PV cycle (Zumthor et al. 2016; Cernikova et al. 2020).
116

In this report, we address open questions concerning *G. lamblia's* PV ultrastructure and
its associated molecular machinery in a comparative approach with one closely and one more
distantly related fornicata and metamonada species, *Spironucleus sp*. and *Tritrichomonas foetu*s,
respectively. Using volumetric electron microscopy and super resolution light microscopy we
developed a classification of PVs based on organelle morphology. Comparative analysis of
*Giardia's* PVs with endocytic compartments of fornicata and metamonada species emphasized
the genus-specific nature of the *Giardia* endocytic system architecture. In addition, using a
combination of co-immunoprecipitation and phylogeny techniques, we provide evidence that   a
proposed diverged clathrin light chain previously named *GlCLC* (Zumthor et al. 2016) is unique
to the *Giardia* genus and evolved *de novo* as structurally analogous to CLC after loss of a *bona*
*fide* CLC in the last Fornicata common ancestor (LFCA). Taken together, the emergence of a
unique and highly polymorphic endocytic system such as the one found in the genus *Giardia* is
linked to the proposed convergent evolution of an independent CLC analogue concomitant with
loss of a mostly conserved CLC orthologue.
131

**RESULTS AND DISCUSSION**
**Complete FIB-SEM rendering of a *G. lamblia* trophozoite reveals a novel landscape of**
**vesicular compartments**
Volumetric scanning electron microscopy (vSEM) is currently considered the gold
standard for the determination of biological ultrastructure (Titze and Genoud 2016). Focused ion
beam electron scanning microscopy (FIB-SEM) uses a beam of gallium ions to mill and image
consecutive layers of an embedded biological sample, resulting in a voxel resolution as low as 1-
2 nm (Kizilyaprak et al. 2014). This technique allows for sectioning and imaging of entire cells
(Wei et al. 2012). It was previously implemented for partial rendering of *G. lamblia* trophozoite
sections (Zumthor et al. 2016) and more recently in (Tůmová et al. 2020). Here we sectioned for

the first time a complete *G. lamblia* trophozoite at a voxel resolution of 125 nm$^3$ (5x5x5 nm) after

high pressure freezing (HPF) and embedding. Images representing the sagittal plane adjacent to

the cell centre (figure 1A, 1D and supplementary figure 1A and B) show all the major cell

compartments such as the nuclei (Figure 1A, N), the endoplasmic reticulum (Figure 1A, ER),

mitosomes (Figure 1A and C, m) and elements of the cytoskeleton: axonemes (Ax), funis (F) and

the ventral disc (VD) (Figure 1D; Dawson 2010). Two different types of small cytoplasmic

organelles are observed: PVs (arrow heads) with heterogenous morphology and smaller and

electron-dense membrane vesicles of uniform size and appearance which we termed Small

Vesicles (SVs; asterisks) (figure 1B and E).

After serial sectioning and alignment with TrakEM (Cardona et al. 2012), we used the

supervised machine learning (ML) tool Ilastik for pixel based image segmentation of PVs and SVs

(Sommer et al. 2011; Berg et al. 2019). The algorithm collection performs supervised learning

and recognition of patterns based on ground truth training provided by the user. Patterns are

sorted into classes. Once the algorithm is trained on a subset of image data, it is used to analyse

complete datasets and assigning features to different classes following a decision tree method

(Sommer and Gerlich 2013; Kan 2017). This process enabled the three-dimensional rendering of

selected trophozoite features: the complete cytoskeleton, the ER, PVs and mitosomes

(supplementary figure 1C). In addition, we were able to calculate the volume of the cell at 138

$\mu$m$^3$ as well as the average volume of mitosome organelles (N=14) at 0.001093±0.0005698$\mu$m$^3$

with a 95% confidence interval between [0.0007643, 0.001422] $\mu$m$^3$ (supplementary figure 1D).

Similarly, supervised ML assisted pixel segmentation and object clustering analysis

allowed identification of two statistically distinct morphological classes of PVs: spherical and

tubular/elongated PVs. Individual PV organelles of both classes (N=467) were rendered in three

dimensions (figure 1F and supplementary video 1). Spherical PVs average a volume of 9.243x10$^{-4}$

±3.322x10$^{-4}$ $\mu$m$^3$ in a 95% confidence interval between [9.022x10$^{-4}$; 9.022x10$^{-4}$] $\mu$m$^3$ while tubular

PVs average a volume of 1.067x10-3±3,322x10$^{-4}$ $\mu$m$^3$ with a 95% confidence interval between

[9.843x10$^{-4}$; 1.150x10$^{-3}$] $\mu$m$^3$, a statistically significant difference (t-student test, (p< 0.0001),

corroborating PV grouping in these two classes.  To further investigate morphological

heterogeneity of PVs, we analysed trophozoite ultrastructure using freeze fracture scanning

electron microscopy. We documented PV heterogeneity and the presence of spherical and tubular

PV forms (Supplementary Figure 2). Additional ultrastructural studies using transmission electron

microscopy (TEM) were consistent with this classification (Supplementary figure 3A and B).

We proceeded with the rendering of 269 SVs – small spherical vesicles, with distinctly

higher electron density than PVs and what could be a coat on the cytoplasmic side of the

delimiting membrane (figure 1G). SVs were also identified by TEM (Supplementary figure 3),
proximal to the PM. SVs average a volume of $2.525 \times 10^{-4} \pm 9.280 \times 10^{-5}$ $\mu m^3$ in a 95% confidence
interval between [$2.414 \times 10^{-4}$; $2.637 \times 10^{-4}$] $\mu m^3$ (figure 1G, box-plot on the left). This equals to an
average diameter of 77.23±9.666 nm in a 95% confidence interval between [76.07; 78.40] nm,
differing significantly from spherical PVs which average 120.1±9.507 nm in a 95% confidence
interval between [119.2; 121] nm (p< 0.0001) (figure 1G, box-plot on the right). Thus, there is
statistical support for SVs as a distinct category of membrane-bounded vesicles (Supplementary
figure 3A and C).
Taken together, these findings lead us to hypothesize that, unlike previously thought, PVs
are morphologically heterogenous and may comprise different functional categories (Poteryaev
et al. 2010; Hipolito et al. 2018; Suresh et al. 2020). However, these data are currently insufficient
to determine whether distinct morphologies correlate with distinct functions.
188
**Combining super-resolution microscopy with ML-assisted image analysis identifies three**
**classes of endocytic compartments in *G. lamblia* trophozoites**
FIB-SEM as a technique is not well-suited to the investigation of large cell numbers, and
TEM cannot readily provide 3D volumetric information on subcellular compartments. Hence, to
address PV heterogeneity in more detail, we continued our investigation of *Giardia* endocytic
compartments by Super-resolution Light Microscopy (SRM) techniques and ML assisted image
analysis of compartment shapes.
The dimensions of *Giardia* endocytic compartments are well below the diffraction limit of
conventional light microscopy (Combs and Shroff 2017). To overcome the Abbe diffraction barrier
we used stimulated emission depletion microscopy (STED), potentially achieving a lateral (x,y)
and axial (z) resolution of 25-50 nm and 60-100 nm, respectively. This technique decreases the
point spread function signal from the illuminated region (Klar et al. 2000; Willig et al. 2006) and
allows for accurate imaging of trophozoite PV lumina loaded with a highly photostable fluid phase
marker (10 kDa Dextran-Alexa Fluor 594) which is readily taken up into PVs via the fluid phase
endocytic pathway (Figure 2A and Supplemental Video 2).  In addition to spherical and tubular
PVs documented in FIB-SEM, using STED we also determined the presence of polymorphic
dextran-labelled organelles i.e., spherical PVs with elongated rods (figure 2A). All labelled PVs
were further analysed using the ML-assisted algorithm of the Ilastik program suite. We first
performed supervised pixel segmentation followed by supervised object classification. In this
second step, we defined and trained the classifier in three organelle morphologies: spherical,
tubular, and polymorphic. The latter comprised characteristics of both vesicular and tubular

classes, generally with spherical centres with attached tubular protrusions (Figure 2B). After
organelle classification we measured their projected areas. Spherical organelles (N =1684) have
an average projected area of 0.0205±0.0169 $\mu m^2$ with a 95% confidence interval between
[0.0197;0.0213] $\mu m^2$. Tubular endocytic organelles (N=835) present an average projected area of
0.0453±0.0278 $\mu m^2$ with a 95% confidence interval between [0.0435;0.0472] $\mu m^2$. Polymorphic
organelles (N=400) have an average projected area of 0.0981±0.0429 $\mu m^2$ with a 95% confidence
interval between [0.0939;0.102] $\mu m^2$. ANOVA analysis reveals that each of the three categories
is indeed significantly distinct ($p<0.0001$) based on projected surface area (Figure 2C). This lends
further support to the possibility that PV morphological heterogeneity may have functional
implications..

Although a STED microscopy-based approach clearly allows resolution of individual
organelles as small as PVs, the distinctly lower axial resolution remains limiting for three-
dimensional rendering of organelles. Therefore, to push the boundaries of resolution and to further
characterize PV morphology, we employed Single Molecule Localization Microscopy (SMLM)
(Kao et al. 1994; Huang et al. 2008; Jones et al. 2011).

*Giardia* PVs in trophozoites were loaded with a 10 kDa Dextran-Alexa Fluor 647 fluid
phase marker with a high degree of photostability to survive  repeated cycles of photoactivation
and excitation in SMLM experiments (Dempsey et al. 2011; Olivier et al. 2013).  After acquisition,
images were reconstructed using the ImageJ plugin ThunderStorm which performs signal centroid
calculation, image reconstruction and output (Schindelin et al. 2012; Ovesný et al. 2014). Dextran
uptake in PVs was confirmed using conventional widefield microscopy (Figure 3A). STORM
image reconstruction shows the subcellular distribution of the fluorescent marker and defines
individual organelle lumina (Figure 3B). A closer inspection revealed the presence of
morphologically distinct endocytic organelles as previously observed in our FIB-SEM and STED
datasets (Figure 3B, ROI and Supplementary Video 3). We again used the supervised ML-
assisted algorithm in Ilastik to classify the different morphologies. After a pixel segmentation
routine, we performed object classification using supervised ground truth training on subsets of
organelles images. Three categories of PVs were defined: spherical, tubular and polymorphic
(Figure 3C). To test whether the morphological categorization was consistent with categorization
based on organelle volume, we calculated the average lumina volumes of >4000 organelles from
the three PV categories. ANOVA testing of organelle volumes for vesicular (0.00507±0.00336
$\mu m^3$, N=1989, 95% confidence interval: [0.00492;0.00522] $\mu m^{3)}$, tubular (0.0103±0.00925 $\mu m^3$
N=838, 95% confidence interval: [0.00967, 0.0109] $\mu m^3$), and polymorphic organelles

(0.0227±0.0214 $\mu m^3$ N=1494, 95% confidence interval: [0.0216, 0.0238] $\mu m^3$ confirmed
statistically significant (p <0.0001) morphological differences (Figure 3D and summarised in
in supplementary table 1) .

Taken together, the data generated using three distinct imaging techniques clearly
demonstrates PV heterogeneity which may be linked to distinct functions and/or maturation states
in this unique endocytic system.  To reflect this novel finding and considering that these endocytic
and peripherally localized organelles are neither proper vesicles nor canonical vacuoles, we
propose renaming PVs to peripheral endocytic compartments (PECs).

**252  Comparative analysis of endocytic and secretory organelles in *Giardia*, *Spironucleus* sp.**
**253  and *T. foetus***

*Giardia* spp. have evolved a unique cell architecture including a dedicated organelle for
attachment to the small intestinal epithelium – the ventral disk (VD) (Dawson 2010; Brown et al.
2016). In turn, this innovation defines a distinct dorso-ventral as well as antero-posterior
polarization of the flagellated trophozoite, marked by swimming directionality.  PVs/PECs localize
exclusively to the dome-shaped dorsal parasite PM except a small circular patch at the centre of
the VD called the bare zone (Zumthor et al. 2016; Cernikova et al. 2020). The result is a maximally
decentralized architecture of the *Giardia* endocytic system forming a single-layer interface of what
we now appreciate as 3  morphologically distinct organelle classes between the cell exterior and
the cytoplasm/ER (Abodeely et al. 2009; Zumthor et al. 2016). We asked whether this type of
decentralized sub-PM localisation and polymorphic morphology of endocytic compartments was
also represented in other tractable  related members of the Diplomonadida and phylogenetically
more distant metamonada lineages which do not have a VD, respectively *Spironucleus vortens*
and *Spironucleus salmonicida* and the parabasalid *Tritrichomonas foetus*

The diplomonads *Spironucleus vortens* and *Spironucleus salmonicida* are amongst the
closest tractable relatives of *G. lamblia* that can be grown axenically under similar conditions
(Paull and Matthews 2001; Jørgensen and Sterud 2007; Kolisko et al. 2008a; Xu et al. 2014; Xu
et al. 2016). Their endocytic compartments and machineries are partially characterized, with some
reports of large vacuolar structures detected by electron microscopy in trophozoites (Sterud and
Poynton 2002; Ástvaldsson et al. 2019). Unlike *Giardia*, both species lack dorso-ventral
polarization but display a distinct antero-posterior axis.  Putative endocytic organelles in *S.*
*vortens* have been detected by fluorescence microscopy of live and fixed cells after incubation
with fluorophore-coupled dextran (Zumthor et al. 2016). To further investigate these endocytic
compartments, we incubated *S. vortens* and *S. salmonicida* trophozoites with a 10 kDa Dextran-

TexasRed fluid phase marker (Figure 4). In stark contrast to the distinctly arrayed PV/PEC

labelling seen in *Giardia lamblia* (Figure 4A), confocal microscopy revealed the presence of

several dispersed labelled organelles in both *S. vortens* (Figure 4B) and *S. salmonicida* (Figure

4C). *Spironucleus spp.* display several relatively large globular membrane compartments, similar

to those observed in well-characterized model organisms (Huotari and Helenius 2011; Day et al.

2018) lacking a fixed subcellular localization. While *S. salmonicida* endocytic compartments

localise mostly at the cell periphery (Figure 4C), *S. vortens* organelles present both peripheral

and central localizations (Figure 4B and Supplementary Video 4). We also assessed endosome

morphology in *T. foetus* using the same labelled dextran-based approach. Similar to *Spironucleus*

species, *T. foetus* presents an antero-posterior axis but no attachment organelle nor dorso-ventral

polarization. Similar to *Spironucleus spp.*, *T. foetus* accumulated the endocytosed fluid phase

maker in several globular endocytic compartments (Figure 4D) consistent with previous reports

on vacuolar structures  identified in *T. foetus* by electron microscopy (Lealda et al. 1986). In our

ultrastructure observations we could not detect multivesicular bodies or vacuoles containing intra-

luminal vesicles in either *Spironucleus* spp.  or *T. foetus*.

Taken together, these data show how, in closely-related protozoa lacking dorso-ventral

polarization and a dedicated attachment organelle, endocytic organelles appear to have no

specific localization. This lends support to the notion that PV organelle architecture  is intimately

associated to the emergence of the VD, both as adaptations to the mammalian small intestine

niche (Zumthor et al., 2016).

To visualize and measure morphological parameters of *Spironucleus* and *T. foetus*

endocytic compartments, we performed 2D-STED imaging and transmission electron microscopy

(TEM). *S. vortens* cells loaded with 10 kDa Dextran-Alexa Fluor 594 showed accumulation of the

fluid phase marker in roughly spherical organelles (Figure 5A). Labelled endocytic vacuoles have

an average diameter of 468±206 nm (95% confidence interval [421;515] nm, N=10) (Figure 5A,

box-plot). Volumetric rendering of 3D reconstructed optical sections document the uniformly

globular morphology of these organelles (Supplementary Video 5). TEM analysis revealed an

ellipsoid shape of endocytic organelles in *S. vortens* with an average maximal diameter of

844±335 nm in a 95% confidence interval between [763;905] nm (Figure 5B, box-plot). The

dimensions measured in TEM represent those of the membrane-delimited organelle. In contrast,

the dimensions measured by STED represent a projection of the fluid phase marker distribution

within the available organelle lumen. The fact that the former (844±335 nm) is larger than the

latter (468±206 nm) indicates that these organelles contain additional cargo which prevents the

endocytosed fluid phase marker to distribute in the complete compartment volume delimited by

the organelle membranes. TEM investigation in *S. salmonicida* cells (Supplemental Figure 4A)

showed the presence of small globular vacuoles (V) close to the PM (Supplemental Figure 4B)

with an average diameter of 205±62.6 nm (N=114) in a 95% confidence interval of [193;217] nm

(Supplemental Figure 4E). These vacuoles are smaller than the ones found in *S. vortens*

(Supplementary Figure 4C,D,F, p-value < 0.0001). In these conditions, neither coated vesicles

nor a stacked Golgi apparatus could be documented in *S. vortens* or *S. salmonicida*.

2D-STED analysis of *T. foetus* cells incubated with 10 kDa Dextran-Alexa Fluor 594

revealed a roughly circular distribution of the marker within endocytic vacuoles (Figure 6A and

Supplemental Video 6) with an average maximal diameter of 517±251 nm (95% confidence

interval [455;580] nm, N=10) (Figure 6A, box-plot). TEM imaging revealed the presence of two

distinct classes of endosome-like vesicles (Figure 6B) based on electron density of the lumen.

Low-density vesicles were identified both at the cell periphery and in central areas termed

vacuoles (V); vesicles of higher electron density were previously identified as digestive vacuoles

(DVs) (Lealda et al. 1986) and contain structured material and membranes.  Analysis of TEM

micrographs showed that DVs are significantly larger than vacuoles, with an average diameter of

764±203 nm (N=50) (95% confidence interval [707;822] nm). Vacuoles in turn have an average

diameter of 246±100 nm (N=153) in (95% confidence interval [230;262] nm) (Figure 6B, box-plot).

Stacked Golgi organelles are abundant in TEM micrographs of *T. foetus* trophozoites, as

documented previously (Rosa et al. 2014) (Supplemental Figure 5A). Consistent with a more

canonical architecture of the membrane trafficking system in *T. foetus,* coated vesicles were

observed in the cytosol particularly in the vicinity of Golgi stacks (Supplemental Figure 5B) (Lealda

et al. 1986; Midlej et al. 2011; Schlacht et al. 2014). These vesicles averaged a diameter of

58.4±13.1 nm (N=128) (95% confidence interval [56.1;60.7] nm) corresponding to the size range

of clathrin coated vesicles (CCVs) (Traub 2011).

Finally, to probe the dynamics of endocytic compartments in *G. lamblia*, *S. vortens*, *S.*

*salmonicida* and *T. foetus*, cells were exposed to 10 kDa Dextran-TexasRed for 5, 10, 20 or 30

minutes, fixed chemically, and imaged by confocal microscopy (Figure 7). The number of *G.*

*lamblia* PECs labelled with the fluid-phase marker increased over time, with the label

accumulating strictly at the cell periphery (Figure 7A). In contrast, endocytic compartments in *S.*

*vortens* were first visualized  at the PM, and were then observed at more central locations of the

cell at later time points. Given the overall increase in fluorescent intensity and the motile nature

of these organelles, it appears there is a constant uptake of dextran over the analysed period

(Figure 7B). In these conditions, *S. salmonicida* and T. foetus vacuoles behave similarly since
both sets of organelles are diffused within the cell cytoplasm, with a steady decrease and then
marked increase  in dextran content (Figure 7C and D).
348
**GICHC foci associate with different classes of giardial PECs with different stoichiometry**
Previously, we established that *Giardia* clathrin heavy chain (*Gl*CHC) associates to
discrete static foci at the dorsal PM of trophozoites, in close proximity to PV/PECs. Further,
*Gl*CHC strongly interacts with a putative albeit highly diverged *Giardia* clathrin light chain
homologue previously named *Gl*CLC (Zumthor et al. 2016; Cernikova et al. 2020). We wondered
whether association to clathrin assemblies holds any relation to the heterogeneity of PV
morphology. To address this question, we used STED microscopy to investigate epitope-tagged
*Gl*CHC (*Gl*CHC-HA) deposition at distinct foci at the dorsal PM and the cell periphery, consistent
with PV/PECs location (Figure 8A). Segmentation of foci using a ML-assisted Ilastik tool, allowed
to determine the dimensions of *Gl*CHC at an average diameter of 134±36.6 nm (N = 4524) (95%
confidence interval [132;135] nm) (Figure 8B). Similar to GlCHC, subcellular distribution of
epitope-tagged *Gl*CLC (*Gl*CLC-HA) showed an identical pattern consistent with its demonstrated
direct interaction with *Gl*CHC (figure 8C) (Zumthor et al. 2016). Segmentation of foci using a ML-
assisted Ilastik tool, determined the dimensions of *Gl*CLC foci at an average diameter of 159±48.8
nm (N = 984) (95% confidence interval [156;162] nm) (Figure 8D),  Notably, the average size of
*Gl*CLC foci is larger than that of *Gl*CHC foci (p< 0.0001, t-student test) (Figure 8E).
Using STED microscopy, we further determined the number of *Gl*CHC foci showing signal
overlap with the 3 classes of  Dextran-Texas Red loaded PV/PECs (figure 8F). By calculating the
degree of signal overlap between *Gl*CHC foci and PEC lumina, we determined that spherical
PECs are associated with at most one *Gl*CHC focus with an average of 0.488±0.159 foci per
spherical PEC (95% confidence interval [0.403;0.573]). Tubular PECs associated with at least
one *Gl*CHC focus with an average of 1.15±0.287 foci per tubular PEC (95% confidence interval
[0.994;1.3]). Polymorphic PECs associated with 3 or more *Gl*CHC foci with an average of
3.85±1.14 foci per PEC (95% confidence interval [3.25;4.46]). Taken together, we find a directly
proportional and statistically-significant ratio of clathrin foci to PEC size and type (ANOVA; p-
value < 0.0001). This is in line with the notion that PV morphological heterogeneity is indeed
correlated with organelle functional diversity, as measured by association to clathrin assemblies.
376
**Pan-eukaryotic searches for CHC and CLC reveal loss of a *bona fide* CLC within the**
**fornicata lineage and the emergence of putative CLC analogues**

We previously proposed that PV/PEC localization at the dorsal PM of trophozoites and
evolution of the adhesive disc attachment organelle are interdependent adaptations to life in direct
contact with the host's gut epithelium (Zumthor et al. 2016). Furthermore, we found a direct
correlation between types of PV/PEC and the number of foci of clathrin assemblies. Given that
the role for clathrin assemblies in Giardia has not been elucidated and that the nature of *Gl*CLC
(*Gl*4259) as a clathrin light chain orthologue is dubious, we sought to shed light on the significance
of this correlation by investigating the distribution of both CHC and CLC orthologues in selected
eukaryotic lineages. To do this, we employed protein homology searches based on Hidden-
Markov Models (HMM) (Eddy 2011) using as query an alignment of canonical and documented
CHC or CLC sequences from several protozoa and metazoan species (Supplementary Tables 2
and 5) (Morgan et al. 2001; Kaksonen et al. 2005; Adung'a et al. 2013; Kirchhausen et al. 2014;
Karnkowska et al. 2016; Karnkowska et al. 2019, Füssy et al., 2021 ). In our search we considered
assembled read data from RNA-seq experiments (transcriptomics) as reliable as genomic
sequence data (Cheon et al. 2020). In this case, we used the reference CHC or CLC sequences
and performed tblastn searches. Nucleotide sequences from each reliable hit (lowest e-value)
were translated and subjected to a reciprocal blast-p analysis to validate protein identity.  We
found CHC homologues in all selected genomes and transcriptomes we searched, highlighting
the likely essential nature of CHC (Figure 9A and supplementary table 4 and 5).
*Gl*CHC is a clearly divergent ortholog compared to its counterpart in eukaryotic model organisms,
with only 24% amino acid identity to human CHC (Marti, Regös, et al. 2003). A domain analysis
of selected CHC sequences (supplementary figure 6 and supplemental table 10) reveals that
*Gl*CHC contains fewer α-helical domains than other analysed CHC sequences, further
highlighting its divergence. We also performed an in-depth search for the CHC triskelion
uncoating "QLMLT" motif which we documented previously to be missing in *Giardia* (Fotin et al.
2004; Rapoport et al. 2008; Zumthor et al. 2016). Notably, this motif appears to be only present
in Metazoa and in the closely related Filastera and Choanoflagellata (King et al. 2008; Fairclough
et al. 2013; Suga et al. 2013). In Fungi only a partial "L(M)TL"motif was identified and we were
unable to detect a conserved uncoating motif in CHC sequences of members of the
Archaeplastida, Amoebozoa or SAR supergroups (supplementary figure 7).
In stark contrast to CHC, the search for *bona fide* CLC sequences did not retrieve any
reliable predictions in available genomes and transcriptomes from species of the Fornicata
lineage, including the lineages Hexamitidae, Retortamonas and Carpediemonas-like organisms
(Xu et al. 2014; Leger et al. 2017; Tanifuji et al. 2018; Füssy et al. 2021; Salas-Leiva et al. 2021).
Importantly, this search did not return the putative, highly diverged *Gl*CLC  (Zumthor et al. 2016).

There are documented CLC orthologues in members of the Discoba, such as *Trypanosoma brucei* CLC (Tb927.10.14760) (Manna et al. 2017) and in the parabasalid *Trichomonas vaginalis* (TVAG_29749) (Carlton et al. 2007; Aurrecoechea et al. 2009). Furthermore, we readily identified a CLC homologue in *T. foetus* (gene accession OHT14195.1, forward HMMer e-value of 1.00E-26 and reverse Blastp e-value of 2.00E-11, returning the human CLC homologue) (Figure 9A). Therefore, while *bona fide* CLC orthologues can be readily identified in Discoba and Euglenozoa members and Preaxostyla – as in the metamonad *Monocercomonoides exilis* (Karnkowska et al. 2016) – no sequence could be found among members of the Fornicata lineage. Further, we were unable to identify a *bona fide* CLC sequence within the newly documented transcriptome of Hemimastigophora (Lax et al. 2018). These observations are in line with the notion that, unlike CHC, CLC is dispensable. This is also supported by failures to identify *bona fide* CLC in chromerids such as *Cryptosporidium parvum* and *Babesia bovis* (Woo et al. 2015) and in some ciliate lineages, such as *Tetrahymena thermophila* and *Paramecium caudatum* (Richardson and Dacks 2022).

Given that *Gl*CLC's predicted 3D structure is reminiscent of CLCs (Zumthor et al., 2016) but it could not be retrieved as related to a *bona fide* CLC, with its only known orthologue found in *Giardia muris* (Figure 9A), *Gl*CLC was further analysed using the HHPred suit, in the attempt to find distantly-related non-Giardia sequences (Zimmermann et al. 2017). This search retrieved no robust prediction for a non-Giardia sequence (supplemental table 9). Given that the degree of divergence is such that no reliable claim to orthology can currently be supported and no orthologue for *Gl*CLC can be found outside the *Giardia* genus, we propose the renaming of GlCLC to *Giardia lamblia* Analogous to Clathrin Light Chain- **GlACLC**- as a CLC structural analogue acquired and retained in the last *Giardia* common ancestor (LGCA). This appears to correlate with loss of a *bona fide* CLC with the last Fornicata common ancestor (LFCA).

To test the extent of environmental pressure on this protein family's evolution, we calculated synonymous vs non-synonymous mutation ratios ($\omega$ = ks/kn) for *Gl*ACLC homologues (supplemental figure 8). Interestingly, known sequences for all *Giardia* isolates present a $\omega < 1$ which indicates that current sequences are not under selective pressure to evolve.

To further investigate the structural analogy of *Gl*ACLC to canonical CLCs, we performed *in silico* modelling of its C-terminal domain () using the new standard in *ab initio* protein structure modelling – AlphaFold - based on deep-learning neural networks (Jumper et al. 2021; Tunyasuvunakool et al. 2021) (Figure 9B and supplemental table 11). Template modelling score independent of sequence (Tm-align) and Root Mean Square Deviation (RMSD) (Zhang and

Skolnick 2005; Kufareva and Abagyan 2012) values provide substantial evidence for structural
analogy of *Gl*ACLC and canonical CLCs, in line with previous observations (Zumthor et al. 2016).
The newly predicted structures for *Gl*ACL*C* have a stronger resemblance to the predicted
structure of a mammalian clathrin light chain (Wilbur et al. 2010). Altogether, the presented *in*
*silico* data strongly suggest *Gl*ACLC to be structural analogue of CLC.

**SsCHC is distributed in the cytosol and interacts with a putative light chain structural**
**analogue**

We hypothesize that *Gl*ACLC  is a Giardia-specific CLC analogue, which arose *de novo*
in the LGCA, possibly to supplant the loss of a *bona fide* CLC, We wondered whether *de novo*
acquisition of a CLC analogue with divergent sequence but preservation of structural and
potentially also functional features had occurred independently in other Diplomonadida lineages.
To address this question, we selected *Spironucleus salmonicida*the closest genetically tractable
and sequenced relative to *Giardia* (Jerlström-Hultqvist et al. 2012; Xu et al. 2014) in which a *bona*
*fide* CLC cannot be detected, We expressed an epitope-tagged variant of the *S. salmonicida* CHC
ortholog (*Ss*CHC-3xHA) and detect it  distributed in a punctate pattern throughout the
trophozoite cytosol . Using  higher resolution confocal microscopy , *Ss*CHC-3xHA was shown in
cytoplasmic structures reminiscent of giardial CHC focal assemblies (Supplemental Figure 9A
and B and Supplemental Video 7).

To probe for the presence of a CLC analogue in *Spironucleus*, we performed native co-
immunoprecipitation experiments (native  co-IP) using the *Ss*CHC-3xHA-expressing *S.*
*salmonicida* transgenic line. *In silico* analysis of the mass spectrometry  dataset focused on the
most abundant proteins pulled down with a minimum of 10 peptide hits using stringent criteria.
This yielded 171 proteins which were either exclusive to the native co-IP sample derived from the
transgenic reporter line, or ≥3-fold enriched in the transgenic line compared to the non-transgenic
parental strain (Supplemental Table 12). We identified several endocytosis-related proteins, with
*Ss*-dynamin being the most abundant (48 hits and exclusive to *Ss*CHC co-IP reaction), together
with *Ss*-β-adaptin, *Ss*-calmodulin and *Ss*Sec7 (Supplemental Figure 9C and D). Despite its
intranuclear localization, annexin 5 is also found to be a putative interactor of *Ss*CHC (Einarsson
et al. 2019).

Since we postulated that a possible CLC analogue would be among the hypothetical
proteins, we probed those hits using the HHPred algorithm (Zimmermann et al. 2017), focusing
on candidates with predicted secondary structures composed of alpha-helical and coiled-coil
domains and a size < 40 kDa, consistent with all CLC documented thus far. One of these, protein

*Ss*50377_11905, was found to be prominently pulled down and contains several coil-coil domains,

at a predicted weight of 39kDa. The 150 amino acid C-terminus of the protein was modelled in

AlphaFold and superimposed with CLC structures (supplementary figure 9E). TM-align values

within structure similarity (0.5 or above), and RMSD values of *circa* 5-6 Å make a compelling case

for *Ss*50377_11905 to be a *S. salmonicida* CLC analogue , similar to *Gl*CLC. Using *Ss*11905, we

also performed homology searches in the available transcriptome of the related diplomonad

*Trepomonas sp.* (Kolisko et al. 2008b; Xu et al. 2016). We found a candidate ortholog,

TPC1_16039 (forward tblastn e-value of  1E-5 and reverse blastp e-value of 4E-12)

(supplementary table 11). Notably, however, searches using *Ss*50377_11905 into the remaining

fornicata representatives failed to retrieve candidate homologues, whether *Gl*ACLC or other (data

not shown). Protein structure modelling with AlphaFold and super-imposition with *Ss*11905,

*Gl*ACLC, *T. brucei* CLC and *H. sapiens* CLC suggest that  TPC1_16039 is also a structural CLC

analogue, orthologous to *Ss*11905 (supplementary figure 10 ).

**DISCUSSION**

***Giardia lamblia's* endocytic organelle system consists of three classes of small acidifying**

**membrane compartments.**

Subsequent to ingestion and excystation, *Giardia* trophozoites attach to the intestinal

lumen, proliferating and encysting on localised foci throughout the mucosa of the small intestine

(Barash et al. 2017). Nutrients required for this propagation are taken up from the environment

through PV/PEC-mediated endocytosis of fluid phase and membrane bound material (Lanfredi-

Rangel et al. 1998; Adam 2001; Abodeely et al. 2009; Carranza and Lujan 2009; Cotton et al.

2011; Zumthor et al. 2016; Touz et al. 2018). Despite the essential nature of these endocytic

organelles, complete resolution of the ultrastructure of the *Giardia* endocytic pathway remains

unsolved. To address this we performed an ultrastructural investigation of *G. lamblia* endocytic

compartments to obtain a nanometric view of their morphology as defined by their membrane as

well as the lumen accessible to fluid phase markers in labelling experiments (Abodeely et al.

2009; Zumthor et al. 2016).

We began by dissecting an entire *G. lamblia* trophozoite using scanning electron

microscopy and focused our analysis on PVs. These structures were segmented and rendered in

three dimensions. Using this method unambiguously detected at least two distinct classes of PV

morphologies, with some being obviously globular in shape while others presenting a more tubular

nature. After expanding our analysis of PVs/PECs to super resolution light microscopy methods

STED and STORM (Jacquemet et al. 2020) we determined that PVs/PECs are present in three

discernible morphologies: spherical, tubular and polymorphic. Thus, we proposed the renaming

of these organelles into peripheral endocytic compartments (PECs).

Furthermore, in line with previous reports (McCaffery et al. 1994; McCaffery and Gillin

1994; Benchimol 2002; Zumthor et al. 2016), we also detected smaller vesicles (SVs) of around

80 nm in radius which appear to be coated, based on their electron-dense surface, and are not

related to CHC foci at the PV-PM interface. Identifying the nature of SV coats may shed light on

their corresponding cargo. For instance, COPI components such as the small GTPase ARF1 and

β'-COPI were found to be located not only at the parasite's ER but also at the cell periphery  (Marti,

Regös, et al. 2003; Stefanic et al. 2006; Stefanic et al. 2009), in line with SV distribution. If indeed

COPI were found to act as coat for these currently uncharacterized membrane carriers, an

intriguing possibility emerges for SVs as vehicles for the trafficking of variant surface proteins

(VSPs) to the *Giardia* cell surface. VSP secretion is compromised by the presence of brefeldin-A,

implicating ARF-GTPase cycles in VSP trafficking (McCaffery et al. 1994; Lujan et al. 1995; Marti,

Li, et al. 2003; Marti, Regös, et al. 2003). Currently, the exact mechanism for VSP translocation

from the ER to the plasma membrane remains unknown, although it has been postulated that

PVs/PECs may be involved. Furthermore, previous reports exclude the presence of an

intermediate VSP trafficking compartment between the ER and the PM detectable by microscopy

(Marti, Li, et al. 2003; Marti, Regös, et al. 2003). Given the estimated diameter of SVs at ca. 80

nm, these compartments would have easily escaped detection in standard light microscopy

experiments. An alternative hypothesis concerning SVs is that they are peroxisome derivatives.

Recently, peroxisomes have been found in *Entamoeba histolytica,* a microaerophile like Giardia*,*

with diameters between 90-100 nm, in the range of *Giardia* SVs (Verner et al. 2021). Furthermore,

reports on immuno-EM detection of peroxisome-like proteins in *Giardia* highlighted the presence

of small dense vesicles with diameters of *circa.* 100 nm (Acosta-Virgen et al. 2018)  . Taken

together, while we favour the hypothesis of SVs being secretory trafficking vesicles considering

their apparent coating, the possibility of SVs corresponding to cryptic peroxisome-like organelles

cannot be excluded.

Compared to endosome-like vacuoles in *Carpediemonas*-Like Organisms (CLOs) (Yubuki

et al. 2013; Yubuki et al. 2016; Hamann et al. 2017) and large vesicular endosome-like structures

observed in *S. salmonicida* and *S. vortens* and the more distantly related Parabasalia member,

*T. foetus*, specific and complete remodelling of endosomes has occurred in the *Giardia* genus. *T.*

*foetus*, except for the presence of endosome-like vesicles, presents digestive vacuoles and a

stacked Golgi apparatus. Coated vesicles, likely CCVs, are observed near the *T. foetus* Golgi

apparatus and the PM. In our analysis, we could not confirm fluid phase material uptake through

the cytostome present in *Spironucleus sp* (Sterud and Poynton 2002). As noted, dextran

accumulated in spherical vesicles of different dimensions and unknown origin, similar to

endosomes. Figures 10A and B summarize the results of our comparative analysis and highlight

the unique endocytic system in *Giardia* where, unlike related species and other excavates, PEC-

mediated uptake is restricted to the dorsal side of the cell (Ebneter and Hehl 2014; Zumthor et al.

2016) while the ventral side is deputed to attachment to host structures. Interestingly, endosome

and lysosome tubulation has been documented in macrophages (Hipolito et al. 2018; Suresh et

al. 2020) and are linked with different physiological states of the organelles and subsequent

function in the cell – like prompting the cell for phagocytosis. This does permit the hypothesis for

the different kinds of PECs present in *G. lamblia* also representing different stages in organelle

maturation and or active role at a given time. In line with this, we provide evidence for different

stochiometric association of CHC foci with different kinds of PECs. Naturally, only further

biochemical dissection of *Giardia* endocytic pathway will help clarify the matter.

**G. lamblia possesses a highly divergent clathrin heavy chain and a newly acquired clathrin**

**light chain analogue**

We performed an in-depth search for CHC homologs within excavates and other key

eukaryotic groups. We found that CHC is conserved in all of these organisms, underlining the vital

role of CHC in eukaryotic organisms. The *s*equence divergence of the giardial CHC protein is

reflected in an overall decrease in the number of α-helical domains which are essential for the

formation of the triskelion leg, and hence necessary for coat assembly (Kirchhausen et al. 2014).

Thus, the reduction in α-helical domains during *Gl*CHC evolution, may have led to a lower

propensity of *Gl*CHC forming triskelion assemblies and membrane coats. So far, none of the many

attempted methods to detect GlCHC in association with small vesicles have been able to show

anything other than an exclusive focal localization at PVs/PEC membrane interfaces (Zumthor et

al. 2016)). Also, the *Gl*CHC protein does not contain the C-terminal uncoating motif "QLMLT" nor

is this motif present in the CHC homologs of any diplomonad. In fact, this motif appears to be only

present in Metazoa and in the closely related Filastera and Choanoflagellata (King et al. 2008;

Fairclough et al. 2013; Suga et al. 2013) despite the documented ability to form and uncoat *bona*

*fide* CCVs in some protozoa (Link et al. 2021). In Fungi only a partial "L(M)TL"motif was identified.

We could not detect a conserved uncoating motif in CHC sequences of members of the

Archaeplastida, Amoebozoa or SAR supergroups (supplementary figure 7). Taken together, this

data indicates the uncoating QLMLT motif is apparently specific to and likely and invention of the

Holozoa lineage. This observation points to as yet uncharacterised uncoating mechanisms are
present in other lineages. For example, clathrin mediated endocytosis is essential in the parasitic
protist *Trypanosoma brucei* and CCVs have been documented in this organism (Morgan et al.
2001; Allen et al. 2003; Adung'a et al. 2013; Link et al. 2021). Clathrin and other coat proteins
associated with CCVs need to be recycled. While HSC70 is documented in *T. brucei* and likely
involved in clathrin uncoating (Rapoport et al. 2008), no *bona fide* uncoating motif has been
documented (Adung'a et al. 2013; Manna et al. 2017; Link et al. 2021).
590
In contrast to *Gl*CHC, evolution of the previously identified putative *Gl*CLC/*Gl*4259 protein,
presents a different and surprising natural history. This protein was identified as the strongest
interactor of GlCHC (Zumthor et al. 2016) and is present in all sequenced *Giardia* lineages.
*Gl*CLC/*Gl*4259 has no measurable sequence conservation but a high degree of structural
similarity to *bona fide* CLCs, warranting its proposed renaming to *Gl*ACLC. Aside from the *Giardia*
genus, we were unable to identify homologues for *Gl*ACLC in any other eukaryotic taxa , nor could
we find any orthologues of CLC in any available Fornicata genome/transcriptome sequence,
suggesting that the last Fornicata common ancestor (LFCA) lacked a canonical CLC. Taken
together, the available data is currently insufficient to decide between two mutually exclusive
evolutionary scenarios: 1) secondary loss of a canonical CLC in the last fornicate common
ancestor, with acquisition of a structurally and functionally related *GlACLC*, or 2) massive
sequence divergence of the original CLC due to significant changes in function particularly in
Giardia where originally dynamic membrane coating machinery has evolved to become a static
structural element supporting interfaces between plasma membrane and the endocytic system.
The discovery of a strong interactor of CHC in the closely related *S. salmonicida Ss*11905, with
structural similarity to *GlACLC* as well as to bona fide CLCs is consistent with both scenarios.
Notably, this protein neither retrieves *GlACLC*  nor CLCs in BLAST searches, leaving no evidence
of direct homology. By contrast, robust predictions for CLC homologues were made for members
of the Preaxostyla, Discoba and Parabasalia lineages. Nonetheless, other lineages appear also
to have lost a bona fide CLC, like *C. parvum* and *T. thermophila* (Woo et al. 2015; Richardson
and Dacks 2022), but perhaps similar investigations to ours of CHC may identify CLC
analogues/divergent homologues. Taken together, this data suggests that the constraints on the
CHC primary structure are higher than on CLC even after massive changes in clathrin coat
function with demonstrated complete losses in some protists.  Members of the *Giardia* genus as
well as *S. salmonicida* have no identifiable *bona fide* CLC, yet, at least the giardial *Gl*ACLC has
retained its function as a CHC interacting partner.

Our data provide a robust understanding of *Giardia, Hexamitidae* members, and
*Tritrichomonas foetus* endocytic pathway organellar ultrastructure. Contrary to *Spironucleus* or
*Tritrichomonas* and other excavates, *Giardia* underwent a complete remodelling of its endocytic
machinery. Our analysis revealed its organelles to be polymorphic in nature, justifying the
proposed name change to peripheral endocytic compartments. Our analysis of *GCHC* sequences
highlights its divergence which is likely due to a massive reorganization of the endocytic pathway
in these species, whilst origin and evolution of CLC structural and to some extent functional
homologs in *Giardia* (*GlACLC*) and in certain *Hexamitidae* members (*S. salmonicida* and
*Trepomonas* sp. PC1) remains uncertain.
626

**MATERIALS AND METHODS**
***Cell culture and transfection***
*Giardia intestinalis* strain WB (clone C6; ATCC catalog number 50803) trophozoites were grown
using standard methods as described in Morf et. al. (Morf et al. 2010)  Episomally- transfected
parasites were obtained via electroporation of the circular pPacV-Integ-based plasmid prepared
in *E. coli* as described in Zumthor et al. (Zumthor et al. 2016) Transfectants were selected using
Puromycin (final conc. 50 µg ml$^{-1}$; InvivoGen). *Spironucleus vortens* and *Spironucleus*
*salmonicida* were cultured as described before (Paull and Matthews 2001; Xu et al. 2014). *S.*
*salmonicida* was transfected using a modified PAC vector and selected with Puromycin (final
conc. 50 µg ml$^{-1}$; InvivoGen) (Jerlström-Hultqvist et al. 2012). *Tritrichomonas foetus* was
axenically grown also as described (Lealda et al. 1986).
638

**Construction of expression vectors**
*Spironucleus salmonicida* CHC sequence (SS50377_14164) was amplified with the primers
<u>ATATTTAATTAAGGCGGATCTATAGTTTCTTGGAATACTAAAATAGGA</u>      (forward)      and
<u>TATGCGGCCGCCACCAGTTATCAGCGGGTGCC</u> (reverse) containing a MluI and a NotI
rectrictyion site respectively. The genomic sequence amplified contained a 5' UTR region of
179bp which encodes a putative promoter. The genomic fragment was inserted in the previously
described vector pSpiro-PAC-3xHA-C (Jerlström-Hultqvist et al. 2012).
646

**Focused Ion Bean Scanning Electron Microscopy (FIB-SEM) of a full *Giardia* trophozoite**
**and image analysis**
Wild type *Giardia lamblia* trophozoites were subject to High Pressure Freezing, and processed
as established in (Zumthor et al. 2016). Ion milling and Imaging was performed in a Auriga 40

Crossbeam system (Zeiss, Oberkochen, Germany) using the FIBICS Nanopatterning engine

(Fibics Inc., Ottawa, Canada) following the aforementioned established protocol. Pixel size was

set to 5 nm, obtaining isotropic imaging. Alignment of the dataset was performed resorting to the

ImageJ plugin Sift (Schindelin et al. 2012). Image segmentation was done using the semi-

autonomous algorithm ilastik (Sommer et al. 2011). The routine of pixel and object classification

and used. Algorithm training was performed in a small representative region of the dataset which

was then applied to the complete dataset. Imaris (Bitplane AG) was used for three-dimensional

rendering and volume measuring.

**Transmission Electron Microscopy analysis of *Giardia lamblia*, *Spironucleus spp*. and**

***Tritrichomonas foetus* cells and analysis**

*G. lamblia*, *S. vortens, S. salmonicida* and *Tritrichomonas foetus* samples were subject to high

pressure freezing and processed as we previously established (Gaechter et al. 2008; Zumthor et

al. 2016). Samples were imaged in a FEI CM100 Transmission Electron Microscope. Pixel size

was assigned to 0.8 nm. Tiles were obtained automatically after determination of focal point. Tiles

were aligned with TrakEM2 (Cardona et al. 2012).

**Immunofluorescence Assays**

Chemically fixed cells for subcellular recombinant protein localization were prepared as previously

described (Konrad et al. 2010). HA-epitope tagged recombinant proteins were detected using a

rat-derived monoclonal anti-HA antibody (dilution 1:200, Roche) followed by a secondary anti-rat

antibody coupled to AlexaFluor 488 fluorophores (dilution 1:200, Invitrogen). Samples were

embedded in Vectashield (VectorLabs) or Prolong Diamond Mounting medium (Invitrogen)

containing 4',6-diamidino-2-phenylindole (DAPI) for nuclear staining.

***Fluid phase marker uptake***

Dextran uptake assays were performed as described in (Gaechter et al. 2008; Zumthor et al.

2016) using Dextran 10kDa at 2mg/mL (Invitrogen). Coupled fluorophore was chosen based on

image technique chosen. Immunostaining was performed as described above with the exception

of using only 0.02% Triton-X100 (Sigma) in 2% BSA (Sigma) for permeabilization, to prevent

leakage and loss of Dextran signal. Intensities were calculated with a costume developed macro

in Fiji/ImageJ (Schindelin et al. 2012), resorting to WEKA algorithms for segmentation (Arganda-
Carreras et al. 2017).

**Laser Scan Confocal Microscopy (LSCM)**

Imaging was performed in an inverted Confocal Laser Scanning Microscope Leica SP8 using
appropriate parameters. Confocal images were subsequently deconvolved using Huygens
Professional (https://svi.nl/Huygens-Professional) and analysed using Fiji/ImageJ (Schindelin et
al. 2012).

**Stimulated Emission Depletion (STED) Microscopy**

Sample preparation was done as described for LSCM. For imaging, samples were mounted in
ProLong Diamond antifade reagent (Thermo Fisher Scientific). Super resolution microscopy was
performed on a LSCM SP8 gSTED 3X Leica (Leica Microsystems) using appropriate gating
settings. Nuclear labelling was omitted due to possible interference with the STED laser. A pulse
depletion laser of 775 nm at 100% strength was used to deplete signal coming from samples
using the fluorophore Alexa Fluor 594. Signal from samples containing Alexa Fluor 488 were
depleted with the depletion laser line 592 nm at 50% strength. Pinhole was kept at 1 AU. Images
were deconvolved using Huygens Professional (https://svi.nl/Huygens-Professional). After
deconvolution, signal was segmented following a pixel and object classification routine in ilastik.
Thresholding was processed in Fiji/ImageJ (Schindelin et al. 2012) with respective calculation of
organelle area.

**Single Molecule Localization Microscopy (SMLM)**

Cells were fixed into a coverslip using a cytospin (6 min, 600 g). Samples were then embedded
in Vectashield based imaging medium (Olivier et al. 2013). Excess buffer was dried up and
samples were sealed. Single Molecule Imaging was performed on a on a Leica SR-GSD 3D
microscope (Leica Microsystems) as described in (Mateos et al. 2016) with a cylindrical lenses,
in order to image the apical cell region, giving a z-depth of about 800 nm.. A minimum of
100 000 events were recorded. Image reconstruction was performed with the ImageJ plugin
Thunderstorm (Ovesný et al. 2014). Reconstructed images were segmented following a pixel and
object classification routine in ilastik (Sommer et al. 2011; Berg et al. 2019). Thresholding and
volume calculation was performed in Imaris (Bitplane AG).

**Native Co-immunoprecipitation of *S. salmonicida* CHC**

Co-immunoprecipitation assays on control wild type *S. salmonicida* and transgenic *S. salmonicida*
bearing the HA-tagged CHC were processed as previously established (Zumthor et al. 2016) in
non-cross-linking conditions agent.
721

**Protein analysis and sample preparation for mass spectrometry (MS)-based protein**
**identification**
SDS-PAGE analysis was performed on 4%-10% polyacrylamide gels under reducing conditions.
Blotting was done as described in (Konrad et al. 2010) using primary rat-derived anti-HA antibody
(dilution 1:500, Roche) followed by anti-rat (dilution 1:2000; Southern Biotech) antibody coupled
to horseradish peroxidase. Gels for mass spectroscopy (MS) analysis were stained with Instant
blue (Expedeon) and de-stained with ultrapure water. MS-based protein identification was
performed as previously reported (Zumthor et al. 2016).
730

**In silico co-immunoprecipitation dataset analysis**
The co-IP datasets derived from transgenic cells expressing epitope-tagged "baits" as affinity
handles were filtered using dedicated control co-IP datasets generated from non-transgenic wild-
type parasites to identify candidate interaction partners unique to bait-specific datasets. This was
done using Scaffold4 (http://www.proteomesoftware.com/products/scaffold/). Unless otherwise
indicated, bait-derived co-IP data was filtered using high stringency parameters (Exclusive
Spectrum Counts at 95-2-95, 0% FDR) and manually curated to rank putative interaction partners
in a semi-quantitative fashion using ESCs as a proxy for relative abundance. Only proteins with
more than 10 hits were considered. Proteins in both datasets were only considered if present 3-
fold in the transgenic line versus the control. *In silico* analysis of hypothetical proteins was mainly
carried        out        using        BLASTp        for        protein        homology        detection
(http://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE=Proteins)                    and                    HHPred
(http://toolkit.tuebingen.mpg.de/hhpred) for protein homology detection based on Hidden Markov
Model (HMM-HMM) comparisons and a cut-off at e-value < 0.05 was implemented to assign *in*
*silico* annotation to otherwise non-annotated proteins of unknown function (Zimmermann et al.
2017).
Protein        structure        was        modelled        with        the        *ab initio*        modelling        tool        AlphaFold
(https://alphafold.ebi.ac.uk/)        from        Alphabet,        powered        by        Google        DeepMind
(https://deepmind.com/) deep learning neural network algorithms   (Jumper et al. 2021;
Tunyasuvunakool et al. 2021). Modelling was done via Google Colab in a Jupyter notebook
environment(//colab.research.google.com/github/deepmind/alphafold/blob/main/notebooks/).

TM-align calculation was performed online in the server: https://zhanggroup.org/TM-score/.
Pymol (The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC.) was used for
protein structure prediction visualisation, superimposing and RMSD calculation using the *cealign*
command.
756

757     **Data Availability**
758         Access to raw mass spectrometry data is provided through the ProteomeXchange
759     Consortium on the PRIDE platform (Perez-Riverol et al. 2019). Data is freely available using
760     project accession number and project DOI. Project DOI/accession number for datasets derived
761     from bait specific and control co-IP MS analyses are as follows: PXD020201.
762

**Homologue search and Phylogenetic analysis and tree construction**
CHC and CLC sequences were probed among several available genomes and transcriptomes
with special focus within the fornicata members. Query protein sequences for CHC and CLC from
several pan-eukaryotic representatives were obtained and aligned using MUSCLE v.3.8.31
(Edgar 2004) (Supplementary Table 2). Resulting alignments were used to generate Hidden
Markov Models using the hmmbuild option and HHMer searches were made on all available
genomes with an e-value cutoff to 0.01 (Eddy 2011). Hits were considered valid if reciprocal
BLASTp returned a Homo sapiens homologue with a e-value < 0.05. Transcriptome searches
were carried out resorting to tBLASTn searches using the *Homo sapiens* and
*Monocercomonoides exilis* respective sequences for CHC or CLC. Once a hit was found it was
translated into an amino acid sequence and was considered valid if it pulled a *Homo sapiens*
homolog with an e-value < 0.05. All found sequences can be found in supplementary tables 2 to
9. Protein domain searches were performed at the Conservate Domain Database (CDD), through
the Pfam database (Lu et al. 2020; Mistry et al. 2021). The interPro and SMART platforms were
also used for domain classification (Letunic and Bork 2017; Mitchell et al. 2019). Synonymous vs
non- synonymous mutation ratio was calculated with an available online software
(http://services.cbu.uib.no/tools/kaks) following maximum likelihood parameters.
780

**Statistical analysis and further used software**
All data was analysed for statistical significance and plotted using Prism 9 (Graphpad,
https://www.graphpad.com/scientific-software/prism/) software. Images were composed using
Affinity Designer software (https://affinity.serif.com/en-gb/). Video processing was made using Da
Vinci Resolve v17.3.

**ACKNOWLEDGMENTS**

ABH and CF are funded by Swiss National Foundation grant 31003A-166437 and
PR00P3_179813, respectively. Imaging and image analysis were performed with equipment from
the Centre of Microscopy and Image Analysis (ZMB) of the University of Zurich. We thank the
following members of the ZMB for technical and scientific support: Dr. Jana Döhner, Dr. Moritz
Kirchmann, Dr. Dominik Hänni, Dr. José Mateos and Dr. Urs Ziegler. Finally we would like to
thank members of the Hehl lab for insightful discussions.

**CONFLICTS OF INTEREST**

No conflicts of interest

**AUTHOR CONTRIBUTIONS**

RS, ABH and CF designed and curated the study. RS performed all experiments and analysed
all ensuing experimental data with the exception of *Spironucleus salmonicida* culturing and uptake
experiments performed by AA and SS, and SEM experiments performed by JPZ. SVP, RS and
JBD performed molecular phylogeny analyses. RS, ABH and CF wrote and revised the
manuscript. All authors read and approved the final manuscript prior to submission.

**REFERENCES**

Abodeely M, DuBois KN, Hehl A, Stefanic S, Sajid M, DeSouza W, Attias M, Engel JC, Hsieh I,
Fetter RD, et al. 2009. A contiguous compartment functions as endoplasmic reticulum and
endosome/lysosome in Giardia lamblia. *Eukaryot. Cell* [Internet] 8:1665–1676. Available
from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2772394&tool=pmcentrez&rend
ertype=abstract
Acosta-Virgen K, Chávez-Munguía B, Talamás-Lara D, Lagunes-Guillén A, Martínez-Higuera A,
Lazcano A, Martínez-Palomo A, Espinosa-Cantellano M. 2018. Giardia lamblia:
Identification of peroxisomal-like proteins. *Exp. Parasitol.* 191:36–43.
Adam RD. 2001. Biology of Giardia lamblia. *Clin. Microbiol. Rev.* 14:447–469.
Adung'a VO, Gadelha C, Field MC. 2013. Proteomic Analysis of Clathrin Interactions in
Trypanosomes Reveals Dynamic Evolution of Endocytosis. *Traffic* 14:440–457.
Allain T, Amat CB, Motta JP, Manko A, Buret AG. 2017. Interactions of Giardia sp. with the

intestinal barrier: Epithelium, mucus, and microbiota. *Tissue Barriers* 5:1–16.

Allen CL, Goulding D, Field MC. 2003. Clathrin-mediated endocytosis is essential in Trypanosoma brucei. *EMBO J.* 22:4991–5002.

Arganda-Carreras I, Kaynig V, Rueden C, Eliceiri KW, Schindelin J, Cardona A, Seung HS. 2017. Trainable Weka Segmentation: A machine learning tool for microscopy pixel classification. *Bioinformatics* 33:2424–2426.

Ástvaldsson Á, Hultenby K, Svärd SG, Jerlström-Hultqvist J. 2019. Proximity Staining Using Enzymatic Protein Tagging in Diplomonads. *mSphere* [Internet] 4:1–15. Available from: http://msphere.asm.org/lookup/doi/10.1128/mSphereDirect.00153-19

Aurrecoechea C, Brestelli J, Brunk BP, Carlton JM, Dommer J, Fischer S, Gajria B, Gao X, Gingle A, Grant G, et al. 2009. GiardiaDB and TrichDB: Integrated genomic resources for the eukaryotic protist pathogens Giardia lamblia and Trichomonas vaginalis. *Nucleic Acids Res.* 37:526–530.

Barash NR, Nosala C, Pham JK, Mcinally SG, Gourguechon S, Dawson SC. 2017. Giardia Colonizes and Encysts in High- Density Foci in the Murine Small Intestine. *mSphere* 2:1–20.

Benchimol M. 2002. A new set of vesicles in Giardia lamblia. *Exp. Parasitol.* [Internet] 102:30–37. Available from: http://linkinghub.elsevier.com/retrieve/pii/S001448940200142X

Benchimol M. 2005. The nuclei of Giardia lamblia - new ultrastructural observations. *Arch. Microbiol.* [Internet] 183:160–168. Available from: http://www.ncbi.nlm.nih.gov/pubmed/15703885

Berg S, Kutra D, Kroeger T, Straehle CN, Kausler BX, Haubold C, Schiegg M, Ales J, Beier T, Rudy M, et al. 2019. Ilastik: Interactive Machine Learning for (Bio)Image Analysis. *Nat. Methods* [Internet] 16:1226–1232. Available from: http://dx.doi.org/10.1038/s41592-019-0582-9

Brown JR, Schwartz CL, Heumann JM, Dawson SC, Hoenger A. 2016. A detailed look at the cytoskeletal architecture of the Giardia lamblia ventral disc. *J. Struct. Biol.* [Internet] 194:38–48. Available from: http://linkinghub.elsevier.com/retrieve/pii/S1047847716300107

Burki F, Roger AJ, Brown MW, Simpson AGB. 2020. The New Tree of Eukaryotes. *Trends Ecol. Evol.* [Internet] 35:43–55. Available from: https://doi.org/10.1016/j.tree.2019.08.008

Caccìo SM, Ryan U. 2008. Molecular epidemiology of giardiasis. *Mol. Biochem. Parasitol.* 160:75–80.

Cardona A, Saalfeld S, Schindelin J, Arganda-Carreras I, Preibisch S, Longair M, Tomancak P, Hartenstein V, Douglas RJ. 2012. TrakEM2 software for neural circuit reconstruction. *PLoS*

*One* 7.

Carlton JM, Hirt RP, Silva JC, Delcher AL, Schatz M, Zhao Q, Wortman JR, Bidwell SL, Alsmark

UCM, Besteiro S, et al. 2007. Draft Genome Sequence of the Sexually Transmitted

Pathogen Trichomonas vaginalis. *Science (80-. ).* 315:207–213.

Carranza PG, Lujan HD. 2009. New insights regarding the biology of Giardia lamblia. *Microbes*

*Infect.* 12:71–80.

Cernikova L, Faso C, Hehl AB. 2020. Phosphoinositide-binding proteins mark, shape and

functionally modulate highly-diverged endocytic compartments in the parasitic protist

Giardia lamblia. Available from: http://dx.doi.org/10.1371/journal.ppat.1008317

Cheon S, Zhang J, Park C. 2020. Is phylotranscriptomics as reliable as phylogenomics? *Mol.*

*Biol. Evol.* 37:3672–3683.

Combs CA, Shroff H. 2017. Fluorescence microscopy: A concise guide to current imaging

methods. *Curr. Protoc. Neurosci.* 2017:2.1.1-2.1.25.

Cotton JA, Beatty JK, Buret AG. 2011. Host parasite interactions and pathophysiology in Giardia

infections. *Int. J. Parasitol.* 41:925–933.

Dacks JB, Field MC. 2018. Evolutionary origins and specialisation of membrane transport. *Curr.*

*Opin. Cell Biol.* [Internet] 53:70–76. Available from:

https://doi.org/10.1016/j.ceb.2018.06.001

Dawson SC. 2010. An insider's guide to the microtubule cytoskeleton of Giardia. *Cell. Microbiol.*

12:588–598.

Day KJ, Casler JC, Glick BS. 2018. Budding Yeast Has a Minimal Endomembrane System.

*Dev. Cell* [Internet] 44:56-72.e4. Available from:

https://doi.org/10.1016/j.devcel.2017.12.014

Dempsey GT, Vaughan JC, Chen KH, Bates M, Zhuang X. 2011. Evaluation of fluorophores for

optimal performance in localization-based super-resolution imaging. *Nat. Methods*

[Internet] 8:1027–1036. Available from:

http://www.nature.com/doifinder/10.1038/nmeth.1768

Ebneter JA, Hehl AB. 2014. The single epsin homolog in Giardia lamblia localizes to the ventral

disk of trophozoites and is not associated with clathrin membrane coats. *Mol. Biochem.*

*Parasitol.* [Internet] 197:24–27. Available from:

http://linkinghub.elsevier.com/retrieve/pii/S0166685114001388

Eddy SR. 2011. Accelerated profile HMM searches. *PLoS Comput. Biol.* 7.

Edgar RC. 2004. MUSCLE: Multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Res.*

Einarsson E, Ástvaldsson Á, Hultenby K, Andersson JO. 2019. Comparative Cell Biology and
Evolution of Annexins in Diplomonads. *mSphere* 1:1–16.

Fairclough SR, Chen Z, Kramer E, Zeng Q, Young S, Robertson HM, Begovic E, Richter DJ,
Russ C, Westbrook MJ, et al. 2013. Premetazoan genome evolution and the regulation of
cell differentiation in the choanoflagellate Salpingoeca rosetta. *Genome Biol.* 14:1–15.

Faso C, Hehl AB. 2011. Membrane trafficking and organelle biogenesis in Giardia lamblia: Use
it or lose it. *Int. J. Parasitol.* 41:471–480.

Fekete E, Allain T, Siddiq A, Sosnowski O, Buret AG. 2021. Giardia spp. and the Gut
Microbiota: Dangerous Liaisons. *Front. Microbiol.* 11.

Fotin A, Cheng Y, Grigorieff N, Walz T, Harrison SC, Kirchhausen T. 2004. Structure of an
auxilin-bound clathrin coat and its implications for the mechanism of uncoating. *Nature*
[Internet] 429:2004. Available from: http://mcb.asm.org/lookup/doi/10.1128/MCB.00448-15

Frontera LS, Moyano S, Quassollo G, Lanfredi-Rangel A, Rópolo AS, Touz MC. 2018.
Lactoferrin and lactoferricin endocytosis halt Giardia cell growth and prevent infective cyst
production. *Sci. Rep.* [Internet] 8:18020. Available from:
http://www.nature.com/articles/s41598-018-36563-1

Füssy Z, Vinopalová M, Treitli SC, Pánek T, Smejkalová P, Čepička I, Doležal P, Hampl V.
2021. Retortamonads from vertebrate hosts share features of anaerobic metabolism and
pre-adaptations to parasitism with diplomonads. *Parasitol. Int.* 82.

Gaechter V, Schraner E, Wild P, Hehl AB. 2008. The single dynamin family protein in the
primitive protozoan giardia lamblia is essential for stage conversion and endocytic
transport. *Traffic* 9:57–71.

Hamann E, Tegetmeyer HE, Riedel Di, Littmann S, Ahmerkamp S, Chen J, Hach PF, Strous M.
2017. Syntrophic linkage between predatory Carpediemonas and specific prokaryotic
populations. *ISME J.* [Internet] 11:1205–1217. Available from:
http://dx.doi.org/10.1038/ismej.2016.197

Hampl V, Hug L, Leigh JW, Dacks JB, Lang BF, Simpson AGB, Roger AJ. 2009. Phylogenomic
analyses support the monophyly of Excavata and resolve relationships among eukaryotic
'"supergroups."' *PNAS* 106:3859–3864.

Heimerl T, Flechsler J, Pickl C, Heinz V, Salecker B, Zweck J, Wanner G, Geimer S, Samson
RY, Bell SD, et al. 2017. A complex endomembrane system in the archaeon Ignicoccus
hospitalis tapped by Nanoarchaeum equitans. *Front. Microbiol.* 8:1–13.

Hipolito VEB, Ospina-Escobar E, Botelho RJ. 2018. Lysosome remodelling and adaptation
during phagocyte activation. *Cell. Microbiol.* 20:1–8.

Huang B, Wang W, Bates M, Zhuang X. 2008. Three-Dimensional Super-Resolution
Reconstruction Microscopy. *Science (80-. ).* [Internet] 319:810–813. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/18174397

Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, Butterfield CN,
Hernsdorf AW, Amano Y, Kotaro I, et al. 2016. A new view of the tree and life's diversity.
*Nat. Microbiol.* 48.

Huotari J, Helenius A. 2011. Endosome maturation. *EMBO J.* [Internet] 30:3481–3500.
Available from: http://dx.doi.org/10.1038/emboj.2011.286

Jackson AP, Otto TD, Aslett M, Armstrong SD, Bringaud F, Schlacht A, Hartley C, Sanders M,
Wastling JM, Dacks JB, et al. 2016. Kinetoplastid Phylogenomics Reveals the Evolutionary
Innovations Associated with the Origins of Parasitism. *Curr. Biol.* [Internet] 26:161–172.
Available from: http://dx.doi.org/10.1016/j.cub.2015.11.055

Jacquemet G, Carisey AF, Hamidi H, Henriques R, Leterrier C. 2020. The cell biologist's guide
to super-resolution microscopy. *J. Cell Sci.* 133.

Jaiswal JK, Rivera VM, Simon SM. 2009. Exocytosis of Post-Golgi Vesicles Is Regulated by
Components of the Endocytic Machinery. *Cell* [Internet] 137:1308–1319. Available from:
http://dx.doi.org/10.1016/j.cell.2009.04.064

Jerlström-Hultqvist J, Einarsson E, Svärd SG. 2012. Stable transfection of the diplomonad
parasite Spironucleus salmonicida. *Eukaryot. Cell* 11:1353–1361.

Jones SA, Shim S-H, He J, Zhuang X. 2011. Fast, three-dimensional super-resolution imaging
of live cells. *Nat. Methods* [Internet] 8:499–505. Available from:
http://www.nature.com/doifinder/10.1038/nmeth.1605

Jørgensen A, Sterud E. 2007. Phylogeny of Spironucleus (Eopharyngia: Diplomonadida:
Hexamitinae). *Protist* 158:247–254.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates
R, Žídek A, Potapenko A, et al. 2021. Highly accurate protein structure prediction with
AlphaFold. *Nature* [Internet] 596:583–589. Available from:
http://dx.doi.org/10.1038/s41586-021-03819-2

Kaksonen M, Roux AAA. 2018. Mechanisms of clathrin-mediated endocytosis. *Nat. Rev. Mol.*
*Cell Biol.* [Internet] 19:313–326. Available from: http://dx.doi.org/10.1038/nrm.2017.132

Kaksonen M, Toret CP, Drubin DG. 2005. A modular design for the clathrin- and actin-mediated
endocytosis machinery. *Cell* 123:305–320.

Kan A. 2017. Machine learning applications in cell image analysis. *Immunol. Cell Biol.* 95:525–
530.

Kao HP, Kao HP, Verkman a S, Verkman a S. 1994. Tracking of Single Fluorescent Particles
in Three Dimensions: Use of Cylindrical Optics to Encode Particle Position. *Biophys. J.*
67:1291–1300.

Karnkowska A, Treitli SC, Brzoň O, Novák L, Vacek V, Soukal P, Barlow LD, Herman EK,
Pipaliya S V, Pánek T, et al. 2019. The Oxymonad Genome Displays Canonical Eukaryotic
Complexity in the Absence of a Mitochondrion. *Mol. Biol. Evol.*

Karnkowska A, Vacek V, Zubáčová Z, Treitli SC, Petrželková R, Eme L, Novák L, Žárský V,
Barlow LD, Herman EK, et al. 2016. A Eukaryote without a Mitochondrial Organelle. *Curr.*
*Biol.* [Internet]:1–11. Available from:
http://www.sciencedirect.com/science/article/pii/S0960982216302639

King N, Westbrook MJ, Young SL, Kuo A, Abedin M, Chapman J, Fairclough S, Hellsten U,
Isogai Y, Letunic I, et al. 2008. The genome of the choanoflagellate Monosiga brevicollis
and the origin of metazoans. *Nature* 451:783–788.

Kirchhausen T, Owen D, Harrison SC. 2014. Molecular Structure, Function, and Dynamics of
Clathrin-Mediated Membrane Traffic. *Perspect. Biol.* 6.

Kizilyaprak C, Daraspe J, Humbel BM. 2014. Focused ion beam scanning electron microscopy
in biology. *J. Microsc.* 254:109–114.

Klar TA, Jakobs S, Dyba M, Egner A, Hell SW. 2000. Fluorescence microscopy with diffraction
resolution barrier broken by stimulated emission. *Proc. Natl. Acad. Sci. U. S. A.* 97:8206–
8210.

Kolisko M, Cepicka I, Hampl V, Leigh J, Roger AJ, Kulda J, Simpson AGB, Flegr J. 2008a.
Molecular phylogeny of diplomonads and enteromonads based on SSU rRNA, alpha-
tubulin and HSP90 genes: Implications for the evolutionary history of the double
karyomastigont of diplomonads. *BMC Evol. Biol.* 8:1–14.

Kolisko M, Cepicka I, Hampl V, Leigh J, Roger AJ, Kulda J, Simpson AGB, Flegr J. 2008b.
Molecular phylogeny of diplomonads and enteromonads based on SSU rRNA, alpha-
tubulin and HSP90 genes: Implications for the evolutionary history of the double
karyomastigont of diplomonads. *BMC Evol. Biol.* 8:1–14.

Konrad C, Spycher C, Hehl AB. 2010. Selective condensation drives partitioning and sequential
secretion of cyst wall proteins in differentiating Giardia lamblia. *PLoS Pathog.* 6:e1000835.

Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, Panchalingam S, Wu Y, Sow SO,
Sur D, Breiman RF, et al. 2013. Burden and aetiology of diarrhoeal disease in infants and
young children in developing countries (the Global Enteric Multicenter Study, GEMS): A
prospective, case-control study. *Lancet* 382:209–222.

Kufareva I, Abagyan R. 2012. Methods of protein structure comparison. *Methods Mol. Biol.*
857:231–257.

Lanata CF, Fischer-Walker CL, Olascoaga AC, Torres CX, Aryee MJ, Black RE, UNICEF  for
the CHERG of the WHO and. 2013. Global Causes of Diarrheal Disease Mortality in
Children <5 Years of Age: A Systematic Review.Sestak K, editor. *PLoS One* [Internet]
8:e72788. Available from: https://dx.plos.org/10.1371/journal.pone.0072788

Lanfredi-Rangel  a, Attias M, de Carvalho TM, Kattenbach WM, De Souza W. 1998. The
peripheral vesicles of trophozoites of the primitive protozoan Giardia lamblia may
correspond to early and late endosomes and to lysosomes. *J. Struct. Biol.* 123:225–235.

Lax G, Eglit Y, Eme L, Bertrand EM, Roger AJ, Simpson AGB. 2018. Hemimastigophora is a
novel supra-kingdom-level lineage of eukaryotes. *Nature* [Internet] 564:410–414. Available
from: http://www.nature.com/articles/s41586-018-0708-8

Lealda N, Silva C, Elias CA. 1986. Tritrichomonas foetus : Ultrastructure of Endocytosis and
Cytochemistry. *Exp. Parasitol.* 62:405–415.

Leger MM, Kolisko M, Kamikawa R, Stairs CW, Kume K, Čepička I, Silberman JD, Andersson
JO, Xu F, Yabuki A, et al. 2017. Organelles that illuminate the origins of Trichomonas
hydrogenosomes and Giardia mitosomes. *Nat. Ecol. Evol.* 1.

Letunic I, Bork P. 2017. 20 years of the SMART protein domain annotation resource. *Nucleic*
*Acids Res.* [Internet] 46:493–496. Available from:
http://academic.oup.com/nar/article/doi/10.1093/nar/gkx922/4429069/20-years-of-the-
SMART-protein-domain-annotation

Link F, Borges AR, Jones NG, Engstler M. 2021. To the Surface and Back: Exo- and Endocytic
Pathways in Trypanosoma brucei. *Front. Cell Dev. Biol.* 9:1–15.

Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, Gwadz M, Hurwitz DI,
Marchler GH, Song JS, et al. 2020. CDD/SPARCLE: The conserved domain database in
2020. *Nucleic Acids Res.* 48:D265–D268.

Lujan HD, Marotta A, Mowatt MR, Sciaky N, Lippincott-Schwartz J, Nash TE. 1995.
Developmental induction of Golgi structure and function in the primitive eukaryote Giardia
lamblia. *J. Biol. Chem.* 270:4612–4618.

Manna PT, Obado SO, Boehm C, Gadelha C, Sali A, Chait BT, Rout MP, Field MC. 2017.
Lineage-specific proteins essential for endocytosis in trypanosomes. *J. Cell Sci.* [Internet]
130:1379–1392. Available from: http://jcs.biologists.org/lookup/doi/10.1242/jcs.191478

Marti M, Li Y, Schraner EM, Wild P, Kohler P, Hehl AB. 2003. The Secretory Apparatus of an
Ancient Eukaryote: Protein Sorting to Separate Export Pathways Occurs Before Formation

of Transient Golgi-like Compartments. *Mol. Biol. Cell* 14:1433–1447.

Marti M, Regös A, Li Y, Schraner EM, Wild P, Müller N, Knopf LG, Hehl AB. 2003. An

Ancestaral Secretory Apparatus in the Protozoan Parasite Giardia intestinalis. *J. Biol.*

*Chem.* 278:24837–24848.

Mateos JM, Guhl B, Doehner J, Barmettler G, Kaech A, Ziegler U. 2016. Topographic contrast

of ultrathin cryo-sections for correlative super-resolution light and electron microscopy. *Sci.*

*Rep.* [Internet] 6:34062. Available from: http://www.nature.com/articles/srep34062

McCaffery JM, Faubert GM, Gillin FD. 1994. Traffic of a Trophozoite Variant Surface Protein

and a Major Cyst Wall Epitope during Growth, Encystation, and Antigenic Switching. *Exp.*

*Parasitol.* 79:236–249.

McCaffery JM, Gillin FD. 1994. Giardia lamblia: Ultrastructural basis of protein transport during

growth and encystation. *Exp. Parasitol.* 79:220–235.

Midlej V, Pereira-Neves A, Kist LW, Bogo MR, Benchimol M. 2011. Ultrastructural features of

Tritrichomonas mobilensis and comparison with Tritrichomonas foetus. *Vet. Parasitol.*

[Internet] 182:171–180. Available from: http://dx.doi.org/10.1016/j.vetpar.2011.05.015

Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto SCE,

Paladin L, Raj S, Richardson LJ, et al. 2021. Pfam: The protein families database in 2021.

*Nucleic Acids Res.* 49:D412–D419.

Mitchell AL, Attwood TK, Babbitt PC, Blum M, Bork P, Bridge A, Brown SD, Chang HY, El-

Gebali S, Fraser MI, et al. 2019. InterPro in 2019: Improving coverage, classification and

access to protein sequence annotations. *Nucleic Acids Res.* 47:D351–D360.

Morf L, Spycher C, Rehrauer H, Fournier CA, Morrison HG, Hehl AB. 2010. The Transcriptional

Response to Encystation Stimuli in Giardia lamblia Is Restricted to a Small Set of Genes.

*Eukaryot. Cell* 9:1566–1576.

Morgan GW, Allen CL, Jeffries TR, Hollinshead M, Field MC. 2001. Developmental and

morphological regulation of clathrin-mediated endocytosis in Trypanosoma brucei. *J. Cell*

*Sci.* 114:2605–2615.

Naslavsky N, Caplan S. 2018. The enigmatic endosome - Sorting the ins and outs of endocytic

trafficking. *J. Cell Sci.* 131.

Olivier N, Keller D, Rajan VS, Gönczy P, Manley S. 2013. Simple buffers for 3D STORM

microscopy. *Biomed. Opt. Express* [Internet] 4:885. Available from:

https://www.osapublishing.org/boe/abstract.cfm?uri=boe-4-6-885

Ovesný M, Křížek P, Borkovec J, Švindrych Z, Hagen GM. 2014. ThunderSTORM: A

comprehensive ImageJ plug-in for PALM and STORM data analysis and super-resolution

imaging. *Bioinformatics* 30:2389–2390.

Paull GC, Matthews RA. 2001. Spironucleus vortens, a possible cause of hole-in-the-head
disease in cichlids. *Dis. Aquat. Organ.* 45:197–202.

Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ, Inuganti A,
Griss J, Mayer G, Eisenacher M, et al. 2019. The PRIDE database and related tools and
resources in 2019: Improving support for quantification data. *Nucleic Acids Res.* 47:D442–
D450.

Pipaliya S V., Santos R, Salas-Leiva D, Balmer EA, Wirdnam CD, Roger AJ, Hehl AB, Faso C,
Dacks JB. 2021. Unexpected organellar locations of ESCRT machinery in Giardia
intestinalis and complex evolutionary dynamics spanning the transition to parasitism in the
lineage Fornicata. *BMC Biol.* 19:1–23.

Poteryaev D, Datta S, Ackema K, Zerial M, Spang A. 2010. Identification of the switch in early-
to-late endosome transition. *Cell* 141:497–508.

Poulin R, Randhawa HS. 2015. Evolution of parasitism along convergent lines: From ecology to
genomics. *Parasitology* 142:S6–S15.

Radulescu AE, Siddhanta A, Shields D. 2007. A Role for Clathrin in Reassembly of the Golgi
Apparatus. *Mol. Biol. Cell* 18:94–105.

Rapoport I, Boll W, Yu A, Bocking T, Kirchhausen T. 2008. A Motif in the Clathrin Heavy Chain
Required for the Hsc70/Auxilin Uncoating Reaction. *Mol. Biol. Cell* 19:3250–3263.

Richardson E, Dacks JB. 2022. Distribution of Membrane Trafficking System Components
Across Ciliate Diversity Highlights Heterogenous Organelle-Associated Machinery. *Traffic*.

Rivero MR, Miras SL, Quiroga R, Rópolo AS, Touz MC. 2011. Giardia lamblia low-density
lipoprotein receptor-related protein is involved in selective lipoprotein endocytosis and
parasite replication. *Mol. Microbiol.* 79:1204–1219.

Robinson MS. 2015. Forty Years of Clathrin-coated Vesicles. *Traffic* 16:1210–1238.

Rosa IDA, Caruso MB, Rodrigues SP, Geraldo RB, Kist LW, Bogo MR, Gonzaga L,
Vasconcelos ATR DE, Morgado-Díaz JA, Zingali RB, et al. 2014. New insights on the Golgi
complex of Tritrichomonas foetus. *Parasitology* 141:241–253.

Salas-Leiva DE, Tromer EC, Curtis BA, Jerlström-Hultqvist J, Kolisko M, Yi Z, Salas-Leiva JS,
Gallot-Lavallée L, Williams SK, Kops GJPL, et al. 2021. Genomic analysis finds no
evidence of canonical eukaryotic DNA processing complexes in a free-living protist. *Nat.*
*Commun.* 12:1–13.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, et al. 2012. Fiji: an open-source platform for biological-

1092   image analysis. *Nat. Methods* [Internet] 9:676–682. Available from:

1093   http://www.nature.com/doifinder/10.1038/nmeth.2019

1094 Schlacht A, Herman EK, Klute MJ, Field MC, Dacks JB. 2014. Missing pieces of an ancient

1095   puzzle: Evolution of the eukaryotic membrane-trafficking system. *Cold Spring Harb.*

1096   *Perspect. Biol.* 6.

1097 Soltys BJ, Falah M, Gupta RS. 1996. Identification of endoplasmic reticulum in the primitive

1098   eukaryote Giardia lamblia using cryoelectron microscopy and antibody to BiP. *J. Cell Sci.*

1099   109:1909–1917.

1100 Sommer C, Gerlich DW. 2013. Machine learning in cell biology – teaching computers to

1101   recognize phenotypes. *J. Cell Sci. 126,* 126:5529–5539.

1102 Sommer C, Straehle C, Kothe U, Hamprecht FA. 2011. Ilastik: Interactive learning and

1103   segmentation toolkit. *Proc. - Int. Symp. Biomed. Imaging*:230–233.

1104 Stefanic S, Morf L, Kulangara C, Regös A, Sonda S, Schraner E, Spycher C, Wild P, Hehl AB.

1105   2009. Neogenesis and maturation of transient Golgi-like cisternae in a simple eukaryote. *J.*

1106   *Cell Sci.* 122:2846–2856.

1107 Stefanic S, Palm D, Svärd SG, Hehl AB. 2006. Organelle Proteomics Reveals Cargo Maturation

1108   Mechanisms Associated with Golgi-like Encystation Vesicles in the Early-diverged

1109   Protozoan Giardia lamblia *. *J. Biol. Chem.* 281:7595–7604.

1110 Sterud E, Poynton S. 2002. Spironucleus vortens (Diplomonadida) in the Ide, Leuciscus idus

1111   (L.) (Cyprinidae): a Warm Water Hexamitid Flagellate Found in Northern Europe. *J.*

1112   *Eukaryot. Microbiol.* 49.

1113 Suga H, Chen Z, De Mendoza A, Sebé-Pedrós A, Brown MW, Kramer E, Carr M, Kerner P,

1114   Vervoort M, Sánchez-Pons N, et al. 2013. The Capsaspora genome reveals a complex

1115   unicellular prehistory of animals. *Nat. Commun.* 4:1–9.

1116 Suresh B, Saminathan A, Chakraborty K, Cui C, Krishnan Y. 2020. Tubular lysosomes harbor

1117   active ion gradients and poise macrophages for phagocytosis. *PNAS* [Internet]

1118   18:2020.12.05.413229. Available from: https://doi.org/10.1101/2020.12.05.413229

1119 Tanifuji G, Takabayashi S, Kume K, Takagi M, Nakayama T, Kamikawa R, Inagaki Y,

1120   Hashimoto T. 2018. The draft genome of Kipferlia bialata reveals reductive genome

1121   evolution in fornicate parasites. *PLoS One*.

1122 Titze B, Genoud C. 2016. Volume scanning electron microscopy for imaging biological

1123   ultrastructure. *Biol. Cell* 108:307–323.

1124 Touz M, Feliziani C, Rópolo A. 2018. Membrane-Associated Proteins in Giardia lamblia. *Genes*

1125   *(Basel).* [Internet] 9:404. Available from: http://www.mdpi.com/2073-4425/9/8/404

Tovar J, León-Avila G, Sánchez LB, Sutak R, Tachezy J, Van Der Giezen M, Hernández M,
Müller M, Lucocq JM. 2003. Mitochondrial remnant organelles of Giardia function in iron-
sulphur protein maturation. *Nature* 426:172–176.

Traub LM. 2011. Regarding the amazing choreography of clathrin coats. *PLoS Biol.* 9:3–7.

Tůmová P, Nohýnková E, Klingl A, Wanner G. 2020. A rapid workflow for the characterization of
small numbers of unicellular eukaryotes by using correlative light and electron microscopy.
*J. Microbiol. Methods* 172.

Tunyasuvunakool K, Adler J, Wu Z, Green T, Zielinski M, Žídek A, Bridgland A, Cowie A, Meyer
C, Laydon A, et al. 2021. Highly accurate protein structure prediction for the human
proteome. *Nature* [Internet] 596:590–596. Available from: http://dx.doi.org/10.1038/s41586-
021-03828-1

Verner Z, Žárský V, Le T, Narayanasamy RK, Rada P, Rozbeský D, Makki A, Belišová D, Hrdý
I, Vancová M, et al. 2021. Anaerobic peroxisomes in Entamoeba histolytica metabolize
myo-inositol.

Wei D, Jacobs S, Modla S, Zhang S, Young CL, Cirino R, Caplan J, Czymmek K. 2012. High-
resolution three-dimensional reconstruction of a whole yeast cell using focused-ion beam
scanning electron microscopy. *Biotechniques* 53:41–48.

Wilbur JD, Hwang PK, Ybe JA, Lane M, Sellers BD, Jacobson MP, Fletterick RJ, Brodsky FM.
2010. Conformation switching of clathrin light chain regulates clathrin lattice assembly.
*Dev. Cell* [Internet] 18:841–848. Available from:
http://dx.doi.org/10.1016/j.devcel.2010.04.007

Willig KI, Rizzoli SO, Westphal V, Jahn R, Hell SW. 2006. STED microscopy reveals that
synaptotagmin remains clustered after synaptic vesicle exocytosis. *Nature* 440:935–939.

Woo YH, Ansari H, Otto TD, Linger CMK, Olisko MK, Michálek J, Saxena A, Shanmugam D,
Tayyrov A, Veluchamy A, et al. 2015. Chromerid genomes reveal the evolutionary path
from photosynthetic algae to obligate intracellular parasites. *Elife* 4:1–41.

Xu F, Jerlström-Hultqvist J, Einarsson E, Ástvaldsson Á, Svärd SG, Andersson JO. 2014. The
Genome of Spironucleus salmonicida Highlights a Fish Pathogen Adapted to Fluctuating
Environments. *PLoS Genet.* 10.

Xu F, Jerlström-Hultqvist J, Kolisko M, Simpson AGB, Roger AJ, Svärd SG, Andersson JO.
2016. On the reversibility of parasitism: adaptation to a free-living lifestyle via gene
acquisitions in the diplomonad Trepomonas sp. PC1. *BMC Biol.* 14:1–15.

Yubuki N, Huang SSCC, Leander BS. 2016. Comparative Ultrastructure of Fornicate Excavates,
Including a Novel Free-living Relative of Diplomonads: Aduncisulcus paluster gen. et sp.

nov. *Protist* [Internet] 167:584–596. Available from:

http://dx.doi.org/10.1016/j.protis.2016.10.001

Yubuki N, Simpson AGB, Leander BS. 2013. Comprehensive Ultrastructure of Kipferlia bialata

Provides Evidence for Character Evolution within the Fornicata (Excavata). *Protist*

[Internet] 164:423–439. Available from: http://dx.doi.org/10.1016/j.protis.2013.02.002

Zhang Y, Skolnick J. 2005. TM-align: A protein structure alignment algorithm based on the TM-

score. *Nucleic Acids Res.* 33:2302–2309.

Zimmermann L, Stephens A, Nam SZ, Rau D, Kübler J, Lozajic M, Gabler F, Söding J, Lupas

AN, Alva V. 2017. A Completely Reimplemented MPI Bioinformatics Toolkit with a New

HHpred Server at its Core. *J. Mol. Biol.* [Internet] 430:1–7. Available from:

https://doi.org/10.1016/j.jmb.2017.12.007

Zumthor JP, Cernikova L, Rout S, Kaech A, Faso C, Hehl AB. 2016. Static Clathrin Assemblies

at the Peripheral Vacuole Plasma Membrane Interface of the Parasitic Protozoan Giardia

lamblia. *PLoS Pathog.* 12:1–33.

1174

**FIGURE LEGENDS**

**Figure 1 – Complete scanning of a *Giardia* trophozoite by focused ion beam scanning**

**electron microscopy (FIB-SEM).** (A and D) A complete *G. lamblia* trophozoite was scanned with

an isotropic resolution of 5 nm. The Nucleus (N), the Endoplasmic Reticulum (ER), cytoskeletal

features such as the Ventral disk (VD), Funis (F) and axonemes (Ax) and mitosomes (m), are

observed. Peripheral vesicles (PV) are marked by arrowheads. Smaller, electron dense vesicles

are also documented (asterisks) – small vesicles (SVs). (B) Region of interest of (A) highlighting

PVs of different morphology with arrowheads. (C) Highlight of mitosomes proximal to ER

membrane. (E) Region of interest of (D) highlighting SVs (asterisks). (F) Full reconstruction of

PVs with ilastik and rendering in Imaris reveals the presence of at least two PV morphologies:

spherical (green) and tubular (violet). Box-plot: 403 spherical and 64 tubular PVs were segmented

out. Spherical PVs average a volume of $0.0009243\pm0.0003322$ $\mu m^3$ with a 95% confidence

interval between [0.0009022; 0.0009022] $\mu m3$. Tubular PVs average a volume of

$0.001067\pm0.0003322\mu m^3$ with a 95% confidence interval between [0.0009843; 0.001150] $\mu m^3$.

(G) Reconstruction of 269 SVs revealed their volumes average $0.0002525\pm9.280x10^{-5}$ $\mu m^3$ in a

95% confidence interval between [0.0002414; 0.0002637] $\mu m^3$ (left box-plot). This equals to an

average diameter of $77.23\pm9.666$ nm in a 95% confidence interval between [76.07; 78.40] nm.

The diameter of spherical PVs averages $120.1\pm9.507$ nm in a 95% confidence interval between

[119.2; 121] nm (right box-plot). The difference in diameter between SVs and spherical PVs is

statistically significant (****; p-value < 0.0001, t-student significance test). Scale bars: (A and D)

2 $\mu$m, (B, C and E) 500 nm. ROI: region of interest.

1196

**Figure 2 – Super Resolution imaging of *Giardia lamblia* peripheral vesicles with Stimulated**

**Emission Depletion (STED).** (A) *Giardia* trophozoites loaded with 10kDa Dextran-AlexaFluor

594 were imaged using Confocal and STED microscopy. Dorsal (upper row) and ventral (lower

row) regions are represented. In contrast to confocal imaging, STED microscopy allows to

separate individual organelles and to visualise different endocytic compartment morphologies.

ROI: region of interest. (B) Organelles segmentation with ilastik distinguishes three Dextran-

labelled PV categories. (C) PV areas were calculated post-segmentation on maximum projections

of the dorsal regions of ≥10 cells, using ilastik. Spherical PVs (green, N =1684) have an average

area of 0.0205±0.0169 $\mu$m$^2$ in a 95% confidence interval between [0.0197;0.0213], ubular PVs

(blue, N=835) have an average area of 0.0453±0.0278 $\mu$m$^2$ in a 95% confidence interval between

[0.0435;0.0472]. and polymorphic PVs (magenta, N=400) have an average area of

0.0981±0.0429 $\mu$m$^2$ in a 95% confidence interval between [0.0939;0.102]. The differences in area

are statistically significant (ANOVA; p-value <0.0001). Scale bars: (A) 5 $\mu$m and 1 $\mu$m (ROIs), (B)

5 $\mu$m.

1211

**Figure 3 - Super Resolution imaging of *Giardia lamblia* PVs by Stochastic Object**

**Reconstruction Microscopy (STORM).** (A) Widefield microscopy-based imaging of a Giardia

trophozoite loaded with 10kDa Dextran-Alexa Fluor 647. (B) Reconstruction of single molecule

events using the Fiji plugin Thunderstorm. As with STED imaging, different PV morphologies are

observed. (C) PVs were segmented with the help of ilastik and volumes were calculated and

plotted in (D). Spherical PVs (green, N=1989) present an average volume of 0.00507±0.00336

$\mu$m$^3$ with a 95% confidence interval between [0.00492;0.00522] $\mu$m$^3$, tubular PVs (blue, N=838)

present an average volume of 0.0103±0.00925 $\mu$m$^3$ in a 95% confidence interval between

[0.00967, 0.0109] $\mu$m$^3$. Polymorphic PVs (magenta, N=1494) present an average volume of

0.0227±0.0214 $\mu$m$^3$ in a 95% confidence interval between [0.0216, 0.0238] $\mu$m$^3$. The differences

in area are statistically significant (ANOVA; p-value <0.0001). Based on these and previously

shown data, the renaming of PVs to Peripheral Endocytic Compartments (PECs), is proposed.

Scale bars: 5 $\mu$m and 1 $\mu$m (ROI),.

1225

**Figure 4 – Uptake of fluorescently-labelled dextran in metamonada and discoba members: *G. lamblia*, *S. vortens*, *S. salmonicida* and *T. foetus* after 30 minutes.**

(A) *G. lamblia* cells present endocytic compartments spread on the cell periphery not resolved by conventional light microscopy. On the other hand, (B) *S. vortens*, (C) *S. salmonicida* and (D) *T. foetus* present vesicular shaped endocytic compartments. Scale bars: 5 $\mu$m (full cells) and 1 $\mu$m (ROI).

**Figure 5 – Super Resolution imaging of *S. vortens* endocytic compartments with STED and Transmission Electron Microscopy (TEM)** (A) Following incubation with dextran-TexasRed for 30 minutes, *S. vortens* display elongated endocytic compartments with an average diameter of 468±206 nm in a 95% confidence interval between [421;515] nm (upper box-plot). (B) TEM imaging detects several endosome-like vacuoles (V) throughout the cell,  with an average diameter of 844±335 nm in a 95% confidence interval between [763;905] nm (lower box-plot). Measurements were done manually. The Endoplasmic Reticulum (ER) and a nucleus (N) are also highlighted in the images. Scale bars: 5 $\mu$m (full field of view) and 1 $\mu$m (ROIs).

**Figure 6 – Super Resolution Imaging of *T. foetus* endocytic compartments with STED and TEM .** (A) 2D-STED analysis of *T. foetus* cells loaded with 10 kDa Dextran-Alexa Fluor 594 illuminate endosomes with globular structures at an average diameter of 517±251 nm in a 95% confidence interval between [455;580] nm (upper box-plot). (B) TEM investigation of *T. foetus* cells reveals two kinds of endosome-like vesicles. Low electron-density vesicles found both at the cell periphery and centre are deemed vacuoles (V) with an average diameter of 246±100 nm (N=153) in a 95% confidence interval of [230;262] nm. Vesicles of higher electron-density and with noticeable content are  Digestive Vacuoles (DVs).  DVs are larger than vacuoles, with an average diameter of 764±203 nm (N=50)  in a 95% confidence interval of [707;822] nm (lower box-plot). Scale bars: 5 $\mu$m (full field of view) and 1 $\mu$m (ROIs).

**Figure 7 – Time course on fluorescent dextran uptake in selected metamonada and discoba specimens:  *G. lamblia*, *S. vortens*, *S. salmonicida* and *T. foetus* after 30 minutes.**

**(A)** PV/PECs in *G. lamblia* start acquiring external material right after 5 minutes of incubation with dextran. Over 30 minutes of dextran incubation, the endocytic marker does not leave PECs. In (B) *S. vortens*, (C) *S. salmonicida* and (D) *T. foetus,* dextran is up taken in small vesicles. In *S. vortens* these small vesicles tend to agglomerate at the centre of the cell while in *S. salmonicida*,

1259    these vesicles seem to stay peripheral. In *T. foetus*, vesicles bearing dextran from at the periphery

1260    of the cell and also migrate to the interior of the cell. All scale bars: 5 μm.

1261

**Figure 8 – *Gl*CHC and *Gl*CLC foci associate to *Giardia* PECs with different stoichiometries.**

(A) An epitope-tagged *Gl*CHC reporter was used to localise foci of GlCHC deposition at the cell

periphery beneath the PM, by STED microscopy .(B) These foci can be clearly resolved with

STED microscopy (left graph). (C) The same was done with an epitope-tagged *Gl*CLC reporter

and (D) foci resolved as in (B) (right graph). (E) Foci were segmented with ilastik. Areas were

determined and diameters calculated in an automatic procedure assuming spherical geometry.

*Gl*CHC foci present an average diameter of 134±36.6 nm (N = 4524) in a 95% confidence interval

between [132;135] nm. *Gl*CLC foci average a diameter of 159±48.8 nm (N = 984) in a 95%

confidence interval between [156;162] nm. *Gl*CLC foci are larger than *Gl*CHC in significant

statistical manner (p-value < 0.0001, t-student test). (F) *Gl*CHC foci associate in different

stoichiometry to different classes of PECs. Spherical PECs are either not associated to clathrin

or with just one focus: mean of 0.488±0.159 foci per spherical in a 95% confidence interval

between [0.403;0.573] foci per spherical organelle. Tubular PECs associate with one focus of

*Gl*CHC: mean of 1.15±0.287 in a 95% confidence interval between [0.994;1.3] foci per tubular

organelle. Polymorphic PECs associate with 3 or more *Gl*CHC foci with an average of 3.85±1.14

in a 95% confidence interval between [3.25;4.46] foci per polymorphic organelle. These

distributions are statistically significative with a p-value < 0.0001 (ANOVA analysis).

1279

**Figure 9 – Homology search for *bona fide* CHC and CLC reveal key importance of CHC and**

**patchy conservation of CLC.** (A) CHC and CLC homology searches demonstrate the

conservation of CHC orthologs in all analysed lineages. However, several lineages appeared to

have lost a readily-detectable *bona fide* CLC including all selected Fornicata, certain Chromerids

(Woo et al. 2015), some ciliates (Richardson and Dacks 2022) and members of the

Hemimastigophora. (B) *Ab initio in silico* protein modelling of *G. lamblia* ACLC (formerly

*Gl*CLC/Gl4529), *T. brucei* CLC, *T. foetus* CLC, *M. exilis* CLC, *D. discoideum* and *H. sapiens* CLC

sequences using AlphaFold, the current standard in *in silico protein* structure modelling. TM-align

and RMSD values were calculated showing close structure analogy between predicted structures

(table).

1290

**Figure 10 –Endosome-like organelle models in *Giardia, Spironucleus* and *Tritrichomonas,***

**and proposed evolution of CLC.** (A) Simplified cartoons of the endocytic and secretory pathway

in *Giardia lamblia* (Giardiinae), *Spironucleus sp.* (Hexamitidae) and *Tritrichomonas foetus*
(Parabasalia). PECs are a hallmark of the Giardia lineage, while more canonical vesicular
endosomes are present in both the Spironucleus lineage and Parabasalia. (B) Simplified
evolutionary model for *bona fide* CLC. The last eukaryotic common ancestor possessed a *bona*
*fide* CLC which was lost at the last Fornicata common ancestor. In at least two derived lineages,
– Giardinae and *Spironucleus* spp. - de *novo protein* analogue to CLC was acquired
independently.

1300

**Supplementary figures**
**Supplementary figure 1 – Rendering of a *G. lamblia* trophozoite scanned with FIB-SEM**
**reveals the cell's inner ultrastructure.** (A) 3D view of acquired FIB-SEM trophozoite data. (B)
Single slice showing inner cellular structures such as cytoskeleton elements at the median body
(MB), the ventral disk (VD), the endoplasmic reticulum (ER), mitosomes (m) and peripheral
vacuoles (PV), highlighted in the region of interest (ROI). (C) Segmentation of different categories
of the dataset: cell volume (138 $\mu m^3$), cytoskeleton, endoplasmic reticulum, peripheral vacuoles,
small vesicles and mitosomes. (D) Mitosome volume (N = 14, box-plot) was determined post
segmentation at an average volume of $0.001093\pm0.0005698\mu m^3$ in a 95% confidence interval
between [0.0007643, 0.001422] $\mu m^3$.

1311

**Supplementary figure 2 – Cryo-SEM of freeze-fractured trophozoites reveals varying**
**vacuolar morphology in *Giardia lamblia*.** (A) Overview of cryo-preserved *Giardia* trophozoites
subjected to freeze-fracture and SEM imaging. Nuclei (N), Endoplasmic Reticulum (ER), Ventral
Disk (VD) and peripheral endocytic compartments (PEC) and plasma membrane (PM) are clearly
identifiable. (B and C) Insets showing different PEC morphology: vesicular (asterisk) and tubular
(hashtag). Scale bar: (A) 2 $\mu m$ and (B and C) 500 nm.

1318

1319 **Supplementary figure 3 – TEM investigation of *Giardia lamblia* endocytic and secretory**
1320 **pathway.** (A) Overview of a trophozoite. Different PEC structures, vesicular and tubular are
1321 observed, together with small vesicles (SV). The N (nucleus) and ER are also highlighted. (B)
1322 Close up on tubular PECs (hashtag). (C) Close up on vesicular PECs (asterisk) and SVs
1323 (arrowhead). Scale bars: (A) 2 $\mu m$, (B) 1 $\mu m$ and (C) 500 nm.

1324

**Supplementary figure 4 – TEM investigation of *S. salmonicida* endocytic and secretory**
**pathway.** (A) *S. salmonicida* presents vacuolar formations close to the plasma membrane. Cells

also present a prominent endoplasmic reticulum (ER; blue-framed inset). (B) Highlight of vacuolar

formations (V) and ER. (C) Second cell displaying an abundance of PV close to its plasma

membrane. (D) Highlight of vacuoles (V) and the prominent ER that connects to the plasma

membrane (asterisk). (E) *S. salmonicida* PVs average a diameter of 205±62.6 nm (N=114) in a

95% confidence interval of [193;217]. (F) *S. vortens* peripheral vacuoles are larger than *S.*

*salmonicida* vacuoles in a statistically significant manner (p-value < 0.0001). Diameters were

manually determined. Scale bars: (A and C) 2 $\mu$m and (B and D) 500 nm.

1334

1335 **Supplementary figure 5 – TEM investigation of *T. foetus* Golgi vesicles**. (A) More than one

1336 Golgi apparatus (G) can be found per cell. These organelles resemble canonical stacked Golgi

1337 releasing small coated vesicles. (B) These vesicles average a diameter of 58.4±13.1 nm (N=128)

1338 in a 95% confidence interval of [56.1;60.7] nm. Scale bar: (A) 500 nm.

1339

**Supplementary figure 6 – Pan-Eukaryotic prediction of clathrin heavy chain protein**

**domains.** Pfam analysis of predicted protein domains for several clathrin heavy chain proteins

sequences from the following species: *Giardia lamblia, Spironucleus vortens, Spironucleus*

*salmonicida, Trepomonas sp., Hexamita inflata, Dysnectes brevis, Kipferlia bialata,*

*Carpediomonas membranifera, Aduncisulcus paluster, Chilomastix cuspidata, Trypanosoma*

*brucei, Naegleria gruberi, Tritrichomonas foetus, Monocercomonoides exilis, Tetrahymena*

*thermophila, Hemimastix kukwestjiik, Chlamydomonas reinhardtii, Dyctiostilium discoideum,*

*Saccharomyces cerevisiae, Caenorhabditis elegans, Homo sapiens, Salpingoeca rosetta,*

*Capsospora owczarzaki* and *Monosiga brevicollis*. A general decrease in domain complexity is

observed in excavates compared with higher eukaryotes. CLOs: Carpediomonas-like organisms.

Diplom: Diplomonada.

1351

1352 **Supplementary figure 7 – The QLMLT motif is exclusive to Holozoa.** Alignment of the C-

1353 terminii of CHC sequences from selected Opisthokonta, Archaeplastida, Amoebozoa and SAR

1354 species highlights the present of the QLMLT uncoating motif only in Holozoa supergroup. The

1355 positioning of the QLMLT is highlighted in blue.

1356

1357 **Supplemental figure 8 – Calculation of *Giardia* ACLC synonymous vs non- synonymous**

1358 **mutation ratio ($\omega$ = ks/kn).** (A) Phylogenetic tree resulting of maximum likelihood analysis of the

1359 *Giardia* ACLC sequences. Each node is represented by a number. (B) Overall $\omega$ < 1 indicating

1360 there is no selective pressure on *Giardia* ACLC.

1361

**Supplemental figure 9 – *S. salmonicida* CHC (*Ss*CHC) is distributed in the cell cytosol in**
**foci and does interact with a structural form of CLC.** (A) *Ss*CHC was tagged C-terminally with
three HA tags. It is found throughout the cell cytosol. Signal is observed in 88% of the analyzed
cells (N = 171). (B) High resolution imaging of *Ss*CHC using confocal imaging reveals CHC foci.
(C) Distribution of the 171 proteins found in higher abundance in the *Ss*CHC native co-IP dataset.
(D) Native co-IP of tagged *Ss*CHC reporter reveals interaction with several members of the
endocytic pathway such as dynamin or beta-adaptin and calmodulin. (E) *Ab initio in silico* protein
modelling with AlphaFold of *Ss*11905, *Gl*ACLC, *Tb*CLC and *Hs*CLC. TM-align and RMSD scores
for predicted structures of *Giardia* ACLC, *Trypanosoma brucei* CLC and Ss11905 with respect to
*Homo sapiens* CLC show overall structural conservation with respect to a *bona fide* CLC. Scale
bars: (A) 20 $\mu$m. (B) 5 $\mu$m.

1373

1374 **Supplementary figure 10 – *Trepomonas* sp. PC1 also harbours a putative CLC analogue.**
1375 *Ab initio* protein modelling of TPC1_16039, orthologous to *Ss*11905 in combination with *Ss*11905,
1376 *Gl*ACLC, *Tb*CLC and *Hs*CLC. RMSD and TM-align cores show overall structural conservation
1377 with respect to a *bona fide* CLC.

1378

1379 **Supplementary Videos**
1380 **Supplementary Video 1 –** Three-dimensional rendering of endocytic compartments in *G. lamblia*
1381 derived from FIB-SEM sectioning and imaging. Scale bar 1 $\mu$m.

1382

1383 **Supplementary Video 2 –** Comparison between confocal and STED imaging of *Giardia* PECs.
1384 Scale bar: 3 $\mu$m.

1385

1386 **Supplementary Video 3 –** Tri-dimensional reconstruction of PV/PECs from STORM data.

1387

1388 **Supplementary Video 4 –** Tri-dimensional confocal imaging of *S. vortens* with Dextran-Texas
1389 Red. Both peripheral and near-nuclear endosome-like vacuoles are observed.

1390

1391 **Supplementary Video 5 -** Tri-dimensional STED imaging of *S. vortens* with Dextran Alexa Fluor
1392 594 reveals endosome-like vacuoles in greater detail.

1393

1394    **Supplementary Video 6 -** Tri-dimensional STED imaging of *T. foetus* with Dextran Alexa Fluor

1395    594 reveals endosome-like vacuoles in greater detail.

1396

1397    **Supplemental Video 7 –** Tri-dimensional high resolution confocal imaging and representation of

1398    *Ss*CHC-3xHA foci in the cell cytoplasm.

1399

1400    **Supplementary Tables**

1401    **Supplementary Table 1-** PECs volume comparison as calculated in FIB-SEM and STORM

1402    experiments

1403

**Supplementary Tables 2-9 (one file)**

Supplementary Table 2 - Queries used for CHC HHM profile building

Supplementary Table  3 - Results from Pan-Eukaryotic search of CHC homologues in available

Proteomes

Supplementary Table 5 - Queries used for CLC HHM profile building

Supplementary Table 6 - Results for GlCLC search in available Giardia genomes

Supplementary Table 7 - Results for bona fide CLC present in other genomes/transcriptomes

Supplementary Table 8 - Comparison of GlCLC with bona fide CLC

Supplementary Table 9 - Ten best hits from HHPred

1413

1414    **Supplementary Table 10** -  Search for domains from Clathrin heavy chain super family repeats

1415

1416    **Supplementary Table 11** - Modelled sequences for CLC and alike

1417

1418    **Supplementary Table 12** - SsCHC co-IP results
1419

# Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

# Figure 8

figure 9

A

| | CHC | bona fide CLC | GACLC |
|---|---|---|---|
| *Giardia lamblia* AWB | ■ | □ | ■ |
| *G. lamblia* ADH | ■ | □ | ■ |
| *G. lamblia* AS175 | ■ | □ | ■ |
| *G. lamblia* EP15 | ■ | □ | ■ |
| *G. lamblia* EP15 | ■ | □ | ■ |
| *G. lamblia* BGS | ■ | □ | ■ |
| *G. lamblia* BGS-B | ■ | □ | ■ |
| *G. lamblia* BAH | ■ | □ | ■ |
| *G. lamblia* Assemblage C | ■ | □ | ■ |
| *G. lamblia* Assemblage D | ■ | □ | ■ |
| *G. muris* | ■ | □ | ■ |

| | CHC | bona fide CLC |
|---|---|---|
| *Spironucleus salmonicida* | ■ | □ |
| *S. barkhanus* | ■ | □ |
| *S. vortens* | ■ | □ |
| *Hexamita inflata* | ■ | □ |
| *Trepomonas sp.* PC1 | ▦ | ▢ |
| *Retortamonas dobelli* | ▦ | ▢ |
| *Retortamonas caviae* | ▦ | ▢ |
| *D. brevis* | ▦ | ▢ |
| *C. cuspidata* | ▦ | ▢ |
| *K. bialata* | ■ | □ |
| *A. paluster* | ▦ | ▢ |
| *M. exilis* | ■ | ■ |
| *T. foetus* | ■ | ■ |
| *Trichomonas vaginalis* | ■ | ■ |
| *Bodo saltans* | ■ | ■ |
| *Trypanosoma brucei* | ■ | ■ |
| *Leishmania donovani* | ■ | ■ |
| *Naegleria gruberi* | ■ | ■ |
| *Naegleria fowleri* | ■ | ■ |

| | CHC | bona fide CLC |
|---|---|---|
| *Cryptosporidium parvum* | ■ | □ |
| *Babesia bovis* | ■ | □ |
| *Toxoplasma gondii* | ■ | ■ |
| *Plasmodium falciparum* | ■ | ■ |
| *Tetrahymena thermophila* | ■ | □ |
| *Paramecium caudatum* | ■ | □ |
| *Hemimastix kukwestjijk* | ▦ | ▢ |
| *Spironema multiciliatum* | ▦ | ▢ |
| *Disctyostelium discodeum* | ■ | ■ |
| *Arabidopsis thaliana* | ■ | ■ |
| *Saccharomyces cerevisiae* | ■ | ■ |
| *Homo sapiens* | ■ | ■ |
| *Capsaspora owczarzaki* | ■ | ■ |
| *Salpingoeca rosetta* | ▦ | ▦ |

■ Present in genome   ▦ Present in transcriptome
□ Absent in genome   ▢ Absent in transcriptome

B

*G. lamblia* ACLC

*M. exilis* CLC

*T. foetus* CLC

*T. brucei* CLC

*D. discodeum* CLC

*H. sapiens* CLC

| Specie/TM-align | *H. sapiens* | *D. discodeum* | *M. exilis* | *T. brucei* | *T. foetus* | *G. lamblia* |
|---|---|---|---|---|---|---|
| *H. sapiens* | 1 | 0.48716 | 0.44078 | 0.41651 | 0.46168 | 0.48174 |
| *D. discodeum* | 0.48716 | 1 | 0.57744 | 0.47487 | 0.49614 | 0.51481 |
| *M. exilis* | 0.44078 | 0.57744 | 1 | 0.4525 | 0.45185 | 0.51904 |
| *T. brucei* | 0.41651 | 0.47487 | 0.4525 | 1 | 0.54812 | 0.47487 |
| *T. foetus* | 0.46168 | 0.49614 | 0.45185 | 0.54812 | 1 | 0.59364 |
| *G. lamblia* | 0.48174 | 0.51481 | 0.51904 | 0.47487 | 0.59364 | 1 |

| Specie/RMSD (Å) | *H. sapiens* | *D. discodeum* | *M. exilis* | *T. brucei* | *T. foetus* | *G. lamblia* |
|---|---|---|---|---|---|---|
| *H. sapiens* | 0 | 4.338 | 6.205 | 4.581 | 5.821 | 4.979 |
| *D. discodeum* | 4.338 | 0 | 3.08 | 10.905 | 3.718 | 6.231 |
| *M. exilis* | 6.205 | 3.08 | 0 | 5.286 | 3.067 | 6.205 |
| *T. brucei* | 4.581 | 10.905 | 5.286 | 0 | 5.492 | 6.173 |
| *T. foetus* | 5.821 | 3.718 | 3.067 | 5.492 | 0 | 4.74 |
| *G. lamblia* | 4.979 | 6.231 | 6.205 | 6.173 | 4.74 | 0 |