IL-33 priming and antigenic stimulation synergistically promote the

transcription of proinflammatory cytokine and chemokine genes in human

skin mast cells

Junfeng Gao[1], Yapeng Li[1], Xiaoyu Guan[1], Zahraa Mohammed[2,3], Gregorio Gomez[4], Yvonne

Hui[2], Dianzheng Zhao[1], Carole A. Oskeritzian[2], Hua Huang[1,5*]

[1] Department of Immunology and Genomic Medicine, National Jewish Health, Denver, CO,

USA; [2]Department of Pathology, Microbiology and Immunology, University of South Carolina

School of Medicine, Columbia, SC, USA; [3]College of Medicine, AI-mustansiriyah university,

Baghdad, Iraq; [4]Department of Biomedical Sciences, University of Houston College of Medicine,

Houston, TX, USA; [5]Department of Immunology and Microbiology, University of Colorado

School of Medicine, Denver, CO, USA.

[*]Address correspondence and reprint requests to Dr. Hua Huang, Department of Immunology

and Genomic Medicine, National Jewish Health, 1400 Jackson Street, Denver, CO 80206. E-mail

address: huangh@njhealth.org

Abstract

Background Antigenic stimulation through cross-linking the IgE receptor and epithelial cell-derived cytokine IL-33 are potent stimuli of mast cell (MC) activation. Moreover, IL-33 primes a variety of cell types, including MCs to respond more vigorously to external stimuli. However, target genes induced by the combined IL-33 priming and antigenic stimulation have not been investigated in human skin mast cells (HSMCs) in a genome-wide manner. Furthermore, epigenetic changes induced by the combined IL-33 priming and antigenic stimulation have not been evaluated.

Results We found that IL-33 priming of HSMCs enhanced their capacity to promote transcriptional synergy of the *IL1B* and *CXCL8* genes by 16- and 3-fold, respectively, in response to combined IL-33 and antigen stimulation compared to without IL-33 priming. We identified the target genes in IL-33-primed HSMCs in response to the combined IL-33 and antigenic stimulation using RNA sequencing (RNA-seq). We found that the majority of genes synergistically upregulated in the IL-33-primed HSMCs in response to the combined IL-33 and antigenic stimulation were predominantly proinflammatory cytokine and chemokine genes. Moreover, the combined IL-33 priming and antigenic stimulation increase chromatin accessibility in the synergy target genes but not synergistically. Transcription factor binding motif analysis revealed more binding sites for NF-κB, AP-1, GABPA, and RAP1 in the induced or increased chromatin accessible regions of the synergy target genes.

Conclusions Our study demonstrates that IL-33 priming greatly potentiates MCs' ability to transcribe proinflammatory cytokine and chemokine genes in response to antigenic stimulation, shining light on how epithelial cell-derived cytokine IL-33 can cause exacerbation of skin MC-mediated allergic inflammation.

Keywords: Human skin mast cells; IgE receptor cross-linking; IL-33 priming; Chromatin accessibility; RNA-seq analysis; Pro-inflammatory cytokine and chemokine genes.

Background

Mast cells (MCs) are long-lived tissue-resident cells with evolutionarily conserved functions in

many inflammatory settings, including allergic inflammation and host defense against pathogen

infection [1-3]. The release of these mediators initiates a cascade of immune responses, resulting

in allergic reactions. MCs are dispersed throughout the boundaries between the external

environment and tissues, such as around blood vessels, in the skin, and at mucosal surfaces.

Because of their strategic locations, MCs are among the first exposed to antigenic stimulations

and cytokines and chemokines secreted by injured epithelial cells and other inflammatory cells at

the site of inflammation. MCs express FcεRI, the high-affinity receptor for IgE, which consists

of one IgE-binding α subunit, one signaling β subunit, and a signaling γ subunit dimer. MCs can

be activated by antigen (Ag) crosslinking of FcεRI-bound Ag-specific IgE (IgECL), which is

produced in an adaptive immune [4]. Ag crosslinking of the IgE/FcεRI complex induces MC

release of histamine, which is pre-made and stored in MC granules, and de novo gene

transcription [5, 6].

Epithelial cell-derived cytokines IL-25, IL-33, and TSLP have been reported to activate

MCs to amplify inflammation [7, 8]. IL-33 is a member of the IL-1 family [9, 10]. It is expressed

in epithelial cells, dendritic cells, and macrophages in tissues such as the skin, colon, lymph

nodes, and lung [11, 12]. Recent research has highlighted the role of interleukin-33 (IL-33) in

modulating skin inflammatory cascades and its potential as a therapeutic target for atopic

dermatitis [13, 14]. IL-33 protein is released from injured epithelial cells upon infections and

inflammation as IL-33 stored in the nucleus of destroyed epithelial cells is set free [10, 15]. MCs

express high levels of IL-33 (IL1RL1) receptor [16-18]. IL-33 activates MAP kinases and

induces NF-κB phosphorylation [12]. IL-33 promotes the differentiation of MC progenitor cells

into mature MCs that produce Th2 cytokines IL-5 and IL-13 [18]. Moreover, it directly activates

MCs to produce pro-inflammatory cytokines and chemokines, including IL-6, IL-10, TNF, GM-

CSF, CXCL8, and CCL1 [7, 19]. It synergizes with TSLP to promote MC differentiation and

activation [20]. IL-33 also synergizes with IgE receptor activation of primary human MCs to

promote cytokine and chemokine production without IL-33 priming [7, 21]. While previous

studies have used qPCR to measure the synergistic effects of antigen and IL-33 stimulations on

mRNA expression of *CCL1*, *CCL2*, *CXCL1*, *CXCL2*, *CXCL8*, *IL1B*, *IL5*, *IL6*, *IL13* genes in

various MCs [7, 19, 21, 22], genome-wide profiling has not been conducted.

Moreover, the role of IL-33 priming in enhancing the transcription of cytokine and

chemokine genes in MCs has not been clearly established. IL-33 primes a variety of cell types,

including MCs to respond more vigorously to external stimuli [23, 24]. IL-33 has been shown to

play a crucial role in promoting IgE-mediated allergic inflammation in vivo [25]. Notably,

studies have demonstrated that IL-4, IL-5, and IL-13 levels are increased in ovalbumin (OVA)-

sensitized mice treated with IL-33, compared to mice treated with IL-33 alone or OVA alone

[26], suggesting that prolonged exposure to IL-33 can further increase MC responses to antigenic

stimulation.

In this study, we sought to determine whether pretreatment with IL-33 for 24 hours can

promote gene expression in response to the combined IL-33 and IgECL stimulation compared to

without IL-33 pretreatment. We demonstrate that IL-33 priming of HSMCs enhanced their

capacity to promote transcriptional synergy of the *IL1B* and *CXCL8* genes in response to antigen

stimulation compared to without IL-33 priming. We identified the target genes in the IL-33-

primed HSMCs upon combined IL-33 and antigenic stimulation using RNA sequencing (RNA-

seq). We found that the majority of genes synergistically upregulated in the IL-33-primed

HSMCs in response to the combined IL-33 and antigen stimulation were predominantly

proinflammatory cytokine and chemokine genes. Moreover, the combined IL-33 and antigen

stimulation increased chromatin accessibility in the synergy genes but not synergistically in the

IL-33-primed HSMCs. Our study shines a light on a mechanism by which IL-33 might cause

exacerbation of skin MC-mediated allergic inflammation.

Results

The combined IL-33 priming and antigenic stimulation promote expression of two clusters of genes

To investigate the synergistic gene transcription promoted by antigen and epithelial cell-derived cytokines IL-33 in human skin-derived mast cells (HSMCs), we first compared whether pretreatment with IL-33 for 24 hours can promote gene expression in response to the combined IL-33 and IgECL stimulation compared to without IL-33 pretreatment. We chose to measure *IL1B* and *CXCL8* mRNA expression, which is known to be upregulated by the combined IgECL and IL-33 stimulation [7, 19-22] by qPCR. We used a published method to analyze transcriptional synergy, which is defined transcriptional synergy as greater than additive effects and is calculated by the formula synergistic $\log_2 [E_{\text{IgECL+IL-33}}/(E_{\text{IgECL}}+E_{\text{IL-33}})]$ [7, 27, 28]. Specifically, log2FC>0 indicates positive transcriptional synergy, while $\log_2$FC<0 represents negative transcriptional synergy. A $\log_2$FC=0 indicates additive. We found that IL-33 pretreatment for 24 hours promoted transcriptional synergy induced by the combined IL-33 and IgECL stimulation ($\log_2$FC *IL1B*=3.5, *CXCL8*=2.9), resulting in 16- and 3-fold increase in transcriptional synergy for *IL1B* and *CXCL8,* respectively, compared to without IL-33 pretreatment ($\log_2$FC *IL1B*=1.5, *CXCL8*=1.9) (Fig. 1). Statistical differences among the single and combined treatments were analyzed using two-way ANOVA as recommended by the published paper [29]. Our data demonstrate that although with or without IL-33 pretreatment, IgECL and IL-33 stimulation can both promote transcriptional synergy for the *IL1B* and *CXCL8* genes, pretreatment with IL-33 dramatically enhances the transcriptional synergy.

Figure 1. IL-33 priming and antigenic stimulation synergistically promote the transcription of *IL1B* and *CXCL8* genes in HSMCs. HSMCs were not treated (UN) or treated with IgE receptor cross-linking (IgECL) for 2 hours, IL-33 for 2 hours or 26 hours (pre-treated for 24 hours prior to the addition of the other stimuli treatment for 2 hours), or in combination of IgECL and IL-33 (2 hours or 26 hours). qPCR analysis of the mRNA expression of *IL1B* and *CXCL8*. Statistical differences were analyzed using two-way ANOVA. Data represents mean ± SEM of four biological samples. *** $P < 0.001$.

Given that many molecular events can occur during the 24-hour pretreatment, we thus

referred to the pretreatment with IL-25, TSLP, or IL-33 for 24 hours in the combined

stimulations as priming. For the combined treatments, we pretreated HSMCs with cytokines IL-

25, TSLP, also produced by epithelial cells, or IL-33 for 24 hours before subjecting them to the

combined cytokines and IgECL stimulation for an additional 2 hours. We found that IgECL

stimulation alone upregulated 501 genes with increased expression greater than 2-fold and 37

genes with increased expression greater than 10-fold (Fig. 2A, 3A and Table S1, sheet 1). We

found that IL-33 priming alone upregulated 259 genes that were expressed 2-fold greater and 22

genes with increased expression greater than 10-fold (Fig. 2B, 3A and Table S1, sheet 2).

Whereas IL-25 priming alone only upregulated nine genes greater than 2-fold with none of the

genes increased expression greater than 10-fold (Fig. 2C, 3A and Table S1, sheet 3). TSLP

upregulated 17 genes greater than 2-fold with one gene increased expression greater than 10-fold

(Fig. 2D, 3A and Table S1, sheet 4). These results indicate that IL-33 is a more potent stimulus

than IL-25 or TSLP in promoting gene expression in HSMCs.

Figure 2. Differential mRNA expression analysis of RNA-seq prepared from HSMCs that were exposed to various treatments. Heatmap representations of top ranked genes after IgE receptor cross-linking treatment alone (A), IL-33 treatment alone (B), IL-25 treatment alone (C), or TSLP treatment alone (D). Legends: un, not treated; E, IgECL; IL-33, IL-33 treatment; IL-25, IL-25 treatment; T, TSLP treatment; $\log_2$FC, $\log_2$ fold change. The numbers indicate RNA reads (RPKM). Data A-D represent two biological samples.

To determine whether any of the combinatorial stimulations may also upregulate HSMC

genes, we used the K mean clustering method to group genes into clusters with similar

expressions. We first determined the optimal K numbers of six using Elbow Plot (Fig. S1).

Heatmap K means clustering analysis revealed that the combined IL-33 and antigen treatment in

the IL-33-primed HSMCs upregulated the expression of two major clusters of genes (cluster four

and cluster six) relative to IgECL or IL-33 single stimulation (Fig. 3B). There were 902 genes in

cluster four and 641 genes in cluster six (Table S2). IL-25 and TSLP, when added with IgECL,

also promoted expression of genes in cluster four. However, the magnitude of upregulation

induced by IL-25 and TSLP was lower compared to IL-33. IL-25 and IgECL upregulated 50

genes compared with IgECL alone (Fig. 3A, S2A and Table S3, sheet 1). TSLP and IgECL

upregulated 27 genes compared with IgECL alone (Fig. 3A, S2B and Table S3, sheet 2). The

addition of IL-25 and TSLP to the IgECL and IL-33 combination did not significantly alter MC

gene expression profiles compared with the IgECL and IL-33 combination (Fig. 3A, S2C and

Table S3, sheet3). These results demonstrated that epithelial cell-derived cytokines IL-33

together with IgECL promoted the expressions of cluster four and six HSMC genes.

A

B

Figure 3. IL-33 priming acts synergically with IgECL to promote gene transcription in clusters four and six. HSMCs were not treated (UN), treated with IgE receptor cross-linking (IgECL) for 2 hours, IL-33 for 26 hours (priming for 24 hours prior to the addition of the other stimuli treatment for 2 hours), IL-25 for 26 hours, TSLP for 26 hours alone (single) or in combinations (combined) as indicated. (**A**) Venn diagram showing the overlap of upregulated differentially expressed genes among the four comparisons of IL-33 vs UN, IL-25 vs UN, IgECL vs UN, TSLP vs UN (left panel), and the comparisons of IgECL+ IL-33 vs IgECL, IgECL+ IL-25 vs IgECL, IgECL+ TSLP vs IgECL (right panel). (**B**) K-means clustering analysis was performed using R package pheatmap. The cluster size numbers indicate the number of genes that are grouped in the cluster. RNA-seq data were generated from two biological samples.

The combined IL-33 and antigen stimulation in the IL-33-primed HSMCs synergistically

induce pro-inflammatory gene transcription

We performed gene ontology (GO) analysis on genes in clusters four and six to determine which

types of genes were enriched among genes that were promoted by the combined IL-33 and

antigen stimulation in the IL-33-primed HSMCs. We found that genes encoding for cytokines

(Cluster four, $p=1.0\times10^{-5}$; cluster six, $p=2.5\times10^{-5}$, Fig. 4A), receptors and receptor binding

(Cluster four, $p=7.9\times10^{-2}$; cluster six, $p=3.6\times10^{-2}$, Fig. 4A), signaling molecules (Cluster four,

$p=9.7\times10^{-6}$; cluster six, $p=7.3\times10^{-3}$, Fig. 4A), transcription factors (Cluster four, $p=1.8\times10^{-2}$;

cluster six, $p=2.6\times10^{-3}$, Fig. 4A), NF-κB signaling pathway (Cluster four: $p=6.5\times10^{-2}$, cluster six:

$p=4.2\times10^{-2}$, Fig. 4A), chemokines (Cluster four: $p=1.1\times10^{-5}$; cluster six, $p=9.7\times10^{-2}$, Fig. 4A),

and molecules involved in cell death (Cluster four, $p=2.6\times10^{-2}$; cluster six: $p=1.7\times10^{-2}$, Fig. 4A)

were significantly enriched in both clusters.

The top-ranked representative genes in clusters four and six are listed in Fig. 4B. Genes

encoding cytokines and chemokines family in cluster four were most significantly enriched.

They include *CXCL8*, *CXCL3*, *TNFSF4*, *IL33*, *IL6*, *CCL20*, *IL23A*, *CCL1*, *CXCL2*, and *IL13*

(Fig. 4B). Genes enriched in the cytokines and chemokines gene family in cluster six include

IL1B, *IL5*, *CSF3*, *CXCL1* and *IL1A*. While *CXCL8*, *IL5*, *IL13*, and *IL6* have been reported to be

upregulated synergistically by the combined IL-33 and antigen stimulation [22, 30, 31], *TNFSF4*,

a member of TNF family, *CXCL3*, *IL33*, *CCL20*, *IL23A*, *CCL1*, *CXCL2*, *CXCL1*, *CSF3*, and

IL1A, have not been previously reported as target genes of the combined IL-33 and antigen

stimulation in HSMCs.

Genes encoding signaling molecules in cluster four were also highly enriched (Fig. 4A).

These genes include *RGS2*, which potently interferes with signaling through receptors that

couple to Gαq [32], and *RGS16*, which inhibits signal transduction through the Gα13-RHO [33].

MTSS1 and *ITK* were enriched in cluster six (Fig. 4B). These genes were not reported previously

as target genes for the synergistic stimulation of IL-33 and antigen. We also found that genes

encoding receptor and receptor binding molecules were significantly enriched in clusters four

and six (Fig. 4A). These genes included *IL7R*, a receptor for IL-7 and TSLP [34-36], *FCGR2A,*

which encodes a cell surface receptor belonging to the family of immunoglobulin Fc receptor,

and *CD274*, encoding PD-L1, a ligand that binds to the receptor PD1 commonly found on T-

cells to block T-cell activation [37, 38] (Fig. 4B).

A

Go analysis

Cluster four

	Gene counts	P-value
Cytokines		1.00E-05
Receptors and receptor binding		7.90E-02
Signaling molecules		9.70E-06
Transcription factors		1.80E-02
NF-κB signaling pathway		6.50E-02
Chemokines		1.10E-05
Cell death		2.60E-02

Cluster six

	Gene counts	P-value
Cytokines		2.50E-05
Receptors and receptor binding		3.60E-02
Signaling molecules		7.30E-03
Transcription factors		2.60E-03
NF-κB signaling pathway		4.20E-02
Chemokines		9.70E-02
Cell death		1.70E-02

B

Representative genes

Cluster four

Cytokines and chemokines

Gene	UN	IL33	IgECL	E+33	Log$_2$FC
CXCL2	0.3	1	1.7	144	5.7
CXCL3	0.4	0.3	2.8	155.9	5.7
CXCL8	15.6	6	185.1	5533.2	4.9
IL23A	0.2	0.3	0.6	10.4	3.5
IL33	0.9	0.5	1.6	10.9	2.4
CCL1	36.9	106	199.6	1173.5	1.9
CCL20	0	0.9	2.6	12.2	1.8
TNFSF4	5.4	2.2	12.7	48.4	1.7
IL6	10.9	17.7	54.7	229	1.7
IL13	1.3	4.4	6.6	21.7	1.0

Receptors and receptor binding

Gene	UN	IL33	IgECL	E+33	Log$_2$FC
CD200	0.1	0.3	8.9	19	1.0
CD274	17.5	13	36.5	89.9	0.9
HEG1	5.6	7.1	7.6	26.2	0.8
DUSP1	9.6	6.5	19.8	46.7	0.8
LRP12	9.8	10.1	13.5	31.5	0.4

Signaling molecules

Gene	UN	IL33	IgECL	E+33	Log$_2$FC
RGS16	3.1	1.9	7.2	25.8	1.5
GADD45A	12.4	17.8	54.1	147.1	1.0
RGS2	126.8	63.6	288.6	596.1	0.8
VCAN	2.9	2.6	4	10.3	0.6
LAPTM4B	12.2	6.5	8.5	17.4	0.2

Transcription factors

Gene	UN	IL33	IgECL	E+33	Log$_2$FC
EGR4	0.4	0.1	4.1	14.3	1.8
CREB5	1.3	0.4	5.9	19.8	1.7
ZFP36L2	82.4	43.3	154	461.2	1.2
MLF1	2	1.7	6.6	18.6	1.2
RFX2	2.6	8	21.1	52.3	0.8
CEBPD	4.1	5.9	13	32.7	0.8

NF-κB signaling pathway

Gene	UN	IL33	IgECL	E+33	Log$_2$FC
REL	10.1	8.7	22.7	67.4	1.1
TNFRSF21	5.5	4	6.7	20.3	0.9
RIPK2	17.8	22.9	27.7	92	0.9
MAP3K8	12.9	19.3	67	146.1	0.8
GBP2	38.4	27.2	82.5	179.3	0.7
NFKB1	20.1	38.6	40.1	112.2	0.5

Cell death

Gene	UN	IL33	IgECL	E+33	Log$_2$FC
BCL2A1	5.6	4.7	41.9	94	1.0
ACKR3	2.3	8.8	9.5	29.1	0.7
BCL6	6	5.7	7.2	17.8	0.5

Cluster six

Cytokines and chemokines

Gene	UN	IL33	IgECL	E+33	Log$_2$FC
CSF3	0.2	1.2	0.9	17.4	3.1
CXCL1	0.6	1.2	1.1	17.4	2.9
IL1B	0.3	53.8	0.6	367.7	2.8
IL5	0.2	6.5	0.4	35.9	2.4
IL1A	1.3	3.9	3.9	29.6	1.9

Receptors and receptor binding

Gene	UN	IL33	IgECL	E+33	Log$_2$FC
MFSD2A	0.8	7.4	3.2	33.3	1.7
IL7R	0.6	21.8	1.5	48.3	1.1
FCGR2A	10	27.9	7.3	57.5	0.7
TNFRSF9	0.4	9.9	2.9	20.9	0.7
HLA-E	343.4	460.3	434.2	991.9	0.1

Signaling molecules

Gene	UN	IL33	IgECL	E+33	Log$_2$FC
ITK	8.3	8.2	7	23.9	0.7
MTSS1	14.6	42	28.8	96.3	0.4
FSTL1	12.2	10.6	10.1	21.7	0.1

Transcription factors

Gene	UN	IL33	IgECL	E+33	Log$_2$FC
SKIL	5.6	14.3	5.3	32.5	0.7
TWIST2	9.6	16.8	14.9	36	0.2
TWIST1	11.3	23.9	21.5	49.5	0.1
YBX3	14.6	20.5	18.3	41.6	0.1

NF-κB signaling pathway

Gene	UN	IL33	IgECL	E+33	Log$_2$FC
TNFAIP3	3.7	28.9	11.6	65.1	0.7
TNFAIP2	4.1	8.3	6.7	23	0.6
NFKBIZ	48.4	76.7	68.2	201	0.5
NFKBIA	173.3	283.5	173	575.5	0.3

Cell death

Gene	UN	IL33	IgECL	E+33	Log$_2$FC
NINJ1	21	59.5	33.4	141	0.6
TNFAIP8	48	43.5	30.3	93.3	0.3
G0S2	14.2	42.4	39.9	95.4	0.2

Intensity: 1 | 25 | 50 | 500 | 1000

one-side Fisher's exact test with the adjustment of Benjamini-Hochberg method. (**B**) Heatmap representation of top ranked genes in each gene family in clusters four and six. The numbers indicate RNA reads (FPKM). $\log_2$FC: $\log_2[E_{IgECL+IL-33}/(E_{IgECL}+E_{IL-33})]$. Data represents the average reads generated from two biological samples.

Many genes encoding molecules involved in cell survival and cell death were identified in

clusters four and six (Fig. 4A). *BCL2A1* is a highly regulated NF-κB target gene that exerts

important pro-survival functions [39]. BCL6 acts to prevent apoptotic cell death in

differentiating myocytes [40]. *ACKR3*, which encodes a receptor for CXCL12 and CXCL11

[41], has been reported to inhibit apoptosis [42]. TNF Alpha Induced Protein 8 (TNFAIP8) was

showed to increase cell survival and decrease apoptosis in hepatocellular carcinoma cells [43].

On the other hand, we also found that IL-33 and IgECL stimulation upregulated genes involved

in apoptosis, including *G0S2,* which promotes apoptosis by binding to BCL2. *NINJ1,* which

encodes NINJURIN-1 [44], has been shown to mediate plasma membrane rupture during lytic

cell death [45] and promote apoptosis in macrophages during early ocular development [46].

Genes encoding transcription factors were found to be significantly enriched in clusters four

and six (Fig. 4A). EGR4, activated by NFAT in response to antigenic stimulation, regulates pro-

inflammatory cytokine gene expression [47], whereas CEBPD results in increased *IL6* gene

transcription [48]. Transcription factors *CREB5*, *MLF1*, *ZFP36L2* and *RFX2* were enriched in

cluster four, while transcription factors *SKIL*, *TWIST2*, *YBX3* and *TWIST1* were enriched in

cluster six (Fig. 4B). Genes encoding NF-κB signaling pathway were also highly enriched in

both clusters four and six (Fig. 4A). These genes included *NFKB1*, encoding p50 subunit of NF-

κB transcription factor complex that can lead to amplified transcription outputs for many of the

NF-κB target genes, such as *IL1B*, *IL6*, *CXCL8*, *TNF*, *MCP1*, and *CXCL1* [49]. *REL*,

TNFRSF21, *MAP3K8*, *GBP2* and *RIPK2* were enriched in cluster four, while *TNFAIP3*,

NFKBIA, *NFKBIZ* and *TNFAIP2* were enriched in cluster six (Fig. 4B). Our results demonstrate

that proinflammatory cytokine and chemokine genes were significantly enriched among genes

highly responsive to the combined IL-33 and antigen stimulation in the IL-33-primed HSMCs as

those genes with $\log_2$FC>1.0.

RNA-seq results were validated by performing a qPCR analysis on selected genes whose

products can promote MC survival, form positive feed-forward loops, or enable MCs to perform

new functions using HSMCs derived from four independent donors. We confirmed the

transcriptional synergy for *CXCL2* ($\log_2$FC=3.9), *CXCL8* ($\log_2$FC=1.9), *IL1B* ($\log_2$FC=2.0),

CCL1 ($\log_2$FC=1.4), *NAMPT* ($\log_2$FC=1.6), *IL7R* ($\log_2$FC=0.9), *CD274* ($\log_2$FC=0.3), *FCGR2A*

($\log_2$FC=0.3), *NFKB1* ($\log_2$FC=0.1) and *TNFAIP8* ($\log_2$FC=0.5) in the IL-33-primed HSMCs in

response to the combined IL-33 and IgECL stimulation (Fig. 5A). Consistent with the

transcriptional synergy, synergy at the protein level was also detected for IL1B ($\log_2$FC=1.9) and

CXCL8 ($\log_2$FC=3.1) (Fig. 5B). CXCL2 protein expression was also measured but was not

detectable using the commercial CXCL2 ELISA kit.

Figure 5. Validation of gene expression promoted by the combined IL-33 priming and IgE receptor cross-linking. HSMCs were not stimulated (UN) or stimulated with IgE receptor cross-linking (IgECL) for 2 hours, IL-33 for 26 hours (priming for 24 hours prior to the addition of the other stimuli treatment for 2 hours), or in combination of IgECL and IL-33. (**A**) qPCR analysis of the mRNA expression of activation-induced genes identified by RNA-seq. (**B**) The same treatments as described under (A) were applied, supernatants were collected after 24 hours and IL-1B and CXCL8 were quantified by ELISA. Statistical differences were analyzed using two-way ANOVA. Data represents mean ± SEM of four biological samples. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

The combined IL-33 and antigen stimulation increases chromatin accessibility in the

synergy genes but not synergistically in the IL-33-primed HSMCs

Enhancers integrate signal inputs and convert them into transcriptional outputs [50]. Thus, it is

likely that the combined IL-33 and antigen treatment activate signal-dependent transcription

factors that, in turn, bind to enhancers to increase their chromatin accessibility in the IL-33-

primed HSMCs. To investigate this, we performed Omni-ATAC-seq analysis on the HSMCs that

were primed with IL-33 for 24 hours and treated with the combined IL-33 and IgECL

stimulation for 2 hours or HSMCs that were treated IL-33 alone for 26 hours or IgECL alone for

2 hours. We used differential bind analysis (Diffbind) to compare ATAC-seq narrow peaks from

the IL-33-primed HSMCs treated with the combined IL-33 and IgECL stimulation or from

HSMCs that were treated with either IL-33 or IgECL alone. We assigned ATAC-seq peaks

located within 100 kb upstream or downstream of the transcription start site (TSS) of a gene

using Binding and Expression Target Analysis (BETA) [51]. Tens of thousands of narrow peaks

(21,451), which were significantly enriched peaks compared to pooled and normalized

background, increased significantly in the IL-33-primed HSMCs treated with the combined IL-

33 and IgECL stimulation compared to IL-33 stimulated alone (Table S4). Fewer narrow peaks

(521) were found to be significantly elevated when compared with IgECL stimulated alone

(Table S4).

Figure 6. IL-33 priming and IgECL increase chromatin accessibility in the synergy target genes. (**A**) The numbers of upregulated ATAC-seq narrow peaks in the combined IL-33 priming and antigenic stimulation relative to IgECL or IL-33 stimulation were subtracted using Diffbind analysis. ATAC-seq was performed on HSMCs under various conditions, IgECL, IL-33 or IgECL plus IL-33. ATAC-seq narrow peaks (NPs) of *CXCL2* (**B**), *CXCL8* (**C**), *IL1B* (**D**) genes in HSMCs treated with IgECL and IL-33 (E+33) are shown (blue bars). Differential NPs generated using DiffBind comparing IgECL+IL-33 with IgECL (E+33/E NPs) or comparing IgECL+IL-33 with IL-33 (E+33/33 NPs) are also shown (blue bars). Candidate enhancers are marked with red bars and E - or + numbers, indicating distances (kb) from the TSS of the indicated genes. RPM, reads per million nucleotides mapped. The IGV tracks are generated from one biological sample, representing two biological replicates with similar patterns.

We analyzed the differences in the number of upregulated narrow peaks in positive

transcriptional synergy genes (referred to as synergy target genes, with log2FC>0) and non-

positive transcriptional synergy genes (referred to as non-synergy target genes, with log2FC≤0,

Table S5). A significant genome-wide difference in upregulated narrow peaks was observed

between synergy target genes and non-synergy target gene in response to IL-33 priming and

antigenic stimulation (Fig. 6A). In Fig. 6B-D, we focused our analysis on the top-ranked *CXCL2*,

CXCL8, and *IL1B* genes and assessed the normalized values of differentially increased ATAC-

seq narrow peaks in the samples primed with IL-33 and treated with combined IL-33 and IgECL

stimulation compared to those treated with either IgECL or IL-33 alone. We observed that the

combined IL-33 and IgECL treatment led to increased chromatin accessibility, as indicated by

higher narrow peak values at several candidate enhancers of the synergy target genes.

Significantly increased narrow peak values were found for *CXCL2* gene at E+65 [IgE+IL-33/IL-

33 (E+33/33) ratio=1.6], E+42 (E+33/33=1.5), PE (E+33/33=1.1), and E-10 (E+33/33=1.5)

compared to IL-33 priming alone (Fig. 6B). Additionally, combined IL-33 and IgECL treatment

led to significantly increased narrow peak values at E-35 (E+33/33=1.3), E-18 (E+33/33=1.5), E-

5 (E+33/33=2.0), PE (E+33/33=1.3), and E+3 (E+33/33=1.4) of the *CXCL8* gene, compared to

IL-33 priming alone (Fig. 6C). Furthermore, significantly increased narrow peak values were

observed at E-18 (E+33/E=1.3), PE (E+33/E=1.3), and E+3 (E+33/E=1.5) of the *CXCL8* gene

when compared to IgECL stimulation alone (Fig. 6C). In the *IL1B* gene, significant increases in

narrow peak values were found at E+86 (E+33/33=1.6), E+71 (E+33/33=1.4), E+55

(E+33/33=1.5), E-3 (E+33/33=1.5), and E-43 (E+33/33=1.4) when compared to the IL-33

priming alone (Fig. 6D). However, no differential increased narrow peaks were observed for

CXCL2 and *IL1B* genes in comparison to IgECL alone (Fig. 6B, D).

Only E-18, PE, and E+3 enhancers of the *CXCL8* showed increased narrow peaks in HSMCs

treated with combined IL-33 and IgECL when compared to either IL-33 treatment or IgECL

stimulation alone. The E+33/33 ratios were 1.5, 1.3, 1.4, for E-18, PE, E+3, respectively.

Additionally, the E+33/E ratios were 1.3, 1.3, 1.5 for E-18, PE, E+3, respectively (Fig. 6C).

These results indicate that the combined IL-33 priming and IgECL stimulation resulted in

increased ATAC-seq peak values at certain enhancers of the synergy genes *CXCL2*, *CXCL8*, and

IL1B in HSMCs but were less than additive compared to IL-33 priming or IgECL stimulation

alone.

We analyzed transcription factor binding motifs in the chromatin accessible areas that were

upregulated by the combined IL-33 and antigen stimulation in the IL-33-primed HSMCs relative

to IgECL stimulation or IL-33 treatment alone using MEME-ChIP [52]. We found that NF-κB,

AP-1, RAP1, and GABPA binding motifs were significantly enriched in the increased narrow

peaks (Fig. 7). Our results suggest that genes responsiveness to the combined IL-33 and antigen

stimulation in the IL-33-primed HSMCs is predetermined by the transcriptional codes embedded

in the gene sequences. Furthermore, the combined IgECL and IL-33 treatment might achieve

greater synergy in promoting gene transcription of the synergy target genes by binding to NF-κB,

AP-1, GABPA and RAP1 binding sites.

Discussion

The results of this study provide direct evidence that IL-33 primes HSMCs with synergically

enhanced capacity to respond to the combined IL-33 and antigen treatment. IL-33 is known to

prime variety of cell types to respond more vigorously to external stimuli [23, 24]. It has been

reported that IL-33 primes mouse MCs with amplified responses to ATP and IgG-mediated

activation [53, 54]. Our study adds a new IL-33 priming capacity in HSMCs to the existing list.

Compared with previous work that used a 15-minute IL-33 pretreatment [7], our work is

designed to directly assess the effect of a 24 hour of IL-33 pretreatment on the upregulation of

genes in HSMCs using a genome-wide and unbiased method. We demonstrate that IL-33

priming resulted in a 16-fold increase in transcriptional synergy for *IL1B* and a 3-fold increase

for *CXCL8*. Similar to findings by the previous report [7], we confirmed a remarkable synergy at

the protein level for CXCL8. Mechanisms by which this transcriptional synergy is not yet clear.

A previous study used rat basophilic leukemia MCs, RBL-2H3 to demonstrate that multiple

copies of NF-κB and NFAT binding sites were critical in antigen and IL-33 stimulation-driven

reporter gene transcription. Our transcription factor motif analysis on chromatin accessible

regions that were promoted by IL-33 priming and the subsequent combined IL-33 and antigen

stimulation indeed established significantly enriched NF-κB binding motif and AP-1 binding

motif, supporting previous conclusion that IL-33 and antigen stimulations synergistically activate

NF-κB and NFAT to promote gene transcription of pro-inflammatory cytokines and chemokines

TNF, *IL6*, *IL13*, *CCL2* (*MCP1*), *CCL7* (*MCP3*) and *CCL3* (*MIP1A*) [22]. However, NF-κB and

NFAT are signal-dependent transcription factors. Activation of these transcription factors is

rapid within minutes of receptor binding [55]. Therefore, the previous finding does not offer a

complete explanation of IL-33 priming action. One best-studied example of cytokine priming is

IL-4 priming for Th2 cell differentiation. IL-4 priming induces upregulation of GATA3, which

then reprograms Th2 cells to acquire the capacity to transcribe the *Il4* gene [56]. It is possible a

major transcription factor that depend on NF-κB and NFAT can be upregulated by IL-33 priming

in HSMCs, conferring dramatically enhanced capacity to transcribe the cytokine and chemokine

genes in the later response to antigenic stimulation or the combined IL-33 and antigen

stimulation.

Our transcription factor motif analysis on IL-33 priming and antigenic stimulation

responsive enhancers also provide some clues to the candidate transcription factors that are

induced by IL-33 priming. Given that *NFKB1* (also known as p50) and *REL* (also known as p68)

are the two subunits of NF-κB transcription factor complex and that GBP2, a guanine-binding

protein, RIPK2, a member of the receptor-interacting protein (RIP) family of serine/threonine

protein kinases, are known potent activator of the NF-κB pathway, the putative transcription

factor whose induction dependent on NF-κB signaling can form a positive feed-forward loops to

enhance HSMCs' capacity to respond a combined secondary antigenic or the combined IL-33

and antigen stimulation. Interestingly, we also found potential negative feedback loops formed

by TNFAIP2, TNFAIP3, NFKB1A, and NFKB1Z whose coding genes were significantly

increased by the combined IL-33 and antigen stimulation. Therefore, upregulation of these gene

transcriptions might provide critical control of NF-κB activation.

In addition to NF-κB binding motif and AP-1 binding motifs, we found more binding

sites in the chromatin accessible regions that were promoted by combined IL-33 priming and

antigen stimulation for GABPA, which binds to GA repeats and is a paralog gene of ETS2.

While GABPA mRNA was expressed at low level revealed by our RNA-seq analysis, several

ETS2 family transcription factor genes *ETS2*, *ELK1*, *ELF1*, *SPI1,* whose gene products can bind

to GABPA binding site, were expressed in HSMCs. Whether these ETS2 family transcription

factors can be recruited to bind GABPA binding sites in response to the combined IL-33 priming

and antigenic stimulation remains to be determined.

Enhancers integrate signal inputs through recruiting signal-dependent transcription factor

to their cognate transcription factor binding sites in the enhancers. Binding of transcription

factors then recruit histone modifying enzymes to change H3K27 acetylation, leading to changes

in chromatin accessibility [50, 57, 58]. We found IL-33 priming and the subsequent combined

IL-33 and antigen stimulation indeed increased chromatin accessibility at the enhancers of

synergy target genes. However, the chromatin accessibility increases induced by these processes

did not reach additive or synergistic levels compared to IL-33 priming or antigen stimulation

alone. These results suggest that the degree of changes in chromatin accessibility at individual

enhancers might not correlate with the degree of transcriptional synergy. Further research

deciphering transcriptional codes contained in the IL-33 priming responsive enhancers of

synergy target genes could provide insights into the mechanisms underlying IL-33 priming for

antigen-mediated MC activation and potentially lead to the development of targeted therapies for

infection-exacerbation of allergic inflammation.

Conclusions

In conclusion, our study provides evidence that IL-33 and IgECL stimulation can promote

transcriptional synergy for mast cell genes, and IL-33 priming significantly enhances this

synergy. We also found that combined IL-33 and antigen stimulation lead to the upregulation of

two gene clusters in IL-33-primed HSMCs, cluster four (902 genes) and cluster six (641 genes),

as indicated by heatmap K means clustering analysis. We demonstrate that proinflammatory

cytokine and chemokine genes were significantly enriched among genes highly responsive to the

combined IL-33 and antigen stimulation in the IL-33-primed HSMCs. Furthermore, analysis of

Chromatin Accessibility revealed that although the combined IL-33 and antigen treatment in the

IL-33-primed HSMCs amplifies chromatin accessibility in top-ranked *CXCL2*, *CXCL8* and *IL1B*

genes, albeit the enhancement is not synergistic in nature. Overall, our study sheds light on the

mechanism by which the epithelial cell-derived cytokine IL-33 may exacerbate skin MC-

mediated allergic inflammation.

Materials and Methods

Human skin mast cell (HSMC) culture

De-identified fresh skin biopsies were obtained after breast reduction or mastectomy from the

Cooperative Human Tissue Network of the National Cancer Institute. Handling human skin

tissues was approved by the University of South Carolina Institutional Review Board. Human

skin MCs were cultured from human skin essentially as described [59, 60]. After removing

subcutaneous fat by blunt dissection, residual tissue was cut into 1- to 2-mm fragments then

digested with type 2 collagenase (1.5 mg/ml), hyaluronidase (0.7 mg/ml), and type 1 DNase

(0.15 mg/ml) in HBSS for 1 h at 37°C, for a total of three cycles of 1 h digestion. After each

cycle, the dispersed cells were collected by filtering through a no. 80 mesh sieve and suspended

in HBSS containing 1% FCS and 10 mM HEPES. Cells were suspended in HBSS, layered over a

Percoll cushion, and centrifuged at $700 \times g$ at room temperature for 20 min. Nucleated cells were

collected from the buffer/Percoll interface. Cells enriched by Percoll density-dependent

sedimentation were adjusted to a concentration of 1×10^6 cells/ml in serum-free X-VIVO^{TM} 15

medium (Lonza BioWhittaker) containing 100 ng/ml recombinant human SCF (Peprotech) in 24-

well plates. The culture medium was changed weekly, and wells were split when cell

concentration doubled. The percentages of mast cells were assessed cytochemical by

metachromatic staining with toluidine blue. Mature mast cells at ~ 100% purity and viability are

obtained by 6-8 weeks of culture, and 8- to 16-week-old mast cells were used in these

experiments. HSMCs were shipped to the Huang laboratory at National Jewish Health for RNA-

seq and qPCR analysis.

HSMCs stimulations

IgE receptor cross-linking (IgECL) on HSMCs was accomplished through sensitization with 1

µg/mL chimeric human IgE anti-NP antibody (MCA333S, BIO-RAD, Hercules, CA) for 24

hours. The sensitized HSMCs were challenged with 100 ng/mL NP-HSA (N-5059, LGC,

Petaluma, CA) for two additional hours prior to performing RNA-seq, ATAC-seq, or

Quantitative RT-PCR analysis. For single or combined cytokine and IgECL treatments, IL-33

(50 ng/mL, 200-33, PEPROTECH, Cranbury, NJ), TSLP (50 ng/mL, 300-62, PEPROTECH,

Cranbury, NJ) or IL-25 (50 ng/mL, 200-24, PEPROTECH, Cranbury, NJ) were added to HSMCs

for a total of 26 hours (pre-treated for 24 hours prior to the addition of the other stimuli treatment

for 2 hours).

RNA-seq

Total RNA was extracted from HSMCs that were untreated or treated with various stimuli using

the RNeasy mini kit (Qiagen, Valencia, CA) according to the manufacturer's protocol. The

RNA-seq libraries were prepared as described before [61]. Briefly, the quality of input RNA was

analyzed by running the RNA sample on the 4150 TapeStation System (Agilent, CA). PolyA

mRNA was isolated using the NEBNext® Poly(A) mRNA magnetic isolation module (E7490,

New England Biolabs, Ipswich, MA). RNA-seq libraries were obtained using the NEB-Next®

Ultra™ II RNA library prep kit for Illumina® (E7770S, New England Biolabs, Ipswich, MA).

The isolated mRNA was fragmented, reverse transcribed into cDNA, end-repaired, and adapter-

ligated. The ligated samples were digested by USER enzyme, amplified by PCR, and purified

using AMPure-XP beads (Beckman Coulter Life Sciences, A63881, Indianapolis, IN). The

quantity and quality of RNA libraries were assessed on an Agilent Technologies 2100

Bioanalyzer. Paired-ended sequencing of the RNA libraries was performed on an Illumina

NovaSEQ6000 platform.

Omni-ATAC-seq

Omni-ATAC-seq, an improved ATAC-seq protocol, was performed according to the published

method [61, 62]. Briefly, 50,000 HSMCs that were untreated, treated with IL-33 alone, or IgECL

with or without IL-33 were spun down and washed once with cold PBS. The cells were

resuspended in 50 μl cold ATAC-RSB-lysis buffer and incubated for 3 minutes. The ATAC-

RSB-lysis buffer was immediately washed out with 1 mL ATAC-RSB buffer. The cell pellet was

resuspended in 50 μl transposition mix and incubated for 30 minutes at 37 °C. The reaction was

stopped by adding 2.5 μl pH 8 0.5 M EDTA. The Qiagen MiniElute PCR purification kit

(Qiagen) was used to purify the transposed DNA. Purified DNA was amplified using the

following conditions: 72°C for 5 min, 98 °C for 30 s, and 13 cycles: 98 °C for 10s, 63 °C for 30

s, 72 °C for 1min. The amplified libraries were purified, size-selected, and the quality and

quantity of libraries were assessed on 4150 TapeStation System (Agilent, CA). The pair-ended

sequencing of DNA libraries was performed on an Illumina NovaSEQ6000 platform.

RNA-seq data analysis

The RNA-seq data was analyzed as described before [61]. Briefly, the raw reads (average 20

million paired end reads, two biological replicates for each treatment) were analyzed and quality

checked by FastQC. The reads were aligned to the hg38 reference genome using the Spliced

Transcripts Alignment to a Reference (STAR, version 2.4.0.1) software. Reads (FPKM) were

assembled into reference transcripts and counted using Cufflinks (version 2.2.1). The average

reads from two biological samples were calculated using Cuffmerge (version 1.0.0). The

differential gene expression between the resting and stimulated samples was analyzed using

Cuffdiff (version 2.2.1). K-means clustering was performed using pheatmap (version 1.0.12).

The optimal numbers of K in the RNA-seq datasets were determined using R packages Tidyverse

(version 1.3.1), Cluster (version 2.1.0), Factoextra (version 1.0.7), and GridExtra (2.3). Genes in

clusters were extracted using pheatmap (version 1.0.12).

Omni-ATAC-seq data analysis

The Omni-ATAC-seq data was analyzed as described before [61]. Briefly, raw sequencing reads

(average 80 million paired-end reads, two biological replicates for each treatment) were aligned

to the hg38 reference genome using Bowtie2 with very-sensitive and -x 2000 parameters. The

read alignments were filtered using SAMtools to remove mitochondrial genome and PCR

duplicates. Narrow peaks were called by MACS2 with the q-value cut-off of 0.05, and the

sequencing data was displayed using Integrative Genomics Viewer (IGV).

Quantitative RT-PCR

For mRNA expression analysis, the HSMCs were untreated, treated with IL-33 or IgECL alone

or in combination. Total RNA from HSMCs was isolated with TRIzol^{TM} Reagent according to

the manufacturer's instructions. cDNA was synthesized by reverse transcription. Quantitative

PCR was performed in a QuantStudio 7 Flex Real-Time PCR System (ThermoFisher, MA). The

sequences of qPCR primers are listed in Table S6. Relative mRNA amounts were calculated as

follows: Relative mRNA amount $= 2^{[Ct(Sample)-Ct(HPRT)]}$.

ELISA measurement of chemokine and cytokine productions

HSMCs were pretreated with IL-33 (50 ng/mL, 200-33, PEPROTECH, Cranbury, NJ), or

sensitized with 1 mg/mL chimeric human IgE anti-NP antibody (MCA333S, BIO-RAD,

Hercules, CA), or in combination for 24 hours. Then washed before stimulation with 100 ng/mL

NP-HSA (N-5059, LGC, Petaluma, CA) or 50 ng/mL IL-33, or in combination for additional 24

hours. Untreated HSMCs were included as a control. The amounts of CXCL8 (88-8086-22,

Thermo Fisher Scientific), IL1B (88-7261-22, Thermo Fisher Scientific), and CXCL2 (900-

M120, Peprotech) protein in the supernatants were measured by ELISA.

DAVID GO analysis

497 Gene ontology enrichment analysis was performed on genes that responded to the synergistic

498 stimulations of IgECL plus IL-33 by using the Database for Annotation, Visualization, and

499 Integrated Discovery (DAVID, version 6.8). The top-ranked genes in each enriched category

500 were selected for heatmap representations using an Excel spreadsheet.

Assigning regulatory regions to genes

Binding and Expression Target Analysis (BETA) minus (Version 1.0.7) used to calculate the

regulatory potential score and assign the regulatory regions identified by Omi-ATAC-seq to their

putative target genes [51] with the following setting: Reference genome, Human: GRCh38;

Distance: within 100 kb upstream and 100 kb downstream from the transcriptional start site of

the gene.

DiffBind

For analysis of activation-induced accessible regions, Omni-ATAC-seq reads of HSMCs

samples that were treated with IL-33 and IgECL were compared with those of HSMCs samples

treated with IL-33 or IgECL using the R package DiffBind (version 2.10.0) [63]. The number of

upregulated peaks in the synergy genes were calculated by adding the number of short peaks

within 100 kb upstream and 100 kb downstream from TSS were significantly increased in

response to the combined IL-33 priming and antigenic stimulation relative to IL-33 or IgECL.

TF binding motif enrichment analysis

Multiple Em for Motif Elicitation (MEME)-ChIP performs comprehensive motif analysis on

large sets of sequences such as those identified by ChIP-seq or CLIP-seq experiments [52].

MEME-ChIP was used to identify the TF binding motifs enriched in accessible regions. To

identify the TF binding motifs enriched in the induced accessible regions, BED files of IgE-

stimulated and IgE+IL-33-stimulated were uploaded to the Find Individual Motif Occurrences

(FIMO) tool of MEME (FIMO) [64]. The number of transcription factor binding sites in the

increased narrow peaks of the synergy genes was calculated by adding the number of

transcription factor binding sites within the narrow peaks. The densities of transcription factor

binding sites in the increased narrow peaks were normalized by the sizes of narrow peaks and

expressed as the number of binding sites per kilobase.

Data availability

All raw and processed sequencing data generated in this study have been submitted to the NCBI

Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/) under accession number

GSE196895. The RNA-seq data can be found in the GEO database: "GSE196862"; The ATAC-

seq data can be found in the GEO database: "GSE196883".

Statistical analysis

One-tail Fisher Exact probability value was used for GO analysis. Statistical differences among

the single and combined treatments were analyzed using two-way ANOVA [29]. Identifying

statistically significantly differentially bound sites in DiffBind used specific statistical methods

built in the Bioconductor packages EdgeR [65] and DEseq2 [66].

Abbreviations

HSMCs: Human skin mast cells; RNA-seq: RNA sequencing; ATAC-seq: Transposase-

accessible chromatin sequencing; IgECL: IgE cross-linking; TSLP: Thymic stromal

lymphopoietin; qRT-PCR: quantitative reverse transcription-PCR; NF-κB: Nuclear factor kappa-

light-chain-enhancer of activated B cells; AP-1: Activator protein 1; GABPA: GA-binding

protein alpha chain; RAP-1: Ras-related protein 1; NFAT: nuclear factor of activated T-cells;

MAP: mitogen-activated protein; NP: Nitrophenyl-hapten; GO: gene ontology; FPKM:

Fragments Per Kilobase of transcript per Million mapped reads; IGV: Integrative Genomics

Viewer; DAVID: Database for Annotation, Visualization, and Integrated Discovery; BETA:

Binding and Expression Target Analysis; MEME: Multiple Em for Motif Elicitation; TSS:

Transcription start site.

Declarations

Ethics approval and consent to participate

The study was performed in accordance with the Declaration of Helsinki on research involving

humans. The skin tissues used to obtain the mast cells for this study were purchased from the

Cooperative Human Tissue Network of the National Cancer Institute, and informed consent was

obtained from all participants. This study was reviewed by the Institutional Review Board (IRB)

of the University of South Carolina. The IRB of the University of South Carolina has determined

that the referenced study (protocol number Pro00020620) meets the Not Human Subject criteria

set forth as defined by the Code of Federal Regulations (45 CFR 46) as the specimens and/or

private information/data were not collected specifically for the current research project through

an interaction/intervention with living individuals and the investigators including collaborators

on the proposed research cannot readily ascertain the identity of the individuals to whom the

coded private information or specimens pertain. All methods and experiments were performed in

accordance with relevant guidelines and regulations.

Consent for publication

Not applicable

Availability of data and materials

The RNA-seq datasets and ATAC-seq datasets generated in this study have been submitted to the

NCBI Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/) under accession

number GSE196895.

Competing interests

The authors declare no competing interests.

Funding

Supported by grants from the National Institutes of Health R01AI107022 and R01AI135194

(H.H.), National Institutes of Health (NIH)/National Institute of General Medical Sciences

(NIGMS) Pilot Project P20GM103641 (C.A.O. and G.G.), a fund provided by a fellowship from

the Higher Committee for Education Development (HCED) and Ministry of Higher Education

and Scientific Research (MOHSR) in Iraq (Z.M.).

Authors' contributions

Conceptualization, J.G. and H.H.; methodology, J.G., Y.L., G.G., C.A.O. and H.H.;

investigation, J.G., Y.L., X.G., Z.M., D.Z. and Y.H.; visualization, J.G. and Y.L.; funding

acquisition, H.H., C.A.O. and G.G.; supervision, H.H., C.A.O. and G.G.; writing-original draft,

J.G. and H.H.; writing-review & editing, J.G., C.A.O. and H.H.

Acknowledgements

We thank lab members for the thoughtful discussions. We are grateful to Dr. Bifeng Gao and the

staff of the Genomics Shared Resource Facility at the University of Colorado Cancer Center for

Next-Generation Sequencing.

Authors' information

[1] Department of Immunology and Genomic Medicine, National Jewish Health, Denver, CO,

USA; [2]Department of Pathology, Microbiology and Immunology, University of South Carolina

School of Medicine, Columbia, SC, USA; [3]College of Medicine, AI-mustansiriyah university,

Baghdad, Iraq; [4]Department of Biomedical Sciences, University of Houston College of Medicine,

Houston, TX, USA; [5]Department of Immunology and Microbiology, University of Colorado

School of Medicine, Denver, CO, USA.

Footnotes

Correspondence address:

Address correspondence and reprint requests to Dr. Hua Huang, Department of Immunology and

Genomic Medicine, National Jewish Health, 1400 Jackson Street, Denver, CO 80206. E-mail

address: huangh@njhealth.org

References

1. Galli SJ, Grimbaldeston M, Tsai M: **Immunomodulatory mast cells: negative, as well as positive, regulators of immunity**. *Nature reviews Immunology* 2008, **8**(6):478-486.

2. Gurish MF, Austen KF: **The diverse roles of mast cells**. *The Journal of experimental medicine* 2001, **194**(1):F1-5.

3. Babina M, Franke K, Bal G: **How "Neuronal" Are Human Skin Mast Cells?** *Int J Mol Sci* 2022, **23**(18).

4. Galli SJ, Gaudenzio N, Tsai M: **Mast Cells in Inflammation and Disease: Recent Progress and Ongoing Concerns**. *Annu Rev Immunol* 2020, **38**:49-77.

5. Potaczek DP, Kabesch M: **Current concepts of IgE regulation and impact of genetic determinants**. *Clinical and experimental allergy : journal of the British Society for Allergy and Clinical Immunology* 2012, **42**(6):852-871.

6. Hellman L: **Regulation of IgE homeostasis, and the identification of potential targets for therapeutic intervention**. *Biomed Pharmacother* 2007, **61**(1):34-49.

7. Franke K, Wang Z, Zuberbier T, Babina M: **Cytokines Stimulated by IL-33 in Human Skin Mast Cells: Involvement of NF-kappaB and p38 at Distinct Levels and Potent Co-Operation with FcepsilonRI and MRGPRX2**. *Int J Mol Sci* 2021, **22**(7).

8. Divekar R, Kita H: **Recent advances in epithelium-derived cytokines (IL-33, IL-25, and thymic stromal lymphopoietin) and allergic inflammation**. *Curr Opin Allergy Clin Immunol* 2015, **15**(1):98-103.

9. Yagami A, Orihara K, Morita H, Futamura K, Hashimoto N, Matsumoto K, Saito H, Matsuda A: **IL-33 mediates inflammatory responses in human lung tissue cells**. *J Immunol* 2010, **185**(10):5743-5750.

10. Martin NT, Martin MU: **Interleukin 33 is a guardian of barriers and a local alarmin**. *Nature immunology* 2016, **17**(2):122-131.

11. Saenz SA, Taylor BC, Artis D: **Welcome to the neighborhood: epithelial cell-derived cytokines license innate and adaptive immune responses at mucosal sites**. *Immunol Rev* 2008, **226**:172-190.

12. Schmitz J, Owyang A, Oldham E, Song Y, Murphy E, McClanahan TK, Zurawski G, Moshrefi M, Qin J, Li X *et al*: **IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines**. *Immunity* 2005, **23**(5):479-490.

13. Chen YL, Gutowska-Owsiak D, Hardman CS, Westmoreland M, MacKenzie T, Cifuentes L, Waithe D, Lloyd-Lavery A, Marquette A, Londei M *et al*: **Proof-of-concept clinical trial of etokimab shows a key role for IL-33 in atopic dermatitis pathogenesis**. *Sci Transl Med* 2019, **11**(515).

14. Sroka-Tomaszewska J, Trzeciak M: **Molecular Mechanisms of Atopic Dermatitis Pathogenesis**. *Int J Mol Sci* 2021, **22**(8).

15. Saluja R, Khan M, Church MK, Maurer M: **The role of IL-33 and mast cells in allergy and inflammation**. *Clin Transl Allergy* 2015, **5**:33.

16. Lohning M, Stroehmann A, Coyle AJ, Grogan JL, Lin S, Gutierrez-Ramos JC, Levinson D, Radbruch A, Kamradt T: **T1/ST2 is preferentially expressed on murine Th2 cells, independent of interleukin 4, interleukin 5, and interleukin 10, and important for Th2**

effector function. *Proceedings of the National Academy of Sciences of the United States*
of America 1998, **95**(12):6930-6935.

17.   Moritz DR, Rodewald HR, Gheyselinck J, Klemenz R: **The IL-1 receptor-related T1 antigen**
is expressed on immature and mature mast cells and on fetal blood mast cell
progenitors. *J Immunol* 1998, **161**(9):4866-4874.

18.   Allakhverdi Z, Smith DE, Comeau MR, Delespesse G: **Cutting edge: The ST2 ligand IL-33**
potently activates and drives maturation of human mast cells. *J Immunol* 2007,
179(4):2051-2054.

19.   Bawazeer MA, Theoharides TC: **IL-33 stimulates human mast cell release of CCL5 and**
CCL2 via MAPK and NF-kappaB, inhibited by methoxyluteolin. *European journal of*
pharmacology 2019, **865**:172760.

20.   Saluja R, Zoltowska A, Ketelaar ME, Nilsson G: **IL-33 and Thymic Stromal Lymphopoietin**
in mast cell functions. *European journal of pharmacology* 2016, **778**:68-76.

21.   Silver MR, Margulis A, Wood N, Goldman SJ, Kasaian M, Chaudhary D: **IL-33 synergizes**
with IgE-dependent and IgE-independent agents to promote mast cell and basophil
activation. *Inflamm Res* 2010, **59**(3):207-218.

22.   Andrade MV, Iwaki S, Ropert C, Gazzinelli RT, Cunha-Melo JR, Beaven MA: **Amplification**
of cytokine production through synergistic activation of NFAT and AP-1 following
stimulation of mast cells with antigen and IL-33. *European journal of immunology* 2011,
41(3):760-772.

23.   Suzukawa M, Iikura M, Koketsu R, Nagase H, Tamura C, Komiya A, Nakae S, Matsushima
K, Ohta K, Yamamoto K *et al*: **An IL-1 cytokine member, IL-33, induces human basophil**
activation via its ST2 receptor. *J Immunol* 2008, **181**(9):5981-5989.

24.   Le HT, Tran VG, Kim W, Kim J, Cho HR, Kwon B: **IL-33 priming regulates multiple steps of**
the neutrophil-mediated anti-Candida albicans response by modulating TLR and
dectin-1 signals. *J Immunol* 2012, **189**(1):287-295.

25.   Kato Y, Morikawa T, Kato E, Yoshida K, Imoto Y, Sakashita M, Osawa Y, Takabayashi T,
Kubo M, Miura K *et al*: **Involvement of Activation of Mast Cells via IgE Signaling and**
Epithelial Cell-Derived Cytokines in the Pathogenesis of Pollen Food Allergy Syndrome
in a Murine Model. *J Immunol* 2021, **206**(12):2791-2802.

26.   Sjoberg LC, Nilsson AZ, Lei Y, Gregory JA, Adner M, Nilsson GP: **Interleukin 33**
exacerbates antigen driven airway hyperresponsiveness, inflammation and
remodeling in a mouse model of asthma. *Sci Rep* 2017, **7**(1):4219.

27.   Schrode N, Seah C, Deans PJM, Hoffman G, Brennand KJ: **Analysis framework and**
experimental design for evaluating synergy-driving gene expression. *Nat Protoc* 2021,
16(2):812-840.

28.   Foucquier J, Guedj M: **Analysis of drug combinations: current methodological**
landscape. *Pharmacol Res Perspect* 2015, **3**(3):e00149.

29.   Slinker BK: **The statistics of synergism**. *J Mol Cell Cardiol* 1998, **30**(4):723-731.

30.   Ndaw VS, Abebayehu D, Spence AJ, Paez PA, Kolawole EM, Taruselli MT, Caslin HL,
Chumanevich AP, Paranjape A, Baker B *et al*: **TGF-beta1 Suppresses IL-33-Induced Mast**
Cell Function. *J Immunol* 2017, **199**(3):866-873.

31.   Paranjape A, Chernushevich O, Qayum AA, Spence AJ, Taruselli MT, Abebayehu D,
Barnstein BO, McLeod JJ, Baker B, Bajaj GS *et al*: **Dexamethasone rapidly suppresses IL-**

33-stimulated mast cell function by blocking transcription factor activity. *J Leukoc Biol*
2016, **100**(6):1395-1404.

32.   Kehrl JH, Sinnarajah S: **RGS2: a multifunctional regulator of G-protein signaling.** *The*
international journal of biochemistry & cell biology 2002, **34**(5):432-438.

33.   Johnson EN, Seasholtz TM, Waheed AA, Kreutz B, Suzuki N, Kozasa T, Jones TL, Brown
JH, Druey KM: **RGS16 inhibits signalling through the G alpha 13-Rho axis.** *Nature cell*
biology 2003, **5**(12):1095-1103.

34.   Noguchi M, Nakamura Y, Russell SM, Ziegler SF, Tsang M, Cao X, Leonard WJ:
Interleukin-2 receptor gamma chain: a functional component of the interleukin-7
receptor. *Science* 1993, **262**(5141):1877-1880.

35.   Park LS, Martin U, Garka K, Gliniak B, Di Santo JP, Muller W, Largaespada DA, Copeland
NG, Jenkins NA, Farr AG *et al*: **Cloning of the murine thymic stromal lymphopoietin**
(TSLP) receptor: Formation of a functional heteromeric complex requires interleukin 7
receptor. *The Journal of experimental medicine* 2000, **192**(5):659-670.

36.   Pandey A, Ozaki K, Baumann H, Levin SD, Puel A, Farr AG, Ziegler SF, Leonard WJ, Lodish
HF: **Cloning of a receptor subunit required for signaling by thymic stromal**
lymphopoietin. *Nature immunology* 2000, **1**(1):59-64.

37.   Keir ME, Liang SC, Guleria I, Latchman YE, Qipo A, Albacker LA, Koulmanda M, Freeman
GJ, Sayegh MH, Sharpe AH: **Tissue expression of PD-L1 mediates peripheral T cell**
tolerance. *The Journal of experimental medicine* 2006, **203**(4):883-895.

38.   Noman MZ, Desantis G, Janji B, Hasmim M, Karray S, Dessen P, Bronte V, Chouaib S: **PD-**
L1 is a novel direct target of HIF-1alpha, and its blockade under hypoxia enhanced
MDSC-mediated T cell activation. *The Journal of experimental medicine* 2014,
211(5):781-790.

39.   Vogler M: **BCL2A1: the underdog in the BCL2 family.** *Cell death and differentiation*
2012, **19**(1):67-74.

40.   Kumagai T, Miki T, Kikuchi M, Fukuda T, Miyasaka N, Kamiyama R, Hirosawa S: **The**
proto-oncogene Bc16 inhibits apoptotic cell death in differentiation-induced mouse
myogenic cells. *Oncogene* 1999, **18**(2):467-475.

41.   Burns JM, Summers BC, Wang Y, Melikian A, Berahovich R, Miao Z, Penfold ME,
Sunshine MJ, Littman DR, Kuo CJ *et al*: **A novel chemokine receptor for SDF-1 and I-TAC**
involved in cell survival, cell adhesion, and tumor development. *The Journal of*
experimental medicine 2006, **203**(9):2201-2213.

42.   Hattermann K, Held-Feindt J, Lucius R, Muerkoster SS, Penfold ME, Schall TJ, Mentlein R:
The chemokine receptor CXCR7 is highly expressed in human glioma cells and
mediates antiapoptotic effects. *Cancer Res* 2010, **70**(8):3299-3308.

43.   Niture S, Gyamfi MA, Lin M, Chimeh U, Dong X, Zheng W, Moore J, Kumar D: **TNFAIP8**
regulates autophagy, cell steatosis, and promotes hepatocellular carcinoma cell
proliferation. *Cell Death Dis* 2020, **11**(3):178.

44.   Araki T, Milbrandt J: **Ninjurin, a novel adhesion molecule, is induced by nerve injury**
and promotes axonal growth. *Neuron* 1996, **17**(2):353-361.

45.   Kayagaki N, Kornfeld OS, Lee BL, Stowe IB, O'Rourke K, Li Q, Sandoval W, Yan D, Kang J,
Xu M *et al*: **NINJ1 mediates plasma membrane rupture during lytic cell death.** *Nature*
2021, **591**(7848):131-136.

46. Lee HJ, Ahn BJ, Shin MW, Jeong JW, Kim JH, Kim KW: **Ninjurin1 mediates macrophage-**
induced programmed cell death during early ocular development. *Cell death and*
differentiation 2009, **16**(10):1395-1407.
47. Decker EL, Nehmann N, Kampen E, Eibel H, Zipfel PF, Skerka C: **Early growth response**
proteins (EGR) and nuclear factors of activated T cells (NFAT) form heterodimers and
regulate proinflammatory cytokine gene expression. *Nucleic Acids Res* 2003, **31**(3):911-
921.
48. Hungness ES, Luo GJ, Pritts TA, Sun X, Robb BW, Hershko D, Hasselgren PO:
Transcription factors C/EBP-beta and -delta regulate IL-6 production in IL-1beta-
stimulated human enterocytes. *Journal of cellular physiology* 2002, **192**(1):64-70.
49. Liu T, Zhang L, Joo D, Sun SC: **NF-kappaB signaling in inflammation**. *Signal Transduct*
Target Ther 2017, **2**.
50. Spitz F, Furlong EE: **Transcription factors: from enhancer binding to developmental**
control. *Nature reviews Genetics* 2012, **13**(9):613-626.
51. Wang S, Sun H, Ma J, Zang C, Wang C, Wang J, Tang Q, Meyer CA, Zhang Y, Liu XS: **Target**
analysis by integration of transcriptome and ChIP-seq data with BETA. *Nat Protoc*
2013, **8**(12):2502-2515.
52. Machanick P, Bailey TL: **MEME-ChIP: motif analysis of large DNA datasets**.
Bioinformatics (Oxford, England) 2011, **27**(12):1696-1697.
53. Straus DB, Pryor D, Haque TT, Kee SA, Dailey JM, Jackson KG, Barnstein BO, Ryan JJ: **IL-33**
priming amplifies ATP-mediated mast cell cytokine production. *Cell Immunol* 2022,
371:104470.
54. Kaieda S, Wang JX, Shnayder R, Fishgal N, Hei H, Lee RT, Stevens RL, Nigrovic PA:
Interleukin-33 primes mast cells for activation by IgG immune complexes. *PloS one*
2012, **7**(10):e47252.
55. Zhang DX, Glass CK: **Towards an understanding of cell-specific functions of signal-**
dependent transcription factors. *Journal of molecular endocrinology* 2013, **51**(3):T37-
50.
56. Zhu J, Yamane H, Cote-Sierra J, Guo L, Paul WE: **GATA-3 promotes Th2 responses**
through three different mechanisms: induction of Th2 cytokine production, selective
growth of Th2 cells and inhibition of Th1 cell-specific factors. *Cell Res* 2006, **16**(1):3-10.
57. Schoenfelder S, Fraser P: **Long-range enhancer-promoter contacts in gene expression**
control. *Nat Rev Genet* 2019, **20**(8):437-455.
58. Reiter F, Wienerroither S, Stark A: **Combinatorial function of transcription factors and**
cofactors. *Current opinion in genetics & development* 2017, **43**:73-81.
59. Kambe N, Kambe M, Kochan JP, Schwartz LB: **Human skin-derived mast cells can**
proliferate while retaining their characteristic functional and protease phenotypes.
Blood 2001, **97**(7):2045-2052.
60. Caslin HL, Taruselli MT, Paranjape A, Kiwanuka K, Haque T, Chumanevich AP, Oskeritzian
CA, Ryan JJ: **The Use of Human and Mouse Mast Cell and Basophil Cultures to Assess**
Type 2 Inflammation. *Methods Mol Biol* 2018, **1799**:81-92.
61. Li Y, Gao J, Kamran M, Harmacek L, Danhorn T, Leach SM, O'Connor BP, Hagman JR,
Huang H: **GATA2 regulates mast cell identity and responsiveness to antigenic**

stimulation by promoting chromatin remodeling at super-enhancers. *Nat Commun*
2021, **12**(1):494.
62. Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna S,
Satpathy AT, Rubin AJ, Montine KS, Wu B *et al*: **An improved ATAC-seq protocol reduces**
background and enables interrogation of frozen tissues. *Nat Methods* 2017,
14(10):959-962.
63. Ross-Innes CS, Stark R, Teschendorff AE, Holmes KA, Ali HR, Dunning MJ, Brown GD,
Gojis O, Ellis IO, Green AR *et al*: **Differential oestrogen receptor binding is associated**
with clinical outcome in breast cancer. *Nature* 2012, **481**(7381):389-393.
64. Grant CE, Bailey TL, Noble WS: **FIMO: scanning for occurrences of a given motif.**
Bioinformatics (Oxford, England) 2011, **27**(7):1017-1018.
65. Robinson MD, McCarthy DJ, Smyth GK: **edgeR: a Bioconductor package for differential**
expression analysis of digital gene expression data. *Bioinformatics (Oxford, England)*
2010, **26**(1):139-140.
66. Love MI, Huber W, Anders S: **Moderated estimation of fold change and dispersion for**
RNA-seq data with DESeq2. *Genome Biol* 2014, **15**(12):550.
815

Figure Legends

Figure 1. IL-33 priming and antigenic stimulation synergistically promote the transcription of

IL1B and *CXCL8* genes in HSMCs. HSMCs were not treated (UN) or treated with IgE receptor

cross-linking (IgECL) for 2 hours, IL-33 for 2 hours or 26 hours (pre-treated for 24 hours prior to

the addition of the other stimuli treatment for 2 hours), or in combination of IgECL and IL-33 (2

hours or 26 hours). qPCR analysis of the mRNA expression of *IL1B* and *CXCL8*. Statistical

differences were analyzed using two-way ANOVA. Data represents mean ± SEM of four

biological samples. *** $P < 0.001$.

Figure 2. Differential mRNA expression analysis of RNA-seq prepared from HSMCs that were

exposed to various treatments. Heatmap representations of top ranked genes after IgE receptor

cross-linking treatment alone (A), IL-33 treatment alone (B), IL-25 treatment alone (C), or TSLP

treatment alone (D). Legends: un, not treated; E, IgECL; IL-33, IL-33 treatment; IL-25, IL-25

treatment; T, TSLP treatment; log$_2$FC, log$_2$ fold change. The numbers indicate RNA reads

(RPKM). Data A-D represent two biological samples.

Figure 3. IL-33 priming acts synergically with IgECL to promote gene transcription in clusters

four and six. HSMCs were not treated (UN), treated with IgE receptor cross-linking (IgECL) for

2 hours, IL-33 for 26 hours (priming for 24 hours prior to the addition of the other stimuli

treatment for 2 hours), IL-25 for 26 hours, TSLP for 26 hours alone (single) or in combinations

(combined) as indicated. (**A**) Venn diagram showing the overlap of upregulated differentially

expressed genes among the four comparisons of IL-33 vs UN, IL-25 vs UN, IgECL vs UN,

TSLP vs UN (left panel), and the comparisons of IgECL+ IL-33 vs IgECL, IgECL+ IL-25 vs

IgECL, IgECL+ TSLP vs IgECL (right panel). (**B**) K-means clustering analysis was performed

using R package pheatmap. The cluster size numbers indicate the number of genes that are

grouped in the cluster. RNA-seq data were generated from two biological samples.

Figure 4. Genes synergically upregulated by IL-33 priming and IgE receptor cross-linking are

enriched in genes encoding pro-inflammatory cytokines and chemokines. (**A**) Gene ontology

(GO) enrichment analysis of the upregulated genes in clusters four and six. *P*-values were

calculated by a one-side Fisher's exact test with the adjustment of Benjamini-Hochberg method.

(**B**) Heatmap representation of top ranked genes in each gene family in clusters four and six. The

numbers indicate RNA reads (FPKM). $\log_2$FC: $\log_2[E_{\text{IgECL+IL-33}}/(E_{\text{IgECL}}+E_{\text{IL-33}})]$. Data represents

the average reads generated from two biological samples.

Figure 5. Validation of gene expression promoted by the combined IL-33 priming and IgE

receptor cross-linking. HSMCs were not stimulated (UN) or stimulated with IgE receptor cross-

linking (IgECL) for 2 hours, IL-33 for 26 hours (priming for 24 hours prior to the addition of the

other stimuli treatment for 2 hours), or in combination of IgECL and IL-33. (**A**) qPCR analysis

of the mRNA expression of activation-induced genes identified by RNA-seq. (**B**) The same

treatments as described under (A) were applied, supernatants were collected after 24 hours and

IL-1B and CXCL8 were quantified by ELISA. Statistical differences were analyzed using two-

way ANOVA. Data represents mean ± SEM of four biological samples. * $P < 0.05$, ** $P < 0.01$,

*** $P < 0.001$.

Figure 6. IL-33 priming and IgECL increase chromatin accessibility in the synergy target genes.

(**A**) The numbers of upregulated ATAC-seq narrow peaks in the combined IL-33 priming and

antigenic stimulation relative to IgECL or IL-33 stimulation were subtracted using Diffbind

analysis. ATAC-seq was performed on HSMCs under various conditions, IgECL, IL-33 or

IgECL plus IL-33. ATAC-seq narrow peaks (NPs) of *CXCL2* (**B**), *CXCL8* (**C**), *IL1B* (**D**) genes in

HSMCs treated with IgECL and IL-33 (E+33) are shown (blue bars). Differential NPs generated

using DiffBind comparing IgECL+IL-33 with IgECL (E+33/E NPs) or comparing IgECL+IL-33

with IL-33 (E+33/33 NPs) are also shown (blue bars). Candidate enhancers are marked with red

bars and E - or + numbers, indicating distances (kb) from the TSS of the indicated genes. RPM,

reads per million nucleotides mapped. The IGV tracks are generated from one biological sample,

representing two biological replicates with similar patterns.

Figure 7. The AP-1, NF-κB, RAP1, and GABPA binding motifs are enriched in the regions with

increased chromatin accessibility in the IL-33 priming and IgE receptor cross-linking-

upregulated genes. Data represents two biological samples.

Figure 1

Figure 2

A IgECL

Gene	un		E	Log₂FC(E/un)
IL31	0.3		78.6	8.2
HBEGF	0.3		15.7	5.6
IL3	0.8		29.7	5.3
INHBA	21.0		751.4	5.2
ADCYAP1	2.5		83.2	5.1
FGL2	13.8		417.8	4.9
HES4	6.5		189.2	4.9
RGCC	14.1		396.3	4.8
LINC01010	0.9		24.1	4.7
DHRS9	2.7		67.3	4.6
IL11	0.7		16.9	4.5
ADAMTS4	0.5		10.1	4.4
HIP1R	0.5		10.8	4.3
CCL3	34.4		615.3	4.2
CXCL8	12.7		211.5	4.1
SH2D2A	3.9		64.4	4.0
ANGPTL4	64.5		1029.4	4.0
LUCAT1	3.1		48.5	4.0
CCL3L1	56.1		862.3	3.9
C17orf107	3.1		47.6	3.9

B IL-33

Gene	un		IL-33	Log₂FC(IL-33/un)
EBI3	0.1		230.5	10.7
IL1B	0.3		53.8	7.5
MAOA	0.2		21.5	6.7
CDH13	0.3		27.5	6.4
CCL5	0.3		24.8	6.2
CCL26	1.8		79.4	5.4
IL7R	0.6		21.8	5.3
MMP9	0.7		24.8	5.1
C1QTNF1	1.0		21.7	4.5
TNFAIP6	5.4		83.2	4.0
PT3X	1.3		18.5	3.8
IL32	2.1		29.4	3.8
ACE	0.8		11.5	3.8
JAK3	2.2		25.7	3.5
C15ORF48	3.2		34.0	3.4
CHST1	2.5		24.2	3.3
NFKB2	21.4		203.8	3.3
ACDYAP1	3.0		27.4	3.2
BATF	62.3		543.3	3.1
C3	5.5		48.1	2.1

C IL-25

Gene	un		IL-25	Log₂FC(IL-25/un)
RMRP	27.5		106.6	2.0
RF02198	11.3		29.0	1.4
RF00100	55.1		128.0	1.2
RPS17P1	5.5		12.3	1.1
KCNN4	5.0		10.8	1.0
HMMP1.HBD	6.1		11.0	0.9
MTND3P25	56.0		101.5	0.9
CD14	31.2		55.9	0.8
IGFBP6	9.5		16.9	0.8
RCC1A73A	11.4		20.0	0.8
NMB	18.1		28.5	0.7
CDH12	7.2		11.2	0.6
CHCHD6	18.5		28.4	0.6
BOD1	13.1		20.0	0.6
MEDAG	10.7		16.1	0.6
TEF	7.9		11.8	0.6
CCN5	27.4		40.3	0.5
HS3ST2	39.1		57.0	0.6
RPL13P12	82.5		117.8	0.5
SFRP2	9.7		13.8	0.5

D TSLP

Gene	un		T	Log₂FC(T/un)
IL1B	0.3		3.7	3.6
MRPS31P5	3.2		18.4	2.5
KCNN4	5.0		13.0	1.4
CXCL8	15.6		38.0	1.3
DLEU1	13.0		24.0	0.9
CACNA1H	10.6		18.8	0.8
RF02196	40.3		70.9	0.8
FO705657.1	6.3		11.1	0.8
ARHGAP33	12.6		21.8	0.8
MS4A6A	14.8		25.4	0.8
RECQL5	7.2		12.4	0.8
IGFBP6	9.5		16.2	0.8
FAM98B	9.0		15.2	0.7
THOC3	14.0		22.8	0.7
NDUFB1P1	12.7		20.7	0.7
MTND3P25	56.2		91.3	0.7
KANSL1.AS1	8.6		13.8	0.7
PDE6B	8.6		13.8	0.7
CBX3P9	6.4		10.1	0.6
MTCO2P2	157.6		264.2	0.6

Intensity

A

Single

IL-33 VS UN   IL-25 VS UN

204
(28.5%)   4
(0.6%)

42
(5.9%)   1
(0.1%)   1
(0.1%)

448
(62.6%)   0
(0%)   1
(0.1%)   2
(0.3%)

0
(0%)   2
(0.3%)

2
(0.3%)

0
(0%)   6
(0.8%)

3
(0.4%)

IgECL VS UN   TSLP VS UN

Combined

IgECL+ IL-33 VS IgECL   IgECL+ IL-25 VS IgECL

242
(79.9%)   19
(6.3%)   15
(5%)

10
(3.3%)

6
(2%)   6
(2%)

5
(1.7%)

IgECL+ IL-25 VS IgECL

B

Cluster 2 Size 2441
Cluster 5 Size 1202
Cluster 1 Size 618
Cluster 4 Size 902
Cluster 3 Size 1138
Cluster 6 Size 641

2
1
0
-1
-2

IL-33
IL-33+IL-25+TSLP
UN
IL-25
TSLP
IgECL+IL-33
IgECL+IL-33
+IL-25+TSLP
IgECL
IgECL+IL-25
IgECL+TSLP

Figure 4

A

Go analysis

Cluster four

Category	Gene counts (bar)	P-value
Cytokines		1.00E-05
Receptors and receptor binding		7.90E-02
Signaling molecules		9.70E-06
Transcription factors		1.80E-02
NF-κB signaling pathway		6.50E-02
Chemokines		1.10E-05
Cell death		2.60E-02

Cluster six

Category	Gene counts (bar)	P-value
Cytokines		2.50E-05
Receptors and receptor binding		3.60E-02
Signaling molecules		7.30E-03
Transcription factors		2.60E-03
NF-κB signaling pathway		4.20E-02
Chemokines		9.70E-02
Cell death		1.70E-02

B

Representative genes

Cluster four

Cytokines and chemokines

Gene	UN	IL33	IgECL	E+33	Log₂FC
CXCL2	0.3	1	1.7	144	5.7
CXCL3	0.4	0.3	2.8	155.9	5.7
CXCL8	15.6	6	185.1	5533.2	4.9
IL23A	0.2	0.3	0.6	10.4	3.5
IL33	0.9	0.5	1.6	10.9	2.4
CCL1	36.9	106	199.6	1173.5	1.9
CCL20	0	0.9	2.6	12.2	1.8
TNFSF4	5.4	2.2	12.7	48.4	1.7
IL6	10.9	17.7	54.7	229	1.7
IL13	1.3	4.4	6.6	21.7	1.0

Receptors and receptor binding

Gene	UN	IL33	IgECL	E+33	Log₂FC
CD200	0.1	0.3	8.9	19	1.0
CD274	17.5	13	36.5	89.9	0.9
HEG1	5.6	7.1	7.6	26.2	0.8
DUSP1	9.6	6.5	19.8	46.7	0.8
LRP12	9.8	10.1	13.5	31.5	0.4

Signaling molecules

Gene	UN	IL33	IgECL	E+33	Log₂FC
RGS16	3.1	1.9	7.2	25.8	1.5
GADD45A	12.4	17.8	54.1	147.1	1.0
RGS2	126.8	63.6	288.6	596.1	0.8
VCAN	2.9	2.6	4	10.3	0.6
LAPTM4B	12.2	6.5	8.5	17.4	0.2

Transcription factors

Gene	UN	IL33	IgECL	E+33	Log₂FC
EGR4	0.4	0.1	4.1	14.3	1.8
CREB5	1.3	0.4	5.9	19.8	1.7
ZFP36L2	82.4	43.3	154	461.2	1.2
MLF1	2	1.7	6.6	18.6	1.2
RFX2	2.6	8	21.1	52.3	0.8
CEBPD	4.1	5.9	13	32.7	0.8

NF-κB signaling pathway

Gene	UN	IL33	IgECL	E+33	Log₂FC
REL	10.1	8.7	22.7	67.4	1.1
TNFRSF21	5.5	4	6.7	20.3	0.9
RIPK2	17.8	22.9	27.7	92	0.9
MAP3K8	12.9	19.3	67	146.1	0.8
GBP2	38.4	27.2	82.5	179.3	0.7
NFKB1	20.1	38.6	40.1	112.2	0.5

Cell death

Gene	UN	IL33	IgECL	E+33	Log₂FC
BCL2A1	5.6	4.7	41.9	94	1.0
ACKR3	2.3	8.8	9.5	29.1	0.7
BCL6	6	5.7	7.2	17.8	0.5

Cluster six

Cytokines and chemokines

Gene	UN	IL33	IgECL	E+33	Log₂FC
CSF3	0.2	1.2	0.9	17.4	3.1
CXCL1	0.6	1.2	1.1	17.4	2.9
IL1B	0.3	53.8	0.6	367.7	2.8
IL5	0.2	6.5	0.4	35.9	2.4
IL1A	1.3	3.9	3.9	29.6	1.9

Receptors and receptor binding

Gene	UN	IL33	IgECL	E+33	Log₂FC
MFSD2A	0.8	7.4	3.2	33.3	1.7
IL7R	0.6	21.8	1.5	48.3	1.1
FCGR2A	10	27.9	7.3	57.5	0.7
TNFRSF9	0.4	9.9	2.9	20.9	0.7
HLA-E	343.4	460.3	434.2	991.9	0.1

Signaling molecules

Gene	UN	IL33	IgECL	E+33	Log₂FC
ITK	8.3	8.2	7	23.9	0.7
MTSS1	14.6	42	28.8	96.3	0.4
FSTL1	12.2	10.6	10.1	21.7	0.1

Transcription factors

Gene	UN	IL33	IgECL	E+33	Log₂FC
SKIL	5.6	14.3	5.3	32.5	0.7
TWIST2	9.6	16.8	14.9	36	0.2
TWIST1	11.3	23.9	21.5	49.5	0.1
YBX3	14.6	20.5	18.3	41.6	0.1

NF-κB signaling pathway

Gene	UN	IL33	IgECL	E+33	Log₂FC
TNFAIP3	3.7	28.9	11.6	65.1	0.7
TNFAIP2	4.1	8.3	6.7	23	0.6
NFKBIZ	48.4	76.7	68.2	201	0.5
NFKBIA	173.3	283.5	173	575.5	0.3

Cell death

Gene	UN	IL33	IgECL	E+33	Log₂FC
NINJ1	21	59.5	33.4	141	0.6
TNFAIP8	48	43.5	30.3	93.3	0.3
G0S2	14.2	42.4	39.9	95.4	0.2

Intensity

1	25	50	500	1000

Figure 5

A

B

Figure 6

Figure 7

TF	Motif	E-value	TF	Motif	E-value
AP-1		1.8e-022	NF-κB		8.5e-022
RAP1		4.1e-006	GABPA		2.0e-003