Cooperative *cis*-interactions between ectodomains of TCRαβ CD3 subunits enable mechanotransduction

Peiwen Cong[1,2†], Aswin Natarajan[3†], Zhou Yuan[1,2,4,a†], Chenghao Ge[1,2,b†], Stefano Travaglino[1,2], Saikiran Beesam[3], Xiang Li[3], Danielle Grazette[3], Michelle Krogsgaard[3‡], Cheng Zhu[1,2,4‡]

[1]Wallace H. Coulter Department of Biomedical Engineering, [2]Petit Institute for Bioengineering and Bioscience, Georgia Institute of Technology, Atlanta, GA, USA; [3]Laura and Isaac Perlmutter Cancer Center and Department of Pathology, New York University Grossman School of Medicine, New York, NY, USA; [4]Georgia W. Woodruff School of Mechanical Engineering, Georgia Institute of Technology, Atlanta, GA, USA

[†]These authors contributed equally.

[‡]Corresponding authors: Michelle.Krogsgaard@nyulangone.org and cheng.zhu@bme.gatech.edu

[a]Present address: Alfred E. Mann Department of Biomedical Engineering, University of

Southern California, Los Angeles, CA 90007, USA

[b]Present address: Amgen Inc., One Amgen Center Dr., Thousand Oaks, CA, USA

ABSTRACT (220 words)

TCR signaling poses a mechanical problem: pMHC binding occurs at the TCRαβ ectodomain

(ECD) head, whereas ITAM phosphorylation occurs on CD3 cytoplasmic tails. Chemistry cannot

bridge this >10 nm gap, requiring the two events to be coupled through the TCR–CD3 interface,

thus involving conformational allostery and being regulatable by force. Although weak ECD *cis*-

interactions between TCRαβ and CD3 have been proposed to contribute to this coupling, their

kinetics and mechanical competence remain elusive. Here, we quantify TCRαβ–CD3 ECD *cis*-

interactions in a pseudo-*cis* configuration using two-dimensional binding and single-bond force

spectroscopy, finding that TCRαβ–CD3γε and TCRαβ–CD3δε interactions have low affinity and

rapid kinetics, yet form catch bonds. Critically, concurrent engagement of CD3γε and CD3δε

produces high *cis*-cooperativity, yielding a stronger and longer-lasting CD3γε–TCRαβ–CD3δε

trimolecular catch bond than the sum of the two dimeric bonds, with force-stabilized lifetimes

matching those of agonist TCR–pMHC *trans*-interaction. Molecular dynamics simulations reveal

an expanded, cooperative, and asymmetric contact network, making CD3δε more force-responsive

and susceptible to conformational change than CD3γε. Interface mutations do not alter force-free

affinities but remodel cooperative *cis*-bond profiles, leading to an inverse correlation with *trans*-

bond profiles and T cell signaling. These results identify cooperative ECD *cis*-interaction as a

mechanically regulatable allosteric coupling element at the TCR–CD3 junction important to

antigen recognition and signal initiation.

INTRODUCTION (main text 8207 words)

Antigen recognition by the T-cell receptor (TCR) results in T cell activation, leading to proliferation, differentiation, and effector functions (*1*). Signaling is initiated by ligation of peptide-major histocompatibility complex (pMHC) with TCR at its extracellular ligand-binding site, where the antigen information is received, then transmitted across the membrane along the octameric TCR–CD3 complex (TCRαβ associated with CD3γε, CD3δε, and CD3ζζ) to the CD3 cytoplasmic tails (*2*). Concurrently, coreceptor CD8 or CD4 places the Src family kinase Lck close to the exposed CD3 tails to phosphorylate their immunoreceptor tyrosine-based activation motifs (ITAMs) (*3*). Phosphorylated ITAMs recruit Zap70, starting downstream signaling cascades (*1, 4*). Despite our detailed knowledge of the structural components of the TCR signaling machinery (*5, 6*), the mechanism of how pMHC binding at the TCRαβ head leads to phosphorylation of CD3 ITAMs, two events separated by ~15 nm in space, precluding the action of local chemistry, remains unclear.

Two of the TCR triggering models—conformational change and mechanosensor models (*7-9*)—share a common requirement: a physical coupling pathway must link ligand binding to ITAM phosphorylation. Structurally, this pathway must traverse the interface between TCRαβ and CD3s that includes ectodomains (ECDs), connecting peptides (CP), transmembrane (TM) bundle, and cytoplasmic tails. Yet identifying this coupling has been challenging because the TCR is subject to two seemingly opposing design constraints. On one hand, the complex must be sufficiently connected to maintain quaternary integrity and transmit information. On the other hand, it must remain sufficiently dynamic to permit state transitions that gate signaling. Resolving how molecular architecture balances connectivity with mobility is a central obstacle to a unified physical mechanism of TCR triggering.

A likely candidate for this coupling is *cis*-interaction (between proteins anchored to the same cell

surface) between the TCRαβ constant domains and the CD3 ECDs. Multiple lines of evidence

support this notion: the functional sidedness of CD3s (*10*) and the importance of the TCRαβ

constant domains (Cα/Cβ) (*11, 12*) indicate that TCR–CD3 *cis*-interactions are crucial for signal

transmission. Disrupting these interactions with anti-CD3 antibodies can change how T cells are

activated (*13-15*). The membrane-proximal ECD "ring" may act as the starting point for signal

propagation and influences downstream elements such as the CPs (*16*), TM bundle (*17, 18*), and

cytosolic tails (*2, 19*). Mutations affecting these *cis*-interactions can impact TCR assembly stability

(*20, 21*), surface expression (*22*), down-modulation (*23*), and cellular responses (*10, 21, 24-27*).

However, direct mechanistic interpretation has been limited by two gaps. First, ECD interactions

appear very weak and transient, raising doubts about their ability to support sustained signal

transmission. Second, the subtle coupling may reside in force-dependent dynamic allostery that is

sensitive to experimental perturbations, rather than in easily-captured large, stable structural

changes, as suggested by discrepant results from different experiments: Early cryo-electron

microscopy (cryo-EM) structures (*28-31*) and crosslinking experiments have shown minimal

stable rearrangements upon ligation (*27*), whereas nuclear magnetic resonance (NMR) (*32, 33*)

and molecular dynamics (MD) (*33, 34*) studies indicate dynamic allostery in the TCR constant

domains and at the CD3 interfaces upon pMHC binding.

More recent cryo-EM structures of the TCR–CD3 complex reconstituted in nanodiscs suggest

unliganded TCR–CD3 adopts "compact/closed" conformations (*35*). Upon pMHC engagement the

complex shifts to an "extended/open" architecture, and restricting ECD opening can impair

activation (*35*). These findings support the plausibility of activation-linked allosteric transitions,

but they do not identify the mechanical mechanism that stabilizes and coordinates this closed-to-

open transition across the TCR–CD3 junction, nor do they establish whether the implicated ECD

interfaces are mechanically competent to support TCR–pMHC *trans*-interaction in the presence of

force (*1, 6, 18, 36-39*).

Mechanical forces have been suggested to induce conformational allostery (*40-42*), regulate

lifetimes of TCR–pMHC *trans*-bonds (*40, 41, 43-46*), and trigger signaling (*43, 47-49*), which, in

turn, modulate effector functions (*41, 43, 45, 50, 51*). T cells exert endogenous forces on the TCR–

CD3 via engaged pMHC (*52-55*), which must be mediated, at least in part, by TCR–CD3 ECD

cis-interactions, therefore allowing force to play a role. Thus, mechanical force naturally fits the

allosteric framework, likely as a regulator rather than an alternative to allostery. T cells exert forces

on engaged ligands, and mechanical load can bias conformational ensembles by stabilizing specific

bound states and pathways. In this view, force is a control parameter that can reveal latent allosteric

couplings that are otherwise too fleeting to detect at equilibrium. If ECD *cis*-interactions

participate in signal transmission, they must satisfy a stringent mechanical criterion: they must

remain connected long enough under physiological loads to couple ligand engagement to

downstream phosphorylation, while permitting the local rearrangements and fluctuations required

for gating. This criterion turns our inquiry into a quantitative question—what are the kinetic and

force-dependent properties of the TCRαβ–CD3 ECD *cis*-bonds, and can cooperative assembly

transform weak pairwise contacts into a durable, force-stabilized coupling element?

Addressing this question has been difficult because TCRαβ–CD3 ECD *cis*-interactions are too

weak and transient to be measured by three-dimensional (3D) methods (*24, 56-58*) such as SPR

(Supplementary Fig. 1A, B). Moreover, most prior characterizations have been performed without

controlling mechanical load, obscuring whether force reshapes the interaction landscape in ways

relevant to mechanotransduction. Thus, despite longstanding hypotheses that ECD *cis*-interactions

link antigen recognition to CD3 signaling, direct measurements of their 2D kinetics, cooperativity,

and force response have been lacking.

Here we combine ultrasensitive biophysical measurements with molecular dynamics (MD)

simulations to define the kinetic and mechanical properties of TCRαβ–CD3 ECD *cis*-interactions.

Using micropipette adhesion frequency (MAF) assay (*59, 60*) and biomembrane force probe (BFP)

thermal fluctuation and force-clamp assays (*43, 61*), we analyzed two-dimensional (2D) binding

kinetics of TCRαβ interactions with either CD3γε, CD3δε, or both in a pseudo-*cis* configuration.

We find that the bimolecular interactions are low-affinity and short-lived yet form catch bonds,

indicating the presence of force-favored bound states. Crucially, co-engagement of CD3γε and

CD3δε produces strong synergy in both bond formation and lifetime, resulting in a long-lived

cooperative CD3γε–TCRαβ–CD3δε trimolecular catch bond. Complementary analyses at the

atomic level using conventional MD (CMD) and steered MD (SMD) simulations corroborate

cooperative stabilization through an expanded contact network and reveal asymmetric force

responses between CD3γε and CD3δε. Finally, targeted mutations at the trimeric interface remodel

force-dependent *cis*-bond profiles without altering force-free affinity, allowing us to relate

mechanical coupling at the TCR–CD3 junction to T cell signaling. Together, these results establish

cooperative ECD *cis*-interaction as a mechanically regulatable allosteric coupling element that can

stabilize and shape information transmission across the TCR–CD3 interface.

RESULTS

2D kinetic measurements reveal weak TCRαβ–CD3 *cis*-interactions.

To directly measure physical interactions among TCRαβ, CD3γε, and CD3δε (ECDs; hereinafter

omitted for simplicity), we utilized the ultra-sensitive MAF assay (*59, 60*) and BFP thermal

fluctuation assay (*60, 62*). To compensate for the anticipated low affinities, we coated high

densities of hybrid 2B4 TCRαβ (mouse 2B4 TCRαβ variable domains fused with human LC13

TCRαβ constant domains) (*26, 63*) and human CD3γε or CD3δε onto opposing surfaces (Fig. 1A,

B). To mimic *cis*-interactions, N-terminal biotinylated TCRαβ was coated on one side (*left*), while

C-terminal biotinylated CD3s were coated on the opposing side (*right*), allowing head-to-tail

alignment for "pseudo-*cis*" contacts. In the MAF assay (Fig. 1A), one red blood cell (RBC) was

driven to cyclically contact the other RBC with a consistent contact area (A_c, in μm^2) and time (t_c,

in s) and separated to detect binding events from visually observed RBC elongations. An adhesion

frequency, P_a (*i.e.*, number of binding events divided by total touches) enumerated from a sequence

of repeated contact cycles was measured. Assuming second-order forward and first-order reverse

reaction between monomeric TCRαβ and CD3γε or CD3δε, we previously showed that $P_a = 1 -$

$\exp(-\langle n \rangle)$, where $\langle n \rangle = m_r m_l A_c K_a [1 - \exp(-k_{off} t_c)]$ is the average number of bonds per

contact. Here, m_r and m_l (in μm^{-2}) are the respective densities of TCRαβ and CD3γε or CD3δε,

K_a (in μm^2) is 2D affinity, and k_{off} (in s^{-1}) is off-rate (*59, 60*). When $k_{off} t_c \gg 1$, $\exp(-k_{off} t_c) \approx 0$,

the effective 2D affinity $A_c K_a$ (in μm^4) can be solved explicitly as $A_c K_a = -\ln(1 - P_a)/m_r m_l$.

The effective affinities of TCRαβ–CD3γε and TCRαβ–CD3δε *cis*-interactions were similar (3.78

and 2.78 × 10^{-6} μm^4, respectively), which are reliable because they are about an order of magnitude

above the detection limit of our current MAF setups (Fig. 1C, *orange* and *blue*) (*46*). For

comparison, we measured the 2B4 TCR–K5:IE^k *trans*-interaction side-by-side using the C-

terminal biotinylated pMHC-coated RBC to test against the intact TCR–CD3 complex expressed

on hybridoma cells. The resulting effective 2D affinity was tens to hundreds of folds higher (~

5.87×10^{-4} μm^4, Fig. 1C, *gray*) even though the hybridoma cell surface is rough and the TCRαβ-

bearing RBC is smooth, which could result in one to two orders of magnitude lower effective 2D

affinity for the rough than smooth cells (*64*). In addition, using C-terminal biotinylated CD3γε and

CD3δε coated on the opposing surfaces, we quantified the CD3γε–CD3δε heterotypic interaction

(~ 1.10×10^{-5} μm^4, Fig. 1C, *green*), which was higher than the TCR–CD3 heterotypic interactions.

Similarly, the effective 2D affinities for the CD3γε–CD3γε, CD3δε–CD3δε, and TCRαβ–TCRαβ

homotypic interactions were measured with lower values or undetectable (Supplementary Fig. 1C).

To further determine the on- and off-rate, we used the BFP thermal fluctuation assay to measure

off-rate, k_{off} (*60, 62*). We attached a functionalized glass bead to the apex of the micropipette-

aspirated ultra-soft hypotonic RBC to form the probe on the left (Fig. 1B), whose thermally driven

fluctuations were monitored in real time with high spatial-temporal precision (few nanometers and

sub-milliseconds) (*65*). The target bead coated with the corresponding binding partner was brought

close to the probe to allow molecular contacts via thermal fluctuations, facilitating bond formation.

Upon binding, the receptor–ligand (*i.e.*, either TCRαβ *vs.* CD3, CD3 *vs.* CD3, or pMHC *vs.* TCR–

CD3) bond acted as an additional spring in parallel with the RBC spring, dampening the confined

Brownian motions (*60, 62*). The sudden reduction in thermal fluctuations indicated bond formation,

while the restoration to the original fluctuation amplitude marked bond dissociation. The durations

in between represent bond lifetimes, which were exponentially distributed, and the reciprocal of

the mean bond lifetime is equal to the k_{off}. The 2D off-rates for TCRαβ–CD3γε, TCRαβ–CD3δε,

and CD3γε–CD3δε *cis*-interactions measured in this way were very rapid (> $10 \, \text{s}^{-1}$) compared to

the slower k_{off} of the 2B4 TCR–K5:IE^k *trans*-interaction plotted alongside in Fig. 1D. These

results also validated the ~ 2 s contact time used in our MAF assay, which was sufficiently long to

ensure the measured adhesion frequencies had reached steady-state. Finally, the effective 2D on-

rates, $A_c k_{on}$, were calculated as $A_c k_{on} = A_c K_a \times k_{off}$ for the corresponding interactions (Fig. 1E).

Together, the very low affinities and fast off-rates explain why SPR failed to detect these weak and

rapid TCRαβ–CD3 interactions, underscoring the ultra-sensitivity of our 2D assays (*66*).

Pairwise *cis*-interactions among TCR–CD3 ECDs form bimolecular catch bonds.

We (*41-44, 46, 50*) and others (*40, 45, 51, 67, 68*) have shown that mechanical force modulates

TCR–pMHC *trans*-interactions and amplifies antigen discrimination (*40, 43, 44*). To test whether

force also modulates dissociation of the TCR–CD3 *cis*-interactions, we used the BFP force-clamp

assay (*41, 43-45*) to measure the force-dependent lifetimes of TCRαβ–CD3γε, TCRαβ–CD3δε,

and CD3γε–CD3δε *cis*-bonds (Fig. 1B). Like in the thermal fluctuation assay, we also used bead-

mounted RBCs as ultra-sensitive force transducers to measure molecular interactions in the force-

clamp assay. Unlike the thermal fluctuation assay, which positioned the target bead at the null

position for bond formation and dissociation at zero force, the force-clamp assay retracted the

target by a pre-set distance to exert a constant tensile force onto the bond. To enhance stability of

force measurement, a dual-edge tracking system was implemented to monitor the positions of both

the BFP micropipette and the probe, allowing us to subtract the micropipette drifts from the force

signals. The target was programmed to move in repeated "approach-touch-retract-hold-return"

cycles, generating a force *vs.* time trace (Fig. 1F) that exemplified both non-binding and binding

events. The compressive (negative) force indicates bead-target contact during which bond

formation can occur. Upon retraction, the compressive force either returns to zero (no binding) or

converts to tensile (positive) force (binding). In the latter case, retraction was halted at the

designated force level until bond dissociation, indicated by the sudden return of force to zero.

Interestingly, BFP measurements revealed that all three *cis*-interactions formed catch bonds, with

mean lifetimes prolonged by low forces (< 7-17 pN, depending on the specific interaction) until

being overpowered by higher forces that turned the bond profiles into slip bonds (*i.e.*, mean bond

lifetimes shortened by forces, Fig. 1G). These findings demonstrate that TCR–CD3 *cis*-interactions

can bear durable force, supporting the notion that information on antigen recognition can be

transmitted mechanically through TCRαβ–CD3 interactions.

Notably, these TCRαβ–CD3 *cis*-interactions exhibited shorter bond lifetimes across the entire

force range than our previously reported TCRαβ–pMHC *trans*-bonds here (replotted from (*42*) in

Fig. 1G, *gray*), suggesting that individual TCRαβ–CD3 bimolecular *cis*-interactions alone may not

persist long enough to fulfill the proposed mechanical transmission requirements. The 1:1:1

stoichiometry of TCRαβ:CD3γε:CD3δε revealed by the cryo-EM structures (*28, 29, 31*) and our

direct kinetic measurements of CD3γε–CD3δε interactions (Fig. 1C-E, G, *green*) prompted us to

ask: Does TCRαβ bind CD3γε and CD3δε concurrently to form stronger bonds? If so, is this

concurrent binding independent or cooperative? If cooperative, how much is the bond stability

enhanced, and does this enhancement meet the mechanical transmission requirements?

CD3δε and CD3γε interact with TCRαβ cooperatively in bond formation and bond lifetime.

To test whether CD3δε and CD3γε interact with TCRαβ cooperatively in *cis*, we applied affinity-

and lifetime-based measurements with MAF and BFP while co-presenting both CD3s. The results

were compared against predictions from a non-cooperative binding model. Significant deviations

from these predictions are taken as evidence for positive *cis*-cooperativity.

To ensure rigorous comparison of adhesion frequencies, a single TCRαβ-bearing RBC was

sequentially brought into contact with a RBC coated with CD3γε, CD3δε, or an equimolar mixture

of both (denoted CD3m) in randomized order (Fig. 2A). Streptavidinylated RBCs were prepared

from the same batch and coated using excess amounts of biotinylated CD3s to ensure saturated

mounting to SA with equivalent density. Adhesion frequencies were measured over 30 touches per

pairing for TCRαβ–CD3γε, TCRαβ–CD3δε, and TCRαβ–CD3m (Fig. 2B, connected by *solid lines*)

before switching to a new TCRαβ-bearing RBCs (specificity checks of all interactions are shown

in Supplementary Fig. 1D). Across eight TCRαβ RBCs tested, the adhesion frequency to CD3m

was consistently higher than either subunit alone. To interpret this result, we transformed P_a into

the average number of bonds per contact, $\langle n \rangle = -\ln(1 - P_a)$, which is additive for independent,

concurrent binding (*69, 70*). Under this non-cooperative assumption, the predicted bond number

for CDm is $\langle \hat{n}_m \rangle = 1/2 \langle n_{\gamma\epsilon} \rangle + 1/2 \langle n_{\delta\epsilon} \rangle$, where $\langle n_{\gamma\epsilon} \rangle$ and $\langle n_{\delta\epsilon} \rangle$ were measured from respective

bimolecular TCRαβ interactions with the single CD3 species indicated by the subscripts (Fig. 2B,

horizontal dotted line). Strikingly, the observed $\langle n_m \rangle$ significantly exceeded the predicted $\langle \hat{n}_m \rangle$,

falsifying the independent binding hypothesis and demonstrating cooperative binding. The degree

of cooperativity can be calculated as $\langle n_m \rangle / \langle \hat{n}_m \rangle - 1$, yielding a value of 170%, substantially

higher than zero (Fig. 2C), confirming pronounced positive *cis*-cooperativity.

Next, we used BFP thermal fluctuation and force-clamp assays to measure the average lifetimes

of TCRαβ–CD3m bonds (denoted as τ_m) in the absence and presence of force. The BFP target

beads were functionalized similarly to the target RBCs in the MAF assay, except that the surface

density of TCRαβ was adjusted to maintain infrequent adhesion events (≤ 20%), ensuring a high

probability (~ 90%) of single specific bonds (*59*) (Supplementary Fig. 1E-H). The lifetime

scattergram was globally fitted by the previously described two-state, two-pathway dissociation

models (Supplementary Fig. 2A-I) (*46*). TCRαβ–CD3m bonds persisted significantly longer than

single-ligand TCRαβ–CD3γε or TCRαβ–CD3δε bonds across all force levels tested (Fig. 2D;

single-ligand curves replotted from Fig. 1G). The τ_m *vs.* f curve peaked at ~ 13 pN with a lifetime

of ~ 1.5 s, whereas the $\tau_{\gamma\epsilon}$ and $\tau_{\delta\epsilon}$ *vs.* f curves peaked at ~ 7 and 15 pN, respectively, with lower

values of ~ 0.6 s for both. Notably, the τ_{m} bond profile exhibited a more pronounced catching

behaviour, closely matching that of the TCRαβ–pMHC *trans*-interaction (Fig. 1G, *gray*; peak at ~

10 pN with a lifetime ~1.6 s). These results indicate that dual ligands are able to provide sufficient

cis-stability for mechanotransmission, whereas single ligands are not.

To investigate the basis of this prolonged dual-ligand lifetime, we performed survival analysis on

bond lifetime distributions (*60, 62, 65*). The survival probability curve of zero-force lifetimes for

TCRαβ–CD3m displayed two distinct decay phases, in stark contrast to the log-linear decay

observed for single-ligand *cis*-interactions (Fig. 2E). The presence of a slow-decaying

subpopulation confirms the existence of the long-lived bond states, bond species, or both.

Similar subpopulations of longer-lasting bonds in dual-ligand were also observed in the

survival curves of low- and high-force lifetimes (Supplementary Fig. 2J, K). These longer-lived

events likely represent trimolecular CD3δε–TCRαβ–CD3γε bonds that substantially prolong τ_{m},

and result in positive cooperativity. To quantify this cooperativity in terms of bond lifetime, we

calculated the average lifetime of TCRαβ–CD3m bonds as the weighted sum of the bond lifetimes

of two bimolecular interactions expected from no cooperativity, $\tau_{\gamma\epsilon}$ and $\tau_{\gamma\epsilon}$, by $\hat{\tau}_{\mathrm{m}} = P_{\gamma\epsilon} \times \tau_{\gamma\epsilon} +$

$P_{\delta\epsilon} \times \tau_{\delta\epsilon}$ (Fig. 2D, *dotted curve*), where the bond formation possibilities, $P_{\gamma\epsilon}$ and $P_{\delta\epsilon}$, equal the

respective fractions of TCRαβ–CD3γε and TCRαβ–CD3δε bond numbers, $\langle n_{\gamma\epsilon} \rangle / (\langle n_{\gamma\epsilon} \rangle + \langle n_{\delta\epsilon} \rangle)$

and $\langle n_{\delta\epsilon} \rangle / (\langle n_{\gamma\epsilon} \rangle + \langle n_{\delta\epsilon} \rangle)$. The assumption of independent binding predicts $\tau_{\mathrm{m}} = \hat{\tau}_{\mathrm{m}}$, but the

cooperative analysis, $\Delta\tau / \hat{\tau}_{\mathrm{m}} = (\tau_{\mathrm{m}} - \hat{\tau}_{\mathrm{m}}) / \hat{\tau}_{\mathrm{m}}$, revealed ~ 74% to 328% increments in the CD3γε–

TCRαβ–CD3δε trimolecular *cis*-bond lifetime over the force range of 0-30 pN (Fig. 2F),

highlighting the strong force-enhanced positive cooperativity.

MD simulations corroborate cooperative interaction of CD3γε and CD3δε with TCRαβ.

To corroborate the experimentally observed *cis*-cooperativity, we performed force-free conventional MD (CMD) simulations (*71*) on TCRαβ–CD3γε, TCRαβ–CD3δε, and CD3γε–TCRαβ–CD3δε complexes. The initial full-atomic model of TCR–CD3 ECDs, including the trimeric and dimeric structures (Fig. 3A and Supplementary Fig. 3A), were extracted from the cryo-EM structure (PDB: 6JXR) (*28*). All equilibrated complexes remained bound after 200 ns CMD simulations. Their binding interfaces were analyzed in terms of contact areas (Fig. 3B and Supplementary Fig. 3B) and binding free energies (Fig. 3C and Supplementary Fig. 3C), which were inversely correlated. Using cooperative analyses similar to Fig. 2C, F, we found these two measures also exhibited positive *cis*-cooperativities (Fig. 3D). Because binding affinity, K_a, is related to free energy change, ΔE by $K_a \propto \exp(-\Delta E / k_B T)$, where k_B is the Boltzmann constant and T is the absolute temperature, these simulations provide independent verification of cooperative 2D affinity measurements.

We next simulated force-induced dissociations of TCRαβ–CD3 complexes using steered MD (SMD) (*72*). A pulling force was applied on the TCRαβ N-terminus, while the C-termini of CD3γε, CD3δε or both were anchored (Fig. 3E, *insets* and Supplementary Fig. 4A). To complete simulation within accessible computational timescales, we used relatively high clamping forces (100-200 pN) to accelerate dissociations, which were detected as sudden increases in the minimum distances between TCRαβ and either or both CD3s (Fig. 3E, *curves* and Supplementary Fig. 4B). The times to dissociate ranked as TCRαβ–CD3δε < TCRαβ–CD3γε < CD3γε–TCRαβ–CD3δε. Simulated force effects on times to dissociation (Fig. 3F) were consistent with experimental lifetime trends, following expected slip behaviors beyond 15 pN (Fig. 2D). The trimolecular CD3γε–TCRαβ–CD3δε bonds dissociated at 626, 404, and 233 ns under 150, 175, and 200 pN,

respectively, which persisted at least an order of magnitude longer than the two dimeric bonds. To

quantify cooperativity, we calculated $\Delta t / \hat{t}_{\delta\gamma}$, which compares observed times to dissociation, $t_{\delta\gamma}$,

to the predicted times to dissociation for the non-cooperative trimolecular complex $\hat{t}_{\delta\gamma} = t_{\gamma\epsilon} +$

$t_{\delta\epsilon} - t_{\gamma\epsilon}t_{\delta\epsilon}/(t_{\gamma\epsilon} + t_{\delta\epsilon})$, calculated using the TCRαβ–CD3γε and TCRαβ–CD3δε dissociation

times, $t_{\gamma\epsilon}$ and $t_{\delta\epsilon}$, respectively. Observed dissociation times exceeded this prediction by from

1,200% at 150 pN to 1,900% at 200 pN, demonstrating strong *cis*-cooperativity and mechanical

reinforcement (Fig. 3G). Force-ramp SMD simulations on partial and full TCR–CD3 assembles

yielded rupture forces in the same order, TCRαβ–CD3δε < TCRαβ–CD3γε < CD3γε–TCRαβ–

CD3δε (Supplementary Fig. 4C, D), further confirming the higher stability of the trimolecular

complex. Together, complementary experimental and computational approaches demonstrate

positive *cis*-cooperativity between CD3δε and CD3γε in their co-binding to TCRαβ.

Structural basis of *cis*-cooperativity quantified by atomic-level contacts.

To elucidate the structural basis of *cis*-cooperativity, we analyzed interface contacts within the

TCRαβ–CD3γε, TCRαβ–CD3δε, and CD3γε–CD3δε dimers and compared them with those within

the CD3γε–TCRαβ–CD3δε trimer. The contact area sizes were preserved from the dimer to trimer

interfaces with only modest alteration upon incorporation of the third subunit. Because CD3γε and

CD3δε bind TCRαβ orthogonally with minimal overlap, their combined contact area in the trimer

was only slightly smaller than the sum of dimers. This slight steric loss was more than offset by

the gain of the CD3γε–CD3δε contact area in the trimer, which substantially increased the total

contact area and lowered the binding free energy (Fig. 3B, C), consistent with *cis*-cooperativity.

Similarly, we analyzed hydrogen (H) bonds within the TCRαβ–CD3γε, TCRαβ–CD3δε, and

CD3γε–CD3δε dimers and compared them with those within the CD3γε–TCRαβ–CD3δε trimer,

finding substantially more H-bonds in the trimer than in the dimers at the start and in the middle,

but not near the end of SMD simulations (Supplementary Fig. 4E). Similar cooperativity

calculations found > 100% and 180% more H-bonds in the trimer without and with force,

respectively (Supplementary Fig. 4F).

We further quantified the level of interfacial contact using dynamic distances from each residue to

its binding partner. We termed this quantification occupancy, defined as the fraction of simulation

time during which the center of mass of that residue remains within 4 Å of any atom of the *cis*-

interacting ECDs. As such, the occupancy of an individual residue ranks from 0% to 100% as

interaction strengthens.

To reveal dynamic binding footprints, we mapped these occupancies onto the molecular surfaces,

with color intensities proportional to occupancy percentages (Fig. 4A). To expose the respective

TCRαβ–CD3γε and TCRαβ–CD3δε interfaces, a pair of the same trimer with different views

formed the mirror images, where CD3γε (Fig. 4A, *left*) and CD3δε (Fig. 4A, *right*) were rotated

outward away from fixed TCRαβ. Differential color coding was used to indicated CMD-derived

occupancy levels on interfaces of CD3γε–TCRαβ–CD3δε trimer (*cyan*), TCRαβ–CD3γε dimer

(*magenta*, after CD3δε removal), and TCRαβ–CDδε dimer (*yellow*, after CD3γε removal),

respectively. Overlay of dual-color maps (*blueish* overlap for TCRαβ–CD3γε; *greenish* overlap for

TCRαβ–CD3δε) highlighted conserved interactions. These analyses revealed the shifts in residue

contacts from residing in the respective dimeric structures to the trimeric structure in the absence

of force. Interestingly, the interface of TCRαβ with CD3γε was nearly unchanged by the presence

of CD3δε (Fig. 4A, *left*, *magenta vs. cyan*), whereas that with CD3δε exhibited a marked

reorientation in the presence of CD3γε (Fig. 4A, *right*, *yellow vs. cyan*). This was consistently

observed in all three independent CMD runs.

To quantify occupancy footprint colocalization between dimer and trimer states, we calculated

Manders' overlap coefficients (*73*), finding significantly greater values for CD3γε than CD3δε (Fig.

4B). The higher Manders' coefficient for γε than δε revealed a higher level of overlapping contact

interfaces between trimer and dimer of TCRαβ with CD3γε than CD3δε, which negatively

correlated with the level of contact adjustments going from dimer to trimer after adding the missing

CD3. This indicates that, structurally, the TCRαβ–CD3γε interface remains unchanged with or

without CD3δε, whereas the TCRαβ–CD3δε interface must reshape to accommodate the

incorporation of CD3γε during trimeric assembly. Asymmetric TCRαβ interactions with two CD3s

may result in unequal contributions to *cis*-cooperativity.

Asymmetric responses of CD3ε under *cis*-cooperativity and mechanostranmission.

To further elucidate the asymmetric roles of CD3s during cooperative binding and force

transmission, we investigated the differential dynamic responses of CD3γε *vs.* CD3δε using

occupancy and fluctuation analyses. We first compared the occupancies determined using SMD

simulations in the presence of force with values determined in previous section using CMD

simulations for every single α, β, γ, ε, δ, and ε' chains (note the omissions of TCR and CD3

designations for simplicity). Hereinafter, we use ε' to denote the ε chain that forms the heterodimer

with the δ chain to distinguish it from the ε chain that forms the heterodimer with the γ chain,

despite their identical sequences. Notwithstanding the average occupancies of ε' and ε chains of

the trimer were similar in the absence of force, ε' had far fewer contact residues with αβ than ε

both in the absence and presence force, indicating asymmetry (Fig. 4C). Interestingly, under

pulling force applied to the TCRαβ, both the number of residues with non-zero values and the

domain average of ε' occupancies declined sharply, whereas those of ε remained stable (Fig. 4C).

These data suggest that ε' is more sensitive to force, whereas ε provides more mechanical support

for *cis*-cooperativity.

To expand the ε and ε' occupancy data (Fig. 4C), we similarly calculated mean occupancies of αβ

residues contacting γ and δ domains from CMD and SMD simulated γε–αβ–δε trimer. We also

performed parallel occupancy calculations using simulated αβ–δε and αβ–γε dimers for all CD3

chains. We then plotted these paired occupancies as points using values obtained from CMD

(without force) and SMD (with force) as *x*- and *y*-coordinates and drew vectors connecting any

two points for the same CD3 chains with the start- and end-point coordinates calculated from

simulated dimeric and trimeric structures, respectively (Fig. 4D). The magnitude of a vector

represents the amount of occupancy change for a single CD3 chain (γ, ε, δ, or ε') from before (as

αβ–γε or αβ–δε dimer) to after (as γε–αβ–δε trimer) incorporating the missing CD3 heterodimer

(δε or γε). Remarkably, the δ (*red*) and ε' (*blue*) vectors were of much larger magnitudes than the

γ (*green*) and ε (*orange*) vectors, revealing much larger cooperativity-induced occupancy changes

in the δ and ε' than the γ and ε chains. This quantitatively depicts the strength of larger non-

overlapping trimer and dimer footprints (*cyan areas*) of δε (Fig. 4A, *right*) than γε (Fig. 4A, *left*),

re-emphasizing that, to form a γε–αβ–δε trimer via cooperative *cis*-interaction among ECDs, it

requires more contact adaptation from incorporating γε into the αβ–δε dimer than incorporating δε

into the αβ–γε dimer.

The directions of these vectors indicate whether the changes are positive (increasing) or negative

(decreasing). Their *x*- and *y*-components represent the changes occurring in the absence and

presence of a 175 pN force, respectively. On Fig. 4D, points below (*light blue*) or above (*light red*)

the diagonal line indicate domain contacts of αβ with CD3 chains that are suppressed or enhanced

by force, respectively. Interestingly, δ switched from a force-suppressed to a slightly force-

enhanced state (moving from below to above the diagonal line), while ε' redirected from slightly

force-enhanced to force-suppressed (moving from above to below the diagonal line); by

comparison, hardly any changes of force-induced contacts, and neither switches of their force-

enhanced (γ) and force-suppressed (ε) states were seen in either γ or ε chain. (Fig. 4D). These

results reveal that δε is more force-sensitive and γε is more mechanically stable, indicating their

different roles in transmitting force across the interfaces between TCRαβ and the two CD3 ECDs

In addition to occupancy analysis, which measures contact stability, we analyzed root-mean-square

fluctuations (RMSF) in the absence and presence of a 175 pN clamp force, which reflect residue

flexibility. By averaging across the entire chain, we assessed the mobility changes of different

chains before and after applying force, revealing another layer of mechano-responsiveness for the

trimer. ε' showed a larger mean RMSF than ε (Fig. 4E), and force-enhanced RMSF in both ε' and

ε residues, but much more for ε' than ε (Fig. 4E). These results inverted the trends of the occupancy

data (Fig. 4C), showing a negative correlation expected intuitively. The differences in RMSF

values calculated in the presence and absence of force (ΔRMSF) were projected onto the TCR–

CD3 quaternary ECD structure, where force-enhanced (*blue*) and force-suppressed (*red*)

fluctuations were indicated by color, and absolute ΔRMSF values in response of force were

indicated by backbone thickness (Fig. 4F). Under pulling, δε exhibited more amplified dynamics

than γε, forming a hinge in δ (*red*) and promoting an out swinging of the semi-detached ε' (*blue*),

further revealing asymmetry between γε and δε.

Identifying residues critical to *cis*-cooperativity and their distinctive force-responses.

To pinpoint the structural determinants of *cis*-cooperativity, we developed a modified occupancy

analysis (*i.e.*, ΣOccupancy) that integrated occupancies from CMD and SMD simulations,

weighted by the number of connections from that residue to all residues in the binder. Briefly,

instead of residue-to-domain occupancy, we quantified residue-to-residue occupancy in a matrix

format (Supplementary Fig. 5A). Subsequently, we summed over all possible contacts with the

interacting domain (Supplementary Fig. 6, *gray*) to account for one-to-many interactions. This

approach broadened the dynamic range of detection and identified ~10 and 14% more contacting

residues in CMD and SMD, respectively, compared with cryo-EM static structure. We used

ΣOccupancy to map the high-contact residues at the TCR–CD3 interface, visualized by sidechain

color-coding on the structure, thereby highlighting candidate residues with strong interface

connectivity (Fig. 4G).

To assess force sensitivity, instead of integrating the CMD and SMD results, we calculated the

difference between the CMD- and SMD-derived occupancy matrix (Supplementary Fig. 5B).

Summing over by row or column yielded ΔOccupancy, in turn identified force-sensitive residues

whose contacts were strengthened (*red*) or weakened (*blue*) by pulling (Supplementary Fig. 6 *red*

and *blue*). We mapped these force-sensitive residues on the TCR–CD3 interface by color-coding

(Fig. 4H). From a top-down view, force preferentially reduced TCRαβ–CD3γε contacts, whereas

TCRαβ–CD3δε contacts were more dynamic. It seems reasonable to suspect that these force-

sensitive residues play key roles in mediating force transmission across the TCR–CD3 interface.

Mutating interfacial residues alters catch bond but not affinity of TCRαβ–CD3 interactions.

Among all *cis*-contact residues, two regions of the TCRβ constant domain (Cβ) stood out by their

strikingly opposite mechano-responses. Both the CC' loop (residues 182-187) and the Helix 4–F

strand segment (residues 221-226) were highly conserved and enriched in high ΣOccupancy

residues; however, the CC' loop showed negative ΔOccupancy and the Helix 4–F strand displayed

positive ΔOccupancy (Fig. 5A). This suggests that force applied from the TCRβ N-terminus to the

Cβ domain elicits residue-specific allostery with opposite effects in adjacent structural regions,

moving the CC' loop away from, and the Helix 4–F strand closer to, the CD3s.

We targeted the Cβ CC' loop and Helix 4–F strand regions, highlighted on the TCRαβ structure

(Fig. 5B), and introduced double-alanine mutations: G182A/K183A (GK), N222A/P223A (NP),

and N225A/H226A (NH), made to alter residues with both strong (large ΣOccupancy) and force-

sensitive (high absolute ΔOccupancy) contacts. As previously reported (26), mutant mouse 2B4

TCRαβ chains assembled normally with CD3 chains in transduced hybridoma cells, yet altered T

cell activation, measured by interleukin-2 (IL-2) secretion upon co-culture with K5:IE^k expressing

CHO cells (Fig. 5C, reformatted IL-2 dose-dependent response curves from Ref. (26)). Among

these, NP acted as a gain-of-function (GOF) mutation, whereas GK and NH behaved as loss-of-

function (LOF) mutations. The IL-2 alterations underscore the functional relevance of TCRαβ–

CD3 cis-interaction in effectively relaying the ligation signals for downstream T cell activation.

To scrutinize how cis-interactions regulate signaling propagation, we measured cis-interactions

between WT and mutant (MT) N-terminally biotinylated 2B4 TCRαβ with C-terminally

biotinylated CD3γε, CD3δε, or both. To ensure the 1:1 ratio of CD3γε and CD3δε constructs, we

engineered flexible amino acid linkers to covalently tether the two CD3 heterodimers, thereby

avoiding heterogeneous CD3 mixtures (i.e., in vitro CD3γε–CD3δε, see Methods). Comparing

MTs to WT TCRαβ, statistically indistinguishable normalized numbers of bonds were observed

for bimolecular interactions with CD3γε ($7.52 - 10.94 \times 10^{-7}$ μm^4; Fig. 5D, first group) and CD3δε

($6.25 - 9.11 \times 10^{-7}$ μm^4; Fig. 5D, second group) and much higher but similarly valued were

observed for trimolecular interactions with both CD3s ($22.75 - 27.82 \times 10^{-7}$ μm^4; Fig. 5D, third

group). The cooperative analysis also revealed similar positive cooperativity measures in WT and

MT trimers (Supplementary Fig. 7A). Furthermore, trans-interactions between complete WT and

MT 2B4 TCR–CD3 complexes expressed on hybridoma cells with C-terminally biotinylated

K5:IE^k pMHC reported orders of magnitude higher than *cis*-counterparts but near-identical values

with each other ($5.59 - 5.87 \times 10^{-4}$ μm^4; Fig. 5D, *fourth group*). These results rule out zero-force

kinetics of TCRαβ–CD3 *cis*-interaction and TCR–pMHC *trans*-interaction as a possible reason for

the altered T cell activation by the mutations.

However, under force, more pronounced kinetic differences emerged in terms of the catch-slip

bond. Bond lifetime *vs.* force curves for WT and MT TCRαβs with CD3γε were similar (Fig. 5E);

so were those with CD3δε, except for the NP mutation (Fig. 5F), which exhibited a markedly

enhanced catch bond. Interestingly, when comparing aganist each other, their force-dependent

bond lifetimes demonstrated asymmetry again, as measured by the optimal forces where bond

lifetime peaks and the catch bond intensities (a metric summarizing force-dependent bond lifetime

changes (*42*)), both showing significantly higher values for four TCRαβ mutants interacting with

CD3δε than CD3γε (Supplementary Fig. 7B, C). This validates the results from structural analysis

and MD simulations (Fig. 4C-F), confirming that CD3δε is more force-responsive than CD3γε.

In the case of co-presenting both CD3s, bond lifetime profiles revealed marked mutational effects

(Fig. 5G). Notably, the NH LOF TCRαβ mutant formed exceptionally stable bonds with CD3γε–

CD3δε, sometimes exceeding 100 seconds, resulting in an average bond lifetime several-fold

longer than the WT TCRαβ. Similarly, the GK LOF mutant exhibited an enhanced catch bond with

longer lifetimes across the entire force range, whereas the NP GOF mutant displayed a bond profile

resembling WT. Recalling our bond measurements for corresponding *trans*-interactions (Fig. 5H,

replotted from Ref. (*42*)), the mutational effects on TCR–pMHC *trans*-interactions and TCRαβ–

CD3 ECD *cis*-interaction seem exhibit a reverse correlation: the longer the *cis*-bond, the shorter

the *trans*-bond and vice versa.

Correlating *cis*- and *trans*-interactions between TCRαβ and CD3s with T cell signaling.

We next inspected possible correlations between IL-2 secretions (Fig. 5C) and the kinetic measures

of the ECD *cis*-interactions of WT and MT TCRαβs with either CD3γε, CD3δε, or both in the

absence and presence of force. For bimolecular *cis*-interactions, as expected from their statistically

indistinguishable values across each group, the effective 2D affinities (Fig. 5D, *first two groups*)

showed poor predictive power for normalized IL-2 AUC, with steep regression slopes and/or low

R^2 values (Fig. 6A). Here, the inverse of the slope indicates the sensitivity-of-prediction, and R^2

reflects the goodness-of-fit (see Methods). Poor predictive powers for T cell function were

similarly observed from lifetimes measured at zero force and optimal force (Fig 5E, F) for both

TCRαβ–CD3γε and TCRαβ–CD3δε bimolecular *cis*-bonds (Fig. 6B, *orange and blue*, respectively;

Supplementary Fig. 7D). Of note, slightly better correlation with T cell signaling was observed for

TCRαβ bonds with CD3δε than CD3γε at both types of lifetime measures, with higher R^2 values

and more gradual slopes (Fig. 6B, *dotted vs. solid lines*), potentially because of the differential

force-responsiveness.

To examine the contribution of cooperativity of the CD3γε–TCRαβ–CD3δε ECD *cis*-interaction

to T cell activation, the normalized bond numbers (Fig. 5D, *third group*) and lifetimes at zero force

(Fig. 5H, 0 pN) of the trimeric interaction were analyzed, but again found lacks of functional

correlations due to their comparable values (Fig. 6C, *red*), similar to their bimolecular counterparts.

In contrast to the poor function correlations with the peak lifetime at optimal force of the two

bimolecular bonds (Fig. 6B, *solid lines*), a significant reverse correlation was observed between

peak lifetime of the cooperative trimolecular bond (Fig. 5H, *optimal force*) and IL-2 production,

with a much higher R^2 value and a much flatter slope (Fig. 6C, *black*).

The sharp contrast between the differential predictive powers at zero and optimal forces prompted

us to analyze the force effect more rigorously by plotting *vs.* IL-2 AUC, the goodness-of-fit and

sensitivity-of-prediction based on WT and MT lifetimes under each force, finding that the negative

correlation persisted across nearly the entire force range tested (Fig. 6D). Because of the negative

correlations, the sensitivity-of-prediction followed the reverse trends of catch bonds, with biphasic

behavior most pronounced at optimal force, where force-dependent lifetimes diverged the most

between TCR WT and MTs. The Pearson coefficient rapidly approached -1 upon application of

force and remained strongly correlated across $\sim$ 5-20 pN. These results underscore a

mechanotransduction role of the cooperative trimeric *cis*-catch bond.

Since IL-2 produced by T cells expressing these WT and MT TCRs correlates with the peak

lifetime of TCR–K5:IE^k *trans*-bonds (*42*), we correlated the peak lifetimes of *trans*- against *cis*-

interactions, finding strong anti-correlations with those of the trimer ($R^2 \sim 0.8$, Fig. 6E, *black*), but

not dimers ($R^2 < 0.5$, Fig. 6E, *orange* and *blue*). To confirm the force effect on *cis*- and *trans*-

coordination, we plotted the force-dependent fitting of *cis*- *vs.* *trans*- bond profiles for WT and NP,

GK, and NH mutants indicated no correlation in the absence of force (Pearson coefficient, $r \sim 0$ at

zero force), and weak correlations in the presence of force for the bimolecular *cis*-bonds ($0.33 < r$

< 0.73 for $\gamma\varepsilon$, and $-0.03 < r < 0.82$ for $\delta\varepsilon$; Supplementary Fig. 7E), but strong negative-correlations

in the presence of force for the trimolecular *cis*-bond (r dropped sharply, reaching $\sim$ -1 at 10 pN

before rebounding; Fig. 6F). Such correlation analysis reveals an interesting synergy between

trans- and *cis*-interactions, suggesting a possible feedback mechanism of the TCR

mechanosensory machine: from the constant domains, the downstream mechanotransduction

region transmitting back to the variable domains, the antigen recognition region.

DISCUSSION

Noncovalent *cis*-interactions between the TCRαβ and CD3 chains are essential because they not

only hold the octameric TCR–CD3 complex together stably but also provide the "wiring" bridging

the >10 nm spatial gap between ligand engagement at the TCRαβ N-termini and ITAM

phosphorylation at the CD3 cytoplasmic tails (Supplementary Fig. 8A, *right*). NMR chemical-shift

perturbation experiments revealed TCRαβ–CD3γε and TCRαβ–CD3δε ECD dimeric interactions

(*26*), but their functional relevance has been difficult to test because these contacts are

exceptionally weak and fast. Using MAF and BFP assays that measure direct physical interactions

in the same membrane-proximal geometry where these interfaces exist, we quantified the kinetics

of TCRαβ–CD3γε and TCRαβ–CD3δε bimolecular binding and CD3γε–TCRαβ–CD3δε

trimolecular binding both in the absence and presence of force, showing that these ECD *cis*-

interactions are not merely structural accessories but are force-bearing, force-regulated bonds. A

key indication that the TCR–CD3 ectodomain junction can function as a mechanically addressable

coupling element—capable of transmitting, filtering, and stabilizing information under load rather

than simply holding the complex together at rest—is the discovery that these interactions form *cis*-

catch bonds (Supplementary Fig. 8A, *left*).

However, the individual TCRαβ–CD3γε and TCRαβ–CD3δε bimolecular catch bonds are too

short-lived to sustain mechanotransmission over the lifetime of an agonist TCR–pMHC interaction.

We showed that the solution to this paradox is cooperativity: When CD3γε and CD3δε engage

TCRαβ concurrently, the resulting trimolecular assembly exhibits strong positive *cis*-cooperativity,

more likely forming a long-lived *cis*-catch bond whose mechanical stability matches that of the

TCR–pMHC *trans*-catch bond (Supplementary Fig. 8B). While early assembly studies suggested

cis-interaction cooperativity among the TM bundles in the TCR–CD3 intra-complex (*17, 74*) since

both CD3γε and CD3δε were required for their enhanced assembly with the TCRαβ (*25*), the

present work qualitatively demonstrated and quantitatively evaluated the trimolecular binding

cooperativity, using a kinetic approach that follows our previous analysis of positive cooperativity

in the TCR–pMHC–CD8 (*50, 75, 76*) and TCR–pMHC–CD4 (*46*) trimolecular interactions, as

well as the negative cooperativity between two distinct sets of immuno-*trans*-interactions: TCR–

pMHC–CD8 and PD-1–PD-ligands (*77*). Note that our measurements are likely underestimated

because the TCR and CD3 ECDs were measured under pseudo-*cis* configurations without TM

bundling, which would place them in the most appropriate orientations, especially when CD3γε

and CD3δε were captured randomly as CD3m on the beads. In addition to biophysical

measurements, CMD and SMD simulations provide multiple lines of corroborative evidence for

strong synergy. These include contact areas, free energies, dissociation times, distinct dynamic

footprints, and contact residues between the ECDs of TCRαβ and two CD3s separately or

concurrently. Regardless of the metrics used, our results definitively demonstrated positive

cooperativity in the TCR–CD3 *cis*-interactions, as the integrated response of the complex exceeded

that of its individual components. Specifically, the whole trimolecular interaction forms more H

bonds with larger footprints and stronger contacts that are mechanically more stable and longer

lasting than even the sum of the two bimolecular interactions. Conceptually, cooperativity converts

two weak, transient tethers into an integrated "mechanical clutch": force preferentially stabilizes

the coupled state, enabling sustained load sharing and extending the time window for downstream

signaling to commit (Supplementary Fig. 8B).

The ability of the cooperative TCR–CD3 ECD *cis*- catch bond to support force for times that match

the lifetime of cognate TCR–pMHC *trans*-catch bond implies their importance in signal relay.

This may be significant in light of our current perspectives on the role of the *trans*-catch bond in

TCR mechanosensing, which enhances the predictive power for T cell response beyond that of

binding affinity in the absence of force (*9, 39, 78*). By comparison, γδTCR–CD3 complexes lack

interactions among ECDs with flexible CPs (*79-81*); correspondingly, γδTCR–ligand *trans*-

interactions form slip bonds and lack mechanosensing (*12*). This contrast highlights the importance

of the cooperative trimolecular *cis*-catch bond revealed here, which must contribute, at least in

part, to the "mechanical conduit" with the needed structural connectivity among various αβTCR–

CD3 chains important for signaling. Such connectivity must be critical for supporting force,

regardless of whether applied externally by pMHC or generated endogenously via CD3 cytosolic

tails. An interrelated dual role may be envisioned for the cooperativity-reinforced *cis*-catch bond:

First, force does not merely test bond strength; it selects bond state. Catch behavior implies an

underlying force-favored bound conformation (or pathway) that is accessed more frequently or

stabilized under loads. Second, because the cooperative *cis*-catch bond is maximally stabilized

within a defined force regime, the TCR–CD3 junction can act as a "force band-pass filter,"

preferentially transmitting signals within the physiological force window and rejecting either

insufficient load (no stabilization) or excessive load (slip-driven rupture). In this view, the

mechanistic role of *cis*-cooperativity is not simply to strengthen the complex, but to shape how

mechanical information is encoded in bond lifetimes and, thus, signaling probability. Therefore,

our findings support the TCR mechanotransduction hypothesis (*9, 39*) and are also consistent with

the allostery hypothesis (*82*).

Further support for this view comes from observed fluctuations that are inversely correlated with

the cooperatively enhanced contact stability, consistent with dynamic allostery rather than purely

elastic linkages within both TCR and CD3 ECDs. NMR (*32, 33*) and CMD (*33, 41*) studies have

observed long-range allosteric communication across the TCRαβ ECD upon ligand engagement,

manifesting as chemical shift changes and increased dynamics at the CD3 binding site, suggesting

regulatory capacity via distant conformational coupling. Dynamics of CD3s have been

hypothesized to be the primary drivers of cross-membrane signal transmission, based on their more

structured CP regions compared to the flexible TCR anchors (*83*). Although early cryo-EM (*29,*

30) and photo-crosslinking (*27*) studies failed to detect a different stable conformation upon pMHC

or activating antibody ligation in the absence of force, a recent cryo-EM study observed distinct

conformations of the TCR–CD3 complex preserved in native states using nanodiscs (*35*). The

conceptual connection is that cooperative *cis*-coupling provides a plausible means to stabilize and

mechanically gate such large-scale ECD rearrangements: weak pairwise contacts would be

insufficient to sustain coordinated motion, whereas a cooperative *cis*-catch bond could maintain

quaternary integrity while allowing force to bias transitions between compact and open ensembles.

Consistent with the mechanically regulated dynamic allostery is the asymmetric responses of the

two CD3 ECDs, arising from their dissimilar molecular architectures and distinct interactions with

the TCR α and β chains (Supplementary Fig. 8A, *right*). The Cβ FG loop has been suggested to be

important for TCR mechanosensing, partly because of its proximity to the CD3γ ECD, which may

restrict domain motions (*40, 84, 85*). Additionally, CD3δε exhibits heavier glycosylation than

CD3γε (*57, 86*), suggesting a tighter association of CD3γε with TCRβ than CD3δε. Here, we found

that across SMD and BFP (but not CMD and MAF) measurements, CD3δε is more force-

responsive and undergoes greater contact reshaping and fluctuation changes, whereas CD3γε

maintains a more invariant interface with TCRαβ and appears mechanically more stable. This

division of labor suggests a useful conceptual model: CD3γε behaves as a stabilizing strut that

preserves the coupled architecture under load, while CD3δε behaves as a compliant hinge that

accommodates force-induced rearrangements. Such asymmetric compliance provides a plausible

way to meet two competing requirements of mechanotransduction: the complex must remain

connected long enough to transmit load, yet must also remain dynamic enough to permit

conformational/dynamical transitions that ultimately gate tail exposure and ITAM phosphorylation.

Indeed, the ECD asymmetry revealed here may be related to the structural (*35*) and functional (*87*)

differences of CD3 chains during TCR triggering. Our findings shed light on the asymmetric

behaviors of ε chains, warranting further investigation: upon TCR–pMHC binding, the

cytoplasmic tail of γ-associated ε is kept in a constrained, inactive conformation by binding to the

inner leaflet of the plasma membrane, while that of δ-associated ε' is exposed to the cytoplasm,

ready to be phosphorylated by Lck to initiate TCR signaling.

Mutagenesis was used to perturb the ECD *cis*-bonds within the TCR–CD3 assembly to further

elucidate their potential mechanotransduction role. Three double mutations at the TCRαβ constant

domains were examined due to the high dynamic contacts of these residues with the CD3s, which

also show GOF (NP) and LOF (GK and NH) effects as measured by IL-2 production in antigen-

stimulated T hybridoma cells (*26*). Interestingly, none of these mutants altered their effective 2D

affinities with either CD3 or the normalized average number of bonds with both CD3s when

measured in the absence of force. A similar lack of effect was observed with these mutants for

bimolecular catch bonds. The only exception was the NP mutation, which forms a much more

pronounced catch bond with CD3δε but not CD3γε. Again, this force-dependent asymmetry may

be related to the asymmetric behavior of the two ε cytoplasmic tails, for the more adaptive TCRαβ–

CD3δε *cis*-bond may serve as the preferred path of force transmission to enable the larger share of

total force through the ε' chain, facilitating the release of its cytoplasmic tail for signal initiation.

Hypothetically, the mechanotransmission of information requires the TCR–CD3 complex to have

a mechanically integrated architecture with directional constraints, achieved by asymmetric

coupling, to allow binding of different ligands to tip the balance among pre-existing states of an

energetically metastable structure, enabling force-regulated conformational changes to propagate

allosterically. Thus, the Cβ mutants further sharpen the mechanistic picture by separating

equilibrium binding from mechanical coupling. Despite leaving force-free *cis*-affinity/bond

number largely unchanged, these substitutions remodel the force-dependent lifetime landscape—

most prominently for the cooperative trimolecular *cis*-catch bond—demonstrating that the

functional lever resides in force-activated dynamics rather than in static affinity. In other words,

these mutations tune the mechanical impedance of the TCR–CD3 junction: they alter how the

interface redistributes force and stabilizes (or destabilizes) mechanically favored bound states.

This also provides a practical design principle: mechanotransduction can be engineered by

targeting residues that modulate the "force-response" of *cis*-coupling, without necessarily

perturbing antigen recognition chemistry.

Surprisingly, the GOF mutation suppressed the trimolecular *cis*-catch bond, whereas both LOF

mutations enhanced it. This apparent paradox is explained by our MD simulation results, showing

that the GOF mutant enables, whereas the LOF mutants disable the TCRαβ synegistic associations

with the CD3γε–CD3δε. These opposing effects enhance and restrict CD3 flexibility, respectively,

thereby amplifying and attenuating downstream signaling, correspondingly. Indeed, this

explanation is consistent with cryo-EM and crosslinking studies, which show that TCR mutants

that strengthen the TCR–CD3 interface paradoxically impair T-cell function (*27*), underscoring

the functional relevance of CD3 motions.

Previously, we found the same mutations produced a positive correlation between T-cell function

and *trans*-bond profile: the GOF mutant TCR (NP) formed a stronger catch bond than WT, whereas

the LOF mutants (GK and NH) formed a weaker catch bond or even a slip bond, supporting the

predictions from our structural and biophysical model of TCR–pMHC bond profile (*42*). The

model suggests that domain stretching, tilting, and rotating near the hinges of the Vα/Vβ and

Cα/Cβ domains facilitate *trans*-catch bond formation. A striking finding of the present work is the

inverse relationship between the cooperative *cis*-catch bond and T-cell function, together with the

anti-correlation between the *cis*- and *trans*-bond profiles (Supplementary Fig. 8C). These trends

suggest a tradeoff between internal complex stability and external ligand sensitivity: excessive

stabilization of the intra-complex *cis*-junction may "over-clutch" the receptor, restricting the

structural compliance that supports optimal *trans*-catch bond formation and antigen discrimination,

whereas a more labile *cis*-junction may permit the conformational freedom needed to maximize

trans mechanosensing—up to the point where connectivity becomes limiting. Thus, the

unexpected results from this work actually validate our previous model, for the weaker the *cis*-

catch bond, the more flexible the TCR–CD3 complex, the stronger the *trans*-catch bond, and the

more potent the T cell function. This view is also consistent with the findings of previous studies,

which show that LOF and GOF mutations in TM residues correspond to those that stabilize and

destabilize the TCR–CD3 complex, respectively (*18, 88*). We therefore propose that TCR

triggering requires not maximal *cis*-stability per se, but a balance between connectivity and

mobility, such that force can be transmitted while motion remains permissible.

MATERIALS AND METHODS

Proteins, cells, and antibodies

A biotinylation sequence (GLNDIFEAQKIEWHE) was added to the N-terminus of ECD constructs of 2B4 TCRαβ and the C-terminus of human CD3γε, CD3δε, and CD3γε–CD3δε subunits through PCR. For the single heterodimer CD3γε and CD3δε, a 26-residue peptide linker, C-GSADDAKKDAAKKDDAKKDDAKKDGS-N, was engineered to connect different CD3 chains (*89*). For the CD3γε–CD3δε heterodimer–heterodimer, a different 25-residue peptide linker, C-GSSPNSASHSGSAPQTSSAPGSQGS-N, was added to join the two dimers (*90*). Proteins were expressed in One Shot™ *E. coli* BL21 (DE3) (Invitrogen) as insoluble inclusion bodies, refolded, and purified as previously described (*26*). Inclusion bodies for IE^k α (with C-terminal biotinylation sequence) and β chains were produced in One Shot™ *E. coli* BL21 (DE3), refolded with K5 peptide (ANERADLIAYFKAATKF), and purified as described previously (*91*). For all proteins, purified fractions were biotinylated using a BirA protein ligase kit (Avidity LLC). The biotinylated proteins were further purified by gel filtration (S200, GE Life Sciences) in PBS (pH 7.4). The proteins were tested for biotinylation efficiency by gel-shift analysis after incubation with excess streptavidin.

Mouse 58$^{-/-}$ T cell hybridoma cells (*92*) expressing mouse CD3 but not TCRαβ were a generous gift from Dr. Bernard Malissen (Centre d'immunologie de Marseille-Luminy, France). 58$^{-/-}$ cells were transfected with mutant mouse 2B4 TCR constructs through retroviral transduction and cultured as described previously (*26*). The transduced cells were stained with PE anti-mouse CD3ε (clone 145-2C11, eBioscience) and allophycocyanin (APC)-conjugated anti-TCRβ (clone H57-597, eBioscience) mAbs, then sorted for dual expression of CD3 and TCR. The sorted cells were expanded for 6 days and quantified for TCR and CD3ε expression.

Flow cytometry and site density

Samples were stained using 10 µg/ml (as indicated below) of antibodies in 100 µl of FACS buffer (PBS or Hanks Ca^{2+}/Mg^{2+} free, 2% FBS, 5 mM EDTA, and 0.1% NaN3) for 30 min at 4 °C. Subsequently, samples were washed twice in 500 µl and then resuspended in 300 µl of FACS buffer. High-performance analysis and sorting were conducted using BD FACSAria™ III, then flow cytometric data were processed using FlowJo™ Software. PE-conjugated anti-mTCRβ (clone H57-597, BD Biosciences), anti-mIE^k (clone 14-4-4S, BD Biosciences), and PE-Cy7-conjugated anti-mCD3ε (clone 145-2C11, BD Biosciences) were used for *trans*-interactions. PE-conjugated anti-mTCR Vα11.1, 11.2 (clone RR8-1, Biolegend), anti-hCD3ε (clone UCHT1, Biolegend), isotype rat IgG2b,κ (clone RTK4530, Biolegend), isotype mouse IgG1,κ (clone MOPC-21, Biolegend) were used for *cis*-interactions.

Micropipette adhesion frequency assay

The MAF assay for measuring 2D affinity has been described previously (*59, 60*). Briefly, human RBCs were isolated from whole blood obtained from healthy donors using a protocol approved by the Institutional Review Board of the Georgia Institute of Technology. RBCs were biotinylated with various concentrations of biotin-X-NHS linker (Millipore/Sigma) to achieve different levels of surface functionalization. Then, the biotinylated RBCs were incubated with a saturating amount of tetrameric streptavidin (Thermo Fisher) and the biotinylated protein of interest for 30 min at room temperature for each step. Based on available biotin sites, the protein was fully coupled to RBCs via biotin-streptavidin-biotin conjugations, and the resulting surface density was quantified by flow cytometry.

An experiment chamber was assembled by sandwiching coverslips onto short-edge metal spacers, leaving long-edge gaps open for inserting solutions and reagents. The channel bridging two metal

spaces was created by filling the center space with Leibovitz's L-15 Medium (Gibco) and sealing

the solution with mineral oil. The chamber was mounted on the stage of an inverted microscope (a

40×/NA 0.75 objective with a 4× TV tube, Nikon, Eclipse Ti) and imaged at 200 frames per second

(Basler, Ace 2). RBCs, beads, and hybridoma cells were loaded into different regions of the

channel; the forged micropipettes (Sutter Instrument, P-97) were inserted from each side of the

openings to form opposing alignments.

Using suction pressures from the manometer systems, a pair of RBCs (or a RBC and a bead or a

hybridoma cell), bearing receptor and ligand, respectively, were captured by two opposing

capillary micropipettes (Fig. 1A). The piezo-driven target RBC on the right was brought into

contact with the stationary probe RBC on the left in the repeated "approach-contact-retract" testing

cycles. Binding *vs.* non-binding events were sampled at a constant contact time and a consistent

contact area: upon retraction, RBC elongations reflect binding as 1, and no deformations indicate

non-binding as 0. The adhesion frequency ($P_a = \langle s \rangle$) is the mean of the obtained Bernoulli

sequence (*i.e.*, $\{s_1, s_2, s_3, \cdots, s_n\}$), in other words, the probability of the occurrence of binding

events.

Due to the ultra-low affinities, the *cis*-bond formations are *i.i.d.* and rare. Thus, the probability of

forming n bonds obeying the Poisson distribution, $p_n = (\langle n \rangle^n / n!) \exp(-\langle n \rangle)$, with a single

parameter, $\langle n \rangle$, the mean number of bonds formed during contact time. As a result, the adhesion

frequency is expressed as,

$$P_a = 1 - p_0 = 1 - \exp(-\langle n \rangle). \tag{1}$$

731   To further relate the characteristic $\langle n \rangle$ to pairwise *cis*-binding kinetics, we established the kinetic

732   differential equation that is governed by the second-order forward and first-order reverse reactions,

$$\frac{d\langle n \rangle}{dt} = m_r m_l A_c k_{on} - k_{off}\langle n \rangle \tag{2}$$

733   , where m_r and m_l (μm^{-2}) are site densities of receptors and ligands on the RBC surfaces, A_c (μm^2)

734   and t (s) are contact area and time; k_{on} ($\mu m^2 s^{-1}$) and k_{off} (s^{-1}) are 2D on-rate and off-rate, whose

735   ratio gives 2D affinit $K_a = k_{on}/k_{off}$ (μm^2). Clearly, one of Eq. 2 solutions is,

$$\langle n \rangle = m_r m_l A_c K_a[1 - \exp(-k_{off}t)] \tag{3}$$

736   , whose integration constant is zero, assuming there is no bond initially. Because of the rapid off-

737   rates of *cis*-interactions, the exponential term vanishes at steady-state with a long contact time (2

738   s). As a result, effective 2D affinity $A_c K_a$ can be simply derived from P_a by,

$$A_c K_a = \frac{\langle n \rangle}{m_r m_l} = \frac{-\ln(1 - P_a)}{m_r m_l}. \tag{4}$$

Here, m_r and m_l were determined by calibrating the mean fluorescence intensity (MFI) of a

protein of interest fully stained with a PE-tagged antibody against the MFI of PE quantitation beads

(BD Biosciences) by flow cytometry.

Biomembrane force probe (BFP)

The BFP thermal fluctuation (*60, 62*) and force-clamp (*43, 50*) assays for measuring bond lifetime

in the absence and presence of force, respectively, have been described in detail previously. Briefly,

isolated human RBCs were treated and functionalized to form ultra-sensitive mechanotransducers.

Probe RBCs were fully biotinylated with a saturated amount of biotin-PEG-NHS (JenKem), then

inflated into hypotonic states by nystatin (ThermoFisher). Next, the probe beads were prepared for

dual roles: to attach to the RBCs and to display the protein of interest. In short, streptavidin-

maleimide (Sigma) was clicked onto thiol-functionalized glass beads (MPTMS, Thermo Fisher)

overnight at 4°C. Afterward, the biotinylated protein-of-interest was partially coupled to the

streptavidin beads (90 minutes at room temperature), leaving enough vacancies for RBC

attachments through biotin-streptavidin interactions. Lastly, either beads, RBCs, or cells were used

as the targets, on which the binding partners were either chemically conjugated or naturally

assembled for the probe to test against.

A similar experiment chamber (see MAF section) was prepared prior to loading the probe RBCs,

probe beads, and targets. In addition to the two opposing micropipettes, a third helper pipette was

inserted to assist probe assembly: capturing smaller probe beads and attaching them to the apex of

RBCs. The microscope was set up in the same way, but aside from Basler visualization, another

high-speed CCD camera (Mako G040B, Allied Vision) was utilized to trace any deformations of

the probe RBCs. To do so, a 60-line strip window across the probe was imaged at 1,200 frames

per second, where a homemade dual-edge tracking program (LabVIEW, Texas Instruments)

scanned the real-time bead positions relative to the reference on the pipette (*i.e.*, the axial length

of probe RBCs). This drift-prone design allows ultra-stable and ultra-sensitive micro-

manipulations with ~ 2 nm spatial and ~ 1 ms temporal resolutions.

1. **Thermal fluctuation assay**

This assay is based on the equipartition theorem (*62, 65*), where the RBCs elastic energy is

quadratic under small deformations,

$$\frac{1}{2}\kappa_x \langle \Delta x^2 \rangle = \frac{1}{2}k_B T \tag{5}$$

, where κ_x is the spring constant of RBC's simple harmonic motion, and k_B is the Boltzmann

constant that scales absolute temperature to energy. The union of the target with the probe, via

bond formation, dampens its Brownian motion, equivalent to adding a parallel spring that stiffens

the system with a ramped-up spring constant. Whereas bond breakage leads to the restoration of

κ_x. Due to the anti-correlation between κ_x and $\langle \Delta x^2 \rangle$, the bond association and dissociation events

were pinpointed by the sudden decreases and increases in the variance of probe displacements,

respectively.

During the course of measurements, we adjusted the manometer system to lightly grab the probe

RBC using the left micropipette, keeping the RBC ultra-soft with small κ_x to promote sensitive

detections. Subsequently, the probe bead was aligned and pushed against the apex of probe RBC,

where biotin-streptavidin firmly fused them to assemble the complete mechano-sensing probe (Fig.

1b, *left*). Then, we brought the target to the proximity of the probe, leaving a tiny gap between the

target's receptor-presenting and the probe bead's ligand-presenting surfaces. Hence, the thermal-

driven Δx occasionally closed the gap to allow the ligand-receptor binding. If bonds occurred, the

durations between the onsets of $\langle \Delta x^2 \rangle$ reductions and the moments of $\langle \Delta x^2 \rangle$ resumptions (*i.e.*,

sequence of events from the exponential random variable, $\{t_1, t_2, t_3, \cdots, t_n\}$) were sampled, whose

mean reports the ligand-receptor lifetime, $\tau = \langle t \rangle$. Note that the adhesion frequency was kept low

enough to ensure limited second bond formation before breaking the first bond.

To calculate the off-rate, k_{off} of pairwise interactions from sampled lifetime events, the kinetic

differential equation of first-order reverse reaction is simplified by setting $k_{\text{on}} = 0$ (Eq. 2) as,

$$\frac{d\langle n \rangle}{dt} = -k_{\text{off}}\langle n \rangle. \tag{6}$$

788   Its solution is scalable with the initial number of bonds, $\langle n_0 \rangle$, and reexpressed as the time-

789   dependent survival probability, $P_{\text{s}}(t)$,

$$P_{\text{s}}(t) = \frac{\langle n \rangle}{\langle n_0 \rangle} = \exp(-k_{\text{off}}t). \tag{7}$$

As a result, $P_{\text{s}}(t)$, the number of events lasting longer than t normalized by total lifetime events,

starts from point (0, 1) and decays exponentially. After rearranging lifetime events chronologically,

the linear regression slope of survival probability *vs.* time in logarithmic scale estimates k_{off}.

2. Force-clamp assay

The force-clamp assay utilized the same BFP micropipette apparatus, probe setups, and coating

strategies as the thermal fluctuation assay (Fig. 1B). Nonetheless, instead of passively monitoring

lifetime events without interference, we precisely quantified and interactively manipulated the

force levels exerted on the ligand-receptor bonds to obtain the force-dependent lifetime, $\tau(f)$. The

probe, as a harmonic oscillator, obeys Hooke's law,

$$f = \kappa_x \Delta x \tag{8}$$

799   , where κ_x converts the axial displacements into the force applied by the spring. To fine-tune probe

800   stiffness, the suction pressure (Δp) on probe RBCs modulates the hypotonic RBC membrane

801   tension, which, in turn, relates to the spring constant by a geometric scaling factor as,

$$\kappa_x(\Delta p) \approx \frac{2\pi}{\ln\left(\frac{4R_0^2}{R_p R_c}\right)} \sigma = \frac{\pi R_p \Delta p}{\left(1 - \frac{R_p}{R_0}\right)\ln\left(\frac{4R_0^2}{R_p R_c}\right)} \tag{9}$$

, where σ is the membrane tension, and the characteristic measures of the probe, R_0, R_p, and R_c,

are radii of the outer RBC sphere, the inner RBC cap, and the bead-RBC circular contact,

respectively. Typically, the designated Δp sets $\kappa_x \sim 0.3$ pN/nm, which establishes the ultra-

sensitive force sensor at a resolution of ± 5 nm in x-deformations.

During experiments, once fine-tuned to the specified stiffness, we utilized a high-precision

piezoelectric linear actuator (Physik Instrumente) to micro-manipulate the target, allowing it to

undergo "approach-contact-retract-hold-return" testing cycles against the stationary probe. Based

on   the   instantaneous

$f(t)$, automated subprograms (LabVIEW, Texas Instruments) made interactive decisions about the

target's   sub-nanometer   movements.   The   target   **approached**   the   probe   till

$f(t)$ dropped below zero. Next, the target was stopped at $f(t) \sim -25$ pN and **contacted** the probe

to facilitate ligand-receptor associations for a fixed period of ~ 0.5 s. At its end, the piezo **retracted**

the target from the probe with two outcomes: a non-binding event was indicated by $\langle f(t) \rangle \sim 0$,

and a binding event was detected by $\langle f(t) \rangle \gg 0$ at the moment of separation (Fig. 1F). If the

ligand-receptor formed a bond and the ramping-up force reached an expected level, f, the target

was **held** in place to maintain the constant Δx, such that the tensile force on the bond was clamped

until the spontaneous dissociation occurred (*i.e.*, the sudden decline of $f(t)$). Later, the piezo

returned the target to its origin to reinitiate the next testing cycle. Among 100s of cycles tested on

a pair of targets and probes, lifetime events (*i.e.*, $\{t_1^f, t_2^f, t_3^f, \cdots, t_n^f\}$) were sampled as the time

intervals between the instants of $f(t)$ ramped up to the desired magnitudes, and the moments of

$f(t)$ suddenly dropped to zero. The mean of lifetime events reports the force-dependent ligand-

receptor lifetime, $\tau^f = \langle t^f \rangle$, where the superscript denotes being subjected to the force-clamping

of f. To ensure that most interactions represented single-bond events, the site densities of the

molecules coated on the beads were adjusted to maintain the adhesion frequency below 20%.

Cooperativity analysis

Multiple metrics were implemented to assess the cooperativity of *cis*-interaction, such as 2D

affinity, force-dependent bond lifetime, *etc*. When CD3γε and CD3δε were co-presented, the

comparisons between the actual measurements and the expected values assuming independence

revealed the level of cooperativity of the TCR and CD3 mixture. To formulate a criterion and a

quantitative index for cooperativity, we derived the general form as,

$$\frac{\Delta \langle Y \rangle}{\langle \widehat{Y}_m \rangle} = \frac{\langle Y_m \rangle - \langle \widehat{Y}_m \rangle}{\langle \widehat{Y}_m \rangle} = \frac{\langle Y_m \rangle}{\langle \widehat{Y}_m \rangle} - 1 \begin{cases} > 0 & \text{positive cooperativity} \\ = 0 & \text{zero cooperativity} \\ < 0 & \text{negative cooperativity} \end{cases} \qquad (10)$$

, where Y is an arbitrary random variable about *cis*-interaction, and Y_m and $\widehat{Y}_m$ are the obtained and

expected values from the CD3 mixture, respectively. Zero cooperativity indicates independent

binding, while positive or negative cooperativity signifies dependent binding, with the magnitude

of cooperativity quantifying the level of dependency.

Three types of cooperativity analysis were employed to evaluate the *cis*-bindings between TCRαβ

and CD3γε, CD3δε mixture using *in vitro* MAF, BFP, and *in silico* SMD dissociation assays. Under

the assumption of independent bindings, the expected values for corresponding assays are listed

below,

MAF
$$\langle \hat{n}_m \rangle = m_{\alpha\beta} m_{\gamma\epsilon} A_c K_a^{\gamma\epsilon} + m_{\alpha\beta} m_{\delta\epsilon} A_c K_a^{\delta\epsilon} \tag{11}$$

BFP
$$\hat{\tau}_m = \frac{m_{\gamma\epsilon} K_a^{\gamma\epsilon}}{m_{\gamma\epsilon} K_a^{\gamma\epsilon} + m_{\delta\epsilon} K_a^{\delta\epsilon}} \tau_{\gamma\epsilon} + \frac{m_{\delta\epsilon} K_a^{\delta\epsilon}}{m_{\gamma\epsilon} K_a^{\gamma\epsilon} + m_{\delta\epsilon} K_a^{\delta\epsilon}} \tau_{\delta\epsilon} \tag{12}$$

SMD
$$\langle \hat{t}_{\delta\gamma} \rangle = \langle t_{\gamma\epsilon} \rangle + \langle t_{\delta\epsilon} \rangle - \frac{\langle t_{\gamma\epsilon} \rangle \langle t_{\delta\epsilon} \rangle}{\langle t_{\gamma\epsilon} \rangle + \langle t_{\delta\epsilon} \rangle}. \tag{13}$$

Here, K_a, τ, and $\langle t \rangle$ in expressions are characteristic kinetic parameters of pairwise *cis*-interactions

quantified from corresponding TCRαβ–CD3γε and TCRαβ–CD3δε assays, while A_c and m are

controllable factors. Note that τ and $\langle t \rangle$ distinguish the quantifications of lifetime using different

approaches, where the former population study reports the ensemble average of multiple

association states (subscript m), and the latter computer study is based on the single trimeric state

(subscript δγ).

Derivations of the expected value of trimeric dissociation lifetime

The expected values for the first two cooperativity analyses are straightforward, and we only show

the derivations for Eq. 13 in detail here. The third cooperativity analysis is based on the waiting

time of dissociations during SMD simulations. Again, this differs from the population mean of

BFP lifetime measurements because we simulated a single-molecular construct at a time, either

TCRαβ–CD3γε or TCRαβ–CD3δε, or CD3γε–TCRαβ–CD3δε complexes. Recall that the survival

probabilities for the pairwise associations of TCRαβ–CD3γε and TCRαβ–CD3δε are,

$$P_s^{\gamma\epsilon} = \exp\left(-\frac{t}{\langle t_{\gamma\epsilon}\rangle}\right) \text{ and } P_s^{\delta\epsilon} = \exp\left(-\frac{t}{\langle t_{\delta\epsilon}\rangle}\right) \tag{14}$$

853 , respectively, where $\langle t_{\gamma\epsilon}\rangle$ and $\langle t_{\delta\epsilon}\rangle$ are their corresponding average waiting time till dissociation.

854 Assuming the CD3γε and CD3δε dissociations from the concurrently bound states are independent,

855 and the presence of high loads prevents any rebinding upon dissociations, the cumulative joint

856 probability for TCRαβ to dissociate from both concurrently during $(0, t)$ is,

$$F_{\delta\gamma}(t) = \left(1 - P_s^{\gamma\epsilon}(t)\right)\left(1 - P_s^{\delta\epsilon}(t)\right) \tag{15}$$

$$= 1 - e^{-t/\langle t_{\gamma\epsilon}\rangle} - e^{-t/\langle t_{\delta\epsilon}\rangle} + e^{-t(1/\langle t_{\gamma\epsilon}\rangle + 1/\langle t_{\delta\epsilon}\rangle)}.$$

857 The corresponding probability density is,

$$\rho_{\delta\gamma}(t) = \frac{dF_{\delta\gamma}(t)}{dt} = \frac{e^{-t/\langle t_{\gamma\epsilon}\rangle}}{\langle t_{\gamma\epsilon}\rangle} + \frac{e^{-t/\langle t_{\delta\epsilon}\rangle}}{\langle t_{\delta\epsilon}\rangle} - \frac{e^{-t(1/\langle t_{\gamma\epsilon}\rangle + 1/\langle t_{\delta\epsilon}\rangle)}}{1/\langle t_{\gamma\epsilon}\rangle + 1/\langle t_{\delta\epsilon}\rangle}. \tag{16}$$

858 Thus, the expected value of trimeric dissociation lifetime under the non-coop assumption is,

$$\langle \hat{t}_{\delta\gamma}\rangle = \int_0^\infty t\rho_{\delta\gamma}(t)dt = \langle t_{\gamma\epsilon}\rangle + \langle t_{\delta\epsilon}\rangle - \frac{\langle t_{\gamma\epsilon}\rangle\langle t_{\delta\epsilon}\rangle}{\langle t_{\gamma\epsilon}\rangle + \langle t_{\delta\epsilon}\rangle}. \tag{17}$$

Kinetic modeling of dual-species interaction

861 To model the dual-species scenario, in addition to the pairwise interactions in the single-species

862 scenario, only the CD3γε–TCRαβ–CD3δε trimolecular interaction is allowed as a concurrent

863 binding state. As such, for simplicity, the total TCRαβ bonds with CD3m include only three

864 subpopulations.

$$\langle n_m\rangle = \langle n_{\gamma\epsilon}^m\rangle + \langle n_{\delta\epsilon}^m\rangle + \langle n_{\delta\gamma}^m\rangle. \tag{18}$$

, where n with different subscriptions denote the number of bonds for the TCRαβ–CD3γε, TCRαβ–

CD3δε bimolecular interactions, and the CD3γε–TCRαβ–CD3δε trimolecular interaction,

sequentially. Here, the concurrent mode exhibits unique kinetics, redistributing the probabilities of

dimeric and trimeric bond formation.

The competitive kinetics between the pairwise and concurrent TCRαβ–CD3s *cis*-interactions

follow the two-step model. In the first step, the TCRαβ interacts with either CD3γε (kinetic rates

k_1 and k_{-1}) or CD3δε (kinetic rates k_2 and k_{-2}) to form TCRαβ–CD3γε or TCRαβ–CD3δε

bimolecular bonds. In the second step, the TCRαβ–CD3δε bond interacts with CD3γε (kinetic rates

k_3 and k_{-3}) or the TCRαβ–CD3γε bond interacts with CD3δε (kinetic rates k_4 and k_{-4}) to form

CD3γε–TCRαβ–CD3δε trimolecular bonds. Refer to Eq. 2, the modified kinetic differential

equations for the steady state are,

$$\frac{d}{dt}\begin{Bmatrix} \langle n_{\gamma\epsilon}^m \rangle \\ \langle n_{\delta\epsilon}^m \rangle \\ \langle n_{\delta\gamma}^m \rangle \end{Bmatrix} + \begin{bmatrix} m_{\delta\epsilon}k_4 + k_{-1} & 0 & -k_{-4} \\ 0 & m_{\gamma\epsilon}k_3 + k_{-2} & -k_{-3} \\ -m_{\delta\epsilon}k_4 & -m_{\gamma\epsilon}k_3 & k_{-3} + k_{-4} \end{bmatrix} \begin{Bmatrix} \langle n_{\gamma\epsilon}^m \rangle \\ \langle n_{\delta\epsilon}^m \rangle \\ \langle n_{\delta\gamma}^m \rangle \end{Bmatrix} = \begin{Bmatrix} m_{\alpha\beta}m_{\gamma\epsilon}A_c k_1 \\ m_{\alpha\beta}m_{\delta\epsilon}A_c k_2 \\ 0 \end{Bmatrix} \tag{19}$$

876   , whose solutions are derived as,

$$\langle n_{\gamma\epsilon}^m \rangle = \frac{m_{\alpha\beta}m_{\gamma\epsilon}A_c}{m_{\delta\epsilon}k_4 + k_{-1}}\left[k_1 + m_{\delta\epsilon}k_{-4}\frac{k_1 k_4(m_{\gamma\epsilon}k_3 + k_{-2}) + k_2 k_3(m_{\delta\epsilon}k_4 + k_{-1})}{k_{-1}k_{-4}(m_{\gamma\epsilon}k_3 + k_{-2}) + k_{-2}k_{-3}(m_{\delta\epsilon}k_4 + k_{-1})}\right] \tag{20}$$

$$\langle n_{\delta\epsilon}^m \rangle = \frac{m_{\alpha\beta}m_{\delta\epsilon}A_c}{m_{\gamma\epsilon}k_3 + k_{-2}}\left[k_2 + m_{\gamma\epsilon}k_{-3}\frac{k_1 k_4(m_{\gamma\epsilon}k_3 + k_{-2}) + k_2 k_{(45)}(m_{\delta\epsilon}k_4 + k_{-1})}{k_{-1}k_{-4}(m_{\gamma\epsilon}k_3 + k_{-2}) + k_{-2}k_{-3}(m_{\delta\epsilon}k_4 + k_{-1})}\right] \tag{21}$$

$$\langle n_{\delta\gamma}^m \rangle = m_{\alpha\beta}m_{\gamma\epsilon}m_{\delta\epsilon}A_c\frac{k_1 k_4(m_{\gamma\epsilon}k_3 + k_{-2}) + k_2 k_3(m_{\delta\epsilon}k_4 + k_{-1})}{k_{-1}k_{-4}(m_{\gamma\epsilon}k_3 + k_{-2}) + k_{-2}k_{-3}(m_{\delta\epsilon}k_4 + k_{-1})} \tag{22}$$

, where *m* with different subscriptions indicates the site densities of TCRαβ, CD3γε, and CD3δε,

respectively.

The cooperativity resides in the second step by allowing the CD3γε–TCRαβ–CD3δε concurrent

cis-engagements. As expected, letting $k_3 = k_4 = 0$ abolishes the trimolecular bonds and reduces

Eq. 20, 21 to the forms of Eq. 3 for the TCRαβ–CD3γε and TCRαβ–CD3δε bimolecular bonds.

Note that the comparison between the gain in $\langle n_{\delta\gamma}^m \rangle$ and the loss in $\langle n_{\gamma\epsilon} \rangle - \langle n_{\gamma\epsilon}^m \rangle$ and $\langle n_{\delta\epsilon} \rangle - \langle n_{\delta\epsilon}^m \rangle$

defines the sign of cooperativity. Furthermore, the reduction of bimolecular bond fraction leads to

the inequality of $\langle n_{\delta\gamma}^{m} \rangle \geq \Delta\langle n \rangle$. From which, the lower bound for the trimolecular bond fraction

can be estimated using solely the value of cooperativity by $\langle n_{\delta\gamma}^{m} \rangle / \langle n_{m} \rangle \geq \Delta\langle n \rangle / (\Delta\langle n \rangle + \langle \hat{n}_{m} \rangle)$ (*i.e.*,

cooperativity divided by one plus cooperativity). For example, based on the bond number

cooperativity of 170% that we measured, the chance of trimolecular bond formation is higher than

62% of the total bonds.

To model the dual-species dissociation processes, similar to the off-rate analysis in the thermal

dynamic assay, by setting the right-hand side of Eq. 19 master equations to zero, we obtained the

two-step dissociation of dual-species following a multi-exponential decay. Again, letting the

second-step kinetic rates $k_3 = k_{-3} = k_4 = k_{-4} = 0$ reduces the master equations to two

uncoupled first-order reverse reactions for the TCRαβ–CD3γε and TCRαβ–CD3δε *cis*-dissociation,

following single-exponential decays.

Kinetic modeling of catch-bonds

The *cis*-bond profiles of TCRαβ interacting with CD3γε, CD3δε, or both all demonstrate catch-

slip behaviors, with bi-exponential-decay survival probabilities, suggesting the presence of two

bound species (*93-95*), particularly under the effects of clamping force. Therefore, for the analysis

of force-dependent dissociations of bimolecular bonds, we assumed that a single bond might

undergo either a fast-dissociation pathway from a weak state or a slow-dissociation pathway from

a strong state (indicated by *), also allowing them to exchange with each other. For the analysis of

the mixture bond data, besides combining the two-pathway models obtained for the pairwise

TCRαβ–CD3γε and TCRαβ–CD3δε dissociations, we added a third population of trimolecular

CD3γε–TCRαβ–CD3δε bonds using the same modeling scheme. The corresponding reaction

diagrams for all scenarios are depicted in Supplementary Fig. 2A.

The force dependency of kinetic rates follows Bell's equation (96),

$$k_{i,j}(f) = k_{i,j}^0 \exp\left(\frac{f x_{i,j}}{k_{\mathrm{B}} T}\right) \qquad (23)$$

, where the subscript i denotes rates of dissociation along fast (-f) or slow (-s) pathways and of

activation into (a) or deactivation from (-a) the strong states; the subscript j denotes different bond

types of TCRαβ–CD3γε (γ), TCRαβ–CD3δε (δ), and CD3γε–TCRαβ–CD3δε (δγ); the superscript

0 denotes the respective force-free kinetic rates. According to transition state theory, the force

linearly modulates the free energy of the transition barrier, $\Delta E^{\ddagger}(f) = \Delta E_0^{\ddagger} - fx$, and subsequently

regulates the kinetic rates, $k(f) = \exp\left(-\left(\Delta E_0^{\ddagger} - fx\right)/k_{\mathrm{B}}T\right)$, werhe x is the characteristic

transition distance, and k^0 is proportional to $\exp\left(-\Delta E_0^{\ddagger}\right)$.

This two-state, two-pathway dissociation model (46) recapitulated the lifetime of catch bonds and

the bi-exponential decay of survival probabilities. Here, the force-free fast dissociation rates for

the dimeric bonds were obtained from the thermal fluctuation assay, and the remaining parameters

were then evaluated by globally fitting the model to the force-lifetime distributions via maximum

likelihood estimation (MLE). The initial proportions of the two states were determined from the

ratios of the kinetic rates, assuming equilibrium. A two-step optimization procedure was employed

to minimize the MLE: an extensive parameter space was explored using differential evolution to

obtain an approximate solution, followed by L-BFGS to hone in on the minimum (Supplementary

Fig. 2E-G). The resultant kinetic rates *vs*. force relationships are presented in Supplementary Fig.

2B-D, and Bell's parameters are summarized in Supplementary Table 2. The standard deviation

for each parameter was obtained by inverting the Hessian matrix at the MLE to approximate the

covariance matrix, whose diagonal entries correspond to the variance of the fitted parameters.

Molecular dynamics simulations

1. Molecular constructs

All simulations were based on the cryo-EM structure (PDB ID: 6JXR) (*28*). The TCRαβ, CD3γε, and CD3δε were truncated near the CPs between extracellular and intracellular regions, leaving the complete TCR–CD3s ECD in its trimeric assembly. The TCRαβ were trimmed at their C-termini, leaving the α-chain Gly22-Glu237 and the β-chain Val26-Glu269. The CD3γε and CD3δε were obtained by truncating the γ-chain after Cys107, ε-chain after Asn121, δ-chain after Glu98, and ε'-chain after Glu124. To minimize potential trimming effects, all chain ends were modified to prevent the exposure of charged residues. An acetyl group was added in front of the N-terminus, and an N-methylamine group capped the C-terminus of every chain. All hydrogen atoms, as well as any missing atoms, were added back through Gromacs utilities (*28*). The disulfide bonds in the original cryo-EM structure were preserved through Gromacs interactive S-S bridge selection procedures (*28*). The resulting CD3γε–TCRαβ–CD3δε trimeric assembly was used to simulate the concurrent *cis*-binding. Then, three partial assemblies were generated to simulate the pairwise *cis*-interactions by removing CD3δε, CD3γε, and TCRαβ individually from the fully-assembled ECDs, yielding dimeric TCRαβ–CD3γε, TCRαβ–CD3δε, and CD3γε–CD3δε, respectively.

2. Unit cell preparation

Unit cells enclosing molecular systems of interest were built with physiologically realistic surroundings and then brought into thermodynamically favorable states. The initial structures were oriented and centered within optimized orthorhombic cells, which accommodated the physical extensions in all directions with a ≥ 1.6 nm buffering zone apart from the box sides to ensure the protein was well-isolated. The molecules in unit cells were solvated using the TIP3P water model (*97*). The negative charges were counterbalanced using sodium ions, and then the ionic strength was set to ~ 150 mM with sodium chloride. To further prepare the system thermodynamically,

energy-unfavorable close contacts were eliminated through the steepest descent and conjugate

gradient energy minimization cycles. While restraining the heavy atoms in macromolecules, the

unit cell was brought to a suitable ensemble employing a 200-ps velocity rescaling simulation to

adjust the temperature and a 200-ps Berendsen Barostat simulation to correct the pressure (300 K,

1.01325 bar). After releasing the heavy-atom restraints, the systems were considered ready for

subsequent equilibration and production runs.

3.   Conventional molecular dynamics

CMD simulations were performed using GROMACS (version 2019.6) (*98-100*) under the recent

AMBER99SB*-ILDNP force field with improved helix-coil transitional and side-chain torsional

dynamics (*101*). From previously prepared unit cells, the set of initial velocities was randomly

sampled from the Maxwell-Boltzmann distribution. The molecular trajectories were self-evolved

under the leap-frog algorithm by integrating Newton's law of motion every 2 fs. For CMD, the

individual simulation runs were executed under periodic boundary conditions to sample the NVT

ensemble under corresponding constant volumes (*i.e.*, CD3γε–TCRαβ–CD3δε: $12.5 \times 10.8 \times 9.9$

nm^3; TCR–CD3γε: $12.6 \times 9.8 \times 8.6$ nm^3; TCR–CD3δε: $11.9 \times 10.3 \times 9.2$ nm^3; CD3γε–CD3δε:

$11.5 \times 8.1 \times 8.0$ nm^3) and 300 K constant temperature. The neighborhood lists were refreshed

every 10 steps using the buffered Verlet scheme, with a 1.2 nm non-bonded cutoff for Coulomb

and van der Waals interactions. The long-range potentials of electrostatic interactions were

computed using the Particle Ewald Mesh summation method. All systems were simulated for 100s

ns with three repeats, where each trajectory consists of at least a 50 ns equilibration stage and the

remaining production run. Conformational snapshots every 0.1 ns during the production stage were

extracted for subsequent analysis.

4.   Steered molecular dynamics

SMD simulations were initiated from the well-equilibrated configurations from corresponding

CMD simulations (*i.e.*, the conformational snapshots at 100 ns). They were prepared according to

the Unit cell preparation section, but with a 20-nm box length to accommodate the molecular

extension and separation under external pulling. The external forces with constant magnitudes (*i.e.*,

force-clamp) or constant velocities (*i.e.*, force-ramp) along the fixed *z*-direction were added to the

trimeric and dimeric complexes. The clamping or ramping forces were applied to the N-terminus

of the TCRα chain to accelerate the dissociations while keeping the C-termini of CD3ε chains

fixed to mimic the anchor effect of their TM domains. For force-clamp simulations, five

independent trajectories under 100, 125, 150, 175, and 200 pN were generated for each of the two

dimeric systems, but only three independent runs under 150, 175, and 200 pN were executed for

the trimeric complex. We aborted the CD3γε–TCRαβ–CD3δε simulations at smaller clamp forces

of 100 and 125 pN due to the inability to observe dissociation events after the exceedingly long

waiting time (> 1000 ns). For force-ramp simulations, a 70 pN/nm pseudo spring was utilized to

exert increasing levels of pulling to accelerate dissociations. One end of the spring was linked to

the α chain N-terminus of the targeting molecule, and the other free end departed in the *z*-direction

at 1 nm/ns. Five repeats each for two dimeric and one trimeric ECD complexes were performed to

assess the corresponding rupture forces as a consistency check.

Occupancy analysis

Several variations of occupancy analysis were employed to dynamically evaluate the extent of a

particular residue in dynamic contact with its binding partner throughout the simulation. At

timestamp i, the residue being analyzed is defined to be in contact (denoted $o_i = 1$) if the center

of mass of this residue is within 4 Å of any atoms of its binding partner; otherwise, it is not in

touch (denoted $o_i = 0$). The occupancy index (%) indicates the fraction of time in contact during

a given period, as $\sum_i o_i/N$, where N is the total number of analyzed frames.

1. Residue-to-residue occupancy (Supplementary Fig. 5)

By setting residues within a subunit as a reference group, this occupancy analysis quantifies the

extent of dynamic interaction between each residue from one subunit and every single residue from

the other subunit over a given period. Thus, it evaluated the level of any pairwise contacts, thereby

forming a correlation matrix where the column indexes the residues from one subunit; the row

indexes the residues from the other subunit, and the elements denote the values of residue-to-

residue relative binding strength, ranging from 0 to 1.

1. Residue-to-domain occupancy (Supplementary Fig. 6)

This occupancy analysis quantifies each residue against all residues within its binding partner,

using the entire subunit as the reference group. It is usually smaller than the sum of residue-to-

residue occupancy over the binding domain because the residue in question may simultaneously

be in contact with multiple residues in the domain under consideration; on the other hand, these

contact periods can only be counted once for residue-to-domain occupancy. To account for this

disadvantage, ΣOccupancy is defined as the integration of the CMD and SMD row-/column-wise

sums of residue-to-residue occupancies; while ΔOccupancy is defined as the differentiation of the

CMD from the SMD row-/column-wise sums of residue-to-residue occupancies, which were

calculated to summarize and discriminate force-free and force-clamp interface dynamics (Fig. 4G,

H).

2. Domain-to-domain occupancy

This occupancy analysis further reduces data presentation to examine the overall binding tendency

against the other domain by averaging over all interacting residues (*i.e.,* positive terms of residue-

to-domain occupancies), providing a measure of the overall domain-to-domain adhesion strength.

Linear regression

Linear regression was conducted to assess the correlations between *cis*-kinetic measures (as x) and

functional outputs or *trans*-kinetic measures (as y). The goodness-of-fit was quantified using

Pearson's correlation coefficient, *i.e.*, $(n \sum xy - \sum x \sum y)/\sqrt{(n \sum x^2 - (\sum x)^2)(n \sum y^2 - (\sum y)^2)}$,

which preserves the sign of correlation, as well as its square, R-squared. The sensitivity-of-

prediction, *i.e.*, $(n \sum x^2 - (\sum x)^2)/(n \sum xy - \sum x \sum y)$, is the inverse of the linear-regression

slope.

ACKNOWLEDGEMENTS

This work was supported by NIH grants R01GM124489 (to M.K.) and R01CA284604 (to M.K. and C.Z.) and U01 CA214354 (to C.Z. and M.K.). The MD simulations were supported by an NSF award MCA08X014 (to C.Z.) in advanced computing infrastructure for the U.S. and performed in the Extreme Science and Engineering Discovery Environment (XSEDE). Cell sorting/flow cytometry technologies were provided by NYU Langone's Cytometry and Cell Sorting Laboratory, which is supported in part by NIH grant P30CA016087. SIYR:H2-K^b protein was made by the NIH Tetramer Core Facility at Emory University. We thank Larissa Doudy for technical assistance, Johannes Huppa for providing the H57 scFv, and Bernard Malissen for providing mouse 58$^{-/-}$ T cell hybridoma cells.

AUTHOR CONTRIBUTIONS

C.Z., P.C., and C.G. conceived the study; C.Z. and M.K. directed the study; P.C., Z.Y., C.G., and A.N. performed experiments; P.C. performed simulations; P.C. and A.N. performed structural analysis; A.N., S.B, and D.G produced proteins; P.C. and S.T. analyzed data using mathematical models; P.C., A.N., Z.Y., M.K., and C.Z. analyzed data and wrote the paper with contributions from other authors.

STATEMENT OF COMPETING INTERESTS

The authors declare no competing interests. M.K. serves on the scientific advisory boards of Genentech and Merck and Co. and received research support from Merck Sharp & Dohme Corp., a subsidiary of Merck and Co., Inc., Genentech, Biogen, Novartis, and the Mark Foundation.

1047

REFERENCES

1049

1. A. H. Courtney, W. L. Lo, A. Weiss, TCR Signaling: Mechanisms of Initiation and Propagation. *Trends Biochem Sci* **43**, 108–123 (2018).

2. C. Xu *et al.*, Regulation of T cell receptor activation by dynamic membrane binding of the CD3epsilon cytoplasmic tyrosine-based motif. *Cell* **135**, 702–713 (2008).

3. L. A. Pitcher, N. S. C. van Oers, T-cell receptor signal transmission: who gives an ITAM? *Trends in Immunology* **24**, 554–560 (2003).

4. B. Malissen, P. Bongrand, Early T Cell Activation: Integrating Biochemical, Structural, and Biophysical Cues. *Annual Review of Immunology* **33**, 539–561 (2015).

5. K. C. Garcia, L. Teyton, I. A. Wilson, Structural basis of T cell recognition. *Annual Review of Immunology* **17**, 369–397 (1999).

6. R. A. Mariuzza, P. Agnihotri, J. Orban, The structural basis of T-cell receptor (TCR) activation: An enduring enigma. *J Biol Chem* **295**, 914–925 (2020).

7. M. Wieczorek *et al.*, Major Histocompatibility Complex (MHC) Class I and MHC Class II Proteins: Conformational Plasticity in Antigen Presentation. *Frontiers in Immunology* **8**, (2017).

8. N. Jeffreys, J. M. Brockman, Y. Zhai, D. E. Ingber, D. J. Mooney, Mechanical forces amplify TCR mechanotransduction in T cell activation and function. *Applied Physics Reviews* **11**, 011304 (2024).

9. S. Travaglino, Y. Jeon, Y. Kim, C. Zhu, H.-K. Choi, Mechanotransduction through T cell receptor: consensus, controversies, and future outlooks. *Experimental & Molecular Medicine*, (2025 In press).

10. M. S. Kuhns *et al.*, Evidence for a functional sidedness to the alphabetaTCR. *Proc Natl Acad Sci U S A* **107**, 5094–5099 (2010).

11. Elaine P. Dopfer *et al.*, The CD3 Conformational Change in the γδ T Cell Receptor Is Not Triggered by Antigens but Can Be Enforced to Enhance Tumor Killing. *Cell Reports* **7**, 1704–1715 (2014).

12. R. J. Mallis *et al.*, Molecular design of the γδT cell receptor ectodomain encodes biologically fit ligand recognition in the absence of mechanosensing. *Proceedings of the National Academy of Sciences* **118**, e2023050118 (2021).

13.   J. P. Van Wauwe, J. G. Goossens, P. C. Beverley, Human T lymphocyte activation by
monoclonal antibodies; OKT3, but not UCHT1, triggers mitogenesis via an interleukin 2-
dependent mechanism. *The Journal of Immunology* **133**, 129–132 (1984).

14.   S. R. Riddell, P. D. Greenberg, The use of anti-CD3 and anti-CD28 monoclonal antibodies
to clone and expand human antigen-specific T cells. *Journal of Immunological Methods*
128, 189–201 (1990).

15.   A. P. Menon *et al.*, Modulating T Cell Responses by Targeting CD3. *Cancers*. 2023
(10.3390/cancers15041189).

16.   Y. Wang *et al.*, A Conserved CXXC Motif in CD3ε Is Critical for T Cell Development and
TCR Signaling. *PLOS Biology* **7**, e1000253 (2009).

17.   M. E. Call, J. Pyrdol, M. Wiedmann, K. W. Wucherpfennig, The Organizing Principle in
the Formation of the T Cell Receptor-CD3 Complex. *Cell* **111**, 967–979 (2002).

18.   A. L. Lanz *et al.*, Allosteric activation of T cell antigen receptor signaling by quaternary
structure relaxation. *Cell Rep* **36**, 109375 (2021).

19.   K. N. Brazin *et al.*, The T Cell Antigen Receptor α Transmembrane Domain Coordinates
Triggering through Regulation of Bilayer Immersion and CD3 Subunit Associations.
Immunity **49**, 829–841.e826 (2018).

20.   Z.-Y. J. Sun, K. S. Kim, G. Wagner, E. L. Reinherz, Mechanisms Contributing to T Cell
Receptor Signaling and Assembly Revealed by the Solution Structure of an Ectodomain
Fragment of the CD3εγ Heterodimer. *Cell* **105**, 913–923 (2001).

21.   N. Martinez-Martin *et al.*, Cooperativity between T cell receptor complexes revealed by
conformational mutants of CD3epsilon. *Sci Signal* **2**, ra43 (2009).

22.   R. A. Fernandes *et al.*, T cell receptors are structures capable of initiating signaling in the
absence of large conformational rearrangements. *J Biol Chem* **287**, 13324–13335 (2012).

23.   H. Liu, M. Rhodes, D. L. Wiest, D. A. A. Vignali, On the Dynamics of TCR:CD3 Complex
Cell Surface Expression and Downmodulation. *Immunity* **13**, 665–675 (2000).

24.   M. S. Kuhns, M. M. Davis, Disruption of extracellular interactions impairs T cell receptor-
CD3 complex stability and signaling. *Immunity* **26**, 357–369 (2007).

25.   Y. He *et al.*, Identification of the Docking Site for CD3 on the T Cell Receptor beta Chain
by Solution NMR. *J Biol Chem* **290**, 19796–19805 (2015).

26.   A. Natarajan *et al.*, Structural Model of the Extracellular Assembly of the TCR-CD3
Complex. *Cell Rep* **14**, 2833–2845 (2016).

27.   A. Natarajan *et al.*, In situ cell-surface conformation of the TCR-CD3 signaling complex.
EMBO reports **25**, 5719–5742–5742 (2024).

28. D. Dong *et al.*, Structural basis of assembly of the human T cell receptor-CD3 complex.
Nature **573**, 546–552 (2019).

29. L. Sušac *et al.*, Structure of a fully assembled tumor-specific T cell receptor ligated by
pMHC. *Cell* **185**, 3201–3213.e3219 (2022).

30. K. Saotome *et al.*, Structural analysis of cancer-relevant TCR-CD3 and peptide-MHC
complexes by cryoEM. *Nature Communications* **14**, 2401 (2023).

31. Y. Chen *et al.*, Cholesterol inhibits TCR signaling by directly restricting TCR-CD3 core
tunnel motility. *Mol Cell* **82**, 1278–1287 e1275 (2022).

32. K. Natarajan *et al.*, An allosteric site in the T-cell receptor Cbeta domain plays a critical
signalling role. *Nat Commun* **8**, 15260 (2017).

33. S. Rangarajan *et al.*, Peptide-MHC (pMHC) binding to a human antiviral T cell receptor
induces long-range allosteric communication between pMHC- and CD3-binding sites. *J*
Biol Chem **293**, 15991–16005 (2018).

34. B. Knapp, P. A. van der Merwe, O. Dushek, C. M. Deane, MHC binding affects the
dynamics of different T-cell receptors in different ways. *PLoS Comput Biol* **15**, e1007338
(2019).

35. R. Q. Notti *et al.*, The resting and ligand-bound states of the membrane-embedded human
T-cell receptor–CD3 complex. *Nature Communications* **16**, 10996 (2025).

36. B. Alarcon, D. Gil, P. Delgado, W. W. Schamel, Initiation of TCR signaling: regulation
within CD3 dimers. *Immunol Rev* **191**, 38–46 (2003).

37. P. A. van der Merwe, O. Dushek, Mechanisms for T cell receptor triggering. *Nat Rev*
Immunol **11**, 47–55 (2011).

38. D. Moogk, A. Natarajan, M. Krogsgaard, T cell receptor signal transduction: affinity, force
and conformational change. *Curr Opin Chem Eng* **19**, 43–50 (2018).

39. C. Zhu, W. Chen, J. Lou, W. Rittase, K. Li, Mechanosensing through immunoreceptors.
Nat Immunol **20**, 1269–1278 (2019).

40. D. K. Das *et al.*, Force-dependent transition in the T-cell receptor beta-subunit
allosterically regulates peptide discrimination and pMHC bond lifetime. *Proc Natl Acad*
Sci U S A **112**, 1517–1522 (2015).

41. P. Wu *et al.*, Mechano-regulation of Peptide-MHC Class I Conformations Determines TCR
Antigen Recognition. *Mol Cell* **73**, 1015–1027 e1017 (2019).

42. H.-K. Choi *et al.*, Catch bond models may explain how force amplifies TCR signaling and
antigen discrimination. *Nature Communications* **14**, 2616 (2023).

43.   B. Liu, W. Chen, B. D. Evavold, C. Zhu, Accumulation of dynamic catch bonds between
1147   TCR and agonist peptide-MHC triggers T cell signaling. *Cell* **157**, 357–368 (2014).

1148   44.   J. Hong *et al.*, Force-Regulated In Situ TCR-Peptide-Bound MHC Class II Kinetics
1149   Determine Functions of CD4+ T Cells. *J Immunol* **195**, 3557–3564 (2015).

1150   45.   L. V. Sibener *et al.*, Isolation of a Structural Mechanism for Uncoupling T Cell Receptor
1151   Signaling from Peptide-MHC Binding. *Cell* **174**, 672–687 e627 (2018).

1152   46.   M. N. Rushdi *et al.*, Cooperative binding of T cell receptor and CD4 to peptide-MHC
1153   enhances antigen sensitivity. *Nat Commun* **13**, 7055 (2022).

1154   47.   S. T. Kim *et al.*, The alphabeta T cell receptor is an anisotropic mechanosensor. *J Biol
1155   Chem* **284**, 31028–31037 (2009).

1156   48.   S. Pryshchep, V. I. Zarnitsyna, J. Hong, B. D. Evavold, C. Zhu, Accumulation of serial
1157   forces on TCR and CD8 frequently applied by agonist antigenic peptides embedded in
1158   MHC molecules triggers calcium in T cells. *J Immunol* **193**, 68–76 (2014).

1159   49.   Y. Feng *et al.*, Mechanosensing drives acuity of alphabeta T-cell recognition. *Proc Natl
1160   Acad Sci U S A* **114**, E8204–E8213 (2017).

1161   50.   J. Hong *et al.*, A TCR mechanotransduction signaling loop induces negative selection in
1162   the thymus. *Nat Immunol* **19**, 1379–1390 (2018).

1163   51.   A. Akitsu *et al.*, Parsing digital or analog TCR performance through piconewton forces.
1164   *Science Advances* **10**, eado4313 (2024).

1165   52.   W. Chen, C. Zhu, Mechanical regulation of T-cell functions. *Immunological Reviews* **256**,
1166   160–176 (2013).

1167   53.   K. H. Hu, M. J. Butte, T cell activation requires force generation. *Journal of Cell Biology*
1168   **213**, 535–542 (2016).

1169   54.   J. Gohring *et al.*, Temporal analysis of T-cell receptor-imposed forces via quantitative
1170   single molecule FRET measurements. *Nat Commun* **12**, 2502 (2021).

1171   55.   D. L. Harrison, Y. Fang, J. Huang, T-Cell Mechanobiology: Force Sensation, Potentiation,
1172   and Translation. *Frontiers in Physics* **Volume 7 - 2019**, (2019).

1173   56.   J. Wang *et al.*, Atomic structure of an alphabeta T cell receptor (TCR) heterodimer in
1174   complex with an anti-TCR fab fragment derived from a mitogenic antibody. *EMBO J* **17**,
1175   10–26 (1998).

1176   57.   Z.-Y. J. Sun *et al.*, Solution structure of the CD3εδ ectodomain and comparison with
1177   CD3εγ as a basis for modeling T cell receptor topology and signaling. *Proceedings of the
1178   National Academy of Sciences* **101**, 16867–16872 (2004).

1179   58.   W. Wang *et al.*, Quantitative Analysis of T Cell Receptor Complex Interaction Sites Using
1180   Genetically Encoded Photo-Cross-Linkers. *ACS Chemical Biology* **9**, 2165–2172 (2014).

1181   59.   S. E. Chesla, P. Selvaraj, C. Zhu, Measuring two-dimensional receptor-ligand binding
1182   kinetics by micropipette. *Biophys J* **75**, 1553–1572 (1998).

1183   60.   J. Huang *et al.*, The kinetics of two-dimensional TCR and pMHC interactions determine
1184   T-cell responsiveness. *Nature* **464**, 932–936 (2010).

1185   61.   E. Evans, K. Ritchie, R. Merkel, Sensitive force technique to probe molecular adhesion
1186   and structural linkages at biological interfaces. *Biophys J* **68**, 2580–2587 (1995).

1187   62.   W. Chen, E. A. Evans, R. P. McEver, C. Zhu, Monitoring receptor-ligand interactions
1188   between surfaces by thermal fluctuations. *Biophys J* **94**, 694–701 (2008).

1189   63.   E. W. Newell *et al.*, Structural basis of specificity and cross-reactivity in T cell receptors
1190   specific for cytochrome c-I-E(k). *J Immunol* **186**, 5823–5832 (2011).

1191   64.   T. E. Williams, S. Nagarajan, P. Selvaraj, C. Zhu, Quantifying the Impact of Membrane
1192   Microtopology on Effective Two-dimensional Affinity*210. *Journal of Biological
1193   Chemistry* **276**, 13283–13288 (2001).

1194   65.   W. Chen, V. I. Zarnitsyna, K. K. Sarangapani, J. Huang, C. Zhu, Measuring Receptor–
1195   Ligand Binding Kinetics on Cell Surfaces: From Adhesion Frequency to Thermal
1196   Fluctuation Methods. *Cellular and Molecular Bioengineering* **1**, 276–288 (2008).

1197   66.   B. Liu *et al.*, 2D TCR-pMHC-CD8 kinetics determines T-cell responses in a self-antigen-
1198   specific TCR system. *Eur J Immunol* **44**, 239–250 (2014).

1199   67.   X. Zhao *et al.*, Tuning T cell receptor sensitivity through catch bond engineering. *Science*
1200   **376**, eabl5282 (2022).

1201   68.   R. Qin *et al.*, TCR catch bonds nonlinearly control CD8 cooperation to shape T cell
1202   specificity. *Cell Research* **35**, 265–283 (2025).

1203   69.   C. Zhu, T. E. Williams, Modeling concurrent binding of multiple molecular species in cell
1204   adhesion. *Biophys J* **79**, 1850–1857 (2000).

1205   70.   T. E. Williams, P. Selvaraj, C. Zhu, Concurrent binding to multiple ligands: kinetic rates
1206   of CD16b for membrane-bound IgG1 and IgG2. *Biophys J* **79**, 1858–1866 (2000).

1207   71.   M. Karplus, G. A. Petsko, Molecular dynamics simulations in biology. *Nature* **347**, 631–
1208   639 (1990).

1209   72.   S. Izrailev *et al.*, in *Computational molecular dynamics: challenges, methods, ideas*.
1210   (Springer, 1999), pp. 39–65.

1211   73.   E. M. M. Manders, J. Stap, G. J. Brakenhoff, R. V. Driel, J. A. Aten, Dynamics of three-
1212   dimensional replication patterns during the s-phase, analysed by double labelling of dna
1213   and confocal microscopy. *Journal of Cell Science* **103**, 857–862 (1992).

1214   74.   N. Manolios, F. Letourneur, J. S. Bonifacino, R. D. Klausner, Pairwise, cooperative and
1215   inhibitory interactions describe the assembly and probable structure of the T-cell antigen
1216   receptor. *EMBO J* **10**, 1643–1651 (1991).

1217   75.   N. Jiang *et al.*, Two-Stage Cooperative T Cell Receptor-Peptide Major Histocompatibility
1218   Complex-CD8 Trimolecular Interactions Amplify Antigen Discrimination. *Immunity* **34**,
1219   13–23 (2011).

1220   76.   B. Liu *et al.*, The cellular environment regulates in situ kinetics of T-cell receptor
1221   interaction with peptide major histocompatibility complex. *European Journal of
1222   Immunology* **45**, 2099–2110 (2015).

1223   77.   K. Li *et al.*, PD-1 suppresses TCR-CD8 cooperativity during T-cell antigen recognition.
1224   *Nat Commun* **12**, 2746 (2021).

1225   78.   H. K. Choi, C. Zhu, Catch Bonds in Immunology. *Annu Rev Immunol* **43**, 641–666 (2025).

1226   79.   W. Xin *et al.*, Structures of human γδ T cell receptor–CD3 complex. *Nature* **630**, 222–229
1227   (2024).

1228   80.   B. S. Gully *et al.*, Structure of a fully assembled γδ T cell antigen receptor. *Nature* **634**,
1229   729–736 (2024).

1230   81.   M. Hoque *et al.*, Structural characterization of two γδ TCR/CD3 complexes. *Nature
1231   Communications* **16**, 318 (2025).

1232   82.   B. Alarcon, W. W. Schamel, Allosteric Changes Underlie the Outside-In Transmission of
1233   Activatory Signals in the TCR. *Immunological Reviews* **329**, e13438 (2025).

1234   83.   J. H. Wang, E. L. Reinherz, The structural basis of alphabeta T-lineage immune
1235   recognition: TCR docking topologies, mechanotransduction, and co-receptor function.
1236   *Immunol Rev* **250**, 102–119 (2012).

1237   84.   T. Sasada *et al.*, Involvement of the TCR Cβ FG Loop in Thymic Selection and T Cell
1238   Function. *Journal of Experimental Medicine* **195**, 1419–1431 (2002).

1239   85.   M. Touma *et al.*, The TCR Cβ FG Loop Regulates αβ T Cell Development1. *The Journal
1240   of Immunology* **176**, 6812–6823 (2006).

1241   86.   S. T. Kim *et al.*, TCR Mechanobiology: Torques and Tunable Structures Linked to Early
1242   T Cell Signaling. *Frontiers in Immunology* **3**, (2012).

1243   87.   H. Li *et al.*, Lipid-regulated phosphorylation hierarchy of the T cell receptor tyrosine
1244   motifs. *Mol Cell* **85**, 3694–3710.e3698 (2025).

88. W. W. Schamel, B. Alarcon, S. Minguet, The TCR is an allosterically regulated
macromolecular machinery changing its conformation while working. *Immunological*
Reviews **291**, 8–25 (2019).

89. L. Kjer-Nielsen *et al.*, Crystal structure of the human T cell receptor CD3 epsilon gamma
heterodimer complexed to the therapeutic mAb OKT3. *Proc Natl Acad Sci U S A* **101**,
7675–7680 (2004).

90. M. Arslan *et al.*, Effect of non-repetitive linker on in vitro and in vivo properties of an anti-
VEGF scFv. *Scientific Reports* **12**, 5449 (2022).

91. M. Krogsgaard *et al.*, Agonist/endogenous peptide-MHC heterodimers drive T cell
activation and sensitivity. *Nature* **434**, 238–243 (2005).

92. F. Letourneur, B. Malissen, Derivation of a T cell hybridoma variant deprived of functional
T cell receptor alpha and beta chain transcripts reveals a nonfunctional alpha-mRNA of
BW5147 origin. *Eur J Immunol* **19**, 2269–2274 (1989).

93. W. Thomas *et al.*, Catch-bond model derived from allostery explains force-activated
bacterial adhesion. *Biophys J* **90**, 753–764 (2006).

94. C. D. Buckley *et al.*, Cell adhesion. The minimal cadherin-catenin complex binds to actin
filaments under force. *Science* **346**, 1254211 (2014).

95. D. L. Huang, N. A. Bax, C. D. Buckley, W. I. Weis, A. R. Dunn, Vinculin forms a
directionally asymmetric catch bond with F-actin. *Science* **357**, 703–706 (2017).

96. G. I. Bell, Models for the specific adhesion of cells to cells. *Science* **200**, 618–627 (1978).

97. W. L. Jorgensen, J. Chandrasekhar, J. D. Madura, R. W. Impey, M. L. Klein, Comparison
of Simple Potential Functions for Simulating Liquid Water. *J Chem Phys* **79**, 926–935
(1983).

98. B. Hess, C. Kutzner, D. van der Spoel, E. Lindahl, GROMACS 4: Algorithms for Highly
Efficient, Load-Balanced, and Scalable Molecular Simulation. *J Chem Theory Comput* **4**,
435–447 (2008).

99. M. J. Abraham *et al.*, GROMACS: High performance molecular simulations through
multi-level parallelism from laptops to supercomputers. *SoftwareX* **1-2**, 19–25 (2015).

100. U. R. Shrestha *et al.*, Generation of the configurational ensemble of an intrinsically
disordered protein from unbiased molecular dynamics simulation. *Proc Natl Acad Sci U S*
A **116**, 20446–20452 (2019).

101. K. Lindorff-Larsen *et al.*, Improved side-chain torsion potentials for the Amber ff99SB
protein force field. *Proteins* **78**, 1950–1958 (2010).

102.   F. Eisenhaber, P. Lijnzaad, P. Argos, C. Sander, M. Scharf, The double cubic lattice
1279   method: Efficient approaches to numerical integration of surface area and volume and to
1280   dot surface contouring of molecular assemblies. *J Comput Chem* **16**, 273–284 (1995).

1281   103.   R. Huey, G. M. Morris, A. J. Olson, D. S. Goodsell, A semiempirical free energy force
1282   field with charge-based desolvation. *Journal of Computational Chemistry* **28**, 1145–1152
1283   (2007).
1284

Figure 1. 2D kinetic quantifications of TCRαβ–CD3 ECD *cis*-interactions.
(A, B) Photomicrographs of the micropipette adhesion frequency (MAF) assay (*A*) and the biomembrane force probe (BFP) assay (*B*), overlain with legends indicating the measurement types. The two micropipettes held the probe (*left*) and the target (*right*), respectively, with each presenting one of the interacting pair. The target image shown in (*B*) is a hybridoma cell expressing full TCR–CD3 complexes, but beads coated with purified protein ECDs were also used, depending on the experiment. The target was brought into repeated contact with the probe to enable and interrogate ligand–receptor interactions. The probe, which was either a pressurized RBC (*A*) or a glass bead attached to the apex of the pressurized RBC (*B*), served a dual role of ligand presenter and force transducer, detecting the binding events to evaluate adhesion probability (*A*) and measuring bond lifetime (*B*), respectively, as indicated in the legends (see Methods). **(C)** Effective 2D affinities (A_cK_a) of pairwise *cis*-interactions between the ECDs of TCRαβ (αβ), CD3δε (δε), and CD3γε (γε) measured by MAF. **(D)** 2D off-rates (k_{off}) of the indicated *cis*-interactions quantified by the thermal fluctuation assay using BFP. **(E)** Calculated effective 2D on-rates ($A_ck_{on} = A_cK_a \times k_{off}$) of the indicated *cis*-interactions. Data are presented as mean ± SEM in (*C*)-(*E*), with the 2D affinity, off-

rate, and on-rate of the *trans*-interaction between cell-surface 2B4 TCR and its cognate pMHC K5:IE^k (*gray*) also plotted in the corresponding panels for comparison. **(F)** Representative force traces of non-binding *vs.* binding events from the BFP force-clamp assay. The target, either a bead or a cell, was driven to approach the probe (~ 0 pN before contact), pressed against it for 1 s (~ − 35 pN), which resulted in either a non-binding (returning to ~ 0 pN) or a binding event (crossing 0 pN to a positive, tensile force) upon target retraction. The latter trajectory illustrates a lifetime event in which the ligand–receptor bond was clamped with a force of ~16 pN for ~0.5 s (*red*) until dissociation, indicated by a sudden drop in force to ~ 0 pN. **(G)** Force-dependent average lifetimes of pairwise αβ–δε, αβ–γε, and δε–γε *cis*-bonds (*points*, mean ± SEM of > 30 lifetime events per force bin) at zero force by thermal fluctuation assay (*closed symbols*) and at non-zero forces by force-clamp assay (*open symbols*). The overlaid smooth curves are the global fits of the two-state dissociation model (Supplementary Fig. 2; see Methods). In contrast to *cis*-bonds, the points and curve in *gray* denote the previously published force-lifetime results of the 2B4–K5:IE^k *trans*-interaction (*42*), which provides a direct comparison.

Figure 2. Experimental demonstration of cooperativity in *cis*-interaction of CD3γε, CD3δε, and TCRαβ ECDs.

(A) Experimental schemes. Purified protein ECDs were captured onto the RBC or bead surfaces through biotin-streptavidin (SA) coupling. N-terminally-biotinylated 2B4 hybrid TCRαβ was coated onto the probe (*left*), while C-terminally-biotinylated human CD3γε (*blue*), CD3δε (*orange*), or a 1:1 mixture of both (CD3m, *red*) was coupled onto the opposing target (*right*). These tail-to-head binding arrangements ensure preferred *cis*-interaction orientations for TCR–CD3 associations. **(B)** Adhesion frequency, P_a (*left ordinate*), and average bond number, $\langle n \rangle = -\ln(1 - P_a)$ (*right ordinate*), measured by the MAF assay. The same TCRαβ-bearing RBC was tested sequentially against three RBCs bearing, respectively, CD3γε, CD3δε, or CD3m in random order (*points connected by solid lines*). These CD3-bearing RBCs were from the same batch of streptavidinylated cells, which were then incubated with saturating amounts of CD3γε, CD3δε, and CD3m, respectively, to equalize site density coating. Each probe-target pair was tested for 30 cycles, with a 2-s contact time per cycle. The predicted bond number of dual-ligand RBCs (*dotted line*), $\langle \hat{n}_m \rangle$, is the sum of the bond numbers of TCRαβ interacting with CD3γε and CD3δε, separately. Thus, $\langle \hat{n}_m \rangle$ equals to the average of the bond numbers measured from the two RBCs bearing single CD3 species because the respective densities of CD3γε and CD3δε on the mixed ligand RBCs are half of the densities of CD3γε and CD3δε on the single-ligand RBCs under the assumption of concurrent and independent binding (see Methods). This no-cooperativity assumption is invalidated by the data, which clearly demonstrate that the actual measurements greatly outnumber the prediction (note that a log scale is used on the right ordinate). Several metrics were used to quantify cooperativity (*C*, *F*, and Fig. 3D, G), which is defined as the difference between the measurement and the prediction, normalized by the expected value assuming no cooperativity. **(C)** Positive 2D binding cooperativity of TCRαβ–CD3m *cis*-interaction.

Here, $\Delta\langle n_m\rangle/\langle\hat{n}_m\rangle$ (*points*) utilized the bond number in (*B*) as the cooperative metric. The case of no cooperation corresponds to a zero value, whereas a greater than zero value reveals 2D binding cooperativity. **(D)** Force-dependent average bond lifetime, $\tau(f)$, for TCRαβ–CD3m *cis*-interactions quantified using BFP. The predicted average bond lifetime of dual-species (*dotted line*), $\hat{\tau}_m$, is the weighted sum of the single-species force-lifetime curves of the αβ–δε and αβ–γε bimolecular bonds (replotted from Fig. 1G for comparison). The weights are fractions of bond formation based on the 2D binding affinities of single-ligand *cis*-interactions (Fig. 1C) and corresponding amounts (1:1 ratio). The overlaid smooth curves are the global fits of the two-state, two-pathway dissociation models (Supplementary Fig. 2A-I; see Methods). **(E)** The bond survival probability for the TCRαβ–CD3m *cis*-interactions derived from zero-force individual bond lifetimes (whose average is $\tau(0)$) measured from the thermal fluctuation assay. The bond survival probability curves of the αβ–δε and αβ–γε bimolecular bonds are shown for comparison (using pairwise lifetime data in Fig. 1D). **(F)** Positive bond lifetime cooperativity of dual-ligand *cis*-interaction, $\Delta\tau/\hat{\tau}_m$, using the bond lifetime in (*D*) as the cooperative metric. Again, the no cooperation case corresponds to a zero value, whereas a greater than zero value reveals cooperativity in force-dependent bond lifetime. In (*D*) and (*F*), data points are presented as mean ± SEM from > 30 lifetime-events per force bin; global-fitting force-lifetime and survival-probability curves (*D*, *E*, and *F*) are plotted with means (*solid line*) and standard deviations (*shaded area*). Significance in (*C*) and (*F*) was assessed using a one-sample t-test against zero.

Figure 3. Computational demonstration of cooperativity in *cis*-interaction of CD3γε, CD3δε, and TCRαβ ECDs.

(A) Initial structures for MD simulations. The ECD of CD3γε–TCRαβ–CD3δε trimolecular complex (*left*) was modeled based on the published cryo-EM structure (*28*), 6JXR. The two TCRαβ–CD3δε (*middle*) and TCRαβ–CD3γε (*right*) bimolecular structures were obtained by removing CD3γε and CD3δε from the trimer, respectively. **(B, C)** Contact areas and interaction energies for the dimers and trimer from CMD simulations. Mean ± SD of contact areas (*B*), defined by solvent accessible surface areas (*101*), and interaction energies (*C*), quantified using empirical free energy scores (*102*), were calculated from the TCRαβ–CD3δε (*blue*), TCRαβ–CD3γε (*orange*), and CD3γε–TCRαβ–CD3δε (*red*) structures (Supplementary Fig. 3B, C). Statistics were obtained by comparing any two groups using unpaired t-tests. **(D)** Cooperative metrics of the δε–αβ–γε trimeric complex in terms of contact areas (*left*) and free energy scores (*right*). The cooperative analyses were conducted similarly to that in Fig. 2C, but were based on contact areas and interaction energies in (*B*) and (*C*) rather than on bond numbers. **(E)** The minimum distances between TCRαβ and corresponding CD3δε (*blue*), CD3γε (*orange*), or both (*red*) *vs.* simulation time, calculated from representative trajectories of SMD-simulated TCR–CD3 ECD dissociation.

Initial molecular constructs of αβ–δε and αβ–γε dimeric assemblies and of δε–αβ–γε trimeric complex are indicated, where TCRαβ N-termini were pulled away from corresponding C-terminal anchored CD3s during a 175-pN force-clamping SMD simulations (*insets*). The sudden rise of the trajectory from the baseline signifies the time point of dissociation (*arrows*). **(F)** *In silico* times required for the complex to dissociate, $t(f)$, under constant forces of 100-200 pN from SMD simulations. The predicted time-to-dissociation of the γε–αβ–δε trimer (*dotted line*), $\hat{t}_{\delta\gamma}$, is the TCR dissociation time from both δε and γε, under the assumption that the two bimolecular bonds are in parallel, and their dissociations are independent and concurrent (see Methods). **(G)** Cooperative metric $\Delta t / \hat{t}_{\delta\gamma}$ *vs.* force of the δε–αβ–γε trimeric complex. The cooperative analyses were conducted similarly to that in Fig. 2F but were based on the SMD simulated time-to-dissociation in (*F*) rather than on experimentally measured bond lifetimes. Significance was assessed using one-sample t-tests against zero in (*D*) and (*G*).

Figure 4. Structural basis of force-dependent cooperative TCR–CD3 ECD *cis*-interactions.

(A) Dynamic "footprints" of αβ–γε (*left*) and αβ–δε (*right*) interfaces are defined by the CMD-derived occupancies, which were projected onto the light gray space-filling representations of TCR–CD3 ECDs. The *insets* depict the placements and orientations of γε (*orange*) and αβ–δε (*gray*), αβ–γε (*gray*) and δε (*blue*) from the far left to the far right: two sets of side-view images mirroring each other to allow CD3s to face outward. The respective γε (*left*) and δε (*right*) binding interfaces with αβ are exposed by flipping the CD3s 180° while keeping the rest fixed (before *transparent*; after *opaque*). The pseudo-color maps are coded for interface footprints in the trimer (*cyan*; only the unobscured areas are visible) as well as in dimers with γε (*magenta*) and δε (*yellow*), with intensity levels matching the occupancy values, ranging from 0% (white) to 100% (fully saturated). When comparing trimer *vs.* dimer, the unobscured *cyan* areas relative to the *magenta* and *yellow* areas represent the increased footprints of αβ with γε (*left two* panels) or δε (*right two* panels) from dimer structures upon assembly of the trimer structure by incorporating the missing CD3. When comparing γε *vs.* δε, the magenta area is larger than the yellow area, indicating a more stable αβ–γε than αβ–δε dimer. Moreover, the overlapping areas are smaller on the right (*cyan* + *magenta*) than on the left (*cyan* + *yellow*), both in absolute values and percentages, indicating smaller cooperative effects due to adding δε to αβ–γε than adding γε to αβ–δε in the trimer and demonstrating asymmetry with less contact adjustment of γε than δε after trimer assembly. **(B)** Quantification of colocalization of occupancies between trimer and dimers by Manders' overlap coefficient, defined by $\sum D_i T_i / (\sum D_i^2 \sum T_i^2)^{1/2}$ where i denotes the residue index of αβ–γε (*orange*) and αβ–δε (*blue*); D_i and T_i denote the i^{th} occupancy values of dimer and trimer, respectively, before and after *cis*-cooperative trimer assembly with the missing CD3. A Student t-test was used to assess significance based on three independent simulation repeats. **(C)** Interfacial stabilities of two ε-chains under 0 and 175 pN force, quantified by mean ± SEM of ε (*left*) or ε' (*right*) chains residues with non-zero occupancies (*points*) interacting with αβ calculated from CMD (0 pN, *open bars*) or SMD (175 pN, *hatched bars*) simulated γε–αβ–δε trimeric structure. Here, γ-associated ε-chain (ε) and δ-associated ε-chain (ε') are distinguished using the prime superscript. **(D)** Mean occupancies of γ (*green*), ε (*orange*), δ (*red*), or ε' (*blue*) chains interacting with αβ quantified by averaging non-zero occupancies calculated from CMD (without force, *x*-axis) or SMD (with force, *y*-axis) simulated αβ–γε and αβ–δε dimer structures (*arrow starting points*) or γε–αβ–δε trimer structure (*arrow ending points*). **(E)** Fluctuations of two ε-chains under 0 and 175 pN force, quantified by mean ± SEM of residue RMSFs of ε (*left*) or ε' (*right*) chains calculated from CMD (0 pN, *open bars*) or SMD (175 pN, *hatched bars*) simulated γε–αβ–δε trimeric structure. **(F)** Change of flexibilities in the TCR–CD3 complex upon pulling. The *inset* shows the top view of the γε–αβ–δε ECD structure under a 175 pN force. Differential RMSF profiles between SMD and CMD (*blue* for enhanced and *red* for suppressed fluctuations) are mapped onto the ribbon view of trimeric ECD. The residues' absolute ΔRMSF values are represented by the thickness of the protein backbone. Statistical significance levels were assessed by Welch's t-test assuming non-equivalent SD. **(G, H)** Critical interacting residues mapped onto the γε–αβ–δε interfaces showing the strong-interacting (*G*) and the force-sensitive (*H*) contacts.

The *inset* in (*G*) is the top-view of the γε–αβ–δε ECD structure. The same top-views of the full ECD assembly are shown in ribbon representation where tightly interacting residues are highlighted by sticks and colors. Only strong-interacting contact residues are shown in (*G*) using sidechains colored from *white* to *red*, based on their ΣOccupancy level. The force-sensitive residues with significant occupancy differences upon applying a 175 pN clamping force are shown in (*H*), with sidechains colored from *white* to *red* for increases and from *white* to *blue* for decreases in their ΔOccupancy level (see Methods).

Figure 5. Effects of mutating Cβ contacting residues on *cis*-interaction affinity and bond lifetime.

(A) TCRβ local segments with distinct force-responses. Residues in the CC' loop (*upper table*) and Helix 4 – F strand (*lower table*) of the Cβ domain both contact CD3 (indicated by ΣOccupancy) but respond differently to 175 pN clamping force (indicated by ΔOccupancy). **(B)** TCRαβ structure with double-mutants listed in (*A*) highlighted on the Cβ CC' loop and Helix 4-F strand, which contribute to the force-dependent cooperative TCR–CD3 *cis*-interaction according to (*A*). **(C)** Differential IL-2 cytokine productions of cells containing Cβ mutations. The hybridoma

expressing 2B4 TCR WT and MTs were stimulated by K5:IE^k displayed on CHO cells. The area under the dose-response curve (AUC), normalized by that of WT, quantifies the relative IL-2 secretion. This data has previously been reported (*26*) and is reformulated here to facilitate experimental design and correlative analysis. **(D)** Normalized average number of bonds of *cis*-interactions between CD3δε (*first group*), CD3γε (*second group*), or CD3δε–CD3γε (*third group*) and TCRαβ WT (*gray*) or β-chain double-mutants NP (*green*), GK (*purple*), or NH (*magenta*) MTs. Corresponding $\langle n \rangle / m_l m_r$ of WT and indicated mutant 2B4 TCRs on hybridomas *trans*-interacting with K5:I-E^k are plotted for comparison (*fourth group*). **(E-G)** Force-dependent bond lifetimes (*points*, mean ± SEM of > 30 lifetime events per force bin; *curves with shades*, mean ± SD of globally fitted bond profiles) of *cis*-interactions of CD3γε (*E*), CD3δε (*F*), or both (*G*) with TCRαβ WT (*gray*), NP (*green*), GK (*purple*), or NH (*magenta*) MTs. **(H)** Corresponding $\tau(f)$ of WT and indicated mutant 2B4 TCRs on hybridomas *trans*-interacting with K5:I-E^k are shown for comparison. This data has previously been reported (*42*) and is replotted here to facilitate correlative analysis and mechanistic understanding.

Figure 6. Correlating ECD *cis*-interaction parameters of TCRαβ with CD3γε, CD3δε, or both with T cell function and *trans*-interaction parameters between TCR and pMHC.

(A) The correlations between the IL-2 secretion (in Fig. 5C) and effective 2D affinity for αβ–γε (*orange*) or αβ–δε (*blue*) *cis*-interaction (in Fig. 4C, *first* and *second group*). **(B)** The correlations between IL-2 secretion and zero-force (*dotted lines*) *vs.* optimal-force (*solid lines*) bond lifetime for αβ–γε (*orange*) or αβ–δε (*blue*) *cis*-interaction (in Fig. 5E, F). *Inset*: Different catchiness and sensitivity of CD3γε (*orange*) *vs.* CD3δε (*blue*). The arrows display the effects of force. **(C)** Correlations between normalized γε–δε–αβ *cis*-kinetics and T-cell signal. The mean ± SEMs of relative IL-2 AUC (in Fig. 5C) are negatively correlated with the peak lifetimes of ECD *cis*-interactions of WT and MT TCRαβs (in Fig. 5G) when both CD3γε, CD3δε co-presented (*black*). The poor correlations between IL-2 production and zero-force bond lifetime (*red dashed line*) or normalized number of bonds (*red solid line*) for γε–δε–αβ *cis*-interactions are also plotted for comparison. **(D)** Goodness-of-fit and sensitivity of force-dependent bond lifetimes of γε–δε *cis*-interacting with WT and MT TCRαβs in predicting downstream IL-2 responses. The Pearson correlation coefficients (*black*) and inverse of the linear-regression slopes (*red*) between

normalized IL-2 AUC and γε–δε–αβ *cis*-catch bonds are plotted for a range of forces (*curves with shades*, mean ± SD). (**E**) Correlations between γε–δε–αβ *cis*- and *trans*-bond lifetimes. The mean ± SEMs of the peak lifetimes of trimolecular *cis*-interactions (in Fig. 5G) are negatively correlated with those of *trans*-interactions from WT and MT 2B4 TCRs expressed on hybridoma cells interacting with K5:IE^k (in Fig. 5H). The poor correlations between the peak lifetimes of bimolecular *cis*-interactions (*orange*, αβ–γε and *blue*, αβ–δε) with the peak lifetimes of the WT and MT 2B4 TCRs *trans*-interacting with K5:IE^k are plotted for comparison (in Fig. 5E, F). (**F**) Anti-correlation between the γε–δε–αβ *cis*-interaction and the TCR–pMHC *trans*-interaction for WT and MT. The Pearson correlation coefficients of the *cis*- *vs.* *trans*-bond lifetime are shown for every single magnitude of force (*curves with shades*, mean ± SD).