High genetic diversity in the pelagic deep-sea fauna of the Atacama

Trench revealed by environmental DNA

Salvador Ramírez-Flandes[1,2,*], Carolina E. González[2], Montserrat Aldunate[2], Julie Poulain[3],
Patrick Wincker[3,4,5], Ronnie N. Glud[6,7], Rubén Escribano[1,2], Sophie Arnaud Haond[8] and
Osvaldo Ulloa[1,2,*]
[1] Departmento de Oceanografía, Universidad de Concepción, 4070386 Concepción, Chile
[2] Instituto Milenio de Oceanografía (IMO), Universidad de Concepción, 4070386 Concepción,
Chile
[3] Commissariat à l'Energie Atomique (CEA), Institut de Génomique (IG), Genoscope, 2 rue
Gaston Crémieux, BP5706, 91057 Evry, France
[4] Centre National de Recherche Scientifique (CNRS), UMR 8030, CP5706, Evry, France
[5] Université d'Evry, UMR 8030, CP5706, Evry, France
[6] Danish Center for Hadal Research, Danish Institute of advanced Science, and Nordcee,
University of Southern Denmark, 5230 Odense M, Denmark
[7] Tokyo University of Marine Science and Technology, 4-5-7 Konan, Minato-ku, Tokyo, 108-
8477, Japan
[8] MARBEC, Université Montpellier, Institut Français de Recherche pour L'Exploitation de la
Mer, CNRS, Institut de recherche pour le développement, 34203 Sète, France
*To whom correspondence should be addressed. Email: sram@udec.cl or oulloa@udec.cl

Statement of authorship: SRF, RE, SAH, and OU designed research. RNG provided access
to study locations, samplings platforms, and general logistics. MA and OU collected samples.
JP, PW, and SAH contributed to sample preparation, sequencing, and analytical tools. SRF
and CEG analyzed the data. SRF, CEG, RE, SAH, and OU interpreted the results. SRF and
OU wrote the first draft of the manuscript, and all authors contributed with revisions to the
final manuscript.

Data accessibility statement: NCBI bioproject PRJNA806167.

Keywords: metazoa, ecology, metabarcoding, Peru-Chile Trench, Kermadec Trench

Running title: High pelagic metazoan diversity in the Atacama Trench.

Abstract

A current paradigm in marine biodiversity states that faunal richness decreases with depth. However, the deep-ocean ecosystem has been significantly under-sampled, hindering a complete view of its biodiversity. This situation is accentuated in ultra-deep waters, where the remote and extreme conditions unfit the traditional sampling methods. Using environmental DNA, we assessed the pelagic metazoan diversity of the Atacama Trench from the high-productive near-surface level down to ~8000 m depth. Our results show that waters deeper than 4000 m contributed up to 50% of the overall genetic diversity. These findings contrast with similar observations in the less-productive Kermadec Trench, where the diversity in deep waters was lower than in shallower waters. Moreover, both deep pelagic ecosystems exhibited some unknown phylogenetic clades within the dominant taxonomic groups: hydrozoans and copepods. The deep-ocean biota may thus contribute to global biodiversity far more than hitherto suggested, especially in zones influenced by high primary production. Our results underline the need for increased effort to study these remote ecosystems and improve our understanding of their contribution to the ecology and biogeochemistry of the deep-sea pelagic and benthic realms.

Introduction

Knowledge of the global magnitude, geographic distribution, and biodiversity loss is essential to ecology and conservation biology. This need has dramatically enhanced the number of biogeographic investigations in the past decades (Gaston, 2000). Current estimates suggest that about two-thirds of the species on Earth have been described so far (Costello and Chaudhary, 2017). The number of species has been estimated at ~0.3 million in the ocean and ~1.7 million on land (including fresh-water ecosystems) (Costello and Chaudhary, 2017), yet the forecast of the remaining undescribed diversity is highly variable, particularly in the marine realm (Appeltans et al., 2012). These estimates are partially based on the fact that only ~16% of currently named species are marine (Grosberg et al., 2012), challenging the classical biogeographic model that relates habitat size to species richness (Costello and Chaudhary, 2017). However, less than 1% of the deep ocean has been explored (Higgs and Attrill, 2015), due primarily to the logistic and technological difficulties involved in sampling the vast and high-hydrostatic-pressure abyssal (4,000—6,000 m) and hadal (6,000—11,000 m) realms (Smith et al., 2009). Moreover, the deep sea is characterized by low biomass production and import, and it has long been expected to be scarcely populated with a handful of scavengers thriving in a cold and biomass-poor world (Ramirez-Llodra et al., 2010). Consequently, our knowledge of the faunal diversity below 1 km depth—almost 80% of the biosphere (Childress, 1995)—is elusive. Indeed, as our comprehension of the latitudinal gradient in marine biodiversity is biased by sampling effort (Chaudhary et al., 2016), that of the vertical distribution of marine biodiversity is also similarly biased. The deep-sea diversity has therefore been suggested to equal or even surpass that present in shallow waters (Costello and Chaudhary, 2017; Valentine and Jablonski, 2015), despite the hostile conditions of the deep sea, which is cold, dark, extremely diluted, and contains limited food sources (Jamieson et al., 2010; Priede and Froese, 2013). Conversely, other studies suggest that this comparatively unbalanced picture of species richness reflects an ecological truth rather than unbalanced sampling efforts—unless many rare and endemic species would be discovered in the deep sea or, for example, if marine parasites were far more diverse than presently documented (Costello and Chaudhary, 2017).

Moreover, hadal trenches sustain elevated benthic biological activity at extreme
hydrostatic pressure due to material focusing (Glud et al., 2013), but their pelagic
faunal realm remains unexplored.

In the last decade, marine ecologists have started to assess marine metazoan
biodiversity using methods based on the analysis of environmental DNA (eDNA)
(Deiner et al., 2017) to complement the traditional methods based on the sampling
and study of entire organisms. This molecular method is based on the fact that
organisms leave traces of their genetic material in the environment through
shedding and waste depositions (Taberlet et al., 2012). The use of eDNA allows the
detection of high levels of biodiversity with a reasonable amount of starting
material, thus offering a way to circumvent the problem of sampling the deep sea.
Since DNA degradation in the water column typically occurs within only a few days
(Andruszkiewicz Allan et al., 2021; Harrison et al., 2019; Thomsen et al., 2012),
this technique can allow estimates of the biodiversity present at the sampling time
or, at least, within a relatively short time window. Thus, eDNA-based tools have
emerged as a promising alternative for rapid, sensitive, and non-invasive
assessment of biodiversity (Harrison et al., 2019). However, to date, marine eDNA
metazoan biodiversity studies have mainly focused on the epipelagic zone (< 200 m
depth) and the ocean seafloor sediments (Cristescu and Hebert, 2018; Sinniger et
al., 2016), leaving unexplored the majority of the water column—from the
mesopelagic zone (200—1000 m) down to the abyssal and hadal zones (4,000—
11,000 m). Recently, the eDNA method has been applied at the basin scale to
assess the pelagic eukaryotic diversity, including the very abundant protists, down
to 5200 m, and compare it with that in the abyssal and bathyal surficial sediments
(Cordier et al., 2022). The composition of pelagic metazoan communities in the
bathypelagic or abyssal, let alone the hadal, realm thus remains poorly known
despite their potential importance for the stability of ecosystems, as for models for
climate change and conservation policies. Here we used eDNA metabarcoding of
two genetic markers, nuclear 18S rRNA and mitochondrial cytochrome c oxidase I
genes, henceforth, 18S and COI, to assess the metazoan biodiversity in the water
column of the Atacama Trench—from 85 m to more than 8000 m depth—and in the

abyssal and hadal waters of the Kermadec Trench (Atacama and Kermadec,
hereinafter; Fig. 1 and Table S1).

Results

The analysis of the combined data (12,093,326 and 11,551,435 18S and COI
metabarcoding sequences, respectively) resulted in 497 metazoan 18S Amplicon
Sequence Variants (ASVs) and 307 COI translated ASVs (tASVs). The taxonomic
diversity of these sequences was mainly associated with the cnidarian class
Hydrozoa and the crustacean subclass Copepoda throughout the water column,
while tunicates, annelids, scyphozoans, placozoans, and sponges were also
prominent in some samples (Figs. 1 and S1; Table S2).

Figure 1. Geographic setting of the sampling of eDNA data in the Kermadec and Atacama
Trenches (left) and taxonomic diversity distribution (presence/absence) of the Amplicon
Sequence Variants (ASVs) of the 18S rRNA gene metabarcoding (right). The names of the
samples are split into three parts, encoding the name of the cruise/station, the water filter
size range ("02" for the range 0.2-3 µm, and "3" for the range 3-20 µm), and the depth
(meters). Detailed information on the samples can be found in Table S1.

The overall genetic richness distribution of 18S ASVs in the Atacama Trench was

slightly higher in the abysso-hadal waters (>4000 m) than in shallower waters and

markedly higher compared to the abysso-hadal waters of the Kermadec Trench

(Figs. S4 and S5). In Atacama, the metazoan diversity of ASVs was distributed

among ~10 metazoan phyla. The majority of these ASVs were found in waters >

4000 m, according to the 18S and COI metabarcoding data (Tables 1 and S2,

respectively). An evaluation of the contribution of each depth level to the overall

water-column genetic richness revealed that the abysso-hadal waters of the

Atacama Trench region harbor ~50% of the total amount of detected ASVs,

suggesting a high level of endemism (Fig. 2A and 2B). In contrast, only <20% of

the ASVs found in the abysso-hadal samples of Kermadec were not observed at 500

m. Moreover, the percentages of exclusive ASVs per depth followed a similar

pattern (Fig. 2C), and consequently, the genetic diversity was observed to be more

equally distributed with depth in Atacama than in Kermadec (Fig. 2d).

Phylum	Aquatic Biomes	General Biomes	<4000m	>4000m	Ubiq.	Total
Annelida (mostly polychaetes)	Pelagic/Benthic	Marine/Freshwater/Terrestrial	1	21	7	29
Arthropoda (mostly copepods)	Pelagic/Benthic	Marine/Freshwater/Terrestrial	38	65	42	145
Brachiopoda	Benthic	Marine	0	0	0	0
Bryozoa	Pelagic/Benthic	Marine/Freshwater	0	0	0	0
Chaetognatha	Pelagic/Benthic	Marine	0	13	4	17
Chordata (mostly tunicates)	Pelagic/Benthic	Marine/Freshwater/Terrestrial	8	18	40	66
Cnidaria (mostly hydrozoans)	Pelagic/Benthic	Marine/Freshwater	21	41	66	128
Ctenophora	Pelagic/Benthic	Marine/Freshwater	0	3	1	4
Cycliophora	Benthic	Marine	0	0	0	0
Echinodermata	Benthic	Marine	1	4	0	5
Entoprocta	Pelagic/Benthic	Marine/Freshwater	0	0	0	0
Gastrotricha	Benthic	Marine/Freshwater	0	0	0	0
Gnathostomulida	Benthic	Marine	0	0	0	0
Hemichordata	Pelagic/Benthic	Marine	0	3	0	3
Micrognathozoa	Pelagic/Benthic	Freshwater	0	0	0	0
Mollusca	Pelagic/Benthic	Marine/Freshwater/Terrestrial	0	5	1	6
Nematoda	Benthic	Marine/Freshwater/Terrestrial	0	0	0	0
Nemertea	Benthic	Marine/Freshwater/Terrestrial	0	0	0	0
Onychophora	NA	Terrestrial	0	0	0	0
Phronida	Benthic	Marine	0	0	0	0
Placozoa	Pelagic/Benthic	Marine	0	0	0	0
Platyhelminthes	Pelagic/Benthic	Marine	0	0	0	0
Porifera (sponges)	Pelagic/Benthic	Marine/Freshwater	0	3	1	4
Priapulida	Benthic	Marine	0	0	0	0
Syndermata	Pelagic/Benthic	Marine/Freshwater	0	0	0	0
Tardigrada	Pelagic/Benthic	Marine/Freshwater/Terrestrial	0	0	0	0
Xenacoelomorpha	Pelagic/Benthic	Marine	0	0	0	0
Total			**69**	**176**	**162**	**407**

146 **Table 1**. Counts of 18S ASVs per metazoan phyla obtained in Atacama above, below, and
147 above-and-below 4000 m (fourth, fifth, and sixth column, respectively). The second and
148 third columns show the biomes documented for each corresponding phylum. A taxonomic
149 affiliation of the ASVs associated with all the groups can be found in Table S3 for Atacama
150 and Table S4 for Kermadec. Table S2 displays similar information for the COI tASVs.

Figure 2. a) Compositional coverage of total metazoan ASVs per sample (excluding those obtained with the filter 3-20 µm for Atacama as this filter was not used in Kermadec) shows that the faunal genetic diversity is more uniformly distributed in the water column of Atacama than in Kermadec. For example, at 500 m, the Kermadec system already covered 50% of its metazoan diversity, while in Atacama this was achieved only at abysso-hadal depths. **b)** The per-sample contribution of ASVs in the water columns of Atacama and Kermadec also shows a different pattern, in which samples from the abysso-hadal pelagic system contribute more to the metazoan biodiversity in Atacama than in Kermadec. In this case, the bar lengths represent the number of ASVs of the 18S genetic marker. **c)** Proportion of shared ASVs vs. exclusive number of ASVs per sample. The circles' sizes represent the total number of ASVs per sample, with the gray slice representing the shared ASVs, and the colored slices depicting the exclusive ASVs (those only present in the corresponding sample.) **d)** The Gini coefficient represents degree of inequality in the distribution of the richness of ASVs between the samples from Atacama and Kermadec. The total and exclusive richness of ASVs was more equally distributed in Atacama than in Kermadec, as this coefficient expresses greater equality with values closer to zero.

The phylogenetic relationships of the 18S ASVs associated with the taxonomic
groups with the highest genetic diversity—namely, Hydrozoa and Copepoda (Figs.
1, S1, S4, and S5)—with the closest related sequences present in Genbank,
revealed multiple well-supported (>95% of probability) clades exclusively
composed of ASVs observed in this study (Figs. 3, S6 and S7). One of these clades
(Hyd_Isedis1, Fig. 3a) contained ASVs exclusively detected in Atacama that could
not be placed within any recognized hydrozoan order, suggesting that it might
represent a novel *incertae sedis* clade within the Hydroidolina subclass. Other such
putative novel hydrozoan clades exclusively comprising ASVs recovered from
Atacama were found within Siphonophorae (Hyd_siph1 and Hyd_siph2, Fig. S6a),
Rhopalonematidae (Hyd_rhop1, Fig. S6b), and Geryonidae (Hyd_gery1, Fig. S6c). A
putative novel hydrozoan clade exclusively composed of ASVs from Kermadec was
found within Clytiidae (Hyd_clyt1, Fig. S6d). Also, novel copepod clades with an
exclusive composition of ASVs recovered from Atacama were found within
Oncaeidae (Cop_onca1 and Cop_onca2, Fig. S7a), Clausocalanidae (Cop_clau1, Fig.
S7b) and Calanidae (Cop_cala1 and Cop_cala2, Fig. S7c).

Figure 3. Phylogenetic trees of the 18S ASVs associated with Hydrozoa (a) and Copepoda
(b). The whole set of reference sequences utilized for these trees can be found in Table S6
and Table S7 for Hydrozoa and Copepoda, respectively. The colored clades in which the
ASVs generated exclusive groups are expanded in Figs. S6 and S7, for Hydrozoa and
Copepoda, respectively.
192

The diversity of the 18S ASVs within Hydrozoa and Copepoda was also notably
higher in Atacama than in Kermadec (Figs. S2 and S3). Regarding Hydrozoa,
although the taxonomic diversity and abundance of eDNA sequences of
siphonophores (pelagic colonial hydrozoans) was predominantly observed in
Atacama and Kermadec, the overall taxonomic diversity at the family level was
richer in Atacama than in Kermadec (Figs. S2a and S3a). Moreover, a significant
proportion of eDNA sequences associated with calycophoran siphonophores were
replaced by physonect siphonophores at hadal depths (Fig. S2a and S3a).

Concerning copepods, calanoids of the families Paracalanidae, Calocalanidae, and
Eucalanidae prevailed over other calanoid and podoplean groups at most depths
(Fig. S2b). In Kermadec, the most prominent calanoid groups were Lucicutiidae,
Clausocalanidae, and Mecynoceridae, along with a strong dominance of
harpacticoids affiliated with Pseudotachidiidae, persistently observed in the two
deepest samples (Figs. S2b and S3b). Pseudotachidiids have been reported in
several abyssal ecosystems (Kim et al., 2019) and geographically diverse deep-sea
sediments (Sinniger et al., 2016), but our study did not detect them in Atacama.
Notably, the eDNA signature of the copepod community in Atacama at 500 m depth
was remarkably different from the signature of those in any other sample of this
work. This sample came from the lower limit of the Anoxic Marine Zone (AMZ)
situated in the eastern South Pacific Ocean (Ulloa et al., 2012)—an oceanographic
feature absent in the Kermadec Trench region. This eDNA AMZ signature pointed to
metridinids and oncaeids, whose adaptations to low-oxygen conditions—most likely
to avoid predation from organisms with higher oxygen demands— have been
described in other hypoxic marine ecosystems (Böttger-Schnack et al., 2008;
Teuber et al., 2013).

Ctenophores (comb jellies) and Tunicates were also conspicuous in Atacama and
Kermadec (Fig. 1). The phylogenetic analysis associated with ctenophores did not
form well-supported clades of ASVs, as it did in the case of tunicates. This analysis
classified the ASVs of tunicates within the Appendicularia and Thaliacea (pelagic
tunicates) classes and none within the Ascidacea (benthic tunicates) class. In the
Appendicularia class, two clades exclusively composed of ASVs from this study were
grouped (Tun_App1 and Tun_App2), and several ASVs remained unattached to any
known family within this class (Fig. S8). The rest of the taxonomic groups mainly
included, among others, polychaetes in Annelida; decapods and euphausiids in
Crustacea; saggitoideans in Chaethognatha; fishes from the families Atherinidae,
Carangidae, Gobiidae, and Sebastidae within chordates from Actinopteri; and
murex snails, tulip snails, and sea butterflies in Mollusca (Tables S3 and S4).

Discussion
The coarse-level taxonomic pattern of the metazoan diversity detected in the
Atacama and Kermadec trenches (Figs. 1 and S1) is consistent with those reported
for other related pelagic ecosystems, which were based on studies conducted on
entire organisms collected with net tows down to 4000 m depth (de Puelles et al.,
2019; Kosobokova and Hirche, 2000; Vereshchaka et al., 2017) or inferred from
molecular taxonomic markers of epi-mesopelagic (< 1000 m) marine snow
(Lundgreen et al., 2019). However, the distribution of the genetic richness of the
18S ASVs in the Atacama Trench was slightly higher in the abysso-hadal waters
(>4000 m) compared to shallower waters and markedly higher compared to the
abysso-hadal waters of the Kermadec Trench (Figs. S4 and S5). This result contrast
with results reported down to mesopelagic depths in other regions using traditional
methods (Costello and Chaudhary, 2017; Jamieson et al., 2010), which generally
have shown that the faunal richness decreases with ocean depth—despite never
having scrutinized metazoan diversity in waters deeper than 5000 m before.

The abysso-hadal realm represents a significant fraction of the biosphere, yet it is
the least-explored environment on Earth (Robison, 2009). This biome is ecologically
relevant as it sustains a crucial pelagic food web with a significant role in energy
export to the ocean's sediments (Steinberg and Landry, 2017). The traditional

techniques for sampling marine metazoan organisms require specialized sampling
gear and lengthy, time-consuming sorting and taxonomic identifications by scarce
experts. They are thus, by themselves, insufficient to cover the vast ocean.
Recently, new eDNA-based methods have been used in deep ocean sediments and
pelagic waters studies. These methods complement the traditional methods as they
provide a rapid and more extensive way to assess biodiversity in these vast and
remote ecosystems.

To this end, our results highlight two main points. First, hydrozoans and copepods
dominate the metazoan diversity throughout these ultra-deep water columns. The
predominance of hydrozoan siphonophores in the hadopelagic waters of the
Atacama Trench was an unexpected result, as these animals were not, thus far,
considered as characteristic of these ecosystems (Jamieson, 2015). This result
should be interpreted with care, though, as mass jelly die-offs—called jelly-falls—
significantly contribute to the pelagic particulate matter that ends on the seabed
(Lebrato et al., 2012; Luo et al., 2020). However, the persistence of this signature
across sites spanning several kilometers and the distinct type of siphonophores
found in hadal (as compared to abyssal) waters in Atacama (Figs. S2a and S3a)
make it more likely that these eDNA signatures come from indigenous organisms.
Moreover, a significant proportion of eDNA sequences associated with calycophoran
siphonophores were replaced by physonect siphonophores at hadal depths (Fig. S2a
and S3a). Unlike calycophores, physonect siphonophores possess pneumatophores,
which provide their colonies with improved motor skills (Castellani and Edwards,
2017). Other prominent hydrozoans were trachymedusae affiliated with the
Rhopalonematidae and Geryoniidae families, whose eDNA sequences generated
multiple well-supported phylogenetic clades of ASVs (Fig. 3a), suggesting the
presence of several undescribed high-rank taxonomic groups.

Secondly, abyssal and hadal waters in systems like Atacama can contribute to a
significant proportion of the overall metazoan genetic diversity in the water column,
suggesting the existence of high diversity and endemism below 4000 m depth. The
main characteristic of this ecosystem is its high primary production in near-surface
waters (Daneri et al., 2000), which might promote a habitat complexity capable of

contributing to the development of low-oxygen zones (Ulloa et al., 2012) and
making the whole water-column eutrophic (Danovaro et al., 2002), thus promoting
biodiversity (Clarke and Gaston, 2006; Grosberg et al., 2012; Vereshchaka et al.,
2017).

The models for climate change predictions need to incorporate the diversity and
biomass of all the trophic levels in the water column, yet there is a large gap in
knowledge about the contribution of metazoans to abyssal and hadal pelagic
environments. Diversity promotes stability in ecosystems (Ives and Carpenter,
2007), and thus it represents necessary information that should guide the
assessment of the impact of environmental perturbations. This information would
improve the output of the ecological models and, consequently, conservation
policies. For instance, a recent study suggests that jellies biomass should be
included to improve the accuracy of biogeochemical models for carbon estimations
(Luo et al., 2020). Moreover, a biodiversity loss in the deep ocean might be
associated with an exponential reduction of its ecosystem functions (Danovaro et
al., 2008).

The output of any eDNA study depends on the geographical coverage and the
primers' specificity for DNA amplification, as some natural taxonomic groups might
have evolved molecular sequences incompatible with the primers. Admittedly, our
findings are based on a limited number of hard-to-get samples and a methodology
that may still be giving an incomplete view of the actual metazoan diversity present
in the environment. Also, the completeness of the available sequence databases
severely constrains this view. Nonetheless, our results provide the first view of the
diversity and distribution of ultra-deep pelagic fauna and suggest that the deep-
ocean biota might contribute to global biodiversity much more than what is
calculated in present-day estimates, especially in general zones of increased habitat
complexity due to the influence of high primary production. These results also
highlight the potential of eDNA-methods to explore marine biodiversity in the three
dimensions of the world's ocean and are consistent with previous sediment
observations for the Atacama Trench as a hotspot of faunal diversity (Danovaro et
al., 2002). Furthermore, they stress the need for developing and improving

standardized approaches to improve our understanding of the ecology of this vast
and still underexplored biome.

Material and Methods

Collecting samples: Seawater samples were collected during three oceanographic
cruises (TAN1711, ATACAMEX and SO261) conducted in the Kermadec and
Atacama trenches of the South Pacific Ocean. The TAN1711 with RV *Tangaroa*
cruise (from November 24th to December 14th, 2017) covered a transect from New
Zealand (41°18'S, 174°47'W) to the Northern Kermadec Trench (21°01'S,
175°12'W), involving four oceanographic stations. Both the ATACAMEX cruise with
the RV *Cabo de hornos* (from January 28th to February 4st, 2018) and SO261 cruise
with RV *Sonne* (March 2nd to April 2nd, 2018) were carried out in the deepest region
of the trench (20°19'S — 24°15'S off northern Chile, Fig. 1, Table S1). Each sample
was filtered across a hydrophilic 20-μm nylon membrane (Merck Millipore,
Massachusetts, USA) and filtered onto a Sterivex fileter using a peristaltic pump.
The membrane filter containing captured eDNA and cellular material from the water
column was filled with lysis buffer and stored at -80°C. **DNA extractions** were
carried out by Genoscope (Évry, France) using the same protocol as described by
Alberti et al. (2017) for Tara Oceans water samples. The protocol is based on the
cryogenic grinding of membrane filters, followed by nucleic acid extraction with
NucleoSpin RNA kits combined with the NucleoSpin DNA buffer set (Macherey–
Nagel, Düren, Germany). **PCR amplification and sequencing:** The 18S-V1V2
metabarcodes were generated using the SSUF04 and SSUR22mod primers. The COI
metabarcodes were generated using the mlCOIintF and jgHCO2198 primers. Paired-
end sequencing (2×250bp) of the libraries was performed on either the HiSeq 4000
or HiSeq 2500 instruments (Illumina, San Diego, CA, USA). **Sequence analysis:**
Paired-end reads were assembled and quality filtered with vsearch v2.17.1, using
parameter of maxEE at 0.5, minimum length of 250 bp, and a maximum length of
500 bp. The primers in the resulting metabarcoding sequences were removed with
pTrimmer v1.3.4. The sequences lacking the primers were discarded. The
sequences from all datasets were then put into a single file, and the unique
sequences were determined with vsearch. Chimeras were then removed with the
same program, and in the same process, the Amplicon Sequence Variants (ASVs)

were created. The 18S ASVs were compared as nucleotide sequences using BLASTN (parameters task dc-megablast, bitscore cutoff of 50) against the SILVA database (138.1) to construct the corresponding taxonomic profiles. The swarm clusters were created with swarm v3.0 with d=1 and fastidious parameters. The COI ASVs were translated into protein sequences (translated ASV or tASV), using the longest translation of the six open reading frames. These tASVs were aligned against a custom reference sequence database of COI sequences with BLASTP (bitscore cutoff of 50). **Phylogenetic analyses**: The phylogenetic trees of 18S ASVs and references sequences were constructed as follows. First, reference sequences for hydrozoans, copepods, and tunicates were collected from the literature and from respective searches in Genbank, generating separate datasets for each taxonomic group (Tables S5, S6, and S7). These datasets were complemented with the ASVs sequences associated with each taxonomic group according to the previously described procedure (comparison with the SILVA database). Second, these datasets of sequences were then aligned using mafft with parameters for high accuracy. The respective alignments were used to create phylogenetic trees with IQ-TREE2. Further details on these methods and corresponding references can be found in the Supplementary Material.

Acknowledgments

This work was supported the Chilean National Agency for Research and Development through the Millenium Science Initiative-ANID Program (grant ICN12 019-IMO) and grant Fondecyt 1191360 (to O.U.) and AUB17002 to Wolfgang Schneider. We thank the captains, crews, and scientific personnel of the RV *Sonne* (SO261; ship time provided by BMBF, Germany, awarded to Frank Wenzhoefer, Mathias Zabel, and Ronnie N. Glud), and RV *Tangaroa* (TAN1711; shiptime partly funded by the Coasts & Oceans Centre of New Zealand's National Institute of Water & Atmospheric Research -NIWA-, awarded to Ashley A. Rowden and Ronnie N. Glud), in the framework of the HADES-ERC Advanced grant "Benthic diagenesis and microbiology of hadal trenches"#669947 awarded to Ronnie N. Glud. Additional support was provided by the Danish National Research Foundation via the Danish Center for Hadal Reserch, grant number DNRF145. This work is part of the "Pourquoi Pas les Abysses?" project funded by Ifremer, and the project eDNAbyss

375 (AP2016 -228) funded by France Génomique (ANR-10-INBS-09) and Genoscope-
376 CEA.

References

Andruszkiewicz Allan E, Zhang WG, C. Lavery A, F. Govindarajan A. 2021.
Environmental DNA shedding and decay rates from diverse animal forms and
thermal regimes. *Environ DNA* **3**:492–514. doi:10.1002/edn3.141

Appeltans W, Ahyong ST, Anderson G, Angel M V., Artois T, Bailly N, Bamber R,
Barber A, Bartsch I, Berta A, Błazewicz-Paszkowycz M, Bock P, Boxshall G,
Boyko CB, Brandão SN, Bray RA, Bruce NL, Cairns SD, Chan TY, Cheng L,
Collins AG, Cribb T, Curini-Galletti M, Dahdouh-Guebas F, Davie PJF, Dawson
MN, De Clerck O, Decock W, De Grave S, De Voogd NJ, Domning DP, Emig CC,
Erséus C, Eschmeyer W, Fauchald K, Fautin DG, Feist SW, Fransen CHJM,
Furuya H, Garcia-Alvarez O, Gerken S, Gibson D, Gittenberger A, Gofas S,
Gómez-Daglio L, Gordon DP, Guiry MD, Hernandez F, Hoeksema BW, Hopcroft
RR, Jaume D, Kirk P, Koedam N, Koenemann S, Kolb JB, Kristensen RM, Kroh
A, Lambert G, Lazarus DB, Lemaitre R, Longshaw M, Lowry J, MacPherson E,
Madin LP, Mah C, Mapstone G, McLaughlin PA, Mees J, Meland K, Messing CG,
Mills CE, Molodtsova TN, Mooi R, Neuhaus B, Ng PKL, Nielsen C, Norenburg J,
Opresko DM, Osawa M, Paulay G, Perrin W, Pilger JF, Poore GCB, Pugh P, Read
GB, Reimer JD, Rius M, Rocha RM, Saiz-Salinas JI, Scarabino V, Schierwater B,
Schmidt-Rhaesa A, Schnabel KE, Schotte M, Schuchert P, Schwabe E, Segers
H, Self-Sullivan C, Shenkar N, Siegel V, Sterrer W, Stöhr S, Swalla B, Tasker
ML, Thuesen E V., Timm T, Todaro MA, Turon X, Tyler S, Uetz P, Van Der Land

J, Vanhoorne B, Van Ofwegen LP, Van Soest RWM, Vanaverbeke J, Walker-

Smith G, Walter TC, Warren A, Williams GC, Wilson SP, Costello MJ. 2012. The

magnitude of global marine species diversity. *Curr Biol* **22**:2189–2202.

doi:10.1016/j.cub.2012.09.036

Böttger-Schnack R, Schnack D, Hagen W. 2008. Microcopepod community structure

in the Gulf of Aqaba and northern Red Sea, with special reference to

Oncaeidae. *J Plankton Res* **30**:529–550. doi:10.1093/plankt/fbn018

Castellani C, Edwards M. 2017. Marine Plankton: A Practical Guide to Ecology,

Methodology, and Taxonomy. Oxford University Press. doi:10.1093

Chaudhary C, Saeedi H, Costello MJ. 2016. Bimodality of Latitudinal Gradients in

Marine Species Richness. *Trends Ecol Evol*. doi:10.1016/j.tree.2016.06.001

Childress JJ. 1995. Are there physiological and biochemical adaptations of

metabolism in deep-sea animals? *Trends Ecol Evol* **10**:30–36.

doi:10.1016/S0169-5347(00)88957-0

Clarke A, Gaston KJ. 2006. Climate, energy and diversity. *Proc R Soc B Biol Sci*

273:2257–2266. doi:10.1098/rspb.2006.3545

Cordier T, Angeles IB, Henry N, Lejzerowicz F. 2022. Patterns of eukaryotic

diversity from the surface to the deep-ocean sediment. *Sci Adv* **9309**:26–28.

Costello MJ, Chaudhary C. 2017. Marine Biodiversity, Biogeography, Deep-Sea

Gradients, and Conservation. *Curr Biol* **27**:R511–R527.

doi:10.1016/j.cub.2017.04.060

Cristescu ME, Hebert PDN. 2018. Uses and misuses of environmental DNA in

biodiversity science and conservation. *Annu Rev Ecol Evol Syst* **49**:209–230.

doi:10.1146/annurev-ecolsys-110617-062306

Daneri G, Dellarossa V, Quiñones R, Jacob B, Montero P, Ulloa O. 2000. Primary

production and community respiration in the Humboldt Current System off

Chile and associated oceanic areas. *Mar Ecol Prog Ser* **197**:41–49.

doi:10.3354/meps197041

Danovaro R, Gambi C, Dell'Anno A, Corinaldesi C, Fraschetti S, Vanreusel A, Vincx

M, Gooday AJ. 2008. Exponential Decline of Deep-Sea Ecosystem Functioning

Linked   to   Benthic   Biodiversity   Loss.   *Curr   Biol*   **18**:1–8.

doi:10.1016/j.cub.2007.11.056

Danovaro R, Gambi C, Della Croce N. 2002. Meiofauna hotspot in the Atacama

Trench, eastern South Pacific Ocean. *Deep Res Part I Oceanogr Res Pap*

49:843–857. doi:10.1016/S0967-0637(01)00084-X

de Puelles MLF, Gazá M, Cabanellas-Reboredo M, Santandreu M del M, Irigoien X,

González-Gordillo JI, Duarte CM, Hernández-León S. 2019. Zooplankton

abundance and diversity in the tropical and subtropical ocean. *Diversity* **11**:1–

22. doi:10.3390/d11110203

Deiner K, Bik HM, Mächler E, Seymour M, Lacoursière-Roussel A, Altermatt F, Creer

S, Bista I, Lodge DM, de Vere N, Pfrender ME, Bernatchez L. 2017.

Environmental DNA metabarcoding: Transforming how we survey animal and

plant communities. *Mol Ecol* **26**:5872–5895. doi:10.1111/mec.14350

Gaston KJ. 2000. Global patterns in biodiversity. *Nature* **405**:220–227.
doi:10.1038/35012228

Glud RN, Wenzhöfer F, Middelboe M, Oguri K, Turnewitsch R, Canfield DE, Kitazato
H. 2013. High rates of microbial carbon turnover in sediments in the deepest
oceanic trench on Earth. *Nat Geosci* **6**:284–288. doi:10.1038/ngeo1773

Grosberg RK, Vermeij GJ, Wainwright PC. 2012. Biodiversity in water and on land.
Curr Biol **22**:R900–R903. doi:10.1016/j.cub.2012.09.050

Harrison JB, Sunday JM, Rogers SM. 2019. Predicting the fate of eDNA in the
environment and implications for studying biodiversity. *Proc R Soc B Biol Sci*
286. doi:10.1098/rspb.2019.1409

Higgs ND, Attrill MJ. 2015. Biases in biodiversity: Wide-ranging species are
discovered first in the deep sea. *Front Mar Sci* **2**:1–8.
doi:10.3389/fmars.2015.00061

Ives AR, Carpenter SR. 2007. Stability and diversity of ecosystems. *Science*
317:58–62. doi:10.1126/science.1133258

Jamieson A. 2015. The Hadal Zone: Life in the deepest oceans, The Hadal Zone:
Life in the Deepest Oceans. doi:10.1017/CBO9781139061384

Jamieson AJ, Fujii T, Mayor DJ, Solan M, Priede IG. 2010. Hadal trenches: the
ecology of the deepest places on Earth. *Trends Ecol Evol* **25**:190–197.
doi:10.1016/j.tree.2009.09.009

Kim JG, Yu OH, Lee J. 2019. Description of two deep-water copepods of the genus

Leptotachidia Becker from the northwestern Pacific (Harpacticoida,

Pseudotachidiidae). *Zookeys* **873**:133–153. doi:10.3897/zookeys.873.34630

Kosobokova K, Hirche HJ. 2000. Zooplankton distribution across the Lomonosov

Ridge, Arctic Ocean: Species inventory, biomass and vertical structure, Deep-

Sea Research Part I: Oceanographic Research Papers. doi:10.1016/S0967-

0637(00)00015-7

Lebrato M, Pitt KA, Sweetman AK, Jones DOB, Cartes JE, Oschlies A, Condon RH,

Molinero JC, Adler L, Gaillard C, Lloris D, Billett DSM. 2012. Jelly-falls historic

and recent observations: A review to drive future research directions.

Hydrobiologia **690**:227–245. doi:10.1007/s10750-012-1046-8

Lundgreen RBC, Jaspers C, Traving SJ, Ayala DJ, Lombard F, Grossart HP, Nielsen

TG, Munk P, Riemann L. 2019. Eukaryotic and cyanobacterial communities

associated with marine snow particles in the oligotrophic Sargasso Sea. *Sci Rep*

9:1–12. doi:10.1038/s41598-019-45146-7

Luo JY, Condon RH, Stock CA, Duarte CM, Lucas CH, Pitt KA, Cowen RK. 2020.

Gelatinous Zooplankton-Mediated Carbon Flows in the Global Oceans: A Data-

Driven Modeling Study. *Global Biogeochem Cycles* **34**:0–2.

doi:10.1029/2020GB006704

Priede IG, Froese R. 2013. Colonization of the deep sea by fishes. *J Fish Biol*

83:1528–1550. doi:10.1111/jfb.12265

Ramirez-Llodra E, Brandt A, Danovaro R, De Mol B, Escobar E, German CR, Levin

LA, Martinez Arbizu P, Menot L, Buhl-Mortensen P, Narayanaswamy BE, Smith

CR, Tittensor DP, Tyler PA, Vanreusel A, Vecchione M. 2010. Deep, diverse and

definitely different: Unique attributes of the world's largest ecosystem.

Biogeosciences **7**:2851–2899. doi:10.5194/bg-7-2851-2010

Robison BH. 2009. Conservation of deep pelagic biodiversity. *Conserv Biol* **23**:847–

858. doi:10.1111/j.1523-1739.2009.01219.x

Sinniger F, Pawlowski J, Harii S, Gooday AJ, Yamamoto H, Chevaldonné P,

Cedhagen T, Carvalho G, Creer S. 2016. Worldwide analysis of sedimentary

DNA reveals major gaps in taxonomic knowledge of deep-sea benthos. *Front*

Mar Sci **3**:1–14. doi:10.3389/fmars.2016.00092

Smith KL, Ruhl HA, Bett BJ, Billett DSM, Lampitt RS, Kaufmann RS. 2009. Climate,

carbon cycling, and deep-ocean ecosystems. *Proc Natl Acad Sci U S A*.

doi:10.1073/pnas.0908322106

Steinberg DK, Landry MR. 2017. Zooplankton and the Ocean Carbon Cycle. *Ann Rev*

Mar Sci **9**:413–444. doi:10.1146/annurev-marine-010814-015924

Taberlet P, Coissac E, Hajibabaei M, Rieseberg LH. 2012. Environmental DNA. *Mol*

Ecol **21**:1789–1793. doi:10.1111/j.1365-294X.2012.05542.x

Teuber L, Schukat A, Hagen W, Auel H. 2013. Distribution and ecophysiology of

calanoid copepods in relation to the oxygen minimum zone in the eastern

tropical atlantic. *PLoS One* **8**:1–13. doi:10.1371/journal.pone.0077590

Thomsen PF, Kielgast J, Iversen LL, Møller PR, Rasmussen M, Willerslev E. 2012.

Detection of a Diverse Marine Fish Fauna Using Environmental DNA from

Seawater Samples. *PLoS One* **7**:1–9. doi:10.1371/journal.pone.0041732

Ulloa O, Canfield DE, DeLong EF, Letelier RM, Stewart FJ. 2012. Microbial

oceanography of anoxic oxygen minimum zones. *Proc Natl Acad Sci U S A*

109:15996–6003. doi:10.1073/pnas.1205009109

Valentine JW, Jablonski D. 2015. A twofold role for global energy gradients in

marine biodiversity trends. *J Biogeogr* **42**:997–1005. doi:10.1111/jbi.12515

Vereshchaka A, Abyzova G, Lunina A, Musaeva E. 2017. The deep-sea zooplankton

of the North, Central, and South Atlantic: Biomass, abundance, diversity. *Deep*

Res Part II Top Stud Oceanogr **137**:89–101. doi:10.1016/j.dsr2.2016.06.017