High total water loss driven by low-fat diet in desert-adapted mice

Danielle M. Blumstein[1], Jocelyn P. Colella[2], Ernst Linder[3], Matthew D. MacManes[1]

1. University of New Hampshire, Molecular, Cellular, and Biomedical Sciences

Department, Durham, NH 03824 (DMB Dani.Blumstein@unh.edu, Orcid ID: 0000-0003-

2129-5541; MDM Matthew.MacManes@unh.edu, Orcid ID: 0000-0002-2368-6960)

2. University of Kansas, Biodiversity Institute, Lawrence, KS 66045 (JPC

colella@ku.edu, Orcid ID: 0000-0003-2463-1029)

3. University of New Hampshire, Department of Mathematics and Statistics, Durham,

NH 03824 (EL ernst.linder@unh.edu)

Abstract

Availability of essential food resources is one of the most important drivers of survival.

Animals must either relocate or adapt to persist in environments where critical resource

abundance is changing. Investigating a species' ability to respond to alternative

conditions can reveal a myriad of responses. An optimal diet balances energy gain,

water regulation, and nutrition. We used flow-through respirometry to characterize

metabolic phenotypes of the desert-adapted cactus mouse (*Peromyscus eremicus*)

under diurnally variable environmental conditions that mimic the environment of the

Sonoran Desert. We treated mice with two different energy-controlled diets, a standard

diet and a low-fat diet, and measured energy expenditure, rate of water loss, respiratory

quotient, weight, and electrolyte levels. Mice fed the low-fat diet lost significantly more

water than those on the standard diet. Our results suggest that cactus mice may have

limited capacity to tolerate water deprivation if optimal foods become less abundant.

Given that climate change is predicted to modify the distribution of food items,

understanding these links may have important implications for long-term population

viability for desert and non-desert adapted animals alike.

Keywords: dietary fat, energy expenditure, *Peromyscus*, physiology, respirometry, total

water loss

Introduction

As the planet undergoes unprecedented climate change, increased desertification is

predicted (Wu et al. 2014; IPCC 2019; Mirzabaev et al. 2019). Desertification leads to

increased aridity, a strong selective force that may result in changes to species

distributions and community composition. Understanding these responses is critically

important for predicting the impacts of climate change on biodiversity. Organismal

responses to climate change vary, but can be classified into three general categories:

geographic relocation to maintain physiological balance (Moritz et al. 2008; Walsh et al.

2016; Riddell et al. 2021), adaptation in place (particularly for species with fragmented

habitats or limited dispersal ability, Reed et al. 2011), or perish, leading to population

extirpation or extinction. Distributional changes in response to changing climate are well

documented (Parmesan and Yohe 2003; Parmesan 2006; Moritz et al. 2008; Van der

Putten et al. 2010; Pecl et al. 2017; Morrison et al. 2018). Adaptive responses can take

two forms: evolutionary change over generational timeframes or phenotypic plasticity

within an animal's lifetime; however, the latter is less well documented. Adaptation *in*

situ may involve morphological, behavioral, or physiological changes. For example,

there are strong directional selections for seasonal adaptations (e.g., seasonal color

camouflage, Jones et al. 2018), but in the absence of standing genetic variation, there is

evidence of increased extinction risk (Spielman et al. 2004).

The Sonoran Desert is hot, dry, and environmentally dynamic. Daily ambient

temperatures in the Summer can reach as high as 50°C during the day and as low as

15°C at night. In the winter, the daytime temperature is typically between 20 – 30°C and

can fall to near freezing at night. Average relative humidity (RH) in the Sonoran Desert

changes by 20% between seasons (Reid 1987; Unland et al. 1996; Sheppard et al.

2002, Boyd Deep Canyon Desert Research Center). High ambient temperatures and

lack of extrinsic water challenge mammalian survival, as mammals rely on water for

fundamental metabolic processes, behavior, and/or evaporative cooling (Reece et al.

2015). Despite these challenges, many mammals thrive in deserts. This has led to

decades of work aimed at one central question: How do desert animals survive

(Bradford 1974; MacMillen and Hinds 1983; Nagy 1987; Cortés et al. 2000; Kordonowy

and MacManes 2017; MacManes 2017; Tigano et al. 2020; Bittner et al. 2021; Colella et

al. 2021b)?

One of the most critical challenges faced by desert mammals includes the

regulation of water balance. Water balance is maintained through a cascade of

hormones that elicit a response designed to increase water retention and decreases

water used in the excretion of nitrogenous waste. In brief, dehydration results in a

decrease in blood volume and an increase in osmolality, the latter, an effect driven

largely by the rise in serum Sodium (Thornton 2010; Leib et al. 2016). The decrease in

blood volume, detected by baroreceptors, signals the heart rate to increase in an

attempt to maintain cardiac output and therefore, blood flow to critical organs

(Kaufmann et al. 2020). In response to the increase in serum osmolality, osmoreceptors

in the hypothalamus are stimulated and antidiuretic hormone (ADH) is released from the

posterior pituitary gland (Thornton 2010; Leib et al. 2016). Then, several things happen;

the kidneys are signaled to recover water (Fuller et al. 2020) and epithelial cells lining

renal collecting tubules are signaled to move aquaporins from the interior of the cell to

the surface. This increases the water permeability of cells and leads to more water

reabsorbed from the tubule system back into the blood stream (Kortenoeven and

Fenton 2014). The renin-angiotensin-aldosterone hormonal system in the kidneys is

also signaled and spits into the active hormone angiotensin II, stimulating the release of

aldosterone from the adrenal glands and ADH from the pituitary gland, resulting in the

retention of sodium in the distal tubules of the nephrons (Goodfriend 2006). While

hormonal regulation of water balance is paramount to survival, behavioral,

physiological, and genomic responses are also critical.

Genomic and transcriptomic studies of desert mammals have found evidence of

positive selection acting on genes involved in energy production, food storage (Jirimutu

et al. 2012; Wu et al. 2014), water reabsorption (Jirimutu et al. 2012; Marra et al. 2012,

2014; Yang et al. 2016; Giorello et al. 2018), and osmoregulation (MacManes and Eisen

2014; Kordonowy and MacManes 2016; Colella et al. 2021b). A strong signature of

positive selection in desert mice on Growth Differentiation Factor 10 (*Gdf10*) may

prevent weight gain when consuming high-fat diets (Hino et al. 2017; Tigano et al.

2020). While this prior research has enhanced our understanding of the molecular

processes underlying adaptation in desert environments, we still do not fully understand

the suite strategies employed by desert animals to manage water efficiently.

Furthermore, animals that have evolved in desert environments are behaviorally

and physiologically adapted to extreme aridity and lack of extrinsic water. Behaviors

such as foraging strategies (Manlick et al. 2021), dietary preference (Frank 1988),

burrowing (Schmidt-Nielsen and Adolph 1964; Louw and Seely 1984; Riddell et al.

2021), group nesting (Schmidt-Nielsen and Adolph 1964), microhabitat use during the

heat of the day (Riddell et al. 2021), and nocturnality (but see Walsberg 2000; Tracy

and Walsberg 2002) may optimize fitness in hot, dry environments by allowing animals

to regulate temperature and water balance in real-time (Frank 1988). Behavioral

changes may be the first line of defense against changing environments, as they are

often dynamic, reversible, and happen in real time. In contrast, physiological

adaptations, such as efficient kidney function or low fecal water content, which are

observed in desert mammals to maintain water balance, can happen in real-time but

typically have evolved over deeper evolutionary timescales (Christian and Matson 1978;

Kordonowy and MacManes 2016).

Dietary intake of food is one important way in which animals may manage water.

There are three main macromolecules that animals ingest as food items; carbohydrates,

fats, and proteins, all of which vary in the amount of available energy and water, as well

as in their cost to metabolize. Energy potential varies greatly, with carbohydrates and

proteins yielding 16.736 J/g while fats yield more than double that (37.656 J/g,

Sánchez-Peña et al. 2017). Metabolic water, the water produced through catabolism of

carbohydrates, fats, or proteins (Schmidt-Nielsen and Adolph 1964; Frank 1988; Orr et

al. 2015) also varies dramatically. The oxidation of carbohydrates yields 0.60 g of

metabolic water per gram, and fats yield 1.07 g per gram due to greater oxygen

requirements during lipid metabolism, while proteins yield the least metabolic water

(0.41 g) and even require water loss for excretion of nitrogenous waste (Mellanby 1942;

Davidson and Passmore 1963; Schmidt-Nielsen and Adolph 1964).

In environments like deserts, where extrinsic water is limited, preformed (dietary)

water and endogenous water production are key to survival. Dietary intake represents

the raw material to produce metabolic water. Food catabolism results in water loss

through respiration, urination, and fecal production (Schmidt-Nielsen 1975). Although

specific dietary composition is often unknown and highly variable, particularly for wild

animals, energy potential (Pyke et al. 1977; Pyke 1984), nutrition, and capacity for water

production (Schmidt-Nielsen and Adolph 1964) all depend on diet composition. In

deserts, vegetation, seeds, and insects comprise the bulk of rodent diets (Reichman

1975) with each varying in fat, protein, and carbohydrate composition and may exhibit

seasonal availability (Wolf and del Rio 2003; Orr et al. 2015). In response to natural

variation in resource availability, animals have the opportunity to make dietary decisions

that may have a substantial impact on their internal water economy.

Mice of the genus *Peromyscus* have the widest distribution of any North

American mammal and are considered model organisms in evolutionary biology due to

their unparalleled habitat diversity (Bedford and Hoekstra 2015), extensive genomic

resources (*e.g.,* Tigano et al. 2020; Colella et al. 2021), abundance of historically and

contemporaneously collected samples in natural history museums (Pergams and Lawler

2009; Pardi et al. 2020), and ability to live and breed under laboratory conditions

(Crossland et al. 2014). The desert specialist cactus mouse, *Peromyscus eremicus*, is

endemic to the southwestern United States and exhibits behavioral (Murie 1961; Veal

and Caire 1979), physiological (Macmillen 1965; Kordonowy et al. 2017; Colella et al.

2021a), and molecular (MacManes 2017; Tigano et al. 2020) adaptations to desert

environments, making it an interesting natural experimental model to examine the

potential for adaptation to warmer, drier environments. *Peromyscus eremicus* are

omnivorous and opportunistic in their diet, utilizing seeds, arthropods, and green

vegetation seasonally (Bradley and Mauer 1973; Meserve 1976) and they shift to

consuming cactus seeds and/or fruit pulp during the summer months (Orr et al. 2015).

We examine the effect of dietary fat content on the metabolic physiology of the

desert-adapted cactus mouse. To accomplish this, we extend previously characterized

circadian metabolic patterns for males and females (Colella et al. 2021a) to rates of

metabolism and water loss in animals fed an experimental diet low in fat, but unchanged

in terms of energy composition and level of macronutrients. This allows us to test the

relationships between dietary fats and energy expenditure and water balance, both of

which have direct connections to fitness. Given that fats yield more water and energy

compared to carbohydrates and proteins, we hypothesized that animals fed an LFD

would perform more poorly than those on the SD.

Methods

Animal Care and Experimental Model

We worked with a captive colony of cactus mice, held at the University of New

Hampshire. Captive animal care procedures followed guidelines established by the

American Society of Mammologists (Sikes et al., 2016) and were approved by the

University of New Hampshire Institutional Animal Care and Use Committee under

protocol number 103092.

Sexually mature, non-reproductive healthy adults (n = 28 males, n = 28 females)

between three and nine months of age, bred from wild-derived lines maintained at the

Peromyscus Genetic Stock Center of the University of South Carolina (Columbia, South

Carolina, USA) were used in this study. Mice were randomly assigned to one of two

feeding groups: a standard diet group (SD - LabDiet® 5015*, 26.101% fat, 19.752%

protein, 54.148% carbohydrates, energy 15.02 kJ/g, food quotient [FQ, the theoretical

RQ produced by the diet based on macronutrient composition, Westerterp, 1993] 0.89

or a low-fat diet group (LFD - Modified LabDiet® 5015 [5G0Z], 6.6% fat, 22.8% protein,

70.6% carbohydrates, energy 14.31 kJ/g, FQ = 0.92). All food was stored in the desert

chamber to match the environmental conditions described below. Animals were housed

individually in 9.5L animal chambers inside a larger, environmentally controlled room

built to simulate the temperature, humidity, and photoperiodic conditions of the Sonoran

Desert (Kordonowy et al. 2017; Colella et al. 2021a). Each animal chamber contained

dried cellulose-based bedding, and animals were provided food and water *ad libitum*.

Mice were acclimated to the LFD for one month and to experimental cages for 24 hours

prior to the beginning of metabolic measurements. Animals were weighed to the nearest

tenth of a gram on a digital scale before being housed in an experimental chamber for

four days during metabolic data collection. Animals were then weighed at the end of the

experiment.

Environmental temperatures and relative humidity (RH) follow a natural, diurnal

pattern. Temperature and RH were fixed at 32°C and 10% RH during the light phase for

11 hours (06:00 to 20:00) and dropped to 24°C and 25% RH during a one-hour

transition starting at 20:00, then remained constant during the dark phase for seven

hours, before the temperature rises again and the RH falls to light phase temperature

during a three-hour transition that starts at 06:00. Photoperiod was regular with 16

hours of light and 8 hours of dark. See Figure 1A for a visual representation of

environmental conditions.

Metabolic phenotyping

Metabolic phenotyping data were collected starting at 10:00 using a previously

described (Colella et al. 2021a) commercially available respirometry system purchased

from Sable Systems International (SSI). Physiological data were collected following the

methods outlined in Whitfield et al. (2015) and Colella et al. (2021a). Oxygen

consumption (VO_2 ml min^{-1}), carbon dioxide production (VCO_2 ml min^{-1}), and water

vapor (WVP, kPa) lost via urine, feces, cutaneous, and respiration were continually

measured during 72-hour trials. This was repeated twice for each sex under each diet

(SD and LFD), resulting in three-day experiments performed sequentially in eight

batches of seven mice (and one empty baseline chamber per three-day experiment) at

a time, with batches alternating between females and males for a total of 28 adult males

and 28 adult females.

Airstreams were pulled from flow-through chambers at a constant rate of 1600 ml

min^{-1} (96 L h^{-1}) by an SSI SS-4 sub-sampler pump (one for each chamber), multiplexed

through the SSI MUXSCAN, sub-sampled at 250 ml min^{-1} into a Field Metabolic System

(FMS) where water vapor, CO_2, and O_2 were analyzed with no scrubbing. The FMS was

zeroed and spanned the sensors between each 72-hour experiment following the SSI

Instrument Settings and Calibration manual using dry gas with known concentrations of

CO_2 and O_2.

To generate accurate individual measurements, chambers were measured in a

pseudorandom order for 120s each, approximately three times every hour. Using

methods described in Lighton (2018), we used the most stable 50% of each 120s period

to produce a single averaged value for the rate of O_2 consumption (ml min^{-1}), rate of

CO_2 production (ml min^{-1}), and rate of water loss (RWL) (mg h^{-1}) per measurement

window. Lag time between the analyzers was corrected so that measurements

represent the same time periods (Lighton 2018). This procedure results in a highly

repeatable set of measurements. VO_2 and VCO_2 were calculated using equations 10.5

and 10.6, respectively, from Lighton (2018). Respiratory quotient (RQ) was calculated

as the ratio of VCO_2 to VO_2 (Lighton 2018). Energy expenditure (EE), kJ hr^{-1}, was

calculated as in Lighton (2018, eq. 9.15). Response variable measurements that were

more than three standard deviations away from the mean, which often represented

times when animal care staff entered the room, were considered outliers, and therefore

were removed from the downstream analysis.

Electrolytes

At 12:00 Eastern Standard Time (EST), immediately following the conclusion of the

experiment, animals were weighed and euthanized with an overdose of inhaled

isoflurane and decapitation. 120 μL of trunk blood was collected within one minute of

death for serum electrolyte measurement using an Abaxis i-STAT® Alinity machine and

CHEM8+ cartridges (Abbott Park, IL, USA, Abbott Point of Care Inc.), and calibrated

automatically with each cartridge run. CHEM8+ cartridges measure the concentration of

Sodium (Na), Potassium (K), Creatinine (Cr), blood urea nitrogen (BUN), Hematocrit

(HCT), Glucose (Glu), and ionized Calcium (iCa), which are expected to vary in

response to hydration status and renal function. Finally, using Sodium, Glucose, and

BUN, we calculated osmolality using the formula in Rasouli (2016). After assessing for

normality, we used a student's two-tailed t-test (stats::t.test) to test for significant ($p <$

0.05) differences in weight and electrolytes within and between the treatments and

sexes for each experimental group.

Statistical Analysis

All statistical analyses were conducted in R v 4.0.3 (R Core Team 2020) unless

otherwise specified. To perform analyses on the shape of the temporally variable

curves, we modeled the effect of time in days, diurnal time, and EE, RQ and/or RWL on

EE, RQ, and RWL using Generalized Additive Mixed Models (GAMMs) through the

function mgcv::gamm (Lin and Zhang 1999; Wood 2017). To avoid violating the

assumption of independence, we used the experimental batches and the mice nested

within the experimental batch as random effects. The R function "gamm" then

simultaneously estimates the variances of these random effects as well as the treatment

effects, thus implicitly adjusting treatment and covariate effects for the effect of

correlated nested groups of measurements. As a result, treatments do not explain

differences between individual mice, but rather the average difference between groups

of mice, because we were interested in the diet treatment and the effect of sex, as well

as in the response over the full 72-hour period and the diurnal cycle. Our models

included two fixed effects, diet and sex, and four nonlinear smoothing regression terms:

time in days as a nonlinear trend, diurnal cycle as a circular spline function, and two of

the three respirometry response variables (EE, RQ, and/or RWL). Using the effective

degrees of freedom (edf), we were able to determine the degree of non-linearity of each

curve (Wood 2017). To test for physiological differences between mice fed each diet

during the light and dark phases, we subsetted data collected during the light phase and

dark phase for each response variable. Using the subsetted data, we used GAMMs as

described above.

To test for differences in the total volume of water lost for each animal over the

course of the experiment, rather than the shape of the curve, we estimated the amount

of water lost for each individual for every hour of the experiment and summed those

values. These values were then used as the response variable in an ANCOVA

(car::anova, Fox and Weisberg 2019). The model consisted of two independent

variables, sex and diet, and the interaction between them. Weight at the start of the

experiment was used as the covariate. Pairwise comparisons were made using the

Tukey separation of adjusted means test (emmeans:: emmeans and

emmeans::contrast, Lenth 2021).

Results

Data from 56 sexually mature (but non-reproductive) male and female mice were
collected as described above. All mice were subjected to standard animal care
procedures before the experiment which included a health assessment conducted by
licensed veterinary staff. No health issues were detected by veterinary staff, nor were
there significant differences in weight between males and females (mean of 22.2 g
[range: 15.40 g - 29.10 g] versus 21.2 g [range: 16.91 g - 29.34 g], respectively; Table
1), or between diet groups.

Figure 1

72 hours of room (A) and respirometry (B-D) data collection spit by sex for 28 adult males and 28 adult females plotted in a 24-hour window to display circadian patterns for each group: Organismal responses to standard diet are in gray and low-fat diet in blue. Vertical blocks shaded in gray indicate the dark phase when animals are active, and unshaded blocks indicate light phase when animals are inactive. **A)** Room temperature ($°C$) is indicated by a solid red line. **B)** 24-hour measurements of energy expenditure (EE kJ hr^{-1}), **C)** rate of water loss (RWL, H_2O g hr^{-1}), and **D)** respiratory quotient (RQ), for females (F, left) and males (M, right). Statistical significance between the fixed effects, diet and sex, for the 24hr GAMMs are denoted with one or more · under the y-axis label, 'ns' indicates non-significance. Statistical significance between the fixed effects, diet and sex, from GAMMs on the subsetted day/night data are denoted with one or more * on the top axis of the plots. The two symbols (*/·) signify the same significance values (* $p <= 0.05$, **: $p <= 0.01$, ***: $p <= 0.001$, ****: $p <= 0.0001$).

Electrolytes

Several serum electrolyte values were significantly different between diets for males and females (Sodium $p = 0.22$ and $p = 0.01$ respectively; Potassium $p = 0.02$ and $p = 0.01$; Blood Urea Nitrogen $p = 0.03$ and 0.71; Hematocrit $p = 0.04$ and 0.02; Ionized Calcium $p = 0.001$ and 0.001; osmolality $p = 0.01$ and 0.05) (Figure 2, Table1). When comparing males and females on the same diet, Sodium ($p = 0.02$) and Osmolality ($p =$

305   0.03) differed significantly for mice on the SD and no electrolyte measurements were

306   differed between males and females on the LFD.

		Weight (g)	Na (mmol/L)	K (mmol/L)	Cr (µmol/L)	BUN (mmol/L)	Hct (% PCV)	iCa (mmol/L)	Osmolality (mmol/L)
		mean (+/- SD)	mean (+/- SD)	mean (+/- SD)	mean (+/- SD)	mean (+/- SD)	mean (+/- SD)	mean (+/- SD)	mean (+/- SD)
Standard diet	Female	22.4+/-2.73	137.5+/-3.58	8.02+/-1.35	0.22+/-0.04	36.25+/-5.33	26.08+/-9.58	1.1+/-0.1	291.15+/-7.34
	Male	22.24+/-2.14	140.71+/-2.89	7.84+/-0.96	0.2+/-0.03	33.21+/-4.08	32.5+/-2.79	1.16+/-0.07	297.54+/-5.82
	All	21.13+/-2.63	139.23+/-3.56	7.92+/-1.14	0.21+/-0.03	34.62+/-4.85	29.54+/-7.42	1.13+/-0.09	295.04+/-7.1
Low fat diet	Female	20.02+/-2.68	143.6+/-4.7	5.98+/-1.49	0.2+/-0	38.3+/-5.5	34.4+/-3.27	1.29+/-0.07	307.83+/-10.3
	Male	22.29+/-4.05	142.5+/-3.63	6.37+/-1.59	0.2+/-0	37+/-3.53	36.2+/-5.57	1.27+/-0.09	304.18+/-9.22
	All	22.35+/-3.38	143.05+/-4.12	6.18+/-1.51	0.2+/-0	37.65+/-4.55	35.3+/-4.54	1.28+/-0.08	306.01+/-9.7

Table 1

Summary statistics for weight (g) and serum electrolyte measurements (Na = Sodium

(mmol/L), K = Potassium (mmol/L), Cr = Creatinine (µmol/L), BUN = Blood Urea

Nitrogen (mmol/L), Hct = Hematocrit (% PCV), iCa = Ionized Calcium (mmol/L), and

osmolality (mmol/L) for female and male *Peromyscus eremicus* for standard and low-fat

diet groups.

Figure 2

Violin plots showing the distribution of serum electrolyte measurements (Na = sodium

(mmol/L), K = potassium (mmol/L), BUN = blood urea nitrogen (mmol/L), Hct =

hematocrit (% PCV), iCa = ionized calcium (mmol/L), Glu = glucose (mmol/L), and

osmolality (mmol/L) for female (F) and male (M) *Peromyscus eremicus* fed standard

(grey) and low fat (blue) diets. Observations (n = 56, 28 for each sex) are represented

by black dots. P-values from pairwise t-tests are reported above the brackets.

Metabolic Phenotyping

In total, we recorded 8,125 120s intervals: 4,091 female, 4,034 male (Supplemental

Table 1). We measured metabolic variables for 4,105 120s intervals under the LFD:

2,053 female and 2,052 male measurements. For the SD, we recorded 4,020 total

observations (2,038 female, 1,982 male). After removing outliers, we retained 3,945

female measurements (1,977 SD and 1,968 LFD) and 3,963 male measurements

(1,945 SD, 2,018 LFD). *Peromyscus eremicus* show physiological patterns that are in

phase with photoperiod and room temperature for both diets (Figure 1), as described in

(Fox and Weisberg 2019; Colella et al., 2021b). All raw data and processed machine-

readable csv files are available on Zenodo:

https://zenodo.org/record/6422231#.YIXSD9PML0o.

Energy Expenditure

Consistent with previous work (Colella et al. 2021a), males and females for both diets

show diurnal patterning of EE, with the highest EE occurring during the dark phase

when environmental temperature is lowest, and animals are active. The lowest EE

occurred during the light phase, when environmental temperature was greatest and

animals were inactive (Figure 1B). Using a GAMM, we modeled EE with diet and sex as

fixed effects, as well as other predictors relevant factors such as day, time, RWL, and

RQ and found that neither sex nor diet influenced EE (diet $p = 0.56$, sex $p = 0.18$,

Supplemental Table 2). When analyzing day and night separately, there were no

differences in EE between males and females nor between the two diets during the light

or dark phase (light phase: diet $p = 0.550$, sex $p = 0.293$; dark phase: diet $p = 0.571$, sex

$p = 0.182$, Supplemental Table 3 and 4). There was a weak nonlinear relationship

between EE and day (Supplemental Figure 2A, Supplemental Table 2, edf = 1.98, $p <$

$2e^{-16}$). Additionally, the relationship between diurnal time and EE was statistically

significant but nonlinear (Supplemental Figure 2F, Supplemental Table 2, edf = 7.98, p

$< 2e^{-16}$).

Rate of Water Loss

Both treatment groups, SD and LFD, and both sexes exhibited diurnal patterning of rate

of water loss and had the highest rate of water loss during the light phase when

environmental temperature is highest, even though animals are inactive (Figure 1C,

Supplemental Figure 1F, Supplemental Table 2, edf = 7.95, $p < 2e^{-16}$). Peak loss

occurred between 09:00 and 11:00. By fitting a 24-hour time-continuous GAMM

(Supplemental Figure 1E-H), we showed that both diet and sex had a significant effect

on the RWL (diet $p = 2.13E^{-05}$, sex p = 0.002, Supplemental Table 2), with both males

and females fed an LFD losing more water than those fed the SD. When modeling

water loss during the light phase only, when animals are inactive, we showed that both

diet and sex had a significant effect on RWL (diet $p = 4.80e^{-15}$, sex $p = 1.69e^{-11}$,

Supplemental Table 3). When modeling water loss only during the dark phase, when

animals are active, RWL was significantly different between males and females (p =

0.015, supplemental table 4), but RWL was not different within sex between different

diets (p = 0.797, Supplemental Table 4). The relationship between RWL and

experimental day was weakly non-linear, but significant (Supplemental Figure 1E,

Supplemental Table 2, edf = 1.99, $p < 2e^{-16}$).

When analyzing the volume of water lost, both diet and sex significantly affect the

amount of water lost over the course of the experiment when controlling for individual

body weight ($p = 5.22e^{-2}$ and p = 0.017, respectively, Supplemental Table 5), but the

interaction term was not significant (p= 0.633, Supplemental Table 5). Further

investigation into the sex:diet interaction showed significant differences in the female

diet comparison (p=0.043, Supplemental Table 5), and also in the male diet comparison

(p=0.005, Supplemental Table 5).

Respiratory Quotient

RQ showed circadian rhythms for both diets and sexes with the highest RQ occurring

during the light phase and the lowest RQ occurring during the dark phase (similar to

Colella et al. 2021b; Figure 1D). By fitting a 24-hour GAMM we found no significant

difference in RQ between the sexes or the two diets (diet p = 0.52, sex p = 0.83,

supplemental 2) but we did find that day and time were highly significant (p = $1.98e^{-07}$, p

< $2.2e^{-16}$ respectively). Time of day had a complex relationship with RQ (Supplemental

Figure 1, edf = 7.69, Supplemental Table 2) and experimental day had a weak non-

linear relationship with RQ (Supplemental Figure 1D, edf = 1.59, Supplemental Table 2).

When modeling RQ during the light phase, RQ rose above the FQ, with females

fed the SD having the highest RQ at the start of the light phase and then stabilizing at

an RQ > 1 around 14:00. We showed that neither sex nor diet were significantly

different for RQ (sex p = 0.740, diet p = 0.543, Supplemental Table 3). When modeling

RQ during the dark phase, RQ values were comparable to the FQ of the respective diet

for both sexes (FQ LFD: 0.92, FQ SD: 0.89) and RQ was significantly different between

the diets (p = 0.017, Supplemental Table 4) but not between males and females (p =

0.739, Supplemental Table 4).

Discussion

In light of current rapid climate change (IPCC 2019), understanding how animals

respond to environmental change is of paramount importance. Species unable to adapt

in place could relocate, but given the magnitude and rate of change and extent of

habitat fragmentation, species may be unable to move fast enough or find suitable

habitat elsewhere (Loarie et al. 2009; Beever et al. 2010; Sgrò et al. 2011; Henry et al.

2012). The only option left is adaptive evolutionary change, local extirpation, or in more

extreme circumstances, extinction. Extreme environments (*e.g.,* high altitude, deep sea,

deserts) and the organisms living in them represent natural experimental models for

understanding dynamic organismal responses. This research capitalizes on the

experimental tractability of a desert specialist rodent to better understand how diet may

influence water economy.

We explored the impacts of differences in dietary fat intake and found that mice

fed a diet lower in fat lost more water than mice fed a diet higher in fat. In environments

where water scarcity is extreme, even small differences in water balance could have

substantial impacts on cognitive function and physical performance, and therefore

survival and ultimately reproductive success (Boogert et al. 2018). Our results may

clarify the physiological consequences of food availability and dietary choices.

Electrolytes

Electrolytes provide information on the overall metabolic state, renal function, and other

core physiological functions (Kutscher 1968), including hydration status (Cheuvront et

al. 2010) or disease state. Therefore, to further understand the impacts of diet on

physiological function of endotherms in desert environments, we collected serum

electrolyte data from male and female cactus mice fed diets with varying levels of fat.

We established diet-specific differences for serum Na, K, Cr, BUN, Hct, iCa, and

osmolality values for *P. eremicus*. Notably, the synthetic markers of pathological renal

impairment, BUN and Cr, were not significantly different between treatments. Previous

studies have shown that the electrolyte-water balance is frequently unaffected when

desert rodents are water-deprived for long periods of time (Heimeier et al. 2004;

Heimeier and Donald 2006; Boumansour et al. 2021), however, other desert rodent

studies have demonstrated evidence of dehydration in the absence of exogenous water

without renal impairment (Kordonowy et al. 2017; Boumansour et al. 2021).

Interestingly, key electrolytes that are sensitive to hydration status were

significantly different between the dietary treatment groups (Figure 2). Serum Na, serum

osmolality, and Hct are all expected to be elevated in clinical dehydration and were

found to be elevated in animals fed the LFD. Though some of the contrasts were not

statistically significant, measurements were consistently higher in both males and

females, and this is a strong sign of differences in hydration. These differences are

particularly interesting because while animals fed an LFD had higher rates of water loss

than those fed the SD, all animals had access to water *ad libitum*. Elevated serum Na,

osmolality, and Hct in the context of normal renal function are evidence of dehydration,

but dehydration in neuro-intact animals with free access to water is unusual. While

dietary fats may be responsible for changes in solute balance (Friedman et al. 2012),

typically the relationship is in the opposite direction to what we found (Abdel-Rahman

2010). One possible hypothesis is related to an impairment of the physiological

mechanisms related to thirst. Thirst is a response to changes in the blood chemistry

(*i.e.,* increase in osmolality Gilman 1937; Wolf 1950; Heimeier et al. 2004) or volume

(Fitzsimons 1961; Stricker 1966) both of which are monitored by the lamina terminalis

circumventricular organs (Bourque 2008; Zimmerman et al. 2017). Activation of these

brain regions motivates an organism to find water. Previous studies have found that the

expression of two genes, *Rxfp1* and *Pdyn* in relevant cell types, correlate with neural

activation under osmotic and hypovolemic thirst in lab mice (Pool et al. 2020). Further

studies should examine these genes and others in desert organisms to understand the

basis of thirst pathways. Regardless of the process, elevated serum Sodium may have

important and complex physiological consequences, which may include elevated blood

pressure and increased fluid retention, both of which have obvious implications for

survival in deserts and deserve future study.

Energy Expenditure

Not all diets provide animals with the same amount of energy per unit mass (Withers

1982). Different macronutrients have different energy potentials, with carbohydrates and

proteins yielding 16.7 kJ/g while fats yield 37.7 kJ/g (Sánchez-Peña et al. 2017). Diet

composition influences EE (Acheson 1990; Westerterp 2004; Secor 2009; Beale et al.

2018), as do other intrinsic (*i.e.,* body mass, both temperature, evolutionary history,

reproductive stage, activity, or energy intake) and extrinsic (*i.e.,* habitat [including

environmental temperature], climate, or social environment; Speakman 1997) variables.

In wild populations, dietary switches tend to coincide with seasonal changes that affect

resource quantity and/or quality (Noble et al. 2019). Previous studies have recorded

cactus mice shifting diet seasonally, consuming arthropods during the winter (Hope and

Parmenter 2007), and transitioning to the consumption of cactus seeds and/or fruits

during the summer (Hope and Parmenter 2007; Orr et al. 2015). Many plants native to

the Sonoran Desert have a fat content comparable to that of the LFD used in this study

(5.49 - 9.88% for leaf tissues and branches, McArthur 1994; 0.90- 3.60% for roots and

seeds, Castle et al. 2020).

In the current study, both the low-fat and standard diets were designed to have

equivalent amounts of energy per gram of weight. We designed the experiment to

control for several intrinsic and extrinsic factors that represent sources of variation in

EE, but not individual activity, to show that experimental manipulation of diet does not

result in differences in EE when food is provided *ad libitum*. However, EE and activity

have been shown to be tightly coupled (Garland et al. 2011; Kaiyala et al. 2012),

therefore we believe there to be, at most, limited differences in activity given limited

differences in EE. For both diets, animals are likely to eat enough to satisfy energy

needs, a hypothesis further supported by the fact that there was no net change in

weight over the course of the experiment. Consistent with other studies, we recorded

higher EE during the dark phase of the 24-hour cycle when animals were active (Colella

et al. 2021a), further supporting that metabolic rate is complex and may be modulated

by the environment, activity, or other extrinsic/intrinsic factors.

Rate of Water Loss

In environments like deserts, where water limits survival, even small differences in water

loss, like those observed here between diets, may have important implications for

survival. Sources of water loss include evaporation, urine, and fecal, the latter two used

for processes related to the elimination of waste products resulting from metabolic

processes.

Thermoregulation and water loss: Evaporative water loss is one way for organisms to

cool themselves as evaporation is accompanied by heat dissipation (*e.g.,* via latent heat

of vaporization) and results in a decrease in body temperature and loss of body mass

(Porter and Gates 1969). While body temperature was not measured as a part of this

study, numerous studies have shown that desert animals reduce respiratory water loss

as a major adaptive physiological mechanism for maintaining water balance in deserts

(Schmidt-Nielsen and Schmidt-Nielsen 1951; Hart 1971; MacMillen 1972). Indeed, a

fine balance must exist between using water for thermoregulation with the use of water

for other critical metabolic processes, and with dehydration. Exaggerated use of water

for thermoregulation may be harmful when water is limited. Instead, other less water-

intensive mechanisms for heat loss such as conduction, and behavioral changes such

as estivation and nocturnality to maintain body-temperature homeostasis may be used

but are not quantified in this study.

Despite the potential consequences related to limited water supply, desert

Peromyscus have shown significant capacity for evaporative cooling for the purposes of

thermoregulation (Ramirez et al. 2022). In our study, we found that animals fed the LFD

had increased water loss compared to animals fed the SD, which has a higher fat

concentration (Figure 1C). While this loss may be related to differences in

thermoregulatory performance, the increased loss of water may further compromise

water balance. Are these patterns suggestive of differences in physiological

performance? Is the consumption of a diet lower in fat related to a reduction in

physiological performance and, consequently, fitness? While the current experiment

cannot definitively answer these questions, there is some evidence (*e.g.,* electrolyte

data) to suggest that physiological performance is impaired in animals consuming the

LFD and that this performance, particularly when water is scarce, may have fitness

consequences.

Potential mechanisms for increased water loss on LFD: Across all comparisons,

there is a strong and statistically significant relationship between the rate of water loss

and the fat content of the ingested diet, with individuals fed the LFD losing more water.

While there are some potential mechanisms underlying these findings, all of which will

require future experiments, several can be ruled out. First, as described above, the

maintenance of water balance is a process regulated by a cascade of hormones

including arginine vasopressin and the renin–angiotensin–aldosterone system pathway.

While some hormones are lipid-based (*e.g.,* aldosterone) and their production and

secretion are linked to dietary composition (*i.e.*, sex hormones and severe low body fat),

the degree to which dietary fats are reduced for animals fed the LFD are not to the level

below which hormonal imbalance should be seen. Second, while there is a known

relationship between ketogenic diets and dehydration (Wheless 2001; Freeman et al.

2006), these diets are associated with enhanced diuresis. We observed the opposite

with animals fed the higher-fat diet losing less water than animals fed the LFD. Third,

diets higher in protein require more water for the removal of nitrogenous wastes (Calder

and Braun 1983), these do not account for the observations herein. There is a small

difference in the amount of dietary protein (18.9% in SD versus 19.5% in LFD). This

small difference is unlikely to result in a difference in water loss and therefore hydration

status.

While food and water intake were not measured, a potential mechanism for the

observed differences could be that animals fed the LFD simply eat more than animals

fed a diet higher in fat. While the diets are closely matched in terms of available energy

per unit mass, the LFD is slightly less energy dense. Additionally, it's possible that there

are substantial differences in the non-nutritive fraction of the diet that could explain the

observations. While the current study does not disentangle this potential aspect, future

work could. These differences suggest that animals fed the LFD may be ingesting more

food mass to maintain nutritional homeostasis and with food intake increased there may

also be increased water intake (Bachmanov et al. 2002). While we have no evidence for

this, an increased rate of consumption would potentially result in an increased rate of

urine production and the production of water-containing fecal material. Given that the

animals fed the LFD do seem to be more dehydrated than animals fed the SD

(increased Sodium, osmolarity, HCT, Figure 2), this hypothesis requires that animals

fed the LFD are consuming more water, but not enough to satisfy their physiological

needs. Given that the animals have free access to water, an unmet need is

irreconcilable without further explanation.

Another intriguing hypothesis linked to the concept of unmet hydrational needs

has to do with thirst pathways. Thirst is a complex phenotype under tight neural control.

In brief, thirst is triggered in response to a reduction in blood volume or an increase in

serum Sodium (Thornton 2010). Osmo- and baroreceptors located in critical parts of the

vasculature are responsible for monitoring water balance, and when triggered, set into

motion a hormonal cascade that simultaneously enhance renal water retention and

stimulate thirst (Thornton 2010; Leib et al. 2016). While the former aspects of the

response are crucial to a large number of physiological processes, the latter response,

thirst, could be responsible for the observations described above. While purely a

hypothesis, there are several examples of species losing parts of their sensory systems

via natural selection such as loss of sight (Gore et al., 2018), decreases in certain taste

sensitivities or loss of taste entirely (Jiang et al. 2012), and loss of specific olfactory

functions (Kishida et al. 2015), resulting in dramatic changes from the habitat in which

they have evolved. For animals in deserts where water is largely absent, could well-

developed pathways leading to thirst be similarly changed? While requiring careful

consideration, perhaps using a comparative genomics approach, this hypothesis could

explain why animals with free access to water remain significantly dehydrated.

Respiratory Quotient

RQ, the ratio of CO_2 produced to O_2 consumed, can inform rates of differential fuel

utilization. RQ values typically fall between 0.7 and 1.0, with the catabolism of fats

yielding 0.7, carbohydrates 1.0, and proteins intermediate values (Kleiber 1975). RQ

values exceed 1.0 when anabolism of fat exceeds catabolism (*i.e.,* DNL Benedict 1937;

Abreu-Vieira et al. 2015; Levin et al. 2017) or during anaerobic exercise (Whipp 2007;

Zagatto et al. 2012). RQ was greater than 1.0 during the inactive period in animals on

both low and standard diets suggesting that *de novo* lipogenesis (DNL) is not limited by

fat intake in this species. Here and in previous experiments on *P. eremicus* (Colella et

al. 2021a) RQ is only greater than 1.0 during the light phase when animals are inactive,

which strongly suggests DNL. While carbohydrates, fats, or proteins are all burned as

fuel, if present in excess, they can also be converted to fat which can be stored for

future use (Barboza et al. 2009). DNL is one mechanism for converting non-fat energy

into a form that can be stored. The conversion of glucose to fats prior to entry into the

citric acid cycle is exothermic and results in an energetic cost, representing a

metabolically inefficient use of dietary substrate in cold habitats (Solinas et al. 2015),

but a potentially efficient mechanism of heat regulation in hot deserts. Further, fat

storage could be a way for desert mammals to buffer against an uncertain future where

water may be less available, as fat oxidation releases water (Mellanby 1942; Davidson

and Passmore 1963; Schmidt-Nielsen and Adolph 1964). Given this, the conversion of

other fuel sources to fats may serve as an important reservoir of metabolic water.

Previous studies have shown that DNL can occur under high-carbohydrate and high-fat

diets (Strable and Ntambi 2010). This is supported by our results, as both groups had

an RQ greater than 1.0 during the light phase and RQ values at or below the FQ during

the dark phase, which indicates a shift to the supplied dietary substrate as the main

oxidative fuel during cooler, dark periods.

Conclusion

Desert-adapted species offer a unique opportunity to study how organisms survive

some of the harshest habitats on Earth. The main challenges to survival in desert

habitats are maintaining water balance and thermoregulation, with dietary composition

being critically important, particularly to the former. Examining the physiological

responses of animals fed diets with different fat content aids in our understanding of

how animals may respond to changing environments where food availability and

temperatures may change. This will ultimately be important in predicting an animal's

response to climate change because diet impacts water balance.

To understand ways in which the desert-adapted cactus mouse, *P. eremicus,*

responds to variation in dietary fat content, reproductively mature mice were subjected

to one of two diets, an SD or an LFD. In general, EE and RWL for mice on the SD

during the light phase (a measurement comparable to the resting metabolic rate) agreed

with earlier studies of *Peromyscus* (Withers 1992; Ramirez et al. 2022), although the

experimental design presented here more closely represents the diurnal conditions

under which animals have evolved. We show that dietary fat is important for water

regulation in the desert-adapted cactus mouse through the analysis of interacting

physiological traits: 1) high rates of total water loss during the warmer, drier light phase

and 2) a shift in serum electrolyte values indicating dehydration. These factors suggest

that a low-fat diet could limit the capacity of desert animals to tolerate limited access to

free water as is common in arid environments.

Although we were able to collect hundreds of measurements for each diet, sex,

and time interval, our work has several limitations. Future studies should further modify

the major diet components (carbohydrates and proteins) and assess the seasonal

availability of such macronutrients in natural environments. Additionally, the

incorporation of food and water intake, activity monitoring, and body temperature will

provide further insight into the capacity of *P. eremicus* to tolerate environmental change.

The identification of short-term, physiological responses of *P. eremicus*

associated with diet composition demonstrates that adaptations that evolved over

longer evolutionary timescales affect survival during the accelerated time scales of

climate change. In light of global climate change and increased desertification,

investigating the range and mechanisms of plastic response of desert-adapted species

with regard to diet composition could provide insights into physiological responses to

increasingly erratic climate.

Acknowledgments

We would like to thank members of the MacManes lab for helpful comments and

support on previous versions of the manuscript; the Animal Resources Office and

veterinary care staff at the University of New Hampshire; A. Gerson at the University of

Massachusetts Amherst, Z. Cheviron at the University of Montana, B. Joos and J. Klok

at Sable Systems International for analytic guidance, technical support, and

respirometry training. This work was supported by the National Institute of Health

National Institute of General Medical Sciences (R35 GM128843 to M.D.M.).

Author Contributions

Conceptualization: M.D.M.; Methodology: D.M.B., M.D.M.; Formal analysis: D.M.B.,

E.L.; Investigation: D.M.B., J.P.C.; Resources: M.D.M.; Writing - original draft: D.M.B.;

Writing - review & editing: D.M.B., J.P.C., E.L., M.D.M.; Visualization: D.M.B., E.L.;

Supervision: M.D.M.; Project administration: M.D.M.; Funding acquisition: M.D.M.

Competing Interests

No competing interests declared.

Data Availability

Raw ExpeData (SSI) files are available through Zenodo:

https://zenodo.org/record/6422231#.YIXSD9PML0o. Macro processing files, raw and

processed respirometry data, and cage sampling scheme files are also available on

Zenodo. All R scripts used in this project are available through GitHub at:

https://github.com/DaniBlumstein/Diet_paper.

References

Abdel-Rahman M.K. 2010. Influence of dietary fat on renal function, lipid profile, sex hormones,
and electrolyte balance in rats. European Journal of Lipid Science and Technology
112:1166–1172.

Abreu-Vieira G., C. Xiao, O. Gavrilova, and M.L. Reitman. 2015. Integration of body
temperature into the analysis of energy expenditure in the mouse. Mol Metab 4:461–
470.

Acheson K.J. 1990. Sympathetic Nervous System in the Regulation of Thermogenesis. Pp. 40–
46 in M.J. Müller, E. Danforth, A.G. Burger, and U. Siedentopp eds. Hormones and
Nutrition in Obesity and Cachexia. Springer, Berlin, Heidelberg.

Bachmanov A.A., D.R. Reed, G.K. Beauchamp, and M.G. Tordoff. 2002. Food Intake, Water
Intake, and Drinking Spout Side Preference of 28 Mouse Strains. Behav Genet 32:435–
443.

Barboza P.S., K.L. Parker, and I.D. Hume, eds. 2009. Lipids: Fatty Acids and Adipose Tissue.
Pp. 119–131 in Integrative Wildlife Nutrition. Springer, Berlin, Heidelberg.

Beale P.K., K.J. Marsh, W.J. Foley, and B.D. Moore. 2018. A hot lunch for herbivores:
physiological effects of elevated temperatures on mammalian feeding ecology. Biol Rev
93:674–692.

Bedford N.L. and H.E. Hoekstra. 2015. Peromyscus mice as a model for studying natural
variation. eLife 4:e06813.

Beever E.A., C. Ray, P.W. Mote, and J.L. Wilkening. 2010. Testing alternative models of
climate-mediated extirpations. Ecological Applications 20:164–178.

Benedict F.G. 1937. Lipogenesis in the animal body, with special reference to the physiology of
the goose. Carnegie institution of Washington.

Bittner N.K.J., K.L. Mack, and M.W. Nachman. 2021. Gene expression plasticity and desert
adaptation in house mice. Evolution 75:1477–1491.

Boogert N.J., J.R. Madden, J. Morand-Ferron, and A. Thornton. 2018. Measuring and
understanding individual differences in cognition. Philos Trans R Soc Lond B Biol Sci
373:20170280.

Boumansour L., N. Benhafri, G. Guillon, M. Corbani, H. Touati, A. Dekar-Madoui, and S. Ouali-
Hassenaoui. 2021. Vasopressin and oxytocin expression in hypothalamic supraoptic
nucleus and plasma electrolytes changes in water-deprived male Meriones libycus.
Animal Cells and Systems 0:1–10.

Bourque C.W. 2008. Central mechanisms of osmosensation and systemic osmoregulation.
Nature Reviews Neuroscience 9:519–531.

Bradford D.F. 1974. Water Stress of Free-Living Peromyscus Truei. Ecology 55:1407–1414.

Bradley W.G. and R.A. Mauer. 1973. Rodents of a Creosote Bush Community in Southern
Nevada. The Southwestern Naturalist 17:333–344.

Calder W.A. and E.J. Braun. 1983. Scaling of osmotic regulation in mammals and birds. Am J
Physiol 244:R601-606.

Castle S.T., N. Allan, D. Clifford, C.M. Aylward, J. Ramsey, A.J. Fascetti, R. Pesapane, et al.
2020. Diet composition analysis provides new management insights for a highly
specialized endangered small mammal. (B.-S. Yue, ed.)PLoS ONE 15:e0240136.

Cheuvront S.N., R.W. Kenefick, S.J. Montain, and M.N. Sawka. 2010. Mechanisms of aerobic
performance impairment with heat stress and dehydration. Journal of Applied Physiology
109:1989–1995.

Christian D.P. and Matson J.O. 1978. Comparative water balance in two species of Liomys
(Rodentia: Heteromyidae). Comparative Biochemistry and Physiology A.

Colella J.P., D.M. Blumstein, and M.D. MacManes. 2021a. Disentangling environmental drivers
of circadian metabolism in desert-adapted mice. Journal of Experimental Biology 224.

Colella J.P., A. Tigano, O. Dudchenko, A.D. Omer, R. Khan, I.D. Bochkov, E.L. Aiden, et al.
2021b. Limited Evidence for Parallel Evolution Among Desert-Adapted Peromyscus
Deer Mice. Journal of Heredity 112:286–302.

Cortés A., M. Rosenmann, and F. Bozinovic. 2000. Water economy in rodents: evaporative
water loss and metabolic water production. Rev chil hist nat 73.

Crossland J.P., M.J. Dewey, S.C. Barlow, P.B. Vrana, M.R. Felder, and G.J. Szalai. 2014.
Caring for Peromyscus spp. in research environments. Lab Anim 43:162–166.

Davidson S. and R. Passmore. 1963. Human nutrition and dietetics. Human nutrition and
dietetics.

Fitzsimons J.T. 1961. Drinking by rats depleted of body fluid without increase in osmotic
pressure. The Journal of Physiology 159:297.

Fox J. and S. Weisberg. 2019. An {R} Companion to Applied Regression (Third.). Sage.

Frank C.L. 1988. Diet Selection by a Heteromyid Rodent: Role of Net Metabolic Water
723   Production. Ecology 69:1943–1951.

724   Freeman J., P. Veggiotti, G. Lanzi, A. Tagliabue, and E. Perucca. 2006. The ketogenic diet:
725   from molecular mechanisms to clinical effects. Epilepsy Res 68:145–80.

726   Friedman A.N., L.G. Ogden, G.D. Foster, S. Klein, R. Stein, B. Miller, J.O. Hill, et al. 2012.
727   Comparative effects of low-carbohydrate high-protein versus low-fat diets on the kidney.
728   Clin J Am Soc Nephrol 7:1103–1111.

729   Fuller A., S.K. Maloney, D. Blache, and C. Cooper. 2020. Endocrine and metabolic
730   consequences of climate change for terrestrial mammals. Current Opinion in Endocrine
731   and Metabolic Research 11:9–14.

732   Garland T. Jr, H. Schutz, M.A. Chappell, B.K. Keeney, T.H. Meek, L.E. Copes, W. Acosta, et al.
733   2011. The biological control of voluntary exercise, spontaneous physical activity and
734   daily energy expenditure in relation to obesity: human and rodent perspectives. Journal
735   of Experimental Biology 214:206–229.

736   Gilman A. 1937. The relation between blood osmotic pressure, fluid distribution and voluntary
737   water intake. American Journal of Physiology-Legacy Content 120:323–328.

738   Giorello F.M., M. Feijoo, G. D'Elía, D.E. Naya, L. Valdez, J.C. Opazo, and E.P. Lessa. 2018. An
739   association between differential expression and genetic divergence in the Patagonian
740   olive mouse (Abrothrix olivacea). Molecular Ecology 27:3274–3286.

741   Goodfriend T.L. 2006. Aldosterone—a hormone of cardiovascular adaptation and
742   maladaptation. The Journal of Clinical Hypertension 8:133–139.

743   Hart J.S. 1971. Rodents. Pp. 1–149 in Mammals. Elsevier.

744   Heimeier R.A., R.C. Bartolo, and J.A. Donald. 2004. The effect of water deprivation on signalling
745   molecules that utilise cGMP in the spinifex hopping mouse Notomys alexis. Aust
746   Mammalogy 26:191–198.

747   Heimeier R.A. and J.A. Donald. 2006. The response of the natriuretic peptide system to water
748   deprivation in the desert rodent, Notomys alexis. Comparative Biochemistry and
749   Physiology Part A: Molecular & Integrative Physiology 143:193–201.

750   Henry P., Z. Sim, and M.A. Russello. 2012. Genetic Evidence for Restricted Dispersal along
751   Continuous Altitudinal Gradients in a Climate Change-Sensitive Mammal: The American
752   Pika. PLOS ONE 7:e39077.

753   Hino J., M. Nakatani, Y. Arai, K. Tsuchida, M. Shirai, M. Miyazato, and K. Kangawa. 2017.
754   Overexpression of bone morphogenetic protein-3b (BMP-3b) in adipose tissues protects
755   against high-fat diet-induced obesity. Int J Obes 41:483–488.

Hope A. and R. Parmenter. 2007. Food Habits of Rodents Inhabiting Arid and Semi-arid
Ecosystems of Central New Mexico. Special Publications.

IPCC. 2019. Global Warming of 1.5 C An IPCC Special Report on the Impacts of Global
Warming of 1.5 C Above Pre-Industrial Levels and Related Global Greenhouse Gas
Emission Pathways, in the Context of Strengthening the Global Response to the Threat
of Climate Change. Sustainable Development, and Efforts to Eradicate Poverty
https://www ipcc ch/sr15/ Accessed 1.

Jiang P., J. Josue, X. Li, D. Glaser, W. Li, J.G. Brand, R.F. Margolskee, et al. 2012. Major taste
loss in carnivorous mammals. Proceedings of the National Academy of Sciences
109:4956–4961.

Jirimutu, Z. Wang, G. Ding, G. Chen, Y. Sun, Z. Sun, H. Zhang, et al. 2012. Genome sequences
of wild and domestic bactrian camels. Nat Commun 3:1202.

Jones M.R., L.S. Mills, P.C. Alves, C.M. Callahan, J.M. Alves, D.J.R. Lafferty, F.M. Jiggins, et
al. 2018. Adaptive introgression underlies polymorphic seasonal camouflage in
snowshoe hares. Science 360:1355–1358.

Kaiyala K.J., G.J. Morton, J.P. Thaler, T.H. Meek, T. Tylee, K. Ogimoto, and B.E. Wisse. 2012.
Acutely Decreased Thermoregulatory Energy Expenditure or Decreased Activity Energy
Expenditure Both Acutely Reduce Food Intake in Mice. PLOS ONE 7:e41473.

Kaufmann H., L. Norcliffe-Kaufmann, and J.-A. Palma. 2020. Baroreflex Dysfunction. New
England Journal of Medicine 382:163–178.

Kishida T., J. Thewissen, T. Hayakawa, H. Imai, and K. Agata. 2015. Aquatic adaptation and the
evolution of smell and taste in whales. Zoological Letters 1:9.

Kleiber M. 1975. The fire of life. An introduction to animal energetics.

Kordonowy L., K.D. Lombardo, H.L. Green, M.D. Dawson, E.A. Bolton, S. LaCourse, and M.D.
MacManes. 2017. Physiological and biochemical changes associated with acute
experimental dehydration in the desert adapted mouse, *Peromyscus eremicus*. Physiol
Rep 5:e13218.

Kordonowy L. and M. MacManes. 2017. Characterizing the reproductive transcriptomic
correlates of acute dehydration in males in the desert-adapted rodent, Peromyscus
eremicus. BMC Genomics 18:473.

Kordonowy L.L. and M.D. MacManes. 2016. Characterization of a male reproductive
transcriptome for *Peromyscus eremicus* (Cactus mouse). PeerJ 4:e2617.

Kortenoeven M.L.A. and R.A. Fenton. 2014. Renal aquaporins and water balance disorders.
Biochimica et Biophysica Acta (BBA) - General Subjects, Aquaporins 1840:1533–1549.

Kutscher C. 1968. Plasma volume change during water-deprivation in gerbils, hamsters, guinea
pigs and rats. Comparative Biochemistry and Physiology 25:929–936.

Leib D.E., C.A. Zimmerman, and Z.A. Knight. 2016. Thirst. Current Biology 26:R1260–R1265.

Levin E., G. Lopez-Martinez, B. Fane, and G. Davidowitz. 2017. Hawkmoths use nectar sugar to
reduce oxidative damage from flight. Science 355:733–735.

Lighton J.R.B. 2018. Measuring Metabolic Rates: A Manual for Scientists (2nd ed.). Oxford
University Press.

Lin X. and D. Zhang. 1999. Inference in Generalized Additive Mixed Models by Using
Smoothing Splines. Journal of the Royal Statistical Society Series B (Statistical
Methodology) 61:381–400.

Loarie S.R., P.B. Duffy, H. Hamilton, G.P. Asner, C.B. Field, and D.D. Ackerly. 2009. The
velocity of climate change. Nature 462:1052–1055.

Louw G. N. and Seely M. K. 1984. Ecology of desert organisms. Vegetatio 57:142–142.

MacManes M.D. 2017. Severe acute dehydration in a desert rodent elicits a transcriptional
response that effectively prevents kidney injury. Renal Physiol 11.

MacManes M.D. and M.B. Eisen. 2014. Characterization of the transcriptome, nucleotide
sequence polymorphism, and natural selection in the desert adapted mouse
Peromyscus eremicus. PeerJ 2:e642.

Macmillen R.E. 1965. Aestivation in the cactus mouse, Peromyscus eremicus. Comparative
Biochemistry and Physiology 16:227–248.

MacMillen R.E. 1972. Water economy of nocturnal desert rodents.

MacMillen R.E. and D.S. Hinds. 1983. Water Regulatory Efficiency in Heteromyid Rodents: A
Model and Its Application. Ecology 64:152–164.

Manlick P.J., K. Maldonado, and S.D. Newsome. 2021. Competition shapes individual foraging
and survival in a desert rodent ensemble. Journal of Animal Ecology n/a.

Marra N.J., S.H. Eo, M.C. Hale, P.M. Waser, and J.A. DeWoody. 2012. A priori and a posteriori
approaches for finding genes of evolutionary interest in non-model species:
Osmoregulatory genes in the kidney transcriptome of the desert rodent Dipodomys
spectabilis (banner-tailed kangaroo rat). Comparative Biochemistry and Physiology Part
D: Genomics and Proteomics 7:328–339.

Marra N.J., A. Romero, and J.A. DeWoody. 2014. Natural selection and the genetic basis of
osmoregulation in heteromyid rodents as revealed by RNA-seq. Molecular Ecology
23:2699–2711.

McArthur E.D. 1994. Nutritive Quality and Mineral Content of Potential Desert Tortoise Food
Plants. Intermountain Research Station.

Mellanby K. 1942. Metabolic Water and Desiccation. Nature 150:21–21.

Meserve P.L. 1976. Habitat and Resource Utilization by Rodents of a California Coastal Sage
827   Scrub Community. Journal of Animal Ecology 45:647–666.

828   Mirzabaev A., J. Wu, J. Evans, F. García-Oliva, I.A.G. Hussein, M.H. Iqbal, J. Kimutai, et al.
829   2019. Desertification. In: Climate Change and Land: an IPCC special report on climate
830   change, desertification, land degradation, sustainable land management, food security,
831   and greenhouse gas fluxes in terrestrial ecosystems.

832   Moritz C., J.L. Patton, C.J. Conroy, J.L. Parra, G.C. White, and S.R. Beissinger. 2008. Impact of
833   a Century of Climate Change on Small-Mammal Communities in Yosemite National
834   Park, USA. Science 322:261–264.

835   Morrison L., A. Estrada, and R. Early. 2018. Species traits suggest European mammals facing
836   the greatest climate change are also least able to colonize new locations. (Y. Wiersma,
837   ed.)Divers Distrib 24:1321–1332.

838   Murie M. 1961. Metabolic Characteristics of Mountain, Desert and Coastal Populations of
839   Perormyscus. Ecology 42:723–740.

840   Nagy K.A. 1987. Field Metabolic Rate and Food Requirement Scaling in Mammals and Birds.
841   Ecological Monographs 57:111–128.

842   Noble J.D., S.L. Collins, A.J. Hallmark, K. Maldonado, B.O. Wolf, and S.D. Newsome. 2019.
843   Foraging strategies of individual silky pocket mice over a boom–bust cycle in a
844   stochastic dryland ecosystem. Oecologia 190:569–578.

845   Orr T.J., S.D. Newsome, and B.O. Wolf. 2015. Cacti supply limited nutrients to a desert rodent
846   community. Oecologia 178:1045–1062.

847   Pardi M.I., R.C. Terry, E.A. Rickart, and R.J. Rowe. 2020. Testing climate tracking of montane
848   rodent distributions over the past century within the Great Basin ecoregion. Global
849   Ecology and Conservation 24:e01238.

850   Parmesan C. 2006. Ecological and Evolutionary Responses to Recent Climate Change. Annual
851   Review of Ecology, Evolution, and Systematics 37:637–669.

852   Parmesan C. and G. Yohe. 2003. A globally coherent fingerprint of climate change impacts
853   across natural systems. Nature 421:37–42.

854   Pecl G.T., M.B. Araújo, J.D. Bell, J. Blanchard, T.C. Bonebrake, I.-C. Chen, T.D. Clark, et al.
855   2017. Biodiversity redistribution under climate change: Impacts on ecosystems and
856   human well-being. Science 355:eaai9214.

857   Pergams O.R.W. and J.J. Lawler. 2009. Recent and Widespread Rapid Morphological Change
858   in Rodents. PLOS ONE 4:e6452.

859   Pool A.-H., T. Wang, D.A. Stafford, R.K. Chance, S. Lee, J. Ngai, and Y. Oka. 2020. The
860   cellular basis of distinct thirst modalities. Nature 588:112–117.

Porter W.P. and D.M. Gates. 1969. Thermodynamic equilibria of animals with environment.
Ecological monographs 39:227–244.

Pyke G.H. 1984. Optimal Foraging Theory: A Critical Review. Annu Rev Ecol Syst 15:523–575.

Pyke G.H., H.R. Pulliam, and E.L. Charnov. 1977. Optimal Foraging: A Selective Review of
Theory and Tests. The Quarterly Review of Biology 52:137–154.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Ramirez R.W., E.A. Riddell, S.R. Beissinger, and B.O. Wolf. 2022. Keeping your cool:
thermoregulatory performance and plasticity in desert cricetid rodents. Journal of
Experimental Biology 225:jeb243131.

Rasouli M. 2016. Basic concepts and practical equations on osmolality: Biochemical approach.
Clinical Biochemistry 49:936–941.

Reece W.O., H.H. Erickson, J.P. Goff, and E.E. Uemura. 2015. Dukes' Physiology of Domestic
Animals. John Wiley & Sons.

Reed T.E., D.E. Schindler, and R.S. Waples. 2011. Interacting Effects of Phenotypic Plasticity
and Evolution on Population Persistence in a Changing Climate. Conservation Biology
25:56–63.

Reichman O.J. 1975. Relation of Desert Rodent Diets to Available Resources. Journal of
Mammalogy 56:731–751.

Reid W.T. 1987. *A critical analysis of Eastern Mojave desert temperatures* (Thesis). California
State University, Northridge.

Riddell E.A., K.J. Iknayan, L. Hargrove, S. Tremor, J.L. Patton, R. Ramirez, B.O. Wolf, et al.
2021. Exposure to climate change drives stability or collapse of desert mammal and bird
communities. Science 371:633–636.

Russell V. Lenth. 2021. emmeans: Estimated Marginal   Means, aka Least-Squares Means. R
package version 1.6.0.   https://CRAN.R-project.org/package=emmeans.

Sánchez-Peña M.J., F. Márquez-Sandoval, A.C. Ramírez-Anguiano, S.F. Velasco-Ramírez, G.
Macedo-Ojeda, and L.J. González-Ortiz. 2017. Calculating the metabolizable energy of
macronutrients: a critical review of Atwater's results. Nutrition Reviews 75:37–48.

Schmidt-Nielsen B. and K. Schmidt-Nielsen. 1951. A complete account of the water metabolism
in kangaroo rats and an experimental verification. Journal of Cellular and Comparative
Physiology 38:165–181.

Schmidt-Nielsen K. 1975. Desert Rodents: Physiological Problems of Desert Life. Pp. 379–388
in I. Prakash and P.K. Ghosh eds. Rodents in Desert Environments, Monographiae
Biologicae. Springer Netherlands, Dordrecht.

Schmidt-Nielsen K. and E.F. Adolph. 1964. Desert Animals: Physiological Problems of Heat and
Water. Physiological Zoology 37:338–339.

Secor S.M. 2009. Specific dynamic action: a review of the postprandial metabolic response.
Journal of Comparative Physiology B 179:1–56.

Sgrò C.M., A.J. Lowe, and A.A. Hoffmann. 2011. Building evolutionary resilience for conserving
biodiversity under climate change. Evolutionary Applications 4:326–337.

Sheppard P.R., A.C. Comrie, G.D. Packin, K. Angersbach, and M.K. Hughes. 2002. The climate
of the US Southwest. Climate Research 21:219–238.

Sikes R.S. and the Animal Care and Use Committee of the American Society of Mammalogists.
2016. 2016 Guidelines of the American Society of Mammalogists for the use of wild
mammals in research and education. J Mammal 97:663–688.

Solinas G., J. Borén, and A.G. Dulloo. 2015. De novo lipogenesis in metabolic homeostasis:
More friend than foe? Molecular Metabolism 4:367–377.

Speakman J. 1997. Factors influencing the daily energy expenditure of small mammals.
Proceedings of the Nutrition Society 56:1119–1136.

Spielman D., B.W. Brook, and R. Frankham. 2004. Most species are not driven to extinction
before genetic factors impact them. PNAS 101:15261–15264.

Strable M.S. and J.M. Ntambi. 2010. Genetic control of de novo lipogenesis: role in diet-induced
obesity. Crit Rev Biochem Mol Biol 45:199–214.

Stricker E.M. 1966. Extracellular fluid volume and thirst. American Journal of Physiology-Legacy
Content 211:232–238.

Thornton S.N. 2010. Thirst and hydration: Physiology and consequences of dysfunction.
Physiology & Behavior, Beverages and Health 100:15–21.

Tigano A., J.P. Colella, and M.D. MacManes. 2020. Comparative and population genomics
approaches reveal the basis of adaptation to deserts in a small rodent. Molecular
Ecology 29:1300–1314.

Tracy R.L. and G.E. Walsberg. 2002. Kangaroo rats revisited: re-evaluating a classic case of
desert survival. Oecologia 133:449–457.

Unland H.E., P.R. Houser, W.J. Shuttleworth, and Z.-L. Yang. 1996. Surface flux measurement
and modeling at a semi-arid Sonoran Desert site. Agricultural and Forest Meteorology
82:119–153.

Van der Putten W.H., M. Macel, and M.E. Visser. 2010. Predicting species distribution and
abundance responses to climate change: why it is essential to include biotic interactions
across trophic levels. Phil Trans R Soc B 365:2025–2034.

Veal R. and W. Caire. 1979. Peromyscus eremicus. Mammalian Species 1–6.

Walsberg G.E. 2000. Small Mammals in Hot Deserts: Some Generalizations Revisited.
BioScience 50:109.

Walsh R.E., A.P. Aprígio Assis, J.L. Patton, G. Marroig, T.E. Dawson, and E.A. Lacey. 2016.
Morphological and dietary responses of chipmunks to a century of climate change.
Global Change Biology 22:3233–3252.

Westerterp K.R. 2004. Diet induced thermogenesis. Nutrition & metabolism 1:1–5.

Wheless J.W. 2001. The ketogenic diet: an effective medical therapy with side effects. J Child
Neurol 16:633–635.

Whipp B.J. 2007. Physiological mechanisms dissociating pulmonary CO_2 and O_2 exchange
dynamics during exercise in humans. Experimental Physiology 92:347–355.

Whitfield M.C., B. Smit, A.E. McKechnie, and B.O. Wolf. 2015. Avian thermoregulation in the
heat: scaling of heat tolerance and evaporative cooling capacity in three southern
African arid-zone passerines. Journal of Experimental Biology 218:1705–1714.

Withers P.C. 1982. Effect of diet and assimilation efficiency on water balance for two desert
rodents. Journal of Arid Environments 5:375–384.

______. 1992. Comparative animal physiology. Saunders College Pub. Philadelphia.

Wolf A.V. 1950. Osmometric analysis of thirst in man and dog. American Journal of Physiology-
Legacy Content 161:75–86.

Wolf B.O. and C.M. del Rio. 2003. How important are columnar cacti as sources of water and
nutrients for desert consumers? A review. Isotopes Environ Health Stud 39:53–67.

Wood S. 2017. Generalized Additive Models: An Introduction with R, Second Edition.

Wu H., X. Guang, M.B. Al-Fageeh, J. Cao, S. Pan, H. Zhou, L. Zhang, et al. 2014. Camelid
genomes reveal evolution and adaptation to desert environments. Nature
Communications 5:1–10.

Yang J., W.-R. Li, F.-H. Lv, S.-G. He, S.-L. Tian, W.-F. Peng, Y.-W. Sun, et al. 2016. Whole-
Genome Sequencing of Native Sheep Provides Insights into Rapid Adaptations to
Extreme Environments. Mol Biol Evol 33:2576–2592.

Zagatto A.M., W.E. Miyagi, R.L. Sakugawa, E.I. Kaminagakura, and M. Papoti. 2012. Aerobic
Endurance Measurement by Respiratory Exchange Ratio during a Cycle Ergometer
Graded Exercise Test. Journal of Exercise Physiology Online 15:49–56.

Zimmerman C.A., D.E. Leib, and Z.A. Knight. 2017. Neural circuits underlying thirst and fluid
homeostasis. Nature Reviews Neuroscience 18:459–469.

Supplemental

Table S1: Number of samples collected per time interval.

time interval	diet	number of 120 intervals male	number 120 intervals female
light	standard	867	879
	low fat	914	927
dark	standard	758	755
	low fat	758	719

967
968

969 Table S2: Generalized additive mixed models (GAMM) 24-hour results. **A)** Statical
970 model and results for rate of water loss (RWL, H_2O g/hr^{-1}), **B)** model and results for
971 measurements of energy expenditure (EE kJ/hr^{-1}), **C)** model and results for respiratory
972 quotient (RQ).
973

A)

H2Og~diet+Sex+s(ctime.exp,k=3)+s(time_in_D,bs=cc)+s(EE_kJH)+s(RQ)

	Estimate	Std.	Error	t value	Pr(>|t|)
(Intercept)	0.121	0.008	15.582	<2.00E-16	***
diet	-0.038	0.009	-4.253	0.000	***
Sex	0.027	0.009	3.004	0.003	**

	edf	Ref.df	F	p-value	
s(ctime.exp)	1.992	1.992	183.99	<2e-16	***
s(time_in_D)	7.948	8	668.73	<2e-16	***
s(EE_kJH)	6.565	6.565	20.69	<2e-16	***
s(RQ)	8.422	8.422	37.87	<2e-16	***

R-sq.(adj) = 0.561
Scale est.=0.0016156   n=7908

B)

log(EE_kJH)~diet+Sex+s(ctime.exp,k=3)+s(time_in_D,bs=cc)+s(H2Og)+s(RQ)

	Estimate	Std.	Error	t value	Pr(>|t|)
(Intercept)	-0.542	0.063	-8.628	<2e-16	***
diet	-0.043	0.073	-0.585	0.558	
SexM	0.097	0.073	1.332	0.183	

	edf	Ref.df	F	p-value	
s(ctime.exp)	1.980	1.980	39.660	<2e-16	***
s(time_in_D)	7.976	8.000	1464.100	<2e-16	***
s(H2Og)	7.114	7.114	28.930	<2e-16	***
s(RQ)	8.136	8.136	178.830	<2e-16	***

R-sq.(adj)=0.678
Scale est.=0.076286   n=7908

C)

log(RQ)~diet+Sex+s(ctime.exp,k=3)+s(time_in_D,bs=cc)+s(H2Og)+s(log(EE_kJH))

	Estimate	Std.	Error	t value	Pr(>|t|)
(Intercept)	-0.024	0.034	-0.706	0.480	
diet	-0.025	0.039	-0.641	0.522	
Sex	-0.008	0.039	-0.210	0.834	

	edf	Ref.df	F	p-value	
s(ctime.exp)	1.593	1.593	27.450	1.98E-07	***
s(time_in_D)	7.693	8.000	44.530	< 2.00E-16	***
s(H2Og)	7.397	7.397	47.360	< 2.00E-16	***
s(log(EE_kJH))	7.160	7.160	359.320	< 2.00E-16	***

R-sq.(adj)=0.393
Scale est.=0.0070625   n=7908

Figure S1: Contributions of model terms to energy expenditure (EE kJ h-1, A-D), rate of
water loss (RWL g h-1, E-H), and respiratory quotient (RQ, I-L) for female and male
Peromyscus eremicus across two diet groups (standard diet and low-fat diet) in general
additive mixed models (GAMM). The smoothing curves for each response variable
included two fixed effects; diet (low fat vs standard) and sex, two random effects; mouse
identification number and date of data collection, and four regression terms: time in
days, diurnal cycle, and two of the three respirometry response variables (EE, RQ,
and/or EWL) as regression terms. For each graph, the y-axis is the effect of the x-axis
on the respirometry response variable estimated by a multivariable GAMM. Blue-
shaded areas are 95% confidence intervals, black points are residuals.
986

987 Table S3: Generalized additive mixed models (GAMM) results for only data collected
988 during the light phase. A) Statical model and results for rate of water loss (RWL, H2O
989 g/hr-1), B) model and results for measurements of energy expenditure (EE kJ/hr-1), C)
990 model and results for respiratory quotient (RQ).
991

A)

H2Og~diet+Sex+s(ctime.exp,k=3)+s(EE_kJH)+s(RQ)

	Estimate	Std.	Error	t value	Pr(>|t|)
(Intercept)	0.15503	0.00756	20.506	<2.00E-16	***
diet	-0.058941	0.008737	-6.746	1.69E-11	***
Sex	0.035545	0.008736	4.069	4.80E-05	***

	edf	Ref.df	F	p-value	
s(ctime.exp)	1.974	1.974	68.25	<2e-16	***
s(EE_kJH)	5.797	5.797	10.8	<2e-16	***
s(RQ)	5.309	5.309	102.25	<2e-16	***

R-sq.(adj)=0.349
Scale est.=0.0023669   n=4918

B)

log(EE_kJH)~diet+Sex+s(ctime.exp,k=3)+s(H2Og)+s(RQ)

	Estimate	Std.	Error	t value	Pr(>|t|)
(Intercept)	-0.92542	0.09277	-9.975	<2e-16	***
diet	-0.06415	0.10738	-0.597	0.55	
Sex	0.11275	0.10719	1.052	0.293	

	edf	Ref.df	F	p-value	
s(ctime.exp)	1	1	6.991	0.00822	**
s(H2Og)	6.797	6.797	7.118	<2.00E-16	***
s(RQ)	5.875	5.875	285.938	<2.00E-16	***

R-sq.(adj)=0.189
Scale est.=0.12779   n=4918

B)

log(RQ)~diet+Sex+s(ctime.exp,k=3)+s(H2Og)+s(log(EE_kJH))

Parametric   coefficients:

	Estimate	Std.	Error	t value	Pr(>|t|)
(Intercept)	0.001688	0.036754	0.046	0.963	

diet	0.025869	0.042478	0.609	0.543
Sex	-0.014092	0.042453	-0.332	0.74

	edf	Ref.df	F	p-value	
s(ctime.exp)	1.43	1.43	18.81	1.27E-05	***
s(H2Og)	7.305	7.305	91.83	<2.00E-16	***
s(log(EE_kJH))	6.539	6.539	362.65	<2.00E-16	***

R-sq.(adj)=0.283

Scale est.=0.0084596   n=4918

Table S4: Generalized additive mixed models (GAMM) results for only data collected
during the dark phase. A) Statical model and results for rate of water loss (RWL, H2O
g/hr-1), B) model and results for measurements of energy expenditure (EE kJ/hr-1), C)
model and results for respiratory quotient (RQ).

A)

H2Og~diet+Sex+s(ctime.exp,k=3)+s(EE_kJH)+s(RQ)

	Estimate	Std.	Error	t value	Pr(>|t|)
(Intercept)	0.053382	0.00923	5.784	8.07E-09	***
diet	0.002749	0.01071	0.257	0.797	
Sex	0.025882	0.010637	2.433	0.015	*

	edf	Ref.df	F	p-value	
s(ctime.exp)	1.968	1.968	19.063	4.29E-07	***
s(EE_kJH)	4.881	4.881	8.185	6.41E-07	***
s(RQ)	7.887	7.887	47.955	<2.00E-16	***

R-sq.(adj)=0.219
Scale est.=0.0018757   n=2990

B)

log(EE_kJH)~experiment+Sex+s(ctime.exp,k=3)+s(H2Og)+s(RQ)

	Estimate	Std.	Error	t value	Pr(>|t|)
(Intercept)	0.08701	0.07403	1.175	0.24	
diet	-0.04856	0.0857	-0.567	0.571	
Sex	0.11419	0.08547	1.336	0.182	

	edf	Ref.df	F	p-value	
s(ctime.exp)	1.97	1.97	21.35	<2e-16	***
s(H2Og)	7.853	7.853	12.85	<2e-16	***
s(RQ)	7.469	7.469	34.63	<2e-16	***

R-sq.(adj)=0.12
Scale est.=0.058025   n=2990

C)

log(RQ)~experiment+Sex+s(ctime.exp,k=3)+s(H2Og)+s(log(EE_kJH))

Parametric   coefficients:

	Estimate	Std.	Error	t value

(Intercept)	-0.07254	0.02994	-2.423	0.0155	*
diet	-0.08263	0.03458	-2.39	0.0169	*
Sex	-0.01154	0.03459	-0.333	0.7388	

	edf	Ref.df	F	p-value	
s(ctime.exp)	1.95	1.95	20.1	<2e-16	***
s(H2Og)	6.454	6.454	45.5	<2e-16	***
s(log(EE_kJH))	1	1	39.76	<2e-16	***

R-sq.(adj)=0.292

Scale est.=0.0032848   n=2990

999
1000

1001   Table S5: ANCOVA results for total water loss.

	Sum	Sq	Df	F value	Pr(>F)
diet	130.36	1	19.5126	5.22E-05	***
sex	40.4	1	6.0471	0.01736	*
weight	7.03	1	1.0529	0.30967	
diet:sex	1.54	1	0.2305	0.63317	
Residuals	340.72	51			

contrast	estimate	SE	df	t.ratio	p.value
F Low Fat - M Low Fat	-2.05	0.977	51	-2.101	0.1667
F Low Fat - F Standard	2.76	1.018	51	2.716	0.0433
F Low Fat - M Standard	1.39	0.977	51	1.421	0.4922
M Low Fat - F Standard	4.82	1.014	51	4.748	0.0001
M Low Fat - M Standard	3.44	0.977	51	3.522	0.0049
F Standard - M Standard	-1.38	1.013	51	-1.359	0.5307

1002

low fat diet   standard diet