**High total water loss driven by low-fat diet in desert-adapted mice**

Danielle M. Blumstein[1], Jocelyn P. Colella[2], Ernst Linder[3], Matthew D. MacManes[1]

1. University of New Hampshire, Molecular, Cellular, and Biomedical Sciences Department,

Durham, NH 03824 (DMB Dani.Blumstein@unh.edu, Orcid ID: 0000-0003-2129-5541; MDM

Matthew.MacManes@unh.edu, Orcid ID: 0000-0002-2368-6960)

2. University of Kansas, Biodiversity Institute and Department of Ecology and Evolutionary

Biology, Lawrence, KS 66045 (JPC colella@ku.edu, Orcid ID: 0000-0003-2463-1029)

3. University of New Hampshire, Department of Mathematics and Statistics, Durham, NH 03824

(EL ernst.linder@unh.edu)

**Abstract**

Availability of food resources is an important driver of survival. Animals must either relocate or

adapt to persist in environments where critical resource abundance is changing.  An optimal diet

balances energy gain, water regulation, and nutrition. We used flow-through respirometry to

characterize metabolic phenotypes of the desert-adapted cactus mouse (*Peromyscus eremicus*)

under diurnally variable environmental conditions that mimic the environment of the Sonoran

Desert. We treated mice with two different energetically equivalent diets, a standard diet and a

low-fat diet, and measured energy expenditure, water loss rate, respiratory quotient, weight, and

electrolyte levels. Mice fed the low-fat diet lost significantly more water than those on the

standard diet. Our results suggest that cactus mice may have limited capacity to tolerate water

deprivation if optimal foods become less abundant. Given that climate change is predicted to

modify the distribution of food items, understanding these links may have important implications

for long-term population viability for desert and non-desert adapted animals alike.

**Keywords**: dietary fat, energy expenditure, *Peromyscus*, physiology, respirometry, total water

loss

**Introduction**

One of the most critical challenges faced by desert mammals is the regulation of water. Water balance is maintained through a cascade of hormones that elicit a response designed to increase water retention and decrease water use in the excretion of nitrogenous wastes. Dehydration, in contrast, is an outcome of inadequate water balance. This results in a decrease in blood volume and an increase in osmolality, the latter, an effect driven largely by the rise in serum sodium (Thornton 2010; Leib et al. 2016) and eventually leading to death if water homeostasis is not achieved. While hormonal regulation of water balance is paramount to survival, behavioral, physiological, and genomic responses are also critical.

Dietary intake of food is one important way in which animals manage water. There are three main macromolecules that animals ingest as food items; carbohydrates, fats, and proteins, all of which vary in their amount of energy and water, as well as in their cost to metabolize. Energy potential varies greatly, with carbohydrates and proteins yielding 16.74 J/g, while fats yield more than double that (37.66 J/g, Sánchez-Peña et al. 2017). Metabolic water, the water produced through endogenous catabolism of carbohydrates, fats, or proteins (Schmidt-Nielsen and Adolph 1964; Frank 1988; Orr et al. 2015 production also varies dramatically. Oxidation of carbohydrates yields 0.60 g of metabolic water per gram and fats yield 1.07 g per gram due to greater oxygen requirements of lipid metabolism, while proteins yield the least metabolic water (0.41 g) of any macronutrient and even require water loss for excretion of nitrogenous waste (Mellanby 1942; Davidson and Passmore 1963; Schmidt-Nielsen and Adolph 1964).

In environments like deserts, where extrinsic water is limited, both preformed (dietary) water and endogenous water production are key to survival. Dietary intake represents the raw material used to produce metabolic water. Food catabolism results in water loss through respiration, urination, and fecal production (Schmidt-Nielsen 1975). Specific dietary composition is often unknown and highly variable, particularly for wild animals, energy potential (Pyke et al. 1977; Pyke 1984), nutrition, and capacity for water production (Schmidt-Nielsen and Adolph 1964) all depend on diet composition. In deserts, vegetation, seeds, and insects comprise the bulk of rodent diets (Reichman 1975) with each varying in fat, protein, and carbohydrate composition and seasonal availability (Wolf and del Rio 2003; Orr et al. 2015). In response to natural variation in resource availability, animals have the opportunity to make dietary decisions that may have a substantial effect on their internal water economy.

Mice of the genus *Peromyscus* have the widest distribution of any North American

mammal and are considered model organisms in evolutionary biology due to their unparalleled

habitat diversity (Bedford and Hoekstra 2015), extensive genomic resources (*e.g.,* Tigano et al.

2020; Colella et al. 2021a), abundance of historical and contemporary collections in natural

history museums (Pergams and Lawler 2009; Pardi et al. 2020), and ability to live and breed

under laboratory conditions (Crossland et al. 2014). The desert specialist cactus mouse,

*Peromyscus eremicus*, is endemic to the southwestern United States and exhibits behavioral

(Murie 1961; Veal and Caire 1979), physiological (Macmillen 1965; Kordonowy et al. 2017;

Colella et al. 2021b), and molecular (MacManes 2017; Tigano et al. 2020) adaptations to desert

environments, making it an interesting natural experimental model to examine mechanisms of

adaptation to warmer, drier environments. *Peromyscus eremicus* are omnivorous and

opportunistic in their diet, utilizing seeds, arthropods, and green vegetation seasonally (Bradley

and Mauer 1973; Meserve 1976) and they shift to consuming cactus seeds and/or fruit pulp

during summer months (Orr et al. 2015).

We examine the effect of dietary fat content on the metabolic physiology of the desert-

adapted cactus mouse to understand the physiological consequences of food availability and

dietary choices. To accomplish this, we extend previously characterized circadian metabolic

patterns for males and females (Colella et al. 2021b) to estimate rates of metabolism and water

loss in animals fed an experimental diet low in fat, but unchanged in terms of energy

composition and level of macronutrients, to those on a standard laboratory diet. This

experimental framework allows us to test the relationships between dietary fats, energy

expenditure, and water balance, which have direct connections to fitness. Given that fat

catabolism yields more water and energy compared to carbohydrates or proteins, we

hypothesized that mice fed a low-fat diet would perform more poorly than those on the standard

diet.

**Methods**

*Animal Care and Experimental Model*

We worked with a captive colony of cactus mice maintained at the University of New

Hampshire. Captive animal care procedures followed guidelines established by the American

Society of Mammalogists (Sikes et al. 2016), American Veterinary Medical Association (Leary

et al. 2013), and were approved by the University of New Hampshire Institutional Animal Care

and Use Committee (IACUC, #103092).

Sexually mature, non-reproductive healthy adult mice (n = 28 males, n = 28 females)

between three and nine months of age, bred from wild-derived lines at the *Peromyscus* Genetic

Stock Center at the University of South Carolina (Columbia, South Carolina, USA) were used in

this study. Mice were randomly assigned to one of two feeding groups: a standard diet group (SD

- LabDiet® 5015*, 26.10% fat, 19.752% protein, 54.15% carbohydrates, energy 15.02 kJ/g, food

quotient [FQ, the theoretical respiratory quotient produced by the diet based on macronutrient

composition, Westerterp, 1993] 0.89) or a low-fat diet group (LFD - Modified LabDiet® 5015

[5G0Z], 6.6% fat, 22.8% protein, 70.6% carbohydrates, energy 14.31 kJ/g, FQ 0.92). All food

was stored in the desert chamber to match the environmental conditions described below to

control for the water content in the diets. Animals were housed individually in 9.5 L animal

chambers inside a larger, environmentally controlled room built to simulate the temperature,

humidity, and photoperiodic conditions of the Sonoran Desert (Kordonowy et al. 2017; Colella et

al. 2021b). Each animal chamber contained dried cellulose-based bedding, and animals were

provided food and water *ad libitum*. Mice were acclimated to the LFD for one month and to

experimental cages for 24 hours prior to the beginning of metabolic measurements. Animals

were weighed to the nearest tenth of a gram on a digital scale before being housed in an

experimental chamber for four days during metabolic data collection. Animals were weighed

again at the end of the experiment.

Environmental temperatures and relative humidity (RH) followed a natural, diurnal

pattern. Temperature and RH were fixed at 32°C and 10% RH during the light phase for 11

hours (06:00 to 20:00) and dropped to 24°C and increased to 25% RH during a one-hour

transition starting at 20:00. Temperature and RH then remained constant during the dark phase

for seven hours, before the temperature increased again and the RH fell to light phase

temperature during a three-hour transition that started at 06:00. Photoperiod was regular with 16

hours of light and 8 hours of dark. See Figure 1A for a visual representation of environmental

conditions.

*Metabolic phenotyping*

Metabolic phenotyping data were collected starting at 12:00 using a previously described

(Colella et al. 2021b) commercially available Field Metabolic System purchased from Sable

Systems International (SSI). Physiological data were collected following the methods outlined in

Whitfield et al. (2015) and Colella et al. (2021a). Oxygen consumption ($VO_2$ ml min$^{-1}$), carbon

dioxide production ($VCO_2$ ml min$^{-1}$), and water vapor pressure (WVP, kPa) lost via urine, feces,

cutaneous, and respiration were continually measured during 72-hour trials. The experiment was

repeated twice for each sex under each diet (SD and LFD), resulting in three-day experiments

performed sequentially for eight batches of seven mice (and one empty, baseline chamber) per

three-day experiment. Batches alternated between females and males for a total of 28 adult males

and 28 adult females.

Airstreams were pulled from flow-through chambers at a constant rate of 1600 ml min$^{-1}$

(96 L h$^{-1}$) by an SSI SS-4 sub-sampler pump (one for each chamber), multiplexed through an

SSI MUXSCAN, sub-sampled at 250 ml min$^{-1}$ into a Field Metabolic System (FMS) where

WVP, $CO_2$, and $O_2$ were analyzed with no scrubbing. The FMS was zeroed, and sensors were

spanned between each 72-hour experiment following the SSI Instrument Settings and Calibration

manual using dry gas of known $CO_2$ and $O_2$ concentrations.

To generate accurate individual measurements, chambers were measured in a

pseudorandom order for 120s each, approximately three times every hour. Using methods

described in Lighton (2018), we used the most stable 50% of each 120s period to produce a

single averaged value for the rate of $O_2$ consumption (ml min$^{-1}$), rate of $CO_2$ production (ml min$^{-1}$

), and water loss rate (WLR) (mg h$^{-1}$) per measurement window. Lag time between the

analyzers was corrected so that measurements represent the same time periods (Lighton 2018).

This procedure results in a highly repeatable set of measurements. $VO_2$ and $VCO_2$ were

calculated using equations 10.5 and 10.6, respectively, from Lighton (2018). Respiratory

quotient (RQ) was calculated as the ratio of $VCO_2$ to $VO_2$ (Lighton 2018). Energy expenditure

(EE), kJ hr$^{-1}$, was calculated as in Lighton (2018, eq. 9.15). Response variable measurements that

were more than three standard deviations away from the mean, which often represented times

when animal care staff entered the room, were considered outliers, and therefore were removed

from downstream analysis.

*Electrolytes*

At 12:00 Eastern Standard Time (EST), immediately following the conclusion of the experiment,
animals were weighed and euthanized with an overdose of inhaled isoflurane and decapitation.
120 µL of trunk blood was collected within one minute of death for serum electrolyte
measurement using an Abaxis i-STAT® Alinity machine and CHEM8+ cartridges (Abbott Park,
IL, USA, Abbott Point of Care Inc.), and calibrated automatically with each cartridge run. We
used CHEM8+ cartridges to measure the concentration of sodium (Na), potassium (K), creatinine
(Cr), blood urea nitrogen (BUN), hematocrit (HCT), glucose (Glu), and ionized calcium (iCa),
which are expected to vary in response to hydration status and renal function. Finally, using
sodium, glucose, and BUN, we calculated osmolality using the formula in Rasouli (2016). After
assessing for normality, we used a student's two-tailed t-test in R (stats::t.test) to test for
significant ($p < 0.05$) differences in weight and electrolytes within and between the treatments
and sexes for each experimental group.

*Statistical Analysis*

All statistical analyses were conducted in R v 4.0.3 (R Core Team 2020) unless otherwise
specified. To perform analyses on the shape of the temporally variable curves, we modeled the
effect of time in days, diurnal time, and EE, RQ and/or WLR on EE, RQ, and WLR using
Generalized Additive Mixed Models (GAMMs) using the gamm function in the mgcv R package
(Lin and Zhang 1999; Wood 2017). To avoid violating the assumption of independence, we used
the experimental batches and mice within each experimental batch as random effects. The R
function gamm then simultaneously estimates the variances of these random effects as well as
the treatment effects, thus implicitly adjusting treatment and covariate effects for the effect of
correlated, nested groups of measurements. As a result, treatments do not explain differences
between individual mice, but rather the average difference between groups of mice, because we
were interested in the diet treatment and the effect of sex, as well as in the response over the full
72-hour period and the diurnal cycle. Our models included two fixed effects, diet and sex, and
four nonlinear smoothing regression terms: time in days as a nonlinear trend, diurnal cycle as a
circular spline function, and two of the three respirometry response variables (EE, RQ, and/or

WLR). Using the effective degrees of freedom (edf), we were able to determine the degree of

non-linearity of each curve (Wood 2017). To test for physiological differences between mice fed

each diet during the light and dark phases, we subsetted data collected during the light phase and

dark phase for each response variable and ran GAMMs as described above.

To test for differences in the total volume of water lost for each animal over the course of

the experiment, rather than the shape of the curve, we estimated the amount of water lost per

individual for every hour of the experiment and summed those values. These values were then

used as the response variable in an ANCOVA (car::anova, Fox and Weisberg 2019). The model

consisted of two independent variables, sex and diet, and the interaction between them. Weight

at the start of the experiment was used as the covariate. Pairwise comparisons were made using

the Tukey separation of adjusted means test (emmeans:: emmeans and emmeans::contrast, Lenth

2021).

**Results**

Data from 56 sexually mature (but non-reproductive) male and female mice were collected as

described above. No health issues were detected by veterinary staff during the duration of the

experiment, nor were there significant differences in weight between males and females (mean of

22.20 g [range: 15.40 - 29.10 g] versus 21.2 g [range: 16.91 - 29.34 g], respectively), or between

diet groups.

*Metabolic Phenotyping*

In total, we recorded 8,125 120s intervals: 4,091 female, 4,034 male (Supplemental Table 1). We

measured metabolic variables for 4,105 120s intervals under the LFD: 2,053 female and 2,052

male measurements. For the SD treatment, we recorded 4,020 total observations (2,038 female,

1,982 male). After removing outliers, we retained 3,945 female measurements (1,977 SD and

1,968 LFD) and 3,963 male measurements (1,945 SD, 2,018 LFD). *Peromyscus eremicus* show

physiological patterns that are in phase with photoperiod and room temperature for both diets

(Figure 1), as described in (Fox and Weisberg 2019; Colella et al. 2021b). Raw data and

processed machine-readable csv files are available on Zenodo:

https://zenodo.org/record/6422231#.YlXSD9PML0o.

**Figure 1**

72 hours of environmental (A) and respirometry (B-D) data split by sex for 28 adult males and
28 adult females plotted over a 24-hour window to display circadian patterns for each group.
Organismal responses to standard diet are in gray and low-fat diet in blue. Vertical blocks shaded
in gray indicate the dark phase when animals are active, and unshaded blocks indicate light phase
when animals are inactive. **A)** Room temperature (degrees Celsius) is indicated by a solid red
line. **B)** 24-hour measurements of energy expenditure (EE kJ hr$^{-1}$), **C)** water loss rate (WLR, $H_2O$
g hr$^{-1}$), and **D)** respiratory quotient (RQ), for females (F, left) and males (M, right). Statistical

significance between the fixed effects, diet and sex, for the 24hr GAMMs are denoted with one

or more · under the y-axis label, 'ns' indicates non-significance. Statistical significance between

the fixed effects, diet and sex, from GAMMs on the subsetted day/night data are denoted with

one or more **\*** on the top axis of the plots. The two symbols (\*/·) signify the same significance

values (\* p <= 0.05, \*\*: p <= 0.01, \*\*\*: p <= 0.001, \*\*\*\*: p <= 0.0001).

*Energy Expenditure*

Consistent with previous work (Colella et al. 2021b), males and females show diurnal patterning

of EE on both diets, with the highest EE occurring during the dark phase when environmental

temperature is lowest and animals are active. The lowest EE occurred during the light phase,

when environmental temperature was greatest and animals were inactive (Figure 1B). Using a

GAMM, we modeled EE with diet and sex as fixed effects, as well as other predictors, such as

day, time, WLR, and RQ, and found that neither sex nor diet influenced EE (diet p = 0.56, sex p

= 0.18, Supplemental Table 2). When analyzing light and dark phases separately, there were no

differences in EE between males and females nor between the two diets (light phase: diet p =

0.550, sex p= 0.293; dark phase: diet p = 0.571, sex p = 0.182, Supplemental Table 3 and 4).

There was a weak nonlinear relationship between EE and light phase (Supplemental Figure 2A,

Supplemental Table 2, edf = 1.98, $p < 2e^{-16}$). Additionally, the relationship between time and EE

was statistically significant but nonlinear (Supplemental Figure 2F, Supplemental Table 2, edf =

7.98, $p < 2e^{-16}$).

*Water Loss Rate*

Both treatment groups, SD and LFD, and both sexes exhibited diurnal patterning of WLR and

had the highest WLR during the light phase when environmental temperature was highest, even

though animals were inactive then (Figure 1C, Supplemental Figure 1F, Supplemental Table 2,

edf = 7.95, $p < 2e^{-16}$). Peak loss occurred between 09:00 and 11:00. By fitting a 24-hour time-

continuous GAMM (Supplemental Figure 1E-H), we showed that both diet and sex had a

significant effect on the WLR (diet p = $2.13E^{-05}$, sex p = 0.002, Supplemental Table 2), with both

males and females fed a LFD losing more water than those fed the SD. When modeling water

loss during the light phase only, we showed that both diet and sex had a significant effect on

WLR (diet p = $4.80e^{-15}$, sex p = $1.69e^{-11}$, Supplemental Table 3). When modeling water loss only

during the dark phase, when animals are active, WLR was significantly different between males

and females (p = 0.015, supplemental table 4), but WLR was not different within sex between

different diets (p = 0.797, Supplemental Table 4). The relationship between WLR and

experimental day was weakly non-linear, but significant (Supplemental Figure 1E, Supplemental

Table 2, edf = 1.99, p < $2e^{-16}$).

Both diet and sex significantly affected the amount of water lost over the course of the

experiment when controlling for individual body weight (p = $5.22e^{-2}$ and p = 0.017, respectively,

Supplemental Table 5), but the interaction term was not significant (p= 0.633, Supplemental

Table 5). Further investigation into the sex:diet interaction showed significant differences in both

the female  (p=0.043, Supplemental Table 5) and male (p=0.005, Supplemental Table 5) diet

comparison.

*Respiratory Quotient*

RQ showed circadian rhythms for both diets and sexes with the highest RQ occurring during the

light phase and the lowest during the dark phase (Colella et al. 2021b; Figure 1D). By fitting a

24-hour GAMM we found no significant difference in RQ between the sexes or diets (sex p =

0.83, diet p = 0.52, Supplemental Table 2) but we did find that day and time were highly

significant (p = $1.98e^{-07}$, p < $2.2e^{-16}$ respectively). Time of day had a complex relationship with

RQ (Supplemental Figure 1, edf = 7.69, Supplemental Table 2) and experimental day had a weak

non-linear relationship with RQ (Supplemental Figure 1D, edf = 1.59, Supplemental Table 2).

During the light phase, RQ rose above the FQ, with females fed the SD having the

highest RQ at the start of the light phase and then stabilizing at an RQ > 1 around 14:00. RQ did

not differ by sex nor diet (sex p = 0.740, diet p = 0.543, Supplemental Table 3). When modeling

RQ during the dark phase, values were comparable to the FQ of the respective diet for both sexes

(FQ LFD: 0.92, FQ SD: 0.89) and RQ was significantly different between the diets (p = 0.017,

Supplemental Table 4) but not between males and females (p = 0.739, Supplemental Table 4).

*Electrolytes*

Several serum electrolyte values were significantly different between diets for males and females (sodium p = 0.22 and p = 0.01 respectively; potassium p = 0.02 and p = 0.01; BUN p = 0.03 and 0.71; hematocrit p = 0.04 and 0.02; ionized calcium p = 0.001 and 0.001; osmolality p = 0.01 and 0.05) (Figure 2, Table1). When comparing males and females on the same diet, sodium (p = 0.02) and Osmolality (p = 0.03) differed significantly for mice on the SD and no electrolyte measurements were differed between males and females on the LFD.

**Figure 2**

Violin plots showing the distribution of serum electrolyte measurements: Na = sodium

(mmol/L), K = potassium (mmol/L), BUN = blood urea nitrogen (mmol/L), Hct = hematocrit (%

PCV), iCa = ionized calcium (mmol/L), Glu = glucose (mmol/L), and osmolality (mmol/L) for

female (F, left) and male (M, right) *Peromyscus eremicus* fed standard (grey) and low-fat (blue)

diets. Observations (28 adult females and 28 adult males, n=14 of each treatment, total n=56) are

represented by black dots. P-values from pairwise t-tests are reported above the brackets.

**Discussion**

We explored the effect of dietary fat intake on water balance in a desert adapted rodent and

found that mice fed a diet lower in fat lost more water than mice fed a diet higher in fat. In

environments where water scarcity is extreme, even small differences in water balance could

have substantial impacts on cognitive function and physical performance, and therefore survival

and reproductive success (Boogert et al. 2018). Our results help clarify the physiological

consequences of food availability and dietary choices.

*Energy Expenditure*

Not all diets provide animals with the same amount of energy per unit mass (Withers 1982).

Different macronutrients have different energy potentials (Sánchez-Peña et al. 2017). Diet

composition influences EE (Acheson 1990; Westerterp 2004; Secor 2009; Beale et al. 2018), as

do other intrinsic (*i.e.,* body mass, temperature, evolutionary history, reproductive stage, activity,

or energy intake) and extrinsic (*i.e.,* habitat [including environmental temperature], climate, or

social environment; Speakman 1997) variables. In wild populations, dietary switches tend to

coincide with seasonal changes that affect resource quantity and/or quality (Noble et al. 2019).

Previous studies have recorded cactus mice shifting diet seasonally, consuming arthropods

during the winter (Hope and Parmenter 2007), and transitioning to the consumption of cactus

seeds and/or fruits during the summer (Hope and Parmenter 2007; Orr et al. 2015). Many plants

native to the Sonoran Desert have a fat content comparable to that of the LFD used in this study

(5.49 - 9.88% for leaf tissues and branches, McArthur 1994; 0.90- 3.60% for roots and seeds,

Castle et al. 2020).

In the current study, both diets were designed to have equivalent amounts of energy per

gram of weight. We designed the experiment to control for several intrinsic and extrinsic factors

that represent sources of variation in EE, but not in individual activity. We show that

experimental manipulation of diet does not result in differences in EE when food is provided *ad*

*libitum*. EE and activity are tightly coupled (Garland et al. 2011; Kaiyala et al. 2012). Therefore

we expect, at most, limited differences in activity given limited differences in EE. For both diets,

animals are likely to eat enough to satisfy energy needs, a hypothesis further supported by the

fact that there was no net change in weight over the course of the experiment. Consistent with

other studies, we recorded higher EE during the dark phase of the 24-hour cycle when animals

were active (Colella et al. 2021b), further supporting that metabolic rate is complex and may be

modulated by the environment, activity, or other extrinsic/intrinsic factors.

*Water Loss Rate*

**<u>Thermoregulation and water loss</u>**: Evaporative water loss is one way for organisms to cool

themselves (*e.g.,* via latent heat of vaporization) and results in a decrease in body temperature

and loss of body mass (Porter and Gates 1969). While body temperature was not measured as a

part of this study, numerous studies have shown that desert animals reduce respiratory water loss

as an adaptive physiological mechanism for maintaining water balance in desert environments

(Schmidt-Nielsen and Schmidt-Nielsen 1951; Hart 1971; MacMillen 1972). Indeed, a fine

balance must exist between using water for thermoregulation, the use of water for other critical

metabolic processes, and dehydration. Exaggerated use of water for thermoregulation may be

harmful when water is limited. Instead, other less water-intensive mechanisms of heat loss such

as conduction, and behavioral changes such as estivation and nocturnality may be used to

maintain a homeostatic body temperature but are not quantified in this study.

Despite the consequences of limited water supply, desert *Peromyscus* have significant

capacity for evaporative cooling for thermoregulation (Ramirez et al. 2022). In our study, we

found that animals fed the LFD had greater water loss compared to animals fed the SD (Figure

1C). While that loss may be related to differences in thermoregulatory performance, it may

further compromise water balance. Are these patterns suggestive of differences in physiological

performance? Is the consumption of a diet lower in fat related to a reduction in physiological

performance and, consequently, fitness? While the current experiment cannot definitively answer

these questions, there is some evidence (*e.g.,* electrolytes, water loss) to suggest that

physiological performance is impaired in animals consuming the LFD and that this performance,

particularly when water is scarce, may have fitness consequences.

**Potential mechanisms for increased water loss on LFD**: Across all comparisons, there is a

strong and statistically significant relationship between the water loss rate and dietary fat content,

with individuals fed the LFD losing more water. While the potential mechanisms underlying

these findings will require future experiments, several can be ruled out. First, the maintenance of

water balance is a process regulated by a cascade of hormones including arginine vasopressin

and the renin–angiotensin–aldosterone system pathway (Aisenbrey et al. 1981; Greenleaf 1992).

While some hormones are lipid-based (*e.g.,* aldosterone) and their production and secretion are

linked to dietary composition (*i.e*., sex hormones and severe low body fat), the degree to which

dietary fats are reduced for mice fed the LFD are not to the level at which hormonal imbalances

should be seen. Second, while there is a known relationship between ketogenic diets and

dehydration (Wheless 2001; Freeman et al. 2006), these diets are associated with enhanced

diuresis. We observed the opposite with animals fed the SD; they lost less water than animals fed

the LFD. Third, diets higher in protein require more water for the removal of nitrogenous wastes

(Calder and Braun 1983), but higher protein does not account for the observations herein. To

approximate equivalent energy availability there is a small difference in the amount of dietary

protein in each dietary treatment (18.9% in SD versus 19.5% in LFD), however, that small

difference is unlikely to result in the observed differences in water loss and hydration status.

While food and water intake were not measured directly, a potential mechanism for the

observed differences could be that animals fed the LFD simply eat more than animals fed a diet

higher in fat. While the diets are closely matched in terms of available energy per unit mass, the

LFD is slightly less energy dense. Additionally, it's possible that there are substantial differences

in the non-nutritive fraction of the diet that could be further explored. While the current study

does not disentangle this potential aspect, future work could. These differences suggest that

animals fed the LFD may be ingesting more food mass to maintain nutritional homeostasis and

with food intake increased there may also be increased water intake (Bachmanov et al. 2002).

While we have no evidence for this, an increased rate of consumption would potentially result in

an increased rate of urine production and the production of water-containing fecal material and

urine. Given that animals fed the LFD seem to be more dehydrated than animals fed the SD

(increased Na, osmolality, Hct, Figure 2), this hypothesis requires that animals fed the LFD

consume more water, but not enough to satisfy their physiological needs. Given that the animals

have free access to water, an unmet need is irreconcilable without further explanation.

Another intriguing hypothesis linked to the concept of unmet hydrational needs has to do

with thirst pathways. Thirst is a complex phenotype under tight neural control. Osmo- and

baroreceptors located in the vasculature are responsible for monitoring water balance and, when

triggered, set into motion a hormonal cascade that simultaneously enhance renal water retention

and stimulate thirst (Thornton 2010; Leib et al. 2016). While the former aspects of the response

are crucial to many physiological processes, the latter response, thirst, could be responsible for

the observations described above. While purely a hypothesis, there are several examples of

species losing aspects of their sensory systems via natural selection, loss of sight (Gore et al.

2018), decreases in certain taste sensitivities or loss of taste entirely (Jiang et al. 2012), and loss

of specific olfactory functions (Kishida et al. 2015). For animals in dry deserts, where water is

scarce, could well-developed pathways leading to thirst be similarly changed? While requiring

careful consideration, perhaps using a comparative genomics approach, this hypothesis could

explain why animals with free access to water remain significantly dehydrated.

*Respiratory Quotient*

RQ, the ratio of $CO_2$ produced to $O_2$ consumed, can inform rates of differential fuel utilization.

RQ values typically fall between 0.7 and 1.0, with the catabolism of fats yielding 0.7,

carbohydrates 1.0, and proteins intermediate values (Kleiber 1975). RQ values can exceed 1.0

when anabolism of fat exceeds catabolism (*i.e.,* de novo lipogenesis (DNL) Benedict 1937;

Abreu-Vieira et al. 2015; Levin et al. 2017) or during anaerobic exercise (Whipp 2007; Zagatto

et al. 2012). While carbohydrates, fats, or proteins are all burned as fuel, if present in excess,

they can also be converted to fat which can be stored for future use (Barboza et al. 2009). DNL is

one mechanism for converting non-fat energy into a form that can be stored. The conversion of

glucose to fats prior to entry into the citric acid cycle is exothermic and results in an energetic

cost, representing a metabolically inefficient use of dietary substrate in cold habitats (Solinas et

al. 2015), but a potentially efficient mechanism of heat regulation in hot deserts. RQ was greater

than 1.0 during the inactive period in animals on both diets suggesting that *de novo* lipogenesis

(DNL) is not limited by fat intake in this species. Here and in previous experiments on *P.*

*eremicus* (Colella et al. 2021b), RQ is only greater than 1.0 during the light phase when animals

are inactive, which strongly suggests DNL is occurring. Further, fat storage could be a way for

desert mammals to buffer against uncertain water availability, as fat oxidation releases metabolic

water (Mellanby 1942; Davidson and Passmore 1963; Schmidt-Nielsen and Adolph 1964). Given

this, the conversion of other fuel sources to fats may serve as an important reservoir of metabolic

water. Previous studies have shown that DNL can occur under high-carbohydrate and high-fat

diets (Strable and Ntambi 2010). This is supported by our results, as both groups had an RQ

greater than 1.0 during the light phase and RQ values at or below the FQ during the dark phase,

which indicates a shift to the supplied dietary substrate as the main oxidative fuel during cooler,

dark periods.

*Electrolytes*

Electrolytes provide information on the overall metabolic state, renal function, and other core

physiological functions of an organism (Kutscher 1968), including hydration status (Cheuvront

et al. 2010) or disease state. Therefore, to further understand the effects of diet on physiological

function of small mammals in desert environments, we examined serum electrolytes from male

and female cactus mice fed diets with different levels of fat. We established diet-specific

differences for serum Na, K, Cr, BUN, Hct, iCa, and osmolality values for *P. eremicus*. Notably,

synthetic markers of pathological renal impairment, BUN and Cr, were not significantly different

between treatments. Previous studies have shown that the electrolyte-water balance is frequently

unaffected when desert rodents are water-deprived for long periods of time (Heimeier et al.

2004; Heimeier and Donald 2006; Boumansour et al. 2021), however, other desert rodent studies

have demonstrated evidence of dehydration in the absence of exogenous water without renal

impairment (Kordonowy et al. 2017; Boumansour et al. 2021). So, even with the significant

water lost through respiration, when free water is available, dietary fat content does not

significantly impair the kidney for *P. eremicus*.

Interestingly, other key electrolytes that are sensitive to hydration status were

significantly different between the dietary treatment groups (Figure 2). Serum Na, osmolality,

and Hct are expected to be elevated in clinical dehydration and were found to be elevated in

animals fed the LFD. Though some contrasts were not statistically significant, measurements

were consistently higher for both males and females, a strong sign of differences in hydration

between treatment groups. These differences are particularly interesting because while animals

fed a LFD had higher rates of water loss than those fed the SD, all animals had access to water

*ad libitum*. Elevated serum Na, osmolality, and Hct in the context of normal renal function are

evidence of dehydration, but dehydration in neuro-intact animals with free access to water is

unusual. While dietary fats may be responsible for changes in solute balance (Friedman et al.

2012), typically the relationship is in the opposite direction to what we found (Abdel-Rahman

2010). One possible hypothesis is that there may be an impairment of the physiological

mechanisms related to thirst. Thirst is a response to changes in the blood chemistry (*i.e.,* increase

in osmolality Gilman 1937; Wolf 1950; Heimeier et al. 2004) or volume (Fitzsimons 1961;

Stricker 1966) both of which are monitored by the lamina terminalis circumventricular organs

(Bourque 2008; Zimmerman et al. 2017). Activation of these brain regions motivates an

organism to find water. Previous studies have found that the expression of two genes, *Rxfp1* and

*Pdyn,* in relevant cell types correlates with neural activation under osmotic and hypovolemic

thirst in lab mice (Pool et al. 2020). Further studies should examine these genes and others in

desert organisms to understand the basis of thirst pathways. Regardless of the process, elevated

serum sodium may have important and complex physiological consequences. For example,

elevated blood pressure and increased fluid retention as a result of elevated sodium levels, both

have obvious implications for survival in desert environments and deserve future study.

**Conclusion**

To understand ways in which the desert-adapted cactus mouse, *P. eremicus,* responds to

variation in dietary fat content, reproductively mature mice were subjected to one of two diets, a

SD or a LFD. Overall, EE and WLR for mice on the SD during the inactive, light phase (a

measure comparable to the resting metabolic rate) agreed with earlier studies of *Peromyscus*

(Withers 1992; Ramirez et al. 2022). Further, we show that dietary fat is important for water

regulation in cactus mice through the analysis of interacting physiological traits: 1) high rates of
total water loss during the warmer, drier light phase and 2) a difference in serum electrolyte
values indicating dehydration. Our results suggest that a LFD could limit the capacity of desert
animals to tolerate limited access to free water as is common in arid environments.

The identification of short-term, physiological responses of *P. eremicus* associated with
dietary fat composition demonstrates that adaptations that evolved over long evolutionary
timescales will affect survival during the accelerated time scales of climate change. In light of
global climate change and increased desertification, investigating the range and mechanisms of
plastic responses employed by desert-adapted species could provide insights into physiological
responses to increasingly erratic climate.

**Acknowledgments**

We would like to thank members of the MacManes Lab for helpful comments and support on
early versions of the manuscript; the Animal Resources Office and veterinary care staff at the
University of New Hampshire; A. Gerson at the University of Massachusetts Amherst, Z.
Cheviron at the University of Montana, B. Joos and J. Klok at Sable Systems International for
analytic guidance, technical support, and respirometry training. This work was supported by the
National Institute of Health National Institute of General Medical Sciences (R35 GM128843 to
M.D.M.).

**Author Contributions**

Conceptualization: M.D.M.; Methodology: D.M.B., M.D.M.; Formal analysis: D.M.B., E.L.;
Investigation: D.M.B., J.P.C.; Resources: M.D.M.; Writing - original draft: D.M.B.; Writing -
review & editing: D.M.B., J.P.C., E.L., M.D.M.; Visualization: D.M.B., E.L.; Supervision:
M.D.M.; Project administration: M.D.M.; Funding acquisition: M.D.M.

**Competing Interests**

No competing interests declared.

**Data Availability**

Raw ExpeData (SSI) files are available through Zenodo:

https://zenodo.org/record/6422231#.YlXSD9PML0o. Macro processing files, raw and processed

respirometry data, and cage sampling scheme files are also available on Zenodo. All R scripts

used in this project are available through GitHub at:

https://github.com/DaniBlumstein/Diet_paper.

**References**

ABDEL-RAHMAN, M. K. 2010. Influence of dietary fat on renal function, lipid profile, sex hormones, and electrolyte balance in rats. European Journal of Lipid Science and Technology 112:1166–1172.

ABREU-VIEIRA, G., C. XIAO, O. GAVRILOVA, AND M. L. REITMAN. 2015. Integration of body temperature into the analysis of energy expenditure in the mouse. Molecular Metabolism 4:461–470.

ACHESON, K. J. 1990. Sympathetic Nervous System in the Regulation of Thermogenesis. Pp. 40–46 in Hormones and Nutrition in Obesity and Cachexia (M. J. Müller, E. Danforth, A. G. Burger & U. Siedentopp, eds.). Springer, Berlin, Heidelberg.

AISENBREY, G. A., W. A. HANDELMAN, P. ARNOLD, M. MANNING, AND R. W. SCHRIER. 1981. Vascular effects of arginine vasopressin during fluid deprivation in the rat. The Journal of Clinical Investigation 67:961–968.

BACHMANOV, A. A., D. R. REED, G. K. BEAUCHAMP, AND M. G. TORDOFF. 2002. Food Intake, Water Intake, and Drinking Spout Side Preference of 28 Mouse Strains. Behavior genetics 32:435–443.

BARBOZA, P. S., K. L. PARKER, AND I. D. HUME (EDS.). 2009. Lipids: Fatty Acids and Adipose Tissue. Pp. 119–131 in Integrative Wildlife Nutrition. Springer, Berlin, Heidelberg.

BEALE, P. K., K. J. MARSH, W. J. FOLEY, AND B. D. MOORE. 2018. A hot lunch for herbivores: physiological effects of elevated temperatures on mammalian feeding ecology. Biological Reviews 93:674–692.

BEDFORD, N. L., AND H. E. HOEKSTRA. 2015. Peromyscus mice as a model for studying natural variation. eLife 4:e06813.

BENEDICT, F. G. 1937. Lipogenesis in the animal body, with special reference to the physiology of the goose. Carnegie institution of Washington.

BOOGERT, N. J., J. R. MADDEN, J. MORAND-FERRON, AND A. THORNTON. 2018. Measuring and understanding individual differences in cognition. Philosophical Transactions of the Royal Society B: Biological Sciences 373:20170280.

BOUMANSOUR, L. ET AL. 2021. Vasopressin and oxytocin expression in hypothalamic supraoptic
nucleus and plasma electrolytes changes in water-deprived male Meriones libycus.
Animal Cells and Systems 0:1–10.

BOURQUE, C. W. 2008. Central mechanisms of osmosensation and systemic osmoregulation.
Nature Reviews Neuroscience 9:519–531.

BRADLEY, W. G., AND R. A. MAUER. 1973. Rodents of a Creosote Bush Community in Southern
Nevada. The Southwestern Naturalist 17:333–344.

CALDER, W. A., AND E. J. BRAUN. 1983. Scaling of osmotic regulation in mammals and birds. The
American Journal of Physiology 244:R601-606.

CASTLE, S. T. ET AL. 2020. Diet composition analysis provides new management insights for a
highly specialized endangered small mammal. PLOS ONE 15:e0240136.

CHEUVRONT, S. N., R. W. KENEFICK, S. J. MONTAIN, AND M. N. SAWKA. 2010. Mechanisms of
aerobic performance impairment with heat stress and dehydration. Journal of Applied
Physiology 109:1989–1995.

COLELLA, J. P. ET AL. 2021a. Limited Evidence for Parallel Evolution Among Desert-Adapted
Peromyscus Deer Mice. Journal of Heredity 112:286–302.

COLELLA, J. P., D. M. BLUMSTEIN, AND M. D. MACMANES. 2021b. Disentangling environmental
drivers of circadian metabolism in desert-adapted mice. Journal of Experimental Biology
224.

CROSSLAND, J. P., M. J. DEWEY, S. C. BARLOW, P. B. VRANA, M. R. FELDER, AND G. J. SZALAI.
2014. Caring for Peromyscus spp. in research environments. Lab Animal 43:162–166.

DAVIDSON, S., AND R. PASSMORE. 1963. Human nutrition and dietetics. Human nutrition and
dietetics.

FITZSIMONS, J. T. 1961. Drinking by rats depleted of body fluid without increase in osmotic
pressure. The Journal of Physiology 159:297.

FOX, J., AND S. WEISBERG. 2019. An {R} Companion to Applied Regression. Third. Sage.

FRANK, C. L. 1988. Diet Selection by a Heteromyid Rodent: Role of Net Metabolic Water
Production. Ecology 69:1943–1951.

FREEMAN, J., P. VEGGIOTTI, G. LANZI, A. TAGLIABUE, AND E. PERUCCA. 2006. The ketogenic diet:
from molecular mechanisms to clinical effects. Epilepsy Res 68:145–80.

FRIEDMAN, A. N. ET AL. 2012. Comparative effects of low-carbohydrate high-protein versus low-
fat diets on the kidney. Clinical journal of the American Society of Nephrology: CJASN
7:1103–1111.

GARLAND, T., JR ET AL. 2011. The biological control of voluntary exercise, spontaneous physical
activity and daily energy expenditure in relation to obesity: human and rodent
perspectives. Journal of Experimental Biology 214:206–229.

GILMAN, A. 1937. The relation between blood osmotic pressure, fluid distribution and voluntary
water intake. American Journal of Physiology-Legacy Content 120:323–328.

GREENLEAF, J. E. 1992. Problem: thirst, drinking behavior, and involuntary dehydration.
Medicine & Science in Sports & Exercise 24:645.

HART, J. S. 1971. Rodents. Pp. 1–149 in Mammals. Elsevier.

HEIMEIER, R. A., R. C. BARTOLO, AND J. A. DONALD. 2004. The effect of water deprivation on
signalling molecules that utilise cGMP in the spinifex hopping mouse Notomys alexis.
Australian Mammalogy 26:191–198.

HEIMEIER, R. A., AND J. A. DONALD. 2006. The response of the natriuretic peptide system to
water deprivation in the desert rodent, Notomys alexis. Comparative Biochemistry and
Physiology Part A: Molecular & Integrative Physiology 143:193–201.

HOPE, A., AND R. PARMENTER. 2007. Food Habits of Rodents Inhabiting Arid and Semi-arid
Ecosystems of Central New Mexico. Special Publications.

JIANG, P. ET AL. 2012. Major taste loss in carnivorous mammals. Proceedings of the National
Academy of Sciences 109:4956–4961.

KAIYALA, K. J. ET AL. 2012. Acutely Decreased Thermoregulatory Energy Expenditure or
Decreased Activity Energy Expenditure Both Acutely Reduce Food Intake in Mice. PLOS
ONE 7:e41473.

KISHIDA, T., J. THEWISSEN, T. HAYAKAWA, H. IMAI, AND K. AGATA. 2015. Aquatic adaptation and
the evolution of smell and taste in whales. Zoological Letters 1:9.

KLEIBER, M. 1975. The fire of life. An introduction to animal energetics.

KORDONOWY, L. ET AL. 2017. Physiological and biochemical changes associated with acute
experimental dehydration in the desert adapted mouse, *Peromyscus eremicus*.
Physiological Reports 5:e13218.

KUTSCHER, C. 1968. Plasma volume change during water-deprivation in gerbils, hamsters,
guinea pigs and rats. Comparative Biochemistry and Physiology 25:929–936.

LEARY, S. L. ET AL. 2013. AVMA guidelines for the euthanasia of animals: 2013 edition.
American Veterinary Medical Association Schaumburg, IL.

LEIB, D. E., C. A. ZIMMERMAN, AND Z. A. KNIGHT. 2016. Thirst. Current Biology 26:R1260–R1265.

LEVIN, E., G. LOPEZ-MARTINEZ, B. FANE, AND G. DAVIDOWITZ. 2017. Hawkmoths use nectar sugar
to reduce oxidative damage from flight. Science 355:733–735.

LIGHTON, J. R. B. 2018. Measuring Metabolic Rates: A Manual for Scientists. 2nd edition. Oxford
University Press.

LIN, X., AND D. ZHANG. 1999. Inference in Generalized Additive Mixed Models by Using
Smoothing Splines. Journal of the Royal Statistical Society. Series B (Statistical
Methodology) 61:381–400.

MACMANES, M. D. 2017. Severe acute dehydration in a desert rodent elicits a transcriptional
response that effectively prevents kidney injury. Renal Physiol:11.

MACMILLEN, R. E. 1965. Aestivation in the cactus mouse, Peromyscus eremicus. Comparative
Biochemistry and Physiology 16:227–248.

MACMILLEN, R. E. 1972. Water economy of nocturnal desert rodents.

MCARTHUR, E. D. 1994. Nutritive Quality and Mineral Content of Potential Desert Tortoise Food
Plants. Intermountain Research Station.

MELLANBY, K. 1942. Metabolic Water and Desiccation. Nature 150:21–21.

MESERVE, P. L. 1976. Habitat and Resource Utilization by Rodents of a California Coastal Sage
Scrub Community. Journal of Animal Ecology 45:647–666.

MURIE, M. 1961. Metabolic Characteristics of Mountain, Desert and Coastal Populations of
Perormyscus. Ecology 42:723–740.

NOBLE, J. D., S. L. COLLINS, A. J. HALLMARK, K. MALDONADO, B. O. WOLF, AND S. D. NEWSOME.
2019. Foraging strategies of individual silky pocket mice over a boom–bust cycle in a
stochastic dryland ecosystem. Oecologia 190:569–578.

ORR, T. J., S. D. NEWSOME, AND B. O. WOLF. 2015. Cacti supply limited nutrients to a desert
rodent community. Oecologia 178:1045–1062.

PARDI, M. I., R. C. TERRY, E. A. RICKART, AND R. J. ROWE. 2020. Testing climate tracking of
montane rodent distributions over the past century within the Great Basin ecoregion.
Global Ecology and Conservation 24:e01238.

PERGAMS, O. R. W., AND J. J. LAWLER. 2009. Recent and Widespread Rapid Morphological
Change in Rodents. PLOS ONE 4:e6452.

POOL, A.-H. ET AL. 2020. The cellular basis of distinct thirst modalities. Nature 588:112–117.

PORTER, W. P., AND D. M. GATES. 1969. Thermodynamic equilibria of animals with environment.
Ecological monographs 39:227–244.

PYKE, G. H. 1984. Optimal Foraging Theory: A Critical Review. Annual Review of Ecology and
Systematics 15:523–575.

PYKE, G. H., H. R. PULLIAM, AND E. L. CHARNOV. 1977. Optimal Foraging: A Selective Review of
Theory and Tests. The Quarterly Review of Biology 52:137–154.

R CORE TEAM. 2020. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

RAMIREZ, R. W., E. A. RIDDELL, S. R. BEISSINGER, AND B. O. WOLF. 2022. Keeping your cool:
thermoregulatory performance and plasticity in desert cricetid rodents. Journal of
Experimental Biology 225:jeb243131.

RASOULI, M. 2016. Basic concepts and practical equations on osmolality: Biochemical approach.
Clinical Biochemistry 49:936–941.

REICHMAN, O. J. 1975. Relation of Desert Rodent Diets to Available Resources. Journal of
Mammalogy 56:731–751.

RUSSELL V. LENTH. 2021. emmeans: Estimated Marginal   Means, aka Least-Squares Means. R
package version 1.6.0.   https://CRAN.R-project.org/package=emmeans.

SÁNCHEZ-PEÑA, M. J., F. MÁRQUEZ-SANDOVAL, A. C. RAMÍREZ-ANGUIANO, S. F. VELASCO-
RAMÍREZ, G. MACEDO-OJEDA, AND L. J. GONZÁLEZ-ORTIZ. 2017. Calculating the
metabolizable energy of macronutrients: a critical review of Atwater's results. Nutrition
Reviews 75:37–48.

SCHMIDT-NIELSEN, B., AND K. SCHMIDT-NIELSEN. 1951. A complete account of the water
metabolism in kangaroo rats and an experimental verification. Journal of Cellular and
Comparative Physiology 38:165–181.

SCHMIDT-NIELSEN, K. 1975. Desert Rodents: Physiological Problems of Desert Life. Pp. 379–
388 in Rodents in Desert Environments (I. Prakash & P. K. Ghosh, eds.). Springer
Netherlands, Dordrecht.

SCHMIDT-NIELSEN, K., AND E. F. ADOLPH. 1964. Desert Animals: Physiological Problems of Heat
and Water. Physiological Zoology 37:338–339.

SECOR, S. M. 2009. Specific dynamic action: a review of the postprandial metabolic response.
Journal of Comparative Physiology B 179:1–56.

SIKES, R. S., AND THE ANIMAL CARE AND USE COMMITTEE OF THE AMERICAN SOCIETY OF
MAMMALOGISTS. 2016. 2016 Guidelines of the American Society of Mammalogists for the
use of wild mammals in research and education. Journal of Mammalogy 97:663–688.

SOLINAS, G., J. BORÉN, AND A. G. DULLOO. 2015. De novo lipogenesis in metabolic homeostasis:
More friend than foe? Molecular Metabolism 4:367–377.

SPEAKMAN, J. 1997. Factors influencing the daily energy expenditure of small mammals.
Proceedings of the Nutrition Society 56:1119–1136.

STRABLE, M. S., AND J. M. NTAMBI. 2010. Genetic control of de novo lipogenesis: role in diet-
induced obesity. Critical reviews in biochemistry and molecular biology 45:199–214.

STRICKER, E. M. 1966. Extracellular fluid volume and thirst. American Journal of Physiology-
Legacy Content 211:232–238.

THORNTON, S. N. 2010. Thirst and hydration: Physiology and consequences of dysfunction.
Physiology & Behavior 100:15–21.

TIGANO, A., J. P. COLELLA, AND M. D. MACMANES. 2020. Comparative and population genomics
approaches reveal the basis of adaptation to deserts in a small rodent. Molecular
Ecology 29:1300–1314.

VEAL, R., AND W. CAIRE. 1979. Peromyscus eremicus. Mammalian Species:1–6.

WESTERTERP, K. R. 2004. Diet induced thermogenesis. Nutrition & metabolism 1:1–5.

WHELESS, J. W. 2001. The ketogenic diet: an effective medical therapy with side effects. Journal
of Child Neurology 16:633–635.

WHIPP, B. J. 2007. Physiological mechanisms dissociating pulmonary $CO_2$ and $O_2$ exchange
dynamics during exercise in humans. Experimental Physiology 92:347–355.

WHITFIELD, M. C., B. SMIT, A. E. MCKECHNIE, AND B. O. WOLF. 2015. Avian thermoregulation in
the heat: scaling of heat tolerance and evaporative cooling capacity in three southern
African arid-zone passerines. Journal of Experimental Biology 218:1705–1714.

WITHERS, P. C. 1982. Effect of diet and assimilation efficiency on water balance for two desert
rodents. Journal of Arid Environments 5:375–384.

WITHERS, P. C. 1992. Comparative animal physiology. Saunders College Pub. Philadelphia.

WOLF, A. V. 1950. Osmometric analysis of thirst in man and dog. American Journal of
Physiology-Legacy Content 161:75–86.

WOLF, B. O., AND C. M. DEL RIO. 2003. How important are columnar cacti as sources of water
and nutrients for desert consumers? A review. Isotopes in Environmental and Health
Studies 39:53–67.

WOOD, S. 2017. Generalized Additive Models: An Introduction with R, Second Edition.

ZAGATTO, A. M., W. E. MIYAGI, R. L. SAKUGAWA, E. I. KAMINAGAKURA, AND M. PAPOTI. 2012.
Aerobic Endurance Measurement by Respiratory Exchange Ratio during a Cycle
Ergometer Graded Exercise Test. Journal of Exercise Physiology Online 15:49–56.

ZIMMERMAN, C. A., D. E. LEIB, AND Z. A. KNIGHT. 2017. Neural circuits underlying thirst and fluid
homeostasis. Nature Reviews Neuroscience 18:459–469.

**Supplemental**

Table S1: Number of samples collected per time interval.

| Photoperiod | Diet | Number of 120s intervals male | Number 120s intervals female |
|---|---|---|---|
| light | standard | 867 | 879 |
| | low-fat | 914 | 927 |
| dark | standard | 758 | 755 |
| | low-fat | 758 | 719 |

703 Table S2: Generalized additive mixed models (GAMM) 24-hour results. Statistical models and

704 results for **A)** water loss rate (WLR, $H_2O$ g/hr$^{-1}$), **B)** energy expenditure (EE kJ/hr$^{-1}$), **C)**

705 respiratory quotient (RQ).

A)

H2Og~diet+Sex+s(ctime.exp,k=3)+s(time_in_D,bs=cc)+s(EE_kJH)+s(RQ)

| | Estimate | Std. | Error | t value | Pr(>|t|) | |
|---|---|---|---|---|---|---|
| (Intercept) | 0.121 | 0.008 | 15.582 | <2.00E-16 | *** | |
| diet | -0.038 | 0.009 | -4.253 | 0.000 | *** | |
| Sex | 0.027 | 0.009 | 3.004 | 0.003 | ** | |

| | edf | Ref.df | F | p-value | |
|---|---|---|---|---|---|
| s(ctime.exp) | 1.992 | 1.992 | 183.99 | <2e-16 | *** |
| s(time_in_D) | 7.948 | 8 | 668.73 | <2e-16 | *** |
| s(EE_kJH) | 6.565 | 6.565 | 20.69 | <2e-16 | *** |
| s(RQ) | 8.422 | 8.422 | 37.87 | <2e-16 | *** |

R-sq.(adj) = 0.561

Scale est.=0.0016156    n=7908

B)

log(EE_kJH)~diet+Sex+s(ctime.exp,k=3)+s(time_in_D,bs=cc)+s(H2Og)+s(RQ)

| | Estimate | Std. | Error | t value | Pr(>|t|) | |
|---|---|---|---|---|---|---|
| (Intercept) | -0.542 | 0.063 | -8.628 | <2e-16 | *** | |
| diet | -0.043 | 0.073 | -0.585 | 0.558 | | |
| SexM | 0.097 | 0.073 | 1.332 | 0.183 | | |

| | edf | Ref.df | F | p-value | |
|---|---|---|---|---|---|
| s(ctime.exp) | 1.980 | 1.980 | 39.660 | <2e-16 | *** |
| s(time_in_D) | 7.976 | 8.000 | 1464.100 | <2e-16 | *** |
| s(H2Og) | 7.114 | 7.114 | 28.930 | <2e-16 | *** |
| s(RQ) | 8.136 | 8.136 | 178.830 | <2e-16 | *** |

R-sq.(adj)=0.678

Scale est.=0.076286　　　n=7908

C)

log(RQ)~diet+Sex+s(ctime.exp,k=3)+s(time_in_D,bs=cc)+s(H2Og)+s(log(EE_kJH))

|  | Estimate | Std. | Error | t value | Pr(>\|t\|) |
|---|---|---|---|---|---|
| (Intercept) | -0.024 | 0.034 | -0.706 | 0.480 | |
| diet | -0.025 | 0.039 | -0.641 | 0.522 | |
| Sex | -0.008 | 0.039 | -0.210 | 0.834 | |

|  | edf | Ref.df | F | p-value | |
|---|---|---|---|---|---|
| s(ctime.exp) | 1.593 | 1.593 | 27.450 | 1.98E-07 | *** |
| s(time_in_D) | 7.693 | 8.000 | 44.530 | < 2.00E-16 | *** |
| s(H2Og) | 7.397 | 7.397 | 47.360 | < 2.00E-16 | *** |
| s(log(EE_kJH)) | 7.160 | 7.160 | 359.320 | < 2.00E-16 | *** |

R-sq.(adj)=0.393

Scale est.=0.0070625　　n=7908

Figure S1: Contributions of model terms to energy expenditure (EE kJ h$^{-1}$, A-D), water loss rate

(WLR g h$^{-1}$, E-H), and respiratory quotient (RQ, I-L) for female and male *Peromyscus eremicus*

across two diet groups (standard diet and low-fat diet) in general additive mixed models

(GAMM). The smoothing curves for each response variable included two fixed effects; diet

(low-fat vs standard) and sex, two random effects; mouse identification number and date of data

collection, and four regression terms: time in days, diurnal cycle, and two of the three

respirometry response variables (EE, RQ, and/or EWL) as regression terms. For each graph, the

y-axis is the effect of the x-axis on the respirometry response variable estimated by a

multivariable GAMM. Blue-shaded areas are 95% confidence intervals, black points are

residuals.

Table S3: Generalized additive mixed models (GAMM) results for only data collected during the

light phase. Statistical models and results for A) water loss rate (WLR, H2O g/hr$^{-1}$), B) energy

expenditure (EE kJ/hr$^{-1}$), C) respiratory quotient (RQ).

A)

H2Og~diet+Sex+s(ctime.exp,k=3)+s(EE_kJH)+s(RQ)

|  | Estimate | Std. | Error | t value | Pr(>\|t\|) |  |
|---|---|---|---|---|---|---|
| (Intercept) | 0.15503 | 0.00756 | 20.506 | <2.00E-16 | *** |  |
| diet | -0.058941 | 0.008737 | -6.746 | 1.69E-11 | *** |  |
| Sex | 0.035545 | 0.008736 | 4.069 | 4.80E-05 | *** |  |

|  | edf | Ref.df | F | p-value |  |
|---|---|---|---|---|---|
| s(ctime.exp) | 1.974 | 1.974 | 68.25 | <2e-16 | *** |
| s(EE_kJH) | 5.797 | 5.797 | 10.8 | <2e-16 | *** |
| s(RQ) | 5.309 | 5.309 | 102.25 | <2e-16 | *** |

R-sq.(adj)=0.349

Scale est.=0.0023669    n=4918

B)

log(EE_kJH)~diet+Sex+s(ctime.exp,k=3)+s(H2Og)+s(RQ)

|  | Estimate | Std. | Error | t value | Pr(>\|t\|) |  |
|---|---|---|---|---|---|---|
| (Intercept) | -0.92542 | 0.09277 | -9.975 | <2e-16 | *** |  |
| diet | -0.06415 | 0.10738 | -0.597 | 0.55 |  |  |
| Sex | 0.11275 | 0.10719 | 1.052 | 0.293 |  |  |

|  | edf | Ref.df | F | p-value |  |
|---|---|---|---|---|---|
| s(ctime.exp) | 1 | 1 | 6.991 | 0.00822 | ** |
| s(H2Og) | 6.797 | 6.797 | 7.118 | <2.00E-16 | *** |
| s(RQ) | 5.875 | 5.875 | 285.938 | <2.00E-16 | *** |

R-sq.(adj)=0.189

Scale est.=0.12779    n=4918

B)

log(RQ)~diet+Sex+s(ctime.exp,k=3)+s(H2Og)+s(log(EE_kJH))

Parametric        coefficients:

|  | Estimate | Std. | Error | t value | Pr(>|t|) |
|---|---|---|---|---|---|
| (Intercept) | 0.001688 | 0.036754 | 0.046 | 0.963 | |
| diet | 0.025869 | 0.042478 | 0.609 | 0.543 | |
| Sex | -0.014092 | 0.042453 | -0.332 | 0.74 | |

|  | edf | Ref.df | F | p-value | |
|---|---|---|---|---|---|
| s(ctime.exp) | 1.43 | 1.43 | 18.81 | 1.27E-05 | *** |
| s(H2Og) | 7.305 | 7.305 | 91.83 | <2.00E-16 | *** |
| s(log(EE_kJH)) | 6.539 | 6.539 | 362.65 | <2.00E-16 | *** |

R-sq.(adj)=0.283

Scale est.=0.0084596    n=4918

726　Table S4: Generalized additive mixed models (GAMM) results for only data collected during the

727　dark phase. Statistical models and results for A) water loss rate (WLR, H2O g/hr $^{-1}$), B) energy

728　expenditure (EE kJ/hr $^{-1}$ ), C) respiratory quotient (RQ).

A)

H2Og~diet+Sex+s(ctime.exp,k=3)+s(EE_kJH)+s(RQ)

| | Estimate | Std. | Error | t value | Pr(>\|t\|) |
|---|---|---|---|---|---|
| (Intercept) | 0.053382 | 0.00923 | 5.784 | 8.07E-09 | *** |
| diet | 0.002749 | 0.01071 | 0.257 | 0.797 | |
| Sex | 0.025882 | 0.010637 | 2.433 | 0.015 | * |

| | edf | Ref.df | F | p-value | |
|---|---|---|---|---|---|
| s(ctime.exp) | 1.968 | 1.968 | 19.063 | 4.29E-07 | *** |
| s(EE_kJH) | 4.881 | 4.881 | 8.185 | 6.41E-07 | *** |
| s(RQ) | 7.887 | 7.887 | 47.955 | <2.00E-16 | *** |

R-sq.(adj)=0.219

Scale est.=0.0018757　　n=2990

B)

log(EE_kJH)~experiment+Sex+s(ctime.exp,k=3)+s(H2Og)+s(RQ)

| | Estimate | Std. | Error | t value | Pr(>\|t\|) |
|---|---|---|---|---|---|
| (Intercept) | 0.08701 | 0.07403 | 1.175 | 0.24 | |
| diet | -0.04856 | 0.0857 | -0.567 | 0.571 | |
| Sex | 0.11419 | 0.08547 | 1.336 | 0.182 | |

| | edf | Ref.df | F | p-value | |
|---|---|---|---|---|---|
| s(ctime.exp) | 1.97 | 1.97 | 21.35 | <2e-16 | *** |
| s(H2Og) | 7.853 | 7.853 | 12.85 | <2e-16 | *** |
| s(RQ) | 7.469 | 7.469 | 34.63 | <2e-16 | *** |

R-sq.(adj)=0.12

Scale est.=0.058025     n=2990

C)

log(RQ)~experiment+Sex+s(ctime.exp,k=3)+s(H2Og)+s(log(EE_kJH))

Parametric coefficients:

| | Estimate | Std. | Error | t value | |
|---|---|---|---|---|---|
| (Intercept) | -0.07254 | 0.02994 | -2.423 | 0.0155 | * |
| diet | -0.08263 | 0.03458 | -2.39 | 0.0169 | * |
| Sex | -0.01154 | 0.03459 | -0.333 | 0.7388 | |

| | edf | Ref.df | F | p-value | |
|---|---|---|---|---|---|
| s(ctime.exp) | 1.95 | 1.95 | 20.1 | <2e-16 | *** |
| s(H2Og) | 6.454 | 6.454 | 45.5 | <2e-16 | *** |
| s(log(EE_kJH)) | 1 | 1 | 39.76 | <2e-16 | *** |

R-sq.(adj)=0.292

Scale est.=0.0032848     n=2990

732    Table S5: ANCOVA results for total water loss.

| | Sum | Sq | Df | F value | Pr(>F) |
|---|---|---|---|---|---|
| diet | 130.36 | 1 | 19.5126 | 5.22E-05 | *** |
| sex | 40.4 | 1 | 6.0471 | 0.01736 | * |
| weight | 7.03 | 1 | 1.0529 | 0.30967 | |
| diet:sex | 1.54 | 1 | 0.2305 | 0.63317 | |
| Residuals | 340.72 | 51 | | | |

| | estimate | SE | df | t.ratio | p.value |
|---|---|---|---|---|---|
| contrast | | | | | |
| F Low-fat - M Low-fat | -2.05 | 0.977 | 51 | -2.101 | 0.1667 |
| F Low-fat - F Standard | 2.76 | 1.018 | 51 | 2.716 | 0.0433 |
| F Low-fat - M Standard | 1.39 | 0.977 | 51 | 1.421 | 0.4922 |
| M Low-fat - F Standard | 4.82 | 1.014 | 51 | 4.748 | 0.0001 |
| M Low-fat - M Standard | 3.44 | 0.977 | 51 | 3.522 | 0.0049 |
| F Standard - M Standard | -1.38 | 1.013 | 51 | -1.359 | 0.5307 |