High total water loss driven by low-fat diet in desert-adapted mice

Danielle M. Blumstein[1], Jocelyn P. Colella[1,2], Ernst Linder[3], Matthew D. MacManes[1]

1. University of New Hampshire, Molecular, Cellular, and Biomedical Sciences

Department, Durham, NH 03824 (DMB Dani.Blumstein@unh.edu, Orcid ID: 0000-0003-

2129-5541; MDM Matthew.MacManes@unh.edu, Orcid ID: 0000-0002-2368-6960)

2. University of Kansas, Biodiversity Institute, Lawrence, KS 66045 (JPC

colella@ku.edu, Orcid ID: 0000-0003-2463-1029)

3. University of New Hampshire, Department of Mathematics and Statistics, Durham,

NH 03824 (EL ernst.linder@unh.edu)

Abstract

Availability of essential food resources is one of the most important drivers of survival

and to persist in environments where critical resource abundance is changing, animals

must either relocate or adapt in place. Testing an animal's ability to respond to

alternative conditions can reveal differences in physiological responses. We used flow-

through respirometry to characterize metabolic phenotypes of the desert-adapted

cactus mouse (*Peromyscus eremicus*) under diurnally variable environmental conditions

that mimic the Sonoran Desert and treated mice with two different diets; a standard diet

and a low-fat diet. We found significant diet-specific differences in the rate of water loss

and serum electrolyte values. Mice fed the low-fat diet lost more water relative to those

eating the standard diet and patterns of *de novo* lipogenesis are not limited by dietary

composition. Our results are consistent with the hypothesis that diet contributes to water

homeostasis. Compared to other desert animals, rodents have limited capacity to

dissipate heat using evaporative cooling, limiting thermoregulatory performance at

higher temperatures. We predict that a mismatch in physiological requirements and

environmental conditions could significantly impact small mammal survival in desert

environments if optimal foods become less abundant, as may be the case given climate

change.

Keywords: dietary fat, energy expenditure, *Peromyscus*, physiology, respirometry, total

evaporative water loss

Introduction

As the planet undergoes unprecedented climate change, increased desertification is

predicted (IPCC, 2019; Mirzabaev et al., 2019; Wu et al., 2014). Indeed, water limitation

is a strong selective force that will result in changes to species distributions and

community composition. Understanding these responses is critically important for

predicting the impacts of rapid climate change on biodiversity. Responses to climate

change vary but can be classified into two general categories: geographic relocation to

maintain physiological balance (Moritz et al., 2008; Riddell et al., 2021; Walsh et al.,

2016) or, for species with fragmented habitats or limited dispersal ability, adaptation in

place (Reed et al., 2011). With regards to the former, several studies have

demonstrated or predicted distributional changes of species in response to changing

climate (Moritz et al., 2008; Morrison et al., 2018; Parmesan, 2006; Parmesan and

Yohe, 2003; Pecl et al., 2017; Van de Peer et al., 2017). The latter response is less well

documented. Adaptation *in situ* may involve morphological, behavioral, or physiological

changes, such as directional selection for adaptive seasonal camouflage (Jones et al.,

2018), but in the absence of genetic variation, there is strong evidence of increased

extinction risk (Spielman et al., 2004).

The Sonoran Desert is hot, dry, and dynamic, temperatures fluctuate by >15℃

within a single day and >30℃ between seasons, and average relative humidity changes

by 20% between seasons (Reid, 1987; Sheppard et al., 2002; Unland et al., 1996). High

ambient temperatures and lack of extrinsic water challenges mammalian survival, as

most mammals rely on evaporative cooling (Reece et al., 2015). Despite these

challenges, many mammals thrive in deserts. This has led to decades of work aimed at

one central question: How do desert animals survive (Bittner et al., 2021; Bradford,

1974; Cortés et al., 2000; Kordonowy et al., 2017; MacManes, 2017; MacMillen and

Hinds, 1983; Nagy, 1987; Tigano et al., 2020)?

Animals that have evolved in desert environments are physiologically adapted to

extreme temperatures and aridity, and therefore represent an interesting natural

experimental model to examine adaptation to these conditions, and the potential for

adaptation given climate change. Behaviors such as foraging strategies (Manlick et al.,

2021), dietary preference (Frank, 1988), burrowing (Schmidt-Nielsen and Adolph, 1964;

Louw and Seely, 1984; Riddell et al., 2021), group nesting (Schmidt-Nielsen and

Adolph, 1964), seeking microhabitat refugia during the heat of the day (Riddell et al.,

2021), and nocturnal habits (but see Walsberg, 2000; Tracy and Walsberg, 2002) may

optimize fitness in hot, dry environments by allowing animals to regulate temperature

and water balance in real-time (Frank, 1988). Behavioral changes may be the first line

of defense against changing environments, as they are often dynamic, reversible, and

happen in real-time. In contrast, physiological adaptation, such as efficient kidney

function or low fecal water content which are observed in desert mammals to maintain

water balance, occur across evolutionary time scales (Christian and Matson, 1978;

Kordonowy and MacManes, 2016). Genomic studies have found evidence of positive

selection in desert mammals, acting on genes involved in energy production, food

storage (Jirimutu et al., 2012; Wu et al., 2014), water reabsorption (Giorello et al., 2018;

Jirimutu et al., 2012; Marra et al., 2012; Marra et al., 2014; Yang et al., 2016), and

osmoregulation (Colella et al., 2021a; Kordonowy and MacManes, 2016; MacManes

and Eisen, 2014). A strong signature of positive selection on Growth Differentiation

Factor 10 (*Gdf10*) has also been found and may prevent weight gain under high-fat

diets (Hino et al., 2017; Tigano et al., 2020). While this prior research has enhanced our

understanding of the physiologic processes underlying adaptation in desert

environments, we still don't fully understand the role that diet plays in water economy

regulation.

In environments like deserts, where extrinsic water is limited, endogenous water

production is likely critically important to survival, and dietary intake represents the raw

material for the production of metabolic water (Frank, 1988; Schmidt-Nielsen and

Adolph, 1964). Metabolic water is the water produced through catabolism of

carbohydrates, fats, or proteins. In deserts, insects, vegetation, and seeds comprise the

bulk of rodent diets (Reichman, 1975). Although specific dietary composition is often

unknown, an optimal diet balances energy gain (Pyke, 1984; Pyke et al., 1977),

nutrition, and capacity for water regulation (Schmidt-Nielsen and Adolph, 1964). Dietary

items exhibit seasonal availability in nature, which if consumed in proportion to

availability can influence water homeostasis (Orr et al., 2015; Wolf and del Rio, 2003).

Additionally, with regards to diet, water potential varies dramatically: oxidation of

carbohydrates yields 0.60g of metabolic water per gram, however, fats yield 1.07g per

gram due to greater oxygen requirements, while proteins yield the least metabolic water

(0.41g) and even require water loss for excretion of nitrogenous waste (Davidson and

Passmore, 1963; Mellanby, 1942; Schmidt-Nielsen and Adolph, 1964). We hypothesize

that a standard diet will maximize water conservation when compared to a low-fat diet.

Mice of the genus *Peromyscus* have the widest distribution of any North

American mammal and are considered a model organism in evolutionary biology due to

their unparalleled habitat diversity (Bedford and Hoekstra, 2015), extensive genomic

resources (*e.g.*, Tigano et al. 2020; Colella et al. 2021a), an abundance of historically

and contemporaneously collected samples in natural history museums (Pardi et al.,

2020; Pergams and Lawler, 2009), and ability to live and breed under laboratory

conditions (Crossland et al., 2014). The desert specialist cactus mouse, *Peromyscus*

eremicus, is endemic to the southwestern United States and exhibits behavioral (Murie,

1961; Veal and Caire, 1979), physiological (Colella et al., 2021b; Kordonowy et al.,

2017; Macmillen, 1965), and molecular (MacManes, 2017) adaptations to desert

environments. *Peromyscus eremicus* are omnivorous and opportunistic in their diet,

utilizing seeds, arthropods, and green vegetation seasonally (Bradley and Mauer, 1973;

Meserve, 1976) and shift to intense cactus consumption during the summer months (Orr

et al., 2015).

Given the relationship between dietary composition and metabolic water

production, the current study examines the effects of dietary fat content on the

metabolic physiology of the desert-adapted cactus mouse. To accomplish this, we

extend previously characterized circadian metabolic patterns for males and females

(Colella et al., 2021b) to rates of metabolism and water loss in animals fed an

experimental diet low in fat, but otherwise unchanged. Thereby testing the role that

dietary fat has on water balance, which has a direct connection to fitness.

Methods

Animal Care and Experimental Model

Animal care procedures followed guidelines established by the American Society of

Mammalogists (Sikes and Gannon, 2011; Sikes et al., 2016) and were approved by the

University of New Hampshire Institutional Animal Care and Use Committee under

protocol number 103092.

Sexually mature, non-reproductive healthy adults (n = 28 males, n = 28 females)

between three and nine months of age, bred from wild-derived lines maintained at the

Peromyscus Genetic Stock Center of the University of South Carolina (Columbia, South

Carolina, USA) were used in this study. Mice were randomly assigned to one of two

feeding groups: A standard diet group (SD - LabDiet® 5015*, 26.101% fat, 19.752%

protein, 54.148% carbohydrates, food quotient [FQ] 0.89, supplemental file 3) or a low-

fat diet group (LFD - Modified LabDiet® 5015 [5G0Z], 6.6% fat, 22.8% protein, 70.6%

carbohydrates, FQ = 0.92, supplemental file 4). Animals were housed individually in

9.5L animal chambers inside a larger, walk-in environmental chamber built to replicate

temperature, humidity, and photoperiodic conditions of the Sonoran Desert (Colella et

al., 2021c; Kordonowy et al., 2017). Each animal chamber contained dried cellulose-

based bedding, and animals were provided food and water *ad libitum*. Mice were

acclimated to the LFD for one month and to experimental cages for 24-hours prior to the

beginning of metabolic measurements. Diurnal temperature and relative humidity (RH)

fluctuation are fixed at 32°C and 10% RH during the light phase for 11 hours (06:00 hr

to 20:00 hr), and drop to 24°C and 25% RH during a 1 hr transition (T1) starting at 20:00

hr, then remains constant during the dark phase for seven hours, before the

temperature rises again and the RH falls to light phase temperature during a three-hour

transition (T2) that starts at 06:00 hr. Photoperiod was constant: 16 hr light, 8 hr dark

(Figure 1). Animals were weighed (g) on a digital scale before being housed in

experimental cages for three days during metabolic data collection and weighed again

at the end of the experiment.

Figure 1

72 hours of data collection plotted in a 24-hour window to display circadian patterns for
each group: organismal responses to standard diet are in blue and low-fat diet in gray.
Vertical blocks shaded in gray indicate the dark phase, and unshaded blocks indicate
light phase. A) Temperature (℃) is indicated by a solid red line. Transition 1 (T1)
occurred during the first shift from dark to light at 07:00. Transition 2 (T2) occurred
during the second shift from light to dark at 20:00. B) 24-hour measurements of energy
expenditure (EE kcal/hr^{-1}), C) rate of water loss (H$_2$O mg/hr^{-1}), and D) respiratory
quotient (RQ), for females (F, left) and males (M, right).

Metabolic phenotyping

Metabolic phenotyping data were collected using a previously described (Colella et al.,

2021b) commercially available respirometry system purchased from Sable Systems

International. Physiological data were collected following the methods outlined in

Whitfield et al. (2015) and Colella et al. (2021b). Oxygen consumption (VO_2 ml min^{-1}),

carbon dioxide production (VCO_2 ml min^{-1}), and water vapor (WVP, kPa) were

continually measured during 72 hr trials. This was repeated twice for each sex under

each diet (SD and LFD), resulting in three-day experiments performed sequentially in

eight batches of seven mice (and one empty baseline chamber per three-day

experiment) at a time, with batches alternating between females and males for a total of

28 adult males and 28 adult females.

Each cage was measured for 120s approximately twice every hour. We used the

most stable 50% of each measurement period to produce a single averaged VO_2 (ml

min^{-1}), VCO_2 (ml min^{-1}), and rate of water loss (RWL) (mg h^{-1}) value for that interval.

VO_2 and VCO_2 were calculated using equations 10.5 and 10.6, respectively, from

Lighton (2018). Respiratory quotient (RQ) was calculated as the ratio of VCO_2 to VO_2

(Lighton 2018). Energy expenditure (EE), kcal/hr^{-1}, was calculated as in Lighton (2018,

eq. 9.15). Values for each response variable that were more than three standard

deviations away from the mean, often representing times when animal care staff

entered the chamber, were considered outliers and removed from the downstream

analysis.

Statistical Analysis

All statistical analyses were conducted in R v 4.0.3 (R Core Team, 2020), unless

otherwise specified. Source code is available at

https://github.com/DaniBlumstein/Diet_paper. We used Generalized Additive Mixed

Models (mgcv::gamm, Lin and Zhang, 1999; Wood, 2017) to model the effect of time in

days, diurnal time, and EE, RQ and/or RWL on EE, RQ, and RWL. To avoid violating

the assumption of independence we used the experimental batches and the mice

nested within the experimental batch as random effects. The R function "gamm" then

simultaneously estimates the variances of these random effects as well as the treatment

effects, thus implicitly adjusting treatment and covariate effects for the effect of

correlated nested groups of measurements. As a result, treatments are not explaining

differences between individual mice, but rather the average differences between groups

of mice because we were interested in the diet treatment and the effect of sex, as well

as time traces over the full 72 hr period and the diurnal cycle. Our models included two

fixed effects, diet and sex, and four nonlinear smoothing regression terms: time in days

as a nonlinear trend, diurnal cycle as a circular spline function, and two of the three

respirometry response variables (EE, RQ, and/or RWL) as regression terms.

Electrolytes

At 12:00 EST, immediately following the conclusion of the experiment, animals were

weighed and euthanized following approved IACUC protocols. 120 uL of trunk blood

was collected within 1 minute of death for serum electrolyte measurement using an

Abaxis i-STAT® Alinity machine with the i-STAT CHEM8+ cartridge (Abbott Park, IL,

USA, Abbott Point of Care Inc). CHEM8+ cartridges measure the concentration of

sodium, potassium, creatinine, blood urea nitrogen, hematocrit, and ionized calcium,

which are expected to vary in response to hydration status and renal function. We used

a student's two-tailed t-test (stats::t.test) to test for significant ($p < 0.05$) differences in

weight after the treatments were applied and electrolytes between the sexes for each

experimental group.

Results

Data from 56 sexually mature (but non-reproductive) male and female mice were

collected as described above. All mice were subjected to standard animal care

procedures before the experiment which included a health assessment conducted by

licensed veterinary staff and weighing the animals. No significant difference in weight

between males and females was detected (mean of 22.2g versus 21.2g, respectively;

Table 1), nor between diet groups. No animals were removed from the study due to

health concerns.

		females			males		
		all	standard diet	low fat diet	all	standard diet	low fat diet
weight	median	21.94	22.35	21.79	21.55	22.55	19.90
	mean	22.35	22.29	22.40	21.13	22.24	20.02
	range	15.40-29.10	15.40-25.20	17.23-29.34	16.91-29.34	18.20-26.60	16.91-29.34
Na	median	143.00	143.00	145.00	140.00	140.50	137.50
	mean	143.05	142.50	143.60	139.23	140.71	137.50
	range	132-151	132-142	135-151	136-149	136-146	136-149
K	median	5.85	6.65	5.70	8.15	7.65	8.55
	mean	6.18	6.37	5.98	7.90	7.82	7.99
	range	4.30-9.00	5.50-9.00	4.30-9.00	3.90-9.00	6.70-9.00	3.90-8.90
Cr	median	0.20	0.20	0.20	0.20	0.20	0.20
	mean	0.20	0.20	0.20	0.21	0.20	0.22
	range	0.20-0.30	0.20-0.30	0.20-0.20	0.19-0.30	0.19-0.30	0.20-0.20
BUN	median	36.00	36.00	36.50	33.50	33.00	37.50
	mean	37.65	37.00	38.30	34.62	33.21	36.25
	range	26-48	26-43	32-48	26-46	26-40	32-46
Hct	median	35.00	35.00	34.50	32.00	32.00	32.50
	mean	35.30	36.20	34.40	32.08	32.50	31.58
	range	23-41	23-36	31-41	27-51	27-38	31-51
iCa	median	1.29	1.30	1.29	1.14	1.16	1.09
	mean	1.28	1.27	1.29	1.13	1.16	1.10
	range	0.95-1.42	0.95-1.28	1.18-1.42	1.04-1.39	1.041.28	1.10-1.39

Table 1

Summary statistics for weight (g) and serum electrolyte measurements (Na = sodium, K

= potassium, Cr = creatinine, BUN = blood urea nitrogen, Hct = hematocrit, iCa =

ionized calcium) for female and male *Peromyscus eremicus* across each diet group,

standard and low fat.

Metabolic Phenotyping

In total, we recorded 8,125 120s intervals: 4,091 female, 4,034 male (supplemental

table 1). We measured metabolic variables for 4,105 120s intervals under the LFD:

2,053 female and 2,052 male measurements. For the SD, we recorded 4,020 total

observations (2,038 female, 1,982 male). After removing outliers, we retained 3,945

female measurements (1,977 SD and 1,968 LFD) and 3,963 male measurements

(1,945 SD, 2,018 LFD). *Peromyscus eremicus* show circadian metabolism in phase with

photoperiod and room temperature for both diets (Figure 1), as described in Colella et

al. (2021b). All raw data (Expedata files) and processed machine-readable csv files are

available on Zenodo: https://zenodo.org/record/6422231#.YIXSD9PML0o.

Figure 2

Contributions of model terms to rate of water loss (RWL, A-D), energy expenditure (EE,

E-H), and C) respiratory quotient (RQ, I-L) for female and male *Peromyscus eremicus*

across two diet groups (standard diet and low fat diet) in general additive mixed models

(GAMM). The smoothing curves for each response variable included two fixed effects;

diet (low fat vs standard) and sex, two random effects; mouse identification number and

date of data collection, and four regression terms: time in days, diurnal cycle, and two of

the three respirometry response variables (EE, RQ, and/or EWL) as regression terms.

For each graph, the y-axis is the effect of the x-axis on the respirometry response

variable estimated by a multivariable GAMM. Blue shaded areas are 95% confidence

bands, black points are residuals.

Rate of Water Loss

Both groups of mice, SD and LFD, and both sexes exhibited diurnal patterning of water

loss and had the highest water loss during the light phase (Figure 1C), with peak loss

occurring between 09:00 and 11:00. During the second transition (light to dark) water

loss dropped sharply until 22:00 hours, before slowly increasing until the first transition

(dark to light). Overall, there were substantial differences in the amount of water lost

between males and females (male RWL mean: -2.37 - 7.35, female RWL mean: 1.56 -

5.66). For both males and females, mice on the low-fat diet lost more water during the

light phase (females LFD mean: 2.74, females SD mean: 1.61, males LFD mean: 3.32,

males SD mean: 2.18). Both diet and sex have a significant effect on RWL (diet p =

1.70e-05, sex p = 0.003, supplemental table 2). Time in days has a weakly negative

relationship with RWL even though it is highly significant (Figure 2A, supplemental 2).

The strongest relationship is seen between diurnal time and RWL (Figure 2B,

supplemental 2). Finally, EE and RQ, while highly significant, have a weak relationship

with RWL (Figure 2C, Figure 2D, supplemental table 2).

Energy Expenditure

Males and females for both diets show diurnal patterning of EE with the highest EE

occurring during the dark phase and the lowest EE during the light phase (Figure 1B).

Interestingly, males and females had different EE measurements during the dark phase

for each diet with females on the SD having higher EE (SD mean: 0.24, LFD mean:

0.28) and males on the LFD diet having higher EE (SD mean: 0.31, LFD mean: 0.29).

During the light phase, EE measurements stabilized to similar measurements for all

groups (females LFD mean: 0.09 females SD mean: 0.09, males LFD mean: 0.09,

males SD mean: 0.09). There was no significant difference in EE between the sexes

and the two diets (diet $p = 0.56$, sex $p = 0.18$, supplemental table 2). While significant,

time in days had no relationship with EE (Figure 2E, supplemental table 2). The

relationship that is also statically significant is seen between diurnal time and EE (Figure

2F, supplemental 2). Finally, RWL had a weak positive relationship with EE and RQ had

a negative relationship with EE (Figure 2G, Figure 2H, supplemental table 2).

Respiratory Quotient

RQ showed circadian rhythms for both diets and sexes with the highest RQ occurring

during the light phase and the lowest RQ occurring during the dark phase. During the

dark phase, RQ values were comparable to the food quotient (FQ) of the diet for both

sexes (LFD males mean: 0.92, LFD females mean: 0.94, SD males mean: 0.85, SD

females mean: 0.86, FQ LFD: 0.92, FQ SD: 0.89). During T1, RQ increased above the

FQ for both diets and sexes, with females on the SD having the highest RQ at the start

of the light phase and then stabilizing at 14:00 hours for the remainder of the light phase

(LFD males mean: 1.03, LFD females mean: 1.04, SD males mean: 1.03, SD females

mean: 1.08). We found no significant difference in RQ between the sexes and the two

diets (diet $p = 0.84$, sex $p = 0.52$, supplemental 2). While time in days, diurnal time,

RWL, and EE were highly significant, only EE has a weak relationship with RQ (Figure

2I-L, supplemental table 2).

Electrolytes

Serum electrolyte levels did not differ between males and females except for creatine

(males/females creatinine 0.21/0.20 $p < 0.05$, Table 1). Between experimental groups,

several serum electrolytes were significantly different (SD/LFD sodium 139.23/143.05 p

< 0.05, potassium 7.90/6.18 $p < 0.05$, creatinine 0.21/0.20 $p < 0.05$, blood urea nitrogen

34.62/37.65 $p > 0.05$, hematocrit 32.62/35.30 $p < 0.05$, ionized calcium 1.13/1.28 p <

0.05) (Figure 3, Table1).

Figure 3

Violin plots showing the distribution of serum electrolyte measurements (Na = sodium, K

= potassium, Crea = creatinine, BUN = blood urea nitrogen, Hct = hematocrit, iCa =

ionized calcium) for female (F) and male (M) *Peromyscus eremicus*. Observations (n =

56, 28 each sex) are represented by black dots and violin colors denot diet: blue for

standard diet and gray for low fat diet. Group means that are statically different are

denoted with one or more * (student's t test, p < 0.05) and insignificant comparisons are

denoted with ns (student's t test, p > 0.05). Note, the Abaxis i-STAT® Alinity analyzer

does not measure creatinine below 0.2 mg/dL, and therefore the range for normal

creatinine is truncated at this value.

Discussion

In light of current rapid climate change (IPCC, 2019), understanding how animals

respond to environmental change is of paramount importance. The ability to adapt in

place to changing climate will depend on a combination of adaptive plasticity and

genetic variation, as well as the intensity of selection and length of generation times.

Species unable to adapt in place could relocate but given the magnitude and rate of

climate change and extent of habitat fragmentation, species may be unable to shift

ranges (Beever et al., 2010; Henry et al., 2012; Sgrò et al., 2011). Extreme

environments (e.g. high altitude, deep sea, deserts) and the organisms living in them

represent natural experimental models, as the inhabitants have evolved behaviorally,

physiologically, and genomically to persist in these challenging habitats. Understanding

the mechanisms of survival under extreme conditions can provide insight into the

potential for other species to respond to increasingly erratic and extreme climatic

patterns. However, anthropogenic climate change is outpacing the evolutionary

timescale on which many of these adaptive mechanisms evolved. Metabolic responses

could aid in anticipating survival in novel scenarios but may not be sufficient for long

term survival.

Water balance is complex and is in part modulated by diet. Different

macronutrients yield different volumes of metabolic water via catabolism (carbohydrates

= 0.6g metabolic water per gram, protein = 0.4g per gram, and fats = 1.07g per gram,

Davidson et al., 1986). Many plants native to the Sonoran Desert have a fat content

comparable to that of the LFD used in this study (5.49 - 9.88% for leaf tissues and

branches, McArthur, 1994; 0.90- 3.60% for roots and seeds, Castle et al., 2020).

Moisture content can influence diet selection in wild desert mammals, which depend on

preformed water in their diet (Frank, 1988; Orr et al., 2015). Because all food was

dehydrated to match desert chamber conditions and by providing water *ad libitum,* we

can clearly analyze the role that dietary fat plays in water balance in *Peromyscus*

eremicus.

Here we expand on the long history of organismal water management studies in

desert taxa (Bradford, 1974; MacMillen and Hinds, 1983; Frank, 1988; Hayes et al.,

1998; Cortés et al., 2000; Albright et al., 2017; MacManes, 2017) to assess the

contribution of diet to physiological adaptation to a dry environment. Like other desert

mammals, *P. eremicus* rely on evaporative cooling for thermoregulation and thus, is

expected to employ a frugal water economy when water intake is limited. We

demonstrate diet-specific differences in the rate of evaporative water loss (RWL) and

serum electrolyte values, and a consistent pattern of daily *de novo* lipogenesis (DNL,

Colella et al., 2021b) that is not limited by dietary composition. These results support

the hypothesis that water conservation is at least partially tied to diet composition and

provide a deeper understanding of how desert animals maintain water homeostasis in

hot, arid environments. Thus, we predict that a mismatch in physiological requirements,

the availability of food type, and environmental conditions could significantly impact *P.*

eremicus survival, and perhaps other desert-dwelling animals.

Rate of Water Loss

In our study, animals fed a low-fat diet lost more water than did animals fed a diet higher

in fat (Figure 1c). Additionally, RWL was correlated with sex, with males losing more

water than females, mostly during their inactive daytime phase, when temperatures are

elevated and relative humidity is lower, and during environmental transition phases

when consuming the LFD. While the effects of temperature (Conley, 1985; Gordon,

1990; MacMillen and Hinds, 1983; Walsberg, 2000; Zhu et al., 2008) and weight

(MacMillen and Hinds, 1983; Van Sant et al., 2012) on RWL in mammals are well

described, with smaller-bodied mammals losing more water through evaporation and

water loss increasing rapidly at high temperatures, less understood is how diet affects

the RWL. Previous studies hypothesized that sexual dimorphism in water loss can be

explained by differences in body size, metabolism, and respiratory rate (Cryan and

Wolf, 2003), however, here we found no statistical difference in weight, consistent with

other investigations of this species (Colella et al., 2021b; Manning et al., 2006).

Breakdown of fats in the SD is expected to result in the formation of more water.

Therefore, in the absence of any source of water loss, the SD should lead to an

increase in hydration compared to the LFD. Dehydration would occur if the amount of

water lost is greater than the cumulative amount of water generated and consumed.

Numerous studies have identified reduced respiratory water loss as a major

physiological adaptation to maintaining positive water balance (Hart, 1971; MacMillen,

1972; Schmidt-Nielsen and Schmidt-Nielsen, 1951); however, not all diets have the

same energy uptake (Withers, 1982). Diets with lower potential for metabolic water

production, such as the LFD diet used in this experiment, would challenge survival

because total evaporative water loss alone could result in water imbalance, specifically

in a negative water balance that could not be compensated for by an increase in

exogenous water consumption. Not much is known about *P. eremicus* foraging

strategies and whether foraging varies between sexes. We predict that if *P. eremicus*

are forced to consume a LFD, they become more dehydrated, which likely has

consequences for survival and ultimately, fitness. In hot, arid deserts, where extrinsic

free water is severely limited, individual diet variation could significantly impact survival,

particularly when diet is constrained.

Energy Expenditure

While studies have shown that diet composition influences EE (Acheson, 1990;

Westerterp, 2004; Secor, 2009; Beale et al., 2018), there were no significant differences

in EE found in our study. Factors that influence EE can be intrinsic (i.e., body mass,

evolutionary history, reproductive stage, activity, or energy uptake) or extrinsic (i.e.,

habitat, climate, or social environment) (Speakman, 1997) and in wild populations,

dietary switches tend to coincide with seasonal changes that affect resource quantity

and/or quality (Noble et al., 2019). In the current study, we were able to isolate diet from

other extrinsic and intrinsic factors to show that diet is likely not responsible for

differences in EE. Consistent with other studies, we recorded a higher EE during the

dark phase of the 24-hour cycle (Colella et al., 2021b) further supporting that metabolic

rate is complex and may be modulated by the environment, activity, or other

extrinsic/intrinsic factors.

Respiratory Quotient

Our results suggest that *de novo* lipogenesis (DNL), as indicated by an RQ > 1.0 during

the light phase, is not limited by fat intake in this species. RQ, the ratio of O_2 consumed

to CO_2 produced, can inform rates of differential fuel consumption. RQ values typically

fall between 0.7 and 1.0, with the catabolism of fats yielding 0.7, carbohydrates 1.0, and

proteins intermediate values (Kleiber, 1975). RQ values exceed 1.0 when anabolism of

fat exceeds catabolism (i.e., DNL) (Abreu-Vieira et al., 2015; Benedict, 1937; Levin et

al., 2017) or during aerobic exercise (Whipp, 2007; Zagatto et al., 2012). In the

experiment described here and in previous experiments on *P. eremicus* (Colella et al.,

2021b)*,* RQ is only greater than 1.0 during the light phase when animals are inactive,

supporting DNL. While carbohydrates, fats, or protein are all burned as fuel, if present in

excess, adipose fat can be easily stored in animals (Barboza et al., 2009). One

mechanism for converting non-fat energy into a form that can be stored is DNL. The

conversion of glucose to fats prior to entry into the citric acid cycle increases heat and

results in an energetic cost, representing a metabolically inefficient use of dietary

substrate in cold habitats (Solinas et al., 2015), but a potentially efficient mechanism of

heat regulation in hot deserts. Previous studies have shown that DNL can occur under

high-carbohydrate and high-fat diets (Strable and Ntambi, 2010). This is supported in

our study as both groups had an RQ greater than 1.0 during the light phase and RQ

values at or below the FQ during the dark phase, which indicates a shift to the supplied

dietary substrate as the main oxidative fuel during cooler, dark periods.

Electrolytes

To further understand the impacts of diet on physiological function of endotherms in

desert environments, we collected serum electrolyte data from males and females fed

diets with varying levels of fat. We established diet-specific differences for serum

sodium (Na), potassium (K), creatinine (Cr), blood urea nitrogen (BUN), hematocrit

(Hct), and ionized calcium (iCa) values for *P. eremicus*. Electrolytes provide information

on the overall metabolic state, renal function, body water content, and other core

physiological functions (Kutscher, 1968), including hydration status (Cheuvront et al.,

2010) or disease state. While some electrolytes were elevated in response to the

experimental manipulation, others were not. Notably, the synthetic markers of

pathological renal impairment, BUN and Creatinine, were not significantly different in

any treatment group. Previous studies have shown that the electrolyte-water balance is

frequently unaffected when desert rodents are water-deprived for long periods of time

(Heimeier et al., 2004; Heimeier and Donald, 2006; Boumansour et al., 2021), however,

other desert rodent studies have demonstrated evidence of physiological stress in the

form of elevated serum electrolytes in the absence of exogenous water (Boumansour et

al., 2021; Kordonowy et al., 2017), though without apparent renal impairment.

Interestingly, we found several significant differences related to diet, several of

which were sexually dimorphic. Sodium was found to be significantly elevated in

females but not males (females SD: 140.50, females LFD: 143.60, $p = 0.004$, males SD:

140.71, males LFD: 137.50, $p = 0.210$) with *ad lib* access to water but fed a low-fat diet.

This is particularly interesting as both males and females fed a low-fat diet had higher

rates of water loss than when fed the standard diet. Elevated serum sodium in the

context of normal renal function is a classic sign of dehydration, but significant

dehydration does not appear to be the case here, as mice were given free access to

water, did not lose weight, and there were no differences in hematocrit for any treatment

group. Instead, dietary fats may themselves be responsible for changes in solute

balance. For instance, in rats, but not humans, differences in dietary fat were linked to

changes in serum sodium (Friedman et al., 2012), but in the opposite direction to what

we found (Abdel-Rahman, 2010). These changes were proposed to be aldosterone-

mediated (Song et al., 2004), a process for which we have no data. Whatever the

process, elevated serum sodium may have important and complex physiological

consequences, which may include elevated blood pressure and increased fluid

retention, both of which have obvious implications for survival in deserts and deserve

future study.

Conclusion

Desert-adapted species offer a unique opportunity to study how organisms survive

some of the harshest habitats on Earth. The main challenges to survival in desert

habitats are maintaining water balance and thermoregulation, with the nature of the diet

being critically important, particularly to the former. Examining the physiological

responses of animals fed experimentally modified diets aids in our understanding of

how animals may respond to changing environments where food availability varies. This

will ultimately be important in predicting an animal's response to climate change

because diet impacts water balance. To begin to understand ways in which the desert-

adapted cactus mouse *P. eremicus* responds to variation in dietary fat content,

reproductively mature mice were subjected to one of two diets, a standard diet or a low-

fat diet. Here, we present physiological measurements that show dietary fat is important

for water regulation in the desert-adapted cactus mouse.

We found that *P. eremicus* had a variety of responses to a decrease in dietary fat

through the analysis of interacting physiological traits: 1) high rates of total water loss

during the warmer, drier light phase and 2) a shift in serum electrolyte values indicating

physiological stress. These factors suggest that a low-fat diet could limit the capacity of

desert animals to tolerate limited access to free water as is common in arid

environments. Although we were able to collect hundreds of measurements for each

diet, sex, and time interval, our work has several limitations. Future studies should

further modify the major diet components (carbohydrates and proteins) and assess the

seasonal availability of such macronutrients in natural environments. Additionally, the

incorporation of both body temperature and activity monitoring can provide further

insight into the capacity of *P. eremicus* to tolerate environmental change.

The identification of short-term, physiological responses of *P. eremicus*

associated with diet composition demonstrates that adaptations that evolved over

longer evolutionary timescales affect survival during the accelerated time scales of

climate change. In light of global climate change and increased desertification,

investigating the range and mechanisms of plastic response of desert-adapted species

with regards to diet composition could provide insights into physiological responses to

increasingly erratic climate.

Acknowledgments

We would like to thank members of the MacManes lab for helpful comments and

support on previous versions of the manuscript; the Animal Resources Office and

veterinary care staff at the University of New Hampshire; A. Gerson at the University of

Massachusetts Amherst, Z. Cheviron at the University of Montana, B. Joos and J. Klok

at Sable Systems International for analytic guidance, technical support, and

respirometry training. This work was supported by the National Institute of Health

National Institute of General Medical Sciences (R35 GM128843 to M.D.M.).

Author Contributions

507 Conceptualization: M.D.M.; Methodology: D.M.B.; Formal analysis: D.M.B., E.L.;

508 Investigation: D.M.B., J.P.C.; Resources: M.D.M.; Writing - original draft: D.M.B.; Writing

509 - review & editing: D.M.B., J.P.C., E.L., M.D.M.; Visualization: D.M.B., E.L.; Supervision:

510 M.D.M.; Project administration: M.D.M.; Funding acquisition: M.D.M.

Competing Interests

513 No competing interests declared.

Data Availability

516 Raw ExpeData (SSI) files are available through Zenodo:

517 https://zenodo.org/record/6422231#.YlXSD9PML0o Macro processing files, raw and

518 processed respirometry data, and cage sampling scheme files are available also on

519 Zenodo. All R scripts used in this project are available through GitHub at:

520 https://github.com/DaniBlumstein/Diet_paper.

References

Abdel-Rahman, M. K. (2010). Influence of dietary fat on renal function, lipid profile, sex
hormones, and electrolyte balance in rats. *Eur. J. Lipid Sci. Technol.* **112**, 1166–1172.
Abreu-Vieira, G., Xiao, C., Gavrilova, O. and Reitman, M. L. (2015). Integration of body
temperature into the analysis of energy expenditure in the mouse. *Mol. Metab.* **4**, 461–
470.
Albright, T. P., Mutiibwa, D., Gerson, A. R., Smith, E. K., Talbot, W. A., O'Neill, J. J.,
McKechnie, A. E. and Wolf, B. O. (2017). Mapping evaporative water loss in desert
passerines reveals an expanding threat of lethal dehydration. *Proc. Natl. Acad. Sci.* **114**,
2283–2288.
Barboza, P. S., Parker, K. L. and Hume, I. D. eds. (2009). Lipids: Fatty Acids and Adipose
Tissue. In *Integrative Wildlife Nutrition*, pp. 119–131. Berlin, Heidelberg: Springer.
Bartholomew, G. A., Hudson, J. W. and Howell, T. R. (1962). Body Temperature, Oxygen

Consumption, Evaporative Water Loss, and Heart Rate in the Poor-Will. *The Condor* **64**, 117–125.

Bedford, N. L. and Hoekstra, H. E. (2015). Peromyscus mice as a model for studying natural variation. *eLife* **4**, e06813.

Beever, E. A., Ray, C., Mote, P. W. and Wilkening, J. L. (2010). Testing alternative models of climate-mediated extirpations. *Ecol. Appl.* **20**, 164–178.

Benedict, F. G. (1937). Lipogenesis in the animal body, with special reference to the physiology of the goose.

Bittner, N. K. J., Mack, K. L. and Nachman, M. W. (2021). Gene expression plasticity and desert adaptation in house mice. *Evol. Int. J. Org. Evol.* **75**, 1477–1491.

Boumansour, L., Benhafri, N., Guillon, G., Corbani, M., Touati, H., Dekar-Madoui, A. and Ouali-Hassenaoui, S. (2021). Vasopressin and oxytocin expression in hypothalamic supraoptic nucleus and plasma electrolytes changes in water-deprived male Meriones libycus. *Anim. Cells Syst.* **0**, 1–10.

Bradford, D. F. (1974). Water Stress of Free-Living Peromyscus Truei. *Ecology* **55**, 1407–1414.

Bradley, W. G. and Mauer, R. A. (1973). Rodents of a Creosote Bush Community in Southern Nevada. *Southwest. Nat.* **17**, 333–344.

Castle, S. T., Allan, N., Clifford, D., Aylward, C. M., Ramsey, J., Fascetti, A. J., Pesapane, R., Roy, A., Statham, M., Sacks, B., et al. (2020). Diet composition analysis provides new management insights for a highly specialized endangered small mammal. *PLOS ONE* **15**, e0240136.

Cheuvront, S. N., Kenefick, R. W., Montain, S. J. and Sawka, M. N. (2010). Mechanisms of aerobic performance impairment with heat stress and dehydration. *J. Appl. Physiol.* **109**, 1989–1995.

Christian D.P. and Matson J.O. (1978). Comparative water balance in two species of Liomys (Rodentia: Heteromyidae). *Comp. Biochem. Physiol. A.*

Colella, J. P., Tigano, A. and MacManes, M. D. (2020). A linked-read approach to museomics: Higher quality de novo genome assemblies from degraded tissues. *Mol. Ecol. Resour.* **20**, 856–870.

Colella, J. P., Tigano, A., Dudchenko, O., Omer, A. D., Khan, R., Bochkov, I. D., Aiden, E. L. and MacManes, M. D. (2021a). Limited Evidence for Parallel Evolution Among Desert-Adapted Peromyscus Deer Mice. *J. Hered.* **112**, 286–302.

Colella, J. P., Blumstein, D. M. and MacManes, M. D. (2021b). Disentangling environmental drivers of circadian metabolism in desert-adapted mice. *J. Exp. Biol.*

Colella, J. P., Blumstein, D. M. and MacManes, M. D. (2021c). Disentangling environmental drivers of circadian metabolism in desert-adapted mice. *J. Exp. Biol.* **224**,.

Conley, K. E. (1985). Evaporative water loss: thermoregulatory requirements and measurements in the deer mouse and white rabbit. *J. Comp. Physiol. B* **155**, 433–436.

Cortés, A., Rosenmann, M. and Bozinovic, F. (2000). Water economy in rodents: evaporative water loss and metabolic water production. *Rev. Chil. Hist. Nat.* **73**,.

Crossland, J. P., Dewey, M. J., Barlow, S. C., Vrana, P. B., Felder, M. R. and Szalai, G. J. (2014). Caring for Peromyscus spp. in research environments. *Lab Anim.* **43**, 162–166.

Cryan, P. M. and Wolf, B. O. (2003). Sex differences in the thermoregulation and evaporative water loss of a heterothermic bat, Lasiurus cinereus, during its spring migration. *J. Exp. Biol.* **206**, 3381–3390.

Davidson, S. and Passmore, R. (1963). Human nutrition and dietetics. *Hum. Nutr. Diet.*

Davidson, L. S. P., Passmore, R. and Eastwood, M. A. (1986). *Davidson and Passmore Human Nutrition and Dietetics.* Subsequent edition. Edinburgh ; New York: Churchill Livingstone.

Frank, C. L. (1988). Diet Selection by a Heteromyid Rodent: Role of Net Metabolic Water Production. *Ecology* **69**, 1943–1951.

Friedman, A. N., Ogden, L. G., Foster, G. D., Klein, S., Stein, R., Miller, B., Hill, J. O., Brill, C., Bailer, B., Rosenbaum, D. R., et al. (2012). Comparative effects of low-carbohydrate high-protein versus low-fat diets on the kidney. *Clin. J. Am. Soc. Nephrol. CJASN* **7**, 1103–1111.

Giorello, F. M., Feijoo, M., D'Elía, G., Naya, D. E., Valdez, L., Opazo, J. C. and Lessa, E. P. (2018). An association between differential expression and genetic divergence in the Patagonian olive mouse (Abrothrix olivacea). *Mol. Ecol.* **27**, 3274–3286.

Gordon, C. J. (1990). Thermal biology of the laboratory rat. *Physiol. Behav.* **47**, 963–991.

Hart, J. S. (1971). Rodents. In *Mammals*, pp. 1–149. Elsevier.

Hayes, J. P., Bible, C. A. and Boone, J. D. (1998). Repeatability of Mammalian Physiology: Evaporative Water Loss and Oxygen Consumption of Dipodomys merriami. *J. Mammal.* **79**, 475–485.

Heimeier, R. A. and Donald, J. A. (2006). The response of the natriuretic peptide system to water deprivation in the desert rodent, Notomys alexis. *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.* **143**, 193–201.

Heimeier, R. A., Bartolo, R. C. and Donald, J. A. (2004). The effect of water deprivation on signalling molecules that utilise cGMP in the spinifex hopping mouse Notomys alexis. *Aust. Mammal.* **26**, 191–198.

Henry, P., Sim, Z. and Russello, M. A. (2012). Genetic evidence for restricted dispersal along continuous altitudinal gradients in a climate change-sensitive mammal: the American pika. *Plos One* **7**, e39077.

Hino, J., Nakatani, M., Arai, Y., Tsuchida, K., Shirai, M., Miyazato, M. and Kangawa, K. (2017). Overexpression of bone morphogenetic protein-3b (BMP-3b) in adipose tissues protects against high-fat diet-induced obesity. *Int. J. Obes.* **41**, 483–488.

IPCC (2019). Global Warming of 1.5 C An IPCC Special Report on the Impacts of Global Warming of 1.5 C Above Pre-Industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change. *Sustain. Dev. Efforts Eradicate Poverty Httpswww Ipcc Chsr15 Accessed* **1**,.

Jirimutu, Wang, Z., Ding, G., Chen, G., Sun, Y., Sun, Z., Zhang, H., Wang, L., Hasi, S., Zhang, Y., et al. (2012). Genome sequences of wild and domestic bactrian camels. *Nat. Commun.* **3**, 1202.

Jones, M. R., Mills, L. S., Alves, P. C., Callahan, C. M., Alves, J. M., Lafferty, D. J. R., Jiggins, F. M., Jensen, J. D., Melo-Ferreira, J. and Good, J. M. (2018). Adaptive introgression underlies polymorphic seasonal camouflage in snowshoe hares. *Science* **360**, 1355–1358.

Kleiber, M. (1975). *The fire of life. An introduction to animal energetics.*

Kordonowy, L. L. and MacManes, M. D. (2016). Characterization of a male reproductive transcriptome for *Peromyscus eremicus* (Cactus mouse). *PeerJ* **4**, e2617.

Kordonowy, L., Lombardo, K. D., Green, H. L., Dawson, M. D., Bolton, E. A., LaCourse, S. and MacManes, M. D. (2017). Physiological and biochemical changes associated with acute experimental dehydration in the desert adapted mouse, *Peromyscus eremicus*. *Physiol. Rep.* **5**, e13218.

Kutscher, C. (1968). Plasma volume change during water-deprivation in gerbils, hamsters, guinea pigs and rats. *Comp. Biochem. Physiol.* **25**, 929–936.

Levin, E., Lopez-Martinez, G., Fane, B. and Davidowitz, G. (2017). Hawkmoths use nectar sugar to reduce oxidative damage from flight. *Science* **355**, 733–735.

Lighton, J. R. B. (2018). *Measuring Metabolic Rates: A Manual for Scientists*. 2nd ed. Oxford University Press.

Lin, X. and Zhang, D. (1999). Inference in Generalized Additive Mixed Models by Using Smoothing Splines. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **61**, 381–400.

Louw G. N. and Seely M. K. (1984). Ecology of desert organisms. *Vegetatio* **57**, 142–142.
MacManes, M. D. (2017). Severe acute dehydration in a desert rodent elicits a transcriptional
response that effectively prevents kidney injury. *Ren. Physiol* 11.
MacManes, M. D. and Eisen, M. B. (2014). Characterization of the transcriptome, nucleotide
sequence polymorphism, and natural selection in the desert adapted mouse
Peromyscus eremicus. *PeerJ* **2**, e642.
Macmillen, R. E. (1965). Aestivation in the cactus mouse, Peromyscus eremicus. *Comp.*
Biochem. Physiol. **16**, 227–248.
MacMillen, R. E. (1972). Water economy of nocturnal desert rodents.
MacMillen, R. E. and Hinds, D. S. (1983). Water Regulatory Efficiency in Heteromyid Rodents:
A Model and Its Application. *Ecology* **64**, 152–164.
Manlick, P. J., Maldonado, K. and Newsome, S. D. (2021). Competition shapes individual
foraging and survival in a desert rodent ensemble. *J. Anim. Ecol.* **n/a**,.
Manning, R. W., Jones, C. and Yancey, F. D. (2006). *Morphometric variation in two*
populations of the cactus mouse (Peromyscus eremicus) from Trans-Pecos Texas.
Lubbock, TX : Museum of Texas Tech University,.
Marra, N. J., Eo, S. H., Hale, M. C., Waser, P. M. and DeWoody, J. A. (2012). A priori and a
posteriori approaches for finding genes of evolutionary interest in non-model species:
Osmoregulatory genes in the kidney transcriptome of the desert rodent Dipodomys
spectabilis (banner-tailed kangaroo rat). *Comp. Biochem. Physiol. Part D Genomics*
Proteomics **7**, 328–339.
Marra, N. J., Romero, A. and DeWoody, J. A. (2014). Natural selection and the genetic basis
of osmoregulation in heteromyid rodents as revealed by RNA-seq. *Mol. Ecol.* **23**, 2699–
2711.
McArthur, E. D. (1994). *Nutritive Quality and Mineral Content of Potential Desert Tortoise Food*
Plants. Intermountain Research Station.
Mellanby, K. (1942). Metabolic Water and Desiccation. *Nature* **150**, 21–21.
Meserve, P. L. (1976). Habitat and Resource Utilization by Rodents of a California Coastal
Sage Scrub Community. *J. Anim. Ecol.* **45**, 647–666.
Mirzabaev, A., Wu, J., J. Evans, F. García-Oliva, I.A.G. Hussein, M.H. Iqbal, J. Kimutai, T.
Knowles, F. Meza, D. Nedjraoui, et al. (2019). Desertification. In: Climate Change and
Land: an IPCC special report on climate change, desertification, land degradation,
sustainable land management, food security, and greenhouse gas fluxes in terrestrial
ecosystems.
Moritz, C., Patton, J. L., Conroy, C. J., Parra, J. L., White, G. C. and Beissinger, S. R.
(2008). Impact of a Century of Climate Change on Small-Mammal Communities in
Yosemite National Park, USA. *Science* **322**, 261–264.
Morrison, L., Estrada, A. and Early, R. (2018). Species traits suggest European mammals
facing the greatest climate change are also least able to colonize new locations. *Divers.*
Distrib. **24**, 1321–1332.
Murie, M. (1961). Metabolic Characteristics of Mountain, Desert and Coastal Populations of
Perormyscus. *Ecology* **42**, 723–740.
Nagy, K. A. (1987). Field Metabolic Rate and Food Requirement Scaling in Mammals and
Birds. *Ecol. Monogr.* **57**, 111–128.
Noble, J. D., Collins, S. L., Hallmark, A. J., Maldonado, K., Wolf, B. O. and Newsome, S. D.
(2019). Foraging strategies of individual silky pocket mice over a boom–bust cycle in a
stochastic dryland ecosystem. *Oecologia* **190**, 569–578.
Orr, T. J., Newsome, S. D. and Wolf, B. O. (2015). Cacti supply limited nutrients to a desert
rodent community. *Oecologia* **178**, 1045–1062.
Pardi, M. I., Terry, R. C., Rickart, E. A. and Rowe, R. J. (2020). Testing climate tracking of
montane rodent distributions over the past century within the Great Basin ecoregion.

688   *Glob. Ecol. Conserv.* **24**, e01238.

689 **Parmesan, C.** (2006). Ecological and Evolutionary Responses to Recent Climate Change.
690   *Annu. Rev. Ecol. Evol. Syst.* **37**, 637–669.

691 **Parmesan, C. and Yohe, G.** (2003). A globally coherent fingerprint of climate change impacts
692   across natural systems. *Nature* **421**, 37–42.

693 **Pecl, G. T., Araújo, M. B., Bell, J. D., Blanchard, J., Bonebrake, T. C., Chen, I.-C., Clark, T.**
694   **D., Colwell, R. K., Danielsen, F., Evengård, B., et al.** (2017). Biodiversity redistribution
695   under climate change: Impacts on ecosystems and human well-being. *Science* **355**,
696   eaai9214.

697 **Pergams, O. R. W. and Lawler, J. J.** (2009). Recent and Widespread Rapid Morphological
698   Change in Rodents. *PLOS ONE* **4**, e6452.

699 **Pyke, G. H.** (1984). Optimal Foraging Theory: A Critical Review. *Annu. Rev. Ecol. Syst.* **15**,
700   523–575.

701 **Pyke, G. H., Pulliam, H. R. and Charnov, E. L.** (1977). Optimal Foraging: A Selective Review
702   of Theory and Tests. *Q. Rev. Biol.* **52**, 137–154.

703 **R Core Team** (2020). *R: A language and environment for statistical computing. R Foundation*
704   *for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.*

705 **Reece, W. O., Erickson, H. H., Goff, J. P. and Uemura, E. E.** (2015). *Dukes' Physiology of*
706   *Domestic Animals.* John Wiley & Sons.

707 **Reed, T. E., Schindler, D. E. and Waples, R. S.** (2011). Interacting Effects of Phenotypic
708   Plasticity and Evolution on Population Persistence in a Changing Climate. *Conserv. Biol.*
709   **25**, 56–63.

710 **Reichman, O. J.** (1975). Relation of Desert Rodent Diets to Available Resources. *J. Mammal.*
711   **56**, 731–751.

712 **Reid, W. T.** (1987). A critical analysis of Eastern Mojave desert temperatures.

713 **Riddell, E. A., Iknayan, K. J., Hargrove, L., Tremor, S., Patton, J. L., Ramirez, R., Wolf, B.**
714   **O. and Beissinger, S. R.** (2021). Exposure to climate change drives stability or collapse
715   of desert mammal and bird communities. *Science* **371**, 633–636.

716 **Schmidt-Nielsen, K. and Adolph, E. F.** (1964). Desert Animals: Physiological Problems of
717   Heat and Water. *Physiol. Zool.* **37**, 338–339.

718 **Schmidt-Nielsen, B. and Schmidt-Nielsen, K.** (1951). A complete account of the water
719   metabolism in kangaroo rats and an experimental verification. *J. Cell. Comp. Physiol.* **38**,
720   165–181.

721 **Sgrò, C. M., Lowe, A. J. and Hoffmann, A. A.** (2011). Building evolutionary resilience for
722   conserving biodiversity under climate change. *Evol. Appl.* **4**, 326–337.

723 **Sheppard, P. R., Comrie, A. C., Packin, G. D., Angersbach, K. and Hughes, M. K.** (2002).
724   The climate of the US Southwest. *Clim. Res.* **21**, 219–238.

725 **Sikes, R. S. and Gannon, W. L.** (2011). Guidelines of the American Society of Mammalogists
726   for the use of wild mammals in research. *J. Mammal.* **92**, 235–253.

727 **Sikes, R. S. and the Animal Care and Use Committee of the American Society of**
728   **Mammalogists** (2016). 2016 Guidelines of the American Society of Mammalogists for
729   the use of wild mammals in research and education. *J. Mammal.* **97**, 663–688.

730 **Solinas, G., Borén, J. and Dulloo, A. G.** (2015). De novo lipogenesis in metabolic
731   homeostasis: More friend than foe? *Mol. Metab.* **4**, 367–377.

732 **Song, J., Hu, X., Shi, M., Knepper, M. A. and Ecelbarger, C. A.** (2004). Effects of dietary fat,
733   NaCl, and fructose on renal sodium and water transporter abundances and systemic
734   blood pressure. *Am. J. Physiol. Renal Physiol.* **287**, F1204-1212.

735 **Speakman, J.** (1997). Factors influencing the daily energy expenditure of small mammals.
736   *Proc. Nutr. Soc.* **56**, 1119–1136.

737 **Spielman, D., Brook, B. W. and Frankham, R.** (2004). Most species are not driven to
738   extinction before genetic factors impact them. *Proc. Natl. Acad. Sci.* **101**, 15261–15264.

Strable, M. S. and Ntambi, J. M. (2010). Genetic control of de novo lipogenesis: role in diet-
induced obesity. *Crit. Rev. Biochem. Mol. Biol.* **45**, 199–214.

Tigano, A., Colella, J. P. and MacManes, M. D. (2020). Comparative and population genomics
approaches reveal the basis of adaptation to deserts in a small rodent. *Mol. Ecol.* **29**,
1300–1314.

Tracy, R. L. and Walsberg, G. E. (2002). Kangaroo rats revisited: re-evaluating a classic case
of desert survival. *Oecologia* **133**, 449–457.

Unland, H. E., Houser, P. R., Shuttleworth, W. J. and Yang, Z.-L. (1996). Surface flux
measurement and modeling at a semi-arid Sonoran Desert site. *Agric. For. Meteorol.* **82**,
119–153.

Van de Peer, Y., Mizrachi, E. and Marchal, K. (2017). The evolutionary significance of
polyploidy. *Nat Rev Genet* **18**, 411–424.

Van Sant, M. J., Oufiero, C. E., Muñoz-Garcia, A., Hammond, K. A. and Williams, J. B.
(2012). A Phylogenetic Approach to Total Evaporative Water Loss in Mammals. *Physiol.*
Biochem. Zool. **85**, 526–532.

Veal, R. and Caire, W. (1979). Peromyscus eremicus. *Mamm. Species* 1–6.

Walsberg, G. E. (2000). Small Mammals in Hot Deserts: Some Generalizations Revisited.
BioScience **50**, 109.

Walsh, R. E., Aprígio Assis, A. P., Patton, J. L., Marroig, G., Dawson, T. E. and Lacey, E.
A. (2016). Morphological and dietary responses of chipmunks to a century of climate
change. *Glob. Change Biol.* **22**, 3233–3252.

Whipp, B. J. (2007). Physiological mechanisms dissociating pulmonary CO2 and O2 exchange
dynamics during exercise in humans. *Exp. Physiol.* **92**, 347–355.

Whitfield, M. C., Smit, B., McKechnie, A. E. and Wolf, B. O. (2015). Avian thermoregulation in
the heat: scaling of heat tolerance and evaporative cooling capacity in three southern
African arid-zone passerines. *J. Exp. Biol.* **218**, 1705–1714.

Withers, P. C. (1982). Effect of diet and assimilation efficiency on water balance for two desert
rodents. *J. Arid Environ.* **5**, 375–384.

Wolf, B. O. and del Rio, C. M. (2003). How important are columnar cacti as sources of water
and nutrients for desert consumers? A review. *Isotopes Environ. Health Stud.* **39**, 53–67.

Wood, S. (2017). Generalized Additive Models: An Introduction with R, Second Edition.

Wu, H., Guang, X., Al-Fageeh, M. B., Cao, J., Pan, S., Zhou, H., Zhang, L., Abutarboush, M.
H., Xing, Y., Xie, Z., et al. (2014). Camelid genomes reveal evolution and adaptation to
desert environments. *Nat. Commun.* **5**, 1–10.

Yang, J., Li, W.-R., Lv, F.-H., He, S.-G., Tian, S.-L., Peng, W.-F., Sun, Y.-W., Zhao, Y.-X., Tu,
X.-L., Zhang, M., et al. (2016). Whole-Genome Sequencing of Native Sheep Provides
Insights into Rapid Adaptations to Extreme Environments. *Mol. Biol. Evol.* **33**, 2576–
2592.

Zagatto, A. M., Miyagi, W. E., Sakugawa, R. L., Kaminagakura, E. I. and Papoti, M. (2012).
Aerobic Endurance Measurement by Respiratory Exchange Ratio during a Cycle
Ergometer Graded Exercise Test. *J. Exerc. Physiol. Online* **15**, 49–56.

Zhu, W.-L., Jia, T., Lian, X. and Wang, Z.-K. (2008). Evaporative water loss and energy
metabolic in two small mammals, voles (Eothenomys miletus) and mice (Apodemus
chevrieri), in Hengduan mountains region. *J. Therm. Biol.* **33**, 324–331.

Supplemental

Number of samples per time interval

time interval	diet	number of 120 intervals male	number 120 intervals female
light	standard	867	879
	low fat	914	927
dark	standard	758	755
	low fat	758	719
T1	standard	233	256
	low fat	259	235
T2	standard	87	87
	low fat	87	87

GAMM tables

H2Omg ~ diet + sex + s(days, k = 3) + s(diurnal, bs = "cc") + s(EE) + s(RQ)

	Estimate Std.	Error	t value	Pr(>|t|)	
intercept	1.372	0.129	10.625	< 2e-16	***
diet	0.642	0.149	4.304	1.70E-05	***
sex	0.448	0.141	3.007	0.003	**

	edf	Ref.df	F	p-value	
s(days)	1.992	1.992	170.560	<2e-16	***
s(diurnal)	7.949	8.000	683.880	<2e-16	***
s(EE)	6.630	6.630	21.190	<2e-16	***
s(RQ)	8.413	8.413	38.190	<2e-16	***

log(EE) ~ diet + sex + s(days, k = 3) + s(diurnal, bs = "cc") + s(H2Omg) + s(RQ)

	Estimate Std.	Error	t value	Pr(>|t|)	
intercept	-2.017	0.063	-31.907	<2e-16	***
diet	0.042	0.073	0.578	0.563	
sex	0.098	0.073	1.341	0.180	

	edf	Ref.df	F	p-value	
s(days)	1.979	1.979	38.860	<2e-16	***
s(diurnal)	7.977	8.000	1482.280	<2e-16	***
s(H2Omg)	7.063	7.063	28.730	<2e-16	***
s(RQ)	8.154	8.154	180.690	<2e-16	***

log(RQ) ~ diet + sex + s(day, k = 3) + s(diurnal, bs = "cc") + s(H2Omg) + s(log(EE))

	Estimate Std.	Error	t value	Pr(>|t|)	
intercept	-0.049	0.034	-1.446	0.148	
diet	0.025	0.039	0.640	0.522	
sex	-0.008	0.039	-0.206	0.837	

	edf	Ref.df	F	p-value	
s(days)	1.547	1.547	25.610	<2e-16	***
s(diurnal)	7.706	8.000	46.520	<2e-16	***
s(H2Omg)	7.387	7.387	48.360	<2e-16	***
s(log(EE))	7.149	7.149	359.650	<2e-16	***