Phylogenomic patterns of divergence and gene flow detail the evolution of reinforcement and

hybrid speciation in *Phlox* wildflowers

Austin G. Garner [1,2*], Benjamin E. Goulet-Scott [1,2], and Robin Hopkins [1,2*]

[1] Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA

021382, USA

[2] The Arnold Arboretum, Harvard University, Boston, MA 02131, USA

[*] Authors to whom correspondence should be addressed: aggarner@g.harvard.edu;

rhopkins@fas.harvard.edu

ABSTRACT

It is becoming increasingly clear that the evolutionary histories of many groups of

organisms are riddled with reticulation. Some clades of organisms appear to be hotspots of

speciation by hybridization and gene flow. For example, there are two cases of reinforcement

and up to nine cases of hypothesized hybrid speciation in the eastern standing *Phlox*

(*Polemoniaceae*) wildflowers. However, the relationships of these taxa and the hypotheses about

their evolution have received limited attention with genomic data. We performed a

comprehensive phylogenomic study of the eastern standing *Phlox* to resolve their evolutionary

relationships and test long-standing hypotheses about if and how hybridization and gene flow

acted as creative forces in their diversification. Our phylogenomic analyses on genome-wide

markers provides resolution into the evolutionary relationships of thirty-two eastern standing

Phlox taxa, including well-supported non-monophyletic relationships across species complexes.

Explicit tests for introgression support gene flow occurring alongside the evolution of one of the

two cases of reinforcement, and patterns of divergence and gene flow support one of the five

hypothesized homoploid hybrid speciation events. Additionally, comparative read mapping

allowed assignment of putative subgenome ancestries for four allotetraploids hybrid species. Our

findings demonstrate the utility and importance of phylogenomics in confirming hypothesized

evolutionary histories of non-model systems and add to the growing evidence that gene flow

across species boundaries can play a role in the generation of novel biodiversity.

Keywords: phylogeny; RADseq; speciation; reinforcement; hybridization; gene flow; hybrid

species; polyploid

INTRODUCTION

Hybridization and gene flow can play important and diverse roles during the formation

and divergence of species (Taylor and Larson 2019). While these processes are often viewed as

homogenizing forces that mix genetic variation across species boundaries, they can also generate

novel variation. For example, hybridization and gene flow can create stable hybrid species and/or

fuel the diversification of adaptive radiations (Abbott et al. 2013; Schumer et al. 2014; Marques

et al. 2019). The act of hybridization itself can also create selective pressures favoring increased

reproductive trait divergence to reduce mating between species, i.e. reinforcement (Howard

1993; Garner et al. 2018). The effect of hybridization and gene flow on speciation has been

investigated across the tree of life, yet we still lack a broader understanding of the frequency and

extent to which these forces impact speciation across clades of organisms (but see (Eaton et al.

2015; Pease et al. 2016)). Characterizing the context and consequences of hybridization and gene

flow across groups of species is necessary for discerning how these forces influence the

generation and maintenance of biodiversity.

Much of the foundational work on hybridization as a creative force in speciation comes

from historic research on non-model plant systems (Anderson 1949; Stebbins 1959; Grant 1981).

Patterns of morphological, physiological, and ecological trait variation across and within plant

species created taxonomic conflict, inspiring biologists to infer evolutionary histories of

hybridization and gene flow between species (Goulet et al. 2017). In particular, populations with

individuals exhibiting intermediate, recombinant, and novel trait combinations where closely

related species come into geographic contact motivated hypotheses about the formation of

polyploid and homoploid hybrid species (Gottlieb 1972; Soltis and Soltis 2009; Yakimowski and

Rieseberg 2014; Barker et al. 2016).

Additionally, patterns of trait divergence in sympatry, where closely related species

coexist, but not in allopatry, motivated hypotheses of reinforcement during plant speciation

(Hopkins 2013). Reinforcement occurs when maladaptive hybridization between species

generates selection for traits that increase reproductive isolation. In plants, reinforcement can

result in divergence in reproductive traits such as flower color, flowering time, and cross-

compatibility in sympatry (McNeilly and Antonovics 1968; Levin 1985; Fishman and Wyatt

1999). Although hybridization drives selection for reinforcement, gene flow and recombination

can impede adaptive divergence by reinforcement (Felsenstein 1981). Therefore, much debate

and theory has focused on if and how reinforcement can evolve if hybridization also causes

interspecific gene flow (Servedio and Noor 2003). Yet, only a few empirical studies have

evaluated the presence and extent of gene flow alongside the evolution of reinforcement in

nature (Kulathinal et al. 2009; Lemmon and Juenger 2017; Roda et al. 2017; Turissini and

Matute 2017; Dyer et al. 2018).

Most hypotheses of hybrid speciation and reinforcement are based on patterns of

phenotypic variation and/or a few genetic markers. However, this evidence can be misleading

(Noor 1999; Goulet-Scott et al. 2021; Poelstra et al. 2021). Therefore, it is important to return to

known and hypothesized cases of reticulate evolution with genomic tests for gene flow to

confirm these evolutionary processes and further understand the context of their evolution

(Goulet et al. 2017). Hybridization and gene flow result in the mixing of genetic variation

between species. Therefore, if these processes coincide with the formation of novel diverging

lineages, admixture of parental species' genetic variation will be present in the novel diverging

lineage's genome.

Traditional phylogenetic methods infer evolution along bifurcating trees. However,

modern analyses can use discordance in allele patterns under these bifurcating trees (Green et al.

2010; Durand et al. 2011) or modeling of allele frequencies at tips (Gutenkunst et al. 2009;

Pickrell and Pritchard 2012; Excoffier et al. 2013) to identify a history of gene flow between

non-sister taxa and/or the formation of new lineages through hybridization (Kong and Kubatko

2021). Under a bifurcating phylogenetic framework, novel lineages evolving by reinforcement

with gene flow or hybrid speciation should be observed sister to one of their parental lineages

and have a significant signature of gene flow with these methods from their secondary parental

lineage. In this way, evolutionary relationships inferred from genomic data alongside

phylogenetic tests for gene flow can evaluate and confirm histories of genetic exchange between

lineages (Hibbins and Hahn 2021). Contemporary genome-wide sequencing methods, such as

restriction site-associated DNA (RAD) sequencing (Andrews et al. 2016) allow application of

these methods to radically clarify the evolutionary histories of reticulate non-model systems

(Eaton and Ree 2013; Eaton et al. 2015; Bombonato et al. 2020; Léveillé-Bourret et al. 2020;

Guo et al. 2021; Suissa et al. 2022).

The eastern standing *Phlox* (*Polemoniaceae*) are an ideal system for exploring the role of

hybridization and gene flow in speciation. They form a monophyletic group (Landis et al. 2018)

and are distinct from the rest of *Phlox* by their upright growth habit and natural occurrence

across mid to eastern North America. The natural ranges of these taxa intertwine and overlap

extensively, generating numerous zones of species contact (Fig. 1). Reproductive barriers

between these species are highly permissive (Levin 1966a) and a mix of morphological,

ecological, and biochemical evidence suggests pervasive gene flow between some taxa

Figure 1. Map of individuals sampled across the interdigitating ranges of the thirty-two eastern standing *Phlox* taxa included for phylogenetic analyses. Each dot represents the locality of a single individual with the color corresponding to an individual's taxonomic identity. Three insets show *P. pilosa* subsp. *sangamonensis* (brown), *P. pilosa* subsp. *fulgida* (blue), and *P. divaricata* subsp. *divaricata* (green) in their natural habitat.

(Anderson and Gage 1952; Erbe and Turner 1962; Levin 1966b; Levin and Smith 1966; Levin

and Kerster 1967; Levin and Schaal 1970a; Levin 1985)**.** This clade has served as testing

grounds of evolutionary theory on how reproductive isolation, hybridization, and gene flow can

influence species divergence, including hybrid speciation and reinforcement. The scientific

legacy of the eastern standing *Phlox* makes it a prime system to use modern genomic methods to

study how species diverge with hybridization and gene flow.

Phlox has been a taxonomically difficult genus (Locklear 2011a). Traditional delineations

of *Phlox* species relied on morphological, ecological, and geographic characteristics, resulting in

a long history of describing, promoting, and demoting species, subspecies, and varieties

(Whitehouse 1945; Wherry 1955, 1965; Erbe and Turner 1962; Turner 1998; Locklear 2011b,

2011a). Conspicuous variation in pistil style length has been used to divide the eastern standing

Phlox into long-styled (10-26mm) and short-styled (1.5-4mm) species groups, with the style

longer or shorter than the sepals respectively (Wherry 1930, 1931, 1932, 1955; Ferguson et al.

1999)**.** However, other morphological traits vary widely within and between species, and

species-level relationships within the group remain either poorly supported or incongruent across

phylogenetic studies (Ferguson et al. 1999; Ferguson and Jansen 2002; Landis et al. 2018;

Goulet-Scott et al. 2021). Resolving the phylogenetic relationship of the eastern standing *Phlox*

will provide a powerful framework for testing evolutionary hypotheses of hybridization and gene

flow during their diversification.

In this small clade, there are five hypothesized cases of homoploid hybrid species (*P.*

argillacea, *P. maculata* subsp. *pyramidalis*, *P. pilosa* subsp. *deamii*, *P. pilosa* subsp. *detonsa*,

and *P. amoena* subsp. *lighthipei*) and at least four cases of allotetraploid hybrid species (*P.*

villosissima subsp. *villosissima, P. villosissima* subsp. *latisepala, P. floridana, P. buckleyi*)

(Table 1). Support for these hypotheses were originally based on the observation that putative

hybrid lineages had reproductive, morphological, and ecophysiological trait values that appear to

be intermediate, recombined, or transgressive with putative parental taxa, making them

potentially reproductively and ecologically distinct from their parental lineages (Levin 1963,

1969; Levin and Smith 1966; Hadley and Levin 1969)**.** Further analysis of karyology (Smith and

Levin 1967; Levin 1968), cross compatibility (Levin 1966a), biochemical profiles (Levin and

Schaal 1970a, 1972; Levy and Levin 1974, 1975), and microsatellite markers (Fehlberg et al.

2014) built support for these hypotheses. Finally, natural hybrid zones and synthetic crosses

between pairs of putative parental taxa generated hybrids with phenotypes similar to the putative

stable hybrid lineages in nature (Levin 1963, 1966b; Levin and Smith 1966). However, other

treatments of these putative hybrids species have assumed them to be diverging varieties without

hybridization defining their formation (Wherry 1956) or found no support for hybrid ancestry

with genome-wide markers (Goulet-Scott et al. 2021).

In this same clade of *Phlox,* there are also two cases of reinforcement. During the

speciation of both *P. drummondii* and *P. pilosa* subsp. *pilosa*, flower color evolution within

sympatric populations decreases hybridization with closely related species (reviewed in (Hopkins

2013), Fig. 3). *P. drummondii* has evolved from being light-blue colored to dark-red to prevent

hybridization with sympatric light-blue colored *P. cuspidata* (Levin 1985; Hopkins and Rausher

2012), and *P. pilosa* subsp. *pilosa* flowers have evolved from pink colored to white to prevent

hybridization where it cooccurs in high frequency with pink colored *P. glaberrima* subsp.

interior near Lake Michigan (Levin and Kerster 1967). Biochemical evidence suggests gene flow

occurs between sympatric *P. pilosa* subsp. *pilosa* and *P. glaberrima* subsp. *interior* (Levin and

Schaal 1972), and genomic analyses have inferred a history of gene flow between sympatric *P.*

Table 1. Hybrid speciation hypotheses in the eastern standing *Phlox*

Hypothesized Hybrid Species	Ploidy	Putative Parent Species 1	Putative Parent Species 2	Motivating Previous Evidence
P. argillacea	2N	*P. glaberrima* subsp. *interior*	*P. pilosa* subsp. *pilosa* North	morphology[A], physiology[A], seed protein polymorphism[B], experimental crosses[C]
P. maculata subsp. *pyramidalis*	2N	*P. glaberrima* subsp. *interior*	*P. maculata* subsp. *maculata*	morphology[D], physiology[E], seed protein polymorphism[F], experimental crosses[C]
P. amoena subsp. *lighthipei*	2N	*P. pilosa* subsp. *pilosa* South	*P. amoena* subsp. *amoena*	morphology[G], seed protein polymorphism[F], experimental crosses[CG]
P. pilosa subsp. *deamii*	2N	*P. pilosa* subsp. *pilosa* South	*P. amoena* subsp. *amoena*	morphology[G], experimental crosses[CG], microsatellites[H]
P. pilosa subsp. *detonsa*	2N	*P. pilosa* subsp. *pilosa* South	*P. carolina*	seed protein polymorphism[F], experimental crosses[C]
P. floridana	4N	*P. pilosa* subsp. *pilosa* South	*P. carolina* subsp. *angusta*	karyology[J], flavonoids[K], seed protein polymorphism[F], experimental crosses[C]
P. villosissima subsp. *villosissima*	4N	*P. pilosa* subsp. *pilosa* South	*P. drummondii*	karyology[J], flavonoids[L], seed protein polymorphism[F], experimental crosses[C]
P. villosissima subsp. *latisepala*	4N	*P. pilosa* subsp. *pilosa* South	*P. drummondii*	karyology[J], flavonoids[L], seed protein polymorphism[F], experimental crosses[C]
P. buckleyi	4N	unknown	unknown	karyology[J]

Supporting Literature: A (Levin 1969), B (Levin an Schaal 1972), C (Levin 1966b), D (Levin 1963), E (Hadley Levin 1969), F (Levin an Schaal 1970), G (Levin and Smith 1966), H (Fehlberg et al. 2014), I (Smith and Levin 1967), J (Levin 1968), K (Levy and Levin 1975), L (Levy and Levin 1974).

drummondii and *P. cuspidata* but analyses were limited to single individuals (Roda et al. 2017).

Reexamining these hypotheses of reinforcement with gene flow using geographically widespread

sampling and extensive genomic sequencing will provide a clearer understanding of the

evolution of reproductive trait divergence in sympatry.

In this study, we apply double-digest RAD (ddRAD) sequencing with phylogenomic

approaches to resolve the evolutionary relationships of the eastern standing *Phlox*. We then

leverage this phylogenetic framework to examine the contribution of hybridization and gene flow

in the diversification of this system. Specifically, we test if gene flow has occurred alongside the

evolution of two cases of reinforcement, and investigate patterns of genetic composition, and

gene flow in multiple hypothesized cases of homoploid and allotetraploid hybrid speciation.

MATERIALS AND METHODS

Sampling, ddRAD Library Sequencing, and Data Processing

We collected wild plants from across the native ranges of the eastern standing *Phlox* in

North America during the summers of 2017-2019. We acquired additional samples from natural

populations preserved in tissue culture at the Ohio State University Ornamental Plant Germplasm

Center (OPGC). Our sampling includes seventeen major recognized eastern standing *Phlox*

species, with representation from thirty-two described subspecies, and three outgroup taxa from

the mat-forming *Phlox* (Fig. 1; Supplementary Table S1). We erred on the side of over-splitting

taxa as described in Locklear 2011a, except for the Texas annuals which we only designated at

species level.

Plants were grown in the greenhouse under controlled conditions until flowering. We

extracted genomic DNA from bud tissue using a hybrid Omega Bio-Tek EZNA DNA kit and

CTAB/Chloroform extraction protocol. Genomic libraries were constructed with a double-digest

restriction-site-associated digestion (ddRAD) sequencing protocol and paired-end sequenced as

described in (Goulet-Scott et al. 2021)**.**

Raw reads were demultiplexed, filtered, and *de novo* assembled as described in (Goulet-

Scott et al. 2021)**.** In brief, raw sequence data was demultiplexed and clone filtered using custom

python scripts and the 'process radtags' and 'clone filtering' functions in STACKS v.2.4

(Catchen et al. 2013)**.** Demultiplexed reads were further filtered for adapter sequences and

trimmed for base quality score using Trimmomatic v.0.39 (Bolger et al. 2014) under default

settings. Reads were *de novo* assembled into loci and variant called using iPyRAD v.0.9.50

(Eaton and Overcast 2020), with a consensus locus clustering threshold of 0.93. Samples with

minimal informative shared loci were removed, resulting in a final dataset of 116 ingroup and 4

outgroup individuals.

For the 120 retained samples, raw reads were filtered to an average of 3,060,421 reads

per individual. Filtered reads were used to construct four datasets with varying levels of

missingness, requiring a locus to be shared among a minimum of N = 4, 10, 20, and 30

individuals, named MS4, MS10, MS20, and MS30 respectively. These assembled datasets

produced matrices with a total of 165,549 to 5,3871 loci and 1,884,423 to 105,363 SNP sites

with a locus missingness rate of 90%, 83%, 72%, and 60% respectively.

Phylogenetic Inferences

We inferred the phylogenetic relationships of our samples using concatenation and

coalescent-based methods. First, whole ddRAD loci from the MS10, MS20 and MS30 datasets

were individually concatenated into supermatrices. Maximum likelihood (ML) phylogenies were

estimated from each supermatrix using IQ-TREE v.1.6.10 (Nguyen et al. 2015) with the GTR +

gamma nucleotide substitution model and 1000 bootstrap replicates using ultrafast bootstrap

approximation (Minh et al. 2013; Hoang et al. 2018).

Second, we inferred a population-level species tree using SVDquartets (Chifman and

Kubatko 2014), as implemented in PAUP v.4.0a166 (Swofford 2002). This method uses

unlinked SNPs to infer relationships among quartets of taxa under a coalescent model and then

estimates a species tree using a quartet assembly method. We conducted the analysis on our

dataset with least missingness, MS30, with the mat-forming species removed and *P. stolonifera*

set as the outgroup. To avoid genetic linkage, we sampled a single SNP-site with the least

amount of missing data from each locus, totaling 5,306 variant sites. We used all 7,160,245

possible quartets and 100 non-parametric bootstrap replicates to assess topological support and to

generate a majority-rule consensus species tree.

Support values between these two inference methods are derived from two different

estimation methods and cannot be compared one to one. Standard bootstrap supports (SBS), as

used with our SVDquartets inferred tree, are conservative and underestimate the likelihood of

relationships; SBS supports of 80% have a 95% likelihood of being correct, so SBS >= 80 are

considered trusted supports. However ultrafast bootstrap supports (UBS), as used for our

concatenation inference method, are unbiased when >70, so a 95% support has a 0.95 probability

of being correct (Minh et al. 2013).

Tests for Gene Flow Alongside Reinforcement

We tested for evidence of gene flow alongside the evolution of reinforcement in *P.*

drummondii and *P. pilosa* subsp. *pilosa*. First, we computed Patterson's D-statistic (Green et al.

2010; Durand et al. 2011) to infer if gene flow occurred between two lineages in a given

phylogeny. This test measures the asymmetry in the ratio of two discordant allele patterns,

ABBA and BABA, across the topology of a four-taxon pectinate tree ((P1, P2), P3), O). Under

stochastic lineage divergence without gene flow between lineages P3 and P1 or P3 and P2, the

proportion of discordant allele patterns from incomplete lineage sorting are unbiased

(ABBA=BABA); however, when introgression has occurred between P3 and P1 or P3 and P2,

one discordant allele pattern is more prevalent than the other (ABBA>BABA, BABA>ABBA).

We defined the P1, P2, P3, and O lineages as a taxa and a "test" as a unique combination

of individuals from these taxa. We subset the complete MS4 data matrix to include the

individuals from a set of four taxa that fit the pectinate topology under our ML phylogenetic

trees. P1 is the taxon that diverged by reinforcement (*P. drummondii* and *P. pilosa* subsp.

pilosa), P2 as a taxon out to P1 but not involved in the reinforcement hypothesis, P3 as the taxon

sympatric with P1 driving divergence by reinforcement (*P. cuspidata* and *P. glaberrima* subsp.

interior), and O as an individual from a proximal lineage out to all other lineages. Tests were

iterated over all possible combinations of all individuals of the P1 and P3 lineages, up to three

randomly sampled individuals of the P2 lineage not observed growing geographically near

populations of P1 or P3, and two individuals of the O lineage. The white flowered *P. pilosa*

subsp. *pilosa* lineage that evolved by reinforcement diverged from the lineage of northern *P.*

pilosa subsp. *pilosa* populations (Fig. 2), therefore, we only used *P. pilosa* subsp. *pilosa* North

individuals in these tests. We were also able to consider two different species as the P2 lineage in

the hypothesis concerning *P. pilosa* subsp. *pilosa* North. For all possible tests, we implemented

the D-statistic test for 1,000 bootstrap iterations with loci resampling with replacement, as

described in (Eaton et al. 2015). Tests with |Z-score| >3 are conservatively statistically

significant.

We complemented the D-statistic results with TreeMix v1.13 (Pickrell and Pritchard

2012), a model-based method for inferring population splits and migration (gene flow) under a

tree framework, as implemented in iPyRAD. We targeted a potential signal of introgression by

modeling a subtree of our total phylogenetic sampling. We subset the complete MS4 data matrix

to include all individuals from all non-hybrid lineages in the clades containing the P1 and P3

lineages, described above, all intermediate clades, and from a lineage out to this subtree.

Individuals were grouped based on species identity with *P. drummondii* and *P. pilosa* subsp.

pilosa North split based on flower color phenotype. We denote these subtrees as "*P. pilosa*

subsp. *pilosa* North" and "*P. drummondii*" respectively. We required SNPs to be shared by at

least 20% of individuals from each species and randomly sampled one unlinked SNP from each

locus. The *P. drummondii* (4,638 unlinked SNPs) and *P. pilosa* subsp. *pilosa* North (3,383

unlinked SNPs) subtrees were modeled in TreeMix with $m = 1$-8 possible migration edges. The

best supported number of migration edges was qualitatively chosen as the value at which the log-

likelihood of the model began to increase only marginally.

Testing Origins of Homoploid Hybrid Species

To evaluate if the five putative homoploid hybrid species in the standing *Phlox* (Table 1)

have genetic support we leveraged our phylogenetic inferences and tests for gene flow to

evaluate four criteria: 1) the putative hybrid sits sister to or nested within one of its putative

parental taxon in our bifurcating phylogenetic reconstructions, 2) the putative hybrid shows a

signal of gene flow with its other parental taxon, 3) the signal of gene flow is exclusive to the

putative hybrid, and 4) the signal of gene flow goes into the putative hybrid.

For putative hybrids sister to or within one of their putative parent lineages, we tested for

the presence of gene flow with the second putative parental lineage using Patterson's D-statistic,

as described above. For each hybrid hypothesis, a four-taxon tree was generated by subsetting

the MS4 dataset to all individuals of the putative hybrid species and putative second parent

species, three individuals from a species lineage between the putative hybrid and putative parent,

and two individuals of a closely related outgroup. The putative hybrid and the putative parent

lineages served as either P1 or P3 based on the structure of our ML phylogenies. We also

considered multiple species as the P2 lineage. For putative hybrid lineages with significant

evidence of gene flow with their parental taxon, we evaluated if the signature of gene flow was

localized to the hybrid lineage by testing for any gene flow between the two putative parental

taxa. All hybrid hypotheses tested with D-statistics were also modeled with TreeMix v.1.13, as

described above. The *P. maculata* subsp. *pyramidalis*, *P. amoena* subsp. *lighthipei*, and *P. pilosa*

subsp. *detonsa* modeled subtrees used 4,490, 4,294, and 1,033 unlinked SNPs respectively. We

used the *P. pilosa* subsp. *pilosa* North modeled subtree for the *P. argillacea* hypothesis given

their identical sampling.

Inferring Origins of Allotetraploid Hybrid Species

We investigated the hypothesized ancestries of the four allotetraploid hybrid species

(Table 1) using a consensus and comparative read alignment approach, similar to (Wang et al.

2021)**.** Sequenced representatives of these taxa were previously confirmed to be tetraploid via

flow cytometry at the OGPC.

First, we created reference consensus sequences from diploid lineages. Whole ddRAD

loci present in at least 60% of diploid taxa from the MS4 dataset were concatenated to generate

taxon-level consensus sequences using the iPyRAD window extractor toolkit, retaining the most

common allele in the consensus sequence for species with multiple samples. We then mapped

the filtered sequence reads for all allotetraploids to the diploid consensus sequences using bwa-

mem (Li 2013). We tallied the proportion of primary read alignments with a mapping quality ≥ 5

for the forward and reverse reads. We reasoned the diploid consensus receiving the highest

proportion of quality read alignments would be most closely related to the ancestor of the

mapped reads. We tested this assumption by also mapping one representative from each diploid

taxon to the consensus sequences and observing the highest rates of mapping to the same taxon

consensus with some carry over to closely related sister taxa. At a minimum, this method

allowed identification of the ancestral clade the reads originated from. Under this logic we

inferred the putative ancestry of the allopolyploid subgenomes to diploid clades that received the

highest rates of read mapping (Wang et al. 2021).

RESULTS

Phylogenomic Inferences of Species Evolutionary Relationships

Phylogenetic analyses of genome-wide nuclear ddRAD loci provided novel resolution of

species relationships and partly challenge the historic view of standing *Phlox* evolution. ML trees

inferred from each of three concatenated datasets were highly congruent, despite variation in

missing data (Fig. 2; Supplementary Fig. S1, S2). Therefore, we focused our results to trends

across phylogenetic inferences with discussion of notable differences in interspecific

relationships across datasets.

The base of the eastern standing *Phlox* is notably composed of long-styled taxa. We recover an

early split between *P. stolonifera* and all other taxa (UFB=100), followed by a split of a well-

supported clade (UFB=100) containing *P. ovata* out to *P. pulchra, P. glaberrima* subsp. *triflora,*

and the allotetraploid *P. buckleyi.* The remaining *P. glaberrima* subspecies (*P. glaberrima* subsp.

glaberrima and *P. glaberrima* subsp. *interior*) are found in a neighboring clade, with subspecies

of *P. carolina* and *P. maculata* (UFB=100). Across all inferences, the *P. glaberrima* and *P.*

carolina subspecies are not monophyletic within their species delineation. The next branch

subtends a well-supported small clade including the final long-styled species, *P. amplifolia* and

P. paniculata (UFB=100), with the placement of *P. amplifolia* within or sister to *P. paniculata*

varying across levels of missingness.

Across all datasets, our analyses inferred strong support for a single transition from long

to short-styled species, with all short-styled taxa forming a monophyletic group (UFB=100).

This short-styled group contains taxa from the *P. pilosa* complex, the *P. divaricata* and *P.*

amoena subspecies, and the three Texas annual species (*P. drummondii*, *P. roemeriana*, and *P.*

cuspidata). Half of our *P. pilosa* subspecies samples, including *P. pilosa* subsp. *fulgida*, *P. pilosa*

subsp. *pilosa*, and *P. pilosa* subsp. *deamii*, form a well-supported clade out to the remaining

short-style species (UFB=100). We refer to this group and the *P. pilosa* subsp. *pilosa* individuals

within it as "North". Notably, the white flower morph of *P. pilosa* subsp. *pilosa* that has evolved

by reinforcement, and sometimes described as the putative hybrid species *P. argillacea*, is found

in this clade, and *P. pilosa* subsp. *fulgida* is found nested within *P. pilosa* subsp. *pilosa* North.

Both *P. divaricata* and *P. amoena* species respectively form strongly supported monophyletic

groups (UFB=100), although *P. divaricata* subspecies are interdigitated and do not form

monophyletic groups.

Figure 2. Maximum likelihood phylogenetic inference of eastern standing *Phlox* from whole

concatenated ddRAD loci. Loci were required to be shared by at least 30 individuals. Numbers

on branches represent bootstrap supports from 100 bootstrap replicates using ultrafast bootstrap

approximation (UFB). Bootstrap supports are shown for all nodes differentiating between taxa.

The remaining *P. pilosa* subspecies, Texas annuals, and the allotetraploids *P. floridana,*

P. villosissima subsp. *villossisma* and *P. villosissima* subsp. *latisepala* form a large, well

supported clade (UFB=100). Along the backbone of this clade we observed reduced branch

supports; however, the three annual species, *P. cuspidata*, *P. drummondii*, and *P. roemeriana,*

are monophyletic (UFB=100), with *P. roemeriana* sister to *P. drummondii,* consistent with

previous genomic inferences (Roda et al. 2017)**.** Adjacent to the annuals lie the remaining *P.*

pilosa subspecies. We refer to this *P. pilosa* group as "South" because of the relative southern

origin of these samples to our other *P. pilosa* subspecies samples. This group contains *P. pilosa*

subsp. *pliosa*, *P. pilosa* subsp. *detonsa*, and *P. pilosa* subsp. *ozarkana.* Notably we also observed

P. pilosa subsp. *sangamonensis* is within this *P. pilosa* South group, despite being from well into

the *P. pilosa* North group's range in Illinois.

Low support for partitioning of the *P. pilosa* South group from the Texas annuals across

our datasets may be due to the presence of individuals from the two allotetraploid subspecies *P.*

villosissima subsp. *latisepala* and *P. villosissima* subsp. *villosissima.* Across datasets, these taxa

are inferred either out to the annuals or out to the *P. pilosa* South group with variable support

(UFB=41-100). Variation in their inferred placement between these groups is consistent with

their theorized ancestry as hybrids between populations of *P. pilosa* subsp. *pilosa* South and *P.*

drummondii (Table 1). Similar uncertainty can also be observed for the allotetraploid *P.*

floridana across datasets, observed either out or nested within the *P. pilosa* South group. This

may also be a result of its putative hybrid ancestry between populations of *P. pilosa* subsp. *pilosa*

South and *P. carolina* subsp. *angusta* (Table 1).

Coalescent-based species tree inference using SVDquartets recovered a species tree with

nearly identical relationships to our ML inferred trees (Supplementary Fig. S3); however, we did

observe differences in branch supports at some interspecific positions. We inferred strong

support for most nodes along the backbone of the tree (SBS > 80) and splits between major

species groupings, except for early splits differentiating relative placement within the long-style

species. *P. pulchra, P. glaberrima* subsp. *triflora,* and *P. buckleyi* remained monophyletic

(SBS=84) and separate from the remaining monophyletic *P. glaberrima*/*P.maculata*/*P.carolina*

subspecies (SBS=100). We also recovered the previously observed polyphyletic relationships of

the *P. pilosa* North and *P. pilosa* South groups. Unlike our ML analyses, our coalescent

inferences showed support for a cleaner partitioning of a monophyletic clade of *P. pilosa* South

and the allotetraploids *P. floridana*, *P. villosissima* subsp. *villosissima* and *P. villosissima* subsp.

latisepala, sister to the Texas Annuals (SBS=89).

Evidence of Gene Flow Alongside Reinforcement

We used D-statistics and TreeMix v.1.13 to test for a history of gene flow in two cases of

reinforcement in *Phlox*. Specifically, we leveraged our sampling to compare if a significant

signature of gene flow exists between the focal sympatric species in each case of reinforcement.

All D-statistic tests were structured so a significant negative D supports a history of introgression

between these focal species.

In *P. drummondii*, reinforcement against hybridization with *P. cuspidata* has favored the

divergence of flower color from light-blue to dark-red in sympatry (Fig. 3a). With D-statistic

tests for gene flow between *P. drummondii* and *P. cuspidata*, we found no evidence of

introgression between light-blue allopatric populations (D values were negative but not

significant); However, tests including dark-red *P. drummondii* individuals sympatric with *P.*

cuspidata (N=48), all tests had a negative D value and 24 out of 48 tests showed a significant

excess of BABA patterns with a |Z-score| > 3 (Table 2). 18 of these significant tests used

386
387
Figure 3. (a and b) Schematics of flower color divergence due to reinforcement in two species

of *Phlox* to prevent maladaptive hybridization with another species in sympatry. (a) *P.*

drummondii (red) has similar light-blue colored flowers to *P. cuspidata* (pink) in allopatry but *P.*

drummondii has evolved dark-red flowers in sympatry to reduce hybridizing with *P. cuspidata*.

(b) In allopatry *P. pilosa* subsp. *pilosa* North (light blue) has similar pink colored flowers to *P.*

glaberrima subsp. *interior* (dark blue), but where they co-occur in high frequency in sympatry,

P. pilosa subsp. *pilosa* North has evolved white flowers to reduce hybridizing with *P.*

glaberrima subsp. *interior*. (c and d) Best fit models from TreeMix inferred gene flow between

P. cuspidata and dark-red *P. drummondii* (c) (best fit m=2) but did not support evidence of gene

flow between *P. glaberrima* subsp. *interior* and either color morph of *P. pilosa* subsp. *pilosa*

North (d) (best fit m=1).

Table 2. Summary results for four taxon D-statistic tests for introgression in two cases of reinforcement

Reinforcement Scenario	P1	P2	P3	Range D	Range $Z^\dagger$	nSig/N‡ -D	nSig/N‡ +D	Test ID
P. drummmondii								
	P. drummmondii (light-blue)	*P. roemariana*	*P. cuspidata*	(-0.12, -0.01)	(0.187, 2.479)	0/24	0/24	T2:1 - T2:24
	P. drummmondii (dark-red)	*P. roemariana*	*P. cuspidata*	(-0.16, -0.01)	(0.341, **4.692**)	**24/48**	0/48	T2:25 - T2:72
P. pilosa subsp. *pilosa* North								
	P. pilosa subsp. *pilosa* North (pink)	*P. divaricata* subsp. *divaricata*	*P. glaberrima* subsp. *interior*	(-0.10, 0.08)	(0.008, 2.771)	0/180	0/180	T2:73 - T2:252
	P. pilosa subsp. *pilosa* North (white)	*P. divaricata* subsp. *divaricata*	*P. glaberrima* subsp. *interior*	(-0.08, 0.09)	(0.016, 2.418)	0/90	0/90	T2:253 - T2:342
	P. pilosa subsp. *pilosa* North (pink)	*P. paniculata*	*P. glaberrima* subsp. *interior*	(-0.09, 0.05)	(0.025, 2.707)	0/180	0/180	T2:343 - T2:522
	P. pilosa subsp. *pilosa* North (white)	*P. paniculata*	*P. glaberrima* subsp. *interior*	(-0.09, 0.10)	(0.026, 2.566)	0/90	0/90	T2:523 - T2:612

P1, P2, P3 refer to the three lineages used for the ABBA-BABA D statistic test (See Methods). Outgroups not shown. All tests can be found in Table S2.

† Z-score to assess significance of whether D deviates from zero. Bold indicates significance at $\alpha=0.01$.

‡ Number of significant tests over all possible sampled individuals. Significant -D identifies gene flow between P1 and P3, while significant +D identifies gene flow between P2 and P3.

topologies with a *P. cuspidata* individual sampled from sympatry. Our best fit model (m=2) of

the *P. drummondii* subtree in TreeMix recovered *P. cuspidata* out to *P. drummondii* and *P.*

roemeriana with a migration event going from *P. cuspidata* into dark-red sympatric *P.*

drummondii (Fig. 3c). This migration edge remained present for all models m ≥2

(Supplementary Fig. S4, S9). These results support the hypothesis of gene flow occurring

between sympatric *P. drummondii* and *P. cuspidata* but do not show evidence of gene flow

between allopatric *P. drummondii* and *P. cuspidata*.

In *P. pilosa* subsp. *pilosa*, reinforcement against hybridization with *P. glaberrima* subsp.

interior has favored the divergence of *P. pilosa* subsp. *pilosa* flower color from pink to white

(Fig. 3b). Our phylogenetic analyses demonstrate the white *P. pilosa* subsp. *pilosa* populations

are monophyletic and nested within a group of pink *P. pilosa* subsp. *pilosa* in the *P. pilosa* North

group and polyphyletic from the rest of *P. pilosa* subsp. *pilosa* South (Fig. 2). This discovery

changes the context of this evolutionary scenario, with *P. pilosa* subsp. *pilosa* North more

closely related with *P. glaberrima* subsp. *interior* than previously thought.

Unlike reinforcement in *P. drummondii*, the ranges of *P. pilosa* subsp. *pilosa* North and

P. glaberrima subsp. *interior* are mosaic and cannot be distinctly divided as geographically

allopatric or sympatric (Levin and Kerster 1967). Therefore, we grouped our tests for

introgression by flower color phenotype of the *P. pilosa* subsp. *pilosa* North individuals, the

ancestral pink flower color and the derived white flower color that evolved by reinforcement,

instead of by geography. Our analyses show both an inconsistency in the direction D deviated

from zero and a lack of significant signature of gene flow between *P. glaberrima interior* with

pink flowered (N=360) and white flowered (N=180) colored *P. pilosa* subsp. *pilosa* North (Table

2). All TreeMix models of the *P. pilosa* subsp. *pilosa* North subtree recovered similar topologies

to our ML phylogenetic inferences but no migration events involving pink or white *P. pilosa*

subsp. *pilosa* North (Fig. 3d; Supplementary Fig. S5, S9).

Evidence of Divergence and Gene Flow Underlying Hypotheses of Putative Homoploid Hybrid

Species

The eastern standing *Phlox* have five hypothesized cases of homoploid hybrid speciation

(Table 1). We leveraged our phylogenetic reconstructions and tests for gene flow to ask if these

hypotheses are supported with phylogenetic evidence; specifically, if putative hybrid species sit

nested within or sister to one of its hypothesized parental species and demonstrating a signature

of gene flow, and therefore hybridization, with its other putative parental species.

Our ML and coalescent based phylogenetic inferences show support for four of the five

putative hybrids sitting nested or within one of their putative parents (Fig. 2, Supplementary Fig.

S1, S2, S3); *P. amoena* subsp. *lighthipei* within *P. amoena* subsp. *amoena*, *P. maculata* subsp.

pyramidalis within *P. maculata* subsp. *maculata*, *P. argillacea* within *P. pilosa* subsp. *pilosa*

North*,* and *P. pilosa* subsp. *detonsa* in *P. pilosa* subsp. *pilosa* South*. P. pilosa* subsp. *deamii* was

not found out or within *P. pilosa* subsp. *pilosa* South or *P. amoena* subsp. *amoena*, consistent

with (Goulet-Scott et al. 2021)*.*

D-statistic tests for gene flow did not find support for gene flow between three of the

putative hybrid species and their putative second parental species, *P. amoena* subsp. *lighthipei*

with *P. pilosa* subsp. *pilosa* South (N=624), *P. argillacea* with *P. glaberrima* subsp. *interior*

(N=270), and *P. pilosa* subsp. *detonsa* with *P. carolina* subspecies (N=90*)* (Table 3). Our

TreeMix analyses generally recapitulated similar topologies of the ML phylogenetic inferences

but found no support for gene flow underlying these hypotheses (Supplementary Fig. S5, S6, S7,

S9).

Table 3. Summary results for four taxon D-statistic tests for introgression in four hypothesized cases of hybrid speciation

Hypothesized Hybrid	P1	P2	P3	Range D	Range $Z^{\dagger}$	nSig/$N^{\ddagger}$ -D	nSig/$N^{\ddagger}$ +D	Test ID
P. argillacea								
	P. argillacea	*P. pilosa* subsp. *pilosa* North	*P. glaberrima* subsp. *interior*	(-0.09, 0.08)	(0.023, 2.115)	0/90	0/90	T3:1 - T3:90
	P. argillacea	*P. divaricata* subsp. *divaricata*	*P. glaberrima* subsp. *interior*	(-0.08, 0.09)	(0.016, 2.418)	0/90	0/90	T3:91 - T3:180
	P. argillacea	*P. paniculata*	*P. glaberrima* subsp. *interior*	(-0.09, 0.10)	(0.026, 2.566)	0/90	0/90	T3:181 - T3:270
P. pilosa subsp. *detonsa*								
	P. pilosa subsp. *detonsa*	*P. pilosa* subsp. *pilosa* South	*P. carolina* subsp. *angusta*	(-0.14, 0.01)	(0.033, 2.383)	0/12	0/12	T3:271 - T3:282
	P. pilosa subsp. *detonsa*	*P. pilosa* subsp. *pilosa* South	*P. carolina* subsp. *alta*	(-0.09, 0.02)	(0.042, 1.520)	0/12	0/12	T3:283 - T3:294
	P. pilosa subsp. *detonsa*	*P. pilosa* subsp. *pilosa* South	*P. carolina* subsp. *carolina*	(-0.13, 0.06)	(0.207, 1.794)	0/6	0/6	T3:295 - T3:300
	P. pilosa subsp. *detonsa*	*P. drummondii*	*P. carolina* subsp. *angusta*	(-0.08, -0.01)	(0.223, 1.880)	0/12	0/12	T3:301 - T3:312
	P. pilosa subsp. *detonsa*	*P. drummondii*	*P. carolina* subsp. *alta*	(-0.05, 0.05)	(0.164, 1.238)	0/12	0/12	T3:313 - T3:324
	P. pilosa subsp. *detonsa*	*P. drummondii*	*P. carolina* subsp. *carolina*	(-0.04, 0.06)	(0.024, 1.147)	0/6	0/6	T3:325 - T3:330
	P. pilosa subsp. *detonsa*	*P. roemeriana*	*P. carolina* subsp. *angusta*	(-0.05, 0.06)	(0.055, 1.266)	0/12	0/12	T3:331 - T3:342
	P. pilosa subsp. *detonsa*	*P. roemeriana*	*P. carolina* subsp. *alta*	(-0.05, 0.13)	(0.024, 2.696)	0/12	0/12	T3:343 - T3:354
	P. pilosa subsp. *detonsa*	*P. roemeriana*	*P. carolina* subsp. *carolina*	(-0.05, 0.02)	(0.188, 1.102)	0/6	0/6	T3:355 - T3:360
P. amoena subsp. *lighthipei*								
	P. amoena subsp. *lighthipei*	*P. amoena amoena*	*P. pilosa* subsp. *pilosa* South	(-0.14, 0.08)	(0.005, 2.887)	0/234	0/234	T3:361 - T3:594
	P. pilosa subsp. *pilosa* South	*P. drummondii*	*P. amoena* subsp. *lighthipei*	(-0.13, 0.06)	(0.007, 2.631)	0/234	0/234	T3:595 - T3:828
	P. pilosa subsp. *pilosa* South	*P. cuspidata*	*P. amoena* subsp. *lighthipei*	(-0.10, 0.13)	(0.007, 2.559)	0/156	0/156	T3:829 - T3:984
P. maculata subsp. *pyramidalis*								
	P. maculata subsp. *pyramidalis*	*P. maculata* subsp. *maculata*	*P. glaberrima* subsp. *interior*	(-0.50, 0.08)	(1.869, **20.718**)	**27**/30	0/30	T3:985 - T3:1014
	P. glaberrima subsp. *interior*	*P. carolina* subsp. *alta*	*P. maculata* subsp. *pyramidalis*	(-0.34, -0.06)	(2.002, **13.515**)	**17**/20	0/20	T3:1015 - T3:1034
	P. glaberrima subsp. *interior*	*P. glaberrima* subsp. *glaberrima*	*P. maculata* subsp. *pyramidalis*	(-0.31, -0.06)	(2.167, **12.277**)	**19**/20	0/20	T3:1035 - T3:1054
	P. glaberrima subsp. *interior*	*P. carolina* subsp. *alta*	*P. maculata* subsp. *maculata*	(0.01, 0.20)	(0.299, **6.323**)	0/80	**62**/80	T3:1055 - T3:1134
	P. glaberrima subsp. *interior*	*P. glaberrima* subsp. *glaberrima*	*P. maculata* subsp. *maculata*	(-0.02, 0.19)	(0.090, **6.313**)	0/80	**59**/80	T3:1135 - T3:1214

P1, P2, P3 refer to the three lineages used for the ABBA-BABA D statistic test (See Methods). Outgroups not shown. All tests can be referenced by Test ID in Table S3.
† Z-score to assess significance of whether D deviates from zero. Bold indicates significance at α=0.01.
‡ Number of significant tests over all possible sampled individuals. Significant -D identifies gene flow between P1 and P3, while significant +D identifies gene flow between P2 and

449
450

Figure 4. Results from TreeMix support the hybrid speciation hypothesis for *P. maculata*

pyramidalis in a pruned subtree (best fit m=4). Modeling of 4,490 unlinked SNPs for individuals

clustered by taxon identity inferred *P. maculata* subsp. *pyramidalis* sister to *P. maculata* subsp.

maculata and receiving a migration event (gene flow) from *P. glaberrima* subsp. *interior*.

However, our D-statistic tests did identify strong support for gene flow between *P. glaberrima*

subsp. *interior* and *P. maculata* subsp. *pyramidalis,* with significantly negative values in 63 of 70

tests. We found no evidence of gene flow between *P. glaberrima* subsp. *interior* and *P. maculta*

subsp. *maculta,* N=160 (Table 3). All models of the pyramidalis subtree in TreeMix (best fit

m=4) confirmed this result, with a migration edge from *P. glaberrima* subsp. *interior* into *P.*

maculta subsp. *pyramidalis* (Fig. 4, Supplementary Fig. S8, S9). These results are

consistent with *P. maculata* subsp. *pyramidalis* as a hybrid lineage arising from *P. maculata*

subsp. *maculata* and *P. glaberrima* subsp. *interior* parental lineages.

Inferring Origins of Allotetraploid Hybrid Species

Hybridization of divergent diploid genomes formed four, reproductively isolated,

tetraploid *Phlox* species (Table 1) (Levin 1966a, 1968; Smith and Levin 1967). *P. villosissima*

subsp. *villosissima* and *P. villosissima* subsp. *latisepala* are thought to be independent polyploid

hybridization events between *P. pilosa* subsp. *pilosa* South and *P. drummondii*. *P. floridana* is

hypothesized to be the product of *P. pilosa* subsp. *pilosa* South and *P. carolina* subsp. *angusta***,**

and finally*, P. buckleyi* contains two discordant diploid karyotypes not observed elsewhere and

the ancestors are hypothesized to be extinct (Wherry 1955)**.** We leveraged comparative read

mapping to infer subgenome ancestry of these taxa and evaluate their hypothesized origins.

We concatenated ddRAD loci to generate reduced consensus sequences for each of the 28

diploid species. Each consensus was 439,271 bp long with a locus missingness rate of 25%. For

one test representative of each diploid species, an average of 37,948 reads mapped with high

quality to the consensus multi-sequence FASTA file. The distribution of mapping rates across

consensus sequences demonstrated a clear ability to assign ancestry of diploid genomes, with the

consensus receiving the highest relative proportion of mapped reads corresponding to the known

identity of the diploid individual (Supplementary Table S4).

We repeated this mapping for each allotetraploid, with an average of 17,737 high-quality

mapped reads per polyploid individual. We observed a more diffuse read mapping rate across

consensus sequences relative to the test diploid taxa, yet clear patterns of elevated read mapping

constrained to specific subclades (Fig. 5). Reads of *P. villosissima* subsp. *vollosissima* and *P.*

villosissima subsp. *latisepala* mapped primarily to the *P. pilosa* South group and the Texas

annuals, with each receiving 30-34% and 22-26% of the mapped reads respectively. Within these

two groups the top recipients of mapped reads were *P. pilosa* subsp. *pilosa* South and *P.*

cuspidata for all three allotetraploid individuals. *P. floridana* showed a similar high mapping rate

to the *P. pilosa* South group (34%), with most reads mapping to *P. pilosa* subsp. *pilosa* South

and second most to *P. pilosa* subsp. *detonsa*. Contrary to the hypothesized origins of this species,

Figure 5. Proportion of reads mapped to diploid species-level consensus sequences for a diploid

representative of *P. drummondii* and all allotetraploid hybrid individuals. Bar colors correspond

to diploid species groups in Figure 2.

we did not observe elevated read mapping to the *P. carolina subspecies* consensus sequence.

Instead, we observed elevated read mapping to *P. amoena* subsp. *amoena*. Finally, *P. buckleyi*

showed high rates of read mapping to taxa within the deeper branch of the eastern standing

Phlox, with highest rate of mapping to the *P. ovata*, *P. pulchra*, *P. glaberrima* subsp. *triflora*

clade (38%), with highest recipients being *P. pulchra* and *P. ovata.*

DISCUSSION

Phylogenomic inferences of species relationships and explicit tests for interlineage gene

flow provide a systematic way for inferring the presence, extent, and context of hybridization

and gene flow across a phylogeny. Here we present well-resolved phylogenetic relationships of

the eastern standing *Phlox*. We demonstrate clear support for most described species

relationships and reveal novel non-monophyletic relationships of subspecies in the historically

taxonomically difficult *P. pilosa* and *P. glaberrima*/ *P. carolina*/ *P. maculata* species complexes.

Using this phylogenetic framework, we find support for one case of hypothesized homoploid

hybrid speciation event, identify the putative ancestries of multiple polyploid species, and find

evidence of gene flow in one of two cases of reinforcement. Our findings demonstrate the utility

and importance of phylogenomics in confirming hypothesized evolutionary histories of non-

model systems and add to the growing evidence that gene flow across species boundaries can

play a role in the generation of novel biodiversity.

Evolutionary Relationships of the Eastern Standing Phlox

For nearly one hundred years, the eastern standing *Phlox* have been a foundational

system for understanding plant evolution, yet despite previous phylogenetic study on this clade

(Ferguson et al. 1999; Ferguson and Jansen 2002; Roda et al. 2017; Landis et al. 2018; Goulet-

Scott et al. 2021), the evolutionary relationships of these taxa have remained unclear. Our

phylogenomic inferences on genome-wide ddRAD sequence data clarifies the evolutionary

relationships of the taxa in this group and provides insight into their diversification.

Historic treatments of the eastern standing *Phlox* have grouped the species by

conspicuous differences in style length (Wherry 1930, 1931, 1932, 1955). Unlike prior

phylogenetic studies, we observe support for a large monophyletic clade of short-styled taxa

subtended by multiple paraphyletic clades of long-styled taxa (Fig. 2). Divergence in style-length

can generate reproductive isolation (Kay 2006; Brothers and Delph 2017), and this transition

from long to short style-length may have reduced competition and stimulated the diversification

of the short-styled clade.

Our phylogenetic analyses recapitulate previously inferred relationship of the Texas

Annuals species, as observed (Roda et al. 2017) but not in other studies (Ferguson et al. 1999;

Ferguson and Jansen 2002; Landis et al. 2018). We also find clear monophyletic support for the

P. amoena and *P. divaricata* species groups.

In the taxonomically more difficult *P. glaberrima*/ *P. carolina*/ *P. maculata* complex, we

find support for non-monophyletic relationships among subspecies (Fig. 2). Notably, *P.*

glaberrima triflora is phylogenetically distinct from the rest of the interdigitated taxa of this

species complex. Additionally, the subspecies of the *P. pilosa* complex form two distinct

polyphyletic groups roughly coinciding with their relative northern or southern location of

origin, as suggested in (Goulet-Scott et al. 2021). The one exception is *P. pilosa* subsp.

sangamonensis which genetically clusters with the southern clade but is found in the north (Fig.

1); A discovery consistent with the allopatric speciation hypothesis of *P. pilosa* subsp.

sangamonensis arising by long-distance dispersal from southern populations of *P. pilosa* subsp.

pilosa (Levin and Smith 1965; Levin 1984). The non-monophyletic relationship among the

northern and southern *P. pilosa* taxa and within *P. glaberrima* were suggested in previous

phylogenetic studies, but remained uncertain due to low phylogenetic supports and reliance on

single genetic loci (Ferguson et al. 1999; Ferguson and Jansen 2002; Landis et al. 2018).

Taxonomic delineations within *Phlox* are largely based on morphological characteristics

(Wherry 1955; Locklear 2011a), yet our phylogenetic inferences demonstrate current taxonomy

may not best reflect the true evolutionary relationships of some taxa within the eastern standing

Phlox group. Discovery of these new phylogenetic relationships also reframe the phylogenetic

context of the historic discussions on hybridization, gene flow, reinforcement, and hybrid

speciation within this system, as discussed below.

Evolution of Reinforcement With and Without Gene Flow

Interspecific hybridization can play an important role in speciation by generating

selection for reinforcement. However, limited empirical study has evaluated if this hybridization

also resulted in reinforcement evolving in the face of gene flow (Garner et al. 2018). We applied

D-statistics and TreeMix to assess if a history of gene flow coincides with the evolution of

flower color divergence by reinforcement in *P. drummondii* and *P. pilosa* subsp. *pilosa* North

(Fig. 3).

Divergence from light-blue to dark-red colored flowers in *P. drummondii* has evolved to

prevent hybridization with *P. cuspidata* (Levin 1985; Hopkins and Rausher 2012)*.* Our

geographically widespread sampling of these species revealed evidence of introgression between

P. cuspidata and multiple dark-red *P. drummondii* individuals from across the range of sympatry

but not between *P. cuspidata* and any allopatric light-blue *P. drummondii.* The significant

signature of gene flow in sympatry but not allopatry mirrors the classic evolutionary model of

divergence due to reinforcement to reduce hybridization in sympatry but not in allopatry (Garner

et al. 2018). This pattern also suggests the signal of gene flow is the product of hybridization in

sympatry and not due to ancient introgression between the *P. cuspidata* and *P. drummondii*

lineages. Our results corroborate previous inferences from transcriptome sequencing of single

individuals of each species (Roda et al. 2017)**.** Although F1 hybrids are highly sterile and have

reduced ability to backcross in the lab (Suni and Hopkins 2018), our results demonstrate hybrids

do backcross in nature, resulting in gene flow between the species. Reinforcement for increased

reproductive isolation in *P. drummondii* evolved despite gene flow in sympatry.

Conversely, our analyses do not support a history of gene flow coinciding with

hybridization and reinforcement between *P. pilosa* subsp. *pilosa* North and *P. glaberrima* subsp.

interior. These two species overlap for most of their ranges but co-flower in higher frequency at

the lower tip of lake Michigan. In this zone of elevated contact, *P. pilosa* subsp. *pilosa* North has

evolved from pink to white colored flowers, reducing the formation of hybrid seed by half in

sympatry, making it more adapted to coexist with *P. glaberrima* subsp. *interior* (Levin 1966a;

Levin and Kerster 1967; Levin and Schaal 1970b, 1970a, 1972)**.** Phylogenetic distance (Fig. 2)

and barriers to reproduction between these taxa are higher than between *P. cuspidata* and *P.*

drummondii, but experimental crosses demonstrate these species can generate hybrid seed, with

P. pilosa subsp. *pilosa* North as the maternal parent (Levin 1966a). Our results do not detect

gene flow between *P. pilosa* subsp. *pilosa* North and *P. glaberrima* subsp. *interior* regardless of

flower color (Fig. 3; Table 2). This lack of gene flow suggests hybrid offspring between these

taxa may be completely inviable, lethally maladapted, or have exceedingly high sterility.

The absence and presence of gene flow in these two case studies of reinforcement yield a

powerful comparative opportunity to investigate the dynamics by which reproductive trait

divergence, and flower color specifically, evolves by selection. For reproductive trait divergence

to evolve by reinforcement, alleles conferring increased assortative mating within a species must

remain associated with alleles causing costly hybridization between species (Servedio and

Kirkpatrick 1997; Kirkpatrick 2000). When gene flow between species is present, recombination

can disassociate these alleles, impeding successful divergence. Yet, we find the same trait

evolving due to similar types of selection in two *Phlox* species despite gene flow occurring

alongside one case but not the other. This result may be the consequence of differing genetic

architectures and genetic linkage underlying the cost of hybridization and flower color between

these cases. For example, strong genetic linkage between loci causing postzygotic

incompatibility and flower color divergence can reduce recombination between these traits and

allow buildup of linkage disequilibrium and the successful evolution of stronger reproductive

isolation in sympatry. Further investigation of the genetic architecture underlying reproductive

isolating barriers across these *Phlox* species will better inform the role of trait architecture in the

evolution of reinforcement.

Varying Support for Hypothesized Homoploid and Allotetraploid Hybrid Species

Within the eastern standing *Phlox* there are five hypothesized homoploid hybrid species

and four hypothesized allopolyploid species (Table 1). We only observed genomic support for *P.*

maculata subsp. *pyramidalis* being a homoploid hybrid species between *P. maculata* subsp.

maculata and *P. glaberrima* subsp. *interior. P. maculata* subsp. *pyramidalis* is found

phylogenetically closely related to multiple *P. maculata* subsp. *maculata* individuals (Fig. 2)

with a strong signal of receiving gene flow from *P. glaberrima* subsp. *interior* (Fig. 4; Table

3.). We can conclude our sampled *P. maculata* subsp. *pyramidalis* is of hybrid origin, however

future work will be necessary to discern if the hybrid origin of *P. maculata* subsp. *pyramidalis*

actually gave rise to reproductive isolation of its lineage from its putative parental species

(Schumer et al. 2018).

The remaining four hypothesized cases of homoploid hybrid speciation were not

supported by our genomic investigation, despite extensive phenotypic and biomolecular

evidence. In the cases of *P. amoena* subsp. *lighthipei*, *P. argillacea,* and *P. pilosa* subsp. *detonsa*

the lineages thought to be hybrid in origin may simply be divergent geographically isolated

populations from one of their hypothesized parental lineages. Yet in the case of *P. pilosa* subsp.

deamii, the lineage is distantly related to both hypothesized parents and may possesses

phenotypic traits resembling these species through convergence or incomplete lineage sorting.

Polyploid hybrid lineages can be significantly easier to identify than homoploid hybrids

through chromosome structure and counts. Despite this advantage, inferring the exact

evolutionary origin of these lineages remains challenging (Rothfels 2021). Although the signals

from our comparative read mapping are coarse, our approach suggests the identity of the

progenitor clades that gave rise to the polyploid hybrid species in this group. We provide further

support that *P. villosissima* subsp. *latisepala* and *P. villosisima* subsp. *villosissima* are derived

from *P. pilosa* (*P. pilosa* subsp. *pilosa* South) and a Texas annual species. *P. drummondii* was

previously hypothesized as the parent but our analyses cannot resolve which of the Texas annual

species or ancestor is the parent. As previously hypothesized, we find support for one parent of

P. floridana to be *P. pilosa* subsp. *pilosa* South, yet we find evidence that *P. amoena* subsp.

amoena may be the second parent instead of the *P. carolina subspecies*. We also observe *P.*

buckleyi is genetically similar to the early-diverging long-styled *Phlox*, but we cannot confirm

the specific parental lineages from this group. Future work using longer haplotype phase may

help resolve the evolutionary origins of these species and better inform the dynamics and

evolutionary trajectory of these allopolyploid species.

Across both the hypothesized homoploid and polyploid hybrid lineages, the *P. pilosa*

group has stood out as a potential and realized hotspot of hybridization activity. Four of the five

hypothesized homoploid hybrid lineages and three of the four polyploid lineages were thought to

include *P. pilosa* subsp. *pilosa* as a parent. While our analyses indicate that none of the

hypothesized homoploid hybrid lineages have hybrid origin with *P. pilosa* subspecies, three of

the four hypothesized polyploid hybrid lineages are derived from *P. pilosa* subsp. *pilosa* South

subspecies ancestry.

What has made *P. pilosa* prime for a role in these hypotheses? Notably the high

occurrence of *P. pilosa* subspecies across their composite broad range has not only led to them

being found in sympatry to many species, but potentially also as indicators of locally adapted

phenotypes across these environments. *P. pilosa*'s developmental biology may also make it

prone to evolutionary innovation, with many populations observed having high rates of

unreduced gametes and whole genome duplications across their range. Future work may help

address what factors underly this developmental instability and if it has an outsized role in the

generation of novel evolutionary lineages relative to other *Phlox* species.

Implications on Evolutionary History of Trait Variation

Now that we have a strong phylogenetic backbone for the evolutionary relationships of

taxa in the eastern standing *Phlox,* we can begin to untangle what forces have contributed to

patterns of phenotypic divergence throughout the clade. Much of the data supporting the role of

hybridization in speciation in this group stemmed from trait variation between distantly related

lineages. Our findings support hybridization has likely contributed to some of the phenotypic

variation observed across taxa in these cases, both as a source of genetic variation and force for

selective divergence. Further study of these confirmed cases may inform how specific traits and

even genes move between species and contribute to reproductive isolation between lineages.

Our findings also indicate that forces other than hybridization may be responsible for

convergent and recombinant patterns of trait variation across *Phlox*. Our lack of support for

many hypothesized cases of hybridization motivates future investigation of what did give rise to

the patterns of trait variation across and within these lineages. Further research into these taxa

and their phenotypic variation may provide new insights into the importance of convergent

evolution and/or incomplete lineage sorting to local adaptation within a species.

CONCLUSION

Hybridization between species can be an important force shaping the patterns of

evolution across clades. For example, decades of research have suggested that hybridization and

gene flow are extensive across the eastern standing *Phlox* and have directly played a role in their

speciation. With the accessibility of genomic data and improved analyses, we are now able to

test these hypotheses. We have built a robust phylogenetic framework using genome-wide

markers to better understand the relationship between these well-studied species and document

the occurrence of gene flow. We have determined hybridization has served as an important

creative force in the generation of diversity in this clade, however it may not be as prevalent as

previously thought. We do find evidence suggesting hybrid speciation and gene flow during

reinforcement, but we also find a lack of support for many hypothesized cases of gene flow in

the formation of evolutionary novel lineages. These lineages may just be the product of trait

divergence within a species separate from interspecific genetic exchange. Our work suggests a

strong need to reevaluate our previous evidence for the existence of gene flow, and to rethink our

expectations of how traits diverge within and between closely related species.

FUNDING

This study was supported by the National Science Foundation (DEB-1844906 to R.H.,

and Graduate Research Fellowship to A.G.G.); the Harvard Graduate School of Arts and

Sciences' Herchel Smith Graduate Research Fellowship in Science (to A.G.G. and B.E.G.S.);

and the Harvard University Herbaria Expedition Grant (to A.G.G.).

ACKNOWLEDGEMENTS

We are grateful to C.J. Ferguson, S.V. Edwards, J.S. Suissa, and the Hopkins Lab for

their helpful comments on this manuscript. We are also thankful to P.S. Jourdan and the OPGC

staff for sharing their *Phlox* collections with us.

CONTRIBUTIONS

A.G.G. and R.H. conceptualized the project. A.G.G., B.E.G.S., and R.H. collected and

analyzed the data. A.G.G. and R.H. wrote the manuscript.

DATA ACCESSIBILITY

Supporting data underlying this article is found on Dryad Digital Repository: [insert rep
ID}
701

REFERENCES

Abbott R., Albach D., Ansell S., Arntzen J.W., Baird S.J.E., Bierne N., Boughman J., Brelsford

A., Buerkle C.A., Buggs R., Butlin R.K., Dieckmann U., Eroukhmanoff F., Grill A.,

Cahan S.H., Hermansen J.S., Hewitt G., Hudson A.G., Jiggins C., Jones J., Keller B.,

Marczewski T., Mallet J., Martinez-Rodriguez P., Möst M., Mullen S., Nichols R., Nolte

A.W., Parisod C., Pfennig K., Rice A.M., Ritchie M.G., Seifert B., Smadja C.M.,

Stelkens R., Szymura J.M., Väinölä R., Wolf J.B.W., Zinner D. 2013. Hybridization and

speciation. J. Evol. Biol. 26:229–246.

Anderson E. 1949. Introgressive hybridization. Introgressive hybridization.

Anderson E., Gage A. 1952. Introgressive hybridization in phlox bifida. Am. J. Bot. 39:399–404.

Andrews K.R., Good J.M., Miller M.R., Luikart G., Hohenlohe P.A. 2016. Harnessing the power

of RADseq for ecological and evolutionary genomics. Nat. Rev. Genet. 17:81–92.

Barker M.S., Arrigo N., Baniaga A.E., Li Z., Levin D.A. 2016. On the relative abundance of

autopolyploids and allopolyploids. New Phytol. 210:391–398.

Bolger A.M., Lohse M., Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics. 30:2114–2120.

Bombonato J.R., do Amaral D.T., Silva G.A.R., Khan G., Moraes E.M., da Silva Andrade S.C.,

Eaton D.A.R., Alonso D.P., Ribolla P.E.M., Taylor N., Zappi D., Franco F.F. 2020. The

potential of genome-wide RAD sequences for resolving rapid radiations: a case study in

Cactaceae. Mol. Phylogenet. Evol. 151:106896.

724 Brothers A.N., Delph L.F. 2017. Divergence in style length and pollen size leads to a

725   postmating-prezygotic reproductive barrier among populations of Silene latifolia.

726   Evolution. 71:1532–1540.

727 Catchen J., Hohenlohe P.A., Bassham S., Amores A., Cresko W.A. 2013. Stacks: an analysis tool

728   set for population genomics. Mol. Ecol. 22:3124–3140.

729 Chifman J., Kubatko L. 2014. Quartet inference from SNP data under the coalescent model.

730   Bioinformatics. 30:3317–3324.

731 Durand E.Y., Patterson N., Reich D., Slatkin M. 2011. Testing for ancient admixture between

732   closely related populations. Mol. Biol. Evol. 28:2239–2252.

733 Dyer K.A., Bewick E.R., White B.E., Bray M.J., Humphreys D.P. 2018. Fine-scale geographic

734   patterns of gene flow and reproductive character displacement in Drosophila subquinaria

735   and Drosophila recens. Mol. Ecol.

736 Eaton D.A.R., Hipp A.L., González-Rodríguez A., Cavender-Bares J. 2015. Historical

737   introgression among the American live oaks and the comparative nature of tests for

738   introgression. Evolution. 69:2587–2601.

739 Eaton D.A.R., Overcast I. 2020. ipyrad: Interactive assembly and analysis of RADseq datasets.

740   Bioinformatics. 36:2592–2594.

741 Eaton D.A.R., Ree R.H. 2013. Inferring phylogeny and introgression using RADseq data: an

742   example from flowering plants (Pedicularis: Orobanchaceae). Syst. Biol. 62:689–706.

Erbe L., Turner B.L. 1962. A biosystematic study of the phlox cuspidata-phlox drummondii

complex. Am. Midl. Nat. 67:257.

Excoffier L., Dupanloup I., Huerta-Sánchez E., Sousa V.C., Foll M. 2013. Robust demographic

inference from genomic and SNP data. PLoS Genet. 9:e1003905.

Fehlberg S.D., Ty M.C., Ferguson C.J. 2014. Reexamination of a putative diploid hybrid taxon

using genetic evidence: The distinctiveness ofPhlox pilosasubsp.Deamii(polemoniaceae).

Int. J. Plant Sci. 175:781–793.

Felsenstein J. 1981. Skepticism Towards Santa Rosalia, or Why are There so Few Kinds of

Animals? Evolution. 35:124.

Ferguson C.J., Jansen R.K. 2002. A chloroplast DNA phylogeny of eastern Phlox

(Polemoniaceae): implications of congruence and incongruence with the ITS phylogeny.

Am. J. Bot. 89:1324–1335.

Ferguson C.J., Kramer F., Jansen R.K. 1999. Relationships of eastern north American phlox

(polemoniaceae) based on ITS sequence data. Syst. Bot. 24:616.

Fishman L., Wyatt R. 1999. Pollinator-mediated competition, reproductive character

displacement, and the evolution of selfing in Arenaria uniflora (Caryophyllaceae).

Evolution. 53:1723–1733.

Garner A.G., Goulet B.E., Farnitano M.C., Molina-Henao Y.F., Hopkins R. 2018. Genomic

Signatures of Reinforcement. Genes . 9.

Gottlieb L.D. 1972. Levels of confidence in the analysis of hybridization in plants. Ann. Mo.

Bot. Gard. 59:435.

Goulet B.E., Roda F., Hopkins R. 2017. Hybridization in plants: Old ideas, new techniques.

Plant Physiol. 173:65–78.

Goulet-Scott B.E., Garner A.G., Hopkins R. 2021. Genomic analyses overturn two long-standing

homoploid hybrid speciation hypotheses. Evolution. 75:1699–1710.

Grant V. 1981. Plant Speciation. New York Chichester, West Sussex: Columbia University

Press.

Green R.E., Krause J., Briggs A.W., Maricic T., Stenzel U., Kircher M., Patterson N., Li H., Zhai

W., Fritz M.H.-Y., Hansen N.F., Durand E.Y., Malaspinas A.-S., Jensen J.D., Marques-

Bonet T., Alkan C., Prüfer K., Meyer M., Burbano H.A., Good J.M., Schultz R., Aximu-

Petri A., Butthof A., Höber B., Höffner B., Siegemund M., Weihmann A., Nusbaum C.,

Lander E.S., Russ C., Novod N., Affourtit J., Egholm M., Verna C., Rudan P., Brajkovic

D., Kucan Ž., Gušic I., Doronichev V.B., Golovanova L.V., Lalueza-Fox C., la Rasilla

M. de, Fortea J., Rosas A., Schmitz R.W., Johnson P.L.F., Eichler E.E., Falush D., Birney

E., Mullikin J.C., Slatkin M., Nielsen R., Kelso J., Lachmann M., Reich D., Pääbo S.

2010. A Draft Sequence of the Neandertal Genome. Science. 328:710–722.

Guo C., Ma P.-F., Yang G.-Q., Ye X.-Y., Guo Y., Liu J.-X., Liu Y.-L., Eaton D.A.R., Guo Z.-H.,

Li D.-Z. 2021. Parallel ddRAD and genome skimming analyses reveal a radiative and

reticulate evolutionary history of the temperate bamboos. Syst. Biol. 70:756–773.

Gutenkunst R.N., Hernandez R.D., Williamson S.H., Bustamante C.D. 2009. Inferring the joint

demographic history of multiple populations from multidimensional SNP frequency data.

PLoS Genet. 5:e1000695.

Hadley E.B., Levin D.A. 1969. Physiological evidence of hybridization and reticulate evolution

in phlox maculata. Am. J. Bot. 56:561–570.

Hibbins M., Hahn M.W. 2021. Phylogenomic approaches to detecting and characterizing

introgression. EcoEvoRxiv.

Hoang D.T., Chernomor O., von Haeseler A., Minh B.Q., Vinh L.S. 2018. UFBoot2: Improving

the ultrafast bootstrap approximation. Mol. Biol. Evol. 35:518–522.

Hopkins R. 2013. Reinforcement in plants. New Phytol. 197:1095–1103.

Hopkins R., Rausher M.D. 2012. Pollinator-mediated selection on flower color allele drives

reinforcement. Science. 335:1090–1092.

Howard D.J. 1993. Reinforcement: origin, dynamics, and fate of an evolutionary hypothesis.

Hybrid zones and the evolutionary process.:46–69.

Kay K.M. 2006. Reproductive isolation between two closely related hummingbird-pollinated

neotropical gingers. Evolution. 60:538–552.

Kirkpatrick M. 2000. Reinforcement and divergence under assortative mating. Proc. Biol. Sci.

267:1649–1655.

Kong S., Kubatko L.S. 2021. Comparative Performance of Popular Methods for Hybrid

Detection using Genomic Data. Syst. Biol. 70:891–907.

Kulathinal R.J., Stevison L.S., Noor M.A.F. 2009. The genomics of speciation in Drosophila:

diversity, divergence, and introgression estimated using low-coverage genome

sequencing. PLoS Genet. 5:e1000550.

Landis J.B., Bell C.D., Hernandez M., Zenil-Ferguson R., McCarthy E.W., Soltis D.E., Soltis

P.S. 2018. Evolution of floral traits and impact of reproductive mode on diversification in

the phlox family (Polemoniaceae). Mol. Phylogenet. Evol. 127:878–890.

Lemmon E.M., Juenger T.E. 2017. Geographic variation in hybridization across a reinforcement

contact zone of chorus frogs (Pseudacris). Ecol. Evol. 7:9485–9502.

Léveillé-Bourret É., Chen B.-H., Garon-Labrecque M.-È., Ford B.A., Starr J.R. 2020. RAD

sequencing resolves the phylogeny, taxonomy and biogeography of Trichophoreae

despite a recent rapid radiation (Cyperaceae). Mol. Phylogenet. Evol. 145:106727.

Levin D.A. 1963. Natural Hybridization Between Phlox maculata and Phlox glaberrima and its

Evolutionary Significance. Am. J. Bot. 50:714.

Levin D.A. 1966a. The phlox pilosa complex: Crossing and chromosome relationships. Brittonia.

18:142.

Levin D.A. 1966b. Chromatographic evidence of hybridization and evolution in phlox maculata.

Am. J. Bot. 53:238–245.

Levin D.A. 1968. The genome constitutions of eastern north American phlox amphiploids.

Evolution. 22:612.

Levin D.A. 1969. The challenge from a related species: A stimulus for saltational speciation.

Am. Nat. 103:316–322.

Levin D.A. 1984. Genetic variation and divergence in a disjunct phlox. Evolution. 38:223–225.

Levin D.A. 1985. Reproductive Character Displacement in Phlox. Evolution. 39:1275.

Levin D.A., Kerster H.W. 1967. Natural selection for reproductive isolation in phlox. Evolution.

21:679–687.

Levin D.A., Schaal B.A. 1970a. Reticulate evolution in phlox as seen through protein

electrophoresis. Am. J. Bot. 57:977–987.

Levin D.A., Schaal B.A. 1970b. Corolla color as an inhibitor of interspecific hybridization in

phlox. Am. Nat. 104:273–283.

Levin D.A., Schaal B.A. 1972. Seed Protein Polymorphism in Phlox pilosa (Polemoniaceae).

Brittonia. 24:46.

Levin D.A., Smith D.M. 1965. An enigmatic phlox from Illinois. Brittonia. 17:254.

Levin D.A., Smith D.M. 1966. Hybridization and Evolution in the Phlox pilosa Complex. Am.

Nat. 100:289–302.

Levy M., Levin D.A. 1974. Novel flavonoids and reticulate evolution in the phlox pilosa–p.

Drummondii complex. Am. J. Bot. 61:156–167.

Levy M., Levin D.A. 1975. THE NOVEL FLAVONOID CHEMISTRY AND

PHYLOGENETIC ORIGIN OF PHLOX FLORIDANA. Evolution. 29:487–499.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv [q-bio.GN].

Locklear J.H. 2011a. Phlox a natural history and gardener's guide. Portland, OR: Timber Press.

Locklear J.H. 2011b. Phlox ovata L. (Polemoniaceae): Clarification of the Nomenclature of the

Allegheny Phlox. Castanea. 76:116–117.

Marques D.A., Meier J.I., Seehausen O. 2019. A combinatorial view on speciation and adaptive

radiation. Trends Ecol. Evol. 34:531–544.

McNeilly T., Antonovics J. 1968. Evolution in closely adjacent plant populations IV. Barriers to

gene flow. Heredity (Edinb.). 23:205–218.

Minh B.Q., Nguyen M.A.T., von Haeseler A. 2013. Ultrafast approximation for phylogenetic

bootstrap. Mol. Biol. Evol. 30:1188–1195.

Nguyen L.-T., Schmidt H.A., von Haeseler A., Minh B.Q. 2015. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. Mol. Biol. Evol.

32:268–274.

Noor M.A.F. 1999. Reinforcement and other consequences of sympatry. Heredity (Edinb.).

83:503–508.

Pease J.B., Haak D.C., Hahn M.W., Moyle L.C. 2016. Phylogenomics Reveals Three Sources of

Adaptive Variation during a Rapid Radiation. PLoS Biol. 14:e1002379.

Pickrell J.K., Pritchard J.K. 2012. Inference of population splits and mixtures from genome-wide

allele frequency data. PLoS Genet. 8:e1002967.

Poelstra J., Montero B.K., Lüdemann J., Yang Z., Rakotondranary S.J., Hohenlohe P., Stetter N.,

Ganzhorn J.U., Yoder A.D. 2021. RADseq data reveal a lack of admixture in a mouse

lemur contact zone contrary to previous microsatellite results. bioRxiv.

Roda F., Mendes F.K., Hahn M.W., Hopkins R. 2017. Genomic evidence of gene flow during

reinforcement in Texas Phlox. Mol. Ecol. 26:2317–2330.

Rothfels C.J. 2021. Polyploid phylogenetics. New Phytol. 230:66–72.

Schumer M., Rosenthal G.G., Andolfatto P. 2014. How common is homoploid hybrid

speciation? Evolution. 68:1553–1560.

Schumer M., Rosenthal G.G., Andolfatto P. 2018. What do we mean when we talk about hybrid

speciation? Heredity (Edinb.). 120:379–382.

Servedio M.R., Kirkpatrick M. 1997. The effects of gene flow on reinforcement. Evolution.

51:1764.

Servedio M.R., Noor M.A.F. 2003. The role of reinforcement in speciation: Theory and data.

Annu. Rev. Ecol. Evol. Syst. 34:339–364.

Smith D.M., Levin D.A. 1967. Karyotypes of eastern north American phlox. Am. J. Bot. 54:324.

Soltis P.S., Soltis D.E. 2009. The role of hybridization in plant speciation. Annu. Rev. Plant Biol.

60:561–588.

Stebbins G.L. 1959. The Role of Hybridization in Evolution. Proc. Am. Philos. Soc. 103:231–

251.

Suissa J.S., Kinosian S.P., Schafran P.W., Bolin J.F., Taylor W.C., Zimmer E.A. 2022.

Homoploid hybrids, allopolyploids, and high ploidy levels characterize the evolutionary

history of a western North American quillwort (Isoëtes) complex. Mol. Phylogenet. Evol.

166:107332.

Suni S.S., Hopkins R. 2018. The relationship between post-mating reproductive isolation and

reinforcement in Phlox. Evolution.

Swofford D.L. 2002. PAUP*. Phylogenetic analysis using parsimony (* and other methods).

Vol. Sinauer Associates, Sunderland, MA.

Taylor S.A., Larson E.L. 2019. Insights from genomes into the evolutionary importance and

prevalence of hybridization in nature. Nat. Ecol. Evol. 3:170–177.

Turissini D.A., Matute D.R. 2017. Fine scale mapping of genomic introgressions within the

Drosophila yakuba clade. PLoS Genet. 13:e1006971.

Turner B.L. 1998. Atlas of the Texas species of Phlox (Polemoniaceae). Phytologia. 85:309–326.

Wang N., Kelly L.J., McAllister H.A., Zohren J., Buggs R.J.A. 2021. Resolving phylogeny and

polyploid parentage using genus-wide genome-wide sequence data from birch trees. Mol.

Phylogenet. Evol. 160:107126.

Wherry E.T. 1930. The Eastern Short-styled Phloxes. Bartonia.:24–53.

Wherry E.T. 1931. The Eastern Long-styled Phloxes, part i. Bartonia.:18–37.

Wherry E.T. 1932. The Eastern Long-styled Phloxes, part 2. Bartonia.:14–26.

Wherry E.T. 1955. The genus Phlox, Monograph III. Morris Arbor, Philadelphia, USA.

Wherry E.T. 1956. The Genus Phlox. Am. Midl. Nat. 56:508.

Wherry E.T. 1965. The Genus Phlox, Ten Years After. Bartonia.:13–16.

Whitehouse E. 1945. Annual Phlox Species. Am. Midl. Nat. 34:388.

Yakimowski S.B., Rieseberg L.H. 2014. The role of homoploid hybridization in evolution: a

century of studies synthesizing genetics and ecology. Am. J. Bot. 101:1247–1258.