Quantitative MHC class-I/-II gene expression levels in *CDK12* mutated prostate cancer reveal

intratumorally T cell adaptive immune response in tumors

William Lautert-Dutra[1], Camila M. Melo[1], Luiz P. Chaves[1], Cheryl Crozier[2],

Fabiano P. Saggioro[3], Rodolfo B. dos Reis[1, 4], Jane Bayani[2,5], Sandro L. Bonatto[6] and Jeremy A. Squire[1, 7, *]

[1] Department of Genetics, Ribeirão Preto School of Medicine, University of São Paulo, Ribeirão Preto 14048-

900, SP, Brazil.

[2] Diagnostic Development, Ontario Institute for Cancer Research, Toronto, ON, Canada.

[3] Department of Pathology, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, SP, Brazil

[4] Division of Urology, Department of Surgery and Anatomy, Ribeirao Preto School of Medicine, University of

São Paulo, Ribeirao Preto 14048-900, SP, Brazil.

[5] Laboratory Medicine and Pathobiology, University of Toronto, Toronto, ON, Canada.

[6] School of Health and Life Sciences, Pontifical Catholic University of Rio Grande do Sul, Av. Ipiranga, 668,

90619-900, Porto Alegre, RS, Brazil

[7] Department of Pathology and Molecular Medicine, Queen's University, Kingston, ON K7L3N6, Canada.

*Corresponding author: Department of Genetics, Ribeirão Preto School of Medicine, University of São Paulo,

Ribeirão Preto 14048-900, SP, Brazil. Tel.: + 55 16 3602-3252; E-mail: squirej@fmrp.usp.br.

Abstract

Background

CDK12 inactivation is a predictive biomarker for immune checkpoint blockers (ICB) treatment response in

advanced prostate cancer (PCa), but some CDK12-altered patients fail to respond to ICB. Downregulation of

MHC expression has been described as a mechanism of intrinsic and acquired resistance to ICB in various can-

cers, but there is little information on whether MHC expression levels are altered in CDK12 defective PCa that

fails to respond to ICB treatments.

Methods

Using genomics data of primary and metastatic prostate cancer from two public domain cohorts and a retrospec-

tive cohort, we investigated variation in the expression of the MHC genes and associated downstream changes

in *CDK12* mutated patients. The findings of public domain data were validated using transcriptomic data from a

53-patient retrospective cohort from our Institute.

Results

Based on the analysis of gene expression quartiles, we divided the tumors into "High" and "Low" expression

levels of MHC-I and -II. *CDK12* defective tumors with increased MHC levels showed the activation of several

pathways associated with the immune system and elevated *PD-L1*, *IDO1*, and *TIM3* expression. These

transcriptomic findings were confirmed using expression analyses of our cohort of 53 primary PCa. There was

an increased composition of CD8+ T cells, B cells, γδ T cells, and M1 Macrophages in *CDK12* mutated tumors

with elevated MHC levels based on digital cytometric analyses. In contrast, *CDK12* defective tumors with de-

creased MHC expression were often subject to loss of heterozygosity (LOH) genomic events affecting MHC-I/-

II and the *HLA* gene cluster on chromosome 6. *CDK12* defective PCa expresses higher levels of classical MHC,

has an active and inflamed tumor microenvironment, and increases the presence of effector T cells.

Conclusions

Reduced MHC expression may be caused by the acquisition of specific somatic genomic events that reduce the

expression of antigen presentation genes. Combining CDK12 mutation, MHC expression levels, and LOH

status may better predict outcomes for ICB-eligible PCa. In addition, these findings draw attention to the need to

investigate therapeutic approaches for enhancing MHC expression in *CDK12* defective PCa to improve ICB

responses.

Keywords: Biomarkers, Immunotherapy, Transcriptomics, TCGA, SU2C, Digital cytometry, Antigen Presenta-

tion, Immune evasion, Adaptive immune system, Loss-of-Heterozygosity (LOH)

Background

Treatment of advanced prostate cancer (PCa) remains a therapeutic challenge. Men with distant metas-

tases at diagnosis have the worst overall survival (OS), with only 30% of patients surviving > 5 years [1]. For

recurrent disease, acquired resistance to androgen deprivation therapy (ADT) and chemotherapy remains a sig-

nificant cause of death [2]. Immune-checkpoint blockade (ICB) therapies, such as PD-L1 inhibition, have shown

only significant benefits in a minority of patients [3]. Thus, it is necessary to discover and characterize the ge-

netic pathways and molecular signatures that could help predict more effective disease progression control in

advanced PCa.

Tumor mutation burden affects the infiltration of immune cells in the tumor microenvironment (TME)

[4]. Because of the high load of neoantigens, tumors with defective DNA damage repair (DDR) pathways are

more suitable for immunotherapy [5]. The biallelic inactivation of Cyclin-dependent Kinase 12, *CDK12*, was

recently described as a novel immune active class of advanced PCa with a more aggressive phenotype, a high

mutation burden derived from focal tandem duplication events, and high inflammatory and immune cell infil-

trate distinct from other defective molecular subtypes [6–9]. Although this new subtype has been proposed as a

predictive biomarker of treatment response to immune-checkpoint blockade (ICB) in advanced PCa, many pa-

tients with *CDK12* alterations still fail to respond to ICB treatment. In a retrospective multi-center study,

Antonarakis *et al.* found that just 33% of the *CDK12*-altered advanced PCa patients had a prostate-specific anti-

gen (PSA) response and an increased progression-free survival of 5.4 months when treated with anti-PD-1 ther-

apy [10]. In another study, Schweizer *et al.* showed that of the 19 advanced patients who received ICB, 11

(59%) showed a response based on a decline in PSA, with two patients (11%) having a 100% PSA decline.

However, the molecular causes of ICB resistance in this molecular subtype of PCa are poorly understood

[10][11].

Tumor cells may develop various escape mechanisms that allow them to avoid recognition and destruc-

tion by the immune system. For example, the expression of checkpoint proteins (PD-1/PD-L1, CTLA-4, LAG3)

by tumors can modulate the activity of immune infiltrate cells [11]. They may also develop intrinsic cancer-cell

signaling that can suppress infiltrating immune cells (WNT/β-catenin) and increase pro-tumorigenic immune

cell infiltration (Tregs, M2 Macrophages) in the TME [12,13]. Another mechanism exploited by tumors to avoid

the recognition of cytotoxic T cells (CD8+) and antigen-presenting cells (APCs) is through loss of major histo-

compatibility complex class-I or class-II (MHC-I and MHC-II) [14].

Changes in MHC expression have been linked to tumor progression, poor prognosis, and reduced re-

sponse to ICB in different malignancies [15,16]. As classified by Garrido et al., the alteration in the MHC ex-

pression can be divided into two major mechanistic groups: tumors with "Soft" alterations are capable of recov-

ering or upregulating MHC antigens after cytokine exposure (e.g., characterized as having regulatory abnormali-

ties); those with "Hard" alterations cannot recover MHC expression (e.g., characterized as having structural

defects) [17]. During tumor evolution, the infiltration of cytotoxic lymphocytes eliminates highly immunogenic

tumor clones, causing a selection of surviving cell populations that have acquired MHC alterations through

either "Soft" or "Hard" mechanistic alterations [18]. Low expression of MHC-I and -II has been associated with

poor prognosis and resistance to anti-CTLA-4 and anti-PD-1, respectively [19,20]. However, in PCa, there is

presently limited information on the role of MHC-I and MHC-II expression and ICB response and if the low

expression of antigen presentation mechanism could better predict the lack of response to ICB in *CDK12*-altered

patients.

In PCa, *CDK12* inactivation is known to increase the immunogenicity of tumor cells, but the relation-

ship between *CDK12* loss and MHC expression has not been investigated. Changes in MHC-I and -II expression

are involved in tumor immune evasion in various types of cancer. There is also evidence that higher expression

of MHC genes can identify tumors likely to respond to ICB [15,16]. However, the molecular and genomic

mechanisms responsible for reducing MHC expression are poorly understood. Hence, we hypothesize that varia-

tion in the expression of the MHC genes could explain the lack of response to ICB in *CDK12* defective tumors.

Our transcriptomic analyses suggest that *CDK12* defective PCa expressing higher MHC levels have an active

and inflamed tumor microenvironment with elevated immunomodulatory pathway expression and increased

presence of effector T cells. These findings were validated using transcriptomic data from a 53-patient cohort

from our Institute. Analysis of public domain data showed that tumors with decreased MHC expression also

showed loss-of-heterozygosity (LOH) of MHC-I/-II and the *HLA* gene cluster on chromosome 6 associated with

MHC expression and antigen presentation. Collectively, these data suggest that implementing a combined

measure between *CDK12* mutation and the presence of MHC expression may better predict outcomes for pros-

tate cancer tumors classified as eligible for ICB treatment based on *CDK12* subtype and normal antigen presen-

tation function.

Methods

The CbioPortal for Cancer Genomics [21] was used to search primary (pPCa) and metastatic castra-

tion-resistant (mCRPC) prostate tumors with *CDK12* mutated sample cohorts with matched clinical information.

The classification of samples having *CDK12* loss-of-function followed previous reports [7–9] based only on the

presence of somatic mutations (non-synonymous mutations, deep deletions, and shallow deletions) in both

CDK12 alleles (*CDK12* defective) [6–8]. Only cohorts containing *CDK12* defective samples WES and RNA-

seq data were selected in the Genotypes and Phenotypes (dbGaP) database and applied for access under project

ID 29255 (Additional file 1: Figure S1, additional table 1).

We used a retrospective cohort of clinical intermediated primary PCa derived from radical prostatec-

tomies (*n*=53) to validate our transcriptomic findings. All 53 samples included in the cohort were primary PCa

collected by radical prostatectomy following National Comprehensive Cancer Network (NCCN) clinical prac-

tice guidelines [22] in the Department of Surgery and Anatomy, Urology Division at Ribeirao Preto Medical

School, Brazil, between 2007 and 2015 (Supplementary Table S1). The patients were classified according to the

presence of biochemical recurrence (BCR), defined as PSA>0.2 ng/ml within six months after radical prostatec-

tomy. Patient outcome data were collected to the last follow-up date (additional table 1). This retrospective

study was approved by the Ethics Committee in Research of Hospital of Ribeirão Preto, São Paulo, Brazil

(HCRP) number CAAE 60032122.8.0000.5440 and the Ethics Board of the University of Toronto (Protocol:

00043323).

RNA/DNA isolation

The RNA was isolated from tissues with tumor-rich areas previously marked by a pathologist (FPS).

Sections were processed at the Ontario Institute for Cancer Research, Toronto, Canada (OICR) using a dual

DNA and RNA extraction as previously described [23,24]. Briefly, hematoxylin and eosin slides were prepared

for all the Formalin-Fixed Paraffin-Embedded (FFPE) tissues. Histologic analysis of all the slides was per-

formed in the pathology department, and all tumor areas were carefully marked by an experienced pathologist

(FPS). The percentage of tumor cells (range 70-95% tumor-rich) within each marked tumor-rich area was esti-

mated and recorded (Table S2). Adjacent slides for each tumor were prepared, and the same areas of interest

were microdissected for RNA extraction.

Transcriptomic analysis

For the pPCa (TCGA-Prostate Adenocarcinoma, $n = 48$), we used the package recount2 to download

summarized experiments objects containing the transcription-level RNA-Seq abundance matrix [25,26]. For the

mCRPC (SU2C, $n = 10$), we downloaded SRA Paired-End (PE) reads using the SRAToolkit. The quality of the

raw reads was then measured using the FASTQC program. We quantified the transcripts using the software

Salmon v1.6.0 [27]. The transcriptome index was built using the reference GRCh38 version of the human ge-

nome and transcriptome, downloaded from ENSEMBL and GENCODE, following the manual instruction. We

then used the Tximport v1.22.0 package to import the transcription-level abundance and estimate raw counts

derived from the quantification step. The count data normalization, expression levels, and differential gene ex-

pression (DEG) analysis for pPCa and mCRPC cohorts were executed using the DESeq2 v1.34.0 [28,29]. We

used the clusterProfiler v4.2.1 to implement equally over-representation (ORA) enrichment analysis and gene

set enrichment analysis (GSEA) of the DEGs and the whole transcriptome profile, respectively, using the Gene

Ontology (GO) and Molecular Signatures Database (MSigDB) [30–35] (Figure S1).

The PanImmune Panel (NanoString Technologies Inc., Seattle, WA, USA) was used to profile the

RNA derived from the FFPE samples from our 53-patient tumor cohort. Raw expression data from the

PanImmune Panel was loaded in nSolver software v4.0 (NanoString Technologies) to perform the quality con-

trol (QC analysis) and to build the transcript matrix for downstream analysis.

MHC expression

We performed a hierarchical clustering analysis using representative MHC genes. We observed two

groups of *CDK12* defective tumors based on MHC expression, as shown in Additional file 1, Figure S1. We

dichotomized expression levels below or above the first quartile for each gene; we then used this classification

to generate the final logical values regarding the patient's MHC status ('High' or 'Low' expressed) (Additional

file 1: Figure S2)[36,37]. For the quartile quantification, we used the normalized expression values of the classi-

cal genes that composed each MHC class (e.g., MHC-I: *HLA-A*, *HLA-B*, *HLA-C*; MHC-II: *HLA-DPA1*, *HLA-*

DPB1, *HLA-DQA1*, *HLA-DQB1*, *HLA-DQB2*, *HLA-DRA*, *HLA-DRB5*, *HLA-DRB6*) [37]. Samples were classi-

fied as having MHC 'Low' expression when at least one gene composing each class was expressed at a low

level. In mCRPC *CDK12*- defective, all tumors classified as MHC-High presented similar high expression of

both MHC-I and -II; thus, we called these tumors just MHC High (henceforth, mCRPC *CDK12* defective MHC

High). The summary values used to classify the tumors are shown in table S1. Our retrospective cohort of low-

intermediate risk primary PCa derived from radical prostatectomies was used to validate our transcriptomic

findings that were derived from public domain cohorts.

Genomic profile

We downloaded SRA Paired-End (PE) reads for both pPCa (n=48) and mCRPC (n=10) and processed

it as previously described. The GRChg38 reference was sorted using the SeqKit [38]. The final fastq files were

then aligned to hg38 using bwa with a penalty for up to 3 mismatches per read [39,40]. Sam files were convert-

ed to bam files and processed using samtools v1.16.1. To determine whether any MHC expression differences

were related to genomic alterations, we used the FACETS (Fraction and Allele-Specific Copy Number Estimat-

ed from Tumor Sequencing) algorithm [41]. This approach uses matched normal-tumor WES and provides

mutant allele-specific copy-number homozygous/heterozygous deletions, copy-number neutral loss-of-

heterozygosity (LOH), allele-specific gain/amp in genomic loci associated with genes involved in antigen

presentation (Additional table 2). Reference and variant allele read counts were extracted from the bam file for

common, polymorphic SNPs downloaded from dbSNP (GRCh38p7) using FACETS *snp_pileup* function with a

minimum threshold for mapping quality, the minimum threshold for the base quality, and minimum read depth

of 15, 20, 20, respectively. The pre-processing followed the suggested recommendations from the manual, and

genomic intervals of 150-250bp were used to avoid hyper-segmentation in high polymorphic neighborhood

regions. Mutant allele-specific copy-number changes were declared when the points changes were greater than a

pre-determined critical value (cval) of 100 compared to constant copy-number regions [42,43].

Digital cytometry

To investigate and quantify the immune cell composition in the TME of tumors having *CDK12* defects,

we used the bulk tissue gene expression profiles (GEP) from RNA-seq data from both pPCa and mCRPC with

the digital cytometry CIBERSORTx [44–46]. CIBERSORTx uses bulk tissue GEP, compares the data with

prior knowledge of expression profiles from purified leukocytes, and estimates a tumor's relative immune abun-

dance composition. We used the 'signature matrix' containing a validated leukocyte GEP of 22 human hemato-

poietic cell phenotypes, leukocyte gene signature matrix (LM22), to estimate the immune cell composition from

the TME.

Computational and statistical analysis

A GNU/Linux environment was used to perform the quality control and quantify the raw reads to the

human transcriptome. Subsequently, downstream analysis was performed in RStudio software (R Foundation

for Statistical Computing, R v4.1.2). Pearson Correlation was used to analyze the normalized expression levels

(coef. level = 0.95). A gene was considered differentially expressed when log2 foldchange > 1 was expressed

from the reference group with a *p*-adjusted value < 0.05. For the enrichment analysis, we used a cutoff value of

0.05.

Results

Transcriptomics

We hypothesize that variation in the expression of the MHC genes among *CDK12* defective tumors

could provide predictive information on poor ICB responses in this molecular subtype of PCa. To investigate

MHC expression levels in PCa further, we compared the transcriptomics of prostate tumors, including *CDK12*

defective (*CDK12* defective) prostate tumors, classified as "MHC High" (top 75% quartile) to the

transcriptomics of tumors classified as having a low expression of MHC genes (bottom 25% quartile).

CDK12 defective MHC-I/-II High-expression tumors showed distinct transcriptome profiles with upregulation

of IFN-γ-responsive genes and an inflamed TME

We first compared the transcriptome of *CDK12* defective tumors classified as MHC-High and MHC-

Low (Figure 1a-c, Additional file 1: Figure S2). The 48 primary PCa *CDK12* defective MHC-I High showed

1319 upregulated and 2153 downregulated genes (Additional table 3, 4) and showed four apparent clusters (Fig-

ure 1a). The upper cluster (I) demonstrates *IGHV* and *IGLV* over-expressed genes linked to *CDK12* MHC I

high. The central clusters (II, III) have four *HLA* genes overexpressed in the MHC I high group, which supports

our classification based on *CDK12* defective MHC expression (e.g., *HLA-A*, *HLA-B*, *HLA-C*). Also, this cluster

showed upregulation of many genes related to antigen presentation and CD8+ T cell activity (Figure 1a). The

lower cluster (IV) possesses 14 downregulated genes when MHC I is highly expressed. Likewise, *CDK12* de-

fective MHC-II High exhibited 1129 upregulated and 580 downregulated genes (Additional tables 3, 5). Our

comparison showed two clusters (Figure 1b). The upper cluster showed the upregulation of genes related to

cytotoxicity, immune cell migration, and immune suppression (Figure 1b).

The analysis of ten mCRPC tumors identified 230 upregulated and 46 downregulated genes differen-

tially expressed in *CDK12* defective MHC-high (Additional tables 3, 6) and showed two distinct clusters (Fig-

ure 1c). The lower cluster showed the upregulation of genes linked to innate and adaptive immune response

CD4+ and CD8+ T cell activity (Figure 1c). In conclusion, for both primary PCa and metastatic CRPC, *CDK12*

defective PCa that express higher MHC levels was associated with transcriptomic changes that indicate a gen-

eral pattern of activation of IFN-γ-responsive and cytotoxic activity genes.

In our institutional validation cohort, we identified 25 upregulated and 14 downregulated genes that

were differentially expressed in the MHC-high group and showed two distinct clusters (Figure 1d). The patients

classified as MHC I high group have *HLA* genes overexpressed, consistent with our classification based on

RNA-seq using both public domain cohorts. Thus, for both primary PCa and metastatic CRPC, *CDK12* defec-

tive PCa that express higher MHC levels was associated with transcriptomic changes that indicate a general

pattern of activation of IFN-γ-responsive and cytotoxic activity genes.

To better understand the transcriptional alterations in *CDK12* defective tumors classified as MHC-I/-II

high-expressed, we used ORA and GSEA analysis to identify potential functional associations of expression

changes associated with differentially expressed genes. In primary PCa, tumors with *CDK12* defective MHC-I

and MHC-II high-expressed showed significative enrichment of pathways related to activation of immune cells

and antigen presentation by MHC-I and MHC-II, as expected (Table 1, Additional table 4, 5). Furthermore,

GSEA results showed activation of many pathways in *CDK12* defective MHC-I high-expressed (e.g., Interferon

Gamma Response, Interferon Alpha Response, TNFA Signaling via NFKB, and IL2/STAT5 Signaling (Figure

2a, Additional table 4). In addition, the *CDK12* defective MHC-II High group had similar pathway activation

but with the addition of Allograft Rejection (Figure 2b, Additional table 5).

The *CDK12* defective metastatic group with MHC high-expressed showed enrichment of cytotoxicity

and adaptive cytotoxicity immune response (Table 1, Additional table 6). In addition, GSEA results showed

activation of various hallmark and Reactome pathways (e.g., Allograft Rejection, Interferon Gamma Response,

Interferon Alpha Response, IL2/STAT5 Signaling, TCR Signaling, Interleukin 10 Signaling, and the suppres-

sive PD1 Signaling pathway) (Figure 2c-d, Additional table 6).

Next, to investigate and understand the involvement of the DEGs in pathways driving tumor progres-

sion, we further used ORA analysis on the DEGs from our validation cohort. As expected, tumors with MHC-I

high-expressed showed activation of several immune-related pathways (Table 1, Additional file 1: Figure S4).

Our transcriptomic analysis suggests that *CDK12* defective PCa expressing high MHC genes display

the activation of various pathways associated with an active and inflamed tumor microenvironment. Also, these

results demonstrate the feasibility of using MHC expression levels to uncover intrinsic inflammatory activity in

PCa.

CDK12 defective MHC-I/-II High groups exhibited tumor microenvironment with high expression of

chemokines and immunomodulatory genes.

We investigated the difference between chemokines and immunomodulatory gene expression to eluci-

date the underlying immunologic signature in the TME of CDK12 defective tumors expressing high MHC

genes. In primary PCa, the *CDK12* defective MHC-I High group exhibited upregulation of many chemokines

linked to APCs and effector T cell migration (Addition file 1, Figure S4) and the immunomodulatory genes

HAVCR2 (*TIM3*) and *IDO1*. (Figure 3a-b). Also, the MHC-II high group presented increased expression of

chemokines related to Natural killer, T effector cell, and B cell infiltration (Addition file 1, Figure S2, Figure

S5), and the immunomodulatory genes *CD274* (*PD-L1*), *HAVCR2* (*TIM3*), and *IDO1* (Figure 3c-e). Based on

Pearson Correlation analysis, we observed a significant positive correlation between MHC-I complex genes and

IDO1 but with no other investigated gene (Addition file 1, Figure S7a). Interestingly, among the MHC-II genes,

correlation analysis showed a significant positive association between the expression of the MHC-II complex

and the immunomodulatory genes *CD274* (*PD-L1*), *IDO1*, and *HAVCR2* (*TIM3*) (Figure S7b).

The *CDK12* defective metastatic group with high expression of MHC showed high expression of

chemokines (Addition file 1, Figure S6) and the immunomodulatory gene *CD274* (*PD-L1*) (Figure 3f). Fur-

thermore, we observed a significant positive association between the immunomodulatory genes *HAVCR2*

(*TIM3*) and *CTLA4*, *LAG3*, and MHC-I and -II genes (Addition file 1, Figure S7c-d). *CDK12* defective tumors'

expression of higher levels of MHC genes shows a pattern of upregulation of chemokine and

immunomodulatory mechanisms.

CDK12 defective MHC-I/-II High CDK12 defective tumors showed distinct immune cell infiltrate in the TME.

Our transcriptomics analysis in the public domain cohort predicted that *CDK12* defective tumors ex-

pressing high levels of MHC-I/-II genes would display enrichment of immune cells with high effector and cyto-

toxic activity and the co-inhibitory expression of inhibitory pathways by tumor cells (Figures 1, 2, 3). We used

in silico cytometry (CIBERSORTx) to test this hypothesis and to estimate the immune cell composition in the

TME of *CDK12* defective tumors expressing high levels of MHC-I/-II genes.

We investigated whether tumors expressing high MHC-I and -II levels affected the variation in im-

mune cell composition in *CDK12* defective tumors in both public domain cohorts. In primary PCa, *CDK12*

defective MHC-I high-expressed showed a significant increase in the composition of Naïve B cells, CD8+ T

cells, and γδ T cells. In contrast, the reduced composition of Mastocytes (Figure 4a). In comparison, the MHC-II

high-expressed group exhibited an increase of γδ T cells and reduced composition of Plasma cells and M0 mac-

rophages (Figure 4b).

Additionally, the *CDK12* defective metastatic group with high expression of MHC showed a significa-

tive increased composition of CD8+ T cell and M1 Macrophages (Figure 4c). Our results suggest that *CDK12*

defective PCa expressing higher levels of MHC displayed increased APCs and effector lymphocytes traffic in

their TME.

Genomic alterations of MHC-I/-II region associated with defective CDK12

During tumor evolution, the infiltration of cytotoxic lymphocytes eliminates highly immunogenic tu-

mor clones, causing a preferential selection and survival of cell populations that have acquired MHC alterations.

To determine whether the low levels of MHC-I/-II genes in *CDK12* defective tumors were derived from somatic

genomic alterations affecting antigen presentation genes, we used an allele-specific copy number algorithm to

estimate the copy number profile of *CDK12* defective tumors expressing low levels of MHC-I/-II genes. The

algorithm incorporates quantitative analysis of RNA-seq data derived from MHC-I/-II and *HLA* genes on chro-

mosome 6 and several other genes, such as *PTEN*, commonly subject to LOH in PCa [47]. Estimates include the

determination of the relative clonality of allele-specific copy number alterations based on the fraction of tumor

cells bearing LOH.

WES of primary PCa revealed both subclonal allele-specific copy-neutral losses of heterozygosity

(CN-LOH) and complete loss of heterozygosity (LOH) events affecting MHC-I/-II and *HLA* genes on chromo-

some 6. Both major clonal and minor subclonal events were detected by comparing the copy number to the

estimated cellular fraction of tumor cells harboring the copy number alteration (Table 2 and Figure S8). One

example is illustrated by tumor TCGA-KK-A8IA in which only one major clonal event was detected with a

cellular fraction = 0.833, and LOH events were detected on chromosomes 2, 5, and 10. While TCGA-YL-A8HO

revealed four subclonal events, capturing biallelic loss, CN-LOH, and LOH in chromosomes 2, 5, 6, 15, and 16.

325 Tumors expressing lower levels of MHC-I/II genes may be associated with the somatic genomic LOH events

326 affecting regional transcriptional regulators of MHC expression and the cis-acting regulatory components of the

327 MHC (Table 2). Two patients also showed subclonal complete loss of the *B2M* locus (ID: TCGA-EJ-8471,

328 cellular fraction = 0.37; TCGA-YL-A8HO, cellular fraction = 0.27. Table 2). Interestingly, *B2M* is an important

329 component of the MHC-I complex, and mutations of this gene have previously been associated with ICB re-

330 sistance [48]. In contrast to primary PCa, metastatic tumors harboring *CDK12* mutation expressing low levels of

331 MHC-I/II showed allele-specific copy number gains in the MHC-I/*II* complex and key regulators of MHC ex-

332 pression (Table 3 and Figure S9). Interestingly, one patient showed a clonal LOH event at the *JAK2* and *B2M*

333 loci (ID: 5115615, cellular fraction = 0.90).

334   These data indicate that low expression of MHC in primary tumor clones appears to be associated with

335 frequent somatic CN-LOH and LOH subclonal genomic alterations in primary tumors harboring *CDK12* muta-

336 tion. While metastatic tumors revealed both LOH and high copy-number gains affecting antigen presentation

337 genes and their regulators. Therefore, these subclonal CN-LOH and LOH events are associated with impaired

338 MHC expression that may contribute to reducing tumor immunogenicity.

340 **Discussion**

341   In PCa, *CDK12* inactivation is known to increase the immunogenicity of tumor cells [9,49–52]. MHC-I

342 and -II expression changes are involved in tumor immune evasion in various cancer types [37,53–55]. Using *in*

343 *silico* two public domain RNA-seq cohorts and transcriptional profiling of our institutional cohort, we observed

344 two groups of *CDK12* defective tumors regarding MHC expression. Tumors expressing higher levels of MHC

345 showed the activation of multiple pathways associated with the immune system, significantly high expression of

346 immunomodulatory genes, and increased CD8+ T cells, B cells, $\gamma\delta$ T cells, and M1 Macrophages composition

347 in their TME. Also, Tumors with low MHC expression showed allele-specific copy-number alteration in genes

348 involved in regulating MHC expression and antigen presentation.

349   Analysis of the clinical data showed that higher expression of MHC genes in pPCa and mCRPC, re-

350 spectively, exhibited a trend towards better overall survival when compared to the MHC lower expression group

351 ($p > 0.05$, Figure S3). Several studies have found similar MHC-I and -II gene expression relationships associat-

352 ed with better clinical outcomes and different resistance mechanisms [20,56,57]. Lack of MHC-I expression in

353 melanoma was linked to higher odds of progressive disease and resistance to anti-CTLA4 but not anti-PD1 [20].

354 In the same study, expression of MHC-II in tumors was associated with tumors more likely to respond to anti-

PD1 treatment. MHC-II expression was linked to a positive outcome in Hodgkin Lymphoma treated with nivolumab (anti-PD1) [56]. Park et al. showed that MHC-II positivity in tumor cells was associated with better disease-free survival in patients with lymph node metastasis [57]. From a practical standpoint, these results suggest that analysis of MHC-I and -II expression levels could be developed to determine whether *CDK12* defective prostate tumors expressing higher levels of MHC genes are more likely to respond to ICB treatments.

Transcriptomic signatures associated with MHC expression variation have been described in different tumor types [37]. These signatures capture the activity of genes and biological pathways related to tumor cells' crosstalk with the TME and appear to correlate with clinical responses to ICB [20,37,58–60]. Ayers *et al.* proposed a gene expression profile with eighteen genes relevant to predicting the clinical outcome of anti-PD1 therapy [58]. This IFN-γ gene signature in pretreatment tumor biopsies was associated with better results in melanoma, head-and-neck squamous cell carcinoma, and gastric cancer treated with pembrolizumab. In our study, the *CDK12* patient tumors expressing high levels of MHC genes showed the presence IFN-γ gene signature, which might reflect a better response to anti-PD1 therapy (Figure S9). Our results also showed the activation of pathways related to IFN-γ response in both the pPCA and mCRPC cohorts (Figure 1b, d, f, 3b, d, Figure 4b-c).

An IFN-γ gene signature can alternatively activate the expression of immunomodulatory mechanisms and promote adaptive resistance to therapeutic anti-PD1 [61,62]. IFN-γ is a central actor in the elimination phase during immunoediting [63]. However, exacerbated levels of IFN-γ are also associated with developing protumor molecular mechanisms leading to an immunosuppressive and tolerogenic TME [62]. We identified the upregulation of immunomodulatory molecules, such as *IDO1*, *TIM3,* and *CD274(PD-L1),* in both prostate tumors expressing higher levels of MHC-I and -II genes (Figure 2). Interestingly, the *CDK12* patient tumors expressing high levels of MHC genes showed the presence of IFN-γ gene signature, including upregulation of a non-classical MHC molecule (*HLA-E*), which is a known regulator of NK cell and CTL [64,65]. We also found a link between *IGHV* and *IGLV* over-expressed genes in a group of the *CDK12* defective pPCa expressing high levels of MHC-I (Figure 1a, upper cluster I). The production of Ig by tumor cells (cancer-derived Ig) is described in various tumors and PCa [66–69]. Also, cancer-derived Ig may act as checkpoint proteins and inhibit effector T cells and NK cells [67,70]. Our findings indicate the presence of an acquired resistance mechanism in a subset of *CDK12* defective prostate tumors expressing high levels of MHC genes. We suggest that measuring the basal expression of MHC genes could be developed as a biomarker to characterize *CDK12* patient samples

with a previously inflamed environment and which would reflect the expression of immunomodulatory genes

linked to resistance to anti-PD1.

The expression of MHC molecules plays a pivotal role in providing the signals necessary to recognize

and activate the immune system against tumor neoantigens and is essential to controlling tumor growth through

cytotoxic activity [63,71]. We found a higher abundance of CD8+ T cells in *CDK12* defective tumors express-

ing high levels of MHC genes (Figure 4). This result may either indicate the dependence of CD8+ T cells in

MHC-I antigen presentation or suggest that enhanced CD4+ Th infiltrate could support the continued accumula-

tion of CD8+ CTLs in the tumor microenvironment [59]. Interestingly, *CDK12* pPCa with high expression of

both MHC-I and -II genes showed increased levels of γδ T cells (Figure 4a-b). The basal effector immune cell

population, such as γδ T cells in pPCa, may contribute to IFN-γ signaling and indicate the dependence of the

MHC-unrestricted recognition role of γδ T cells and its presence in the TME [20,72].

The loss of MHC expression could be derived from two different disruption means [73]. First, regula-

tory abnormalities downregulate the expression of MHC genes through mechanisms that do not affect the ge-

nomic structure of HLA genes. In such cases, specific T-cells can recover the MHC expression-mediated re-

sponse (e.g., IFN-γ signaling). Second, during tumor evolution, the infiltration of cytotoxic lymphocytes elimi-

nates tumor clones high immunogenic, causing a selection of clones that may present structural alteration, or

"Hard" lesions, in the MHC loci or other genomic regions (e.g., *B2M*, *IFN*, *STAT*) distinct from the derived

clone [48].

Genomic studies of chromosome 6 in various cancers suggest reduced MHC expression is often associ-

ated with genetic and LOH aberrations that may result in reduced antigen presentation and, thus, facilitate im-

mune evasion [15,74]. In keeping with these data, our analysis of WES from pPCa and mCRPC revealed CN-

LOH and LOH in *CDK12* defective tumors expressing low levels of MHC-I/II genes at known regulators of

MHC expression and the components of the MHC (Table 2-3). Two pPCa patients also showed complete loss of

the *B2M* locus, while one mCRPC exhibited a LOH event at the *B2M*, an important component of the MHC-I

[48]. These results might indicate that after selection, the low expression of MHC genes is derived from tumor

clones harboring "Hard" genomics alterations comprising loci involved in normal MHC expression. Therefore,

these clones will be invisible to the immune system and contribute to tumor progression [18].

This study has some limitations. First, future studies in a large number of patients harboring *CDK12*

mutations are needed to address potential statistical bias regarding our low number of patients. Second, we did

not address the potential molecular mechanism underlying the alteration of the low expression of MHC in

CDK12 defective PCa, including epigenetic alterations and miRNA activity. Further studies are needed to ap-

proach the causes of MHC disruption derived from "Soft" alterations that occur in *CDK12* defective prostate

cancer. Third, because the non-tumor cell content was not defined in this study, we could not establish a limit

for the contribution of the *CDK12* defective tumor cells and TME to the MHC expression since the RNA-seq

relies on data from bulk tissue [75].

Conclusion

We distinguished two sets of *CDK12* defective tumors based on differential MHC expression levels.

Our data suggest that *CDK12* defective PCa expressing higher levels of classical MHC genes have an active and

inflamed tumor microenvironment with elevated immunomodulatory pathway expression and increased pres-

ence of effector T cells. Tumors with decreased MHC expression showed copy-number alteration in genes that

regulate MHC expression and antigen presentation. More extensive *in vitro* and *in vivo* investigations are re-

quired to relate this molecular subtype to potential actionable immunomodulatory mechanisms or possible ther-

apeutic targets in PCa. Depending on the mechanism, the downregulation of MHC expression can sometimes be

therapeutically restored to improve anti-tumor immunity [76]. These findings, therefore, support further investi-

gations into the clinical use of MHC expression modulation in *CDK12* defective prostate cancer to identify

tumors likely to respond to ICB treatment.

List of abbreviations

ADT: Androgen Deprivation therapy

APC: Antigen-presenting cells

CDK12: Cyclin-dependent Kinase 12

CN-LOH: Copy-neutral loss of heterozygosity

dbGaP: The database of Genotypes and Phenotypes

DDR: **DNA damage repair**

DEGs: Differentially expressed genes

FACETS: Fraction and Allele-Specific Copy Number Estimated from Tumor Sequencing

FDR: False discovery rate

GEP: Gene expression profile

GO: Gene ontology

ICB: Immune-checkpoint blockade

LM22: Leukocyte gene signature matrix

LOH: loss of heterozygosity

mCRPC: Metastatic Castration-Resistant Prostate Cancer

MHC: Major histocompatibility complex

MSigDB: Molecular Signature Database

NCCN: National Comprehensive Cancer Network

ORA: Over-representation analyses

PCa: Prostate Cancer

PE: Paired-end

pPCa: Primary prostate cancer

PRAD: Prostate Adenocarcinoma

PSA: Prostate-specific antigen

TCGA-PRAD: The Cancer Genome Atlas

TME: Tumor microenvironment

LOH: Loss-of-Heterozygosity

Ethics approval

This retrospective study was approved by the Ethics Committee in Research of Hospital of Ribeirão Preto, São

Paulo, Brazil (HCRP) number CAAE 60032122.8.0000.5440 and the Ethics Board of the University of Toronto

(Protocol: 00043323).

Consent to publish

Not applicable.

Availability of data and materials

The original dataset is deposited in the database of Genomic and Phenotypes (dbGaP) under accession numbers

phs000178.v11.p8.c1, phs000915.v2.p2.c1. The original clinical information of both cohorts is deposited in the

CbioPortal and GDC Portal. All other data supporting the conclusions of this article are included within the

article and its additional files. Additional files related to our retrospective cohort are available at

https://github.com/WilliamLautertD/CDK12_PCa.

Competing interests

J.B. has the patent application "A Molecular Classifier for Personalized Risk Stratification for Patients with

Prostate Cancer" under consideration. Status: PCT, Filing date: June 18, 2021, International Application No.:

PCT/CA2021/050837, PCT Application Title: Molecular Classifiers for Prostate Cancer. Previous US Provi-

sional Status: Filing Date: June 18, 2020, US Provisional Patent No. 63/040.692, US Provisional Application

Title: Use of Molecular Classifiers to Diagnose, Treat, and Prognose Prostate Cancer.

Funding

This work is partly funded by *Fundação de Amparo à Pesquisa de São Paulo (FAPESP)* (2019/22912-8).

W.L.D. is funded by the *FAPESP* (2021/12271-5). L.P.C. is funded by *FAPESP* (2020/12816-9) and by the

Government of Ontario S. L. B. and J.A.S. are Research Career Awardees of *Conselho Nacional de*

Desenvolvimento Científico e Tecnológico (CNPq).

Authors' contributions

Conception: W. L. D. and J. A. S

Interpretation or analysis of data: W. L. D., S. L. B., and J. A. S

Data acquisition, tissue handling, QC, performing assays: A.E.S., E.W., L.P.C., C.M.M., J.B.

Preparation of the manuscript: W. L. D. and J. A. S.

Revision for important intellectual content: All authors

Supervision: S. L. B. and J. A. S.

Acknowledgments

The authors acknowledge the contributions of the Department of Surgery and Anatomy, the Department of

Pathology, the Medicine School of Ribeirao Preto, and members of Diagnostic Development at the Ontario

Institute for Cancer Research. The authors are grateful to L. A. F. B. and S. P. F. B. for the informatic technical

support in the early stage of the project, Genomic and Molecular laboratory/PUCRS for technical assistance,

and dbGap and the Cbioportal for archiving and distributing the data from the above-cited studies.

References

1.   Rebello RJ, Oing C, Knudsen KE, Loeb S, Johnson DC, Reiter RE, et al. Prostate cancer.
Nat Rev Dis Primers. 2021 Dec 1;7(1).
2.   Wong MCS, Goggins WB, Wang HHX, Fung FDH, Leung C, Wong SYS, et al. Global Inci-
dence and Mortality for Prostate Cancer: Analysis of Temporal Patterns and Trends in
36 Countries. Eur Urol. 2016 Nov;70(5):862–74.
3.   Binnewies M, Roberts EW, Kersten K, Chan V, Fearon DF, Merad M, et al. Understand-
ing the tumor immune microenvironment (TIME) for effective therapy. Nat Med.
2018 May 1;24(5):541–50.
4.   Melo CM, Vidotto T, Chaves LP, Lautert-Dutra W, dos Reis RB, Squire JA. The role of
somatic mutations on the immune response of the tumor microenvironment in pros-
tate cancer. Int J Mol Sci. 2021;22(17).
5.   Schumacher TN, Schreiber RD. Neoantigens in cancer immunotherapy. Vol. 348, Sci-
ence. American Association for the Advancement of Science; 2015. p. 69–74.
6.   Wu YM, Cieślik M, Lonigro RJ, Vats P, Reimers MA, Cao X, et al. Inactivation of CDK12
Delineates a Distinct Immunogenic Class of Advanced Prostate Cancer. Cell. 2018 Jun
14;173(7):1770-1782.e14.
7.   Nguyen B, Mota JM, Nandakumar S, Stopsack KH, Weg E, Rathkopf D, et al. Pan-
cancer Analysis of CDK12 Alterations Identifies a Subset of Prostate Cancers with Dis-
tinct Genomic and Clinical Characteristics. Eur Urol. 2020 Nov 1;78(5):671–9.
8.   Rescigno P, Gurel B, Pereira R, Crespo M, Rekowski J, Rediti M, et al. Characterizing
CDK12-mutated prostate cancers A C. Clinical Cancer Research. 2021 Jan
15;27(2):566–74.
9.   Sokol ES, Pavlick D, Frampton GM, Ross JS, Miller VA, Ali SM, et al. Pan-Cancer Analy-
sis of CDK12 Loss-of-Function Alterations and Their Association with the Focal Tan-
dem-Duplicator Phenotype . Oncologist. 2019 Dec 1;24(12):1526–33.
10.   Antonarakis ES, Pedro ;, Velho I, Fu W, Wang H, Agarwal N, et al. CDK12-Altered Pros-
tate Cancer: Clinical Features and Therapeutic Outcomes to Standard Systemic Ther-
apies, Poly (ADP-Ribose) Polymerase Inhibitors, and PD-1 Inhibitors [Internet]. Vol. 4,
JCO Precis Oncol. 2020. Available from: https://doi.org/10.
11.   Spranger S, Gajewski TF. Mechanisms of Tumor Cell-Intrinsic Immune Evasion. Annu
Rev Cancer Biol [Internet]. 2018;2:213–28. Available from:
https://doi.org/10.1146/annurev-cancerbio-
12.   Li X, Xiang Y, Li F, Yin C, Li B, Ke X. WNT/β-catenin signaling pathway regulating T cell-
inflammation in the tumor microenvironment. Vol. 10, Frontiers in Immunology.
Frontiers Media S.A.; 2019.
13.   Horton BL, Fessenden TB, Spranger S. Tissue Site and the Cancer Immunity Cycle. Vol.
5, Trends in Cancer. Cell Press; 2019. p. 593–603.
14.   McGranahan N, Rosenthal R, Hiley CT, Rowan AJ, Watkins TBK, Wilson GA, et al. Al-
lele-Specific HLA Loss and Immune Escape in Lung Cancer Evolution. Cell. 2017 Nov
30;171(6):1259-1271.e11.
15.   Dhatchinamoorthy K, Colbert JD, Rock KL. Cancer Immune Evasion Through Loss of
MHC Class I Antigen Presentation. Front Immunol. 2021 Mar 9;12.

16.   Thibodeau J, Bourgeois-Daigneault MC, Lapointe R. Targeting the MHC Class II antigen
549   presentation pathway in cancer immunotherapy. Oncoimmunology. 2012 Sep
550   28;1(6):908–16.
551   17.   Garrido F, Aptsiauri N. Cancer immune escape: MHC expression in primary tumours
552   versus metastases. Vol. 158, Immunology. Blackwell Publishing Ltd; 2019. p. 255–66.
553   18.   Algarra I, Garrido F, Garcia-Lora AM. MHC heterogeneity and response of metastases
554   to immunotherapy. Vol. 40, Cancer and Metastasis Reviews. Springer; 2021. p. 501–
555   17.
556   19.   Watson NFS, Ramage JM, Madjd Z, Spendlove I, Ellis IO, Scholefield JH, et al.
557   Immunosurveillance is active in colorectal cancer as downregulation but not com-
558   plete loss of MHC class I expression correlates with a poor prognosis. Int J Cancer.
559   2006 Jan 1;118(1):6–10.
560   20.   Rodig SJ, Gusenleitner D, Jackson DG, Gjini E, Giobbie-Hurder A, Jin C, et al. MHC pro-
561   teins confer differential sensitivity to CTLA-4 and PD-1 blockade in untreated meta-
562   static melanoma [Internet]. Vol. 10, Sci. Transl. Med. 2018. Available from:
563   https://www.science.org
564   21.   Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative analy-
565   sis of complex cancer genomics and clinical profiles using the cBioPortal. Sci Signal.
566   2013 Apr 2;6(269).
567   22.   Mohler JL, Antonarakis ES, Armstrong AJ, D'Amico A v., Davis BJ, Dorff T, et al. Pros-
568   tate Cancer, Version 2.2019, NCCN Clinical Practice Guidelines in Oncology. Journal of
569   the National Comprehensive Cancer Network. 2019 May;17(5):479–505.
570   23.   Bayani J, Yao CQ, Quintayo MA, Yan F, Haider S, D'Costa A, et al. Molecular stratifica-
571   tion of early breast cancer identifies drug targets to drive stratified medicine. NPJ
572   Breast Cancer. 2017 Dec 15;3(1):3.
573   24.   Patel PG, Selvarajah S, Guérard KP, Bartlett JMS, Lapointe J, Berman DM, et al. Relia-
574   bility and performance of commercial RNA and DNA extraction kits for FFPE tissue
575   cores. PLoS One. 2017 Jun 22;12(6):e0179732.
576   25.   Imada EL, Sanchez DF, Collado-Torres L, Wilks C, Matam T, Dinalankara W, et al. Re-
577   counting the FANTOM Cage Associated Transcriptome. bioRxiv. 2019;
578   26.   Abeshouse A, Ahn J, Akbani R, Ally A, Amin S, Andry CD, et al. The Molecular Taxono-
579   my of Primary Prostate Cancer. Cell. 2015 Nov 5;163(4):1011–25.
580   27.   Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-
581   aware quantification of transcript expression. Nat Methods. 2017;14(4):417–9.
582   28.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
583   RNA-seq data with DESeq2. Genome Biol. 2014 Dec 5;15(12).
584   29.   Zhu A, Ibrahim JG, Love MI. Heavy-Tailed prior distributions for sequence count data:
585   Removing the noise and preserving large differences. Bioinformatics. 2019 Jun
586   1;35(12):2084–92.
587   30.   Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrich-
588   ment tool for interpreting omics data. The Innovation. 2021 Aug 28;2(3).
589   31.   Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The Molecu-
590   lar Signatures Database Hallmark Gene Set Collection. Cell Syst. 2015 Dec
591   23;1(6):417–25.
592   32.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al.
593   Gene set enrichment analysis: A knowledge-based approach for interpreting genome-

wide expression profiles [Internet]. 2005. Available from:
www.pnas.orgcgidoi10.1073pnas.0506580102
33. Khatri P, Sirota M, Butte AJ. Ten years of pathway analysis: Current approaches and
outstanding challenges. Vol. 8, PLoS Computational Biology. 2012.
34. Carbon S, Douglass E, Dunn N, Good B, Harris NL, Lewis SE, et al. The Gene Ontology
Resource: 20 years and still GOing strong. Nucleic Acids Res. 2019 Jan 8;47(D1):D330–
8.
35. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level
estimates improve gene-level inferences. F1000Res. 2015 Dec 30;4:1521.
36. Koopman LA, Corver WE, van der Slik AR, Giphart MJ, Fleuren GJ. Multiple Genetic
Alterations Cause Frequent and Heterogeneous Human Histocompatibility Leukocyte
Antigen Class I Loss in Cervical Cancer. Journal of Experimental Medicine. 2000 Mar
20;191(6):961–76.
37. Schaafsma E, Fugle CM, Wang X, Cheng C. Pan-cancer association of HLA gene expres-
sion with cancer prognosis and immunotherapy efficacy. Br J Cancer. 2021 Aug
3;125(3):422–32.
38. Shen W, Le S, Li Y, Hu F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q
File Manipulation. PLoS One. 2016 Oct 5;11(10):e0163962.
39. Bailey MH, Tokheim C, Porta-Pardo E, Sengupta S, Bertrand D, Weerasinghe A, et al.
Comprehensive Characterization of Cancer Driver Genes and Mutations. Cell. 2018
Apr;173(2):371-385.e18.
40. Montesion M, Murugesan K, Jin DX, Sharaf R, Sanchez N, Guria A, et al. Somatic HLA
Class I Loss Is a Widespread Mechanism of Immune Evasion Which Refines the Use of
Tumor Mutational Burden as a Biomarker of Checkpoint Inhibitor Response. Cancer
Discov. 2021 Feb 1;11(2):282–92.
41. Shen R, Seshan VE. FACETS: allele-specific copy number and clonal heterogeneity
analysis tool for high-throughput DNA sequencing. Nucleic Acids Res. 2016 Sep
19;44(16):e131–e131.
42. Villarroya-Beltri C, Osorio A, Torres-Ruiz R, Gómez-Sánchez D, Trakala M, Sánchez-
Belmonte A, et al. Biallelic germline mutations in MAD1L1 induce a syndrome of an-
euploidy with high tumor susceptibility. Sci Adv. 2022 Nov 4;8(44).
43. Murciano-Goroff YR, Schram AM, Rosen EY, Won H, Gong Y, Noronha AM, et al. Re-
version mutations in germline BRCA1/2-mutant tumors reveal a BRCA-mediated phe-
notype in non-canonical histologies. Nat Commun. 2022 Nov 23;13(1):7182.
44. Chen B, Khodadoust MS, Liu CL, Newman AM, Alizadeh AA. Profiling tumor infiltrating
immune cells with CIBERSORT. In: Methods in Molecular Biology. Humana Press Inc.;
2018. p. 243–59.
45. Newman AM, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, et al. Robust enumeration
of cell subsets from tissue expression profiles. Nat Methods. 2015 Apr 29;12(5):453–
7.
46. CIBERSORTx. CIBERSORTx [Internet]. CIBERSORTx. [cited 2022 May 11]. Available
from: https://cibersortx.stanford.edu/
47. Jamaspishvili T, Berman DM, Ross AE, Scher HI, de Marzo AM, Squire JA, et al. Clinical
implications of PTEN loss in prostate cancer. Nat Rev Urol. 2018 Apr 20;15(4):222–34.
48. Zaretsky JM, Garcia-Diaz A, Shin DS, Escuin-Ordinas H, Hugo W, Hu-Lieskovan S, et al.
Mutations Associated with Acquired Resistance to PD-1 Blockade in Melanoma. New
England Journal of Medicine. 2016 Sep;375(9):819–29.

49. Lotan TL, Antonarakis ES. CDK12 deficiency and the immune microenvironment in
prostate cancer. Clinical Cancer Research. 2021 Jan 15;27(2):380–2.

50. Li Y, Zhang H, Li Q, Zou P, Huang X, Wu C, et al. CDK12/13 inhibition induces immuno-
genic cell death and enhances anti-PD-1 anticancer activity in breast cancer. Cancer
Lett. 2020 Dec 28;495:12–21.

51. Dubbury SJ, Boutz PL, Sharp PA. CDK12 regulates DNA repair genes by suppressing
intronic polyadenylation. Nature. 2018 Dec 6;564(7734):141–5.

52. Juan HC, Lin Y, Chen HR, Fann MJ. Cdk12 is essential for embryonic development and
the maintenance of genomic stability. Cell Death Differ. 2016 Jun 1;23(6):1038–48.

53. Garrido F, Ruiz-Cabello F, Aptsiauri N. Rejection versus escape: the tumor MHC di-
lemma. Vol. 66, Cancer Immunology, Immunotherapy. Springer Science and Business
Media Deutschland GmbH; 2017. p. 259–71.

54. Aptsiauri N, Ruiz-Cabello F, Garrido F. The transition from HLA-I positive to HLA-I neg-
ative primary tumors: the road to escape from T-cell responses. Vol. 51, Current
Opinion in Immunology. Elsevier Ltd; 2018. p. 123–32.

55. Montesion M, Murugesan K, Jin DX, Sharaf R, Sanchez N, Guria A, et al. Somatic hla
class i loss is a widespread mechanism of immune evasion which refines the use of
tumor mutational burden as a biomarker of checkpoint inhibitor response. Cancer
Discov. 2021 Feb 1;11(2):282–92.

56. Roemer MGM, Redd RA, Cader Z, Pak CJ, Abdelrahman S, Ouyang J, et al. JOURNAL
OF CLINICAL ONCOLOGY Major Histocompatibility Complex Class II and Programmed
Death Ligand 1 Expression Predict Outcome After Programmed Death 1 Blockade in
Classic Hodgkin Lymphoma. 2018; Available from: https://doi.org/10.1200/JCO.2017.

57. Park IA, Hwang SH, Song IH, Heo SH, Kim YA, Bang WS, et al. Expression of the MHC
class II in triple-negative breast cancer is associated with tumor-infiltrating lympho-
cytes and interferon signaling. PLoS One. 2017 Aug 17;12(8):e0182786.

58. Ayers M, Lunceford J, Nebozhyn M, Murphy E, Loboda A, Kaufman DR, et al. IFN-γ-
related mRNA profile predicts clinical response to PD-1 blockade. Journal of Clinical
Investigation. 2017 Aug 1;127(8):2930–40.

59. Johnson DB, Estrada M v., Salgado R, Sanchez V, Doxie DB, Opalenik SR, et al. Mela-
noma-specific MHC-II expression represents a tumour-autonomous phenotype and
predicts response to anti-PD-1/PD-L1 therapy. Nat Commun. 2016 Jan 29;7.

60. Johnson DB, Bordeaux J, Kim JY, Vaupel C, Rimm DL, Ho TH, et al. Quantitative spatial
profiling of PD-1/PD-L1 interaction and HLA-DR/IDO-1 predicts improved outcomes of
anti-PD-1 therapies in metastatic melanoma. Clinical Cancer Research. 2018 Nov
1;24(21):5250–60.

61. Koyama S, Akbay EA, Li YY, Herter-Sprie GS, Buczkowski KA, Richards WG, et al. Adap-
tive resistance to therapeutic PD-1 blockade is associated with upregulation of alter-
native immune checkpoints. Nat Commun. 2016 Feb 17;7.

62. Zaidi MR, Merlino G. The Two Faces of Interferon-γ in Cancer. Clinical Cancer Re-
search. 2011 Oct 1;17(19):6118–24.

63. Schreiber RD, Old LJ, Smyth MJ. Cancer Immunoediting: Integrating Immunity's Roles
in Cancer Suppression and Promotion [Internet]. 2011. Available from:
www.sciencemag.org

64. Salomé B, Sfakianos JP, Ranti D, Daza J, Bieber C, Charap A, et al. NKG2A and HLA-E
define an alternative immune checkpoint axis in bladder cancer. Cancer Cell. 2022
Sep;40(9):1027-1043.e9.

65.   Borst L, van der Burg SH, van Hall T. The NKG2A–HLA-E Axis as a Novel Checkpoint in
689   the Tumor Microenvironment. Clinical Cancer Research. 2020 Nov 1;26(21):5549–56.
690   66.   Cui M, Huang J, Zhang S, Liu Q, Liao Q, Qiu X. Immunoglobulin Expression in Cancer
691   Cells and Its Critical Roles in Tumorigenesis. Front Immunol. 2021 Mar 24;12.
692   67.   Li X, Ni R, Chen J, Liu Z, Xiao M, Jiang F, et al. The Presence of IGHG1 in Human Pan-
693   creatic Carcinomas Is Associated With Immune Evasion Mechanisms. Pancreas. 2011
694   Jul;40(5):753–61.
695   68.   Pan B, Zheng S, Liu C, Xu Y. Suppression of IGHG1 gene expression by siRNA leads to
696   growth inhibition and apoptosis induction in human prostate cancer cell. Mol Biol
697   Rep. 2013 Jan 2;40(1):27–33.
698   69.   Qin C, Sheng Z, Huang X, Tang J, Liu Y, Xu T, et al. Cancer-driven IgG promotes the
699   development of prostate cancer though the SOX2-CIgG pathway. Prostate. 2020 Sep
700   6;80(13):1134–44.
701   70.   Wang Z, Geng Z, Shao W, Liu E, Zhang J, Tang J, et al. Cancer-derived sialylated IgG
702   promotes tumor immune escape by binding to Siglecs on effector T cells. Cell Mol
703   Immunol. 2020 Nov 21;17(11):1148–62.
704   71.   Qin H, Zhou C, Wang D, Ma W, Liang X, Lin C, et al. Specific antitumor immune re-
705   sponse induced by a novel DNA vaccine composed of multiple CTL and T helper cell
706   epitopes of prostate cancer associated antigens. Immunol Lett. 2005 Jun 15;99(1):85–
707   93.
708   72.   Born WK, Reardon CL, O'Brien RL. The function of γδ T cells in innate immunity. Vol.
709   18, Current Opinion in Immunology. 2006. p. 31–8.
710   73.   Garrido F, Cabrera T, Aptsiauri N. "Hard" and "soft" lesions underlying the HLA class I
711   alterations in cancer cells: Implications for immunotherapy. Vol. 127, International
712   Journal of Cancer. 2010. p. 249–56.
713   74.   Garrido MA, Perea F, Vilchez JR, Rodríguez T, Anderson P, Garrido F, et al. Copy neu-
714   tral loh affecting the entire chromosome 6 is a frequent mechanism of hla class i al-
715   terations in cancer. Cancers (Basel). 2021 Oct 1;13(20).
716   75.   van den Berge K, Hembach KM, Soneson C, Tiberi S, Clement L, Love MI, et al. RNA
717   Sequencing Data: Hitchhiker's Guide to Expression Analysis. Annu Rev Biomed Data
718   Sci. 2019 Jul 20;2(1):139–73.
719   76.   Taylor BC, Balko JM. Mechanisms of MHC-I Downregulation and Role in Immunother-
720   apy Response. Front Immunol. 2022 Feb 28;13.
721

Legends

Figure 1. MHC-I/-II high expression in CDK12 defective prostate cancer showed distinct transcriptome

activity. (a) Transcriptome heatmap exhibiting clustering of top 50 DEGs in pPCa CDK12-Mut MHC-I High

group (n = 30). The upper cluster (I) has IGHV and IGLV over-expressed genes linked to *CDK12* MHC I high.

The central clusters (II and III) have four HLA genes overexpressed in the MHC I high group, supporting our

CDK12 mutated MHC expression classification. The lower cluster (IV) has 14 downregulated genes when

MHC I is highly expressed. **(b)** Transcriptome heatmap exhibiting clustering of top 50 DEGs in the pPCa

CDK12-Mut MHC-II High group (n = 23). The upper cluster (I) has upregulation of genes related to cytotoxici-

ty, immune cell migration, and immune suppression. We could not observe clinical features associated with

MHC I or II clusters. **(c)** Transcriptome heatmap exhibiting clustering of top 50 DEGs in the mCRPC CDK12-

Mut MHC High group (*n* = 6). The combined MHC I and II showed two distinct clusters. The lower cluster

showed the upregulation of genes linked to innate and adaptive immune response and CD4+ and CD8+ T cell

activity. **(d)** Transcriptome heatmap exhibiting clustering of the DEGs from our validation cohort (RP cohort).

The DEGs are relative to the CDK12-Mut 'Low' group. Clinical information is displayed on top of the heatmap

for each patient. The color scale in the heatmap represents the Z-score of the normalized read counts for each

gene, where the red scale indicates upregulated and blue low-expressed genes.

Figure 2. MHC high expressed *CDK12* defective prostate tumors are associated with activation of im-

mune-related pathways. (a) Gene Set Enrichment Analysis of Hallmark's pathways in pPCa *CDK12-Mut*

MHC-I High (*n* = 30)**. (b)** Gene Set Enrichment Analysis of Hallmark pathways in pPCa *CDK12-Mut* MHC-II

High (*n* = 23). **(c-d)** Gene Set Enrichment Analysis of Hallmarks and Reactome pathways in mCRPC *CDK12-*

Mut MHC High (*n* = 10). A normalized enrichment score indicates each factor's positive or negative association

with the condition of interest, which means activation or suppression of the pathway. Enrichment analyses were

performed using *clusterProfiler* and *p*-adjusted value = 0.05 as the cutoff.

Figure 3. MHC high expressed CDK12 defective prostate cancer showed enhanced expression of

immunomodulatory genes. RNA-sequencing expression level of immunomodulatory genes **(a)** *IDO1* ($p <$

0.018), **(b)** *HAVCR2* (*TIM3*) ($p < 0.0022$) in *CDK12-Mut* MHC-I High. *n* = 48; **(c)** *CD274* (*PD-L1*) ($p <$

0.0028), **(d)** *HAVCR2* (*TIM3*) ($p < 0.00012$)*,* **(e)** *IDO1* ($p < 0.001$) in *CDK12-Mut* MHC-II High. *n* = 48; **(f)**

CD274 (*PD-L1*) ($p < 0.0095$). *n* = 10; *: *p*-value < 0.05; **: *p*-value < 0.01, by Mann-Whitney test. *IDO1*,

Indoleamine 2,3-Dioxygenase 1; *HAVCR3* (*TIM3*), T-Cell Immunoglobulin Mucin Receptor 3; *CD274* (*PD-L1*),

Programmed Cell Death 1 Ligand 1.

Figure 4. MHC high expressed *CDK12* defective is associated with enhanced T cell recruitment of pros-

tate tumors. (a) CIBERSORT-derived immune cell abundance of 22 distinct cell subsets based on *CDK12*

defective MHC-I High pPCa (*n* = 30)**. (b)** CIBERSORT-derived immune cell abundance of 22 distinct cell

subsets based on *CDK12* defective MHC-II High pPCa (*n* = 23). **(c)** CIBERSORT-derived immune cell abun-

dance of 22 distinct cell subsets based on *CDK12* defective MHC High mCRPC ($n = 10$). *: *p*-value < 0.05; **:

p-value < 0.01, by Mann-Whitney test.

Table 1. Enrichment results. Enriched GO (BP) pathways of the upregulated genes from TCGA *CDK12*-Mut

MHC-I and MHC-II High, mCRPC group MHC High, and for our validation cohort (RP) MHC-I High group.

Enrichment analyses were performed using *clusterProfiler* and *p*-adjusted value = 0.05 as the cutoff.

Table 2. *CDK12* defective in primary MHC low expressed tumors. Integral results of FACETS analysis of

whole-exome sequencing data from *CDK12* defective primary tumors expressing low levels of MHC-I/II. The

table shows the integer copy number for the allelic ratio of the heterozygous SNPs in each patient's tu-

mor/normal pair. The first column is the patient ID, and the second column is the MHC region loci and other

genes commonly subject to LOH in PCa. The last two columns represent the total copy number and the minor

copy number, respectively. Tumors with a 2:1 ratio are considered normal for each position, while 2:0 and 1:0

represent CN-LOH and LOH events. The cellular fraction represents the estimated number of cells harboring the

genotype. Primary PCa revealed subclonal CN-LOH and LOH at *JAK1, STAT1, IRF2, IRF1, MHC-I/II, TAP1,*

TAP2, IFNR1, JAK2, B2M, CIITA, NLRC5, and *IFNR1*. Two patients showed biallelic loss at the *B2M* locus

(Total copy number/minor copy number ratio=0:0). *MHC-I = *HLA-A, HLA-B,* and *HLA-C;* MHC-II = *HLA-*

DRA, HLA-DRB5, HLA-DRB6, HLA-DPA1, HLA-DPB1, HLA-DQA1, HLA-DQB1, and *HLA-DQB2*.

Table 3. *CDK12* defective in metastatic MHC low expressed tumors. Integral results of FACETS analysis of

whole-exome sequencing data from *CDK12* defective metastatic tumors expressing low levels of MHC-I/II. The

same parameters shown in Table 1 were used to characterize LOH for each gene of interest. Metastatic PCa

showed allele-specific copy number gains at *JAK1, STAT1, IRF2, IRF1, MHC-I/II, TAP1, TAP2, IFNR1, JAK2,*

B2M, CIITA, NLRC5, and *IFNR1*. One patient showed a clonal LOH event at the *JAK2* and *B2M* loci (Total

copy number/minor copy number ratio=1:0). *MHC-I = *HLA-A, HLA-B,* and *HLA-C;* MHC-II = *HLA-DRA,*

HLA-DRB5, HLA-DRB6, HLA-DPA1, HLA-DPB1, HLA-DQA1, HLA-DQB1, and *HLA-DQB2*.

A)

B)

C)

D)

A)

B)

C)

	ID	Description	p.adjust
TCGA-PRAD_MHC-I	GO:0051249	regulation of lymphocyte activation	1,48E-19
	GO:0002460	adaptive immune response based on somatic recombination of immune receptors built from immunoglobulin superfamily domains	1,48E-19
	GO:0050867	positive regulation of cell activation	2,78E-20
	GO:0002449	lymphocyte mediated immunity	1,16E-18
	GO:0006958	complement activation, classical pathway	2,77E-14
	GO:0006959	humoral immune response	1,08E-09
	GO:0050853	B cell receptor signaling pathway	2,20E-08
	GO:0002440	production of molecular mediator of immune response	7,15E-08
	GO:0042742	defense response to bacterium	1,39E-07
	GO:0006910	phagocytosis, recognition	5,66E-08
TCGA-PRAD_MHC-II	GO:0051249	regulation of lymphocyte activation	2,00E-25
	GO:0042110	T cell activation	5,84E-25
	GO:0050867	positive regulation of cell activation	9,60E-23
	GO:0007159	leukocyte cell-cell adhesion	2,50E-20
	GO:0002460	adaptive immune response based on somatic recombination of immune receptors built from immunoglobulin superfamily domains	2,80E-20
	GO:0002443	leukocyte mediated immunity	3,53E-19
	GO:1903037	regulation of leukocyte cell-cell adhesion	8,17E-19
	GO:0070661	leukocyte proliferation	1,75E-15
	GO:0046651	lymphocyte proliferation	5,77E-14
	GO:0070663	regulation of leukocyte proliferation	1,64E-12
mCRPC_MHC	GO:0051251	positive regulation of lymphocyte activation	4,87E-15
	GO:0002449	lymphocyte mediated immunity	6,43E-15
	GO:0002253	activation of immune response	6,67E-15
	GO:0042110	T cell activation	3,27E-11
	GO:0002460	adaptive immune response based on somatic recombination of immune receptors built from immunoglobulin superfamily domains	1,76E-09
	GO:0006959	humoral immune response	1,21E-06
	GO:0019724	B cell mediated immunity	2,55E-06
	GO:0050863	regulation of T cell activation	9,19E-07
	GO:0007159	leukocyte cell-cell adhesion	3,27E-04
	GO:1903037	regulation of leukocyte cell-cell adhesion	1,75E-03
RP_MHC	GO:0019221	cytokine-mediated signaling pathway	1,32E+11
	GO:0022409	positive regulation of cell-cell adhesion	1,32E+11
	GO:0002474	antigen processing and presentation of peptide antigen via MHC class I	1,21E+12
	GO:0072182	regulation of nephron tubule epithelial cell differentiation	1,21E+12
	GO:0060333	interferon-gamma-mediated signaling pathway	0.001
	GO:0002479	antigen processing and presentation of exogenous peptide antigen via MHC class I, TAP-dependent	0.001
	GO:0042127	regulation of cell population proliferation	0.001
	GO:0042590	antigen processing and presentation of exogenous peptide antigen via MHC class I	0.001
	GO:0010719	negative regulation of epithelial to mesenchymal transition	0.002
	GO:0048640	negative regulation of developmental growth	0.002

Patient ID	Gene	chrom	start	end	cellular fraction	Total copy number	Minor copy number
KK.A8IA	*STAT1*	2	134713128	192194260	0.833	1	0
KK.A8IA	*IRF1*	5	55231421	143147410	0.833	1	0
KK.A8IA	*PTEN*	10	87552401	89139399	0.833	1	0
KK.A8IA	*B2M*	15	41191425	77615398	0.208	0	0
KK.A8IG	*IFNGR1*	6	65302925	170584366	0.668	1	0
KK.A8I6	*PTEN*	10	73243437	104454566	0.332	0	0
KK.A8I6	*NLRC5*	16	49396608	90096324	0.11	0	0
XJ.A9DX	*NLRC5*	16	49280867	57722991	0.62	1	0
ZG.A9LM	*STAT1*	2	134317491	192057773	0.47	2	0
ZG.A9LM	*B2M*	15	19882284	101922415	0.47	2	0
ZG.A9LM	*NLRC5*	16	56276055	89228602	0.47	2	0
EJ.8472	*JAK1*	1	21619903	67705610	0.37	1	0
EJ.8472	*B2M*	15	42751857	97971259	0.37	2	0
XQ.A8TA	*IRF2*	4	177976002	189984351	0.92	1	0
XQ.A8TA	*IRF1*	5	128464930		0.92	2	0
XQ.A8TA	*PTEN*	10	87790275	89599433	0.92	0	0
YL.A8HO	*STAT1*	2	119157673	199659594	0.12	0	0
YL.A8HO	*IRF1*	5	143082	181260211	0.38	2	0
YL.A8HO	*MHC-I/II, TAP1/2*	6	304637	170584276	0.27	1	0
YL.A8HO	*IFNGR1*	6	304637	170584276	0.27	1	0
YL.A8HO	*B2M*	15	19882294	101922415	0.27	1	0
YL.A8HO	*CIITA*	16	8644541	58196892	0.22	2	0
YL.A8HO	*NLRC5*	16	8644541	58196892	0.22	2	0
XK.AAJA	*PTEN*	10	7243639	12235132	0.54	1	0
KK.A7AP	*B2M*	15	33311111	57091863	0.87	1	0
KK.A7AP	*NLRC5*	16	48347484	90177833	0.87	1	0
HC.7744	*B2M*	15	42515191	45178103	0.12	0	0
KK.A8IG	*IFNGR1*	6	65302925	170584366	0.668	1	0
EJ.7784	*JAK2*	9	14859	18718116	0.437	1	0
HC.A48F	*JAK1*	1	41513578	80451865	0.86	1	0
HC.A48F	*IRF2*	4	85671	189984257	0.21	1	0
HC.A48F	*IRF1*	5	54400082	171456817	0.86	1	0
HC.A48F	*MHC-I/II, TAP1/2*	6	304637	170584366	0.31	1	0
HC.A48F	*IFNGR1*	6	304637	170584366	0.31	1	0
HC.A48F	*PTEN*	10	49168049	118015066	0.21	1	0
HC.A48F	*B2M*	15	44380814	50749712	0.86	1	0
HC.A48F	*NLRC5*	16	35085998	90177833	0.86	1	0

Table 2

Patient ID	Gene	chrom	start	end	cellular fraction	Total copy number	Minor copy number
5115412	JAK1	1	33804897	68440771	0.49	5	2
5115412	STAT1	2	187346385	205615949	0.54	5	2
5115412	IRF2	4	182891198	185757023	0.87	6	2
5115412	IRF1	5	112892242	139881181	0.81	4	2
5115412	MHC-I	6	29555893	32256053	0.87	4	1
5115412	MHC-II	6	32288409	38345160	0.30	6	2
5115412	TAP-1	6	32288409	38345160	0.30	6	2
5115412	TAP-2	6	32288409	38345160	0.30	6	2
5115412	JAK2	9	2804198	15486990	0.59	4	2
5115412	PTEN	10	80253502	93958843	0.30	6	2
5115412	B2M	15	30626570	59138344	0.59	4	2
5115412	CIITA	16	198972	17358048	0.59	4	2
5115412	NLRC5	16	57372390	71634040	0.52	4	2
5115412	IFGRR2	21	33230307	34889712	0.87	5	2
1115161	IRF2	4	184116877	186709443	0.51	6	3
1115161	HLA-A	6	29330997	30628358	0.79	3	1
1115161	HLA-B	6	30635016	31382266	0.59	5	2
1115161	HLA-C	6	30635016	31382266	0.59	5	2
1115161	JAK2	9	52538	33113788	0.20	3	1
5115615	JAK1	1	63571201	65432064	0.912	3	1
5115615	STAT1	2	188522965	192194738	0.912	3	1
5115615	MHC-I	6	29944749	30415099	0.899	3	1
5115615	MHC-I/II	6	30469283	32407241	0.902	4	2
5115615	MHC-II	6	32439760	32666486	0.912	3	1
5115615	MHC-II/TAP1/2	6	32666737	34289022	0.902	4	2
5115615	JAK2	9	117666	9091026	0.906	1	0
5115615	B2M	15	41691484	46828068	0.906	1	0
5115615	CIITA	16	9938304	11178578	0.912	3	1
5115615	IFGRR2	21	33335636	37018290	0.899	3	1
5115412	JAK1	1	33804897	68440771	0.49	5	2
5115412	STAT1	2	187346385	205615949	0.54	5	2
5115412	IRF2	4	182891198	185757023	0.87	6	2
5115412	IRF1	5	112892242	139881181	0.81	4	2
5115412	MHC-I	6	29555893	32256053	0.87	4	1
5115412	MHC-II	6	32288409	38345160	0.30	6	2
5115412	TAP-1	6	32288409	38345160	0.30	6	2
5115412	TAP-2	6	32288409	38345160	0.30	6	2

Table 3