Calculating ATP production rates from oxidative phosphorylation and glycolysis during cell activation

Brandon R. Desousa[1], Kristen K.O. Kim[1], Wei Y. Hsieh[2], Anthony E. Jones[1], Pamela Swain[3], Danielle H. Morrow[1], David A. Ferrick[3], Orian S. Shirihai[4], Andrew Neilson[3], David A. Nathanson[1], George W. Rogers[3], Brian P. Dranka[3], Anne N. Murphy[5], Charles Affourtit[6], Steven J. Bensinger[2], Linsey Stiles[1,4], Natalia Romero[3], and Ajit S. Divakaruni[1,7]

[1]Department of Molecular and Medical Pharmacology, University of California, Los Angeles, Los Angeles, CA 90095, USA

[2]Department of Microbiology, Immunology, and Molecular Genetics, University of California, Los Angeles, Los Angeles, CA 90095, USA

[3]Agilent Technologies, 5301 Stevens Creek Boulevard, Santa Clara, CA 95051, USA

[4]Department of Medicine, University of California, Los Angeles, Los Angeles, CA 90095, USA

[5]Cytokinetics Inc. 350 Oyster Point Blvd., South San Francisco, CA 94080, USA

[6]School of Biomedical Sciences, University of Plymouth, Plymouth PL6 8BU, United Kingdom

[7]Lead contact

Correspondence: adivakaruni@mednet.ucla.edu

ABSTRACT

The balance between oxidative phosphorylation and glycolysis is shifted in a variety of cell types and tissues upon activation, proliferation, and differentiation. However, description of these metabolic switches has been mostly qualitative, particularly upon acute activation signals. We therefore applied a method to quantify rates of ATP production from oxygen consumption and extracellular acidification data obtained from the Seahorse XF Analyzer to study cell growth and activation. The method detects substantive changes in ATP utilization upon neuronal depolarization and T cell receptor activation that are not evident from steady-state ATP measurements. Furthermore, quantifying ATP utilization during pro-inflammatory macrophage activation reveals the hallmark increase in glycolysis is not merely a compensatory response to decreased oxidative phosphorylation upon LPS treatment. Rather, these changes occur independently, and the profound increases in glycolysis and total ATP utilization upon activation is largely driven by MyD88-linked signaling and independent of oxidative phosphorylation.

INTRODUCTION

Snapshot measurements of cellular ATP content are routinely used to measure changes in energy metabolism, but are generally informative only in extreme cases of energetic dysfunction such as ischemia[1]. During physiological changes such as cell activation, cells readily meet an increased ATP demand without appreciably changing steady-state ATP levels. Biochemical and mass-spectrometry-based approaches can measure the ATP:ADP ratio – a far more useful metric – but their utility is limited. As destructive measurements conducted on cell lysates, they are unable to measure real-time changes and are prone to experimental noise that masks subtle but meaningful differences. Kinetic measurements of energy status are therefore extraordinarily specialized, requiring transfection of a genetically engineered ATP:ADP sensor[2].

Importantly, these approaches lack information about whether the ATP is derived from oxidative phosphorylation or glycolysis. A redistribution between ATP-generating pathways can characterize – and in some cases determine – cell function and fate[3–6]. Glycolysis and offshoot pathways play essential roles in generating precursors for macromolecule biosynthesis[3], supporting one-carbon metabolism[7], and fueling cytoplasmic reducing power[8]. Biological processes such as stem cell pluripotency[9], oncogenesis[4], growth factor signaling[10], and immune cell activation[11] are therefore frequently associated with a preferential shift towards glycolysis. Similarly, terminal differentiation is often accompanied by an increased capacity for oxidative phosphorylation and the complete oxidation of glucose[12–15].

Fluorescence-based, multi-well technologies such as Seahorse XF Analysis can characterize these shifts with simultaneous measurements of the oxygen consumption rate (OCR) and extracellular acidification rate (ECAR), providing relative approximations of the balance between glycolysis and oxidative phosphorylation[16]. However, direct assessment of the total rate of ATP production and the balance between these two pathways in not immediately straightforward. The units are reported on different scales (OCR in pmol O_2/min; ECAR in mpH/min) that preclude direct comparison. Additionally, although the main driver of extracellular pH changes in two-dimensional cell culture systems is often glycolysis[17], acidification of the experimental medium itself cannot quantify lactate efflux[14,18,19].

The utility of these relative measurements is further limited because the maximum amount of ATP that can be generated from the two pathways varies substantially: for a molecule of glucose, the maximum glycolytic yield is 2 ATP as opposed to 33.45 ATP for oxidative phosphorylation[20]. As such, a large increase in acidification of the extracellular medium may not necessarily indicate a rebalancing of ATP production if it is matched with even a marginal increase in oxygen consumption.

We therefore developed a method to transform OCR and ECAR into rates of ATP production from oxidative phosphorylation and glycolysis. This involves three principal steps: (i) converting ECAR into rates of H^+ production (pmol H^+/min), (ii) accounting for non-glycolytic acidification so

changes in H^+ production quantitatively reflect lactate efflux, and (iii) stoichiometric conversion of pmol O_2/min and pmol lactate/min into pmol ATP/min. A similar strategy has been applied previously[20,21], but the values empirically obtained here using 17-21 primary and immortalized cell types sharply contrast previous work. Applying this analysis to models of neuronal depolarization, cancer cell growth, and immune cell activation provides information about cellular bioenergetic status that extends far beyond either static ATP measurements or relative measurements of oxidative phosphorylation and glycolysis.

MATERIALS AND METHODS

Chemicals

All chemicals were purchased from Sigma-Aldrich unless otherwise stated.

Animals

All animal procedures were performed in accordance with the NIH Guide for the Care and Use of Laboratory Animals, and approved by the UCLA Animal Research Committee (ARC). Genetically modified animals [$Myd88^{-/-}$ (009088) and $Trif^{-/-}$ (005037)] were purchased from The Jackson Laboratory.

Murine bone marrow-derived macrophages (BMDMs)

BMDMs were cultured as previously described[22]. Briefly, macrophages were isolated from femurs of male C57BL/6 mice and maintained at 37°C in a humidified 5% CO_2 incubator. Cells were treated with 3 mLs RBC lysis buffer (Sigma-Aldrich) to remove red blood cells for 5 minutes, centrifuged at 386g for 5 minutes, and resuspended in BMDM culture medium. Culture medium consisted of high-glucose DMEM (Gibco 11965) supplemented with 10% heat-inactivated fetal bovine serum (FBS), 2 mM L-glutamine, 100 U/ml penicillin, 100 μg/mL streptomycin, 500 μM sodium pyruvate, and 5% (v/v) conditioned medium containing macrophage colony stimulating factor (M-CSF) produced by CMG cells to induce differentiation to BMDMs. BMDMs were allowed to differentiate for 6 days, changing medium after four days, and then plated for experiments.

Rat cortical neurons

Cortical neurons were generated from embryonic day 18 Sprague-Dawley rats as described previously[23] and maintained at 37°C in a humidified 5% CO_2 incubator. Cells were plated onto poly-d-lysine-coated wells of black-walled 96-well plates for enzymatic assays or Seahorse XF96 plates. Cells for 96-well plate-based assays were seeded at 2.5×10^4 cells/well and maintained in maintenance medium composed of 200 μL Neurobasal (Thermo Fisher Scientific) medium supplemented with 1X B27 serum-free supplement (Thermo Fisher Scientific), 2 mM GlutaMAX

(Thermo Fisher Scientific), 100 U/ml penicillin, and 100 µg/mL streptomycin. Half the medium was

replaced every 3–4 d. Experiments in 96-well plates were conducted at day in vitro (DIV) 13–17.

Human T cells

Frozen stocks of human naïve CD4$^+$ T cells were purchased from Lonza. Cells were thawed

and permitted to recover in RPMI 1640 (Gibco, #11875-085) supplemented with 10%(v/v) FBS at

37°C, 5% CO_2 for 18-24 hr. After recovery, cells were spun at 250g for 10 min and resuspended to a

final density of 4 x 10^6 cells/mL in Agilent Seahorse XF RPMI Medium pH 7.4 (# 103576-100)

supplemented with 10 mM glucose, 2 mM glutamine and 1 mM pyruvate (XF Assay Media). 50 µL of

the naïve T-cell suspension (2 x 10^5 cells) were plated in each well of a poly-D-lysine-coated Seahorse

XF96 microplate. Plates were spun at 200g for 1 min to adhere cells to the plate. 130 µL of warmed

XF Assay Media was then added to each well, and plates were either run in the XF Analyzer or used

to measure intracellular ATP concentration with the Cell-Titer Glo Assay (Promega). ATP

measurements were taken before and after activation with magnetic Dynabeads coated with anti-

CD3/CD28 antibodies (Thermofisher, 4:1 beads/cell).

Primary cells used for aggregate H$^+$:O$_2$ values

Neonatal rat ventricular myocytes[24] and rat cortical astrocytes[25] were isolated as previously

described. Murine splenocytes were isolated from dissected spleens from C57BL/6 mice. Tissue was

mechanically dissociated in HBSS and passed through a 40 µm strainer. Cells were washed and

resuspended in red blood cells lysis buffer (Stemcell # 07850) and incubated on ice for 10 min. Cell

were washed again in HBSS, passed through a 70 µm filter, counted, and used immediately. For

Seahorse XF experiments cells were resuspended in XF Assay Media and seeded at 1.5 x 10^5

cells/well.

Mitochondrial Isolation

Hearts from C57BL/6 mice were drained of blood and homogenized with a hand-held Polytron

followed by a Potter-Elvehjem Teflon-on-glass tissue grinder in ice-cold MSHE buffer [70 mM

sucrose, 210 mM mannitol, 5 mM HEPES, 1 mM EGTA, and 0.5% (w/v) fatty acid-free BSA][26]. The

homogenate was centrifuged at 500g for 10 minutes at 4°C. The supernatant was then collected and

centrifuged at 12,000 g for 10 minutes at 4°C to obtain a mitochondrial pellet. The pellet was washed

in MSHE and centrifuged at 12,000 g for 10 minutes at 4°C, and this wash step was repeated. Finally,

the pellet was resuspended in a small volume (~20 µL) of MSHE, and the protein concentration was

determined by Bradford assay.

Cultured cells

All cell lines were obtained from American Type Culture Collection (ATCC) and cultured as suggested

by the supplier except when otherwise notified. Each formulation of medium was supplemented with

10% (v/v) fetal bovine serum (FBS), 2 mM GlutaMAX, 100 U/mL penicillin, and 100 µg/mL

streptomycin unless otherwise specified. All cells were maintained at 37°C in a humidified 5% CO_2 incubator. Cell lines were maintained in the medium listed parenthetically: A549 [DMEM/F12 (Gibco 11330); HepG2 [MEM (Gibco 11095)]; HCT116 [RPMI (Gibco 11875), C2C12 [DMEM (Gibco 11960 + 2 mM glutamine + 1 mM pyruvate)], A431 [DMEM (Gibco 11960) + 2 mM glutamine + 1 mM pyruvate), BAEC [DMEM (Gibco 11054)], BT474 [ATCC Hybri-Care Cat#46-x], H460 [RPMI (Gibco 11870), 1 mM HEPES, 4 mM glutamine, 5% FBS), HUVEC [EGM Endothelial Cell Growth Medium (Lonza CC-3124)], Jurkat,[RPMI (Gibco A10491-01)], MCF7 [RPMI (Gibco 21870 + 2 mM glutamine + 1 mM pyruvate)], MCF-10A [MEBM supplemented with Kit CC-3150 (Lonza) and 100 ng/mL cholera toxin (Sigma)], MDA-MB231 [RPMI (Gibco 21870 + 2 mM glutamine + 1 mM pyruvate)], PC12 [RPMI (ATCC 30-2001 + 2mM glutamine + 10% Horse Serum + 5% FBS], Raw 264.7 [DMEM (Gibco 11960) + 2 mM glutamine + 1 mM pyruvate].

3T3-L1 pre-adipocytes and differentiation

3T3-L1 pre-adipocytes were maintained below ~70% confluency in DMEM (Gibco 11965) supplemented with 10% (v/v) bovine calf serum (BCS), 100 U/mL penicillin, and 100 µg/mL streptomycin. For differentiation, cells were plated on Day -2 and were allowed to reach confluency over 48 hr. Differentiation was induced on Day 0 by addition of fresh medium with FBS replacing BCS and supplemented with the following: 0.5 mM IBMX, 0.25 µM dexamethasone, 1 µg/ml insulin, and 100 nM rosiglitazone. On Day 2, medium was replaced to DMEM + FBS supplemented with only 1 µg/ml insulin, and thereafter insulin was removed when replacing medium every two days.

Seahorse XF Analysis

All oxygen consumption and extracellular acidification measurements were conducted using an Agilent Seahorse XF96 or XF^{e}96 Analyzer. Experiments were conducted at 37°C and at pH 7.4 (intact cells) or 7.2 (isolated mitochondria). For experiments with cultured cells, only the inner 60 wells were used, and the outer rim was filled with 200 µL of PBS throughout the incubation to minimize variance in temperature and evaporative effects across the plate[27]. All respiratory parameters were corrected for non-mitochondrial respiration and background signal from the instrument with addition of 200 nM rotenone and 1 µM antimycin A. Where appropriate, oligomycin was used at 2 µM unless otherwise specified, and FCCP concentrations were titrated to determine an optimal concentration for a given experiment. Unbuffered DMEM assay medium is composed of DMEM (Sigma #5030; pH 7.4) supplemented with 31.6 mM NaCl, 3 mg/L phenol red, and 5 mM HEPES unless otherwise indicated.

XF analysis primary cortical neurons

All experiments with primary cortical neurons were conducted in a custom, modified Neurobasal-type medium[28] rather than DMEM to maintain appropriate osmolarity. Glutamine was omitted from the assay medium to minimize excitotoxic injury.

Acid injections and calculation of buffering power

Sequential injections of 62.5 nmol H^+ (25 µL of 1.25 mM H_2SO_4) were made using the
Seahorse XF injector ports into unbuffered DMEM assay medium (see above) supplemented with 8
mM glucose, 2 mM glutamine, 2 mM pyruvate and HEPES as indicated in the text. Values were
corroborated with matched experiments using 2.5 mM HCl made from freshly purchased stocks no
more than 2 months old. Calculations were made using a measurement microchamber volume of
2.28 µL. Step-by-step sample calculations for using the buffering power of the medium to convert
ECAR (mpH/min) to pmol H^+/min are provided in the Supplemental Worksheet.

Calculation of lactate:H^+ and H^+:O_2

Rates of medium acidification without contribution from respiration were calculated as follows:
DMEM assay medium was supplemented with 5 mM glucose and varying concentrations of 2-
deoxyglucose (2-DG) and 5 mM iodoacetate (IA) where indicated in the figure. Medium was also
supplemented with 200 nM rotenone (to block complex I), 1 µM antimycin A (to block complex III),
and 2 µM oligomycin (to block ATP hydrolysis and prevent an energy crisis). Rates were calculated
for 75 min, and the spent medium was immediately harvested from each plate and frozen at -80°C
for later calculation of lactate efflux by enzymatic assay[21].

Rates of medium acidification in the absence of glycolysis were calculated as follows: DMEM
assay medium was supplemented with 5 mM pyruvate and 5 mM glutamine (no glucose was
provided) along with 2 mM 2-DG and 50 µM IA (to block residual glycolysis and glycogenolysis). 500
U/mL carbonic anhydrase was added to ensure rapid CO_2 hydration, though it was later determined
to have a negligible effect on rates of acidification. Respiratory rates were titrated by addition of
oligomycin, FCCP (500 nM or 1 µM), and rotenone with antimycin A. The medium was also
supplemented with varying concentrations of both UK5099 and BPTES (3 nM - 3 µM for both) to
further titrate the respiratory rate. Experiments with primary cortical neurons were conducted in
custom medium lacking glutamine as described earlier.

Patient-derived glioblastoma cultures

182   Patient-derived high-grade gliomas were isolated, purified, and maintained as cell lines or
183   implanted as orthotopic xenografts according to previously published protocols[29]. Cultured cells (*in
184   vitro*) and tumor cells acutely purified from xenografts (*ex vivo*) were seeded onto Cell-Tak-coated
185   XF96 plates (25 µg/well; Corning). Cells were plated at either 4.0×10^4 or 1.0×10^5 based on cell size
186   and the initial oxygen consumption rate, and spun at 600*g* for 5 min to adhere cells to the well.

Cell activation assays

Cortical neurons were plated at 2.5×10^4 cells/well and acutely activated with 1.5 µM veratridine
added via the injector port of the instrument[30]. Where indicated, cells were pretreated with 2 nM
ouabain immediately prior to the assay. When activating the NMDA receptor, 100 µM NMDA was
acutely given via the injector port to neurons with or without 15 min. pre-treatment of 10 µM MK-801.
For these experiments, the custom assay medium was supplemented with 2 mM β-hydroxybutyrate

as substrate, as pyruvate could not be used due to complications with measuring glutamate in the
medium by a linked enzymatic assay to confirm depolarization and glutamate efflux (data not shown).
For acute, T cell activation, murine anti-CD3/CD28 Dynabeads (Gibco) were added via the injector
port as previously described[31].

TLR-mediated activation of BMDMs

BMDMs were plated at 5.0×10^4 cells/well for 48 hr. prior to cytokine treatment, which lasted
for 24 hr. unless otherwise specified. All cytokines were purchased from PeproTech, Inc. (New
Jersey, USA) and provided at the following concentrations[32]: lipopolysaccharide (LPS, 50 ng/mL
unless otherwise specified), Pam3CSK4 (Pam3, 50 ng/mL), Poly(I:C) (1 µg/mL), CL307 (100 nM),
and ODN1668 (100 nM).

$H^+:O_2$ in isolated heart mitochondria

Isolated heart mitochondria were plated in XF96 plates (1.5 µg/well) and spun 2000g for 20
min to adhere mitochondria to the well. Assays were conducted in MAS medium ([220 mM mannitol,
70 mM sucrose, 10 mM KH_2PO_4, 5 mM $MgCl_2$, 1 mM EGTA, and 2 mM HEPES (pH 7.2)]
supplemented with 0.2% (w/v) fatty acid-free BSA. For all conditions, uncoupler-stimulated respiration
was measured in the presence of 2 µM oligomycin and 4 µM FCCP to eliminate effects of ATP
synthesis or hydrolysis on the acidification rate. Rates of respiration and acidification were measured
under three conditions: (i) pyruvate/malate (P/M): 10 mM pyruvate with 1 mM malate and 2 mM
dichloroacetate; (ii) pyruvate/malate with an inhibitor mix to block various enzymes to run a truncated
TCA cycle (P/M/truncated): 10 mM pyruvate with 1 mM malate, 2 mM dichloroacetate, 60 µM
fluorocitrate (to block isocitrate dehydrogenase), 2 mM malonate (to block succinate dehydrogenase),
and 1 mM aminooxyacetate (to block transaminase activity), and (iii) succinate/rotenone (S/R): 10
mM succinate with 2 µM rotenone to consume oxygen without release of CO_2 to acidify the
experimental medium[21].

Normalization of Seahorse XF assays to total protein or cell number

Where indicated on the axis, Seahorse XF assays were normalized either to total protein or
cell number determined by *post hoc* high-content imaging. When measuring total protein, cells were
lysed in RIPA buffer [50 mM Tris, pH 7.4, 150 mM NaCl,1% (w/v) NP-40, 0.5% (w/v) sodium
deoxycholate, and 0.1% (w/v) sodium dodecyl sulfate], boiled, and assayed using the bicinchoninic
assay method (ThermoFisher).

When conducting *post hoc* cell counts, Seahorse XF plates were collected after assays,
washed with PBS, and fixed overnight with 2% (w/v) paraformaldehyde in PBS. Cells were left fixed
up to 2 weeks. On the day prior to counting, 10 ng/mL Hoescht 33342 (ThermoFisher) was added
and incubated overnight at 4°C. Cells were counted using a PerkinElmer Operetta high-content, wide-
field, fluorescence imaging system (ex./em. 361/497 nm) coupled to Columbus software (Operetta;
PerkinElmer, Waltham, MA, USA).

Enzymatic lactate measurements

Enzymatic lactate measurements were determined in the spent medium recovered after an Agilent Seahorse XF assay. Medium was brought to room temperature if previously frozen, and mixed 1:1 with a 2X assay solution of 40 U/mL lactate dehydrogenase (Sigma-Aldrich L3916), 1 M Tris (pH 9.8), 20 mM EDTA, 40 mM hydrazine (Sigma-Aldrich 309400), and 4 mM NAD^+. The reaction velocity was given by NADH fluorescence (ex./em. 340/460) and measured using a Tecan Spark multimode plate reader[21]. Measurements were taken after 2-4 min (readings were taken every 30 sec. to determine when stable values were achieved), and calibrated against known lactate concentrations (Sigma-Aldrich L7022). Importantly, lactate standards were made in assay medium matching the experimental medium to account for any effects from millimolar concentrations of pyruvate in some experimental medium. Standard curves were calculated between 0.2 and 100.0 nmol lactate and modeled with a Michaelis-Menten curve using GraphPad Prism 5.0.

Intracellular lactate measurements by GC/MS

Intracellular lactate was measured as previously described[33] using an Agilent 7890B gas chromatograph coupled to a 5977B mass spectrometer with an Agilent J&W DB-35ms column (30 m x 0.25 mm i.d. x 0.25 μm). "Day 6" BMDMs were seeded at 3.3×10^5 cells/well in 6-well tissue culture dishes. After 48 hr., cells were mock-treated or treated with TLR ligands for 24 hr. At the time of harvest and extraction, wells were quickly washed twice with ice-cold 0.9% (w/v) saline. Polar metabolites were extracted with a Folch-like extraction of 40% MeOH, 40% $CHCl_3$, and 20% H_2O with 1μg norvaline as an internal standard. The polar phase was collected and dried overnight in a refrigerated benchtop vacuum concentrator (LabConco). Dried samples were prepared for GC/MS analysis by first treating with 20 μL of 2% (w/v) methoxyamine hydrochloride ("MOX") in pyridine and incubated for 45 minutes at 37°C, followed by addition of 20 μL N-tert-butyldimethylsiyl-nmethyltrifluoroacetamide with 1% tert-butyldimethylchlorosilane for 45 minutes at 37°C to form methoxyamine-tertbutyldimethylsilyl (MOX-tBDMS) derivatives.

The GC oven was held at 100°C for 2 min, increased to 255°C at 3.5°C min^{-1}, then immediately increased to 320°C at 15°C min^{-1} and held for 3 min. Lactate was measured using selected ion monitoring (SIM) mode and integrating fragment ions (m/z 261-265) at a retention time of 12.6 min. Samples were calibrated against a standard curve (0.07 – 50 nmol; Sigma L7022). Cell numbers were quantified using identically treated matched plates. At the time of harvest for GC/MS, the matched plates were fixed with 4% paraformaldehyde for 20 min at room temperature and refrigerated. Two days later, cells were stained with Hoescht 33342 (Thermo Fisher) at 10 ng/mL overnight at 4°C. The following day, cell counts were obtained using a PerkinElmer Operetta high-content, wide-field, fluorescence imaging system (ex./em. 361/497 nm) coupled to Columbus software (Operetta; PerkinElmer, Waltham, MA, USA).

ATP Content

Cellular ATP content was measured in lysed cells with the Cell Titer Glo assay (Promega) in

96-well plates according to the manufacturer's specifications.

Quantification and statistical analysis

All statistical parameters, including the number of replicates (n), can be found in the figure

legends. Statistical analyses were performed using Graph Pad Prism 5 software for Mac OS X. Data

are presented as the mean ± SEM unless otherwise specified. Individual pairwise comparisons were

performed using two-tailed Student's t-test. For experiments involving two or more factors, data were

analyzed by one-way, repeated measures ANOVA followed by Dunnett post-hoc multiple

comparisons tests. Data were assumed to follow a normal distribution (no tests were performed).

Values denoted as follows were considered significant: *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

RESULTS

ECAR (in mpH/min) is readily converted into pmol H^+/min

A strategy to convert rates of oxygen consumption and extracellular acidification into ATP

produced from oxidative phosphorylation and glycolysis includes three sequential steps:

(1) Unit conversion to transform ECAR to pmol H^+/min so rates are directly comparable with

282   OCR measurements (in pmol O_2/min): mpH/min $\rightarrow$ pmol H^+/min;

283   (2) Adjustment of H^+ production rates to remove non-glycolytic acidification and quantitatively

284   reflect lactate efflux: pmol H^+/min $\rightarrow$ pmol lactate/min;

285   (3) Stoichiometric conversion of oxygen consumption and lactate efflux rates into ATP

286   produced from oxidative phosphorylation and glycolysis: pmol O_2/min $\rightarrow$ pmol

287   ATP_{OxPhos}/min; pmol lactate/min $\rightarrow$ pmol ATP_{Glyco}/min

First, converting ECAR into values of pmol H^+/min required calculating the buffering power of

the experimental medium. Buffering power is defined here as the pH change in the measurement

microchamber elicited by a given amount of H^+ (Fig. 1A). Importantly, sequential additional of H_2SO_4

showed the change in pH was directly proportional to the added H^+ over a range greater than 0.5 pH

units (Fig. 1B). Predictably, as the concentration of HEPES in the medium increased, there was a

smaller drop in pH elicited from the added H^+. The results demonstrate that extracellular acidification

measurements in mpH/min are readily converted to values of pmol H^+/min by applying a scalar factor

accounting for the buffering power of the medium and the volume of the measurement microchamber.

Sample calculations and step-by-step instructions for transforming ECAR into values of pmol H^+/min

are provided in the first tab of the Supplemental Worksheet.

In the absence of respiration, glycolysis is the exclusive source of measurable extracellular acidification

We then determined whether measurements of H^+ efflux could accurately measure lactate efflux after correcting for non-glycolytic acidification. In principle, there could be several reactions which cause a meaningful net pH change in the experimental medium. However, we began by testing the hypothesis that respiratory acidification (CO_2 evolution from mitochondrial dehydrogenases linked to O_2 consumption) would be the only acidifying process with large enough flux to affect the measurements on a scale comparable with glycolysis.

If respiratory acidification is indeed the only measurable source of non-glycolytic acidification, then any remaining acid production should be attributable to glycolysis when respiration is inhibited. To test this, we measured whether lactate efflux was directly proportional to H^+ production after chemical inhibition of the respiratory chain. In the presence of electron transport chain inhibitors, cellular H^+ production was gradually reduced with increasing concentrations of glycolytic inhibitors in A549 cells (Fig. 1C). We then measured the lactate content in the spent experimental medium from these samples using an enzymatic lactate assay (Fig 1D) to determine the relationship between lactate efflux and acid production in the absence of respiration (Fig. 1E). As a negative control, no measurable acidification was observed in culture pancreatic β-cells[34], which do not express appreciable levels of lactate dehydrogenase or monocarboxylate transporter-1. Using 17 different cell types comprised of primary and immortalized cultures, this analysis yielded an average lactate:H^+ ratio of 1.53 ± 0.12 (Fig. 1F; Supplemental Table 1).

Unexpectedly, this value deviated substantially from our expectations. In principle, the lactate:H^+ ratio should never be greater than 1, since the lactate efflux should never exceed the total H^+ efflux given their 1:1 stoichiometry. In fact, the theoretical lactate:H^+ should be less than 1 due to acidifying CO_2 evolution from the pentose phosphate pathway and pyruvate efflux that should contribute to total acid production but not generate lactate. We reasoned that this discrepancy could be somewhat due to the incomplete coverage of the measurement well by the fluorometric sensor of the Seahorse XF cartridge: although the enzymatic lactate measurements used medium collected from the entire well, outer portions of the well are likely undetected by the fluorometric sensor spots during the acidification measurements[35].

To test this, we scraped cells from the measurement well from either the center of the three risers directly under the sensor (creating a "donut" shape of cells in the well) or on the rim of the plate outside the three risers (leaving cells in the center untouched) (Fig. S1A). As predicted, both OCR and ECAR were significantly reduced when cells in the middle of the well were removed, but losing cells along the outer rim did not significantly affect the measurements (Fig. S1B). To ensure the results were not due to changes in cell number, the lactate:H^+ ratio was calculated for these conditions. Consistent with the sensor not seeing the outer edges of the well, lactate:H^+ was lowered

335   close to 1 when cells were removed from the outer rim, and increased beyond 8 when cells were

336   removed from center of the plate (Fig. S1C). Rather than a systemic error with the buffer capacity or

337   enzymatic measurements, the result demonstrates that incomplete sensor coverage of the

338   measurement well is a main driver of why lactate:H^+ values deviate from theoretical expectations.

340   In the absence of glycolysis, respiration is the exclusive source of measurable medium

341   acidification

Figure 1 demonstrates that in the absence of respiration, the remaining medium acidification

is tightly correlated with glycolysis. We then tested whether medium acidification would be tightly

correlated to respiration in the absence of glycolysis. We measured acidification rates in HepG2 cells

offered glucose-free medium containing pyruvate, glutamine, and the glycolytic inhibitors 2-

deoxyglucose and iodoacetate. The rates of respiration and acidification were titrated by adjusting

flux through the electron transport chain (with oligomycin, FCCP, and rotenone with antimycin A) and

blocking substrate uptake and oxidation (with varying concentrations of the mitochondrial pyruvate

carrier inhibitor UK5099 and the glutaminase inhibitor BPTES) (Figs. 2A&B). We observed a

proportional relationship between respiration and medium acidification in the absence of glycolysis

(Fig. 2C). We again extended this analysis to a panel of 21 different primary and immortalized cell

preparations to yield an average H^+:O_2 ratio of 0.38 $\pm$ 0.05 (Fig. 2D; Supplemental Table 2).

To better understand the factors that shape this value, we measured whether experimental

perturbations would change respiratory acidification in expected ways. As the complete oxidation of

different substrates yields varying H^+/O_2 ratios, we tested whether we could detect a difference in

measured H^+:O_2 from oxidation of either pyruvate (theoretical H^+:O_2 = 0.8) or glutamine (theoretical

H^+:O_2 = 0.67)[21]. Indeed, the H^+/O_2 ratio resulting from pure glutamine oxidation was lower than pure

pyruvate oxidation (Figs. S2A-C).

Experiments with isolated mitochondria further confirmed that respiratory substrates

(qualitatively) influence acidification rates in an expected manner. We tested acidification from

uncoupled respiration in rat heart mitochondria offered (i) pyruvate with malate (P/M), (ii) P/M and

inhibitors of multiple enzymes creating a truncated TCA cycle[21], and (iii) succinate with rotenone

(S/R). The results qualitatively exhibited the expected behavior: the H^+/O_2 ratio for mitochondria

oxidizing P/M increased in the presence of inhibitors which blocked downstream TCA cycle reactions

(e.g. the alkalinizing succinyl CoA synthetase reaction). Moreover, succinate oxidation in the

presence of rotenone, which will support oxygen consumption without generating CO_2, shows robust

respiratory rates with negligible changes in pH (Figs. S2 D,E).

Finally, we further validated our measurements by testing whether the forced efflux of organic

acids others than lactate would alter the H^+:O_2 ratio. Indeed, NMDA-induced efflux of glutamate in

primary cortical neurons caused a pronounced spike in the acidification rate (Fig. S2F) and H^+/O_2

ratio (Fig. S2G) that was sensitive to the NMDA-receptor antagonist MK801[28]. The result was not

attributable to glycolysis, as cells were offered exclusively β-hydroxybutyrate with glycolytic inhibitors

and lactate efflux was negligible (Fig. S2H). The demonstration that $H^+:O_2$ ratios respond in expected

ways to changes in substrate oxidation and non-glycolytic acid efflux provides additional confidence

in the measured values.

Average lactate:H^+ and $H^+:O_2$ ratios can be used to estimate cellular ATP production rates

The results from Figures 1 and 2 demonstrate that rates of H^+ production can quantify lactate

efflux by correcting for respiratory acidification and accounting for the sensor coverage of the plate

well. We therefore transformed H^+ production rates into lactate efflux rates as follows:

Eq. 1: pmol lactate/min = 1.53 * (H^+_{total} – 0.38 * $OCR_{mito.}$)

where H^+_{total} = H^+ production rates read by the instrument; 0.38 = average $H^+:O_2$ value

obtained in Fig. 2; $OCR_{mito.}$ = basal mitochondrial respiration; 1.53 = average lactate:H^+ value

obtained in Fig. 1.

To validate whether these average values could be broadly applied across different cell types, we

compared lactate efflux calculated using the XF Analyzer against an enzymatic assay in A549

adenocarcinoma cells, C2C12 myoblasts, and murine BMDMs (Fig. 3A). Prior to adjusting the

measurements as in Eq. 1, the H^+ production rate underestimated the rate of lactate efflux (comparing

leftmost bars in each graph with rightmost). But as expected, correcting Seahorse XF measurements

for respiratory acidification and sensor coverage resulted in lactate efflux values (middle bars) that

matched enzymatic measurements.

The ability to quantify lactate efflux with rates of H^+ production allows the transformation of

OCR and ECAR into rates of ATP produced by oxidative phosphorylation and glycolysis. All

necessary parameters are obtained during standard respirometry profiling (Figs. 3B & C), and sample

calculations with representative data sets are available in the Supplemental Worksheet.

The rate of mitochondrial ATP production can be estimated by applying a P/O ratio (moles of

ADP phosphorylated per mole of atomic oxygen consumed) to the respiration that drives ATP

synthesis[36], which is determined using sensitivity to the ATP synthase inhibitor oligomycin. As

detailed elsewhere, oligomycin-sensitive respiration will slightly underestimate ATP-linked respiration

(due to overestimating proton leak), though this is a small, ~10% overestimate which can be adjusted

($0.1*OCR_{LEAK}$ in Eq. 2)[37]. Moreover, assumptions must be made for the selection of P/O ratio based

on which substrate is being oxidized. Here, we use a value of P/O = 2.73 (P/O_2 = 5.45), the maximum

value for complex I-driven respiration[38] and within the range of values for complete glucose

oxidation[20]. The rate of ATP from oxidative phosphorylation can therefore be estimated as in Eq. 2:

Eq. 2: ATP_{OxPhos} = 5.45*(OCR_{ATP} + $0.1*OCR_{LEAK}$)

The rate of glycolytic ATP production can be estimated with 1:1 stoichiometry to the lactate
efflux rate (Eq. 3), and the total ATP production rate is given by the sum of Eqs. 2 & 3.

Eq. 3: $ATP_{Glyco.} = 1.53 \cdot (H^+_{TOTAL} - 0.38 \cdot OCR_{MITO})$

Eq. 4: $ATP_{Total} = ATP_{OxPhos} + ATP_{Glyco.}$

Notably, our calculation will slightly underestimate rates of glycolytic ATP, as it fails to account
for the ATP generated during glycolysis when glucose-derived pyruvate is funneled into mitochondria
for complete oxidation[20]. Accounting for this would require knowing (or assuming) the fractional
contribution of glucose oxidation to oxidative phosphorylation relative to other substrates. To test
whether this omission would appreciably affect experimental conclusions, we measured rates of ATP
production in A549 cells offered complete medium or a simple salts medium supplemented with only
glucose. Offering cells only glucose allowed calculation of this additional glycolytic ATP by applying
known stoichiometry and assuming respiration is exclusively driven by glucose oxidation with no
contribution from endogenous substrate stores. However, rates of both oxidative phosphorylation and
uncoupler-stimulated respiration were substantially limited when cells were offered medium
containing solely glucose (Figs. 3D&E). Additionally, rates of glycolytic ATP production between
groups were only marginally altered despite accounting for the cytoplasmic ATP production linked to
mitochondrial glucose oxidation (Fig. 3D). The method therefore ignores this component, favoring –
on balance – an approach where cells are offered complete DMEM to measure rates of oxidative
phosphorylation and uncoupled respiration that are not artificially restricted by substrate limitation.
425

ATP production rates show patient-derived glioma xenografts acquire an artificial glycolytic
phenotype during in vitro cell culture
A clear benefit of measuring ATP production rates is the ability to extend beyond commonly
used OCR:ECAR ratios to directly quantify changes in the balance between oxidative phosphorylation
and glycolysis. To illustrate this, we analyzed tumor cells from patient-derived glioma xenografts
either immediately harvested and purified from mice (~3 hr. isolation) or matched cells propagated in
in vitro gliomaspheres (>1 week culture period). *In vivo* evidence suggests that the enhanced aerobic
glycolysis observed in some glioma cell culture models may be driven, in part, by artifacts of cell
selection or *in vitro* growth conditions[39]. To explore this, we again demonstrated that our
measurements of lactate using H^+ efflux matched enzymatic measurements for these patient-derived
cultures (Fig. 4A). We then examined the information that can be gained from ATP production rates
as opposed to OCR:ECAR ratios. In five different cell lines, *in vitro* cultured glioma cells showed a
significantly reduced OCR:ECAR ratio compared to cells assayed immediately after purification from
corresponding murine xenografts (*ex vivo*), indicating a relative shift towards glycolysis in *in vitro*
cultured cells (Figs. 4B&C).

Quantifying ATP production rates, however, extends this qualitative analysis. Glioma cells
analyzed *ex vivo* just after isolation and purification are highly dependent on oxidative phosphorylation
(~70% of total ATP). When cultured *in vitro*, however, cells undergo a remarkable shift in metabolism,
increasing their energy demand and switching to glycolysis as the dominant ATP-generating pathway
(Figs. 4D&E). Importantly, there is only a slight trend towards increased oxidative phosphorylation,
suggesting a specific reprogramming of glycolysis acquired upon *in vitro* culture and not simply an
increased energy demand met by both pathways. The result shows that quantifying ATP production
from oxidative phosphorylation and glycolysis yields insights lost upon qualitative, ratio-based
analyses, and further supports an *in vivo* bioenergetic role of mitochondria in glioblastoma[40,41].
450
ATP production rates can measure healthy bioenergetic changes during cell activation that
are lost with static ATP measurements
We next applied this analysis to models of acute cell activation to highlight the differences
between ATP production rates and measurements of ATP content. Healthy cell activation such as
neuronal plasma membrane depolarization or T cell receptor activation should minimally affect
snapshot-in-time intracellular ATP levels, as cells will readily adapt to a new steady-state of increased
ATP turnover[1]. In primary cortical neurons, for example, addition of the Na^+-channel activator
veratridine stimulates Na^+/K^+-ATPase activity (Fig. 5A, top). Both oxygen consumption and
extracellular acidification rates increased in response to acute veratridine addition[30] (Fig. 5A, bottom).
Quantifying ATP production rates showed almost a three-fold increase in the overall rate of ATP
turnover upon depolarization, with the balance between oxidative phosphorylation and glycolysis
broadly unchanged upon activation (Fig. 5B). To demonstrate the sensitivity of the assay, we used a
sub-saturating amount of the Na^+/K^+-ATPase inhibitor ouabain (2 nM) to demonstrate the assay can
pick up small but meaningful changes in the cellular ATP demand. As expected, this substantial
change in steady-state ATP utilization did not appreciably change static ATP levels, though levels
were predictably altered in response to metabolic inhibitors (Fig. 5C).
To further demonstrate the principle that ATP production rates provide information during cell
activation lost with steady-state measurements, we measured human $CD4^+$ T cell activation in
response to anti-CD3/CD28 stimulation. Here again, acute cell activation increased rates of both OCR
and ECAR (Fig. 5D). As is well documented, T cell activation caused a pronounced increase in
glycolysis to meet the energetic and biosynthetic demands of proliferation, cytokine production, and
clonal expansion[42,43]. Nonetheless, despite a four-fold increase in glycolytic ATP production,
comparing both pathways showed oxidative phosphorylation remained the primary source of cellular
ATP production, reinforcing an important role for mitochondria during early T cell activation[44] (Fig.
5E). As was demonstrated previously for neurons, a healthy increase in the ATP demand that is
readily met by T cells did not manifest in substantially altered ATP levels (Fig. 5F). In total, the results

show real-time ATP production rates can reveal activation-associated changes in energy metabolism

that static ATP levels cannot.

ATP production rates demonstrate oxidative phosphorylation and glycolysis are independently regulated during pro-inflammatory macrophage activation

Finally, we applied our calculation to study pro-inflammatory macrophage activation, another

form of immune cell activation associated with substantial metabolic alterations. Upon classical, "type

I" activation [LPS ($\pm$ interferon-γ)], murine BMDMs profoundly remodel energy metabolism away from

oxidative phosphorylation[45] (Fig. 6A). ATP production is shifted to glycolysis and mitochondria are

repurposed to generate signals that support the pro-inflammatory response[46–48]. By measuring this

switch with a qualitative OCR:ECAR ratio, metabolic reprogramming with LPS was indistinguishable

from respiratory chain inhibition with complex I inhibitor rotenone (Fig. 6B, inset). However,

quantifying ATP production rates per cell revealed an increased energy demand associated with LPS

activation, discriminating between mitochondrial dysfunction and mitochondrial repurposing during

macrophage activation (Fig. 6B; Fig. S3).

Quantifying ATP production rates over a range of LPS concentrations further highlights the

depth of information gained from the measurements. Qualitative OCR:ECAR ratios in response to

increasing LPS concentrations (1-30 ng/mL) suggest a uniform, coordinated shift away from oxidative

phosphorylation towards glycolysis to meet ATP demands upon activation (Fig. 6C, inset). Cellular

ATP production rates, on the other hand, surprisingly reveal these two pathways are controlled

independently. Increases in glycolysis and overall ATP utilization are observed upon treatments as

low as 1 ng/mL LPS, whereas rates of mitochondrial ATP production are not compromised until [LPS]

> 10 ng/mL (Fig. 6C). Additionally, rates of maximal respiration are not compromised at low

concentrations of LPS that substantially increase glycolysis, further demonstrating that the increase

in glycolytic ATP is not simply a response to impaired mitochondrial oxidative metabolism (Fig. 6D).

Indeed, the time-dependence of energetic activation further reveals that regulation of

mitochondrial and glycolytic ATP production are disengaged: glycolysis increases within two hours of

activation while rates of oxidative phosphorylation and maximal respiration are maintained up to at

least four hours (Figs. 6E&F). The results show the hallmark metabolic changes upon pro-

inflammatory macrophage activation are a composite phenotype where increases in ATP demand

and glycolysis are controlled distinctly from reductions in oxidative phosphorylation.

Genetic and pharmacologic manipulation of MyD88 confirms glycolysis is regulated independently of oxidative phosphorylation during pro-inflammatory macrophage activation

After conducting a time-course and concentration-response profile with LPS to demonstrate

independent changes in glycolysis and oxidative phosphorylation, we then sought to examine

whether other TLR agonists could selectively enhance glycolysis. LPS is a toll-like receptor 4 (TLR4)

agonist that is linked to two distinct TLR adaptor proteins: myeloid differentiation factor 88 (MyD88)

and TIR-domain-containing adapter-inducing interferon-β (TRIF). The cell surface-localized MyD88

and endosomal TRIF link recognition of microbial products to the activation of broad genetic programs

such as the NF-κB (MyD88) or interferon regulatory factor (IRF) family of transcription factors[49]. To

better understand which of these programs are associated with specific changes in energy

metabolism, we measured ATP production rates in response to Pam3-CSK4, a triacylated lipopeptide

that activates MyD88-associated TLR2, or Poly I:C, a dsRNA analog that will activate TRIF-

associated TLR3 (Fig. 7A).

TLR2 agonism increased the glycolytic rate and the overall ATP production rate substantially

relative to both matched controls and the slight increase observed with TLR3 agonism (Figs. 7B&C).

Here again, neither TLR2 nor TLR3 compromised rates of oxidative phosphorylation (Fig. 7B, blue

bars), further demonstrating that changes to glycolysis during inflammatory activation are discrete

from changes to mitochondrial energy metabolism. TLR2-linked changes in glycolysis occurred with

a magnitude and time-scale similar to LPS (Figs. 7D & 6E), suggesting the robust increase in

glycolysis and ATP production during classical macrophage activation were primarily associated with

MyD88.

To substantiate this, we measured bioenergetics in response to TLR activation in BMDMs

harvested from $Myd88^{-/-}$ mice. As expected, the increase in glycolysis and total ATP production with

Pam3-CSK4 were entirely absent in MyD88-null BMDMs (Fig. 7E). Agonists for TLR7 and TLR9 –

both of which use the MyD88 as an adaptor protein – also elicited large changes in glycolysis that

were absent in $Myd88^{-/-}$ BMDMs (Fig. 7F). Additionally, LPS-induced increases in glycolysis were

unchanged in $Trif^{-/-}$ BMDMs but were significantly reduced in the absence of $Myd88$ (Fig. 7G), with

residual rates likely due to TRIF agonism. Taken together, the data show that MyD88-linked

transcriptional programs largely control the increase in ATP demand and glycolysis associated with

pro-inflammatory macrophage activation in a manner independent of oxidative phosphorylation. More

broadly, they highlight how quantitative ATP production rates yields insights regarding the interplay

of oxidative phosphorylation and glycolysis beyond traditional measurements of OCR and ECAR.

DISCUSSION

In line with previous reports, our empirical approach determined that glycolytic and respiratory

acidification were the only dominant contributors to the extracellular acidification rate under normal

assay conditions. Other reactions that could result in a net pH change of the medium and alter the H$^+$

efflux rate either (i) did not have with fluxes comparable to those of respiratory and glycolytic

acidification, or (ii) occurred proportionally with respiratory CO_2 evolution or lactate efflux (e.g. CO_2

evolution from the pentose phosphate pathway) and fell within a reasonable margin of error. Indeed,

the observed acidification from forced glutamate efflux upon NMDA-induced neuronal depolarization
demonstrates that our analysis would have identified if other acidifying reactions occurred at a
significant and disproportionate rate relative to lactate efflux or CO_2 evolution.

Although we demonstrated that the H^+ efflux rate can quantitatively measure lactate efflux
after accounting for respiratory CO_2, the empirical values obtained did not follow the expected
stoichiometry for either lactate:H^+ or H^+:O_2. These deviations from expected stoichiometry, however,
have theoretical and experimental support. The value of 1.53 ± 0.12 for the lactate:H^+ can be
explained in part by incomplete coverage of the plate well by the fluorometric sensor in the XF
cartridge, as cells in the center of well have a far greater effect on the readings than those on the rim.

In fact, the expected lactate:H^+ value of ~1 is obtained when cells are only within the circle
connecting the three microplate risers, roughly corresponding to the XF96 fluorometric sensor area.
The demonstration that cells directly under the measurement sensor are preferentially, if not
exclusively, read by the cartridge suggests that there is no systematic error with the enzymatic lactate
measurements. Additionally, the result implies a 1:1 lactate:H^+ ratio should never have been
expected, particularly in older, 24-well instruments where the fluorometric sensors consist of two small
spots[35] and not the larger smears found in newer 96-well instruments. Seahorse XF Wave software
now incorporates this correction into the Proton Efflux Rate (PER) parameter, accounting for the
buffering power of the medium and incorporating the geometric corrections obtained from this work.

The H^+:O_2 values from our analysis also deviate from theoretical stoichiometries expected
from the complete oxidation of known substrates. Nonetheless, the obtained values change with the
appropriate directionality and magnitude in response to oxidizing varying energy substrates in both
intact cells and isolated mitochondria. Moreover, lactate efflux rates calculated from the XF Analyzer
using an H^+:O_2 of 0.38 ± 0.05 to adjust for respiratory CO_2 matched enzymatic lactate assays and
proportionally tracked with intracellular lactate levels obtained via GC/MS, suggesting there was no
systemic error in measuring buffering power of the experimental medium.

Notably, the discrepancy between the obtained and expected H^+:O_2 values persists in isolated
mitochondria, where the experiments are highly controlled and the metabolic reactions are well
defined. In this system, the apparent H^+:O_2 values for a truncated reaction where mitochondria are
offered pyruvate and malate to generate citrate (theoretical H^+:O_2 = 1[21]) are more than 3-fold lower
than what is expected in principle. The result is likely due, in part, to the different ways the XF Wave
software calculates rates of H^+ efflux and O_2 consumption. While the reported rates of H^+ production
are a straightforward measurement, the instrument reports rates of O_2 consumption only after an
empirical correction to deconvolute the effects of back-diffusion of ambient oxygen into the
measurement microchamber[35]. This includes use of an 'apparent chamber volume' that is 3-times
greater than the actual measurement chamber volume to account for oxygen storage in the walls of
the polystyrene microplate, and the algorithmic constants were determined by matching respiration

rates obtained from synaptoneurosomes in the XF Analyzer with values obtained from a platinum-
based oxygen electrode[35].

Crucially, these corrections would necessarily account, at least partially, for respiration in
material along the rim of the plate unseen by the measurement sensor because they are clamped to
values obtained from a Clark-type electrode. However, no such corrections are made for rates of
medium acidification. As such, it should be expected that rates of H^+ production are consistently
underestimated relative to rates of O_2 consumption, and that the observed $H^+:O_2$ ratios fall below
theoretical predictions. Put simply, while software corrections adjust respiratory rates to represent
oxygen consumption reflective of the full well, this is not the case when measuring H^+ production.

Notwithstanding the deviation from expected values, the consensus values obtained for
lactate:H^+ and the $H^+:O_2$ accurately adjust for respiratory CO_2 and can be used to quantitatively reflect
lactate efflux. They can therefore be used to convert oxygen consumption and H^+ efflux rates into
estimates of ATP production as detailed in the Supplemental Worksheet. Two approaches were
considered: (i) precise calculation of P/O values by strictly defining the medium composition to better
assume what is being oxidized[20], or (ii) conduct the measurements in a more physiologically relevant
medium with multiple oxidizable substrates, an approximate P/O value, and a necessary
underestimation of glycolytic ATP as described in the results. The latter approach was chosen after
demonstrating that it does not significantly alter measurements of glycolysis but substantially
increases rates of oxidative phosphorylation and respiratory capacity, and is incorporated in the
Seahorse Analytical Software Tools.

The ability to quantify ATP production rates from each pathway provides additional insight into
how ATP utilization is distributed during cell adaptation and activation. For example, *in vivo* stable
isotope tracing has shown that *in vitro* models of glioblastoma and other cancers can overestimate
the contribution of glycolysis to energy metabolism[39,50,51]. In support of this, matched samples of
patient-derived glioblastoma cells clearly show the acquisition of an artificially high ATP demand
driven almost entirely by a five-fold increase in glycolysis in cells cultured *in vitro* relative to those
assayed immediately after isolation from murine orthotopic xenografts.

Additionally, the assay reveals that cells can adjust the balance of ATP production differently
depending on the type of activation and biological context. As an example, acute activation of the
Na^+/K^+-ATPase in primary cortical neurons substantially increases the ATP demand in a way that
does not change the proportion of ATP produced by oxidative phosphorylation and glycolysis. Despite
an acute, 2.5-fold increase in ATP production, the balance between the two pathways remained
roughly constant with about 85% of the ATP produced by mitochondria. It is sometimes stated that
the Na^+/K^+-ATPase and other plasma membrane ATPases preferentially rely on ATP produced by
glycolysis[52,53]. However, our results do not support this relationship, at least in two-dimensional cell
culture models not subject to microenvironmental gradients.

Unlike the instantaneous increase in ATP demand elicited by plasma membrane
depolarization, immune cell activation shows that cells can redistribute the balance between oxidative
phosphorylation and glycolysis when activating appropriate signaling cascades. Both T cell antigen
receptor (TCR)-linked activation in human T cells as well as TLR-mediated activation of murine
macrophages demonstrated that the increase in ATP utilization upon cell activation was increasingly
met by glycolysis relative to the naïve state. In T cells this increased glycolytic flux supports
generation of biosynthetic intermediates necessary for cell growth and expansion[42]. Given that many
types of TLR activation cause growth arrest in murine macrophages, the observed increase in
glycolysis serves a different role. For example, the increase in glycolysis is likely associated with
increased flux through the pentose phosphate pathways to support cytoplasmic NADPH production
and fuel the anti-microbial NADPH oxidase[54].

In fact, we surprisingly find that regulation of glycolysis and oxidative phosphorylation are
disengaged during the pro-inflammatory activation of macrophages. Rather than a coordinated
process where mitochondrial ATP production decreases alongside a matched increase in glycolysis,
pro-inflammatory stimuli can substantially increase glycolysis without changing rates of oxidative
phosphorylation. Genetic and pharmacologic models show that these substantive increases in
glycolysis and ATP utilization are almost entirely driven by MyD88. Understanding the MyD88-linked
molecular mechanisms by which glucose-derived pyruvate is selectively channeled away from
oxidative phosphorylation despite otherwise healthy mitochondria could aid in our understanding of
how metabolism instructs innate immunity[55].

ACKNOWLEDGEMENTS
A.S.D. is supported by National Institutes of Health (NIH) Grants R35GM138003, P30DK063491, and
P50CA092131, as well as the W.M. Keck Foundation. B.R.D. is supported by the U.S. Department
of Defense National Defense Science and Engineering (NDSEG) Fellowship Program. A.E.J. was
supported by the UCLA Tumor Cell Biology Training Program (T32 CA009056) and a Eugene V. Cota
Robles Scholarship. S.J.B. is supported by NIH Grants P01HL146358 and R01HL157710.

AUTHOR CONTRIBUTIONS
A.S.D. and N.R. supervised all aspects of the project. A.S.D., N.R., S.J.B., A.N.M., B.P.D., D.A.N.,
A.N., D.A.F., C.A. and O.S. conceptualized the study and/or designed the research plan. A.S.D.,
B.R.D., K.K.O., W.Y.H., A.E.J., P.S., D.M., L.S., G.W.R., and N.R. conducted experiments. A.S.D.,
B.R.D., K.K.O., W.Y.H., A.E.J., P.S., and N.R. analyzed the data. A.S.D., B.R.D., and N.R. prepared
the figures and wrote the manuscript with editing and approval from all authors.

DISCLOSURES

FIGURE LEGENDS

Figure 1 – Extracellular acidification is directly proportional to lactate efflux in the absence of respiration

All data are mean $\pm$ S.E.M. unless otherwise specified.

(A) Sample trace of H_2SO_4 injections to determine buffer capacity of Seahorse experimental medium. Error bars are obscured by the symbol. (n=16 technical replicates from a single experiment)

(B) Changes in pH from sequential additions of 62.5 nmol H^+ via the injector ports. The experimental medium is DMEM supplemented with 8 mM glucose, 2 mM glutamine, 2 mM pyruvate, and the indicated concentration of HEPES (pH 7.4). The microchamber volume of the 96-well XF Analyzer is 2.28 μL. Step-by-step calculations are available in the Supplemental Worksheet. (n=6 experimental replicates)

(C) The H^+ production rate is given for A549 cells treated acutely (20 minutes prior to assay) with varying concentrations of 2-deoxyglucose (2DG) or 2DG with iodoacetic acid (IAA) to block glycogenolysis. The experimental medium contains 5 mM glucose with 5 mM HEPES, 200 nM rotenone (complex I inhibitor), 1 μM antimycin (complex III inhibitor), and 2 μM oligomycin (complex V inhibitor to block 'reverse' hydrolysis of ATP upon respiratory chain inhibition). (n=6 biological replicates)

(D) Sample trace of standard curve used to measure lactate collected from experimental medium. (n=3 technical replicates)

(E) Plot of lactate efflux rates measured as in (D) against H^+ production rates measured in the Seahorse XF Analyzer. Experimental conditions and titration of the rate with glycolytic inhibitors are as in (C). Parenthetical values given in the key are the fitted slopes for each individual cell type. (n=6 biological replicates)

(F) Aggregate lactate:H^+ ratios, calculated as in (E) for 17 different cell types, yielding an average value of 1.53 $\pm$ 0.12 (mean $\pm$ standard deviation).

Figure 2 – Extracellular acidification is directly proportional to respiration in the absence of glycolysis

All data are mean $\pm$ S.E.M. unless otherwise specified.

(A) Sample kinetic trace of oxygen consumption in HepG2 cells in medium containing 5 mM pyruvate, 5 mM glutamine, 5 mM HEPES, 2 mM 2-deoxyglucose, and 50 μM iodoacetate. Where indicated, the respiratory rate was modulated with addition of UK5099 (MPC inhibitor to block pyruvate oxidation) and BPTES (glutaminase inhibitor to block glutamine oxidation). (n=8 technical replicates from a single experiment)

(B) Sample kinetic trace of H^+ production rates measured from (A). (n=8 technical replicates from a
single experiment)

(C) Plot of H^+ production rates against oxygen consumption rates for experiments conducted as in (A
& B). The rates are titrated with mitochondrial effectors (oligomycin, varying concentrations of
FCCP, rotenone/antimycin A) or combined treatment with UK5099 and BPTES. Individual points
are taken as the average values for all measurements for a particular treatment (e.g. average of
three measurements in response to initial FCCP addition). Parenthetical values given in the key
are the fitted slopes for each individual cell type. (n=6 biological replicates)

(D) Aggregate H^+:O_2 ratios, calculated as in (C) for 21 different cell types, yielding an average value
of 0.38 ± 0.05 (mean $\pm$ standard deviation).

Figure 3 – Transformation of ECAR into lactate efflux allows calculation of ATP production
rates

All data are mean $\pm$ S.E.M. unless otherwise specified.

(A) The H^+ production rate from the XF Analyzer (left bar in each graph), the calculated lactate efflux
rate from the XF Analyzer (center bar), and the lactate efflux rate measured from an enzymatic
assay (right bar) are given for A549 cells, C2C12 cells, and primary bone marrow-derived
macrophages (BMDMs). The shaded component of the H^+ production rates for each cell type is
the calculated contribution from respiratory acidification. (n$\geq$4 biological replicates)

(B) The oxygen consumption rate from a representative experiment with A549 cells is presented.
Cells are offered 8 mM glucose, 2 mM pyruvate, and 2mM glutamine in experimental medium
supplemented with 5 mM HEPES. The parameters needed to calculate the ATP production rate
are indicated with in the figure. (n=6 technical replicates)

(C) The H^+ production rate from the kinetic trace in (B) is presented. Only the initial rates are shown,
as these are the only measurements required to calculate the ATP production rate. (n=6 technical
replicates)

(D) ATP production rates calculated for A549 cells in DMEM supplemented with 8 mM glucose, 2 mM
glutamine, and 2 mM pyruvate and Krebs-Henseleit buffer supplemented with only 10 mM
glucose. (n=4 biological replicates)

(E) Maximal respiratory rates measured in response to oligomycin and FCCP for treatments as in
(D). (n=4 biological replicates)

Figure 4 – ATP production rates capture changes OCR:ECAR ratios cannot in primary cancer
cell models

All data are mean $\pm$ S.E.M. unless otherwise specified. *, $p<0.05$; **, $p<0.01$.

(A) Lactate as measured by the XF Analyzer matches enzymatic lactate measurements for a panel of patient-derived, gliomasphere (GS) cell lines after *in vitro* culture. (n≥2 biological replicates for each line)

(B) OCR:ECAR ratios for patient-derived glioblastoma cells cultured *in vitro* or assayed immediately after harvesting from mouse xenographs (*ex vivo*). (n≥2 biological replicates for each line)

(C) Collated data from (B) for statistical analysis.

(D) ATP production rates calculated from cells as in (B).

(E) Collated data from (D) for statistical analysis.

Figure 5 – ATP production rates capture healthy changes in cell activation that static ATP measurements cannot

All data are mean ± S.E.M. unless otherwise specified. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

(A) (*Top*) Schematic depicting increase in Na^+/K^+-ATPase activity by veratridine, which will depolarize the plasma membrane by increasing sodium influx, thereby triggering ouabain-sensitive increases in ATP production. (*Bottom*) Sample kinetic traces of OCR and ECAR in primary neurons upon acute addition of 1.5 μM veratridine. (n=10 technical replicates)

(B) ATP production rates from oxidative phosphorylation ($ATP_{Ox.Phos.}$, blue) and glycolysis ($ATP_{Glyco.}$, red) in response to (*left to right*) vehicle, 1.5 μM veratridine, veratridine plus 2 nM ouabain to demonstrate sensitivity of the assay, and veratridine plus 1 μM antimycin, 5 mM 2-deoxyglucose, and 1 mM iodoacetic acid to cause an energy crisis and deplete ATP levels. (n=4 biological replicates)

(C) Total ATP levels measured by luminescence after 15 min treatments as in (B). (n=4 biological replicates)

(D) Sample kinetic traces of OCR and ECAR in murine T cells acutely activated with anti-CD3/CD28. (n=4 technical replicates)

(E) ATP production rates from oxidative phosphorylation and glycolysis in response to anti-CD3/CD28. (n=3 biological replicates)

(F) Total ATP levels measured by luminescence after 30 min activation as in (E). (n=3 biological replicates)

Figure 6 – ATP production rates demonstrate glycolysis and oxidative phosphorylation are independently regulated during LPS activation of macrophages

All data are mean ± S.E.M. unless otherwise specified. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

(A) The oxygen consumption rate and H^+ production rate from a representative experiment with BMDMs in response to treatment with 50 ng/mL LPS for 24 hr. (n=5 technical replicates)

(B) The ATP production rate is calculated for BMDMs in response to 50 ng/mL LPS or 50 nM rotenone for 24 hr. (*Inset*) The OCR:ECAR ratio is presented for each condition. (n=6 biological replicates) Representative images used for cell counts are provided in the supplementary material.

(C) The ATP production rate is calculated for BMDMs in response to varying concentrations of LPS ranging from 1-30 ng/mL for 24 hr. (*Inset*) The OCR:ECAR ratio is presented for each condition. (n=5 biological replicates)

(D) FCCP-stimulated, maximal respiratory rates from treatments as in (C). (n=5 biological replicates)

(E) The ATP production rate is calculated for BMDMs in response to treatment for 2 or 4 hr. with 50 ng/mL LPS. (*Inset*) The OCR:ECAR ratio is presented for each condition. (n=4 biological replicates)

(F) FCCP-stimulated, maximal respiratory rates from treatments as in (E). (n=4 biological replicates)

Figure 7 – Calculation of lactate efflux and ATP production rates demonstrate MyD88 increases glycolysis independently of oxidative phosphorylation

All data are mean $\pm$ S.E.M. unless otherwise specified. *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$.

(A) Sample kinetic trace of oxygen consumption (*left*) and H^+ production (*right*) for macrophages treated with the TLR2 agonist Pam3CSK4 ("Pam3," 50 ng/mL) or the TLR3 agonist polyinosine-polycytidylic acid ("Poly I:C," 1 μg/mL) for 24 hr. (n=5 technical replicates)

(B) Aggregate ATP production rates for treatments as in (A). (n=8 biological replicates)

(C) Intracellular lactate as measured by GC/MS for treatments as in (A). (n=5 biological replicates)

(D) Lactate efflux rates measured by the XF Analyzer in response to 50 ng/mL Pam3 for 2 and 4 hr. treatment. (n=4 biological replicates)

(E) ATP production rates from BMDMs isolated from control or *Myd88$^{-/-}$* mice in response to treatment with 50 ng/mL Pam3 for 24 hr. (n=4 biological replicates)

(F) Lactate efflux rates in response to 24 hr. treatment with 100 nM CL307 (a TLR7 agonist) and 100 nM ODN1668 (a TLR9 agonist) in BMDMs isolated from control or *Myd88$^{-/-}$* mice. (n=4 biological replicates)

(G) Lactate efflux rates in BMDMs isolated from control, *Myd88$^{-/-}$, or Trif$^{-/-}$* mice in response to 50 ng/mL LPS (n=4 biological replicates).

REFERENCES

1. Brand, M. D. & Nicholls, D. G. Assessing mitochondrial dysfunction in cells. *Biochemical*
Journal (2011) doi:10.1042/BJ20110162.

2. Tantama, M. & Yellen, G. Imaging changes in the cytosolic ATP-to-ADP ratio. *Methods*
Enzymol. **547**, 355–371 (2014).

3. Lunt, S. Y. & Vander Heiden, M. G. Aerobic glycolysis: meeting the metabolic requirements
of cell proliferation. *Annu. Rev. Cell Dev. Biol.* **27**, 441–464 (2011).

4. DeBerardinis, R. J. & Chandel, N. S. We need to talk about the Warburg effect. *Nat. Metab.*
2, 127–129 (2020).

5. Sullivan, W. J. *et al.* Extracellular Matrix Remodeling Regulates Glucose Metabolism through
TXNIP Destabilization. *Cell* **175**, 117-132.e21 (2018).

6. Shin, B. *et al.* Mitochondrial Oxidative Phosphorylation Regulates the Fate Decision between
Pathogenic Th17 and Regulatory T Cells. *Cell Rep.* **30**, 1898-1909.e4 (2020).

7. Ducker, G. S. & Rabinowitz, J. D. One-Carbon Metabolism in Health and Disease. *Cell*
Metab. **25**, 27 (2017).

8. Lewis, C. A. *et al.* Tracing compartmentalized NADPH metabolism in the cytosol and
mitochondria of mammalian cells. *Mol. Cell* **55**, 253–263 (2014).

9. Gu, W. *et al.* Glycolytic Metabolism Plays a Functional Role in Regulating Human Pluripotent
Stem Cell State. *Cell Stem Cell* **19**, 476–490 (2016).

10. Pavlova, N. N. & Thompson, C. B. The Emerging Hallmarks of Cancer Metabolism. *Cell*
Metab. **23**, 27–47 (2016).

11. Van den Bossche, J., O'Neill, L. A. & Menon, D. Macrophage Immunometabolism: Where
Are We (Going)? *Trends in Immunology* (2017) doi:10.1016/j.it.2017.03.001.

12. Chen, C.-T., Shih, Y.-R. V., Kuo, T. K., Lee, O. K. & Wei, Y.-H. Coordinated changes of
mitochondrial biogenesis and antioxidant enzymes during osteogenic differentiation of human
mesenchymal stem cells. *Stem Cells* **26**, 960–968 (2008).

13. Wilson-Fritch, L. *et al.* Mitochondrial biogenesis and remodeling during adipogenesis and in
response to the insulin sensitizer rosiglitazone. *Mol. Cell. Biol.* **23**, 1085–1094 (2003).

14. Divakaruni, A. S., Paradyse, A., Ferrick, D. A., Murphy, A. N. & Jastroch, M. Analysis and
interpretation of microplate-based oxygen consumption and pH data. in *Methods in*
Enzymology (2014). doi:10.1016/B978-0-12-801415-8.00016-3.

15. Moyes, C. D., Mathieu-Costello, O. A., Tsuchiya, N., Filburn, C. & Hansford, R. G.
Mitochondrial biogenesis during cellular differentiation.
https://doi.org/10.1152/ajpcell.1997.272.4.C1345 **272**, (1997).

16. Pelletier, M., Billingham, L. K., Ramaswamy, M. & Siegel, R. M. Extracellular flux analysis to
monitor glycolytic rates and mitochondrial oxygen consumption. in *Methods in Enzymology*

(2014). doi:10.1016/B978-0-12-416618-9.00007-8.

17.   McConnell, H. M. *et al.* The cytosensor microphysiometer: biological applications of silicon

834   technology. *Science* **257**, 1906–1912 (1992).

835   18.   Newell, K., Franchi, A., Pouyssegur, J. & Tannock, I. Studies with glycolysis-deficient cells

836   suggest that production of lactic acid is not the only cause of tumor acidity. *Proc. Natl. Acad.*

837   *Sci. U. S. A.* **90**, 1127–1131 (1993).

838   19.   Gillies, R. J., Robey, I. & Gatenby, R. A. Causes and consequences of increased glucose

839   metabolism of cancers. *J. Nucl. Med.* **49 Suppl 2**, (2008).

840   20.   Mookerjee, S. A., Gerencser, A. A., Nicholls, D. G. & Brand, M. D. Quantifying intracellular

841   rates of glycolytic and oxidative ATP production and consumption using extracellular flux

842   measurements. *J. Biol. Chem.* (2017) doi:10.1074/jbc.M116.774471.

843   21.   Mookerjee, S. A., Goncalves, R. L. S., Gerencser, A. A., Nicholls, D. G. & Brand, M. D. The

844   contributions of respiration and glycolysis to extracellular acid production. *Biochim. Biophys.*

845   *Acta* **1847**, 171–181 (2015).

846   22.   Hsieh, W. Y., Williams, K. J., Su, B. & Bensinger, S. J. Profiling of mouse macrophage

847   lipidome using direct infusion shotgun mass spectrometry. *STAR Protoc.* **2**, (2020).

848   23.   Kushnareva, Y. E., Wiley, S. E., Ward, M. W., Andreyev, A. Y. & Murphy, A. N. Excitotoxic

849   injury to mitochondria isolated from cultured neurons. *J. Biol. Chem.* **280**, 28894–28902

850   (2005).

851   24.   Rubio, M. *et al.* Cardioprotective stimuli mediate phosphoinositide 3-kinase and

852   phosphoinositide dependent kinase 1 nuclear accumulation in cardiomyocytes. *J. Mol. Cell.*

853   *Cardiol.* **47**, 96–103 (2009).

854   25.   Kim, H. J. & Magrané, J. Isolation and culture of neurons and astrocytes from the mouse

855   brain cortex. *Methods Mol. Biol.* **793**, 63–75 (2011).

856   26.   Kubli, D. A. *et al.* Parkin protein deficiency exacerbates cardiac injury and reduces survival

857   following myocardial infarction. *J. Biol. Chem.* **288**, 915–926 (2013).

858   27.   Lundholt, B. K., Scudder, K. M. & Pagliaro, L. A simple technique for reducing edge effect in

859   cell-based assays. *J. Biomol. Screen.* **8**, 566–570 (2003).

860   28.   Divakaruni, A. S. *et al.* Inhibition of the mitochondrial pyruvate carrier protects from

861   excitotoxic neuronal death. *J. Cell Biol.* (2017) doi:10.1083/jcb.201612067.

862   29.   Gosa, L., Ta, L. & Nathanson, D. A. Processing of Primary Patient Tumors and Subsequent

863   Generation of Primary Cell Lines. *Methods Mol. Biol.* **1897**, 425–431 (2019).

864   30.   Choi, S. W., Gerencser, A. A. & Nicholls, D. G. Bioenergetic analysis of isolated

865   cerebrocortical nerve terminals on a microgram scale: Spare respiratory capacity and

866   stochastic mitochondrial failure. *J. Neurochem.* (2009) doi:10.1111/j.1471-

867   4159.2009.06055.x.

31. Gubser, P. M. *et al.* Rapid effector function of memory CD8+ T cells requires an immediate-
early glycolytic switch. *Nat. Immunol.* **14**, 1064–1072 (2013).

32. Hsieh, W. Y. *et al.* Toll-Like Receptors Induce Signal-Specific Reprogramming of the
Macrophage Lipidome. *Cell Metab.* (2020) doi:10.1016/j.cmet.2020.05.003.

33. Cordes, T. & Metallo, C. M. Quantifying intermediary metabolism and lipogenesis in cultured
mammalian cells using stable isotope tracing and mass spectrometry. in *Methods in*
Molecular Biology (2019). doi:10.1007/978-1-4939-9236-2_14.

34. Zhao, C., Wilson, M. C., Schuit, F., Halestrap, A. P. & Rutter, G. A. Expression and
distribution of lactate/monocarboxylate transporter isoforms in pancreatic islets and the
exocrine pancreas. *Diabetes* **50**, 361–366 (2001).

35. Gerencser, A. A. *et al.* Quantitative microplate-based respirometry with correction for oxygen
diffusion. *Anal. Chem.* (2009) doi:10.1021/ac900881z.

36. Brand, M. D. The efficiency and plasticity of mitochondrial energy transduction. *Biochem.*
Soc. Trans. **33**, 897–904 (2005).

37. Affourtit, C. & Brand, M. D. Measuring mitochondrial bioenergetics in INS-1E insulinoma
cells. *Methods Enzymol.* **457**, 405–424 (2009).

38. Watt, I. N., Montgomery, M. G., Runswick, M. J., Leslie, A. G. W. & Walker, J. E.
Bioenergetic cost of making an adenosine triphosphate molecule in animal mitochondria.
Proc. Natl. Acad. Sci. U. S. A. **107**, 16823–16827 (2010).

39. Marin-Valencia, I. *et al.* Analysis of Tumor Metabolism Reveals Mitochondrial Glucose
Oxidation in Genetically Diverse Human Glioblastomas in the Mouse Brain In Vivo. *Cell*
Metab. **15**, 827–837 (2012).

40. Maher, E. A. *et al.* Metabolism of [U-13 C]glucose in human brain tumors in vivo. *NMR*
Biomed. **25**, 1234–1244 (2012).

41. Mashimo, T. *et al.* Acetate is a bioenergetic substrate for human glioblastoma and brain
metastases. *Cell* **159**, 1603–1614 (2014).

42. Maciver, N. J., Michalek, R. D. & Rathmell, J. C. Metabolic Regulation of T Lymphocytes.
http://dx.doi.org/10.1146/annurev-immunol-032712-095956 **31**, 259–283 (2013).

43. Buck, M. D., Sowell, R. T., Kaech, S. M. & Pearce, E. L. Metabolic Instruction of Immunity.
Cell (2017) doi:10.1016/j.cell.2017.04.004.

44. Sena, L. A. *et al.* Mitochondria are required for antigen-specific T cell activation through
reactive oxygen species signaling. *Immunity* **38**, 225–236 (2013).

45. Van den Bossche, J. *et al.* Mitochondrial Dysfunction Prevents Repolarization of
Inflammatory Macrophages. *Cell Rep.* (2016) doi:10.1016/j.celrep.2016.09.008.

46. Mills, E. L. *et al.* Succinate Dehydrogenase Supports Metabolic Repurposing of Mitochondria
to Drive Inflammatory Macrophages. *Cell* (2016) doi:10.1016/j.cell.2016.08.064.

47.   West, A. P. *et al.* TLR signalling augments macrophage bactericidal activity through
mitochondrial ROS. *Nature* (2011) doi:10.1038/nature09973.

48.   Lampropoulou, V. *et al.* Itaconate Links Inhibition of Succinate Dehydrogenase with
Macrophage Metabolic Remodeling and Regulation of Inflammation. *Cell Metab.* (2016)
doi:10.1016/j.cmet.2016.06.004.

49.   O'Neill, L. A. J., Golenbock, D. & Bowie, A. G. The history of Toll-like receptors-redefining
innate immunity. *Nature Reviews Immunology* (2013) doi:10.1038/nri3446.

50.   Faubert, B. *et al.* Lactate Metabolism in Human Lung Tumors. *Cell* **171**, 358-371.e9 (2017).

51.   Momcilovic, M. *et al.* In vivo imaging of mitochondrial membrane potential in non-small-cell
lung cancer. *Nature* (2019) doi:10.1038/s41586-019-1715-0.

52.   Epstein, T., Xu, L., Gillies, R. J. & Gatenby, R. A. Separation of metabolic supply and
demand: aerobic glycolysis as a normal physiological response to fluctuating energetic
demands in the membrane. *Cancer Metab.* **2**, (2014).

53.   Sepp, M. *et al.* Tight coupling of Na+/K+-ATPase with glycolysis demonstrated in
permeabilized rat cardiomyocytes. *PLoS One* **9**, (2014).

54.   Cathcart, M. K. Regulation of Superoxide Anion Production by NADPH Oxidase in
Monocytes/Macrophages: Contributions to Atherosclerosis. *Arteriosclerosis, Thrombosis, and*
Vascular Biology (2004) doi:10.1161/01.ATV.0000097769.47306.12.

55.   Palsson-Mcdermott, E. M. *et al.* Pyruvate Kinase M2 Regulates Hif-1α Activity and IL-1β
Induction and Is a Critical Determinant of the Warburg Effect in LPS-Activated Macrophages.
Cell Metab. **21**, 65–80 (2015).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7