Multi-marker DNA metabarcoding detects suites of environmental gradients
from an urban harbour

Chloe V. Robinson[1,2], Teresita M. Porter[1], Katie M. McGee[3], Megan McCusker[3],
Michael T.G. Wright[1] & Mehrdad Hajibabaei[1]*

[1] Centre for Biodiversity Genomics and Department of Integrative Biology,
University of Guelph, Guelph, Ontario, Canada N1G 2W1

[2] Whales Initiative, Ocean Wise Conservation Association, Victoria, British
Columbia, Canada V8V 4Z9

[3] Environment and Climate Change Canada, Burlington, Ontario, Canada L7S
1A1

*Corresponding author: Mehrdad Hajibabaei

Email: mhajibab@uoguelph.ca

Abstract

There is increasing need for biodiversity monitoring, especially in places where
potential anthropogenic disturbance may significantly impact ecosystem health.
We employed a combination of traditional morphological and bulk
macroinvertebrate metabarcoding analyses to benthic samples collected from
Toronto Harbour (Ontario, Canada) to compare taxonomic and functional
diversity of macroinvertebrates and their responses to environmental gradients.
At the species rank, sites assessed using COI metabarcoding showed more
variation than sites assessed using morphological methods. Depending on the
assessment method, we detected gradients in magnesium (morphological taxa),
ammonia (morphological taxa, COI sequence variants), pH (18S sequence
variants) as well as gradients in contaminants such as metals (COI & 18S
sequence variants) and organochlorines (COI sequence variants). Observed
responses to contaminants such as aromatic hydrocarbons and metals align with
known patchy distributions in harbour sediments. We determined that the
morphological approach may limit the detection of macroinvertebrate responses
to lake environmental conditions due to the effort needed to obtain fine level
taxonomic assignments necessary to investigate responses. DNA
metabarcoding, however, need not be limited to macroinvertebrates, can be
automated, and taxonomic assignments are associated with a certain level of
accuracy from sequence variants to named taxonomic groups. The capacity to
detect change using a scalable approach such as metabarcoding is critical for
addressing challenges associated with biodiversity monitoring and ecological
investigations.

62 **Keywords:** Biomonitoring, metabarcoding, eDNA, bioindicators, COI mtDNA,
63 18S rRNA

Introduction

Ecosystem degradation is one of the leading causes of biodiversity decline in aquatic realms [1]. Freshwater degradation can be observed as physical changes to habitat morphology, hydrological alterations and changes to biogeochemistry of water and sediment [2,3]. Lakes in particular are more susceptible to the consequences of shoreline developments and loading of nutrients, including phosphorous and nitrogen, than other freshwater habitats [4,5]. These lake stressors often cause a reduction in littoral habitats for submerged macrophytes, changes to sediment composition, and increased levels of eutrophication [4–7], resulting in loss of lake biodiversity [2]. Restoring degraded lake ecosystems typically involves changing the ecological fate of the system towards an ecologically-sound status, where function is retained [1]. To achieve this, stressors need to be first identified and then managed via intervention and ecosystem management techniques to enable recovery and restoration [1,5,8].

Toronto Harbour is located on the north shore of Lake Ontario, directly south of the City of Toronto and receives water from one of the most highly urbanized and industrialized areas in the Great Lakes. Contaminants from stormwater runoff, spills, and chemical input to sewers from industries and residences have contributed to severely degraded water and environmental health [9–12]. In 1994, eight beneficial uses in Toronto Harbour were identified as impaired (beneficial use impairments; BUIs [12],[13]. In 1987, Toronto Harbour was designated as an Area of Concern (AOC) [13] and since then, there has been an increase in biological monitoring in the harbour [9,14,15].

As of 2016, results of Remedial Action Plan (RAP) activities resulted in re-classification of two of the eight identified BUIs ('Not Impaired'), however "Loss of Fish and Wildlife Habitat' remains impaired [13,16]. Over the last 30 years, water quality, sediment quality and the quantity and condition of terrestrial and aquatic habitats have improved considerably, resulting in the preparation to delist

Toronto Harbour as an AOC [13]. In Toronto Harbour, the RAP investigates trends

in benthic macroinvertebrate diversity by equating a higher diversity to a healthier

system [16], with diversity metrics derived solely from morphological analyses [17].

The challenges of morphological approaches compared to a DNA-based

identification approach for monitoring macroinvertebrates have been highlighted

previously (e.g. low taxonomic resolution and high processing costs) [18–23].

DNA extracted from bulk benthos samples has been transformative for routine

benthic macroinvertebrate biomonitoring [24] and for determining gradients in

freshwater environmental conditions [18,25–28]. When taxonomic lists and

corresponding the Hilsenhoff Biotic Index (HBI) tolerance values are paired with

ecological information, such as functional feeding guilds (FFG), metabarcoding

analysis can provide in-depth understanding of ecological function and

responses of macroinvertebrates to different environmental conditions

[19,28,29,30,31].

In this study, we hypothesized that the addition of eDNA metabarcoding will

provide a more taxonomically comprehensive biodiversity measure for focal

bioindicator macroinvertebrates, which will be reflected in finer-scale

environmental assessment. By comparing diversity and functional

macroinvertebrate metrics using eDNA metabarcoding and morphological

assessment methods we assess the responses of these metrics across

environmental gradients. Through this investigation, we aim to determine the

most effective approach for long-term monitoring of benthic communities in

Toronto Harbour and similar systems.

Results

A map showing sampling stations in Toronto Harbour and associated

gradients in water physical-chemical and sediment contaminants is shown in

Figure 1 (Table S1). From these samples, a total of 18,790,832 x 2 paired-end sequence reads were generated for 75 samples and 16 controls. After read pairing, trimming, and denoising (sequence error, chimera, and pseudogene removal for COI) we generated a total of 9,329 exact sequence variants (ESVs) (Table S2).

Figure 1. Toronto Harbour and associated abiotic and biotic gradients. Combined sewage outflows (black triangles) and the Don River (blue triangle) flows into the Toronto Harbour (Ontario, Canada). Sampling stations are shown in the legend. Inset map (top left) shows map area (black box) within respect to the Great Lakes region. Scale bar is in kilometers. Three of the most influential environmental variables are shown along with diversity metrics with the largest responses. Abbreviations: water physical-chemical features (W), sediment contaminants (S), macroinvertebrate (mi). Map tiles by Stamen Design, under CC BY 3.0. Data by OpenStreetMap, under ODbL.

Overall, we found that taxonomic assignment resolution was finer using COI metabarcoding compared with other methods (Figure S1a). Sampling effort assessed using rarefaction found that sequencing depth was appropriate for the metabarcoding samples, and all curves reached saturation, however for samples identified by morphology, curves continue to rise indicating that further sampling would have identified additional families (Figure S1b).

Macroinvertebrate diversity metrics measured using morphology or metabarcoding

Biodiversity was compared between morphological and metabarcoding approaches (Figure 2a). Macroinvertebrate species richness was higher using COI metabarcoding (median 15 species per station) compared with morphological methods (9 species per station) (Wilcox test, p-value > 0.00027) (Figure 2a). 18S metabarcoding detected an even higher diversity of genera (82 genera per station).

Figure 2. Alpha and gamma diversity detected is higher using DNA metabarcoding compared with morphological methods. In a) alpha diversity is compared across methods showing genus richness for (non-macroinvertebrate) eukaryotes sampled using 18S metabarcoding and species richness for macroinvertebrate sampling using COI and morphological sampling. In b) gamma diversity is compared showing the total number of 18S eukaryote genera and total number of species sampled using COI and morphological methods in Toronto Harbour.

The variance between the macroinvertebrate species detected using COI and morphological methods was similar for many stations, however, COI metabarcoding detected divergent communities at several stations (Figure 3a). The first two axes of the PCA plot explains 42.8% of the variance in these communities. These differences were correlated with the relative read abundance of *Limnodrilus hoffmeisteri*, *Tubifex tubifex*, and *Potamothrix vejdovskyi*. When we compared community composition using Bray Curtis dissimilarities, most locations appeared similar to each other when using morphological taxa (stress=0.09, R^2=0,98) and ammonia (r = 0.31, p = 0.035) and magnesium (r = 0.35, p = 0.026) help explain the variance of these communities (Figure 3b). When we compared communities using COI sequence variants, locations showed more separation with Toronto Island and Bathurst Quay locations clustering separately from the rest (stress = 0.14, R^2 = 0.86) and ammonia (r = 0.51, p = 0.002), metals (r = 0.38, p = 0.005), and organochlorines (r = 0.38, p = 0.005) help explain the variance of these communities (Figure 3c). When we compared communities using 18S sequence variants, most locations

appeared similar, but Bathurst Quay and Keating Channel locations clustered
separately (stress = 0.06, R2 = 0.99) and pH (r = 0.43, p = 0.004) and metals (r =
0.40, p = 0.005) help explain the variance of these communities (Figure 3d).

Figure 3.   DNA metabarcoding helps distinguish between sampling
locations in Toronto Harbour. Communities are compared using a) principal
components analysis (PCA) at the species rank for COI and morphological
sampling, b) non-metric multidimensional scaling (NMDS) using morphological
taxa identified to a species rank when possible, c) NMDS using
macroinvertebrate COI sequence variants, and d) NMDS using eukaryote (non-
macroinvertebrate) 18S sequence variants. For PCA, COI samples are shown in
grey circles and morphological samples in white squares and the most strongly
correlated species vectors were added. For NMDS plots, centroids for each
location are plotted for each sampling method using Bray Curtis dissimilarities,
and environmental vectors were added if correlations were at least 0.30 with a p-
value < 0.05.

We also compared the top 5 most abundant (non-macroinvertebrate) eukaryote families using 18S metabarcoding with the top 5 most abundant families detected using both morphological methods and COI metabarcoding (Figure 4a). Although we compare the relative abundance of individuals sampled using morphological methods with the relative read abundance of sequence

variants, it's important to acknowledge that read abundances reflect the specificity of our primers, specimen size, as well as abundance in samples. In terms of detections, COI metabarcoding detects some but not all the macroinvertebrate families detected using morphology. The median relative abundance of Naididae, the most abundant group using either method, was higher using morphological methods (90.6%) compared to COI metabarcoding (31.8%) (Wilcox test, p-value = 3.6e-10). The median relative abundance of Chironomidae was higher using COI (9.2%) compared to morphological methods (4.6%) (Wilcox test, p-value = 0.04). Overall, macroinvertebrate communities at the family rank were positively correlated (Pearson, 70.1%,p-value ~ 0, 95% confidence interval 61% - 78%).

Figure 4. Comparison of major taxonomic and functional groups detected across sampling methods. We compare a) the 5 most abundant families of eukaryotes (non-vertebrate 18S) and macroinvertebrates (COI and morphology) and b) macroinvertebrate functional feeding guilds. Relative read abundance is presented for DNA-derived data and relative abundance of individuals is presented for morphology-derived data. Abbreviations: predators (PR), collector-gatherer (CG), herbivore (HB), shredders (SH), collector-filterers (CF), parasites (PA).

Overall, multi-marker metabarcoding recovers a greater diversity of taxa than morphological methods even when only considering macroinvertebrate taxa in the phyla Arthropoda, Annelida, Mollusca, Cnidaria, and Platyhelminthes for a fair comparison with morphological methods (Table S3). Overall, we detected 47 and 88 unique macroinvertebrate species (using morphological methods versus

COI metabarcoding), 77 and 79 genera, and 30 and 61 families in total across all sampled stations. When considering other (non-macroinvertebrate) eukaryotes detected using 18S metabarcoding, we detected a further 271 genera and 111 families from 36 eukaryote phyla with the most diverse phyla detected from the Ciliophora (105 taxa), Cercozoa (64), Ascomycota (Fungi, 61), Basidiomycota (Fungi, 52 taxa), and the Chytridiomycota (42 taxa).

Macroinvertebrate functional metrics using metabarcoding and morphological methods

Functional feeding groups, a classification system based on how macroinvertebrates acquire food, allowed us to assess whether ecological function varies independently of taxonomic shifts. At each station, the proportion of macroinvertebrate reads (COI metabarcoding) or individuals (morphological) assigned to a family with a particular functional feeding guild(s) was assessed (Figure 4b). The relative abundance of FFGs between methods were significantly different for parasites (Wilcox test, = 0. 3.8e-05), predators (p.adj = 0. 1.8e-05), and 'other' (p.adj = 0.014). The median relative abundance of parasites was ~0% using morphological methods and 0.03% using COI metabarcoding. The median relative abundance of predators was 97% using morphological methods and 11% using COI metabarcoding. The mean relative abundance of 'other' was 2% using morphological methods and 0.2% using COI metabarcoding. Functional communities detected using both methods were not found to be correlated (Pearson, 0.08, p-value = 0.31, 95% confidence interval -0.07 - 0.22), though this may be due to our difficulty assigning function to macroinvertebrate families detected using COI metabarcoding methods. The three most abundant predator families using COI metabarcoding was Naididae (422 sequence variants, 284,851 reads), Chironomidae (81 sequence variants, 96,261 reads), and Bosminidae (34 sequence variants, 23,495 reads); and using

morphological methods was Naididae (30 taxa, 30,163 individuals), Chironomidae (41 taxa, 322 individuals), Dreissenidae (3 taxa, 2241 individuals).

We used hierarchical partitioning to identify environmental variables that explain the variance in diversity and functional metrics (Table S5, Table S6). Sediment contaminants explain more variation in diversity metrics and water physical-chemical features tend to explain more variation in functional metrics (Figure 5). The most influential environmental parameters are metals, aromatic hydrocarbons, and ammonia (Figure 1). Metals make a significant independent contribution explaining 62% of the variation in macroinvertebrate species richness (COI), 55% of the variation in the relative abundance of Thoracosphaeraceae (18S), and 45% of the variation in Haptoria (18S). Aromatic hydrocarbons make a significant independent contribution explaining 62% of the variation in Candonidae (COI), 57% of the variation in Cyprididae (COI), and 45% of the variation in Limnesiidae (morphological). Ammonia makes a significant independent contribution explaining 56% of the variation in Bosminidae (COI), 56% of the variation in Naididae (COI), 50% of the variation in Candonidae (COI), 49% of the variation in Thoracosphaeraceae (18S), and 47% of the variation in Hydridae (morphological). Only water physical-chemical features explain variation in functional metrics. Ammonia makes a significant independent contribution explaining 57% of the variance in parasites detected using morphological methods. Temperature makes significant independent contributions explaining 30-33% of the variation in collector-gatherers and shredders.

Figure 5. Sediment contaminants explain more variation in diversity metrics and water physical chemical features explain more variation in functional metrics. Predictors that make significant (p-value < 0.05)

independent contributions explaining variance in diversity metric response variables (left panel) and functional metric response variables (right panel).

Discussion

The process of monitoring benthic macroinvertebrate communities has been the focus of freshwater health assessments across the world. Morphological-based approaches, which were once the only method of determining richness, are rapidly being complemented with DNA-based methodologies such as metabarcoding, greatly increasing the taxonomic information obtained from benthic samples [21,21,32–34]. In this study, we have demonstrated the benefits of

employing eDNA metabarcoding to lake-based macroinvertebrate biodiversity
assessments.

Previously, morphological-based biomonitoring in Toronto Harbour had focused
on overall richness as the determining factor of lake health [15,16,35]. While this can
be informative as to the overall quantity of macroinvertebrate taxa present in the
lake, it neglects details on the composition of taxa and proportions of bioindicator
groups in relation to other taxa. Species diversity often reduces with degradation
of aquatic environments, with more tolerant species dominating species
compositions and composition of macroinvertebrates [36]. Therefore, the concept
of higher total richness translating to improvements in water quality and lake
health are misleading, as measures of community composition, structure and
diversity are also required to understand changes in environmental gradients [36].
COI metabarcoding also provides finer level taxonomic assignments compared
with morphological methods as has been shown in a previous study for fishes [37].

In addition, the confidence levels for morphological macroinvertebrate
identification are scarcely reported and classifications can vary between
taxonomists [18,38], meaning likelihood of misidentified or missed taxa is not taken
into consideration [21]. Both DNA metabarcoding (i.e., finer resolution taxonomic
assignments, ability to detect microscopic taxa) and morphological methods (i.e.,
absolute abundance counts) provide complementary information that can
contribute and support macroinvertebrate community information for further
ecological analyses. In addition to the typical benthic macroinvertebrates
identified using morphology, DNA metabarcoding enabled accurate detection of

additional phyla, using more inclusive markers as used in this study. In this study,

the detection of unique taxa via 18S sequencing enabled us to clearly distinguish

stations, highlighting that non-macroinvertebrate groups can also provide insights

towards understanding of spatial community dynamics in lake systems [39].

The 18S marker enabled us to detect fungal groups expected to be found from

aquatic environments such as the Chytridiomycota, Blastocladiomycota, and

Cryptomycota (Rozellomycota). We even detected members of the

Archaeorhizomycota (12 sequence variants, 117 reads), a group of fungi with a

global distribution largely known only from marker gene studies of soil with only

two species in culture [40,41]. The application of this marker also resulted in

detection of several fungal taxa that we would expect to find from benthic

samples such as known parasites of arthropods/insects: *Cordyceps*,

Coelomomyces, Labulomycetales, and Cryptomycota.

Compared to DNA methods, morphological methods detected negligible amounts

of aquatic bioindicator taxa, specifically Ephemeroptera (HBI= 0-2; excellent) and

Trichoptera (HBI= 0-4; very good). COI metabarcoding detected Ephemeroptera

and Trichoptera more often, but the relative abundance of these taxa were

relatively low. Environmentally tolerant taxa (i.e. those with HBI score of 6-10),

were detected more so with DNA metabarcoding methods compared to

morphology. For example, species/genera including, *Limnodrilus hoffmeisteri*

(HBI= 10; very poor), and *Tubifex tubifex* (HBI = 10; very poor), and *Hydra* (HBI =

5; poor) were detected exclusively by DNA methods, whereas *Potamothrix*

vejdovskyi (HBI= 8; poor) was detected exclusively with morphological methods.

Naididae as a family are highly tolerant to organic pollution, scoring between 6

and 10 on the HBI index [42]. More families, genera and species from extreme

ends of the tolerance index were detected using DNA metabarcoding,

highlighting the ability of this method to detect presence of these important taxa.

Beyond taxonomic metrics, mapping functional diversity across lake
environments can improve our understanding of lake ecosystem integrity [19,43–45].
For example, the presence of diverse macroinvertebrate predators detected in
Toronto Harbour may indicate that there are sufficient resources to support a
stable multi-level food web [43]. Overall, morphological methods performed very
well with regards to their ability to identify a diverse array of functional groups,
but it was challenging to use automated methods to annotate function to many of
the less abundant but diverse macroinvertebrate families detected using COI
metabarcoding. Despite metabarcoding methods facilitating genus- and species-
level taxonomic assignments, databases of functional metrics for
macroinvertebrates at this taxonomic level are still incomplete [46]. The influence of
different environmental variables (e.g. Metals and aromatic hydrocarbons in
sediment as well as ammonia in lake water) on the relative abundance of
functional feeding groups has implications on the trophic system of the lake as a
whole. An increase or decrease in sediment and water pollutant loads will likely
influence a shift in the taxa exhibiting different FFG guilds. Assigning FFG at
more conservative taxonomic ranks (i.e., family) enables easy integration of both
metabarcoding and morphological taxonomic lists whilst maximizing functional
assignments.

By using a multi-marker approach, we can begin to close knowledge gaps
regarding how water quality and loading of elements in sediment affect taxa at
various trophic levels [47,48]. In this study, metals were found to be one of the more
influential sediment contaminants explaining variation in overall
macroinvertebrate richness using COI metabarcoding as well as variation in the
ciliate group Haptoria. In Toronto Harbour, metals such as cadmium, chromium,
copper, lead, mercury, and zinc were found to have levels that exceeded the
Canadian Federal Probable Effect Level. In this study, we have samples from 5
stations from inner harbour centre that represent a clear gradient of increasing

metal concentrations, decreasing macroinvertebrate richness detected using COI
metabarcoding, and increasing relative read abundance of the ciliate group
Haptoria using 18S metabarcoding. Ciliates, despite their ubiquity in freshwater
systems, the key position they play in trophic webs (feed on bacteria, algae,
other protists while also being consumed by other meiofauna) and known
sensitivity to a range of water quality types (low to highly polluted), they are often
a neglected in water quality assessments [49].

Conclusion

DNA methods can readily be applied for both routine lake community
biomonitoring and targeted monitoring for invasive, threatened, and/or exploited
lake species, integrated with periodic morphological assessments to supplement
routine DNA monitoring with abundance measures. Overall, for taxonomically
comprehensive assessments of lake communities, it is imperative to apply robust
high-throughput methods, such as DNA metabarcoding, to increase the
resolution of biodiversity data and to understand species-specific responses to
environmental gradients in the face of various perturbations and climate warming
effects.

399
400
401 **Methods**
402

Study stations and sampling design

Sample stations in Toronto Harbour have been previously established by
the Ministry of the Environment and Climate Change Resources in 1971, to track
improvements in sediment contaminant concentrations and benthic invertebrate
species composition and density [50] (Table S1; Figure S1).

In October 2018, water, sediment and benthos samples were collected
from Toronto's Inner Harbour over a three-day period, from a subset of 25
stations [17]. At each station, environmental data was collected, followed by the
collection of surface and overlying water, sediment and benthos samples in
order. Samples were collected following the Canadian Aquatic Biomonitoring
Network (CABIN) Open Water sampling protocols for collection of benthic
samples [51,52].

Water samples were analyzed for major ions, nutrients, temperature,
conductivity, pH and dissolved oxygen (see supporting information for technical
details). Sediment samples were analyzed for trace metals, PCB aroclors, total
PCBs and PCB congeners, chlorinated pesticides, chlorobenzenes, technical
toxaphene (insecticide) and toxaphene congeners/parlars. See supporting
information for technical details on water, sediment and benthos sampling and
collection for both DNA and morphological identification.

Morphology-based taxonomic identification of benthos

Invertebrates in the benthic community samples were sorted, identified to
the family level, and counted by EcoAnalysts, Moscow, ID.

DNA sample homogenization and DNA extraction of benthos samples

Samples were transferred from whirl-packs to 50mL conical tubes, using
molecular biology grade water to rinse whirl packs to ensure the entire sample
was transferred. Samples were centrifuged to collect sediment at the bottom, and
excess water was removed. Approximately 0.3g was then directly subsampled
into 2mL bead tubes included within the Qiagen PowerSoil kit. Samples were
extracted according to manufacturer's protocol eluting with 50uL Buffer C6. Each
batch (~95 samples) included one negative extraction control where no tissue
was included. The remaining mass was stored in the Falcon tubes at -20°C as a
voucher.

Library preparation and high-throughput sequencing

Two fragments within the standard COI DNA barcode region and one fragment within the 18S (eukaryote) region were amplified with the following previously optimized and validated primer sets: (LCO1490/230_R [~230 bp] called F230R, mICOIintF/jgHCO2198 [~313 bp] called ml-jg and Uni18SF/ Uni18SR [~600 bp] called Uni18S [53–57], using a two-step PCR amplification regime. The first PCR used COI and 18S specific primers and the second PCR involved Illumina-tailed primers. Uni18S PCR cycling conditions are as follows: 95°C for 3 minutes, followed by 35 cycles of 94°C for 30 seconds, 52°C for 30 seconds, 72°C for 90 seconds, and a final extension of 72°C for 8 minutes. COI PCR cycling conditions for both fragments were: 95°C for 5 minutes, followed by 35 cycles of 94°C for 40 seconds, 46°C for 1 minute, 72°C for 30 seconds, and a final extension of 72°C for 5 minutes. Amplification was visually confirmed through a 1.5% agarose gel electrophoresis. Amplicons were purified with a MinElute PCR purification kit (Qiagen), quantified with fluorometry using a QuantIT PicoGreen dsDNA assay kit (Invitrogen), and normalized to the same concentration prior to dual indexing with a Nextera Index Kit (Illumina). Indexed samples were pooled and purified through magnetic bead purification. The library was then quantified with the QuantIT PicoGreen dsDNA assay kit, and average fragment length was determined on an Agilent Bioanalyzer 2100 with a DNA 7500 chip. The library was then diluted to 4nM based on the concentration and average fragment length and sequenced on an Illumina MiSeq using a v3 chemistry kit (2 x 300 cycles). A 10% PhiX control spike-in was used to ensure sequence diversity.

Bioinformatic processing

Illumina MiSeq paired-end reads were processed using the MetaWorks-1.0.0 pipeline available from https://github.com/terrimporter/MetaWorks [58]. MetaWorks is an automated Snakemake [59] bioinformatic pipeline that runs in a

conda [60] environment. Details of the sequence processing steps of
METAWORKS is described in the Supplementary Material. Further analysis of
leave-one-sequence-out testing with the RDP Classifier conducted with our COI
and 18S reference sets allowed us to assess expected taxonomic assignment
accuracy according to metabarcode length and taxonomic rank [61–63]. Using a
leave one sequence out approach during classifier validation, we determined that
for a ~ 200 bp COI metabarcode, taxonomic assignments are ~ 90% correct at
the species rank, 95%+ correct at the genus - family ranks using the appropriate
bootstrap support cutoffs, and 99%+ correct at more inclusive ranks (e.g. order -
kingdom) assuming the query taxa are present in the reference database [61].
Using a ~ 200 bp 18S metabarcode, taxonomic assignments are ~ 80%+ correct
at the genus - order ranks using the appropriate bootstrap support cutoffs and
about 95%+ correct at more inclusive ranks (class - domain) assuming the query
taxa are present in the reference database.

Statistical analyses

All statistical analyses were conducted in RStudio v 1.1.456 running R v
3.5.1 [64] and plots were created with ggplot2 [65]. Sequence variants recovered
from field blanks comprised 2.7% of unique COI macroinvertebrate sequence
variants and 5% of unique 18S eukaryote sequence variants and were removed
from all subsequent analyses.

Resolution of taxonomic assignments were compared by recoding
taxonomic assignments to: species=1, genus=2, family=3, etc. Sampling effort in
metabarcoding and morphological samples were assessed using rarefaction.
Relative abundance were compared by calculating the proportion or percentage
of reads per taxon per station for metabarcoding data and by calculating the
number of individuals per taxon per station for morphological methods. Prior to
NMDS or PCA matrices were converted to proportions or standardized using a
Hellinger transformation.

For a fair comparison of macroinvertebrate metrics using COI metabarcoding and morphological methods, we limited COI results to taxa in the phyla normally detected using morphological methods: Arthropoda, Annelida, Mollusca, Cnidaria, and Platyhelminthes. We compared several macroinvertebrate diversity metrics for morphology-based and metabarcoding methods such as richness at the species level for macroinvertebrate COI and morphological samples, richness at the genus level for (non-macroinvertebrate) eukaryote 18S samples and read abundance of the top 5 most abundant families per sampling method.

We used principal components analysis (PCA) 'rda' function in the vegan package to assess the variance between macroinvertebrate COI metabarcode and morphological samples [66]. Only vectors for the most strongly correlated species were plotted for clarity. Also using vegan, we used non-metric multi-dimensional scaling (NMDS) analyses to visualize beta diversity based on Bray Curtis dissimilarities using the 'metaMDS' function using 3 dimensions and fitted correlated environmental variables using the 'envfit' function using 999 permutations if correlations were greater than 0.30 and the p-value < 0.05. Goodness of fit calculations and Shephard's curve were calculated using the vegan 'goodness' and 'stressplot' functions.

Water physical-chemical features were measured from samples collected at the bottom of the harbour and contaminants were measured from sediment samples. For each set of measurements for organic contaminants/metals, values for each individual chemical/element was summed across each major contaminant group [17]. We tested each predictor for normality using a Shapiro-Wilk Test using the 'shapiro.test' function in R. Skewness was checked using the 'skewness' function from the moments package [67]. Each individual predictor variable was then transformed (square root, log10, 1/x) as needed to better meet normality assumptions in hierarchical partitioning using a Gaussian model. Predictors were standardized using z-scores and centered prior to analysis. We

checked for collinearity among water and sediment variables using Pearson
correlation coefficients with a cutoff of 0.70, using the 'rcorr' function from the
Hmisc package (Figure S2)[68].

For each family, we added primary functional feeding guild (FFG)
annotations based on the EPA Freshwater Biological Traits Database [69]. This
system recognizes 7 feeding modes: collector-filterer (CF), collector-gatherer
(CG), herbivore (scraper) (HB), parasite (PA), predator (piercer, engulfer) (PR),
shredder (SH), and Other. Remaining missing family annotations were added
using information compiled from the Taxa and Autecology Database for
Freshwater Organisms available from freshwaterecology.info [70]. For this
database, preference was given to feeding type annotations by Moog [71]. Any
further missing annotations were then added by checking feeding habit
annotations [72]. Since individuals were identified to the family level, species in a
family can exhibit more than one feeding type and the presence of multiple
feeding types per family were recorded when this was the case. We multiplied a
family x station matrix containing read/individual counts by each feeding mode in
a family x FFG matrix containing 1's or 0's indicating the presence of the feeding
mode in a family. For each feeding mode, total read counts per station were
recorded and combined into a single FFG x station matrix. Out of 61 families
detected using metabarcoding, 23 (38%) were assigned feeding types. Out of 30
families detected using morphology, 24 (80%) were assigned feeding types.
Further analyses using FFGs represent the proportion of all reads/individuals that
could be both taxonomically assigned to family and functionally assigned a
feeding mode.

We used hierarchical partitioning to regress each macroinvertebrate
metric, individually, on the environmental predictors to identify which water
physical-chemical features or sediment contaminants explain the most variance
independently of the others. This was done using the 'hier.part' package in R [73].

Nutrient/major ion predictors measured from water collected at the bottom of the harbour were analyzed separately from sediment contaminants. Macroinvertebrate diversity metrics included macroinvertebrate species richness based on COI and morphological methods, genus richness based on (non-macroinvertebrate) eukaryote 18S metabarcoding, as well as the relative abundance of the top 5 families based on COI, morphological, and 18S. Functional metrics included the FFGs calculated for macroinvertebrate families using COI and morphological methods. All metrics were standardized using a Hellinger transformation using the 'decostand' function in vegan. Hierarchical partitioning significance was assessed using a randomization test, 1000 replicates, and calculating a goodness of fit measure based on log-likelihood.

References

1. Breed, M. F. *et al.* The potential of genomics for restoring ecosystems and biodiversity. *Nat. Rev. Genet.* **20**, 615–628 (2019).

2. Carpenter, S. R., Stanley, E. H. & Vander Zanden, M. J. State of the world's freshwater ecosystems: Physical, chemical, and biological changes. *Annu. Rev. Environ. Resour.* **36**, 75–99 (2011).

3. Geist, J. Integrative freshwater ecology and biodiversity conservation. *Ecol. Indic.* **11**, 1507–1516 (2011).

4. Jeppesen, E., Søndergaard, M., Meerhoff, M., Lauridsen, T. L. & Jensen, J. P. Shallow lake restoration by nutrient loading reduction--some recent findings and challenges ahead. *Hydrobiologia* (2007).

5. Søndergaard, M. & Jeppesen, E. Anthropogenic impacts on lake and stream

ecosystems, and approaches to restoration. *J. Appl. Ecol.* **44**, 1089–1094

(2007).

6. Marburg, A. E., Turner, M. G. & Kratz, T. K. Natural and anthropogenic

variation in coarse wood among and within lakes. *J. Ecol.* **94**, 558–568

(2006).

7. Schindler, D. W. Recent advances in the understanding and management of

eutrophication. *Limnol. Oceanogr.* **51**, 356–363 (2006).

8. Lau, S. S. S. & Lane, S. N. Continuity and change in environmental systems:

the case of shallow lake ecosystems. *Prog. Phys. Geogr. Earth Environ.* **25**,

178–202 (2001).

9. Brinkhurst, R. O. Distribution and abundance of Tubificid (Oligochaeta)

species in Toronto harbour, Lake Ontario. *J. Fish. Res. Board Can.* **27**, 1961–

1969 (1970).

10. Wood, L. W. & Chua, K. E. Glucose flux at the sediment-water interface of

Toronto Harbour, Lake Ontario, with reference to pollution stress. *Can. J.*

Microbiol. **19**, 413–420 (1973).

11. Nriagu, J. O., Wong, H. K. T. & Snodgrass, W. J. Historical records of metal

pollution in sediments of Toronto and Hamilton harbours. *J. Gt. Lakes Res.*

(1983).

12. Toronto & Region Remedial Action Plan. *Metro Toronto and Region*

Remedial Action Plan. (1989).

13. Dahmer, S. C., Matos, L. & Morley, A. Restoring Toronto's waters: Progress

toward delisting the Toronto and Region Area of Concern. *Aquat. Ecosyst.*

Health Manag. **21**, 229–233 (2018).

14. Munawar, M., Norwood, W., McCarthy, L. & Mayfield, C. In situ

bioassessment of dredging and disposal activities in a contaminated

ecosystem: Toronto Harbour. (1989) doi:10.1007/978-94-009-1896-2_62.

15. Dahmer, S. C., Matos, L. & Jarvie, S. Assessment of the degradation of

aesthetics Beneficial Use Impairment in the Toronto and region Area of

Concern. *Aquat. Ecosyst. Health Manag.* **21**, 276–284 (2018).

16. Metro Toronto and Region Remedial Action Plan. *Within Reach: 2015*

Toronto an Region Remedial Action Plan Progress Report. (2016).

17. Burniston, D. & Waltho, J. *Report on Sediment Quality in the Toronto Inner*

Harbour 2007. (2011).

18. Elbrecht, V., Vamos, E. E., Meissner, K., Aroviita, J. & Leese, F. Assessing

strengths and weaknesses of DNA metabarcoding-based macroinvertebrate

identification for routine stream monitoring. *Methods Ecol. Evol.* **8**, 1265–

1275 (2017).

19. Emilson, C. E. *et al.* DNA metabarcoding and morphological

macroinvertebrate metrics reveal the same changes in boreal watersheds

across an environmental gradient. *Sci. Rep.* **7**, 12777 (2017).

20. Aylagas, E., Borja, Á., Muxika, I. & Rodríguez-Ezpeleta, N. Adapting

metabarcoding-based benthic biomonitoring into routine marine ecological

status assessment networks. *Ecol. Indic.* **95**, 194–202 (2018).

21. Bush, A. *et al.* Studying ecosystems with DNA metabarcoding: lessons from

biomonitoring of aquatic macroinvertebrates. *Front. Ecol. Evol.* **7**, 434 (2019).

22. Serrana, J. M., Miyake, Y., Gamboa, M. & Watanabe, K. Comparison of DNA

metabarcoding and morphological identification for stream macroinvertebrate

biodiversity assessment and monitoring. *Ecol. Indic.* (2019).

23. Fernández, S., Rodríguez-Martínez, S., Martínez, J. L., Garcia-Vazquez, E. &

Ardura, A. How can eDNA contribute in riverine macroinvertebrate

assessment? A metabarcoding approach in the Nalón River (Asturias,

Northern Spain). *Environ. DNA* **1**, 385–401 (2019).

24. Hajibabaei, M. *et al.* Watered-down biodiversity? A comparison of

metabarcoding results from DNA extracted from matched water and bulk

tissue biomonitoring samples. *PLOS ONE* **14**, e0225409 (2019).

25. Baird, D. J. & Hajibabaei, M. Biomonitoring 2.0: a new paradigm in

ecosystem assessment made possible by next-generation DNA sequencing.

Mol. Ecol. **21**, 2039–2044 (2012).

26. Hajibabaei, M., Baird, D. J., Fahner, N. A., Beiko, R. & Golding, G. B. A new

way to contemplate Darwin's tangled bank: how DNA barcodes are

reconnecting biodiversity science and biomonitoring. *Philos. Trans. R. Soc. B*

Biol. Sci. **371**, 20150330 (2016).

27. Beermann, A. J., Zizka, V. M. A., Elbrecht, V., Baranov, V. & Leese, F. DNA

metabarcoding reveals the complex and hidden responses of chironomids to

multiple stressors. *Environ. Sci. Eur.* **30**, 26 (2018).

28. Bush, A. *et al.* DNA metabarcoding reveals metacommunity dynamics in a

threatened boreal wetland wilderness. *Proc. Natl. Acad. Sci.* **117**, 8539–8545

(2020).

29. Compson, Z. G. *et al.* Chapter two - Linking DNA metabarcoding and text

mining to create network-based biomonitoring tools: a case study on boreal

wetland macroinvertebrate communities. in *Advances in Ecological Research*

(eds. Bohan, D. A., Dumbrell, A. J., Woodward, G. & Jackson, M.) vol. 59 33–

74 (Academic Press, 2018).

30. Fernandes, K. *et al.* DNA metabarcoding—a new approach to fauna

monitoring in mine site restoration. *Restor. Ecol.* **26**, 1098–1107 (2018).

31. Fernandes, K. *et al.* Invertebrate DNA metabarcoding reveals changes in

communities across mine site restoration chronosequences. *Restor. Ecol.* **27**,

1177–1186 (2019).

32. Poikane, S. *et al.* Benthic macroinvertebrates in lake ecological assessment:

A review of methods, intercalibration and practical recommendations. *Sci.*

Total Environ. **543**, 123–134 (2016).

33. Macher, J.-N. *et al.* Comparison of environmental DNA and bulk-sample

metabarcoding using highly degenerate cytochrome c oxidase I primers. *Mol.*

Ecol. Resour. **18**, 1456–1468 (2018).

34. Marshall, N. T. & Stepien, C. A. Macroinvertebrate community diversity and

habitat quality relationships along a large river from targeted eDNA

metabarcode assays. *Environ. DNA* **2**, 572–586 (2020).

35. Metro Toronto and Region Remedial Action Plan. *Updates on Actions 2013-*
2014. (2013).

36. López-López, E. & Sedeño-Díaz, J. E. Biological indicators of water quality:
The role of fish and macroinvertebrates as indicators of water quality. in
Environmental Indicators (eds. Armon, R. H. & Hänninen, O.) 643–661
(Springer Netherlands, 2015). doi:10.1007/978-94-017-9499-2_37.

37. Berry, O. *et al.* A comparison of morphological and DNA metabarcoding
analysis of diets in exploited marine fishes. (2015).

38. Sweeney, B. W., Battle, J. M., Jackson, J. K. & Dapkey, T. Can DNA
barcodes of stream macroinvertebrates improve descriptions of community
structure and water quality? *J. North Am. Benthol. Soc.* **30**, 195–216 (2011).

39. Banerji, A. *et al.* Spatial and temporal dynamics of a freshwater eukaryotic
plankton community revealed via 18S rRNA gene metabarcoding.
Hydrobiologia **818**, 71–86 (2018).

40. Porter, T. M. *et al.* Widespread occurrence and phylogenetic placement of a
soil clone group adds a prominent new branch to the fungal tree of life. *Mol.*
Phylogenet. Evol. **46**, 635–644 (2008).

41. Rosling, A. *et al.* Archaeorhizomycetes: unearthing an ancient class of
ubiquitous soil fungi. *Science* **333**, 876–879 (2011).

42. Mandaville, S. M. *Benthic Macroinvertebrates in Freshwaters- Taxa*
Tolerance Values, Metrics, and Protocols. 128 http://lakes.chebucto.org/H-
1/tolerance.pdf (2002).

43. Trzcinski, M. K. *et al.* The effects of food web structure on ecosystem function

exceeds those of precipitation. *J. Anim. Ecol.* **85**, 1147–1160 (2016).

44. Liu, X. & Wang, H. Contrasting patterns and drivers in taxonomic versus

functional diversity, and community assembly of aquatic plants in subtropical

lakes. *Biodivers. Conserv.* (2018).

45. Kovalenko, K. E., Brady, V. J., Ciborowski, J. J. H., Ilyushkin, S. & Johnson,

L. B. Functional changes in littoral macroinvertebrate communities in

response to watershed-level anthropogenic stress. *PLOS ONE* **9**, e101499

(2014).

46. Luiza-Andrade, A., Montag, L. F. de A. & Juen, L. Functional diversity in

studies of aquatic macroinvertebrates community. *Scientometrics* **111**, 1643–

1656 (2017).

47. MacMillan, G. A., Chételat, J., Heath, J. P., Mickpegak, R. & Amyot, M. Rare

earth elements in freshwater, marine, and terrestrial ecosystems in the

eastern Canadian Arctic. *Environ. Sci. Process. Impacts* **19**, 1336–1345

(2017).

48. Pastorino, P. *et al.* Macrobenthic invertebrates as tracers of rare earth

elements in freshwater watercourses. *Sci. Total Environ.* **698**, 134282 (2020).

49. Kulaš, A. *et al.* Ciliates (Alveolata, Ciliophora) as bioindicators of

environmental pressure: A karstic river case. *Ecol. Indic.* **124**, 107430 (2021).

50. Persaud, D., Lomas, T., Boyd, D. & Mathai, S. *Historical development and*

quality of the Toronto waterfront sediments. (1985).

51. Milani, D. & Grapentine, L. *Assessment of sediment quality in the Bay of*

Quinte Area Of Concern. (2000).

52. Reynoldson, T. B., Bailey, R. C., Day, K. E. & Norris, R. H. Biological

guidelines for freshwater sediment based on BEnthic Assessment of

SedimenT (the BEAST) using a multivariate approach for predicting biological

state. *Aust. J. Ecol.* (1995).

53. Geller, J., Meyer, C., Parker, M. & Hawk, H. Redesign of PCR primers for

mitochondrial cytochrome c oxidase subunit I for marine invertebrates and

application in all-taxa biotic surveys. *Mol. Ecol. Resour.* **13**, 851–861 (2013).

54. Leray, M. *et al.* A new versatile primer set targeting a short fragment of the

mitochondrial COI region for metabarcoding metazoan diversity: application

for characterizing coral reef fish gut contents. *Front. Zool.* **10**, 34 (2013).

55. Zhan, A. *et al.* High sensitivity of 454 pyrosequencing for detection of rare

species in aquatic communities. *Methods Ecol. Evol.* **4**, 558–565 (2013).

56. Gibson, J. *et al.* Simultaneous assessment of the macrobiome and

microbiome in a bulk sample of tropical arthropods through DNA

metasystematics. *Proc. Natl. Acad. Sci.* **111**, 8007–8012 (2014).

57. Gibson, J. F. *et al.* Large-scale biomonitoring of remote and threatened

ecosystems via high-throughput sequencing. *PLOS ONE* **10**, e0138432

(2015).

58. Porter, T. M. & Hajibabaei, M. METAWORKS: A flexible, scalable

bioinformatic pipeline for multi-marker biodiversity assessments. *bioRxiv*

2020.07.14.202960 (2020).

59. Köster, J. & Rahmann, S. Snakemake—a scalable bioinformatics workflow
engine. *Bioinformatics* **28**, 2520–2522 (2012).

60. Anon. *Conda.* (2016).

61. Porter, T. M. & Hajibabaei, M. Automated high throughput animal CO1
metabarcode classification. *Sci. Rep.* **8**, 4226 (2018).

62. Porter, T. M. *Eukaryote CO1 Reference Set For The RDP Classifier.*
(Zenodo, 2017). doi:10.5281/zenodo.4741447.

63. Porter, T. M. *SILVA 18S Reference Set For The RDP Classifier.* (Zenodo,
2018). doi:10.5281/zenodo.4741433.

64. R Core Team. *R: A language and environment for statistical computing.*
(2020).

65. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis.* (Springer-Verlag,
2009). doi:10.1007/978-0-387-98141-3.

66. Oksanen, J. *et al. vegan: Community Ecology Package.* (2020).

67. Komsta, L. & Novomestky, F. *moments: Moments, cumulants, skewness,*
kurtosis and related tests. (2015).

68. Harrell, F. & functions), C. D. (contributed several functions and maintains
latex. *Hmisc.* (2021).

69. U.S. Environmental Protection Agency. *Freshwater Biological Traits*
Database. (2012).

70. Schmidt-Kloiber, A. & Hering, D. www.freshwaterecology.info – An online tool
that unifies, standardises and codifies more than 20,000 European freshwater
organisms and their ecological preferences. *Ecol. Indic.* **53**, 271–282 (2015).

71. Moog, O. Fauna Aquatica Austriaca - Catalogue for autecological

classification of Austrian aquatic organisms. (1995).

72. Usseglio-Polatera, P., Tachet, H., Richoux, P. & Bournaud, M. *Invertébrés*

d'eau douce - systématique, biologie, écologie. (2000).

73. Nally, R. M. & Walsh, C. J. Hierarchical Partitioning Public-domain Software.

Biodivers. Conserv. (2004) doi:10.1023/B:BIOC.0000009515.11717.0b.

757
Acknowledgments

We would like to thank Adam Morden, Jonathan Paynter, Tina Mamone, and

Jesse Baillargeon from ECCC for field support and members of the Hajibabaei

lab for help processing samples. Funding for sampling and data collection for this

project was provided by Environment and Climate Change Canada's Great

Lakes Action Plan in conjunction with the government of Canada through

Genome Canada and Ontario Genomics.

765
766
Author contributions

MM and MH designed the study, MM collected samples, MTGW processed and

sequenced samples, CVR and TMP conducted bioinformatic and statistical

analyses with assistance from KMM, CVR wrote the manuscript with help from all

co-authors.

Data availability

Raw sequences will be available from NCBI SRA on acceptance. The

bioinformatic pipeline MetaWorks v1 is available from GitHub at

https://github.com/terrimporter/MetaWorks/releases/tag/v1.0.0, the COI Classifier

v4 and the 18S Classifier v4.1 we used are available from GitHub at

https://github.com/terrimporter/CO1Classifier and

https://github.com/terrimporter/18SClassifier respectively. The code used to

generate figures, including infiles, will be available from GitHub on acceptance.

Additional information

The authors declare that they have no known competing financial interests or

personal relationships that could have appeared to influence the work reported in

this paper.